Genetic Analysis of F1 Cluster Phages that Infect *Mycobacterium smegmatis* Identifies Two Distinct Holin-Like Proteins that Regulate the Host Lysis Event

Running Title: Two Distinct Phage Holin Proteins Regulate the Host Lysis Event in *M. smegmatis*

Richard S. Pollenz[a#], Kira Ruiz-Houston[b], Wynter Dean[a] and Loc Nguyen[a]

[a]Department of Molecular Bioscience, University of South Florida, Tampa, FL

[b]Department of Microbiology, University of Wisconsin, Madison, WI

[#]Corresponding Author:

Richard S Pollenz

Department of Molecular Biosciences

University of South Florida

4202 E Fowler Ave ISA2015

Tampa, FL 33620

pollenz@usf.edu

813-767-3984

ABSTRACT

Phages Girr and NormanBulbieJr (NBJ) infect Gram-positive *Mycobacterium smegmatis* mc^2

155. Both phages contain conserved lysis cassettes that harbor two endolysin genes (*lysin A*

and *lysin B*) and two genes encoding transmembrane domain (TMD) holin-like proteins. The

first holin-like protein, termed LysF1a is 88 amino acids, has two TMDs and a predicted N-in-C-

in membrane topology. The second, termed LysF1b, has a single N-terminal TMD and a

predicted N-out-C-in topology making it distinct from the 1TMD type III holins or spanins in size

and membrane topology. Deletion of either *lysF1a* or *lysF1b* results in lysis defect phenotypes

manifest by reduced plaque size and changes to lysis timing in liquid culture. Deletion of both

lysF1a and *lysF1b* genes is lethal to phage propagation supporting holin function. Phages

expressing only the LysF1b protein trigger lysis prematurely when exposed to energy poisons

while phages expressing only LysF1a do not show early triggering. Lysis recovery mutants were

isolated from phages lacking the *lysF1b* gene and these mutants generated wild type plaque

size but triggered lysis prematurely and showed ~65% reductions in burst size. Genome

sequencing identified different point mutations that mapped to TMD1 or the C-terminal region of

the *lysF1a* gene. Finally, infection of an *M. smegmatis* strain that does not produce lipomannan

and lipoarabinomannan by either wild type phages or phages carrying the *lysF1b* deletion

showed modest plaque size increases and did not fully complement the lysis defect of phages

lacking the *lysF1b* gene. Collectively, the findings show that both LysF1a and LysF1b function

as holins and are required for efficient bacterial lysis. LysF1a does not function as an antiholin

but requires the expression of the LysF1b protein for efficient lysis functioning. The requirement

of two holins with distinct membrane topologies for host lysis has not been described previously.

INTRODUCTION

With an estimated 10^{31} different bacteriophage particles in the world, the pathway whereby dsDNA phages lyse their bacterial hosts after infection and replication is one of the most common cell fates on earth [1]. Extensive biochemical and genetic work with dsDNA phages that infect Gram-negative hosts has established a general three-step model for the triggering of the lysis pathway that involves three distinct sets of proteins. The first are endolysin/s that have defined catalytic activity to lyse the peptidoglycan (PG) layer [2]. The second are holin or antiholin proteins that have between 1-4 transmembrane domain helices (TMD). These proteins work together to, 1) control the timing of the lysis event following phage infection [3, 4], 2) assist with the destabilization of the membrane proton motive force (PMF) [3, 4] and 3) in the canonical holin models, generate large 1μm "pores" in the inner membrane (IM) to allow the exit of the endolysin enzymes to the periplasmic space where they can access and cleave the PG layer [5, 6]. Finally, there are secreted membrane proteins and lipoproteins termed spanins that are associated with the IM and outer membrane (OM). There proteins interact after the PG layer is cleaved to generates the lesion across the IM and OM that allows for the efficient release of the phage progeny [7-9].

dsDNA phages such as lambda, P2, and phage 21 all have genomes < 50,00bp and have a clustered lysis cassette that contain all the genes necessary to mediate the bacterial lysis event [10-12]. Phage lambda encodes the canonical type I holin protein that has 3TMDs and forms large oligomers that create 1μm holes in the IM to allow exit of the endolysin from the cytoplasm to the periplasmic space [12]. Phage 21 defines the type II pinholin lysis pathway and utilizes a 2TMD holin protein that forms ~1,000 heptamers in the IM. The pinholins create only nanometer size holes that are not big enough to allow exit of the endolysin, and they function to disrupt the PMF and trigger lysis. Thus, phages that encode pinholins also encode endolysins that carry secretion-arrest-release (SAR) signals and the SAR-endolysin is held in an inactive

state in the IM prior to triggering. The disruption of the PMF causes the release of the IM

tethered SAR-endolysin to the periplasmic space where it cleaves the PG [4, 13, 14]. In both

lambda and phage 21, a single open reading frame with two in-frame start codons (dual start

motif) encodes two holin proteins that only differ by the addition of 1-2 N-terminal amino acids

[10, 11, 14, 15]. The longer version of each holin functions as the inactive antiholin and in both

cases a percentage of the holin forms dimers with the antiholin that holds that holin in an

inactive state prior to triggering. Upon triggering, the antiholin assumes active holin function

greatly increasing the amount of active holin protein available for hole formation. In phages with

large genomes, such as P1 and T4, the holin, antiholin, endolysin and spanin genes can be

bioinformatically characterized throughout the genome and are not in a defined cassette. The

type III holin from phage T4 has a single N-terminal TMD with an N-in-C-out topology with the

bulk of the sequence oriented in the periplasmic space [16, 17]. T4 phage encodes two different

antiholin proteins that modulate the holin lysis timing and participate in the lysis inhibition activity

of phage T4 [16, 17]. Thus, a common theme in the phage-mediated lysis pathway of lambda,

P1, P2 and T4 is the utilization of holins and antiholins to regulate the lysis event. However,

another variation of the lysis pathway has recently been shown for the Enterococcal phage Mu

[18]. Mu does not utilize canonical holins and antiholins in the bacterial lysis pathway and

instead uses a novel 1TMD protein termed a releasin protein to release the SAR-endolysin from

the IM so it can cleave the PG [18]. As with lambda and phage 21, Mu contains canonical

spanins to create the final lesion for progeny release once the PG has been cleaved. It is

speculated that Mu does not utilize the holin/antiholin regulated lysis mechanism because it

copies its genome via replicative transposition during the lytic cycle and has less selective

pressure due to transposition immunity [18]. This pathway adds yet another variation in phage-

mediated bacterial lysis and it is striking that all the *E. coli* phages that have been rigorously

studied utilize a similar set of tools to complete the lysis pathway through distinct molecular

mechanisms.

There are 1000s of annotated dsDNA phages that infect Gram-positive hosts, yet the

molecular mechanism that these pages use to lyse their hosts is not resolved to the level of the

E. coli phages. Although it is hypothesized that the lysis mechanism in these phages will utilize

a similar set of proteins as described for phages that infect Gram-negative bacteria, the bulk of

biochemical and genetic work has focused more on the enzymes that target the cell wall than on

the holins, possibly because these enzymes may be employed as antibiotic agents [19-21].

Gram-positive bacteria lack a formal OM and instead have a much more complex PG layer and

an extended structural cell wall components beyond the PG such as arabinogalactans and

mycolic acids [22-25]. Thus, addition of an endolysin to Gram-positive bacteria can result in cell

death. Bioinformatic, biochemical and genetic analysis has identified several types of phage-

encoded endolysins that target different regions of the cell wall. The first type of endolysin is

termed lysin A and targets distinct domains of the peptidoglycan [20, 26]. These enzymes are

highly modular, do not contain canonical SAR domains, and deletion studies show that the lysin

A is essential to phage propagation [27, 28]. It is unknown if the lysin A is released to the

peptidoglycan layer through holins since some studies show that exogenous expression of

selected lysin A enzymes can be cytotoxic to *M. smegmatis* [20, 29-32]. Many phages that infect

Gram-positive hosts also contain a second endolysin termed lysin B that targets the linkages in

the mycolic acid layer of the cell wall [27, 30, 33, 34]. It has been hypothesized that lysin B

provides a fitness advantage for optimal bacterial lysis and phage release since deletion of lysin

B results in viable phages with reduced plaque and burst size [27].

In phages that infect Gram-positive hosts the presence of genes that encode holin-like

proteins has primarily been characterized by bioinformatic analysis and many of the phages

contain more than one gene encoding a TMD protein that could serve as a holin or antiholin.

Currently there are reports on host lysis by phages that infect *Staphylococcus aureus* [35-37],

Bacillus sp. [38], *Lactococcal sp.* [39], *Oenococcus oeni* [40, 41], *Streptococcus sp* [42],

Gordonia rubripertincta [43-45] and *Mycobacterium sp.* [46-53]. The holin function for the

encoded TMD proteins in most of these reports is based primarily on the proximity of the TMD

gene/s near an endolysin. Where biochemical work is reported, the majority has been carried

out using exogenous expression of the TMD genes in *E. coli*. However, there are several

reports on phages that infect *Mycobacterium smegmatis* where gene deletion has been utilized

to evaluate lysis function of the predicted holin proteins.

The first study evaluated two genes encoding TMD proteins that are found directly

downstream of the *lysin A* and *lysin B* in the F1 cluster phage Ms6 [48-50]. Ms6 gene *26*

encodes an 88 amino acid protein with 2TMDs while gene *27* encodes a 124 amino acid protein

with 1TMD. Deletion of gene *26* resulted in reduced lysis timing, reduced plaque size but wild

type burst size, while deletion of gene *27* resulted in delayed lysis timing, increased plaque size

and increased burst size. However, there are limitations to these findings because holin function

was not verified using energy poisons and the mutant phages were not tested in liquid lysis

assays in the *M. smegmatis* host versus *E. coli*. In addition, the experiments to assess function

and interaction of the two proteins were carried out using recombinant *E. coli* expression

systems that are not physiologic. Thus, while it appears that both proteins are needed for the

lysis event, it is unclear how each is contributing to the lysis of the *M. smegmatis* host.

The second set of studies evaluated phage D29, a cluster A2 phage that also infects *M.*

smegmatis [51-54]. D29 has a lysis cassette with a *lysin A* (gene *10*), 2TMD holin (gene *11*) and

a *lysin B* (gene *12*). It is important to note that the 2TMD holin of D29 is not an ortholog to the

2TMD protein in Ms6 and differs significantly in amino acid sequence and size. The lysis

cassette that is also distinct from most other actinobacteriophages in that it has no other genes

encoding holin-like TMD proteins within or near the putative lysis cassette. Deletion of gene *11*

results in a viable phage with a different phenotype that the 2TMD deletion in Ms6 [51].

D29_Δ*11* phage shows a modest 30% reduced plaque size, slightly delayed lysis timing and

reduced burst size [51]. The lack of a more severe lysis phenotype is surprising given that gp11

is the only defined holin, especially since the D29 lysin A does not have a defined signal peptide

or SAR-like domain for export [31, 32]. However, it has been reported that the D29 lysin A may

be exported prior to lysis in a yet to be defined pathway that appears to be independent of the

holin [31, 32]. Alternatively, the modest lysis phenotype could also indicate that D29 has

additional holin-like genes that are not located within the defined "lysis cassette" region. Such a

scenario has recently been described in a bioinformatic study of the lysis cassettes of 77

phages that infect the Gram-positive *Gordonia rubripertincta* strain [44]. This study showed that

there was a high level of diversity of the lysis cassette gene organization and that putative holin-

like genes could be identified in genomic regions distal to the endolysin containing lysis

cassettes [44]. This study also showed that most of the phages evaluated had at least two holin-

like genes.

In summary, there is limited biochemical and genetic analysis of host lysis pathways by

phages that infect Gram-positive hosts, especially within the host bacteria. Consistent findings

suggest: 1) the putative holin genes do not appear to have dual start motifs like those identified

in lambda and phage 21; 2) that most annotated phages have multiple genes predicted to

encode holin-like TMD proteins; 3) that the TMD proteins are predicted to have 1, 2, or 4 TMDs;

and 4) that the putative holin-like genes can be bioinformatically identified outside of lysis

cassettes even in phages that have compact genomes of < 60,000bp. Of these findings, one of

the most intriguing is the presence of multiple genes that encode predicted holin-like TMD

proteins. Since antiholins have not been experimentally identified in phages that infect Gram-

positive hosts, one interpretation of these data are that holins and antiholins are expressed from

distinct genes as shown for *E. coli* phages P2 and T4 [16, 17, 55, 56]. It is also possible that

some of the TMD genes encode proteins with spanin-like functions that interact with the

complex cell wall of the Gram-positive bacteria. This hypothesis is supported by a report that

identified a 177 amino acid protein termed LysZ, that is encoded by phages that infect

Corynebacterium glutamicum and may function to disrupt the integrity of the bacterial cell wall

[57]. The current study aims to evaluate how two predicted holin-like TMD proteins encoded by

phages Girr and NormanBulbieJr (NBJ) function in the lysis pathway of *Mycobacterium*

smegmatis mc^2 155 by utilizing recombineered phages and multiple lysis assays under

physiological growth conditions. The findings show that both proteins termed, LysF1a and

LysF1b function as holins and are required for efficient bacterial lysis. The results also show that

the expression of the LysF1b protein is required for efficient functioning of the LysF1a protein

and that the LysF1a is not an antiholin. These results suggest a novel model for host lysis that

requires two holins with distinct membrane topologies.

MATERIALS AND METHODS

Bioinformatic analysis of transmembrane domains and AlphaFold3 analysis

The identification of transmembrane domains (TMD) and protein topologies was carried

out essentially as described by Pollenz et al., 2024 [44] utilizing Deep TMHMM (v1.0.24) [58,

TOPCONS (v2.0) [59], SignalP-5.0 and SignalP-6.0 [60], LipoP 1.0 [61] and HHpred

(databases: PDB_mmCIF70_30_Mar; Pfam-A_v37) [62]. All membrane proteins described in

this report had 100% consensus predictions of TMD domain location, number and protein

topology in all programs. Structural predictions of proteins and protein-protein interactions were

determined using AlphFold3 [63]. For single proteins, pTM scores >50% indicate high

confidence in the model predictions. For protein-protein interactions, ipTM scores <90% are

considered low confidence and should be used with caution [63]. All amino acid alignments

were performed using Clustal Omega [64].

Bacterial strains, growth and plaque assay

The list of bacterial strains, phage and plasmids utilized in this study is presented in

Table 1. *Mycobacterium smegmatis* mc^{2}155 [65] was used as wild type host for all phage

related experiments. *M. smegmatis* ΔmtpA and *M. smegmatis* ΔmtpA-comp are described in

[66]. *M. smegmatis* strains were propagated on Middlebrook 7H10 agar supplemented with 10%

Albumen Dextrose Complex (ADC; 5% albumen, 2% dextrose, 154mM NaCl) at 37^{o}C. To

generate saturated stock cultures, a single colony of *M. smegmatis* was propagated in 25ml of

Middlebrook 7H9 media supplemented with 10% ADC, 1mM CaCl$_2$, 0.5% glycerol and 0.05%

Table 1. Bacterial Strains, Phage and Plasmids

Bacterial Strains	Genotype/Features	Reference/Source
M. smegmatis	*Mycobacterium smegmatis* mc^2 155	65
M. smegmatis ΔmptA	*Mycobacterium smegmatis* mc^2 155 lacking the mptA enzyme	66
M. smegmatis ΔmptA-comp	*ΔmptA* strain carrying complementing mtpA expression plasmid	66
E. coli	*Escherichia coli* NEB5α F'I^Q	New England Biolabs
Bacteriophage	**Genotype/Features**	**Source**
Phage Girr_WT	Mycobacteriophage Girr	68. PhagesDB: https://phagesdb.org/phages/Girr
Phage Girr_*Δ29*-20	Girr_WT with deletion of gene *29*	This study
Phage Girr_*Δ29*-25	Girr_WT with deletion of gene *29*	This study
Phage Girr_*ΔlysF1a*	Girr_WT with deletion of gene *34*	This study
Phage Girr_*ΔlysF1b*	Girr_WT with deletion of gene *35*	This study
Phage NBJ_WT	Mycobacteriophage NormanBulbieJr	69. PhagesDB: https://phagesdb.org/phages/NormanBulbieJr
Phage NBJ_*Δ29*	NBJ_WT with deletion of gene *29*	This study
Phage NBJ_*ΔlysF1a*	NBJ_WT with deletion of gene *33*	69
Phage NBJ_*ΔlysF1b*	NBJ_WT with deletion of gene *34*	This study
Plasmids	**Features**	**Source**
pExTra-Girr_*29*	pExTra01::(Ptet-Girr_*34*-mcherry); Kan^R	68
pExTra-Girr_*34*	pExTra01::(Ptet-Girr_*34*-mcherry); Kan^R	68
pExTra-Girr_*35*	pExTra01::(Ptet-Girr_*35*-mcherry); Kan^R	68
pExTra-NBJ_*29*	pExTra01::(Ptet-NBJ_*29*-mcherry); Kan^R	69
pExTra-NBJ_*32*	pExTra01::(Ptet-NBJ_*32*-mcherry); Kan^R	69
pExTra-NBJ_*33*	pExTra01::(Ptet-NBJ_*33*-mcherry); Kan^R	69
pExTra-Waterfoul_*32*	pExTra01::(Ptet-Waterfoul_*32*-mcherry);	67

	KanᴿR	
pExTra-Waterfoul_32ΔN	pExTra01::(Ptet-Waterfoul_*32(truncation of N-terminal TMD)*-mcherry); Kan^R	This study
pExTra-Waterfoul_32ΔC-40	pExTra0 ::(Ptet-Waterfoul_*32(truncation of C-terminal 40aa)*-mcherry); Kan^R	This study
pJV53	Acetamide inducible plasmid with Ched9c gene *60-61*. Kan^R	70

Tween-80 with shaking (225rpm) at 37^{o}C. For all liquid lysis and one-step experiments, 100ul of

saturated stock was added to 250ml baffled flasks containing 55ml of supplemented 7H9

without Tween. Cultures were shaken (225rpm) overnight at 37^{o}C and utilized in experiments

the next day while in log phase growth (OD$_{600}$ <0.8). Titering experiments showed that an *M.*

smegmatis culture at an OD$_{600}$ of 1.0 was equal to ~6.5 x 10^7 CFU/ml. This value was utilized

when phages were infected at different MOIs. Plaque assays were completed by mixing the

appropriate amount of phage with 250ul of saturated *M. smegmatis* in Middlebrook 7H9 media

supplemented with 10% ADC, 1mM CaCl$_2$ and 0.5% glycerol. The mixture was vortexed and

incubated at 22^{o}C for 10 minutes. Following the infection, the sample was mixed with 4-5ml of

top agar (Middlebrook 7H9, 1mM CaCl$_2$, 0.5% agar) and plated immediately onto 7H10 agar

plates. Plates were allowed to harden and placed in a 37^{o}C bacterial incubator for 16-48 hrs.

Competent *M. smegmatis* cells for recombineering or for plasmid transfection and

complementation studies were prepared as described previously [28, 67-69]. *Escherichia coli*

NEB5α F′I^Q (New England Biolabs) was used for plasmid amplification and purification and

grown in LB broth or on LB agar supplemented with 50ug/ml kanamycin sulfate and.

Plasmid constructs, *M. smegmatis* transformation and complementation analysis

All PCR primers utilized in this study are shown in Table S1. To generate the inducible

expression plasmids pExTra_Waterfoul_*32ΔN*, pExTra_Waterfoul_*32ΔC40*,

pExTra_Waterfoul_32 was PCR amplified with Q5 DNA polymerase (New England Biolabs)

220 using Waterfoul specific primers containing with pExTra homology regions (Table S1). Purified

221 PCR products (were ligated into linearized pExTra01 using isothermal assembly (NEB HiFi 2x

222 Master Mix) as described [65-67]. Recombinant plasmids were recovered by transformation of

223 *E. coli* NEB5α F'I^Q and colonies selected on LB agar supplemented with 50µg/µl kanamycin

224 sulfate. Plasmid DNA was purified using the GeneJet kit (Fisher) and all plasmids were

225 sequence-verified by Sanger sequencing using the pExTra Seq primer (Azenta). For generating

226 complementing strains, competent *M. smegmatis* cells were prepared as described [67-69] and

227 40µl mixed with 100ng plasmid DNA in a 1mm electroporation cuvette. The sample was

228 electroporated with 1800V and the time constant set to 5.0 msec. Following electroporation, the

229 sample was mixed with 950µl of supplemented 7H9 media without Tween and shaken (225rpm)

230 at 37^{o}C for 120 minutes. Following the recovery period, 10µl-100µl aliquots were spread on

231 7H10 agar plates supplemented with 10µg/µl kanamycin sulfate and incubated at 37^{o}C for 3-5

232 days. To generate cultures for plaque assay, single colonies of *M. smegmatis* carrying the

233 appropriate plasmid were harvested into 25ml of supplemented 7H9 media containing 10µg/µl

234 kanamycin sulfate. Cultures were shaken (225rpm) at 37^{o}C for 24-48 hours. Saturated cultures

235 at OD_{600} > 1.5 were centrifuged at 6200rpm for 10 minutes and the pellets resuspended in

236 supplemented 7H9 media (without Tween) to a final OD_{600} of 2.0. These cells were utilized for

237 phage infection and then mixed with 4.5ml top agar supplemented with 10ng/µl kanamycin

238 sulfate with or without 100ng/µl anhydrotetracycline (aTc). The mixture was added to 7H10 agar

239 containing 10µg/µl kanamycin sulfate supplemented with or without 100ng/µl aTc and incubated

240 at 37^{o}C for the times indicated in the results section.

241 **Bacteriophage recombineering and genome sequencing**

242  All gblocks and PCR primers utilized in this study are shown in Table S1. The deletion of

243 genes in phage Girr and NBJ were carried out using the Bacteriophage Recombineering of

244 Electroporated DNA (BRED) procedure as described [28, 70], with the following modifications.

Phage genomic DNA was purified using the Wizard DNA Clean up kit (Promega). dsDNA

recombineering substrates were synthesized as gblocks (Integrated DNA Technologies). To

generate μg quantities of dsDNA substrates, gblocks were solubilized in water to 10ng/μl and

PCR amplified with gblock specific primers using 2X Q5 master mix (New England Biolabs).

PCR products were purified using Zymoclean DNA recovery (Zymoresearch). To generate

recombinant phage, 200-400ng dsDNA substrate was mixed with 100ng phage genomic DNA

and incubated on ice for 10min. The DNA mixture was added to 100μl of recombineering *M.*

smegmatis in a 2mm electroporation cuvette and electroporated at 2500V with the time constant

set to 20.0 msec. Following electroporation, 900μl of supplemented 7H9 media was added to

the cells and they were shaken (225rpm) at 37^{o}C for 90 minutes. Following the recovery period,

the sample was mixed with 250μl of saturated *M. smegmatis* and plated with 5ml 7H9 top agar

onto 7H10 agar plates. Plates were incubated at 37^{o}C for 18-24 hours to recover primary

plaques. Primary plaques were picked into 120μl of phage buffer (10 mM Tris-HCl, pH 7.5; 10

mM MgSO$_4$, 68.5 mM NaCl; 1mM CaCl$_2$), vortexed, and incubated at 22^{o}C for 2-8 hrs. The

presence of mutant phage genomes was determined by Deletion Amplification Detection Assay

(DADA) PCR [28] using a 20μl reaction with: 1μl of plaque sample, 2μl of 0.2 μM DADA primer

mix (1:1 mix of For and Rev primers), 5μl glycerol (40%), 2μl water and 10μl OneTaq 2x Master

Mix (New England Biolabs). PCR conditions were: 94^{o}C for 1 minute followed by 35 cycles of

94^{o}C-1min/66^{o}C 1min and a final 66^{o}C step for 5 minutes. Positive primary plaque samples

were serially diluted and plated on wild type or complementing *M. smegmatis* as indicated.

Secondary plaques were picked as above and screened by standard PCR using gene specific

flanking primers. Pure populations of mutant phage were evaluated by full genome sequencing

as described [71, 72]. Briefly, Genomic DNA was used to create sequencing libraries with the

NEB XLEP-P1 kit. Sequencing was performed by the Pittsburgh Bacteriophage Institute and the

library run on an Illumina NextSeq 1000 instrument. Raw reads were trimmed with cutadapt 4.7

(using the option: –nextseq-trim 30) and filtered with skewer 0.2.2 (using the options: -q 20 -Q

30 -n -l 50) prior to assembly. The resulting sequences were assembled using Consed (v29.0)

with Unicycler (v5.0) and contigs checked for completeness, accuracy, and genome termini.

Final phage contigs were aligned to wild type phage to identify deletions and point mutations.

Plaque size analysis

Plaques were generated using the plaque assay procedure described above. For

experiments where plaque size would be evaluated between several different phages, identical

bacterial samples, incubation times, phage concentrations and top agar amounts were utilized.

Plaque size was evaluated using ImageJ software [73] and by manual measurement using

photos embedded in PowerPoint and custom rulers. In most experiments, results are presented

for an average of 50 plaques and the statistical significance of measurements determined by

unpaired t-test (https://www.graphpad.com/quickcalcs/ttest1/). Plaque volume was calculated by

the formula $\pi r^2 h$ where r is the radius of the plaque and h was set to 1mm for the top agar

thickness.

Liquid lysis assay

Liquid lysis assays were carried out essentially as described by Payne et al., 2009 [27]

with the following modifications. Briefly, overnight cultures of *M. smegmatis* in log phase growth

were diluted to an OD_{600} of 0.25-0.30 in supplemented 7H9 without Tween and 40ml of culture

was added to 250ml baffled flasks. High titer phage lysate was added to the cultures to create

an MOI of 10 and the mixture set at 37^oC for 30 minutes without shaking. T_0 for all experiments

was the time of phage addition. After 30 minutes (T_{30}), the cultures were constantly shaken a

225rpm and aliquots of cells removed with a syringe and sterile needle from the shaking culture

without stopping the incubator. The OD_{600} of the sample was immediately recorded for each

aliquot using a GENESYS spectrophotometer (Fisher Scientific). In some experiments, KCN

(1M in water), or $CHCl_3$ (100%) was added directly to the shaking culture to produce a final

concentration of 10mM or 1%, respectively. All liquid lysis experiments were repeated at least

five separate times and representative experiments are presented.

ATP assay

The presence of ATP in culture supernatants was determined utilizing the ENLIGHTEN

kit (Promega). Briefly, liquid lysis assays were carried out as described above. A 500ul aliquot of

culture was removed at indicated time points and immediately passed through a 0.2μ syringe

filter to remove bacteria. The sample was frozen at -20^{o}C prior to analysis. The presence of ATP

in culture supernatants was determined by mixing 5μl with 50μl ENLIGHTEN reagent

(Promega), mixing for 5 seconds and then reading the integrated luminescence over 10

seconds using a GloMax luminometer (Promega). To assure that the readings were within the

linear range of the instrument an ATP standard curve was utilized.

One-step growth curves and burst size determination

Several experimental procedures have been utilized for one-step growth curves with

Gram-positive bacteria [32, 47, 49, 51]. In our hands, none of these procedures generated

consistent results or provided the ability to directly measure burst size. Thus, we developed a

one-step protocol that is illustrated in Figure S1 and described below. To prepare the cells, the

OD$_{600}$ of fresh overnight cultures of *M. smegmatis* in log growth were determined and 10ml

centrifuged at 3,000rpm for 10 minutes. The cells were resuspended in supplemented 7H9 to a

final concentration equal to an OD$_{600}$ of 3.2/ml. 250μl of cells (~5 x 10^{7} cfu) were added to

microfuge tubes and ~5 x 10^{7} phage added to the cells to create an MOI of 1. The same amount

of phage were also added to 250μl of supplemented 7H9 media without cells to verify the titer

and determine the level of phage adsorption. The mixture was vortexed and set at 37^{o}C for 30

minutes without shaking. T$_{0}$ for all experiments was the time of phage addition to the cells. After

30 minutes, 750μl of supplemented 7H9 was added to all samples. Samples were vortexed and

centrifuged at 6,200rpm for 4 minutes to separate unabsorbed phage. The supernatant was

saved for analysis of titer and adsorption, and the cell pellet immediately resuspended in 1ml

supplemented 7H9, vortexed and diluted 1:10,000 into 70ml of supplemented 7H9 in baffled

250ml flasks. An aliquot was removed and 5μl, 12.5μl and 25μl used for plaque assay to

determine the number of infectious centers to determine the precise number of infected cells. A

second aliquot was removed and filtered through a 0.2µ syringe filter and 150µl utilized for

plaque assay to test for unabsorbed (free) phage at T_{45} minutes (the first one-step data point).

The flask was placed in a $37^{o}C$ incubator and constantly shaken at 225rpm for the duration of

the experiment. At specified time points, 500µl aliquots were removed from the shaking flasks

and each sample immediately passed through a 0.2µ syringe filter to remove cells and collect

free phage. The amount of phage in each sample was determined by performing plaque assays

using 150µl of serially diluted samples (usually 10^{0} to 10^{-4} dilutions were sufficient). The phage

titer in the flask, expressed as PFU/ml was determined at each time point by counting the

plaques on each plate, dividing by the volume of phage used in the infection (150µl), multiplying

by the dilution factor and then multiplying by 10^{3}. The number of infectious centers in the

infected 70ml culture was determined by counting the plaques, dividing by the volume used for

the plaque assay and multiplying by 70,000. The original phage titer and the percentage of

adsorbed phage was determined by spot titers and plaque assay of the serially diluted

supernatants collected from the samples of phage with and without the added cells. Since the

number of cells and phage used in these experiments was held constant, this procedure

consistently produced ~3.5-5.5 x 10^{7} infectious centers. This procedure also resulted in

essentially zero unabsorbed phage in the culture at the start of the one-step assay. Burst size at

all time points could be directly determined by dividing the total phage in the 70ml culture by the

number of infectious centers. All phages were evaluated through a minimum of three separate

assays. A representative data set showing the raw data for adsorption, infections centers and a

one-step assay is provided in Figures S2 and S3.

RESULTS AND DISCUSSION

Topologies of Holin-like TMD Proteins Within the Lysis Cassettes of F1 Cluster Phages Girr and NormaBulbieJr.

It was first important to bioinformatically evaluate the predicted holin-like proteins in the lysis of *M. smegmatis*, it was desirable to begin work in phages that had a defined lysis cassette and prior wet-lab data. F1 cluster phages Girr and NormanBulbieJr (NBJ) were selected since both phages have a define lysis cassette and a full genome cytotoxicity screen has been reported for both phages [68, 69]. Girr and NBJ have an average nucleotide identity of 89.86% but only share 69 of 102 genes. Both phages have a defined lysis cassette organization located downstream of the *tape measure* and *minor tail* genes and encode a Lysin A enzyme predicted to target the peptidoglycan layer and a Lysin B that targets the linkage of the arabinogalactins to the mycolic acids. The *lysin B* gene is directly followed by two genes encoding proteins that have predicted TMDs (Girr *34* and *35* and NBJ *32* and *33*).

It was first important to bioinformatically evaluate the predicted topologies of these TMD proteins and determine how they compared to previously characterized holins or other phage proteins involved in bacterial lysis. Girr *34* and NBJ *32* encode small 80aa and 77aa proteins, respectively, that have two predicted TMD regions and an N-in-C-in topology by all TMD prediction programs (Figure 2A). The proteins are 97% identical and have cytoplasmic N and C-terminal domains that each contain four charged amino acids. The presence of two TMDs and the N-in-C-in topology is like the phage 21 S[21] type II pinholin (Figure 2BC). However, the presence of a pinholin is typically associated with a SAR-endolysin that is secreted to the periplasmic space prior to lysis triggering [10, 11, 13]. Analysis of the Girr or NBJ Lysin A proteins by the various TMD prediction programs and AlpahFold3, show no evidence of SAR domains or signal sequences (Figures S4-S6), and this is consistent with previous analysis of Lysin A proteins from phages that infect *M. smegmatis* [20]. Thus, it is critical to contrast several structural differences of gp32 and gp34 to the canonical S[21]68 pinholin and S[21]71 antiholin.

First, the $S^{21}68$ holin protein has only 3 N-terminal amino acids prior to the TMD1 and

this feature is a critical to its function as a holin and allows the TMD1 to become cytoplasmic

(Figure 2BC) [11-15]. Second, the S^{21} gene has a dual start motif that produces both the $S^{21}68$

holin protein and the $S^{21}71$ antiholin after phage infection [11-15]. The difference in the two

proteins is the addition of the extra basic K residue to the N-terminal cytoplasmic domain of

$S^{21}71$ that limits the TMD1 from exiting the membrane to the same degree as the $S^{21}68$ holin

(Figure 2BC; [11-15]). Thus, the expression of $S^{21}71$ keeps a percentage of the $S^{21}68$ in an

inactive dimer until triggering at which time the $S^{21}71$ converts to an active holin [13-15, 74, 75].

Third, the $S^{21}68_{IRS}$ protein is an engineered version of $S^{21}68$ that adds the MRYIRS motif and 4

charged residues in the N-terminal cytoplasmic domain that creates a dominant negative

version of the antiholin that is poorly triggered to an active holin [74]. Neither Girr gene *34* nor

NBJ gene *32* have a dual start motif that generates multiple products of the same open reading

frame. Genes *34* and *32* encode a single protein product that has a predicted N-terminal

cytoplasmic domain of 13-16 amino acids of which four are charged and two others are polar.

Therefore, the N-terminal topology and the number of charged amino acids in Girr gp34 and

NBJ gp32 are more similar with the $S^{21}71$ antiholin and $S^{21}68_{IRS}$ and would predict that TMD1 of

these proteins remains embedded in the membrane. In addition, the TMDs of S^{21} both contain

GxxxG motifs that are predicted to contribute to homotypic and/or heterotypic helix interactions

of the monomers that facilitate the formation of the pinholes [74, 75]. While TMD1 and TMD2 of

Girr and NBJ have a high percentage of GA residues (Figure 2A), they lack defined GxxxG

domains found in S^{21}. Finally, Girr gp34 can't be modeled with high confidence by AlphaFold3

into either a monomer or a heptamer (Figure S7). Taken together, this data suggests that Girr

gp34 or NBJ gp32 are unlikely to function as a type II holin. Based on the literature we will

designate these proteins as LysF1a due to a predicted function in lysis (lys) and being first

identified in F1 cluster phages.

Since there is a second gene encoding a protein with a single TMD directly downstream

of the *lysF1a* gene, we hypothesized that this protein may also be involved in the lysis pathway.

Girr *35* and NBJ *33* both have -4bp ATGA overlaps to the upstream *lysF1a* gene and encode

98% identical 124aa proteins with a single N-terminal TMD and a predicted N-out-C-in topology

by all TMD prediction programs (Figure 3A). Both proteins have a 22 amino acid extracellular

domain followed by the single TMD. There are four basic residues following the TMD and

AlphaFold3 predicts an extended helical cytoplasmic C-terminal domain that contains ~50%

KRDE charged residues typical of holins [10, 11]. Figure 3B shows the AlphaFold3 models and

predicted topologies of representative proteins that have a single TMD and have been shown to

be involved in the bacterial lysis pathway for comparison gp33 and gp35. Most pertinent is the

type III holin typified by the T protein of phage T4 that forms large 1um holes like canonical type

I holins [16, 17]. However, the type III holin has <20% amino acid identity to gp33 and gp35, is

much larger at 221 amino acids and has the opposite membrane topology with a globular

extracellular domain critical to its holin function (Figure 3B). It noteworthy that spanins also have

a single TMD and model with an extended helical prediction, but like the type III holins, these

proteins are also oriented in a N-in-C-out topology that has been shown experimentally and is

also predicted by the TMD prediction programs (Figure 3B; [10, 11]). Interestingly, there is a

recent report regarding phages that infect the Gram-positive *Corynebacterium glutamicum*

strain. The authors have characterized a protein with 1 TMD that is termed LysZ and is

proposed to interact with lipid-anchored glycans on the extracellular side of the IM to promote

efficient lysis [57]. However, like the type III holin, LysZ is much larger than the Girr and NBJ

proteins and it is proposed to have a N-in-C-out topology that allows it to interact with the lipid-

anchored glycans [57]. Thus, Girr gp35 and NBJ gp33 are most similar in structure and

membrane topology to the recently identified phage Mu releasin protein (Figure 3B) [18].

Releasin however, is named due to its function to release the Mu SAR-endolysin from the

membrane in a holin independent manner [18]. Mu also has <20% amino acid identity to gp33

and gp35 and does not possess the highly charged cytoplasmic domain. Thus, of all the

characterized proteins involved in lysis that have 1 TMD, gp35 and gp33 appear to represent a

novel class of proteins and will be termed LysF1b since the gene is downstream of the *lysF1a*.

Deletion of Individual *lysF1a* and *lysF1b* Genes in Phage Girr are Viable While Deletion of

Both is Lethal

To evaluate the function of the *lysF1a* and *lysF1b* genes, the BRED recombineering

method was utilized to delete them in both Girr and NBJ as described in Methods. Several

primary plaques were positive for each deletion and pure populations of Girr_*ΔlysF1a*,

Girr_*ΔlysF1b*, NBJ_*ΔlysF1a*, and NBJ_*ΔlysF1b* were recovered from secondary plaques. Figure

4 shows the lysis cassette organizations of the Girr mutants and the PCR verification of the

gene deletions from clonal high titer lysates. The data for NBJ is shown in Figure S8 and

deletion of gene *33* is also presented in Wise et al. [69]. The genome structure of all deletions

was confirmed by sequencing the immediate region of the deletion and by whole genome

sequencing to validate that there were no point mutations outside of the targeted gene deletion

region. Since deletion of either of the individual TMD genes resulted in viable phage, it was of

interest to test whether a double deletion of both TMD genes was viable. For these studies

Girr_*ΔlysF1a* DNA was utilized to delete *lysF1b* and a positive primary plaque was picked,

resuspended in buffer and plated to recover ~250-1000 isolated plaques on complementing and

non-complementing *M. smegmatis* as shown in the protocol schematic in Figure 5A. Since

BRED was being used, positive primary plaques will have a mixture of the phage used for

recombineering and the mutant. Lysates representing all the phage from secondary plaques

were recovered and evaluated by DADA PCR to test for the presence of the double mutant.

Figure 5B shows that the double mutant genotype is not recovered on wild type *M. smegmatis*,

even when 800PFU were plated (gel lanes 3-6). However, when >250 phage particles are

plated on the complementing *M. smegmatis* strain transfected with pExTra-Girr*34* expressing

LysF1a, a positive DADA band is observed (Figure 5B, gel lane 8). The inability to recover a

phage with the deletion of both *lysF1a* and *lysF1b* is highly supportive that the expression of

both Girr LysF1a and LysF1b are required for Girr viability. A lethal phenotype is observed when

holin genes are deleted in *E coli.* phages that encode canonical holins required for endolysin

release [10, 11].

Deletion of Individual *lysF1a* and *lysF1b* Genes in F1 Phages Result in Reduced Plaque

Size

456   To assess the phenotypes of the mutant phage, equal concentrations of Girr_WT,

Girr_*ΔlysF1a* and Girr_*ΔlysF1b* were used to infect *M. smegmatis* and the total number of

recovered plaques and plaque sizes evaluated. All phages formed plaques with equivalent

efficiencies under the standard assay conditions described in Methods suggesting no defects in

adsorption (Figure S9). However, both Girr_*ΔlysF1a* and Girr_*ΔlysF1b* presented with

significantly smaller plaques than Girr_WT (Figure 6) as did NBJ_*ΔlysF1a* and NBJr_*ΔlysF1a*

(Figure S10). A summary of the plaque size data for both Girr and NBJ is presented in Table 2.

Deletion of the *lysF1a* gene from Girr or NBJ results in ~40% reduction in plaque diameter and

Table 2 Plaque sizes in WT Girr and *lysF1a* and *lysF1b* deletion mutants.

PHAGE	Plaque size (mm)	% of WT (diam.)	Plaque (vol, mm^3)	% of WT (plaque vol)
NBJ_WT	3.49 +/- 0.533	100	11.479	100
NBJ_*ΔlysF1b*	1.23 +/- 0.225*	35.24*	1.425*	12.42*
NBJ_*ΔlysF1a*	2.05 +/- 0.660*	58.73*	3.96*	34.50*
Girr_WT	3.88 +/- 0.731	100	14.188	100
Girr_*ΔlysF1b*	1.28 +/- 0.376*	32.98*	1.544*	10.88*
Girr_*ΔlysF1a*	2.31 +/- 0.719*	59.53*	5.029*	35.44*

466   *M. smegmatis* was infected with identical numbers of phage and plaque size determined after
467   36 hrs. of growth at 37^{o}C. Plaque diameter was measured using ImageJ. Data is presented as
468   average +/- standard deviation for a minimum of 50 individual plaques. * = statistically significant
469   from Girr or NBJ WT phage *p<0.0001.*
470

a 65% reduction in plaque volume, while deletion of the *lysF1b* is more severe showing a ~65%

reduction in plaque diameter and near 90% reduction in plaque volume. To validate that these

plaque size reductions were directly related to the loss of *lysF1a* or *lysF1b* genes, *M. smegmatis*

containing complementing expression plasmids were infected with the mutant phages and

plaque size evaluated as described in Methods. Since both Girr LysF1b and NBJ LysF1b are

highly toxic when exogenously expressed in *M. smegmatis* [68, 69], complementation of the

lysF1b deletions first utilized Waterfoul gp32. Waterfoul is a K5 cluster phage with a canonical

lysis cassette and gene *32* encodes a 1TMD with the same general size and membrane

topology as Girr LysF1b (Figure 3C) and is not toxic when induced by aTc in *M. smegmatis* [67].

When M*. smegmatis* transfected with Waterfoul gp32 is infected with Girr_*ΔlysF1b* in the

presence of aTc, there is a significant increase in plaque size compared to untreated cells

(Figure 6A). Importantly, the increase in plaque size was not observed when the N-terminal

TMD domain or the highly charged C-terminal 40 amino acids of Waterfoul gp32 were truncated

(Figure 6A). Thus, the complementation by Waterfoul gp32 requires the full-length protein.

Recently it has been reported that the pTet promoter in pExtra is leaky and that it is possible to

complete complementation analysis in the absence of aTc [68]. Thus, *M. smegmatis* was

transfected with pExTra-Girr*35* and pExTra-NBJ*33* and the resulting cells infected with

Girr_Δ*lys1b* as described in Methods. Figure 6B shows that there is a significant increase in

plaque size when Girr_Δ*lys1b* is plated on the complementing strains expressing the LysF1b

proteins compared to cells with the base pExTra01 vector. These results support that the

reduced plaque size of Girr_*ΔlysF1b* is directly attributable to the loss of *lysF1b.* Similarly, the

wild type plaque size is restored when *M. smegmatis* harboring the pExTra-Girr_*34* plasmid is

infected with Girr_*ΔlysF1a* (Figure 8). Thus, the loss of either *lysF1* gene appears to create a

lysis defect phenotype that is like the phenotype when the *lysin B* is deleted in phage Giles and

the cell wall components remain a barrier to efficient phage release [27].

Deletion of Individual *lysF1* Genes Result in Distinct *M. smegmatis* Liquid Lysis

Phenotypes and ATP Release.

A main function of holins is the control of the lysis timing by the destruction of the proton

motive force (PMF) [10, 11]. To assess lysis timing of Girr, NBJ and the *lysF1* mutants, liquid

lysis assays were performed where the optical density (OD_{600}) of infected log growth *M.*

smegmatis cultures is measured over time. Importantly, in all studies, once phage had been

allowed to absorb for 30 minutes, cultures were continually shaken even while samples were

withdrawn so that lysis would not be prematurely triggered by changing the culture conditions

[11, 76]. The data show that cultures infected with Girr_WT grow steadily until ~110 minutes

post infection. Beginning at 110 minutes, there is a significant drop in the OD_{600} of the culture

(triggering event) until it reaches a baseline of OD_{600} 0.10 at about 200 minutes (Figure 9A). In

contrast, Girr_*ΔlysF1a* grows until ~130 minutes and then shows a reduction in OD_{600} until 300

minutes that does not reach the same OD_{600} baseline as Girr_WT even after 480 minutes of

incubation (Figure 9A). Strikingly, cultures infected with Girr_*ΔlysF1b* do not show a drop in

OD_{600} but appear to cease growing after ~200 minutes with only a slight decrease in OD_{600} over

the remaining growth period out to 480 minutes (Figure 9A). For NBJ_WT, OD_{600} begins to drop

after 130 minutes and like Girr_WT reaches a baseline of around OD_{600} 0.10 (Figure 9B).

NBJ_*ΔlysF1a* shows a delay in lysis timing like Girr_*ΔlysF1a*, but the delay is more pronounced

as the OD_{600} does not begin to drop until ~180 minutes post infection (Figure 9B). Just like

Girr_*ΔlysF1b*, NBJ_*ΔlysF1b* ceases to grow after ~200 minutes and does not show a significant

reduction in OD_{600} even after 480 minutes of growth (Figure 9B). Thus, in both phages, deletion

of *lysF1a* results in a triggering delay while deletion of *lysF1b* results in no defined triggering

event in this assay. It is interesting that Girr_WT and NBJ_WT show a ~20-minute difference in

triggering time and that a difference is also observed between Girr_*ΔlysF1a* and NBJ_*ΔlysF1a.*

These results may be due to differences in how the lysis cassette genes are expressed since

Girr has an HNH endonuclease gene between the *lysin A* and *lysin B* (Figure 1) and the

intergenic region between the *lysin B* gene and the *lysF1a* is 8bp in NBJ but 38bp in Girr (Figure

S11).

ATP release has been used as a measure of cell viability and membrane integrity

following phage infection [27, 51]. Figure 9C shows the ATP release from the same experiment

that is presented in Figure 9AB. As expected, ATP release for Girr_WT, NBJ_WT, Girr_*ΔlysF1a*

and NBJ_*ΔlysF1a* generally mirrors the time at which lysis is triggered and the culture OD_{600}

begin to decline. ATP levels in the media also peak when the OD_{600} baseline is reached at about

300 minutes (Figure 9C). In contrast, there is delayed but gradual increase in the amount of

ATP released from the cultures infected with Girr_*ΔlysF1b* and NBJ_*ΔlysF1b* out to 300

minutes. Although the ATP level does not reach that of Girr_WT or NBJ_WT, the release of ATP

confirms that cell integrity has been comprised after infection with the *lysF1b* mutants.

Collectively, the important finding from this data set is that an F1 phage with only the 1TMD

LysF1b protein can trigger the lysis event in the absence of the 2TMD lysF1a protein, but that

the lysF1a protein does not support significant bacterial lysis as measured by a drop in OD_{600}. In

fact, the liquid lysis results for Girr_*ΔlysF1b* and NBJ_*ΔlysF1b* are like those observed for the

lethal *lysin A* deletion in phage Giles where the cultures stopped growing and the OD_{600}

plateaued but did not decline [27]. The similarity in these results suggest that the lysis defects in

the *ΔlysF1b* phage may be related to the inability of the endolysins to efficiently cleave the cell

wall components because they are either not efficiently released from the infected cell or cannot

be activated in some manner.

Energy Poisons Cause Early Triggering of F 1Cluster Phages with Deletion of the *lysF1a*

Gene but not the *lysF1b* Gene in Liquid Lysis Assays

The results presented so far indicate that the expression of both TMD proteins are

required for phage viability, but phages that express only the LysF1b protein can support

bacterial lysis while phages that express only LysF1a are viable but severely lysis

compromised. The diagnostic assay for holin function is the ability to cause early triggering of an

infected culture by the addition of energy poisons (e.g., cyanide or the protonophore

dinitrophenol) due to the disruption of the proton motive force (PMF) [11, 76]. This type of assay

has been utilized in nearly all the pioneering work on holin function of phages that infect *E. coli.*

[10, 11, 18]. *M. smegmatis* was infected with Girr_WT or Girr_*ΔlysF1a* and the OD_{600} of the

culture measured as described in Methods. At 110 minutes after infection, KCN (final

concentration of 10mM) was added directly to one culture of Girr_*ΔlysF1a* and to a culture of

uninfected cells. Figure 10A shows that Girr_WT triggers at ~110 minutes while Girr_*ΔlysF1a*

triggers at 130 minutes just as in Figure 9A. However, when KCN is added to a culture infected

with Girr_*ΔlysF1a*, there is a drop in OD_{600} beginning at ~120 minutes that clearly precedes the

drop in the untreated Girr_*ΔlysF1a* culture. KCN addition to uninfected cells results in immediate

cessation of growth, but not a significant drop in OD_{600}. To validate these results, studies were

repeated with the NBJ mutants. NBJ_WT triggers later than Girr_WT at ~130 minutes and

declines to OD_{600} 0.10 by 300 minutes as expected (Figure 10B). NBJ_*ΔlysF1a* shows delayed

triggering just as in Figure 9B and addition of KCN to a culture infected with NBJ_*ΔlysF1a* at

120 minutes results in lysis triggering by 130 minutes with a gradual reduction in OD_{600} that

mirrors the NBJ_WT. The ability of Girr_*ΔlysF1a* and NBJ_*ΔlysF1a* to trigger lysis, identifies the

1TMD LysF1b protein as a putative holin and not a releasin, since energy poisons do not trigger

lysis prematurely in phage Mu [18].

The next set of studies focused on the phages with deletion of the 1TMD *lysF1b* genes

to determine if they could also be triggered by the addition of KCN. Figure 10C shows the

results for Girr_*ΔlysF1b*. As previously shown in Figure 9A, *M. smegmatis* infected with

Girr_*ΔlysF1b* does not exhibit a defined triggering event as the cells simply cease growing with

a slight reduction in OD_{600} out to 480 minutes. Addition of KCN to the Girr_*ΔlysF1b* culture at

180 minutes results in a modest 30% reduction of OD_{600} between 180-240 minutes compared to

the Girr_$\Delta lysF1b$ untreated culture. However, following this initial decline, the level of change in

OD_{600} generally mirrors that of the untreated Girr_$\Delta lysF1b$ culture and the KCN treated *M.*

smegmatis control. In contrast, addition of 1% $CHCl_3$ to the Girr_$\Delta lysF1b$ culture at 180 minutes,

results in an immediate precipitous drop in OD_{600} to the level of Girr_WT indicating that the

endolysin enzymes are present and active in the cells. The results for NBL_$\Delta lysF1b$ are shown

in Figure 10D and exhibit near identical results to Girr_$\Delta lysF1b$ in that addition of KCN cause a

modest drop during the first hour after addition, and then an OD_{600} decline that mirrors the

untreated NBJ_$\Delta lysF1b$ culture and the KCN treated *M smegmatis*. Addition of $CHCl_3$ results in

the same type of sharp decline in OD_{600} to the level of the NBJ_WT baseline. Collectively, these

results show that phages expressing only the 2TMD LysF1a protein are not efficiently triggered

following disruption of the PMF but that the cells have the requisite enzymes for lysis since lysis

occurs immediately after addition of 1% $CHCl_3$. The inability of the LysF1a proteins to be

triggered by KCN is like the $S^{21}71$ or $S^{21}68_{IRS}$ antiholins [13, 74, 75].

F1 Cluster Phage Deleted for *lysF1a* have Delayed Burst Timing While Phage Deleted for

***lysF1b* Deletion have Reduced Timing and Greatly Reduced Burst Size**

The previous results show that infection of *M. smegmatis* with either Girr_$\Delta lysF1a$ or

NBJ_$\Delta lysF1a$ results in a defined triggering (lysis) event that can be prematurely initiated by

addition of KCN. However, the results do not establish if phage have been efficiently released

from the lysed cells, the timing of the burst event and the burst size. Since it is not possible to

utilize lysogens to study phage release in *M. smegmatis*, one-step grow curves were utilized to

evaluate the various *lysF1a* and *lysF1b* mutant phages. The experimental set up for these

studies is fully described in the methods and representative raw data sets are also shown in

Figures S2 and S3. The one-step results from three different experiments utilizing Girr phages

are presented in Figure 11. Phage release from Girr_WT shows a lag time of ~180 minutes from

the time of infection and then a gradual phage release for 90 minutes before reaching a plateau

at ~300 minutes (Figure 11A). The slow release of phage over a 90-minute period is consistent

with the liquid lysis assays that showed an ~60-minute timeline of OD_{600} decline after triggering.

This is an important observation because there should not be a sharp, immediate lysis of all

cells and accompanying phage release since the phage infections are not synchronized but

carried out over a 30-minute period and the efficient release of phage from Gram-positive

bacteria may also be hampered by the extensive cell wall. Girr_*ΔlysF1a* shows a delayed lag

time by ~20 minutes compared to Girr_WT and a more prolonged period of phage release over

150 minutes before reaching a plateau. The overall burst size for Girr_WT over three

experiments was 277 +/- 81 PFU/cell. Girr_*ΔlysF1a* showed a lower burst of 255 +/- 75 PFU/cell

but this was not statistically different from the for Girr_WT. In contrast, Girr_*ΔlysF1b* shows a

very small but defined burst at ~ 210 minutes that is best illustrated when the data is plotted

independent of the other phages (Figure 11B). The overall burst size for Girr_*ΔlysF1b* is only

~29.2 +/- 36 PFU/cell suggesting very poor efficiency of phage release from the infected cells.

Overall, this data is consistent with the plaque size and liquid lysis results and supports the

hypothesis that *M. smegmatis* integrity is compromised when infected with Girr_*ΔlysF1b* but the

release of phage progeny is severely impacted. In addition, the near wild type burst size

observed in cells infected with Girr_*ΔlysF1a*, further support the holin function of the LysF1b

protein.

Lysis Recovery Mutants Isolated from Girr_*ΔlysF1b* or NBJ_*ΔlysF1b* Show Wild Type

Plaque Size and have Point Mutations in the *lysF1a* Gene

Since both Girr_*ΔlysF1b* and NBJ_*ΔlysF1b* show significant plaque size defects it was

possible to screen for lysis recovery mutants (LRM) by assessing changes to the small plaque

size following multiple rounds of infection and replication. Figure 12A shows the presence of a

large plaque in a Girr_*ΔlysF1b* plaque assay after the second round of harvest and infection.

Table 3. Lysis Recovery Mutants (LRM) from Δ*lysF1b* Phages.

Phage	Type of mutation	Genetic location	Mutation Result
Girr_Δ*lysF1b*-LRM1	point mutation-nucleotide substitution	gene *35 (lysF1a)*	G66D
Girr_Δ*lysF1b*-LRM2	point mutation-nucleotide substitution	gene *35 (lysF1a)*	R19H
Girr_Δ*lysF1b*-LRM6	point mutation-nucleotide deletion	gene *35 (lysF1a)*	missense coding after A63 and premature stop at aa74: A63APRQPRNGFGD.STOP
Girr_Δ*lysF1b*-LRM7	point mutation-nucleotide substitution	gene *35 (lysF1a)*	G66D
Girr_Δ*lysF1b*-LRM8	point mutation-nucleotide insertion	gene *35 (lysF1a)*	stop after G66
Girr_Δ*lysF1b*-LRM11	point mutation-nucleotide insertion	gene *35 (lysF1a)*	stop after G66
Girr_Δ*lysF1b*-LRM12	point mutation-nucleotide insertion	gene *35 (lysF1a)*	stop after G66
Girr_Δ*lysF1b*-LRM14	point mutation-nucleotide substitution	gene *35 (lysF1a)*	G66D
Girr_Δ*lysF1b*-LRM15	point mutation-nucleotide insertion	gene *35 (lysF1a)*	stop after G66
Girr_Δ*lysF1b*-LRM16	point mutation-nucleotide substitution	gene *35 (lysF1a)*	A31T
NBJ_Δ*lysF1b*-LRM1	point mutation-nucleotide substitution	gene *33* (*lysF1a*)	A68E
NBJ_Δ*lysF1b*-LRM2	point mutation-nucleotide substitution	gene *33* (*lysF1a*)	A68E
NBJ_Δ*lysF1b*-LRM3	point mutation-nucleotide substitution	gene *33* (*lysF1a*)	A68E
NBJ_Δ*lysF1b*-LRM7	point mutation-nucleotide substitution	gene *33* (*lysF1a*)	D9A

When phages were recovered from the large plaque, serial diluted and replated, the wild type

plaque size was recovered across the full phage population (Figure 12B). This type of

experiment was repeated multiple times with Girr_*ΔlysF1b* and NBJ_*ΔlysF1b* and ten

independent Girr_*ΔlysF1b*-LRMs and four independent NBJ_*ΔlysF1b*-LRMs were recovered. All

Girr LRM mutants showed a large plaque phenotype compared to the parent Girr_*ΔlysF1b*

(Figure S11 and RSP, data not shown) and PCR analysis of the *lysF1a-lysF1b* genomic region

confirmed that all the LRM mutants contained the expected Δ*lysF1b* deletion (Figure S12). To

determine the location of any nucleotide mutations within the genome of the LRM mutants, DNA

was extracted and the genomes were sequenced as detailed in Methods. In all cases, the

Girr_Δ*lysF1b*-LRMs and NBJ_Δ*lysF1b*-LRMs contained the expected *lysF1b* deletion but all

phage also contained either nucleotide insertions, deletions or substitutions within the *lysF1a*

open reading frame that altered the amino acid sequence of the LysF1a protein. A summary of

all LRM mutants is provided in Table 3 and a schematic showing the amino acid changes to the

different Girr_lysF1a-LRMs is presented in Figure S13.

Six different mutations were identified in the ten Girr_Δ*lysF1b*-LRMs and two in the four

NBJ_Δ*lysF1b*-LRMs.The mutations mapped to three different domains of the LysF1a protein:

the N-terminal cytoplasmic domain, the first TMD or the beginning of the cytoplasmic C-terminal

domain. No mutants were recovered that impacted the second TMD or distal regions of the

cytoplasmic C-terminal domain. In the Girr LRM mutants, two different missense mutations were

found in TMD1 of LysF1a that result in R19H and A31T while one of the NBL LRM mutants

changed the charged cytoplasmic D9 to A. In addition, three G66D mutants were recovered in

the Girr LRMs, and three A68E mutants were recovered in the NBJ LRMs. All of these change a

small hydrophobic amino acid to a negatively charged amino acid just after the end of the TMD2

at the beginning of the cytoplasmic C-terminal domain. Finally, four mutants generate a

nonsense mutation that either truncates the Girr LysF1a protein after G66 or create a frame shift

after A63 that results in missense coding and a premature stop after amino acid 73. All LRM

mutations result in changes to the LyF1a protein that likely alter its conformation, thus, the

findings validate that the LysF1a protein can efficiently function in the lysis pathway and that the

protein is not exclusively an antiholin.

Lysis Recovery Mutants Exhibit Early Lysis and Reduced Burst Size

654   Since the LRMs generate a wild type plaque size (Figure 12A, Figure S12), it was

pertinent to evaluate the lysis timing in liquid culture and determine lag time and burst size. *M.*

smegmatis was infected with Girr_WT, Girr_*ΔlysF1b* and three different Girr_*ΔlysF1b*-LRMs and

the OD_{600} of the culture measured as described in Methods. Figure 13A shows that Girr_WT

triggers lysis as expected at ~110-minutes while Girr_*ΔlysF1b* causes reduced culture growth

but no defined triggering as also shown in Figures 9 and 10. In stark contrast, Girr_*ΔlysF1b*-

LRM1*,* Girr_*ΔlysF1b*-LRM2 and Girr_*ΔlysF1b*-LRM16, all trigger lysis prematurely at ~90

minutes and ultimately reduce the OD_{600} to the same baseline as Girr_WT. When NBJ_WT,

NBJ_*ΔlysF1b*, NBJ_*ΔlysF1b*-LRM1 and NBJ_*ΔlysF1b*-LRM7 were tested in the liquid lysis

assay, the NBJ-LRMs also trigger lysis prematurely and reduce the OD_{600} to the same level as

NBJ_WT (Figure 13B). To validate the early triggering time and assess its impact on burst size,

one-step growth curves were completed as described in Methods. Figure 13C shows a

representative one-step experiment with Girr_WT and two LRM mutants. Phage release from

Girr_*ΔlysF1b*-LRM1 and Girr_Girr_*ΔlysF1b*-LRM2 is detected by 150 minutes and plateaus by

200-minutes, a time point at which Girr_WT has achieved <10% burst size. The two Girr LMRs

average burst size is also only 95.5 +/- 12.5 PFU/cell compared to 289.6 +/- 65 PFU/cell for the

Girr_WT. The reduced burst size is likely the result of lysis triggering well before all phage

progeny have been full assembled and is consistent with reports that have evaluated early

triggering in *E. coli* phages [11, 12, 74]. The current results also indicate that plaque size in our

experimental system is not a function of burst size since a >65% reduction in burst size is still

able to generate wild type-sized plaques (Figure 12A,Figure S12). These findings add further

support to the hypothesis that the small plaque phenotypes of the various LysF1 mutants is the

result of inefficient cell wall lysis that hampers phage release.

Analysis of *lysF1b* Phage Plaque Sizes in *M. smegmatis* Lacking the Lipoarabinomannan

Layer of the Cell Wall

A recent study proposes a 1TMD protein termed LysZ may function as a spanin-like protein to disrupt the integrity of the Gram-positive *Corynebacterium glutamicum* cell wall by interacting with the lipoarabinomannan (LAM) layer [57]. In these studies, deletion of *lysZ* from phages CL31 or Cog results in a small plaque phenotype that is rescued if phage were plated on a *Corynebacteriumg glutamicum* strain that is defective in the synthesis the LAM. Since both LysZ and LysF1b have a single TMD and deletion of the genes results in small plaques, it was pertinent to determine whether an *M. smegmatis* strain lacking the mtpA enzyme required to synthesize LAM (*M. smegmatis* ΔmtpA) [66], would complement the small plaque phenotype of Girr_Δ*lysF1b*. Wild type *M. smegmatis*, *M. smegmatis* ΔmtpA, and *M. smegmatis* ΔmtpA-comp (ΔmtpA carrying complementing mtpA expression plasmid), were grown to saturation and set to OD_{600} of 2.0 prior to infection. Equal amounts of phage were used to infect and cells and plaques evaluated after incubation for 48 hrs. at $37^{o}C$. Figure 14A shows that both wild type Girr and Girr_Δ*lysF1b* exhibit a modest but significant increase in plaque size when infected into the *M. smegmatis* ΔmtpA strain compared to the wild type *M. smegmatis* or the *M. smegmatis* ΔmtpA-comp strain. Girr_WT plaques increased by 1.4-fold from 2.8mm to 3.9mm while Girr_Δ*lysF1b* increased by 1.8-fold from 0.93mm to 1.7mm (Figure 14B). To account for the possibility that the change is plaque size were caused by reduced cell growth and less *M. smegmatis* ΔmtpA bacteria growing over the courses of the experiment, studies were repeated using high density cultures at 2×10^8 CFU/infection. This level of bacteria will exacerbate small plaque phenotypes since the increased cells enhance absorption and reduce phage diffusion [27; 81-83]. In these studies, both Girr_WT and Girr_Δ*lysF1b* showed similar plaques size increases as the low-density experiments verifying that *M. smegmatis* lacking a formal LAM layer is generally less restrictive to lysis and phage release (RSP, data not shown). Since both

Girr_WT and Girr_Δ*lysF1b* show increased plaque size and the plaques of Girr_Δ*lysF1b* do not

reach Girr_WT size, the results indicate that the lack of the LAM layer does not effectively

complement the deletion of *lyF1b*. This is consistent with the proposed holin function of the

LysF1b and supports the hypothesis that the LysF1b and LysZ proteins have distinct functions

in the lysis pathway in their respective bacterial hosts although holin function for LysZ has not

been formally evaluated.

Analysis of Additional Genes Proposed to be Involved in the Lysis Pathway of F1 Cluster

Phages

It is currently unknown how the various endolysins encoded by actinobacteriophages

gain access to the peptidoglycan layer following phage infection. Although exogenous

expression of the lysin A from some phages is cytotoxic to *M. smegmatis*, there is no evidence

of signal sequences or SAR domains in these proteins [20, 29-32]. However, *Mycobacterium sp.*

appear to have several different secretion systems that do not require a signal sequence [77-

80]. Additionally, it has been proposed that phage Ms6 encodes a small chaperone protein

(gp23) required for the export of Lysin A because the deletion of the gene results in a small

plaque phenotype [48, 50]. Since both phage Girr and NBJ encode a homolog to gp23 in the

same general genomic area upstream of the Lysin A (gene *29*; Figure 15A), it was of interest to

determine if deletion caused a lysis defect in Girr and NBJ. Figure 14D shows that Girr gp29 is

73 amino acids and shares 76.2% identity to Ms6 gp23 while NBJ gp29 shares 76 of 77 amino

acids with gp23. BRED was utilized to delete gene *29* from Girr and two independent Girr_Δ*29*

phage mutants were isolated (Figure 15B). Girr_Δ*29-20* and Girr__Δ*29-25* showed no lysis

defect when compared to Girr_WT in the liquid lysis assay (Figure 15C) and showed identical

plaque size to Girr_WT (Figure 15E). BRED deletion of NBJ *29* (NBJ_Δ*29*) also showed wild

type plaque size (Figure 15E) and no difference in liquid lysis when compared to NBJ_WT

(RSP, data not shown). To explore the possible interaction of the Lysin A and gp29 further, the

proteins were modeled using AlphaFold3. The results presented in Figures S14-16 show that all

models produced data with ipTM values <0.4 indicating a very low confidence in the interactions

of the proteins. Thus, these results suggest that gp29 from phage Girr or NBJ is not involved in

the lysis pathway. These finding are consistent with the lethal phenotype observed when both

lysF1a and *lysF1b* are deleted since deletion of holin genes is lethal in phages that express a

cytoplasmic endolysin that must exit through the holin [10-12]. If there was another gene that

was absolutely required for the lysis pathway in Girr and NBJ, especially one that could

enhance the secretion of the Lysin A in a holin-independent manner, it is unlikely that deletion of

lysF1a and *lysF1b* would be lethal. A plausible explanation for the lysis defects observed in the

Ms6 studies is that the method of deletion of gene *23* somehow impacted the expression of

Lysin A (gene *24*) to generate a lysis phenotype. Full genome sequencing data of the Ms6

mutants was not presented or discussed [48, 50] and the requirement of a phage encoded

chaperone has not been identified in any other phage.

CONCLUSIONS AND IMPLICATIONS

The results presented in this report identify the novel LysF1a and LysF1b proteins as

required holin-like components of *M. smegmatis* lysis by F1 cluster phages. The data support

that LysF1a and LysF1b function together as holins to time the lysis event and disrupt the PMF,

however, it appears that LysF1a requires the presence of the LysF1b protein to be active as a

holin. Holin function is based on several key findings. First, the deletion of both *lysF1a* and

lysF1b genes is lethal to phage propagation (Figure 5). This is consistent with the results from

E. coli phages where deletion of canonical holins from phages that express cytoplasmic

endolysins are lethal [10-12]. In addition, deletion of both the *lysF1a* and *lysF1b* genes in the F1

cluster phage Ms6 also appears to be lethal as no viable phage were isolated from secondary

plaques [48]. Second, holin function for the LysF1b protein is directly demonstrated by the ability

of energy poisons to trigger premature lysis in phages that do not express LysF1a (Figure 10).

Third, holin function is supported for LysF1b instead of releasin function because there is no

defined SAR-endolysins in F1 cluster phages and the energy poisons do not trigger early lysis

by releasin in phage Mu [18]. Finally, holin function is supported for LysF1a by the ability to

recover LRM mutants in the Δ*lysF1b* background of both Girr and NBJ that map to the LysF1a

protein and create phage that trigger lysis early but are fully lysis competent (Figure 13).

The 1TMD LysF1b proteins represent a new class of holins and appear to be found

predominantly in phages that infect Gram positive hosts. HHpred analysis shows that the

LysF1b proteins have a high probability hit to Pfam PF10874 (DUF2746) that contains 906

proteins. Within this Pfam are hits to phages that infect *Mycobacterium sp, Gordonia sp,* and

Streptomyces sp and hits to *Micromonospora sp, Rothia sp, Saccharothrix sp, Nocardia*

terpenica, Propionicmonas paludicola, Microbacterium talmoniae, Xylanimonas oleitrophica and

Actinacidiphilia glaucinigra where the later hits are likely to proteins that are encoded by

prophages within the host bacteria. To assess whether additional phages in the

Actinobacteriophage Database at PhagesDB.org express a LysF1-b-like homolog [84], phages

infecting *M. smegmatis* were bioinformatically evaluated as described in Methods to identify the

lysis cassette and associated genes that encode proteins with 1TMD. Of the ~2,600 phages in

the database, 80% encoded a LysF1b-like protein with an N-out-C-in membrane topology and

the proteins were grouped to 20 different phams based on their amino acid identity [85, 86].

Representative proteins from each pham are presented in Figure S17. The proteins differ in size

from 92-150 amino acids, have several basic amino acids after the TMD as highlighted in Figure

3 and have between 30%-50% DEKRH charged amino acids in the C-terminal cytoplasmic

domain (Figure S17). Multiple sequence alignment indicates minimal amino acid identity

between the proteins (Figure S18) and the alignment also failed to identify any small regions of

high identity that might serve as functional domains. Thus, while most phages that infect *M.*

smegmatis in the PhagesDB.org database express a LysF1b-like protein, it is unclear whether

they all function like the LysF1b proteins from Girr and NBJ.

Since deletion of both *lysF1a* and *lysF1b* genes in F1 phages is lethal and the LysF1b

protein clearly can function independent of the LysF1a, what then is the function of the 2TMD

LysF1a proteins in the lysis pathway? As discussed throughout this report, a role of LysF1a as a

type II pinholin is not supported by the sequence data (Figure 3), or by AlphaFold3 models

(Figure S7). In fact, the amino acid sequence data and lack of lysis function in the absence of

LysF1b, better supports LysF1a as an inactive antiholin. However, deletion of the *lysF1a* gene

results in a lysis triggering *delay* in both Girr_Δ*lysF1b* and NBJ_ Δ*lysF1b* with the later delayed

over 60 minutes compared to NBJ_WT (Figure 8). A lysis delay is not consistent with the

function of LysF1a as a true antiholin to the LysF1b, since it would be expected that deletion of

the antiholin would result in *early* host lysis by Girr_Δ*lysF1a* or Girr_Δ*lysF1a* since the LysF1b

would not be inhibited from forming active multimers as shown for lambda S_{105} and phage 21

$S^{21}68$ [10-12]. The data also demonstrate that expression of Girr or NBJ LysF1a in the absence

of the LysF1b results in cell death with release of ATP, but that actual lysis and shape change of

the bacteria is not detected by changes to the OD_{600} (Figure 8). These results suggest that

expression of Girr or NBJ LysF1a also will ultimately disrupt the membrane to destroy the PMF,

but whatever lesion is produced cannot effectively facilitate the release of the endolysins and/or

phage particles. In this context, the key findings are 1) that destabilization of the PMF alone is

not sufficient to "activate" LysF1a into a lesion that supports wild type lysis or phage release

(Figure 10), and 2) that LRMs can be recovered in the Δ*lysF1b* background that have point

mutations LysF1a that result in a fully lysis competent LysF1a protein that happens to trigger

premature (i.e. unregulated) lysis (Figure 13). These findings suggest that amino acid changes

to the protein may induce conformational changes to the LysF1a that allow it to fully support

bacterial lysis and phage release in the absence of the LysF1b. Thus, an attractive working

model is that LysF1a is not an antiholin to LysF1b but simply is not lysis active without LysF1b.

Therefore, as expression of both proteins increases following infection, they reach a

concentration at which they can formally interact, and this interaction facilitates conformational

changes in the LysF1a that allow it to become active in lysis. The two proteins then function

together to facilitate the release of endolysins and/or phage particles. Such a model would be

like the holin/antiholin system in phage lambda and phage 21 where the phages express both

active holins and inactive antiholins and triggering results in an instantaneous conversion of all

proteins to active holins [10-12].

Currently, it is unclear how the LysF1a and LysF1b proteins interact and whether both

proteins together actively participate in the creation a heteromeric lesion the inner membrane.

Clearly, LysF1b alone can trigger lysis and facilitate significant phage release, but as stated

above, lysis timing and the efficiency of phage release are still not optimal. There is also no

precedent for two structurally distinct TMD proteins functioning together to make a lesion. The

lysis pathway of phage P2 utilizes two distinct proteins, holin Y and antiholin LysA, but LysA

protein appears to be fully regulatory to holin Y and is not converted into an active holin like the

antiholins S_{107} or $S^{21}71$ [11, 55, 56]. Thus, the interaction of two distinct holin monomers that

have different membrane topologies is unprecedented and will require direct study of LysF1a

and LysF1b proteins within membranes to resolve the structure. Such work could also directly

address whether the LysF1b also has interactions with cell wall components to facilitate phage

release as suggested for the LysZ proteins of *Corynebacterium* phages [57]. Whether the other

Mycobacteriophages that express different variants of the LysF1a and LysF1b proteins also

utilize them in the same way as Girr and NBJ will require additional deletion and

complementation studies. The current study therefore provides a framework to evaluate lysis

under physiological conditions in Gram-positive hosts. It is exciting to speculate that due to the

presence of *lysF1*-like and *lysZ*-like genes in so many phages, it may be that the phages that

infect Gram-positive hosts use a similar set of tools to complete the lysis pathway through

distinct molecular mechanisms just as observed for phages that infect *E. coli.*

DATA AVAILABILITY STATEMENT

All recombinant phage and expression plasmids used in this study are available upon request. The authors affirm that all data necessary for confirming the conclusions of this article are represented fully within the article and its tables and figures. Extended data, including plasmid Sanger sequencing data and confirmatory cytotoxicity assay data for all Girr and NBJ pExTra constructs can be found at the SEA-GENES project open access database GenesDB (https://genesdb.org). To access these data on genesDB.org, any user can register for a free account, and once logged in to this account, navigate from the home page to Cluster F1 phage Girr or NBJ. Full genome sequences of Girr and NBJ are available in GenBank through accession numbers MH669003 and MH399784, respectively.

ACKNOWLEDGMENTS

We are grateful to the members of the Science Education Alliance for their invaluable research support and for supplying phage and plasmid reagents, particularly Dainelle Heller, Viknesh Sivanathan, Graham Hatfull, Deborah Jacobs-Sera, Becky Garlena and Dan Russell. We also thank the Pittsburgh Bacteriophage Institute for genome sequencing of the phages described in this study. We want to thank A. McKitterick for helpful discussions regarding LysZ. We want to thank Y. Morita for generously sending the ΔmtpA *M. smegmatis* strains.

FUNDING

This project was funded in part by an Undergraduate Grants in Aid for Research from USF Chapter of Sigma Xi to KRH and a USF Internal Proposal Enhancement Grant to RSP.

REFERENCS

1. Mushegian AR. Are There 10(31) Virus Particles on Earth, or More, or Fewer? J Bacteriol. 2020;202(9).

2. Oliveira H, Melo LD, Santos SB, Nobrega FL, Ferreira EC, Cerca N, et al. Molecular aspects and comparative genomics of bacteriophage endolysins. J Virol. 2013;87(8):4558-70.

3. Ryan GL, Rutenberg AD. Clocking out: modeling phage-induced lysis of Escherichia coli. J

Bacteriol. 2007;189(13):4749-55.

4. Pang T, Fleming TC, Pogliano K, Young R. Visualization of pinholin lesions in vivo. Proc Natl

Acad Sci U S A. 2013;110(22):E2054-63.

5. Savva CG, Dewey JS, Deaton J, White RL, Struck DK, Holzenburg A, et al. The holin of

bacteriophage lambda forms rings with large diameter. Mol Microbiol. 2008;69(4):784-

93.

6. Savva CG, Dewey JS, Moussa SH, To KH, Holzenburg A, Young R. Stable micron-scale

holes are a general feature of canonical holins. Mol Microbiol. 2014;91(1):57-65.

7. Summer EJ, Berry J, Tran TA, Niu L, Struck DK, Young R. Rz/Rz1 lysis gene equivalents in

phages of Gram-negative hosts. J Mol Biol. 2007;373(5):1098-112.

8. Rajaure M, Berry J, Kongari R, Cahill J, Young R. Membrane fusion during phage lysis. Proc

Natl Acad Sci U S A. 2015;112(17):5497-502.

9. Berry J, Rajaure M, Pang T, Young R. The spanin complex is essential for lambda lysis. J

Bacteriol. 2012;194(20):5667-74.

10. Young R. Phage lysis: three steps, three choices, one outcome. J Microbiol.

2014;52(3):243-58.

11. Cahill J, Young R. Phage Lysis: Multiple Genes for Multiple Barriers. Adv Virus Res.

2019;103:33-70.

12. Cahill J, Holt A, Theodore M, Moreland R, O'Leary C, Martin C, et al. Spatial and temporal

control of lysis by the lambda holin. mBio. 2024;15(2):e0129023.

13. Park T, Struck DK, Dankenbring CA, Young R. The pinholin of lambdoid phage 21: control

of lysis by membrane depolarization. J Bacteriol. 2007;189(24):9135-9.

14. Barenboim M, Chang CY, dib Hajj F, Young R. Characterization of the dual start motif of a

class II holin gene. Mol Microbiol. 1999;32(4):715-27.

15. Blasi U, Young R. Two beginnings for a single purpose: the dual-start holins in the

880   regulation of phage lysis. Mol Microbiol. 1996;21(4):675-82.

881   16. Ramanculov E, Young R. An ancient player unmasked: T4 rl encodes a t-specific antiholin.

882   Mol Microbiol. 2001;41(3):575-83.

883   17. Ramanculov E, Young R. Genetic analysis of the T4 holin: timing and topology. Gene.

884   2001;265(1-2):25-36.

885   18. Chamblee JS, Ramsey J, Chen Y, Maddox LT, Ross C, To KH, et al. Endolysin Regulation

886   in Phage Mu Lysis. mBio. 2022;13(3):e0081322.

887   19. Raman SK, Siva Reddy DV, Jain V, Bajpai U, Misra A, Singh AK. Mycobacteriophages:

888   therapeutic approach for mycobacterial infections. Drug Discov Today.

889   2024;29(7):104049.

890   20. Payne KM, Hatfull GF. Mycobacteriophage endolysins: diverse and modular enzymes with

891   multiple catalytic activities. PLoS One. 2012;7(3):e34052.

892   21. Schmelcher M, Donovan DM, Loessner MJ. Bacteriophage endolysins as novel

893   antimicrobials. Future Microbiol. 2012;7(10):1147-71.

894   22. Arenskotter M, Broker D, Steinbuchel A. Biology of the metabolically diverse genus

895   Gordonia. Appl Environ Microbiol. 2004;70(6):3195-204.

896   23. Goodfellow M, Jones AL. Corynebacteriales ord. nov. Bergey's Manual of Systematics of

897   Archaea and Bacteria. 2015. p. 1-14.

898   24. Brennan PJ. Structure, function, and biogenesis of the cell wall of Mycobacterium

899   tuberculosis. Tuberculosis (Edinb). 2003;83(1-3):91-7.

900   25. Hart EM, Bernhardt TG. The mycomembrane. Curr Biol. 2025;35(3):R85-R6.

901   26. Catalao MJ, Pimentel M. Mycobacteriophage Lysis Enzymes: Targeting the Mycobacterial

902   Cell Envelope. Viruses. 2018;10(8).

903   27. Payne K, Sun Q, Sacchettini J, Hatfull GF. Mycobacteriophage Lysin B is a novel

904   mycolylarabinogalactan esterase. Mol Microbiol. 2009;73(3):367-81.

28. Marinelli LJ, Piuri M, Swigonova Z, Balachandran A, Oldfield LM, van Kessel JC, et al.

BRED: a simple and powerful tool for constructing mutant and recombinant

bacteriophage genomes. PLoS One. 2008;3(12):e3957.

29. Joshi H, Seniya SP, Suryanarayanan V, Patidar ND, Singh SK, Jain V. Dissecting the

structure-function relationship in lysozyme domain of mycobacteriophage D29-encoded

peptidoglycan hydrolase. FEBS Lett. 2017;591(20):3276-87.

30. Singh AK, Gangakhedkar R, Thakur HS, Raman SK, Patil SA, Jain V. Mycobacteriophage

D29 Lysin B exhibits promising anti-mycobacterial activity against drug-resistant

Mycobacterium tuberculosis. Microbiol Spectr. 2023;11(6):e0459722.

31. Pohane AA, Joshi H, Jain V. Molecular dissection of phage endolysin: an interdomain

interaction confers host specificity in Lysin A of Mycobacterium phage D29. J Biol Chem.

2014;289(17):12085-95.

32. Nair G, Jain V. An intramolecular cross-talk in D29 mycobacteriophage endolysin governs the

lytic cycle and phage-host population dynamics. Sci. Adv.2024;10(6):

33. Gil F, Catalao MJ, Moniz-Pereira J, Leandro P, McNeil M, Pimentel M. The lytic cassette of

mycobacteriophage Ms6 encodes an enzyme with lipolytic activity. Microbiology

(Reading). 2008;154(Pt 5):1364-71.

34. Gil F, Grzegorzewicz AE, Catalao MJ, Vital J, McNeil MR, Pimentel M. Mycobacteriophage

Ms6 LysB specifically targets the outer membrane of Mycobacterium smegmatis.

Microbiology (Reading). 2010;156(Pt 5):1497-504.

35. Loessner MJ, Gaeng S, Wendlinger G, Maier SK, Scherer S. The two-component lysis

system of Staphylococcus aureus bacteriophage Twort: a large TTG-start holin and an

associated amidase endolysin. FEMS Microbiol Lett. 1998;162(2):265-74.

36. Kwan T, Liu J, DuBow M, Gros P, Pelletier J. The complete genomes and proteomes of 27

Staphylococcus aureus bacteriophages. Proc Natl Acad Sci U S A. 2005;102(14):5174-

9.

37. Vybiral D, Takac M, Loessner M, Witte A, von Ahsen U, Blasi U. Complete nucleotide

sequence and molecular characterization of two lytic Staphylococcus aureus phages:

44AHJD and P68. FEMS Microbiol Lett. 2003;219(2):275-83.

38. Godinho LM, El Sadek Fadel M, Monniot C, Jakutyte L, Auzat I, Labarde A, et al. The

Revisited Genome of Bacillus subtilis Bacteriophage SPP1. Viruses. 2018;10(12).

39. Escobedo S, Campelo AB, Wegmann U, Garcia P, Rodriguez A, Martinez B. Insight into the

Lytic Functions of the Lactococcal Prophage TP712. Viruses. 2019 ;11(10).

40. Sao-Jose C, Parreira R, Vieira G, Santos MA. The N-terminal region of the Oenococcus

oeni bacteriophage fOg44 lysin behaves as a bona fide signal peptide in Escherichia coli

and as a cis-inhibitory element, preventing lytic activity on oenococcal cells. J Bacteriol.

2000;182(20):5823-31.

41. Gindreau E, Lonvaud-Funel A. Molecular analysis of the region encoding the lytic system

from Oenococcus oeni temperate bacteriophage phi 10MC. FEMS Microbiol Lett.

1999;171(2):231-8.

42. Nelson D, Schuch R, Zhu S, Tscherne DM, Fischetti VA. Genomic sequence of C1, the first

streptococcal phage. J Bacteriol. 2003;185(11):3325-32.

43. Pope WH, Mavrich TN, Garlena RA, Guerrero-Bustamante CA, Jacobs-Sera D,

Montgomery MT, et al. Bacteriophages of Gordonia spp. Display a Spectrum of Diversity

and Genetic Relationships. mBio. 2017;8(4).

44. Pollenz RS, Bland J, Pope WH. Bioinformatic characterization of endolysins and holin-like

membrane proteins in the lysis cassette of phages that infect Gordonia rubripertincta.

PLoS One. 2022;17(11):e0276603.

45. McGarrah CEE, Algarin-Martinez ED, Cavasini MED, Correa V, Danielson DF, Dean WR, et

al. Isolation and Annotation of Azira, a CT Cluster Phage That Infects Gordonia

rubripertincta. Microbiol Resour Announc. 2023;12(7):e0034723.

46. Dedrick RM, Marinelli LJ, Newton GL, Pogliano K, Pogliano J, Hatfull GF. Functional

requirements for bacteriophage growth: gene essentiality and expression in

mycobacteriophage Giles. Mol Microbiol. 2013;88(3):577-89.

47. Sandoval AM, Abram AM, Alhabib ZM, Antonyan AS, Brikho SM, Buhay SI, et al. Complete

Genome Sequences of Cluster G Mycobacteriophage Darionha, Cluster A

Mycobacteriophage Salz, and Cluster J Mycobacteriophage ThreeRngTarjay. Microbiol

Resour Announc. 2020;9(20).

48. Catalao MJ, Gil F, Moniz-Pereira J, Pimentel M. Functional analysis of the holin-like proteins

of mycobacteriophage Ms6. J Bacteriol. 2011;193(11):2793-803.

49. Garcia M, Pimentel M, Moniz-Pereira J. Expression of Mycobacteriophage Ms6 lysis genes

is driven by two sigma(70)-like promoters and is dependent on a transcription

termination signal present in the leader RNA. J Bacteriol. 2002;184(11):3034-43.

50. Catalao MJ, Gil F, Moniz-Pereira J, Pimentel M. The mycobacteriophage Ms6 encodes a

chaperone-like protein involved in the endolysin delivery to the peptidoglycan. Mol

Microbiol. 2010;77(3):672-86.

51. Bavda VR, Jain V. Deciphering the Role of Holin in Mycobacteriophage D29 Physiology.

Front Microbiol. 2020;11:883.

52. Bavda VR, Yadav A, Jain V. Decoding the molecular properties of mycobacteriophage D29

Holin provides insights into Holin engineering. J Virol. 2021;95(10).

53. Kamilla S, Jain V. Mycobacteriophage D29 holin C-terminal region functionally assists in

holin aggregation and bacterial cell death. FEBS J. 2016;283(1):173-90.

54. Ford ME, Sarkis GJ, Belanger AE, Hendrix RW, Hatfull GF. Genome structure of

mycobacteriophage D29: implications for phage evolution. J Mol Biol. 1998;279(1):143-

64.

55. To KH, Dewey J, Weaver J, Park T, Young R. Functional analysis of a class I holin, P2 Y. J

Bacteriol. 2013;195(6):1346-55.

56. Christie GE, Calendar R. Bacteriophage P2. Bacteriophage. 2016;6(1):e1145782.

57. McKitterick AC, Lyerly EW, Bernhardt TG. Bacteriophages target membrane-anchored

glycopolymers to promote host cell lysis and progeny release. bioRxiv.

2025:2025.06.24.661397.

58. Hallgren J, Tsirigos KD, Pedersen MD, Almagro Armenteros JJ, Marcatili P, Nielsen H, et al.

DeepTMHMM predicts alpha and beta transmembrane proteins using deep neural

networks. bioRxiv. 2022:2022.04.08.487609.

59. Tsirigos KD, Peters C, Shu N, Kall L, Elofsson A. The TOPCONS web server for consensus

prediction of membrane protein topology and signal peptides. Nucleic Acids Res.

2015;43(W1):W401-7.

60. Almagro Armenteros JJ, Tsirigos KD, Sonderby CK, Petersen TN, Winther O, Brunak S, et

al. SignalP 5.0 improves signal peptide predictions using deep neural networks. Nat

Biotechnol. 2019;37(4):420-3.

61. Juncker AS, Willenbrock H, Von Heijne G, Brunak S, Nielsen H, Krogh A. Prediction of

lipoprotein signal peptides in Gram-negative bacteria. Protein Sci. 2003;12(8):1652-62.

62. Soding J, Biegert A, Lupas AN. The HHpred interactive server for protein homology

detection and structure prediction. Nucleic Acids Res. 2005;33(Web Server

issue):W244-8.

63. Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, et al. Accurate structure

prediction of biomolecular interactions with AlphaFold 3. Nature. 2024;630(8016):493-

500.

64. Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez, R,

Butcher S. The EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024.

Nucleic acids research. 2024;52(W1):W521–W525.

1006

65. Snapper SB, Melton RE, Mustafa S, Kieser T, Jacobs WR, Jr. Isolation and characterization

of efficient plasmid transformation mutants of Mycobacterium smegmatis. Mol Microbiol.

1990;4(11):1911-9.

66. Sparks IL, Kado T, Prithviraj M, Nijjer J, Yan J, Morita YS. Lipoarabinomannan mediates

localized cell wall integrity during division in mycobacteria. Nat Commun.

2024;15(1):2191.

67. Heller D, Amaya I, Mohamed A, Ali I, Mavrodi D, Deighan P, et al. Systematic

overexpression of genes encoded by mycobacteriophage Waterfoul reveals novel

inhibitors of mycobacterial growth. G3 (Bethesda). 2022;12(8).

68. Pollenz RS, Barnhill K, Biggs A, Bland J, Carter V, Chase M, et al. A genome-wide

cytotoxicity screen of cluster F1 mycobacteriophage Girr reveals novel inhibitors of

Mycobacterium smegmatis growth. G3 (Bethesda). 2024;14(5).

69. Wise B, Edwards K, Jirsa C, [student authors], Kanther M, Payne K, Pollenz RS, Heller DM.

Identification and characterization of host-modulating effectors encoded by the Cluster F1

mycobacteriophage NormanBulbieJr. bioRxiv 2025.

70. van Kessel JC, Hatfull GF. Mycobacterial recombineering. Methods Mol Biol. 2008;435:203-

15.

71. Gordon D, Abajian C, Green P. Consed: a graphical tool for sequence finishing. Genome

Res. 1998;8(3):195-202.

72. Russell DA. Sequencing, Assembling, and Finishing Complete Bacteriophage Genomes.

Methods Mol Biol. 2018;1681:109-25.

73. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.

Nat Methods. 2012;9(7):671-5.

74. Pang T, Park T, Young R. Mutational analysis of the S21 pinholin. Mol Microbiol.

2010;76(1):68-77.

75. Park T, Struck DK, Deaton JF, Young R. Topological dynamics of holins in programmed

bacterial lysis. Proc Natl Acad Sci U S A. 2006;103(52):19713-8.

76. Doermann AH. The intracellular growth of bacteriophages. I. Liberation of intracellular

bacteriophage T4 by premature lysis with another phage or with cyanide. J Gen Physiol.

1952;35(4):645-56.

77. Rigel NW, Gibbons HS, McCann JR, McDonough JA, Kurtz S, Braunstein M. The Accessory

SecA2 System of Mycobacteria Requires ATP Binding and the Canonical SecA1. J Biol

Chem. 2009;284(15):9927-9936.

78. Abdallah A, Gey van Pittius N, DiGiuseppe Champion P, Cox J, Luirink J, et. al. Type VII

secretion — mycobacteria show the way. Nat Rev Microbiol. 2007;5:883–891.

79. Bendtsen JD, Kiemer L, Fausbøll A, Brunak S. Non-classical protein secretion in bacteria.

BMC Microbiol. 2005;5:58.

80. Zhang Z, Dong L, Li X, Deng T, Wang Q. The PE/PPE family proteins of Mycobacterium

tuberculosis: evolution, function, and prospects for tuberculosis control. Front. Immunol.

2025;16:1606311.

81. Ameh EM, Tyrrel S, Harris JA, Pawlett M, Orlova EV, Ignatiou A, et al. Lysis Performance of

Bacteriophages with Different Plaque Sizes and Comparison of Lysis Kinetics After

Simultaneous and Sequential Phage Addition. Phage (New Rochelle). 2020;1(3):149-57.

82. Abedon ST, Yin J. Bacteriophage plaques: theory and analysis. Methods Mol Biol.

2009;501:161-74.

83. Dennehy JJ, Abedon ST, Turner PE. Host density impacts relative fitness of bacteriophage

Phi6 genotypes in structured habitats. Evolution. 2007;61(11):2516-27.

84. Russell DA, Hatfull GF. PhagesDB: the actinobacteriophage database. Bioinformatics.

2017;33(5):784-6.

85. Gauthier CH, Cresawn SG, Hatfull GF. 2022. PhaMMseqs: a new pipeline for constructing

phage gene phamilies using MMseqs2. G3 Genes Genomes Genetics.

2022;12(11):jkac233.

86. Gauthier CH, Hatfull GF. PhamClust: a phage genome clustering tool using proteomic

equivalence. mSystems. 2023;8(5):e00443-23.

 FIGURE LEGANDS

 Figure 1. Lysis cassette organization of Girr and NormanBulbieJr (NBJ). Predicted protein

 coding genes are shown as colored boxes. The genomic location is represented by the ruler,

 and the numbers represent the kilobase from nucleotide 1. Both lysis cassette regions are

 located upstream of the *tape measure* and *minor tail* genes. Genes above the genome are

 transcribed in the forward direct and genes below in the reverse direction. Genes with the same

 color encode proteins that are grouped to the same pham. Pink shading indicated nucleotide

 identity >90% in the shaded region. The *lysin A, lysin B,* 2TMD *lysF1a and* 1TMD *lysF1b* genes

 are identified. NKF = genes with no know function; *HNH* = HNH endonuclease.

 Figure 2. Amino acid sequence and topology of Girr and NBJ LysF1a proteins and

 comparison to phage 21 S^{21} pinholin. **A.** The amino acid sequences of Girr and NBJ LysF1a

 showing the location of the predicted TMDs and charged amino acids. **B.** The amino acid

 sequence of phage 21 $S^{21}68$ holin, $S^{21}71$ antiholin and $S^{21}68_{IRS}$ showing the location of the

 predicted TMDs and charged amino acids. **C.** Predicted membrane topology of Girr LysF1a,

 $S^{21}68_{IRS}$ and the proposed topology of the $S^{21}68$ holin homodimer with cytoplasmic TMD1.

 Figure 3. Amino acid sequence and topology of lysis proteins with 1TMD. **A.** The amino

 acid sequences of Girr LysF1b, NBJ LysF1b, Waterfoul gp32 and Mu releasin showing the

 location of the predicted TMD and the charged amino acids. **B.** AlphaFold3 structural

 predictions and membrane topologies of 1TMD proteins involved in lysis. pTM values: Mu, 0.51;

 NBJ LysF1b, 0.40; Girr LysF1b, 0.41; Rz, 0.57; T, 0.74. **C.** Predicted topologies of Girr LysF1b

 and Waterfoul gp32.

Figure 4. Deletion of Girr *lysF1a* and *lysF1b*. A. Genomic structure and PCR primer set

locations in Girr gene deletion mutants. Primer set 1 (P1) uses primers Girr-FL_*lysF1a* (F and

R) and generates a 755bp fragment in Girr_WT and 487bp with *lysF1a* deletion. Primer set 2

(P2) uses primers Girr-FL_*lysF1b* (F and R) and generates a 933bp fragment in Girr_WT and

594bp with *lysF1b* deletion. Primer set 3 (P3) uses the forward Girr-FL_*lysF1a* and reverse Girr-

FL_*lysF1b* and generates a 1189bp fragment in Girr_WT and either 921bp with *lysF1a* deletion

or 818b with *lysF1b* deletion. **B.** PCR verification of high titer lysates of Girr_WT, Girr_Δ*lysF1a*

and Girr_Δ*lysF1b* using the indicated primer sets. PCR band sizes are identified and show

correct deletions of the indicated genes.

1095

Figure 5. Deletion of both Girr *lysF1a* and *lysF1b*. A. Schematic showing experimental

procedure. **B.** DADA PCR was carried out on the indicated samples as described in Methods.

Negative control Girr_lysF1a shown in lanes 1 and 7. Positive control is the 1^o plaque pick with

the Girr_lysF1a/lysF1b phage and 325bp DADA band (+DADA) is shown in lanes 2 and 11.

Lanes 3-6 show the DADA PCR screen of 2^o lysates harvested from non-complementing *M.*

smegmatis that contained the indicated number of plaques (PFU). Lanes 8-10 show DADA PCR

screen of 2^o lysates harvested from complementing *M. smegmatis* carrying pExTra-Girr*34*. Note

the lack of a positive DADA band in all samples plated on non-complementing *M. smegmatis*.

1104

1105 **Figure 6 Plaque sizes in Girr_WT, Girr_Δ*lysF1a* and Girr_Δ*lysF1b*.** *M. smegmatis* was

1106 infected with identical numbers of phage and detailed in Methods and plaque size determined

1107 after 36 hrs. of growth at 37^oC. Scale = 1cm. Plaque volume schematics are presented to

1108 illustrate the plaque size differences.

1109

Figure 7. Complementation of plaque phenotype in Girr_Δ*lysF1b*. A. Plaque assays were

completed with *M. smegmatis* using Girr_WT or Girr_Δ*lysF1b* as described in Methods. Plaque

assays were completed with *M. smegmatis* complementing strains carrying the indicated

pExTra-Waterfoul expression plasmids using Girr_Δ*lysF1b* as described in Methods. Plaques

were evaluated following incubation at 37^{o}C for 36 hrs. Plaques sizes were quantified using

ImageJ and the average diameter +/- standard deviation for > 50 plaques is shown. * =

statistically different from Girr_WT, p *<0.0001*. ** = statistically different from Girr_Δ*lysF1b*

infected on *M. smegmatis* with pExTra Waterfoul_*32* with no aTc, p *<0.0001*. Scale = 1cm. **B.**

Plaque assays were completed with *M. smegmatis* using Girr_WT or Girr_Δ*lysF1b* as described

in Methods. Plaque assays were completed with *M. smegmatis* complementing strains carrying

the indicated pExTra-Waterfoul expression plasmids using Girr_Δ*lysF1b* as described in

Methods. Plaques were evaluated following incubation at 37^{o}C for 48 hrs. Plaques sizes were

quantified using ImageJ and the average diameter +/- standard deviation for > 50 plaques is

shown * = statistically different from Girr_WT, *p <0.0001*. ** = statistically different from

Girr_Δ*lysF1b* infected on *M. smegmatis* with pExTra01, *p <0.0001*. Scale = 1cm.

1125

Figure 8. Complementation of plaque phenotype in Girr_Δ*lysF1a*. Plaque assays were

completed with *M. smegmatis* using Girr_WT or Girr_Δ*lysF1a* as described in Methods. Plaque

assays were completed with *M. smegmatis* complementing strains carrying pExTra-Girr_*34*

expression plasmid using Girr_Δ*lysF1a*as described in Methods. Plaques were evaluated

following incubation at 37^{o}C for 48 hrs. Plaques sizes were quantified using ImageJ and the

average diameter +/- standard deviation for > 50 plaques is shown. * = statistically different from

Girr_WT, *p <0.0001*. ** = statistically different from Girr_Δ*lysF1a* infected on *M. smegmatis* with

pExTra-Girr_*34* with no aTc, *p <0.0001*. Scale = 1cm.

1134

Figure 9. Liquid lysis assay and ATP release from *M. smegmatis* after infection. At T_0 a log

growth culture of *M. smegmatis* at an OD_{600} of ~0.25 was infected with the indicated phage at

an MOI of 10:1 and incubated at $37^{\circ}C$ for 30 minutes without shaking. Control cultures (no

phage) did not receive phage. At T_{30} minutes the culture was constantly shaken a 225rpm,

aliquots removed at the indicated times and bacterial growth evaluated at OD_{600}. Open white

arrow shows the 30-minute infection time. **A.** Representative results following infection with

Girr_WT, Girr_$\Delta lysF1a$ and Girr_$\Delta lysF1b$. **B**. Representative results following infection with

NBJ_WT, NBJ_$\Delta lysF1a$ and NBJ_$\Delta lysF1b.$ **C.** Samples of culture were removed from the

cultures represented in A and B at the times indicated by the solid black triangle on the x axis.

The presence of ATP was evaluated as described in Methods.

1145

Figure 10. Energy poisons trigger early lysis in phage expressing LysF1b but not LysF1a.

At T_0 a log growth culture of *M. smegmatis* at an OD_{600} of ~0.25 was infected with the indicated

phage at an MOI of 10:1 and incubated at $37^{\circ}C$ for 30 minutes without shaking. Control cultures

(no phage) did not receive phage. At T_{30} minutes the culture was constantly shaken a 225rpm,

aliquots removed at the indicated times and bacterial growth evaluated at OD_{600}. **A.**

Representative results following infection with Girr_WT, and Girr_$\Delta lysF1a$. KCN (final

concentration of 10mM) was added to a culture infected with Girr_$\Delta lysF1a$ and an uninfected

culture at 110 minutes (open arrow). **B**. Representative results following infection with NBJ_WT,

and NBJ_$\Delta lysF1a.$ KCN (final concentration of 10mM) was added to a culture infected with

NBJ_$\Delta lysF1a$ and uninfected culture at 120 minutes (open arrow). **C.** Representative results

following infection with Girr_WT, and Girr_$\Delta lysF1b$. KCN (final concentration of 10mM) or $CHCl_3$

(final concentration 1%) was added to cultures infected with Girr_$\Delta lysF1b$ and uninfected

cultures at 180 minutes (open arrow). **D**. Representative results following infection with

NBJ_WT, and NBJ_$\Delta lysF1b.$ KCN (final concentration of 10mM) or $CHCl_3$ (final concentration

1%) was added to cultures infected with NBJ_Δ*lysF1b* and uninfected cultures at 180 minutes

(open arrow).

1162

Figure 11. One-step growth curves of *M. smegmatis* infected with Girr_Δ*lysF1a* and

Girr_Δ*lysF1b*. A. One-step growth curves using Girr_WT, Girr_Δ*lysF1a* and Girr_Δ*lysF1b* were

completed as described in Methods. Each time point represents the average and standard

deviation from 3 different experiments. **B**. The data for Girr_Δ*lysF1b* from panel A replotted with

a different Y axis than used in A. **C**. Burst size was determined as described in Methods. The

average bust size +/- standard deviation is from three different experiments and represents the

average of all points once the curve reached a plateau.

1170

Figure 12. Isolation of Girr_Δ*lysF1b*-LRM mutants. **A**. *M. smegmatis* was infected with

GirrΔ*lysF1b*, plate lysates recovered from the plaque assay and *M. smegmatis* infected with the

recovered phage. Panel A shows a representative plaque assay following the second round of

infection. A large plaque is circled. **B**. Phage were isolated from the large plaque in A, serial

diluted and used for plaque assay. Plates were incubated at 37^{o}C for 36 hrs. **C**. Plaque assay

from Girr_WT phage. Plates were incubated at 37^{o}C for 36 hrs. Scale bar = 1cm in all images.

1177

Figure 13. Liquid lysis and one-step assay of Girr and NBJ LRM mutants. At T_0 a log

growth culture of *M. smegmatis* at an OD_{600} of ~0.25 was infected with the indicated phage at

MOI 10:1 for 30 minutes at 37^{o}C without shaking. Control cultures (no phage) did not receive

phage. At T_{30} minutes the culture was constantly shaken a 225rpm and aliquots removed over

time. *M. smegmatis* growth was evaluated at OD_{600}. **A**. Representative results following infection

with Girr_WT and indicated LRM mutants. **B**. Representative results following infection with

NBJ_WT and indicated LRM mutants **C.** One-step growth curves using Girr_WT and

Girr_Δ*lysF1b*-LRM1 and Girr_Δ*lysF1b*-LRM2 were completed as described in Methods. Each

time point represents the average and standard deviation from 3 different experiments.

1187

Figure 14. Analysis of *lysF1b* phage plaque sizes in *M. smegmatis* lacking the

lipoarabinomannan layer. A. Representative plaque assays using Wild type *M. smegmatis, M.*

smegmatis ΔmtpA-comp or *M. smegmatis* ΔmtpA cultures infected with the same amount of

Girr_WT or Girr_Δ*lysF1b* and incubated for 48 hrs. at 37^oC. Ruler = 1cm. **B**. Plaque sizes were

determined as described in Methods. The average plaque diameter +/- standard deviation for 25

plaques is shown.

1194

Figure 15. Analysis of NKF gene *29*. A. Gene organization of Girr, NBJ and Ms6 lysis cassette

region. Predicted protein coding genes are shown as colored boxes. The genomic location is

represented by the ruler, and the numbers represent the kilobase from nucleotide 1. Genes

above the genome are transcribed in the forward direct and genes below in the reverse

direction. Genes with the same color encode proteins that are grouped to the same pham. Pink

shading indicated nucleotide identity >90% in the shaded region. *NKF* = Girr_*29*, NBJ_*29* and

Ms6_*23*. **B**. PCR analysis of Girr_Δ*29*-20 and Girr_Δ*29*-25. **C**. Liquid lysis assay of the indicated

phages was carried out as described in Methods. **D**. Amino acid alignment of Girr gp29, NBJ

gp29 and Ms6 gp23 by Clustal Omega. **E**. Plaque assay of Girr_WT, Girr_Δ*29-20*, Girr Δ*29-25*,

NBJ_WT, and NBJ_Δ*29.* Ruler = 1cm. Plaque sizes were determined as described in Methods.

The average plaque diameter +/- standard deviation for 25 plaques is shown.

1206

1207

1208

A.

amino acid #

| NBJ_gp32 LysF1a | MLTRSFWIDAAER**AIRTFAQTAIATLGAGAVDLMTTDWISV**LSVSGGAAVVSLLMSIGAERRGNPGTASATRAVTAA |
| Girr_gp34 LysF1a | MQAMLTRSFWIDAAER**AARTFAQTAIATLGAGAVDLLATDWVV**LSVSGGAAVVSLLMSIGAERRGNPGTASATRAVTAA |

B.

amino acid #

Phage 21 S²¹68	MDK**ISTGIAYGTSAGSAGYWFLQWL**DQVSPSQ**WAAIGVLGSLVLGFLTYLTNLYF**KIREDRRKAARGE
Phage 21 S²¹71	MKSMDK**ISTGIAYGTSAGSAGYWFLQWL**DQVSPSQ**WAAIGVLGSLVLGFLTYLTNLYF**KIREDRRKAARGE
Phage 21 S²¹68ᵢᵣₛ	MRYIRSDK**ISTGIAYGTSAGSAGYWFLQWL**DQVSPSQ**WAAIGVLGSLVLGFLTYLTNLYF**KIREDRRKAARGE

| ▇ alternative start codon | ▇ G in helix | **__Underline = TMD domains__** |
| ▇ charged residue not in helix | ▇ polar residue | |

C.

A.

MIWESVREAVDAAYQPDDGIDL**IGLLIIGLPSTIAAIGTGIVGV**LTV**R**GQ**RK**G**R**ERARQIDAKTDEIHEQTVNTHDTNMRDDLDEIRDLVRDGFKQIQRDIGGLREELRTERLERIEGDKRRD Girr gp35 LysF1b

MIWESVREAVDAAYQPDDGIDL**IGLLIIGLPSTIAAIGTGIVGV**LTV**R**GQ**RK**G**R**ERARRIDAKTYEIHEQTVNTHDTNMRDDLDEIRDLVRDGFKQIQRDIGGLREELRTERLERIEGDKRRD NBJ gp33 LysF1b

MTGSAVSALPSD**WYGLAGLAVVNAAGLLALFM**K**A**R**GTHKQVANSH**STNLRDDIDGLRDHIVELGRDIGGMRTDVGQLRGEVRDDRTHTHHRFRRIEDKLDELAGRGARESG Waterfoul gp32

M**DLISVLALWPYLLPVVAGGAVWA**RRSFASTERVERLENRMTEMETRYASIPGTEDVHEMRLQIAELSGDIRVLSQRVQSFSHQLELLLENAVNRSNS Mu releasin

NH₂ — TMD1 — basic — KRHDE residues — COOH

Green = TMD domain   Basic region after TMD   + Charged   - Charged

B.

Mu releasin

NBJ gp33 LysF1b

Girr gp35 LysF1b

Lambda Rz spanin

T4 T Type III holin

24/75 (32%) charged

40/79 (50%) charged

40/79 (50%) charged

C.

Girr gp35 LysF1b (124aa)

Waterfoul gp32 (111aa)

extracellular

(1)M   (22)

M(1)   (12)

TMD

TMD

cytoplasmic

(45)   (36)

COOH (124)   COOH (111)

40/79 (50%) charged   36/75 (48%) charged

A.

Girr_WT

lysin A — 31
HNH — 32
lysin B — 33
lysF1a — 34
lysF1b — 35
NKF — 36
27   28   29   30

Girr_ΔlysF1a

lysin A — 31
HNH — 32
lysin B — 33
lysF1b — 35
NKF — 36
27   28   29   30

Girr_ΔlysF1b

lysin A — 31
HNH — 32
lysin B — 33
lysF1a — 34
NKF — 36
27   28   29   30

▶ ◀ Primer set 1 (P1): Girr-FL_lysF1a F & R

▶ ◀ Primer set 2 (P2): Girr-FL_lysF1b F & R

▶ ◀ Primer set 3 (P3): Girr-FL_lysF1a-F and Girr-FL_lysF1b-R

B.

Girr_WT   Girr_ΔlysF1a   Girr_ΔlysF1b
M   P1   P2   P3   P1   P3   P2   P3

3000bp ▷
1500bp ▷
1000bp ▷
100bp ▷

755bp   963bp   1189bp   487bp   921bp   594bp   818bp

Girr_WT

Plaque volume comparisons

Girr_WT

Girr_ΔlysF1a

Girr_ΔlysF1b

10mm

Girr_ΔlysF1a

Girr_ΔlysF1b

Girr_WT
M. Smegmatis
4.56 +/- 0.65 mm

Girr_*ΔlysF1a*
M. Smegmatis
pExTra-Girr*34* + *aTc*
**4.15 +/- 0.55 mm

Girr_*ΔlysF1a*
M. Smegmatis
*2.79 +/- 0.55mm

Girr_*ΔlysF1a*
M. Smegmatis
pExTra-Girr*34 no aTc*
2.83 +/- 0.353 mm

A.

B.

C.

phage	Burst Size PFU/cell
Girr_WT	277 +/- 81
Girr_ΔlysF1a	255 +/- 75
Girr_ΔlysF1b	*29.2 +/- 36

*statistically significant from WT phage *p<0.0001.*

A. Girr_*ΔlysF1b*

B. Girr_*ΔlysF1b*-LRM
3.01 +/- 0.26mm

C. Girr_WT
2.94 +/- 0.25mm

A.

| *M. Smegmatis* Girr_WT | ΔmtpA-Comp Girr_WT | ΔmtpA Girr_WT |

| *M. Smegmatis* Girr_Δ*lysF1b* | ΔmtpA-Comp Girr_Δ*lysF1b* | ΔmtpA Girr_Δ*lysF1b* |

B.

* *p <0.0001* to WT
** *p <0.0001* to *M smeg* WT
*** *p <0.0001* to *M smeg* Δ*lysF1b*

A.

Girr

NKF NKF NKF *NKF* NKF

27 30 *lysin A* *HNH* *lysin B* *lysF1a* *lysF1b* NKF

26 28 29 31 32 33 34 35 36

25 26 27 28 29 30

Ms6

NKF NKF NKF NKF *NKF* *lysin A* *lysF1a* *lysF1b* NKF

19 21 20 22 23 24 *lysin B* 26 27 28

20 21 22 23 24 25

NBJ

NKF NKF NKF *NKF* *lysin A* *lysin B* *lysF1a* *lysF1b* NKF

24 27 26 28 29 30 31 32 33 34

NKF

25 26 27 28 29

B.

M   P4   P3   P4   P3   P3   P4

1000bp
500bp
100bp

Girr_WT   Girr_WT   Girr_Δ29-20   Girr_Δ29-20   Girr_Δ29-25   Girr_Δ29-25

▶◀ Primer set 3 (P3): Girr-FL_*lysF1a* F and Girr-FL_*lysF1b* R

▶◀ Primer set 4 (P4): Girr-FL_*29* F & R

C.

○ No phage
● Girr_WT
▲ Girr_Δ29-20
△ Girr_Δ29-25

OD$_{600}$ vs Time After Start of Infection (minutes)

D.

```
Girr_29   MDRLGIILLKLLGPLADRIADRIADRITENLPDLSDLDDQIVAKLPDLTNLPEQVINIIGRLPRFPFLLGGKP
Ms6_23    MDRLGIVLLKLFGPLADRIADRIADRIAENLPDLSNLDDQIVAKLPDLTNLPAQVMDIIDSALRSIPVLGGILGSKR
NBJ_29    MDRLGIILLKLFGPLADRIADRIADRITENLPDLSNLDDQIVAKLPDLTNLPAQVMDIIDGALRSIPVLGGILGSKR
```

E.

Girr_WT
3.11 +/- 0.12 nm

Girr_Δ29-20
2.98 +/- 0.16 nm

Girr_Δ29-25
3.02 +/- 0.14 nm

NBJ_WT
2.25 +/- 0.25 nm

NBJ_Δ29
2.15 +/- 0.09 nm