**Title:** Preoptic Neurons that Regulate Hibernation Entry

**Authors:** Adrian J. Martinez[1,2]†, Christopher M. Reid[1]†, Aurora J. Lavin-Peter[1], Wenhui

Li[1], Andrew S. Lee[1], Eric C. Griffith[3], Sinisa Hrvatin[1,4,5]*

**Affiliations:**

[1]Whitehead Institute for Biomedical Research, Cambridge, MA 02142, USA

[2]Department of Brain and Cognitive Sciences, Massachusetts Institute of

Technology, Cambridge, MA 02142, USA

[3]Department of Neurobiology, Harvard Medical School, Boston, MA 02115, USA

[4]Department of Biology, Massachusetts Institute of Technology, Cambridge, MA

02142, USA

[5]Howard Hughes Medical Institute, Massachusetts Institute of Technology,

Cambridge, MA, USA

*Corresponding author. Email shrvatin@wi.mit.edu (S.H.)

† These authors contributed equally to this work

**Abstract:** Seasonal hibernation allows species to survive harsh environments by entering a profound hypometabolic and hypothermic state called deep torpor (DT). However, the neural mechanisms controlling entry into this state remain poorly understood. Here, we map neuronal activity across hibernation stages in the Syrian hamster and identify DT-associated activity in the hypothalamic anterior preoptic area (aPOA) as essential for hibernation entry. Single-nucleus multiomic profiling enabled the design of a novel enhancer adeno-associated virus selectively targeting candidate glutamatergic neuronal subpopulations. Using this tool, we demonstrate that activity of a subpopulation of *Samd3*-positive glutamatergic neurons in the hamster aPOA is both necessary and sufficient for DT induction. Together, our findings identify a neuronal population regulating hibernation entry and provide the first genetic access to manipulate this complex behavior.

**Main Text:** The ability to maintain body temperature ($T_b$) within a narrow range is a

defining feature of homeotherm physiology[1,2]. However, to survive extreme environmental

conditions, many mammals can initiate hypothermic and hypometabolic energy-

conserving strategies such as hibernation or daily torpor[3–8]. While $T_b$, heart rate, and

metabolic rate decreases observed in daily torpor last only several hours[9,10], seasonal

hibernation is characterized by deeper, repeated multi-day bouts of torpor periodically

interrupted by arousals to euthermic $T_b$[5,11–15]. Decades of study have characterized the

homeostatic thermoregulatory circuitry that detects and responds to internal and external

thermal cues[1,2,16–27]. However, despite evidence that ambient and internal temperature[3],

circadian clocks[28,29], adenosine, melatonin, and thyroid hormone signaling[30–39], and the

peripheral nervous system[40] modulate hibernation dynamics, the neural mechanisms that

initiate and sustain deep torpor (DT) in hibernating animals remain largely unexplored.

To probe the neural mechanisms that control entry into hibernation-associated deep

torpor, we established a laboratory paradigm of hibernation in the Syrian hamster

(*Mesocricetus auratus*), implanting each animal with telemetric temperature probes for

continuous $T_b$ monitoring. To induce hibernation, adult (5-6-week-old) animals housed at

room temperature (~22ºC) with standard photoperiod (12 hours light (L): 12 hours dark

(D)) were first shifted to a short photoperiod (8L:16D) for 8-10 weeks, followed by transfer

to 4-5°C ambient temperature conditions to simulate seasonal changes (Fig. 1a, b,

Extended Data Fig. 1a). While this paradigm induced hibernation in both males and

females, female hamsters were initially found to hibernate more readily (data not shown),

with ~80% initiating torpor bouts within six weeks of ambient temperature reduction. As a

result, we restricted our subsequent analyses to females.

While typical torpor-associated $T_b$ decreases followed an exponential decay curve (n

= 1262 torpor bouts, Extended Data Fig. 1b), animals exhibited some variability in torpor

dynamics. Altogether, 69% of animals (n = 271) achieved DT (minimal $T_b$ = 5.5 ± 0.5°C,

bout duration 60.3 ± 13.4 h) and 11% (n = 42) exhibited only shallow torpor bouts (minimal

$T_b$ = 12.2 ± 3.1°C, bout duration 6.8 ± 1.8 h), with 20% (n = 81) failing to initiate torpor

during the 6-week observation period (Fig. 1c-f, Extended Data Fig. 1c). Periods of

interbout euthermia (IBE) between DT bouts lasted 15.3 ± 2.8 h (n = 2471 IBE periods,

Fig. 1e). Moreover, consistent with prior reports[41], torpor entry occurred preferentially

during subjective night (~ZT17-18) (n = 3,595 torpor bouts, Fig. 1g) and was

characterized by a stereotyped nesting behavior with curled posture and absence of gross

motor movement, food, or water intake[5,15]. By contrast, arousal from DT exhibited no clear

circadian pattern (Fig. 1h), consonant with disrupted circadian clock oscillations at

temperatures below 15°C[41,42].

**POA activity regulates hamster entry into deep torpor**

Having established laboratory conditions for efficient hibernation induction, we

employed this system to investigate neural activity changes associated with DT onset. To

this end, we collected brain samples during DT and IBE, as well as from non-hibernating

control animals, and performed immunostaining for FOS, a marker of neuronal activity

(Fig. 2a). Given prior reports of neuronal activity marker induction in hypothalamic regions

during DT bouts[41,43,44], and classic studies showing that hypothalamic lesions disrupt $T_b$

regulation during hibernation[45], we focused our analysis primarily on the hypothalamus.

Among the 32 distinct regions analyzed, we observed DT-active regions associated with

circadian rhythm modulation (Suprachiasmatic nucleus)[46–48], energy homeostasis

(Arcuate nucleus)[49–51], and arousal (Paraventricular nucleus of the thalamus)[52,53].

Notably, the anteroventral periventricular nucleus (AVPe) within the anterior POA

exhibited the highest degree of FOS induction in DT relative to IBE or non-hibernating

control samples (14.1% vs. 6.9% and 0.8%, $P = 0.0685$ and $P < 0.001$, respectively) (Fig.

2b-d, Supplementary Tables 1, 2, Extended Data Fig. 2a, b).

To test directly whether hypothalamic neuronal activity is required for the initiation and

maintenance of hibernation, we stereotaxically injected adeno-associated virus (AAV)

expressing an inhibitory Gi-coupled DREADD (Designer Receptor Exclusively Activated

by Designer Drug; AAV-hSyn-hM4D(Gi)-mCherry)[54] into hibernating hamsters (Fig. 2e),

systematically targeting different areas of the anterior hypothalamus that exhibited

substantial DT-associated activity. After allowing animals to recover and resume DT

bouts, we administered the DREADD agonist clozapine (CLZ) via drinking water. As

animals consumed CLZ during their IBE periods, we observed a cessation of torpor bouts

in a subset (4/13) of the tested animals, while all animals that underwent sham surgery

and received CLZ (5/5) readily re-entered torpor (Fig. 2f-h, Extended Data Fig. 2c).

Correlation of the pattern of Gi-DREADD expression across animals with the degree of

DT perturbation identified subregions of the anterior POA (aPOA), including the medial

preoptic area (MPO), medial preoptic nucleus (MPN), and AVPe, as regions whose

silencing was associated with perturbed hibernation (Fig. 2h-j, Supplementary Table 3).

Together, these findings implicate the anterior preoptic area as a key region whose

activity is necessary for the initiation of DT bouts during hibernation.

**Cross-species transcriptomic profiling identifies DT-active neuronal populations**

The aPOA has been implicated in the regulation of fasting-induced daily torpor (FIT) in mice[43,55–59]. Our chemogenetic silencing studies in hamster therefore raised the possibility that this region plays a conserved role in the control of hypometabolic states across mammals. To gain further insight into the diversity of neural cell types in the hamster POA and their relation to the cellular composition of the murine POA, we dissected and profiled the POA of 7 adult hamsters (4 females and 3 males) using single-nucleus paired RNA- and ATAC- sequencing (multiome) (Fig. 3a). We annotated the hamster POA cells by aligning and integrating the resulting data with recently published mouse POA multiome and spatial transcriptomic datasets[60,61], generating the first multiomic cross-species POA cellular taxonomy (Extended Data Fig. 3-5).

Since activity of brain regions within the median and medial POA was essential for DT entry (Fig. 2i), we focused our analysis on these regions. Graph-based clustering identified major neuronal and non-neuronal cell classes, with further subclustering of neuronal cells ($n$ = 26,761 cells) identifying 15 distinct GABAergic and 11 glutamatergic neuronal cell types shared between hamster and mouse (Fig. 3b-d, Extended Data Fig. 4). Importantly, every neuronal cluster contained both mouse and hamster cells (Fig. 3c), and pair-wise transcriptional similarity analysis (cosine similarity score) between all hamster and murine neuronal cell clusters revealed consistent 1:1 cross-species relationships, further supporting the validity of our cross-species atlas alignment (Fig. 3e). Moreover, with the exception of a few cell types, the relative cell type abundance across POA neuronal clusters was largely preserved between hamster and mouse (Extended

Data Fig. 5e). Thus, our analyses reveal broad cellular homology between hamster and

mouse POA.

Focusing more closely on POA neuronal populations implicated in FIT in the mouse,

we found that the expression patterns of molecular markers of mouse FIT-associated

neurons (*Slc17a6/Vglut2, Adcyap1, Esr1,* and *Ptger3*) were broadly conserved across

hamster and mouse POA cell types (Fig. 3f, g). Moreover, *in situ* hybridization (ISH)

analysis indicated that *Vglut2* and *Adcyap1*, two broad markers of FIT-regulating cells,

labeled neurons in similar anatomical locations in the mouse and hamster aPOA,

including the AVPe, MPO, and MnPO (animals = 4, Fig. 3h, Extended Data Fig. 6).

Leveraging these findings, we carried out combined immunostaining and ISH on aPOA

samples from hamsters (n = 3) in deep torpor and found that the majority of DT-active

(FOS-positive) neurons express *Slc17a6/Vglut2* (65.2 ± 4.2%, animals = 3), with a

considerable portion expressing *Adcyap1* (34.0 ± 1.4%, animals = 3), and relatively few

expressing *Slc32a1/Vgat* (12.8 ± 1.5%, animals = 3, Fig. 3i, j, Extended Data Fig. 7a, b).

Together, these findings suggest that homologous aPOA glutamatergic neuronal

populations may be active during mouse FIT and hamster DT.

To further test this idea, we leveraged anatomical information from our transcriptomic

and anatomical atlas to identify Glut-04, -06, -07, -09, -10, and -11 clusters as

corresponding to FIT-regulating neurons (hereafter collectively termed aPOA$^{Vglut2}$

neurons; Fig. 3k, Extended Data Fig. 7c), and expression of the *Samd3* gene was found

to exclusively mark this population (Fig. 3l). Indeed, a substantial fraction of hamster DT-

active (FOS+) neurons also expressed *Samd3* (46.4 ± 4.1% of FOS+ AVPe neurons are

positive for *Samd3* mRNA, Fig. 3m, n), further supporting the idea that aPOA$^{Vglut2}$ neurons

are active during DT entry and might regulate entry into this state. Together, our findings

establish the first cross-species atlas of the aPOA, a key region whose activity is

necessary for the initiation of DT bouts during hibernation, uncovering broad homology

between mouse and hamster cell types and implicating aPOA$^{Vglut2}$ neurons as promising

candidate drivers of DT entry in the hamster.

**149 Enhancer-driven viral tools to access aPOA$^{Vglut2}$ neurons**

To assess the functional role of aPOA$^{Vglut2}$ neurons in hamster DT entry, we required

the ability to selectively target and manipulate these cell populations *in vivo*. Transgenic

options in the hamster remain limited, and a *CamkIIa* promoter-driven AAV exhibits poor

specificity for glutamatergic neurons in the hamster POA (42.6 ± 2.0%, animals = 4,

Extended Data Fig. 8). Recently, AAV constructs incorporating compact cell-type-specific

enhancer gene regulatory elements (GREs) have offered a promising alternative to

traditional transgenic technologies in nonstandard species[62–68]. We therefore sought to

generate new enhancer-AAV tools driving restricted payload expression in the target

aPOA$^{Vglut2}$ cell population.

To develop a viral approach to selectively target aPOA$^{Vglut2}$ neurons, we leveraged the

extensive murine single-nucleus chromatin accessibility (snATAC-seq) data included in

our multiomic cross-species atlas to identify conserved genomic elements exhibiting

selectively enhanced accessibility in aPOA$^{Vglut2}$ neuron cell populations (Fig. 4a). First,

we employed a modeling-based approach to identify cell-type-specific nucleosome-

depleted gene regulatory regions. Subsequently, we performed differential accessibility

analysis to identify putative GREs with enriched accessibility in aPOA$^{Vglut2}$ neuronal

populations relative to other neuronal subtypes (Fig. 4b, Extended Data Fig. 9), and

filtered GREs based on proximity to aPOA$^{Vglut2}$ neuron-restricted marker genes (Fig. 4c).

Notably, using this approach, we identified a conserved 500 bp element proximal to the

mouse *Samd3* gene, hereafter termed *Samd3E*, that exhibited the highest accessibility

enrichment in aPOA$^{Vglut2}$ neurons relative to caudal POA$^{Vglut2}$ neurons and POA$^{Vgat}$

neurons (FDR < $10^{-300}$, $\log_2$Fold Enrichment = 3.1, Fig. 4b-e), with similar aPOA$^{Vglut2}$

neuron-enriched accessibility also observed in the hamster (P = $10^{-173}$, $\log_2$Fold

Enrichment versus other populations = 3.2, Fig. 4f, Extended Data Fig. 9).

The resulting candidate mouse GRE sequences were synthesized and cloned into an

AAV backbone upstream of a minimal beta-globin promoter sequence and a red

fluorescent protein (RFP) reporter transgene prior to packaging into functional AAV8

particles for *in vivo* evaluation. Notably, the *Samd3E* element conferred remarkable

specificity, restricting reporter expression almost exclusively to *Vglut2*+ neurons in the

hamster aPOA (95.2 ± 1.2%; animals = 3), with substantial co-labeling of *Samd3*+ cells

(58.7 ± 3.2%, animals = 3, Fig. 4g, h, Extended Data Fig. 10). Taken together, these

findings highlight the power of this approach to generate viral tools to study brain circuits

across species, including in hibernating animals, and establish the *Samd3E*-driven AAV

(Samd3E-AAV) as an effective tool for achieving targeted genetic access to glutamatergic

neurons of the anterior POA.

**aPOA$^{Vglut2}$ neuronal activity is necessary and sufficient for deep torpor entry**

Employing our newly developed Samd3E-AAV, we set out to address the role of

aPOA$^{Vglut2}$ neuronal activity in hibernation and DT by expressing an inhibitory Gi-

DREADD (AAV-Samd3E-hM4D(Gi)-EGFP) in hamster aPOA$^{Vglut2}$ neurons (Fig. 5a).

Neuronal inhibition via CLZ administration had no significant effect on basal homeostatic

$T_b$ or circadian $T_b$ rhythmicity in control non-hibernating Gi-DREADD-expressing

hamsters (Extended Data Fig. 11a, b). However, actively hibernating Gi-DREADD-

expressing (n = 12) animals showed a significant delay in the resumption of DT bouts

following CLZ treatment compared to when these same animals were administered a

water-only vehicle control (114.9 ± 45.4 hours vs. 64.1 ± 24.5 hours, P = 0.0003). Indeed,

a majority (7/12) of CLZ-treated Gi-DREADD-expressing animals failed to re-enter DT

during the full 6-day course of CLZ treatment, whereas none (0/12) of the animals failed

to resume DT during treatment with the vehicle control. Moreover, animals (n = 5)

expressing a control reporter payload (AAV-Samd3E-EGFP-Synaptophysin-mRuby)

exhibited no significant difference in the time to DT resumption following CLZ or vehicle

treatment (61.6 ± 26.5 hours vs. 47.4 ± 10.2 hours, respectively. P = 0.6167), with all

animals resuming torpor bouts within the CLZ treatment window (Fig. 5b, c). Together,

these results strongly suggest that aPOA$^{Vglut2}$ neuronal activity is required for entry into

hibernation-associated DT.

To determine the extent to which aPOA$^{Vglut2}$ neuron activity is sufficient to drive DT

entry, we selectively introduced an excitatory Gq-DREADD into hamster aPOA$^{Vglut2}$

neurons via our Samd3E-AAV (AAV-Samd3E-hM3D(Gq)-EGFP, Fig. 5d). CLZ

administration led to a rapid, significant drop in $T_b$ in Gq-DREADD-expressing hamsters,

with the animals reaching a minimum body temperature of 13.9 ± 1.0ºC (P < 0.0001

relative to normothermia) and maintaining a reduced body temperature for 56.9 ± 13.1

hours, consistent with the induction of a prolonged torpor-like state (Fig. 5e-g). By

contrast, these same animals exhibited no discernable $T_b$ change upon control PBS

administration (minimal $T_b$ = 33.3 ± 2.4ºC, P = 0.8959 relative to normothermia). In

addition, CLZ treatment of Gq-DREADD-expressing animals (n = 6), but not PBS

treatment of the same animals (n = 6) or PBS/CLZ treatment of hamsters injected with a

control viral construct (n = 3), induced behaviors consonant with torpor entry, including a

curled posture and prolonged inactivity in the nest (P = 0.0168, Fig. 5h, i, Extended Data

Fig. 12, Supplementary Video 1). Moreover, follow-up analyses revealed that the depth

of CLZ-induced torpor across animals correlated with levels of aPOA Gq-DREADD

expression, suggesting that more extensive aPOA$^{Vglut2}$ neuron activation drives deeper

torpor-like bouts (r = -0.9577, P = 0.0104, Extended Data Fig. 11c, d). Together, these

findings show that aPOA$^{Vglut2}$ neuronal activity is both necessary and sufficient to initiate

a profound $T_b$ decrease and DT-associated nesting behaviors in the hamster.

Finally, given that aPOA$^{Vglut2}$ neurons are present in both the Syrian hamster (a

hibernator) and mouse (non-hibernator), we investigated whether virus-mediated

chemogenetic activation of this neuronal population in mice housed at low ambient

temperatures would drive a DT-like decrease in $T_b$. Indeed, CLZ treatment of Samd3E-

Gq-DREADD-transduced mice yielded a rapid and sustained drop in body temperature,

although markedly less than that observed in hamsters (Extended Data Fig. 13). Thus,

aPOA$^{Vglut2}$ neuronal activation is sufficient to elicit acute, pronounced reductions in core

temperature in both hibernating and non-hibernating species, with the difference in

magnitude suggestive of potential hibernator-specific evolutionary modifications in

downstream effector circuitry.

**Discussion**

Hibernation is a conserved, recurrently evolved energy-conserving strategy across the mammalian lineage[5,6]. Entry into this state is orchestrated by a complex interplay of environmental cues and endocrine signals[69]. However, despite prior reports of hibernation-associated neural activity changes and classic lesion studies implicating the POA[41,43–45], the specific neuronal populations regulating entry into hibernation-associated deep torpor remained unknown. Here, using a Syrian hamster model, we combine neural activity mapping with chemogenetic silencing to show that aPOA activity is essential for deep torpor entry.

Our single-nucleus transcriptional and epigenomic profiling of the hamster POA revealed strong cellular homology with the mouse, with no evidence of distinct cell populations. Instead, we observed that aPOA glutamatergic neurons homologous to FIT-regulating neuronal cell types in the mouse exhibit elevated activity upon hibernation-associated DT in the hamster. Through the development and use of a novel enhancer-driven AAV we show that aPOA[Vglut2] neuron silencing blocks hibernation-associated DT entry, whereas chemogenetic activation of this population drives a rapid fall in body temperature and hibernation-like nesting behavior. Together, these studies establish Samd3E-AAV-targeted aPOA[Vglut2] neurons as key regulators of hibernation and provide the first genetic access to hibernation-regulating circuits. Moreover, this Samd3E-AAV, as well as the broader enhancer-driven viral targeting strategy, should be broadly applicable to target these and related cell populations in other hibernating species.

Although mouse FIT and seasonal hibernation both involve profound reductions in $T_b$, metabolic rate, and heart rate, these states differ markedly in their depth, duration, and

ethological significance. FIT represents a short-term response to food scarcity, lasting

only hours, whereas hibernation is typically a seasonal behavior involving deeper, multi-

day torpor bouts punctuated by brief periods of interbout euthermia. Our findings raise

the possibility that these differences reflect distinct connectivity and/or activity patterns of

aPOA$^{Vglut2}$ neurons in the two species, which might derive in part from remaining cross-

species gene expression differences observed in these cell types (Extended Data Fig.

7d). Moreover, chemogenetic activation of the aPOA$^{Vglut2}$ neurons in hamsters drives

substantially deeper torpor bouts than in mice, which might reflect limited efficiency in

fully targeting the aPOA$^{Vglut2}$ neuronal population (Extended Data Fig. 10) or alternatively

might indicate a requirement for additional circuit elements for the full expression of deep

torpor in the hamster. Similarly, in the future it will be important to determine whether this

aPOA$^{Vglut2}$ neuronal population, or aspects of the broader hibernation-regulating circuitry,

undergoes active remodeling during the extended pre-hibernation conditioning period to

support prolonged bouts of deep torpor.

More broadly, our findings in hamster and mouse further support for the idea that

regulated heterothermy is an ancestral trait in mammals[70,71] and that core elements of

the underlying torpor circuitry may be broadly conserved—potentially not only across

torpid and hibernating mammals, but also in species that do not naturally exhibit these

states. Indeed, *in situ* hybridization data from the marmoset (*Callithrix jacchus*), a non-

hibernating primate, indicate that neurons expressing deep torpor-associated markers

are present in the homologous regions within the aPOA (Extended Data Fig. 14)[72–74].

Our study thus lays a foundation for future efforts to define broader hibernation-

regulating circuitry, including upstream inputs conveying seasonal cues, downstream

pathways mediating systemic adaptations, and the cellular mechanisms that allow

neurons to function at near-freezing temperatures. Beyond advancing basic

understanding of hibernation, these insights may eventually inform approaches to

harness the mechanisms behind this amazing adaption to advance medicine in areas

such as therapeutic hypothermia, organ preservation, central metabolic control,

suppression of muscle atrophy, and aging.

**Methods**

**Animals**

Animal experiments were approved by the Massachusetts Institute of Technology Committee of Animal Care (Protocol 2503000788), following ethical guidelines described in the US National Institutes of Health Guide for the Care and Use of Laboratory Animals and the Animal Welfare Act.

**Hamster.** For all experiments, unless otherwise noted, adult female (5-6-week-old) Syrian hamsters (Inotiv, Strain HsdHan:AURA, Stock 089) were used. Hamsters were singly caged and initially housed at 22°C under a standard 12-hour light/dark cycle (12L:12D) cycle in temperature- and humidity-controlled rooms and had *ad libitum* access to food and water.

**Mouse.** Adult female (8-week-old) C57BL/6J (JAX #000664) mice were used. All mice were group housed initially at 22°C under a standard 12L:12D cycle in temperature- and humidity-controlled rooms and had *ad libitum* access to food and water.

No statistical methods were used to predetermine the sample size. Hamsters were randomly assigned to experimental groups before surgery. Where possible, investigators were blinded during analysis.

**Hibernation Induction**

Utilizing a modified hibernation paradigm[15,75] female (5-6-week-old) hamsters were singly caged and housed at 22°C under a standard 12L:12D cycle for a short period to acclimate to the animal facility. Upon completion of acclimation, hamsters were transferred to a reduced 8L:16D cycle with an ambient temperature of 22°C. After 7-8

weeks, hamsters were placed in temperature-controlled chambers with 8L:16D cycle and

ambient temperatures of 4-5°C. Initiation of hibernation was confirmed by changes in $T_b$

measured by telemetric temperature probes. In all cases, hamsters had *ad libitum* access

to food and water.

**Telemetric monitoring of core body temperature**

Within the 7-8-week period of housing at 22°C in 8L:16D cycle, hamsters were

anesthetized with 3% isoflurane and implanted with telemetric/logger temperature probes

(Star-Oddi, DST microRF-T) in the abdominal cavity. Hamsters were allowed to recover

for a minimum of three days before additional procedures or experiments were performed.

After recovery and at the end of the 7-8-week period, hamsters were moved to

temperature-controlled chambers, and core body temperature was monitored in home

cages placed onto a radio frequency receiver antenna (Star-Oddi, Antenna & RF-box).

Upon euthanasia, probes were removed and complete logged data downloaded. Core

body temperature was logged every 20-30 min.

**Torpor bout analysis**

**Classification.** To classify torpor bouts, we employed a Gaussian Mixture Model

in R. The classification was based on minimum body temperature and bout duration. Each

bout was assigned to the cluster in which it had the highest posterior probability of

membership. A cluster characterized by lower $T_b$ and longer duration was designated as

"Deep", whereas the other cluster was classified as "Shallow". Furthermore, periods of

between bouts above the set $T_b$ threshold were classified as IBE.

**Circadian preference.** In R, circadian preference for torpor and IBE initiation was

quantified using the Raleigh test of circular uniformity.

**Exponential decay.** In R, elapsed time in torpor for each body temperature

reading from torpor bouts classified as deep were determined for the first 30 hours. Next,

using the mean of temperatures at each time across animals, a nonlinear model was fit

to the data using the minimum temperature within the 30-hour period as a fixed asymptotic

value.

**Viral constructs**

AAV8-hSyn-hM4D(Gi)-mCherry (Addgene, 50475-AAV8) was obtained from

Addgene.　　AAV8-Samd3E-dTomato-nlsdTomato,　　AAV8-Samd3E-hM4D(Gi)-P2A-

nlsGFP,　　AAV8-Samd3E-EGFP-P2A-Synaptophysin-mRuby,　　and　　AAV8-Samd3E-

hM3D(Gq)-P2A-nlsGFP　were　produced　in-house　using　the　method　previously

described[76] and titered via digital PCR (QIAGEN, QIAcuity). All viruses were diluted with

PBS to a final titer of 1E13 vg/mL before stereotaxic delivery into the hamster brain. For

mice, AAV8-Samd3E-hM3D(Gq)-P2A-nlsGFP with a final titer of 5E13 vg/mL was used.

**Stereotaxic viral injection**

**Hamster.** Hamsters were anaesthetized with 3% isoflurane and placed in a

stereotaxic head frame (David Kopf Instruments, model 1900). If hamsters were actively

hibernating, they were aroused from hibernation and placed at 22°C the day prior to

surgery.  An air-based injection system connected to a pulled glass needle was used to

infuse ~100 nL of virus bilaterally at regions of interest. Hypothalamic regions were

targeted by referencing the hamster stereotaxic atlas[77]. When targeting the aPOA,

coordinates +1.2 mm AP, ±0.4 mm ML, -7.5mm DV were used. Hamsters were allowed

to recover at 22°C for a minimum of 3 days before a return to 4°C chambers. Animals

were afforded a recovery period of at least two weeks post-surgery prior to further

experimental manipulation to allow for viral expression and, if appropriate, resumption of

hibernation.

**Mouse.** Mice were anesthetized with 3% isoflurane and placed in a stereotaxic

head frame (RWD, Guangdong, China). A Nanoject III (Drummond Scientific, Broomall,

PA) system connected to a pulled glass needle was used to infuse ~100 nL of virus

bilaterally into the aPOA (distance from Bregma, AP: +0.6 mm, ML: ±0.4 mm, DV: -5.2

mm). Mice were monitored for three days post-surgery to ensure full recovery, with a

waiting period of at least two weeks to allow sufficient viral expression.

**Chemogenetic manipulations**

**Hamster.** Clozapine dihydrochloride (Hello Bio, HB6129) was either dissolved in

sterile water or PBS at a concentration of 10 mg/mL and then stored at -20°C until use.

For administration via drinking water, CLZ dissolved in sterile water was diluted in drinking

water at a dose of 0.1 mL/kg (1 mg/kg) in a final volume of 250 mL. Animals were

maintained on CLZ water for 3-7 days. For intraperitoneal (I.P.) administration, CLZ

dissolved in PBS was administered at a dose of 0.5 mL/kg (5 mg/kg). In each case when

treating actively hibernating hamsters, they were aroused from hibernation the morning

of each experiment and treatment was administered in the evening.

**Mouse.** Mice injected with AAV8-Samd3E-hM3D(Gq)-P2A-nlsGFP were single

housed and placed in a temperature-controlled chamber at 8°C for three days. After three

days, mice were I.P. injected with CLZ at a dose of 5 mg/kg and surface body temperature

was measured using a thermal camera (Teledyne FLIR C5, Teledyne Flir, Wilsonville,

OR). Images were taken before CLZ injection, and 30 min, 60 min, 3 hours, 6 hours and

24 hours after CLZ injection. Analysis of surface body temperature and representative

images were generated using the Flir Ignite software (https://ignite.flir.com/).

**Immunofluorescence**

Hamsters were deeply anesthetized with isoflurane and then euthanized by

transcardial perfusion of 50 mL ice-cold 1X PBS followed by 50 mL of 4%

paraformaldehyde (PFA). Brains were extracted, post-fixed overnight with 4% PFA at

4°C, and then embedded in 1X PBS with 3% agarose. Brains were sliced on a vibratome

into 50 μm coronal sections and stored at -20°C in cryoprotectant buffer (30% sucrose,

1% polyvinyl-pyrrolidone (PVP-40), 30% ethylene glycol, in 0.1 M phosphate buffer).

Free-floating coronal sections were washed three times with 1X PBS and then blocked

for 1 hour at room temperature with 1X PBS containing 10% donkey serum and 0.3%

Triton X-100. Sections were incubated overnight at 4°C with primary antibodies (Rabbit

anti-FOS, Synaptic Systems, 226-008; Rabbit anti-RFP, Rockland, 600-401-379; Chicken

anti-GFP, Aves Labs, GFP-1020) diluted in 1X PBS containing 5% donkey serum and

0.3% Triton X-100, washed again three times with 1X PBS, and incubated for 2 hours at

room temperature with secondary antibodies in 1X PBS containing 5% donkey serum and

0.3% Triton X-100. After washing three times in 1X PBS, samples were counter-stained

for 15 minutes at room temperature with 1 µg/mL DAPI, washed again three times in 1X

PBS, and then mounted onto SuperFrost Plus glass slides and coverslipped using

Fluoromount-G.

**Determining deep torpor-active brain regions**

Brains were collected in torpor, IBE, and from non-hibernating control animals and

were processed as indicated in the section 'Immunofluorescence' using the following

reagent for the detection of FOS: Rabbit anti-FOS, Synaptic Systems, 226-008.

**Sample imaging.** Sections were imaged with a TissueFAXS SL whole-slide

scanning system equipped with a Zeiss Axio Imager Z2 fluorescence microscope and

TissueFAXS image acquisition software (Ragon Institute Microscopy Core).

**Image analysis**. Using TissueQuest analysis software (TissueGnostics GmbH,

Vienna, Austria), brain regions were manually annotated and nuclear DAPI signal

segmented. Segmented cells containing putative FOS signal were determined to be FOS-

positive based on diameter and mean fluorescence intensity. In R, regional cell counts

were analyzed and percent active neurons determined by the fraction of FOS-positive

and total DAPI-positive cells. Significance was calculated using a mixed-effects ANOVA

with post-hoc pairwise comparisons. Pairwise comparison p-values were multiple tests

corrected using the Bonferonni method.

**Mapping of virally injected hypothalamic nuclei and correlation with**

**chemogenetically induced torpor perturbation**

Hibernating hamsters were bilaterally injected with AAV8-hSyn-hM4D(Gi)-

mCherry. Upon resumption of hibernation, hamsters were administered CLZ water for ~3

days (74 hours) to inhibit virally injected regions, and their core body temperature were

recorded.

**Temperature data analysis.** Upon CLZ water treatment, hamsters reinitiating

deep torpor within 48 hours were classified as unperturbed, and those failing to reinitiate

(remaining euthermic) within this time were classified as perturbed. Sham surgery control

animals treated with CLZ water for 7 days informed this threshold (Extended Data Fig.

2c). Absolute elapsed time of euthermia from start of treatment was used for further

analysis.

**Sample imaging.** Given the unavoidable variability in hypothalamic region

targeting and viral spread, all hamsters were euthanized for post hoc assessment of

transduced brain regions. Collected brains were processed as indicated in the section

'Immunofluorescence' to detect endogenous mCherry fluorescence. Sections were

imaged with a TissueFAXS SL whole-slide scanning system equipped with a Zeiss Axio

Imager Z2 fluorescence microscope and TissueFAXS image acquisition software (Ragon

Institute Microscopy Core).

**Image and correlation analysis.** Viral mCherry expression was quantified across

70 hypothalamic regions in a semi quantitative manner, where the viral expression was

scored 0 (none), 1 (minimal), 2 (partial), or 3 (complete) in each region and hemisphere.

The analysis was performed blinded to whether perturbation of torpor was observed in

each hamster. To assign a single numeric value for each hypothalamic region in each

hamster, we added the semiquantitative expression values from each hemisphere. Next,

a Pearson correlation was calculated for each region across all hamsters between the

viral expression in that region and the length of time spent euthermic.

**Nuclei dissociation and single-cell analysis**

Preoptic area tissue was dissected from adult male and female Syrian hamsters

and immediately flash-frozen. The 10X genomics nuclei isolation kit (1000448) was used

to obtain dissociated nuclei. Tissue was homogenized on ice in a detergent-based nuclei

isolation buffer, passed through a 40-μm filter, and purified by centrifugation. Nuclei were

washed, and counted before loading. Single-nucleus multiome libraries were prepared

using the 10x Genomics Chromium Single Cell Multiome ATAC + Gene Expression

workflow following the manufacturer's protocol. Constructed libraries were quantified,

pooled, and sequenced on an Illumina platform to recommended depth for both

modalities. The raw data were processed in parallel with cellranger-arc to generate gene-

expression count matrices, ATAC fragment files, and peak-by-cell accessibility matrices.

A recently published mouse POA multiome dataset generated with the same 10x system

was obtained to build a reference dataset. A combination of scVI and Scanpy[78,79]

pipelines were used to process and integrate the two datasets, including quality control

(filtering by gene counts, UMI counts, and mitochondrial RNA content), normalization, log-

transformation, identification of highly variable genes, dimensionality reduction,

neighborhood graph construction, and Leiden clustering.

To ensure that downstream analyses focused on POA-associated neuronal

populations, we used the MERFISH dataset as a spatial reference[61]. To this end, mouse

spatial transcriptomic data was subset to only include neurons found in the mPOA and

co-clustered with mouse neuronal snRNA-seq. Canonical POA markers and spatial

annotations were used to confirm that the identified clusters in both species corresponded

to POA-localized neuronal subtypes. Subsequently, the reference mouse dataset was

used to identify homologous hamster neurons through integration and label transferring

with scANVI[80]. Chromatin accessibility matrices were analyzed with Pycistopic[81] including

psuedobulking, peak calling and filtering, identification of co-accessible regions and

differential accessibility analysis.

**Double fluorescent *in situ* hybridization and immunofluorescence**

Hamsters were deeply anesthetized with isoflurane and then euthanized by

transcardial perfusion of 50 mL ice-cold 1X PBS followed by 50 mL of 4% PFA. Brains

were extracted, post-fixed overnight with 4% PFA at 4°C, and then incubated in 1X PBS

with 30% sucrose at 4°C until fully infiltrated and sunk to the bottom of the chamber (72-

96 hours). Brains were embedded in Optimal Cutting Temperature (O.C.T.) compound,

frozen on dry ice, and stored at -80°C until use. Brains were sliced on a cryostat (Leica

CM3050 S) into 10-30 µm coronal sections, placed in cryoprotectant buffer, and stored at

−20°C until use. Free-floating coronal sections were washed three times with 1X PBS,

mounted onto Superfrost Plus slides, and then baked for 45 minutes at 60°C. Subsequent

steps followed the standard fluorescent *in situ* hybridization protocol for the RNAscope

Fluorescent Multiplex V2 Assay using the following probes: Mau-*Samd3*-C1 (1826261),

Mau-*Slc17a6*-C2 (1176381), Mau-*Adcyap1*-O1-C3 (1808861), Mau-*Slc32a1*-C4

(1176381).

Upon completion of *in situ* hybridization, samples immediately were processed

according to the ACD RNAscope Fluorescent Multiplex V2 Assay Combined with

Immunofluorescence manual with the following modifications: Slide-mounted sections

were washed twice in 1X TBS with 0.005% Tween-20 (TBST), blocked for 1 hour at room

temperature with 1X TBS containing 10% donkey serum, and incubated overnight at 4°C

with primary antibodies (Rabbit anti-FOS, Synaptic Systems, 226-008; Rabbit anti-RFP,

Rockland, 600-401-379; Chicken anti-GFP, Aves Labs, GFP-1020) diluted in 1X TBS

containing 5% donkey serum. Slides were washed three times with TBST, incubated for

2 hours at room temperature with secondary antibodies in 1X TBS containing 5% donkey

serum, washed again three times with TBST, counter-stained for 15 minutes at room

temperature with 1 µg/mL DAPI, washed three times with 1X TBS, and finally mounted

on coverslips using ProLong Gold.

**Sample imaging.** Sections were either imaged with a TissueFAXS SL whole-slide

scanning system equipped with a Zeiss Axio Imager Z2 fluorescence microscope and

TissueFAXS image acquisition software (Koch Institute Swanson Biotechnology

Center/Microscopy Core) or a Leica Stellaris 8 FALCON confocal microscope (Whitehead

Institute W.M. Keck Facility for Biological Imaging).

**Image analysis**. Using QuPath software[82], colocalization analysis of RNAscope

and immunofluorescent signals was performed. After manual annotation of regions of

interest, the InstanSeg extension[83] was utilized for cell segmentation, considering each

fluorescent signal. Next, segmented cells containing putative RNAscope and

immunofluorescent signals were determined to be positive or negative for associated

markers based on diameter and mean fluorescence intensity. Subsequent quantification

of counts was performed in R, while plotting and statistical analysis was performed in

GraphPad Prism software (Boston, MA).

**Inhibition of aPOA$^{Vglut2}$ neurons**

Hibernating hamsters were bilaterally injected with either AAV8-Samd3E-

hM4D(Gi)-P2A-nlsGFP or AAV8-Samd3E-EGFP-P2A-Synaptophysin-mRuby. On the

day of treatment, hamsters were aroused in the morning, I.P. injected with CLZ and

provided with CLZ water in the evening, I.P. injected again 16 hours later, and core body

temperature recorded. This regimen accounted for variability in water intake within the

first 24 hours of the experiment and maximized inhibition of our targeted population. A

CLZ water treatment window of 6 days (144 hours) was used. For PBS control treatment,

an identical regimen was followed.

**Temperature data analysis.** In GraphPad Prism (Boston, MA), absolute elapsed

time of euthermia from start of treatment was used to determine differences within groups.

The treatment window duration (144 hours) was applied as an upper bound for statistical

analysis. In R, temperature data from each virus group and treatment condition was

mapped to circadian time. In GraphPad Prism (Boston, MA), statistical analysis was

performed on temperatures in the light and dark cycles within virus groups.

**Excitation of aPOA$^{Vglut2}$ neurons**

Hamsters were bilaterally injected with either AAV8-Samd3E-hM3D(Gq)-P2A-

nlsGFP or AAV8-Samd3E-EGFP-P2A-Synaptophysin-mRuby and housed at 4°C to

acclimate. On the day of treatment, hamsters were aroused in the morning, I.P. injected

with CLZ in the evening, and core body temperature recorded. For PBS control treatment,

an identical regimen was followed.

**Temperature data analysis.** In R, temperature data was analyzed using the

method described in section 'Torpor bout analysis' to determine if the CLZ induced drop

in body temperature clustered with either deep or shallow torpor. In GraphPad Prism

(Boston, MA), statistical analysis was performed on minimum temperatures across each

condition (euthermia, torpor, CLZ, and PBS).

**Sample imaging.** Collected brains were processed as indicated in the section

'Immunofluorescence' using the following reagent for the detection of AAV8-Samd3E-

hM3D(Gq)-P2A-nlsGFP: Chicken anti-GFP, Aves Labs, GFP-1020. Sections were

imaged with a TissueFAXS SL whole-slide scanning system equipped with a Zeiss Axio

Imager Z2 fluorescence microscope and TissueFAXS image acquisition software (Koch

Institute Swanson Biotechnology Center/Microscopy Core).

**Image and correlation analysis.** aPOA GFP expression was quantified in a semi

quantitative manner where expression was scored 0 (none), 1 (minimal), 2 (partial), or 3

(complete) in each hemisphere. The analysis was performed blinded to change in body

temperature observed in each hamster. Hemispheric values were summed within

animals. In GraphPad Prism (Boston, MA), a Pearson correlation was performed between

GFP viral expression and the minimum body temperature.

**Behavioral analysis**

Hamsters housed at 4°C and injected with AAV8-Samd3E-hM3D(Gq)-P2A-

nlsGFP or AAV8-Samd3E-EGFP-P2A-Synaptophysin-mRuby were I.P. injected with PBS

(0.5 mL/kg) after a one-hour baseline video recording of homecage behavior and an

additional hour recording post injection. Following at least a week recovery in 4°C

chambers, animals were injected with CLZ (5 mg/kg) and videos were recorded for one

hour before and post drug administration in 4°C chambers. Homecage behaviors were

manually coded using BORIS software[84]. Behavior was coded 30 min after PBS or CLZ

injection over a 10 min window.

**References**

1.  Tan, C. L. & Knight, Z. A. Regulation of Body Temperature by the Nervous System. *Neuron* **98**, 31–48 (2018).

2.  Morrison, S. F. & Nakamura, K. Central Mechanisms for Thermoregulation. *Annu. Rev. Physiol.* **81**, 285–308 (2019).

3.  Heller, H. C. & Hammel, H. T. CNS control of body temperature during hibernation. *Comp. Biochem. Physiol. A Physiol.* **41**, 349–359 (1972).

4.  Sunagawa, G. A. & Takahashi, M. Hypometabolism during Daily Torpor in Mice is Dominated by Reduction in the Sensitivity of the Thermoregulatory System. *Sci. Rep.* **6**, 1–14 (2016).

5.  Geiser, F. Metabolic Rate and Body Temperature Reduction During Hibernation and Daily Torpor. *Annu. Rev. Physiol.* **66**, 239–274 (2004).

6.  Geiser, F. *Ecological Physiology of Daily Torpor and Hibernation*. (Springer, Cham, Switzerland, 2021).

7.  Drew, K. L. *et al.* Central nervous system regulation of mammalian hibernation: implications for metabolic suppression and ischemia tolerance. *J Neurochem* **102**, 1713–1726 (2007).

8.  Heller, H. C. Hibernation: neural aspects. *Annu. Rev. Physiol.* **41**, 305–321 (1979).

9.  Morton, S. R. Torpor and Nest-Sharing in Free-Living Sminthopsis crassicaudata (Marsupialia) and Mus musculus (Rodentia). *J. Mammal.* **59**, 569–575 (1978).

10. Swoap, S. J. & Gutilla, M. J. Cardiovascular changes during daily torpor in the laboratory mouse. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.* **297**, R769–R774 (2009).

11. Barnes, B. M. Freeze avoidance in a mammal: body temperatures below 0 degree C

in an Arctic hibernator. *Science* **244**, 1593–1595 (1989).

12. Heldmaier, G., Ortmann, S. & Elvert, R. Natural hypometabolism during hibernation

and daily torpor in mammals. *Respir. Physiol. Neurobiol.* **141**, 317–329 (2004).

13. Revel, F. G. *et al.* The circadian clock stops ticking during deep hibernation in the

European hamster. *Proc. Natl. Acad. Sci.* **104**, 13816–13820 (2007).

14. Mohr, S. M., Bagriantsev, S. N. & Gracheva, E. O. Cellular, Molecular, and

Physiological Adaptations of Hibernation: The Solution to Environmental

Challenges. *Annu. Rev. Cell Dev. Biol.* **36**, 315–338 (2020).

15. Chayama, Y., Ando, L., Tamura, Y., Miura, M. & Yamaguchi, Y. Decreases in body

temperature and body mass constitute pre-hibernation remodelling in the Syrian

golden hamster, a facultative mammalian hibernator. *R. Soc. Open Sci.* **3**, 160002

(2016).

16. Teague, R. S. & Ranson, S. W. The role of the anterior hypothalamus in

temperature regulation. *Am. J. Physiol.-Leg. Content* **117**, 562–570 (1936).

17. Hammel, H. T. & Pierce, J. B. Regulation of Internal Body Temperature. *Annu. Rev.*

*Physiol.* **30**, 641–710 (1968).

18. Zaretskaia, M. V., Zaretsky, D. V., Shekhar, A. & DiMicco, J. A. Chemical stimulation

of the dorsomedial hypothalamus evokes non-shivering thermogenesis in

anesthetized rats. *Brain Res.* **928**, 113–125 (2002).

19. Nakamura, K. & Morrison, S. F. A thermosensory pathway that controls body

temperature. *Nat Neurosci* **11**, 62–71 (2008).

20. Hensel, H. Neural processes in thermoregulation. *Physiol. Rev.* **53**, 948–1017 (1973).

21. Jessen, C. Thermal afferents in the control of body temperature. *Pharmacol. Ther.* **28**, 107–134 (1985).

22. Hammel, H. T., Hardy, J. D. & Fusco, M. M. Thermoregulatory responses to hypothalamic cooling in unanesthetized dogs. *Am J Physiol* **198**, 481–6 (1960).

23. Magoun, H. W., Harrison, F., Brobeck, J. R. & Ranson, S. W. Activation of heat loss mechanisms by local heating of the brain. *J. Neurophysiol.* **1**, 101–114 (1938).

24. Tan, C. L. *et al.* Warm-Sensitive Neurons that Control Body Temperature. *Cell* **167**, 47-59.e15 (2016).

25. Song, K. *et al.* The TRPM2 channel is a hypothalamic heat sensor that limits fever and can drive hypothermia. *Science* **353**, 1393–1398 (2016).

26. Yu, S. *et al.* Glutamatergic Preoptic Area Neurons That Express Leptin Receptors Drive Temperature-Dependent Body Weight Homeostasis. *J Neurosci* **36**, 5034–46 (2016).

27. Zhou, Q. *et al.* Hypothalamic warm-sensitive neurons require TRPC4 channel for detecting internal warmth and regulating body temperature in mice. *Neuron* **111**, 387-404 e8 (2023).

28. Vitale, P. M., Darrow, J. M., Duncan, M. J., Shustak, C. A. & Goldman, B. D. Effects of photoperiod, pinealectomy and castration on body weight and daily torpor in Djungarian hamsters (Phodopus sungorus). *J. Endocrinol.* **106**, 367–375 (1985).

29. Paul, M. J., Kauffman, A. S. & Zucker, I. Feeding schedule controls circadian timing

of daily torpor in SCN-ablated Siberian hamsters. *J. Biol. Rhythms* **19**, 226–237

(2004).

30. Magariños, A. M., McEwen, B. S., Saboureau, M. & Pevet, P. Rapid and reversible

changes in intrahippocampal connectivity during the course of hibernation in

European hamsters. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 18775–18780 (2006).

31. Korhonen, T., Mustonen, A.-M., Nieminen, P. & Saarela, S. Effects of cold exposure,

exogenous melatonin and short-day treatment on the weight-regulation and body

temperature of the Siberian hamster (Phodopus sungorus). *Regul. Pept.* **149**, 60–66

(2008).

32. Stanton, T. L., Daley, J. C. & Salzman, S. K. Prolongation of hibernation bout

duration by continuous intracerebroventricular infusion of melatonin in hibernating

ground squirrels. *Brain Res.* **413**, 350–355 (1987).

33. Chmura, H. E. *et al.* Hypothalamic remodeling of thyroid hormone signaling during

hibernation in the arctic ground squirrel. *Commun. Biol.* **5**, 492 (2022).

34. Tamura, Y., Shintani, M., Nakamura, A., Monden, M. & Shiomi, H. Phase-specific

central regulatory systems of hibernation in Syrian hamsters. *Brain Res.* **1045**, 88–

96 (2005).

35. Olson, J. M. *et al.* Circannual rhythm in body temperature, torpor, and sensitivity to

$A_1$ adenosine receptor agonist in arctic ground squirrels. *J. Biol. Rhythms* **28**, 201–

207 (2013).

36. Jinka, T. R., Tøien, Ø. & Drew, K. L. Season primes the brain in an arctic hibernator

to facilitate entrance into torpor mediated by adenosine A(1) receptors. *J. Neurosci.*

*Off. J. Soc. Neurosci.* **31**, 10752–10758 (2011).

37. Frare, C., Jenkins, M. E., Soldin, S. J. & Drew, K. L. The Raphe Pallidus and the

Hypothalamic-Pituitary-Thyroid Axis Gate Seasonal Changes in Thermoregulation in

the Hibernating Arctic Ground Squirrel (Urocitellus parryii). *Front. Physiol.* **9**, 1747

(2018).

38. Stanton, T. L., Winokur, A. & Beckman, A. L. Seasonal variation in thyrotropin-

releasing hormone (TRH) content of different brain regions and the pineal in the

mammalian hibernator, Citellus lateralis. *Regul. Pept.* **3**, 135–144 (1982).

39. Mohr, S. M. *et al.* Hypothalamic hormone deficiency enables physiological anorexia

in ground squirrels during hibernation. *Nat. Commun.* **15**, 5803 (2024).

40. Harris, M. B. & Milsom, W. K. Parasympathetic influence on heart rate in euthermic

and hibernating ground squirrels. *J. Exp. Biol.* **198**, 931–937 (1995).

41. Markussen, F. A. F., Cazarez-Marquez, F., Melum, V. J., Hazlerigg, D. G. & Wood,

S. H. c-fos induction in the choroid plexus, tanycytes and pars tuberalis is an early

indicator of spontaneous arousal from torpor in a deep hibernator. *J Exp Biol* **227**,

(2024).

42. Enoki, R. *et al.* Cold-induced suspension and resetting of Ca2+ and transcriptional

rhythms in the suprachiasmatic nucleus neurons. *iScience* **26**, 108390 (2023).

43. Cheng, M. *et al.* Brainstem catecholaminergic neurons induce torpor during fasting

by orchestrating cardiovascular and thermoregulation changes. *Nat. Commun.* **16**,

5954 (2025).

44. Bratincsak, A. *et al.* Spatial and temporal activation of brain regions in hibernation: c-

fos expression during the hibernation bout in thirteen-lined ground squirrel. *J Comp*

*Neurol* **505**, 443–58 (2007).

45. Weidler, D. J., Earle, A. M., Myers, G. G. & Sieck, G. C. Effect of hypothalamic

lesions on temperature regulation in hibernating ground squirrels. *Brain Res.* **65**,

175–179 (1974).

46. Hastings, M. H., Maywood, E. S. & Brancaccio, M. Generation of circadian rhythms

in the suprachiasmatic nucleus. *Nat. Rev. Neurosci.* **19**, 453–469 (2018).

47. Welsh, D. K., Takahashi, J. S. & Kay, S. A. Suprachiasmatic Nucleus: Cell

Autonomy and Network Properties. *Annu. Rev. Physiol.* **72**, 551–577 (2010).

48. Weaver, D. R. The Suprachiasmatic Nucleus: A 25-Year Retrospective. *J. Biol.*

*Rhythms* **13**, 100–112 (1998).

49. Andermann, M. L. & Lowell, B. B. Toward a Wiring Diagram Understanding of

Appetite Control. *Neuron* **95**, 757–778 (2017).

50. Cone, R. D. *et al.* The arcuate nucleus as a conduit for diverse signals relevant to

energy homeostasis. *Int. J. Obes.* **25**, S63–S67 (2001).

51. Jais, A. & Brüning, J. C. Arcuate Nucleus-Dependent Regulation of Metabolism—

Pathways to Obesity and Diabetes Mellitus. *Endocr. Rev.* **43**, 314–328 (2022).

52. Van der Werf, Y. D., Witter, M. P. & Groenewegen, H. J. The intralaminar and

midline nuclei of the thalamus. Anatomical and functional evidence for participation

in processes of arousal and awareness. *Brain Res. Rev.* **39**, 107–140 (2002).

53. Jones, B. E. Arousal systems. *Front. Biosci. J. Virtual Libr.* **8**, s438-451 (2003).

54. Roth, B. L. DREADDs for Neuroscientists. *Neuron* **89**, 683–694 (2016).

55. Hrvatin, S. *et al.* Neurons that regulate mouse torpor. *Nature* **583**, 115–121 (2020).

56. Takahashi, T. M. *et al.* A discrete neuronal circuit induces a hibernation-like state in

rodents. *Nature* **583**, 109–114 (2020).

57. Zhang, Z. *et al.* Estrogen-sensitive medial preoptic area neurons coordinate torpor in

mice. *Nat. Commun.* **11**, 6378 (2020).

58. Machado, N. L. S. *et al.* Preoptic EP3R neurons constitute a two-way switch for

fever and torpor. *Nature* 1–10 (2025) doi:10.1038/s41586-025-09056-1.

59. Ambler, M., Hitrec, T., Wilson, A., Cerri, M. & Pickering, A. Neurons in the

Dorsomedial Hypothalamus Promote, Prolong, and Deepen Torpor in the Mouse. *J.*

*Neurosci. Off. J. Soc. Neurosci.* **42**, 4267–4277 (2022).

60. Kaplan, H. S. *et al.* Sensory input, sex and function shape hypothalamic cell type

development. *Nature* https://doi.org/10.1038/s41586-025-08603-0 (2025)

doi:10.1038/s41586-025-08603-0.

61. Zhang, M. *et al.* Molecularly defined and spatially resolved cell atlas of the whole

mouse brain. *Nature* **624**, 343–354 (2023).

62. Dimidschstein, J. *et al.* A viral strategy for targeting and manipulating interneurons

across vertebrate species. *Nat. Neurosci.* **19**, 1743–1749 (2016).

63. Cronin, T. *et al.* Efficient transduction and optogenetic stimulation of retinal bipolar

cells by a synthetic adeno-associated virus capsid and promoter. *EMBO Mol. Med.*

**6**, 1175–1190 (2014).

64. Ben-Simon, Y. *et al.* A suite of enhancer AAVs and transgenic mouse lines for

genetic access to cortical cell types. *Cell* **188**, 3045-3064 e23 (2025).

65. Graybuck, L. T. *et al.* Enhancer viruses for combinatorial cell-subclass-specific

labeling. *Neuron* **109**, 1449-1464.e13 (2021).

66. Hrvatin, S. *et al.* A scalable platform for the development of cell-type-specific viral

drivers. *eLife* **8**, e48089 (2019).

67. Nagy, M. A. *et al.* Cis-regulatory elements driving motor neuron-selective viral

payload expression within the mammalian spinal cord. *Proc. Natl. Acad. Sci.* **121**,

e2418024121 (2024).

68. Hunker, A. C. *et al.* Enhancer AAV toolbox for accessing and perturbing striatal cell

types and circuits. *Neuron* **113**, 1507-1524 e17 (2025).

69. Junkins, M. S., Bagriantsev, S. N. & Gracheva, E. O. Towards understanding the

neural origins of hibernation. *J. Exp. Biol.* **225**, jeb229542 (2022).

70. Lovegrove, B. G. A phenology of the evolution of endothermy in birds and mammals.

*Biol Rev Camb Philos Soc* **92**, 1213–1240 (2017).

71. Geiser, F., Stawski, C., Wacker, C. B. & Nowack, J. Phoenix from the Ashes: Fire,

Torpor, and the Evolution of Mammalian Endothermy. *Front Physiol* **8**, 842 (2017).

72. Kita, Y. *et al.* Cellular-resolution gene expression profiling in the neonatal marmoset

brain reveals dynamic species- and region-specific differences. *Proc. Natl. Acad.*

*Sci.* **118**, e2020125118 (2021).

73. Shimogori, T. *et al.* Digital gene atlas of neonate common marmoset brain.

*Neurosci. Res.* **128**, 1–13 (2018).

74. Gene Structural Search | Marmoset Gene Atlas. https://gene-

atlas.brainminds.jp/gene-structural/.

75. Trefna, M. *et al.* The influence of sex and diet on the characteristics of hibernation in

Syrian hamsters. *J. Comp. Physiol. [B]* **187**, 725–734 (2017).

76. Negrini, M., Wang, G., Heuer, A., Björklund, T. & Davidsson, M. AAV Production

Everywhere: A Simple, Fast, and Reliable Protocol for In-house AAV Vector

Production Based on Chloroform Extraction. *Curr. Protoc. Neurosci.* **93**, e103

(2020).

77. Wood, R. I. & Morin, L. P. *A Stereotaxic Atlas of the Golden Hamster Brain*.

(Elsevier Science, 2001).

78. Lopez, R., Regier, J., Cole, M. B., Jordan, M. I. & Yosef, N. Deep generative

modeling for single-cell transcriptomics. *Nat. Methods* **15**, 1053–1058 (2018).

79. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene

expression data analysis. *Genome Biol.* **19**, 15 (2018).

80. Xu, C. *et al.* Probabilistic harmonization and annotation of single-cell transcriptomics

data with deep generative models. *Mol. Syst. Biol.* **17**, e9620 (2021).

81. Bravo González-Blas, C. *et al.* cisTopic: cis-regulatory topic modeling on single-cell

ATAC-seq data. *Nat. Methods* **16**, 397–400 (2019).

82. Bankhead, P. *et al.* QuPath: Open source software for digital pathology image

analysis. *Sci. Rep.* **7**, 16878 (2017).

83. Goldsborough, T. *et al.* A novel channel invariant architecture for the segmentation

of cells and nuclei in multiplexed images using InstanSeg. 2024.09.04.611150

Preprint at https://doi.org/10.1101/2024.09.04.611150 (2024).

84. Friard, O. & Gamba, M. BORIS: a free, versatile open-source event-logging software

for video/audio coding and live observations. *Methods Ecol. Evol.* **7**, 1325–1330

(2016).

85. Lein, E. S. *et al.* Genome-wide atlas of gene expression in the adult mouse brain.

*Nature* **445**, 168–176 (2007).

**Acknowledgments:** We thank M. Alkire, M.M. Chen, J.A. Cheng, and N.T. Nigrin for their

assistance with mouse experiments. We also thank additional Hrvatin lab members for

their thoughtful input on this project and manuscript. The snRNA-seq and snATAC-seq

were performed with the Whitehead Institute Genome Technology Core. Imaging was

performed either at the Ragon Institute Microscopy Core, Koch Institute Swanson

Biotechnology Center/Microscopy Core, or the Whitehead Institute W.M. Keck Facility for

Biological Imaging. This work was supported by the Mathers Foundation Grant, Searle

Scholars Program, Pew Charitable Trust, Longevity Impetus Grant from Norn Group,

Hevolution Foundation and Rosenkranz Foundation, MIT Research Support Committee

Grant, and Howard Hughes Medical Institute Hrabowski Scholar Program.

**Author Contributions:** A.J.M. and S.H. conceptualized the study and designed

experiments. A.J.M, C.M.R, A.J.L.P, W.L., A.S.L. performed and analyzed experiments.

E.C.G. and S.H. advised on the study and obtained funding for the research. A.J.M.,

E.C.G., and S.H. wrote the manuscript.

**Competing Interests:** Authors declare that they have no competing interests.

**Data and materials availability:** Data will be made available on GEO. All other data

needed to evaluate the conclusions are available in the main text, the extended data,

supplementary materials, and available upon request.

**Fig. 1: A laboratory model of Syrian hamster hibernation.**

**a**, A laboratory paradigm for hibernation induction in the Syrian hamster, transitioning

animals to an altered light cycle and ambient temperature to simulate seasonal changes.

**b**, Representative body temperature trace of a hibernating hamster, consisting of repeated bouts of deep torpor (DT) and interbout euthermia (IBE) (left). Thermal images of a hibernating hamster during euthermic and deep torpor intervals (right).

**c**, Classification of torpor bouts as shallow (n = 1317) or deep (n = 2289) based on minimum temperature and bout duration.

**d,e**, Violin plots of minimum body temperature (**d**), and duration (**e**) of shallow and deep torpor bouts as well as IBE periods (n = 2471).

**f**, Percentage of animals undergoing DT, only shallow torpor, or no torpor bouts within 6 weeks of initiating the hibernation period (n = 394 animals).

**g**, Circadian representation (Raleigh plot) of the time of entry into shallow (light blue, n = 1312) or deep (dark blue, n = 2283) torpor bouts. Light and dark cycle phases are represented by yellow and grey, respectively. Red arrow direction and length indicate mean time of initiation and concentration around the mean, respectively. Red lines flanking the arrow indicate s.d.

**h**, Circadian representation as in (**g**) of entry into IBE following deep (left, n = 1496) or shallow (right, n = 1011) torpor bouts.

**Fig. 2: Identification and functional assessment of deep torpor-active brain regions.**

**a**, Strategy for identifying brain regions with hibernation state-dependent activity. Arrows

indicate relative sample collection points for torpor, IBE, and non-hibernating animals.

**b**,**c**, Hypothalamic and thalamic regions demonstrating an increase in FOS-positive cells

in torpor (n = 4 animals) as compared to both IBE (n = 3 animals) and non-hibernating (n

= 4 animals) controls (*P-adjusted < 0.05, **P-adjusted < 0.01, ***P-adjusted < 0.001).

**d**, FOS immunostaining in the hamster POA across hibernation states. AVPe, MPO, and

MnPO denoted by dashed lines.

**e**, Strategy for identifying hypothalamic subregions regulating deep torpor entry.

**f**, $T_b$ traces showing CLZ-mediated disruption of DT in one Gi-DREADD-mCherry-

transduced hamster (solid red) but not the other (dashed black).

**g**, Coronal section and inset showing Gi-mCherry expression in the hypothalamus.

**h**, Quantification of regional viral Gi-mCherry expression across injected animals.

Hypothalamic regions (n = 70) plotted relative to bregma anterior-posterior coordinates.

**i**, Correlation of regional viral expression across animals with the degree of torpor

perturbation identifies the anterior POA (aPOA) as a key DT-regulating region.

**j**, Higher viral Gi-mCherry expression in aPOA regions from animals exhibiting CLZ-

dependent torpor disruption (n = 4) compared to unaffected animals (n = 13). Mean ±

s.e.m.

**Fig. 3: Cell taxonomy and activity mapping implicate glutamatergic aPOA neuronal**

**cell types in DT regulation.**

**a**, Workflow to generate a cross-species POA cellular taxonomy.

**b,c**, Uniform manifold approximation and projection (UMAP) plot showing 18,406 nuclei

from mice (n = 11) and 8,355 nuclei from Syrian Hamsters (n = 7), indicating 15

GABAergic and 11 glutamatergic neuronal subtypes (**b**) or species of origin (**c**).

**d**, Hierarchical clustering of the neuronal subtypes.

**e**, Cosine similarity analysis showing pairwise transcriptional relatedness between

hamster and mouse neuronal subtypes based on mean gene expression.

**f**, Dot plot showing FIT-associated marker gene expression across mouse and hamster

neuronal subtypes.

**g**, UMAP feature plots depicting marker gene expression in the indicated species.

**h**, ISH for select FIT-associated marker genes in hamster aPOA.

**i,j**, Combined FOS immunostaining and ISH analysis for *Slc17a6* (*Vglut2*) and *Adcyap1*

from aPOA of hamsters (n = 3) in DT, with accompanying quantification. Mean ± s.e.m.

**k**, UMAP plot annotating hamster and mouse POA neurons by regional location based

on mouse MERFISH data.

**l**, UMAP feature plots depicting *Samd3* expression in the indicated species.

**m,n**, Combined FOS immunostaining and *Samd3* ISH analysis of aPOA from hamsters

(n = 3) in DT, with accompanying quantification. Mean ± s.e.m.

**a**

Differentially accessible GRE analysis

Discovery of GREs (enhancers) proximal to cell type specific markers

Packaging select enhancers into adeno-associated virus (AAVs)

*In vivo* evaluation of enhancer AAVs

GRE A    GRE B    GRE C

Marker genes

Cell type A

Cell type B

Cell type C

ITR — Enhancer A — dTomato — ITR

**b**

Mouse aPOA^Vglut2 differentially accessible regions

Mean accessibility

log2Fold Enrichment (aPOA^Vglut2/other cells)

**c**

Proximal genes for top enriched GREs

| GRE | Proximal gene |
|---|---|
| GRE 2 | Emx2 |
| GRE 4 | Gm32338 |
| GRE 1 | Samd3 |
| GRE 12 | Pcsk1 |
| GRE 10 | Gm22769 |
| GRE 7 | Gm32358 |
| GRE 8 | Gm49514 |
| GRE 9 | C330002G04Rik |
| GRE 3 | Gm26362 |
| GRE 6 | Gm4812 |
| GRE 11 | E030025P04Rik |
| GRE 5 | 4930405D11Rik |

Distance to nearest TSS (bp)

**d**

Mouse genome accessibility

Samd3E          Samd3

2kb

Chromatin accessibility

aPOA^Vglut2

cPOA^Vglut2

aPOA^Vgat

cPOA^Vgat

Astro

Oligo

Mammalian Conservation

**e**

Samd3 expression and Samd3E accessibility

Hamster: Slc17a6, Adcyap1, Esr1, Ptger3, Samd3

Positive cells [%]: 90, 70, 50, 30, 10

Mouse: Slc17a6, Adcyap1, Esr1, Ptger3, Samd3

Mean expression: 1.0, 0.5, 0.0

Samd3E

Mean accessibility: 1.0, 0.5, 0.0

GABA-04, GABA-10, GABA-03, GABA-07, GABA-02, GABA-01, GABA-11, GABA-15, GABA-14, GABA-05, Glut-03, Glut-01, GABA-09, GABA-06, GABA-13, GABA-08, GABA-12, Glut-08, Glut-02, Glut-05, Glut-11, Glut-09, Glut-10, Glut-04, Glut-06, Glut-07

aPOA^Vglut2 clusters

**f**

Hamster genome accessibility

Samd3E          Samd3

2kb

Chromatin accessibility

aPOA^Vglut2

cPOA^Vglut2

aPOA^Vgat

cPOA^Vgat

Astro

Oligo

**g**

AAV8-Samd3E-dTomato expression in the aPOA

*Vglut2*
Samd3E-dTomato

*Samd3*
Samd3E-dTomato

50µm

500µm

**h**

% Specificity dTomato⁺Marker⁺/dTomato⁺

*Vglut2*    *Samd3*

**Fig. 4: A novel Enhancer-AAV specifically targeting hamster aPOA$^{Vglut2}$ neurons.**

**a**, Workflow for generation of aPOA$^{Vglut2}$ neuron-selective enhancer-AAVs.

**b**, Scatter plot of genomic elements (n = 15,208) based on differential accessibility in aPOA$^{Vglut2}$ neurons and mean aPOA$^{Vglut2}$ accessibility signal. Top 10 most differentially accessible elements (FDR-corrected q < 10$^{-300}$) highlighted in red.

**c**, Distance to proximal gene transcription start site for the top 10 putative GREs from (**b**).

**d**, ATAC-seq genome browser tracks (normalized counts per location) for the mouse *Samd3*-proximal putative GRE (*Samd3E,* red box) in the indicated cell populations in mouse, with accompanying sequence conservation below.

**e**, Dotplot depicting *Samd3* gene expression across mouse and hamster POA neuronal subtypes, along with *Samd3E* accessibility in the mouse.

**f**, ATAC-seq genome browser tracks (normalized counts per location) for the hamster *Samd3*-proximal putative GRE (*Samd3E*, red box) in the indicated cell populations in hamster.

**g**,**h**, Combined immunostaining for viral dTomato and ISH analysis for *Slc17a6* (*Vglut2*) and *Samd3* from aPOA of hamsters injected with AAV8-Samd3E-dTomato, with accompanying quantification. Mean ± s.e.m. (n = 3 animals).

**Fig. 5: aPOA^{Vglut2} neuronal activity is necessary and sufficient for hamster deep**

**torpor entry.**

**a**, Strategy for testing the necessity of aPOA^{Vglut2} neuronal activity for DT reentry.

**b**, $T_b$ traces demonstrating perturbation of torpor resumption upon CLZ administration in

Samd3E-Gi-DREADD- but not control Samd3E-EGFP-transduced hamsters.

**c**, Samd3E-Gi-DREADD-transduced hamsters (n = 12), but not control Samd3E-EGFP-

transduced animals (n = 5), show significant perturbations in torpor initiation when

administered with CLZ but not water (Šídák's multiple comparisons test).

**d**, Strategy for testing the sufficiency of aPOA^{Vglut2} neuronal activity for DT entry.

**e**, $T_b$ traces from Samd3E-Gq-DREADD-injected animals illustrating natural torpor bouts

as well as artificial bouts elicited by CLZ administration.

**f**, Quantification of minimum $T_b$ reached in Samd3E-Gq-DREADD-transduced animals

during euthermic periods, natural deep torpor, and following administration of CLZ or

control PBS (n = 3, Tukey's multiple comparisons test).

**g**, Coronal section showing Samd3E-Gq-DREADD-EGFP expression in the hamster

aPOA.

**h**, Images of a Samd3E-Gq-DREADD-transduced hamster following PBS or CLZ

treatment.

**i**, Percentage of time Samd3E-Gq-DREADD-transduced animals (n = 6) spent inactive in

the nest over a 10 min observation window CLZ or control PBS administration (paired t-

test).

****P-adjusted < 0.0001, ***P-adjusted < 0.001, **P-adjusted < 0.01, *P-adjusted < 0.05,

ns = not significant).

**Extended Data**

**Extended Data Fig. 1 Core body temperature dynamics during hibernation.**

**a**, Representative core body temperature trace from a male hamster across its

hibernation season.

**b**, Initial $T_b$ decreases during torpor onset follow an exponential decay (animals = 188,

torpor bouts = 1262).

**c**, Body temperature traces from three animals within a 6-week observation period,

demonstrating the range of $T_b$ changes.

**Extended Data Fig. 2 Hibernation-associated neuronal activity and disruption of torpor entry during targeted silencing.**

**a**, Bar plots showing the percentage of FOS-expressing neurons across all surveyed regions (n = 32) and hibernation stages (non-hibernating = 3 animals, IBE = 3 animals, torpor = 4 animals, p-adjusted < 0.05, **p-adjusted < 0.01, ***p-adjusted < 0.001). Data presented as mean ± s.e.m.

**b**, Representative FOS immunostaining images of an animal in torpor. The top four deep torpor-active regions after the AVPe are shown: Paraventricular thalamic nucleus anterior (PVA), zona incerta (ZIM), suprachiasmatic nucleus (SCN), and paraventricular thalamic nucleus posterior (PVP).

**c**, Hibernating hamsters (n = 13) were injected with AAV-hSyn-hM4D(Gi)-mCherry, targeting different areas of the anterior hypothalamus. After recovery of DT bouts, clozapine (CLZ, 5 mg/kg) was administered via drinking water. Hamster core body temperature traces are shown, demonstrating CLZ-dependent perturbation of torpor activity in a subset of animals (n = 4) while torpor entry remains unaffected in other animals (n = 9).

**Extended Data Fig. 3 snRNA-seq analysis of hamster and mouse POA and separation of major neuronal and non-neuronal cell types.**

Our snRNA-seq and snATAC-seq data from hamster POA samples were first aligned with the mouse POA multiome dataset from Kaplan et. al[60], separating major neuronal and non-neuronal cell types from both datasets.

**a,b**, Violin plots showing the number of detected genes (**a**) and unique transcripts (UMIs) (**b**) per cell in each hamster snRNA-seq sample.

**c,d**, UMAP plots showing 43,243 nuclei across 7 hamster samples and their Leiden clustering.

**e,f**, UMAP plots showing an integrated dataset containing 45,553 hamster and mouse nuclei, and their separation into major cell types.

929    **g**, Dot plot showing the expression of select gene markers of each major neuronal and

930    non-neuronal cell type. Color indicates gene expression scaled across different cell types.

931    Circle size indicates the percentage of cells expressing each marker gene.

**Extended Data Fig. 4 Creation of a mouse mPOA reference transcriptomic dataset.**

A mouse reference transcriptomic dataset restricted to cell types in our target regions –

the medial and median POA (mPOA) – was generated. To this end, mouse spatial

transcriptomic data (MERFISH)[61] was subset to only include neurons found in the mPOA

and co-clustered with mouse neuronal snRNA-seq populations from Extended Data Fig.

3f, g, enabling the annotation of 15 GABAergic and 11 glutamatergic mPOA neuronal

populations in the mouse snRNA-seq dataset.

**a**, Coronal brain sections indicating subregions of the mPOA. MERFISH data was subset

to only include neurons within the indicated mPOA regions.

**b**, UMAP plot showing the integrated MERFISH mPOA and mouse snRNA-seq dataset.

**c,d**, UMAP plot showing Leiden clustering (**c**) and regional annotation (**d**) of the integrated

MERFISH mPOA and mouse snRNA-seq dataset.

**e**, UMAP plot of only mPOA neurons from (d) colored by Leiden cluster number (left), or

annotated clusters (right). Annotation comprises cell class (Glut = glutamatergic, GABA

= GABAergic), followed by key marker genes. These cells comprise the mouse mPOA

reference transcriptomic dataset.

**f**, UMAP plot showing expression of select marker genes in mouse mPOA.

**g**, Dot plot showing the expression of cell-type-specific marker genes in mouse mPOA.

**Extended Data Fig. 5 Integration of hamster snRNA-seq with the mouse mPOA**

**reference transcriptomic dataset.**

Leveraging the annotations from the mouse mPOA reference transcriptomic dataset

(Extended Data Fig. 4), we annotated neurons from the hamster POA snRNA-seq dataset

to generate the final cross-species atlas.

**a,b**, UMAP plot showing integrated hamster and mouse neuronal snRNA-seq data,

colored by species of origin (**a**), or Leiden cluster (**b**).

**c**, UMAP plot highlighting the location of mouse mPOA neurons, colored based on

clusters from the mouse mPOA reference transcriptomic dataset.

**d**, UMAP plot as in Fig. 3 displaying all mouse and hamster mPOA neurons from the

snRNA-seq dataset, colored by the clusters defined in the mouse mPOA reference

transcriptomic dataset.

**e**, Relative abundance of each cell type in the mouse and hamster POA dataset. Three

cell types show significant differences in abundance (* $P < 0.05$, ** $P < 0.01$, Mann-

Whitney U test).

**f**, Constellation plot showing the relationships between subtypes based on their latent

space.

**g**, Dot plot showing the expression of cell-type-specific marker genes for hamster

neurons.

**Extended Data Fig. 6 Anatomical distribution of mouse FIT-associated neurons.**

**a,b**, Example coronal sections of the mouse brain containing the anterior POA, showing

ISH signal for marker genes *Slc17a6/Vglut2* (**a**) and *Adycap1* (**b**). Data from mouse.brain-

map.org[85].

**Extended Data Fig. 7 Marker expression of DT-active neurons; anatomical**

**localization and cross-species differential gene expression of POA neuronal**

**populations.**

**a**, Example coronal sections showing the aPOA of a Syrian hamster collected during DT

onset. Immunofluorescent staining against FOS indicates the location of active neurons

(yellow), whereas ISH indicates the expression of the marker gene *Slc32a1* (*Vgat*). High-

magnification images are shown as insets with example FOS+ *Slc32a1*+ cells circled.

**b**, Quantification of the fraction of FOS+ neurons during torpor that express *Slc32a1*

(12.8% ± 1.5, animals = 3) and dual-expressing *Slc17a6/Slc32a1* (1.8% ± 0.3, n = 3)

neurons.  Data presented as mean ± s.e.m.

**c**, MERFISH plot showing the spatial distribution of neuronal subtypes in the mouse POA

separated by their major cell types and anatomical location.

**d**, Differential gene expression between hamster and mouse neurons across the four

major neuron groups.

**Extended Data Fig. 8** *CamkIIa* **promoter-based AAVs lack specificity for glutamatergic neurons in the hamster aPOA.**

Hamsters (n = 4) were stereotaxically administered AAV8-CamkIIa-GFP and harvested ~2 weeks later for immunostaining and ISH.

**a**, Example coronal sections showing the hamster POA comprising AVPe, MPA, and MnPO. Immunofluorescent staining against GFP indicates the location of virally labeled cells (yellow), whereas ISH indicates the expression of genes *Slc17a6* (*Vglut2; magenta*) and *Slc32a1 (Vgat; cyan)*. High-magnification images are shown on the right with example GFP+ *Marker*+ cells circled.

**b**, Quantification of the fraction of AAV-CamkIIa-GFP+ neurons that express *Slc17a6* (42.6% ± 2.0, animals = 4), and *Slc32a1* (47.5 ± 5.0%, animals = 4). Data presented as mean ± s.e.m.

**a**

**b**

| | Genomic coordinates (mm10) | Nearest gene |
|---|---|---|
| GRE 1 | chr10:26228096-26228596 | Samd3 |
| GRE 2 | chr19:59457295-59457795 | Emx2 |
| GRE 3 | chr6:111965822-111966322 | Gm26362 |
| GRE 4 | chr10:41139206-41139706 | Gm32338 |
| GRE 5 | chr11:91056877-91057377 | 4930405D11Rik |
| GRE 6 | chr13:69317802-69318302 | Gm4812 |
| GRE 7 | chr16:96450604-96451104 | Gm32358 |
| GRE 8 | chr16:23505809-23506309 | Gm49514 |
| GRE 9 | chr19:23085045-23085545 | C330002G04Rik |
| GRE 10 | chr16:63743408-63743908 | Gm22769 |
| GRE 11 | chr11:109106715-109107215 | E030025P04Rik |
| GRE 12 | chr13:75113945-75114445 | Pcsk1 |

**d**

**Mouse Samd3E accessibility**

chr10:26228096-26228596

**e**

**Hamster Samd3E accessibility**

NW_024429195.1:17389041-017389541

**Extended Data Fig. 9 snATAC-seq analysis reveals *Samd3E* as tool to target POA**

**neurons.**

**a**, Differential accessibility analyses of genomic elements in aPOA$^{Vglut2}$ neurons vs. all

other neurons in mice; top regions (log2FC > 0.5) are marked in red for further analysis.

**b**, Genomic location and nearest gene of the top candidate putative GREs.

**c**, ATAC-seq genome browser tracks (normalized counts per location) of individual mouse

neuronal subtypes showing the mouse *Samd3*-proximal putative GRE (*Samd3E,* red

box), with accompanying sequence conservation below.

**d,e**, UMAP feature plots depicting *Samd3E* accessibility in the indicated species.

1014

**Extended Data Fig. 10 Samd3E-AAV transduction efficiency.**

**a**, Efficiency of targeting aPOA$^{Vglut2}$ and aPOA$^{Samd3}$ neurons with AAV8-Samd3E-

dTomato. Syrian hamsters (n = 3) were injected with AAV8-Samd3E-dTomato.

Immunofluorescent staining against dTomato was used to identify virally labeled cells

(yellow), whereas ISH indicates the expression of the marker gene *Slc17a6* (*Vglut2*,

magenta) and *Samd3* (cyan), as in Fig. 4g.

**b**, Quantification of the fraction of *Slc17a6*+ cells that also express dTomato+ (20.6 ±

3.9%, animals = 3) and *Samd3*+ cells that also express dTomato+ (23.3 ± 2.8%, animals

= 3). Given limited viral spread, this quantification was limited to the area of viral

expression within the aPOA. These results indicate that AAV8-Samd3E-dTomato

achieves high specificity and modest efficiency for targeting aPOA$^{Vglut2}$ and aPOA$^{Samd3}$

neurons.

**a**

**b**

**c**

**d**

1027

**Extended Data Fig. 11 Circadian $T_b$ patterns and correlation between viral aPOA**

**expression and $T_b$ decrease.**

**a**, Circadian fluctuation of $T_b$ in Samd3E-EGFP- and Samd3E-Gi-DREADD-injected

hamsters following administration of CLZ (5 mg/kg) or control PBS. Light and dark cycle

phases are represented by yellow and grey, respectively. Data presented as mean ± s.d.

of core body temperatures during each hour.

**b**, Violin plot quantification of body temperature range between light (yellow) and dark

(grey) phases of Samd3E-EGFP- (n = 5 animals) or Samd3E-Gi-DREADD-injected

animals (n = 12) following CLZ or control PBS administration (Tukey's multiple

comparisons test; *P-adjusted < 0.05, ****P-adjusted < 0.0001, ns = not significant P >

0.05).

**c**, Immunofluorescence images in the aPOA of additional Samd3E-Gq-DREADD-EGFP-

injected animals.

**d**, Correlation between minimum $T_b$ reached following CLZ administration and viral

expression within the aPOA (n = 5 animals, r = -0.95, P = 0.01).

1043

**Extended Data Fig. 12 Changes in behavioral repertoire after aPOA$^{Vglut2}$ neuronal activation.**

Percentage of time in behavioral states over a 10 min observation window post-injection with PBS or CLZ across all hamsters injected with AAV-Samd3E-EGFP-P2A-Synaptophysin-mRuby (n = 3) and AAV-Samd3E-hM3D(Gq)-nlsGFP (n=6).

1049

**Extended Data Fig. 13 Stimulation of aPOA$^{Vglut2}$ neurons in mice and resulting body temperature change.**

**a**, Schematic of experimental design.

**b**, Timeline of experiment and analysis. Mice were acclimated to 8°C for 3 days before injection with CLZ (5 mg/kg) and imaged with a thermal camera at baseline, 30 min, 60 min, 3 hours, 6 hours, and 24 hours post injection at 8°C.

**c**, Representative thermographic images obtained by chemogenetic stimulation of aPOA$^{Vglut2}$ neurons.

**d**, Surface body temperature before (0 min) and after (30 min to 24 h) CLZ injection (n=7 mice). Data are presented as mean ± s.e.m.

a

b

1060

**Extended Data Fig. 14** *In situ* **hybridization for marker of torpor-associated genes**

**in marmosets.**

**a**, Example coronal sections of the marmoset brain containing the anterior POA showing

ISH signal for marker genes *Slc17a6/Vglut2* and *Adycap1*.

**b**, Higher resolution images from (a) displaying the anterior POA. Data from https://gene-

atlas.brainminds.jp/[72–74].