1 **<u>Prediction variability in physiologically based pharmacokinetic modeling of tissue disposition</u>**
2 **<u>under deep uncertainty</u>**
3

Mustafa Farahat[1], DT Flaherty[1], Zachary R Fox[2], Belinda S Akpa[1,3,4]*

[1]Department of Chemical and Biomolecular Engineering, University of Tennessee-Knoxville, Knoxville,
Tennessee, USA

[2]Computational Sciences and Engineering Division, Oak Ridge National Laboratory, Oak Ridge,
Tennessee, USA

[3]National Institute for Modeling Biological Systems, Knoxville, Tennessee, USA

[4]Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

*Corresponding author & Lead Contact

E-mail: bakpa1@utk.edu

Short title: Prediction variability in ML-informed pharmacokinetic modeling

Notice: This manuscript has been authored by UT-Battelle, LLC, under contract DE-AC05-
00OR22725 with the US Department of Energy (DOE). The US government retains and the
publisher, by accepting the article for publication, acknowledges that the US government retains a
nonexclusive, paid-up, irrevocable worldwide license to publish or reproduce the published form of
this manuscript, or allow others to do so, for US government purposes. DOE will provide public
access to these results of federally sponsored research in accordance with the DOE Public Access
Plan (http://energy.gov/downloads/does-public-access-plan).

Abstract

Physiologically based pharmacokinetic (PBPK) models are increasingly invoked in virtual screening workflows for therapeutics. These mechanistic models project pharmacokinetic outcomes from molecular properties, with data-driven models acting as intermediaries to map molecular structure to PBPK input parameters. Errors in predicted parameters and unvalidated assumptions within PBPK models expose PK predictions to deep uncertainty. Herein, we examine how these uncertainties affect the prediction variability of dynamic, tissue-specific exposure. We validated four PBPK models against 1,854 experimental datapoints – to establish their predictive fidelity before introducing parameter uncertainty typical of property-prediction models. Depending on molecule properties and model choice, the coefficient of variation under parameter uncertainty ranged from 10^{-6} to 31 for predicted PK statistics. Further, we identified notable model disagreement for a subset of drug-like chemical space characterized by lipophilic, protonated molecules. Uncertainty quantification revealed biophysicochemical properties and parameter interactions that drove disagreement and highlighted model assumptions that exacerbated prediction variance. Our findings delineate the challenges presented by deep epistemic uncertainty in PBPK modeling.

Keywords: PBPK modeling, epistemic uncertainty, uncertainty propagation, prediction variability.

Introduction

AI-driven drug discovery is evolving rapidly as academic groups, industry, and start-ups develop computational tools to generate and evaluate candidate molecules[1–4]. Computational approaches can shift screening paradigms from sequential experimentation to parallel exploration[5–7], making it possible to evaluate candidate structures through workflows that integratively account for molecular activity[8,9], identify off-target interactions[10–12], and discriminate toxicity liabilities[13,14] – amongst other therapeutic criteria. Predicting these properties directly from chemical structures is a key step that supports high-throughput screening before synthesis, empirical development, and *in vivo* testing[15,16]. However, whether such predictions accelerate discovery depends on the ability of constituent models to capture the features that make a molecule a successful therapeutic. For a molecule to function as a drug, it must engage in desired molecular interactions, reach its site of action within the body, and persist there at concentrations sufficient to modulate biological processes[17–19]. Many efforts in the domain of AI-driven discovery focus on data-driven[20,21] and surrogate models[22,23] that predict molecular-scale phenomena governing candidate potency and safety. What may be neglected in these workflows are the dynamic features of human physiology that determine whether a compound can reach its target *in vivo*, at high enough concentrations, and for long enough to influence disease processes[17,24,25]. Physiologically based pharmacokinetic (PBPK) modeling can address these issues by mapping molecular properties to chemical fate in the human body.

PBPK models are systems of ordinary differential equations that mechanistically describe the body as an assembly of organ compartments connected by circulating blood[26–30]. Each equation takes the form of a material balance capturing the influence of tissue-specific composition on molecule distribution and metabolism. Solving the system of equations thus yields tissue-specific, time-resolved predictions of drug exposure – pharmacokinetic outcomes directly relevant to therapeutic efficacy. As a well-established tool in later stages of drug development[28,31–33], PBPK modeling supports *post hoc* rationalization of therapeutic outcomes and helps refine therapeutic regimens[34,35]. The models may be used to prospectively explore

clinical dosing strategies, assess drug-drug interactions, or predict outcomes for special populations such
as pediatric patients or those with impaired organ function[36–39]. Their informational value in these late-
stage applications has driven growing efforts to deploy PBPK models in virtual screening. Emerging
screening workflows now combine machine learning (ML) models that predict molecular properties with
PBPK frameworks to project *in vivo* drug behavior directly from chemical structures[40–46], and possibly
augment the decision-making value of novel alternative methods[47–50]. However, applying PBPK models
in this way introduces new challenges. When applied at the clinical stage or later, these models rely on
experimentally measured parameters and can be validated against drug- and species-specific observations.
When PBPK models are instead used in ML-informed workflows, establishing their credibility becomes
more difficult. A primary issue is the impact of uncertainty arising from two sources.

The first source is structural. Any PBPK model is an abstraction of the true physiological system[51].
When applied broadly across diverse chemical space, such assumptions may misrepresent key
physiological processes in ways that are not easily detected without experimental validation[26,52,53]. For
practical decision-making in virtual screens, a single model – or small set of models – would need to
produce informative predictions across a wide range of chemical structures. The second source of
uncertainty is parametric. For virtual compounds, drug properties cannot be determined experimentally
and instead drug-specific input parameters for PBPK modeling can be obtained from data-driven
quantitative structure-activity relationship (QSAR) models[54,55]. These approaches are often trained on
sparse or biased datasets and can lead to large, compound-specific prediction errors. As a result, both the
PBPK framework and its input parameters produce a scenario of deep epistemic uncertainty.
Understanding how this uncertainty propagates through PBPK frameworks is key to judging whether
model predictions are useful in early virtual-screening workflows.

In this study, we assess the confidence associated with PBPK predictions for virtual compounds,
where both model assumptions and input parameters are uncertain. To this end, we examined: model
prediction fidelity (for known drugs with empirically characterized input parameters) and prediction
variance (for virtual compounds whose input parameters are perturbed to reflect the uncertainty typical of
published QSAR models). We quantified prediction fidelity by evaluating each model's ability to predict
plasma-tissue partitioning coefficients (Kp) and volume of distribution (Vss) – both equilibrium PK
parameters with available empirical data for model validation. Then, with ML-informed PBPK workflows
in mind, we introduced parameter uncertainty representative of the mean absolute error reported in
published property-prediction models[56–58]. We propagated this uncertainty through four PBPK models: (1)
our own model, modified from the Rodgers & Rowland (R&R)-Lukacova framework, (2) a second R&R-
Lukacova based model reported by Mathew *et al.*[59], and (3 & 4) the calibrated and uncalibrated variants
of the Schmitt-based model reported by Pearce *et al.*[60]. Using Monte Carlo simulations, we quantified
tissue-specific PK statistics relevant to intracellular drug exposure for a synthetic dataset of 10^4 virtual
molecules. Specifically, for each molecule, we recorded steady-state volume distribution (Vss), maximum
unbound concentration ($Cmax,u$), area under the unbound concentration-time curve ($AUCu_{0-\infty}$), and the
duration of time for which unbound concentration exceeded 0.1 μM ($T_{0.1\mu M}$). We thereby quantified the
impact of epistemic uncertainty on prediction variance of model-derived PK outcomes.

Our analysis of prediction variance addressed three core questions: (1) Do PBPK models embedding
different mathematical abstractions for tissue partitioning exhibit different susceptibility to parameter
uncertainty? (2) If so, where in chemical space do these predictions differ most, and is that divergence
evident for nominal predictions or specifically induced under parameter uncertainty? (3) Which drug-
specific input parameters contribute most to prediction variability and model disagreement, and where
would efforts to improve property prediction models be most impactful?

To answer these questions, we (1) evaluated model agreement, (2) identified regions of chemical
space with divergent predictions, and (3) applied global sensitivity analysis to quantify the contribution of
each input parameter to prediction uncertainty. We have thereby shed light on challenges to consider
when using ML-informed PBPK workflows to guide compound assessment, particularly when parameter
uncertainty is high and empirical validation is not feasible.

Results and Discussion

Mechanistic models have been developed to predict one critical class of drug-specific PBPK
parameters: plasma-tissue partition coefficients (Kp). These constants describe the equilibrium
partitioning of drug between the blood plasma and a given tissue and can be predicted as a function of
tissue composition. Partitioning models may have an explicitly or implicitly bounded domain of
applicability or contain terms that must be selected based on molecular properties. Some work well for
acids and zwitterions, others for strong bases, and still others for highly lipophilic compounds[61–64]. Most
are designed to address known drug-like compounds, while others aim to be predictive over a broader
range of molecules from drugs to chemicals of concern for environmental exposure[65]. In all cases,
prediction of Kp requires biophysicochemical property inputs for the molecule of interest. For PBPK
models that are used early in the discovery pipeline, QSAR models provide a data-driven mapping of
molecule structure to molecular properties.

In this study, we interrogate PBPK models based on three tissue-partitioning models: 2 canonical
approaches and a modified variant introduced herein. As reference approaches, we consider the model of
Mathew *et al.*[59] and that of Pearce *et al.*[60]. These represent, respectively, modifications of the
R&R-Lukacova framework and the Schmitt framework for predicting tissue-plasma partitioning. The
model we introduce herein is based on the R&R-Lukacova framework, adopting alternative assumptions
for drug association with proteins (interstitial and intracellular), neutral lipids, and phospholipids. The
distinguishing features of the three Kp models are presented in the Methods section (**Table 1**). In
considering a modified formulation for tissue partitioning relationships, we aimed to improve the model's
ability to generate predictions of similar quality across drug-like chemical space.

Model fidelity-to-data

To assess model performance, we quantified four common statistics that capture fidelity with respect
to experimentally determined equilibrium PK quantities – namely, tissue partitioning coefficients and
volume of distribution. We subsequently quantified each model's: (1) *accuracy*, as indicated by the
fraction of drugs with PK predictions within 2-fold, 3-fold, and 10-fold error; (2) *precision*, as indicated
by the root mean squared error of log-transformed values, RMSLE; (3) *bias*, as indicated by average fold
error, AFE; and (4) *agreement*, as indicated by the concordance correlation coefficient, CCC. CCC
describes the extent to which predictions fall on the identity line of a parity plot (*i.e.*, when predictions are
plotted against measured quantities). As with other correlation coefficients, CCC ranges between -1 and
1. We provide equations for each of these performance metrics in the Methods section.

Data sources

PBPK models have three classes of parameter: (1) those that reflect the nature of the biological
system (*e.g.*, tissue volumes and composition), (2) those that solely reflect the physicochemical properties
of the drug (*e.g.*, distribution coefficient, *logD*, and acid dissociation constants, *pKa*), and (3) those that

capture interactions between drug chemistry and biology (*e.g.*, metabolic clearance, CL_{int}, and plasma

protein binding, *fu*). **Table S1** summarizes each dataset used herein, its source, and its application within

this study.

<u>System-specific data</u>

System-specific parameters for rat and human physiology were taken from the Httk *physiology.data*

table[66]. Likewise, tissue composition information was adapted from Httk (*tissuedata.data*)[66]. Data

describing tissue vascular fraction and tissue albumin and lipoprotein concentrations were drawn from

Edginton *et al*. and Rodgers *et al.*, respectively[67,62]

<u>Drug-specific data</u>

To assess the fidelity of *Kp* predictions, we sourced rat tissue partitioning data from the dataset

collated by Pearce *et al.*[60]. This dataset includes 157 distinct compounds over 11 tissues, with a total of

N = 992 data points. For *Vss*, we used the human *in vivo* values reported by Obach *et al.*[68] – limiting our

predictions to the 862 compounds for which those authors present the required set of drug-specific input

parameters. **Figure S1** shows the distributions of these molecular properties in the Pearce and Obach

datasets.

Model calibration

A key feature of the Pearce model is the use of an empirical correction to adjust predicted values of

Kp[60]. These corrections were derived from linear regressions of predicted versus measured tissue-partition

coefficients for 157 pharmaceutical compounds across 11 tissues. The resulting tissue-specific linear

corrections yield a calibrated set of *Kp* values that are intended to improve overall prediction fidelity.

To assess this model's fidelity on equal footing with the other *Kp* models (*i.e.*, those that have not

been "tuned" to that empirical dataset), we evaluated its performance based on the uncalibrated *Kp*

predictions. We then used the calibrated form of the Pearce model when assessing the credibility of

human *Vss* predictions.

Canonical models exhibit similar fidelity to rat *Kp* and human *Vss* data

When tallied over all 157 drugs and 11 tissues in the *Kp* dataset, the Mathew model underperformed

in predicting rat tissue partitioning coefficients (**Figure 1A**). Less than 40% of *Kp* values were predicted

within a 2-fold error. The Mathew model also failed to exceed the 90% mark for predictions within a 10-

fold error. Although our model shares an underlying framework with Mathew (*i.e.*, that of R&R-

Lukacova), it performed with accuracy comparable to that of uncalibrated Pearce. When we

disaggregated the drug molecules by ion class, it became clear that the overall performance of the

Mathew model was disproportionately impacted by poor predictions for zwitterions – for which less than

20% were predicted within a 2-fold error. Across acids, bases, and neutral compounds, we observed more

parity in performance among the three models. However, the Mathew model did not yield greater

accuracy than our model or Pearce's for any of the ion classes. The detailed performance statistics for rat

Kp across ion classes are presented in **Table S2**.

Note that these fold-error quantities should be interpreted in the context of the number of

experimental measurements reported for compounds in each ion class. By far the largest group was bases

(N=584), followed by acids (N=258), zwitterions (N=92), and neutrals (N=58). At the two-fold error level

our model appears to perform comparatively well on neutral compounds. However, the ~14% difference

between our model and Pearce's uncalibrated model corresponds to only 8 data points. As the validation
data does not provide a measurement for every tissue across the full set of drugs, this difference of 8 data
points could represent 8 different compounds, with empirical measurements within the same tissue, or the
same compound measured over 8 different tissues (or, indeed, distinct compounds across distinct tissues).
Thus, we interpreted the relative accuracy of the *Kp* predictions with caution.

Figure 1. Fold error statistics for each model and ion class. (A) Rat *Kp*. Statistics summarize the quality of 992 predictions per model, spanning 157 molecules and 11 tissues. (B) Human *Vss*. Statistics summarize the quality of 862 predictions per model – one *Vss* prediction per molecule.

When we inspected human *Vss* predictions, we again noted similar accuracy between our model and
that of Pearce (now the calibrated model), with the Mathew model lagging in performance (**Figure 1B**).
When disaggregated, however, the Mathew model maintained something closer to parity across all ion
classes. There is comparatively poorer performance again for zwitterions, but it is less marked than in the
Kp evaluation. For *Vss*, the underperformance with respect to zwitterions is likely dampened by the fact
that this PK metric is a composite quantity, computed as a *Kp*-weighted sum over the volumes of
individual tissues (**Equation 19**). Tissues for which equilibrium partitioning constants are poor reflections

of relevant biological transport phenomena[69–72] (*e.g.*, brain, liver) represent relatively small contributions
by volume[66,73]. By contrast, tissues that represent disproportionately large volumes (*e.g.*, muscle, adipose)
are relatively well described by equilibrium partitioning[74,75]. The full set of *Vss* performance statistics for
each ion class is summarized in **Table S3**.

In **Figure 2**, we compare disaggregated *Kp* predictions for the brain, adipose, and muscle
compartments. These tissues represent ~2%, ~24%, and ~40% of the body's volume, respectively[66,73]. As
expected, the predictive accuracy for brain tissue was relatively poor, with CCC below 0.6 in all models.
Correlations for adipose *Kp* predictions were the best out of all tissues, for all models. This is despite the
models invoking different approaches to describe neutral lipid partitioning in adipose tissue. Our model
and that of Mathew follow the practice of adopting the vegetable-oil-water partition coefficient to capture
the high triglyceride content of that tissue. The Pearce model does not make this adjustment. The Pearce
model does, however, use *logD* rather than *logP* to describe lipid partitioning – a feature that is shared by
our model. Mathew differs in that *logP* is retained, but the partitioning is only associated with the fraction
of molecules predicted to be in the uncharged state.

Figure 2. Measured vs. predicted rat tissue partitioning coefficients (*Kp*). Left: This model. Center: Mathew model. Right: Uncalibrated Pearce model. Lines indicate the identity line, two-fold, three-fold, and ten-fold error intervals.

16   **Figure 3A** summarizes the RMSLE, AFE, and CCC performance statistics for the complete set of *Kp*
17   predictions. Across all ion classes except bases, our model exhibits the lowest RMSLE (*i.e.*, highest
18   precision), along with a consistent tendency to underpredict tissue partitioning (*i.e.*, log$_2$AFE < 0). This

1 tendency is shared by the other R&R-Lukacova model, *i.e.*, that of Mathew, which yields the greatest

2 prediction biases. Overall *Kp* prediction bias is least pronounced for the Pearce (uncalibrated) model. This

3 pattern is not replicated across the *Vss* predictions (**Figure 3B**). Here, for all ion classes, prediction bias is

4 greatest for the calibrated Pearce model, where *Vss* tends to be overpredicted.

Figure 3. Precision, bias, and correlation statistics for each model and ion class. (A) Rat *Kp*. Statistics summarize the quality of 992 predictions per model, spanning 157 molecules and 11 tissues. (B) Human *Vss*. Statistics summarize the quality of 862 predictions per model – one *Vss* prediction per molecule.

In short, despite the non-trivial differences in underlying framework and assumptions, all three

models yield predictions of broadly similar fidelity to data. Differences are, however, observed at the

level of individual tissues. These tissue-level errors are obscured at the level of *Vss*, as *Kp* predictions for

the largest two tissues have high fidelity. Tissue-specific errors will, however, be of concern for

evaluating any tissue-specific exposure criterion. Further differences emerge when results are

disaggregated by ionization class. Our analysis demonstrates that specific assumptions modulate error and

bias within a PBPK framework and the choice of underlying framework from amongst canonical

approaches is not a defining determinant of model credibility. This conclusion, however, assumes perfect

knowledge of each drug's properties. Within an ML-informed PBPK workflow, drug-specific input

parameters (*e.g.*, RMM, *fu*, CL_{int}, *logD*, donor and acceptor *pKa*) are obtained from QSAR models that

map chemical structure to the molecular properties required for PBPK simulation. Data reflecting human

biology can be limited and highly variable, so data-driven prediction of parameters *fu* and CL_{int} proves

particularly recalcitrant[76–81]. Identifying informative molecular features and predictive modeling strategies

to generate these drug-specific parameters is a foundational requirement for reliable predictions – and

thus an active area of research[82–85]. In the following sections, we examine the impact that prediction errors
typical of existing property-prediction models would have on the uncertainty of PBPK predictions.

Prediction variance under epistemic uncertainty

To be useful as a tool for decision making in an ML-informed PBPK workflow, a PBPK model would
need to exhibit two key properties: (1) the model should be adequately sensitive to variations in molecular
properties to reflect the impact of those variations on PK outcomes of therapeutic consequence, and (2)
despite uncertainty in predicted parameter values, those PK predictions should remain informative about
the relative PK performance of virtual candidates. The prior section addressed the first concern by
evaluating PBPK model fidelity for known compounds with measured biophysicochemical parameters –
considering the epistemic uncertainty associated with model formulation. We now turn to the second
concern by examining how PBPK predictions respond to uncertainty in their input parameters.
Specifically, we assess whether predictions retain their informative value once uncertainty is introduced
and whether parameter uncertainty has disparate impacts on model predictions in particular regions of
chemical space.

To do so, we generated 10^4 pseudomolecules by correlated random sampling of RMM, fu, CL_{int}, $logD$,
and pKa from property distributions characterizing the compounds in the Obach dataset (see pseudocode
in **Algorithm 1**). We then evaluated the impact of parameter uncertainty by performing Monte Carlo
simulations using perturbed molecular properties. To this end, for each pseudomolecule, we assumed a
parametric distribution centered on the molecule's "true" property value and scaled its spread to match
the mean absolute error (MAE) reported for published QSAR models[56–58] (see **Table 2**). $LogD$ was
sampled from a normal distribution, fu and pKa from beta distributions, and CL_{int} from a lognormal
distribution. We ran 10^3 Monte Carlo realizations per pseudomolecule through each of the four PBPK
models. The resulting unbound concentration-time profiles allowed us to compute outcome distributions
for Vss and dynamic tissue-specific PK metrics $Cmax,u$, $AUCu$, and $T_{0.1\mu M}$.

Model calibration limits Vss prediction variance

We began our analysis by examining the variance of Vss predictions, as this PK outcome provides
direct continuity with the previous fidelity analysis. As there are no experimental Vss values to provide
ground-truth for our pseudomolecules, the following comparisons are strictly between models. We first
examined the variances of the Vss distributions produced by propagating identical parameter uncertainty
through the four PBPK models. Over the 10^4 pseudomolecules, median prediction variances were broadly
comparable ($\sigma^2 \sim 10^{-1}$, **Table S4**). The one clear exception was the uncalibrated Pearce model, which
exhibited a median variance one order of magnitude greater than that of the other three models. Once
calibrated, the median variance for the Pearce model fell more in line with the other PBPK models (95%
interval 0.16-0.17 L^2/kg^2), and this model's prediction variance became fairly uniform across the
pseudomolecule population. Indeed, it appears that model calibration markedly limits the spread of Pearce
Vss predictions (**Figure S2A**). By contrast, the Mathew model yielded predictions with variance as large
as $10^{16} L^2/kg^2$, depending on the pseudomolecule in question. Notably, there are several outliers with
variance greater than $10^7 L^2/kg^2$ (**Figure S2B**). The only other model yielding several pseudomolecule
outliers was the calibrated form of the Pearce model, which yielded very narrow distributions ($\sim 10^{-3}$
L^2/kg^2) for its outliers – indicative of predictions that are relatively insensitive to parameter uncertainty.

Figure 4. Mathew model exhibits PK outcome prediction discordance for a subset of 751 pseudomolecules (of 10,000 total). (A) Distributions of PK outcomes resulting from parameter uncertainty, as predicted by each of the PBPK models. Distributions shown are prototypical of molecules assigned to: Top - Cluster 1. Bottom - Cluster 2. **(B)** Concordance correlation coefficients for predicted PK outcomes, computed across all pseudomolecules in each cluster (Top - Cluster 1: 751 pseudomolecules. Bottom - Cluster 2: 9249 pseudomolecules). Analysis reflects 10^3 Monte Carlo realizations for each of the 10^4 pseudomolecules and thus summarizes the outcomes of 10^7 simulations per PBPK model. Each heatmap shows pairwise agreement between models (T: this model, M: Mathew, PC: Pearce, calibrated, PU: Pearce, uncalibrated).

Model predictions diverge for a subset of chemical space representing highly protonated,
lipophilic molecules

We next quantified the extent to which model predictions agreed under uncertainty and, thereby,
sought to identify which regions of chemical space provoke model disagreement. We also broadened our
analysis to include dynamic PK outcomes relevant to local tissue exposure. Specifically, we examined
$Cmax,u$, $AUCu_{0-\infty}$, and $T_{0.1\mu M}$ in muscle intracellular water. For each pseudomolecule, PK prediction
distributions were log-transformed, standardized with a whitening transformation, and contrasted by
calculating a Wasserstein distance – a metric that captures differences in both location and shape of paired
distributions[86–88] (see pseudocode in **Algorithm 2**). This analysis produced six distances per PK outcome.
Across the four PK outcomes, this resulted in 24 features that summarized model-to-model agreement
(predictions from four models, compared pairwise). To distinguish patterns of model agreement across
chemical space, we applied K-means clustering to group molecules by these 24 metrics – using a
silhouette statistic to select the number of clusters. We thereby identified two groups among the 10^4
pseudomolecules: a smaller subset (751 molecules; 7.5%) characterized by greater prediction discordance
(*i.e.*, larger Wasserstein distances) and a larger subset (9249 molecules; 92.5%) exhibiting broad
concordance across models. A heatmap of pairwise distances (**Figure S4**) illustrates the segregation
across clusters, and **Figure 4A** depicts predicted distributions for one prototypical pseudomolecule from
each cluster.

To quantify the degree of agreement across all molecules within a given cluster, we evaluated the
cluster-specific CCC for each PK outcome (**Figure 4B**). As for the examples shown in **Figure 4A**
(bottom), predicted distributions for molecules in Cluster 2 largely concur across all outcomes (median
$CCC_{cluster\ 2} = 0.95$; 95%CI:[0.61, 1.00]). The discordance that arises in Cluster 1 (median
$CCC_{cluster\ 1} = 0.76$; 95%CI:[0.029, 1.00]), is most evident for prediction of Vss ($CCC_{cluster\ 1,Vss} = 0.51$;
95%CI:[0.083, 1.00]), and $T_{0.1\mu M}$ ($CCC_{cluster\ 1,T0.1\mu M} = 0.40$; 95%CI:[0.024, 1.00]). The Mathew model
appears to be the principal outlier here, presenting pronounced disagreement with all other models for
Vss, $Cmax,u$, and $T_{0.1\mu M}$.

As a first step to determine what distinguishes molecules whose predictions are discordant across
models, we examined the biophysicochemical properties of the pseudomolecules assigned to each cluster
(**Figure S5**). We found that molecules in Cluster 1 are exclusively bases (78%) and zwitterions (22%),
while molecules in Cluster 2 were more evenly distributed across ion classes. Furthermore, as judged by
two-sample Kolmogorov-Smirnov tests, molecules in Cluster 1 are more lipophilic (*logD*: $D_{KS} = 0.67$,
Cohen's d = 1.67) and more likely to be highly protonated (acceptor *pKa*: $D_{KS} = 0.70$, Cohen's d = 1.92)
than those in Cluster 2 ($D_{critical} = 0.063$, $\alpha = 0.0083$; **Figure S6**).

For highly protonated, lipophilic molecules, prediction discordance arises from model structure
and is exacerbated by parameter uncertainty

We next sought to determine: (1) whether Cluster 1 represents a distinct sub-region of chemical space
within which the four PBPK models will make discordant predictions; (2) whether the four models make
distinct nominal predictions for molecules in this region or, alternatively, (3) whether the disagreement
between models arises due to the simulated parameter uncertainty. Further, if nominal predictions
happened to diverge, we established whether one model provided objectively more credible predictions
than another. To assess credibility, we needed to compare model predictions to ground-truth, which
motivated a return to simulating PK outcomes for molecules in the Obach data set, for which we have
empirical data for Vss. We performed Monte Carlo simulations for the Obach molecules and used the

resulting Wasserstein distance features to map these real drug molecules to the clusters we defined using
the pseudomolecules. By examining fidelity of nominal *Vss* predictions alongside the model agreement
features, we were able to distinguish whether divergent predictions stemmed from differences in
structural assumptions or from the impact of parameter uncertainty.

Figure 5. For protonated, lipophilic molecules, the two R&R models yield nominal predictions with greater discordance, and that discordance is exacerbated under uncertainty. Axes report nominal *Vss* prediction fidelity for: X-axis - This model (TM). Y-axis - Mathew model (M). Colormaps indicate the Wasserstein distances between *Vss* distributions predicted under parameter uncertainty. Colormaps are overlaid with markers indicating the locations of individual compounds. Circles: molecules for which the two models' nominal predictions differ by less than a factor of two. Triangles: compounds for which predictions differ by a factor greater than two. **(A)** Cluster 1: 240 molecules, ≤ 2-fold difference: 108 molecules, > 2-fold difference: 132 molecules. **(B)** Cluster 2: 622 molecules, ≤ 2-fold difference: 397 molecules, > 2-fold difference: 225 molecules. **(C)** Summary of the Cohen's d effect sizes distinguishing the properties of molecules in each sub-group. Asterisks indicate those features with a statistically significant difference between the near-diagonal and off-diagonal molecules – as determined by Kolmogorov-Smirnov statistic, D_{KS}, exceeding the critical values of $D_{critical}$ = 0.21 and 0.14, respectively, for clusters 1 and 2. **Table S6** reports full results of the statistical comparison.

Twenty-eight percent (N = 240) of the 862 Obach compounds mapped to pseudomolecule Cluster 1 –
the cluster corresponding to discordant PBPK predictions under uncertainty. To explore the nature of this
discordance, we compared nominal *Vss* predictions between pairs of models. **Figure 5** shows a
representative result for the two R&R-based models, our model and that of Mathew. Markers indicate the
location of Obach molecules, positioned according to the fidelity of their nominal predictions. As the axes
report *Vss* prediction fidelity with respect to ground truth, molecules located along the diagonal will be
those for which the two models make the same nominal prediction – irrespective of whether that
prediction matches the experimentally observed value of *Vss*. The underlying colormap indicates the
Wasserstein distance between Monte Carlo-derived *Vss* distributions for the two models.

We divided the 240 compounds in Cluster 1 into two groups based on their nominal *Vss* predictions:
(a) a near-diagonal set (N = 108), representing compounds whose predictions from the two models
differed by less than a factor of two, and (b) an off-diagonal set (N = 132), representing compounds with
larger fold-differences in nominal predictions. In **Figure 5A**, we found – perhaps unsurprisingly – that
agreement between nominal predictions (*i.e.*, near-diagonal compounds) appeared to coincide with
regions of less discordance under uncertainty – as determined by comparing Wasserstein distances for
near diagonal and off-diagonal molecules (Cohen's d effect size = -1.6, D_{KS} = 0.77, $D_{critical}$ = 0.18). Where
nominal predictions disagreed (*i.e.*, off-diagonal markers), the underlying colormap indicated greater
discordance under uncertainty. This suggests that much of the observed discordance is due to differences
in model structure. Parameter uncertainty did, however, promote greater discordance between models for
molecules whose nominal predictions generally concurred (Cohen's d = 0.94, D_{KS} = 0.35, $D_{critical}$ = 0.15,
when comparing Wasserstein distances for near-diagonal molecules in the two clusters). Thus, parameter
uncertainty appears to promote greater divergence in model predictions for the type of molecule that
appears in Cluster 1.

Figure 6. Effect sizes characterizing property differences between Obach molecules assigned to clusters 1 and 2. (A) Gardner-Altman plots of properties for molecules in each cluster. Second y-axis indicates the effect size, calculated as Cohen's d values, with bootstrapped confidence intervals. (B) Summary of the Cohen's d effect sizes across all six features. Asterisks indicate those features with a statistically significant difference between the two clusters (as determined by Kolmogorov-Smirnov statistic, D_{KS}, exceeding the critical value $D_{critical}$ = 0.13). Number of molecules per cluster: N_1=240, N_2=622.

We next examined the molecular properties of the two subsets within Cluster 1 to assess whether they
represent distinct regions of chemical space. Each molecule in the dataset is described by six properties
(*RMM, CL$_{int}$, fu, logD,* donor and acceptor *pKa*). Pairwise comparisons of the property distributions for
the two subsets, using two-sample Kolmogorov-Smirnov tests, showed statistically significant
segregation of medium effect or greater for two properties ($D_{KS} > D_{critical}$ for Bonferroni-corrected
confidence level $\alpha = 0.0083$ and Cohen's d > 0.5; **Table S6; Figure S7**). In order of effect size
magnitude, these were *logD* and acceptor *pKa* (**Figure 5C**). Accordingly, off-diagonal molecules were
more likely to be lipophilic and protonated. A similar analysis of Cluster 2 identified three distinguishing
features of medium effect or greater. Off-diagonal molecules in Cluster 2 were characterized by greater
acceptor *pKa* and *fu,* but lower *logD*.

Adding these observations to the fact that acceptor *pKa* globally distinguishes clusters 1 and 2
(Cohen's d = 1.2, $\alpha = 0.0083$; **Figure 6**), we posit that the properties typical of molecules whose *Vss*
predictions are highly discordant between our model and that of Mathew are: (1) high degree of
protonation coupled with (2) high lipophilicity or limited plasma protein binding. Despite their similar
underlying framework, the specific assumptions these two models make about the roles of lipophilicity
and ionization in determining tissue partitioning produce markedly divergent predictions for molecules of
this type. That divergence is thus associated with uncertainty regarding model structure. However, the
greater Wasserstein distances observed for the subgroup of molecules for which our model and Mathew's
make more similar predictions indicates that the entirety of Cluster 1 exhibits greater susceptibility to
parameter uncertainty than predictions made for molecules in Cluster 2.

Subtle changes in model formulation and model calibration markedly impact PK outcome
sensitivity to parameter uncertainty

To explore how drug-specific parameters drive prediction variability in the four different models, we
performed a Sobol' global sensitivity analysis. Sobol' analysis is a variance-based approach that returns
indices apportioning the variance of a model's output between the model's input parameters. In these
analyses, it is common to report the first order and total effect sensitivity indices. These indicate,
respectively, the independent parameter contributions to output variance and the total contributions that
result from adding the effect of parameter interactions. When input parameters exhibit correlations – as
our biophysicochemical properties do (see **Equation 25** for complete correlation matrix) – the
interpretation of the sensitivity indices changes, as the correlations between parameter values act as
another axis of parameter interaction[89–91]. In this case, the sensitivity analysis can yield indices separating
the correlative interactions from the independent interactions.

Using the Monte Carlo estimator for correlated inputs developed by Kucherenko[89], we evaluated the
total correlated sensitivity index, S_i^{TC}, and the total uncorrelated sensitivity index, S_i^{TU}, for each of the
PBPK model's drug input parameters. S_i^{TC} reflects the sum of parameter *i*'s main effect on output
variance (*i.e.*, independent of parameter interactions) and the interactions associated with parameter
correlations. S_i^{TU} reflects the sum of a given parameter's main effect and the correlation-independent
parameter interactions. Given these definitions, S_i^{TC} and S_i^{TU} may be greater than or equal to zero, with
S_i^{TU} tending to zero as $|\rho_{i,\sim i}| \rightarrow 1$ (where, $\sim i$ denotes any parameter other than parameter *i*). In the
absence of parameter correlations, *i.e.*, $|\rho_{i,\sim i}| \rightarrow 0$, S_i^{TC} is reduced to the first order sensitivity index, and
$\sum_i S_i^{TC}$ cannot exceed 1. A corollary conclusion is that sums exceeding 1 must reflect correlation effects.
Details on our implementation are provided in the Methods section and in the code accompanying this
manuscript.

(A)

(B)

Figure 7. Sobol' global sensitivity indices for prediction of human PBPK statistics. Paired bars indicate the total correlated (first order + correlative interactions, left) and total uncorrelated (first order + non-correlative interactions, right) indices, respectively. Results are shown separately for property distributions consistent with: **(A)** Cluster 1. Sample sizes for each model were: $N_{samples}$ = 1,310,720 (calibrated Pearce); $N_{samples}$ = 3,407,870 (uncalibrated Pearce); 11,796,500 (our model); and 12,058,600 (Mathew). **(B)** Cluster 2. Sample sizes for each model were $N_{samples}$ = 65,536 (our model and Pearce models); and 25,690,100 (Mathew), respectively. For cluster 2, the upper limits on the confidence intervals for $S_{Vss,Mathew}$ are 9.2, 8.4, and 5.6 for donor *pKa*, acceptor *pKa*, and *logD*, respectively. The total number of simulations executed for each model's Sobol' analysis was $4 \times N_{samples}$. Sample sizes were chosen based on a convergence analysis (see Methods and **Figures S10 & S11**), and error bars indicate bootstrapped 95% confidence intervals.

We assessed the properties of drugs in the Obach dataset[68] and found statistically significant linear
correlation coefficients among the six PBPK input parameters. Amongst these, the strongest correlation
was between $logD$ and fu ($\rho = -0.56$, p < 0.001), followed closely by $logD$ and donor pKa ($\rho = 0.49$,
p < 0.001). **Figure 7** reports the results of the sensitivity analysis for each of the PBPK models and for
each cluster identified in the agreement analysis.

<u>Impact of model abstractions on parameter attribution for *Vss*</u>

As lipophilicity is a primary descriptor of tissue partitioning across all models, one might expect its
influence on *Vss* to be prominent – perhaps even more so for the subset of chemical space captured by
pseudomolecules in Cluster 1, which tend to be of greater lipophilicity (**Figure S6**). However, our model
is the only one in which $logD$ emerges as the primary driver of *Vss* variance in Cluster 1 (**Figure 7A**).
The influence of $logD$ on the *Vss* predictions of the Mathew model is matched or exceeded by that of
pKa, with a non-trivial contribution from fu. By contrast, for Cluster 1, the Pearce model *Vss* predictions
are most sensitive to fu, and the influence of $logD$ appears particularly small in the calibrated model. As
the Sobol' index evaluated here characterizes variability only *within* the interrogated chemical space, and
it reflects apportionment of variance rather than absolute variance, these results should not be interpreted
as indicating little contribution of $logD$ to *Vss* predictions. Rather, the indices tell us that – given the
shared lipophilic, protonated nature of molecules in Cluster 1 – the primary drivers of *Vss* variation
between molecules shift according to the structure of the PBPK model.

In the case of the Mathew model, this shift exhibits a distinctive qualitative trait, in that the influences
of pKa, fu, and $logD$ are entirely attributed to uncorrelated effects. All other models exhibit some degree
of correlated sensitivity. This is a direct consequence of the different assumptions each model makes in
capturing the role of lipophilicity. Our model and Pearce's represent lipophilicity by the distribution
coefficient, $logD$, which describes the pH-dependent partitioning of neutral and ionized forms of each
species. Mathew instead uses $logP$, which does not reflect the impact of ionization. In keeping with this
choice, the Mathew implementation limits neutral lipid and neutral phospholipid partitioning to the un-
ionized fraction of molecules. These distinct abstractions lead to important differences in the treatment of
phospholipid association. As per **Equation 4**, and as demonstrated in **Figure S8**, neutral phospholipid
association ($log\ Knpl$) increases linearly with $logD$ in our model ($logMA = f(logD)$) and that of Pearce
($logMA = f(logP)$). The Mathew model instead leverages the approximation employed by Rodgers and
Rowland[63] ($Knpl = 0.3P + 0.7$). This yields a relationship between $log\ Knpl$, and input parameter $logD$ that
is non-linear – tending to a constant at low values of $logD$ and then rising steeply as $logD$ approaches
values more typical of drug-like molecules. Consequently, as $logD$ increases, the Mathew estimates for
$log\ Knpl$ grow to be orders of magnitude greater than the $log\ Knpl$ estimates appearing in the other
models.

Whether large estimates for $Knpl$ inflate Kpu depends on how each model describes the interplay
between ionization and lipophilicity. In the Mathew model, neutral phospholipid partitioning only
contributes to predicted Kpu if molecules are not ionized. In Cluster 1 of the synthetic dataset, only 6%
(48/751) of the molecules possess a neutral fraction of > 5%. Thus, for Cluster 1 molecules, the influence
of pKa prevents large estimates for $Knpl$ from impacting Kpu through neutral phospholipids. Instead, the
effect of $Knpl$ will primarily appear via acidic phospholipid (AP) association, which the Mathew model
defines as a characteristic of any molecule possessing a protonated site (*i.e.*, cationic and zwitterionic
fractions), irrespective of net molecular charge. Eighty-five percent (640/751) of protonatable sites within
this cluster of molecules are expected to be highly ionized (defined as having at least 95% of the
population protonated at pH 7.4). Coupled with the fact that Cluster 1 molecules tend to be more

lipophilic than those in Cluster 2 ($logD_{\text{cluster 1}}$ = 3.57, 95% CI:[1.56, 7.72] vs. $logD_{\text{cluster 2}}$ = 0.70, 95%
CI:[−2.41, 5.06]), one expects the Mathew model to yield markedly greater predictions for AP association
than the other models we considered. Our model and that of Pearce adopt Schmitt's correlation to
describe AP association (**Equation 8**). This formulation allows all species to associate with acidic
phospholipids – and to do so with association coefficients that are referenced to $logMA$. By using $logD$ to
predict $logMA$, our model further tempers AP partitioning for ionized species.

These structural choices explain the Sobol' results for prediction of Vss. In the Mathew model, our
sensitivity analysis attributes parameter sensitivity almost entirely to uncorrelated effects across both
clusters. This implies the absence of first order effects, as these would otherwise show up as part of the
correlated sensitivity index. Instead, each parameter's influence emerges through non-correlative
parameter interactions. The strongest contributors were $logD$ and pKa – the same parameters whose
collective impact we identified as jointly determining AP association. While the other models were also
sensitive to these input parameters, the emergent variance observed for Vss predictions was split between
total correlated and total uncorrelated indices. This pattern recurs for Cluster 2 (**Figure 7B**). One practical
implication of the Mathew model's strong parameter interactions is that the variance in the predicted PK
outcomes cannot be reduced by making high-precision predictions for just the most influential of the
drug-specific input parameters. Uncertainty would need to be reduced for all interacting parameters –
thereby increasing the performance demands for data-driven property-predictions.

<u>Model calibration reduces the influence of $logD$ and increases that of fu on dynamic PK statistics, with</u>
<u>notable impact on the relevance of hepatic metabolism</u>

Across all models and both clusters, $Cmax,u$ showed negligible sensitivity to intrinsic clearance
(CL_{int}). This is to be expected for a short IV infusion, where the early distribution dynamics are not
impacted by hepatic clearance, as would be the case for an oral route of administration. The influence of
CL_{int} instead dominates the variance of $AUCu$ and is a pronounced contributor to that of $T_{0.1\mu M}$ – both PK
statistics that reflect timescales long enough to be influenced by hepatic metabolism.

While $Cmax,u$ predictions for molecules in both clusters are largely insensitive to pKa, Mathew
model predictions for molecules in Cluster 1 again exhibit a pronounced, uncorrelated sensitivity to the
ionization parameters and to $logD$. For all other models, $logD$ dominates the prediction variance, and the
similar magnitudes of correlated and uncorrelated S_{logD} indices suggest a primarily first-order influence of
lipophilicity. For Cluster 2, the Mathew model's $S_{i,cmaxu}$ indices mimic those of other models, with
correlated indices exceeding uncorrelated sensitivity and variance influences of similar magnitude across
RMM, fu and $logD$.

A difference between the R&R and Pearce model formulations becomes evident in the influences of
$logD$ and pKa on the prediction of $T_{0.1\mu M}$ for Cluster 1. Both parameters have a stronger influence on the
predictions from the R&R models (This Model and Mathew). Furthermore, in the absence of calibrations,
the Pearce predictions of $T_{0.1\mu M}$ are relatively insensitive to CL_{int}, while the R&R models exhibit a
pronounced uncorrelated sensitivity. By contrast, predictions of $T_{0.1\mu M}$ for molecules in Cluster 2 vary
primarily with CL_{int}, with indices of similar magnitude across all four models. Influences of $logD$, pKa,
and fu are uniformly small.

In addition to inducing CL_{int} sensitivity for $AUCu$ and $T_{0.1\mu M}$ predictions in Cluster 1, Pearce model
calibrations appear to induce sensitivity to fu and reduce the influence of $logD$ (**Figure 8**). This suggests
that the core Pearce model tends to overestimate the impact of lipophilicity on tissue partitioning. That
effect is then corrected by calibration of Kp prior to simulation. Indeed, the Obach compounds whose Kpu
predictions are most impacted are typically highly lipophilic (**Table S7**), and for these molecules, the

tissue most impacted by calibration is adipose (**Table S8**). While reducing $Kpu_{adipose}$ by ~93%, model

calibrations increased Kpu_{liver} by ~42%. Thus, for molecules with high lipophilicity, calibration causes the

simulation to render otherwise-inaccessible compounds available for metabolism by reducing hold-up in

non-metabolic tissues and increasing exposure in hepatic tissue – producing PK dynamics that respond to

CL_{int} variation. For lipophilic molecules, we thus anticipate that model calibration may create a

pronounced qualitative change in PK predictions by 'gate-keeping' the relevance of hepatic metabolism.

Figure 8. Calibration of the Pearce model reduces the influence of *logD* and increases that of *fu* on the variability of most PK outcomes. Bars indicate how Sobol sensitivity indices change upon calibration of the Pearce model. Paired bars indicate the change in total correlated (first order + correlative interactions, left) and total uncorrelated (first order + non-correlative interactions, right) indices, respectively.

Conclusion

This study sheds light on challenges to consider in early drug design workflows where input

parameters are ML-derived, model assumptions may not be validated, uncertainty is considerable, and

decisions may be made without experimental validation. Across experimental datasets, all four PBPK

models considered achieved broadly similar fidelity to observed *Vss* and *Kp*, when evaluated with

measured input parameters. However, once uncertainty was introduced for the pseudomolecules,

clustering of simulation outputs revealed pronounced differences in prediction variance across models and

across chemical space. We found that modeling choices that improved nominal fidelity may render

models inflexible, thereby risking poor extrapolation to novel regions of chemical space. Modeling

choices that invoke ionization state as a filter for lipid and protein association may produce parameter

interactions that degrade prediction fidelity and robustness to uncertainty. Furthermore, models tend to

diverge in their predictions for protonated, lipophilic molecules, which raises questions about whether any

single model will be most credible for decision-making in broad, virtual screens. Our findings do,

however, indicate PBPK modeling strategies that may mitigate (or at least avoid exacerbating) issues

arising from parameter uncertainty.

To limit prediction variability, model abstractions should not binarize the inclusion of specific terms as a function of molecule properties

The two R&R models (our model and that of Mathew) illustrate how distinct assumptions within the same nominal modeling framework can lead to very different outputs. The Mathew model consistently exhibited a bias toward underestimation of Kp and V_{ss} and produced greater prediction errors for zwitterions and bases. Classifying a molecule as cationic in the Mathew model heavily weights acidic phospholipid interactions. Furthermore, by adjusting for lipid association in plasma, the assumptions in the Mathew model effectively exclude plasma and interstitial protein binding for protonated molecules that are also classified as lipophilic and highly bound. Consequently, the Mathew model's predictions are heavily influenced by parameter interactions between $logD$ and pKa. These modeling choices cause large parameter-driven variance in V_{ss} and $T_{0.1\mu M}$ once uncertainty is propagated.

Model calibrations limit prediction flexibility within drug-like chemical space

With respect to Kp, the fidelity of the Pearce model is bolstered by calibrating its outputs to a library of pharmaceutical compounds with measured tissue partitioning. However, this calibration has been reported to reduce prediction flexibility as chemical properties vary – a result we also observe herein. Others have postulated this to be an artefact of model's pharmaceutical-based calibrations being applied to predict outcomes for non-pharmaceutical compounds[92]. Our work suggests that the constraining effect of calibration also limits variability within drug-like chemical space. Alternatively, the calibrations, being based on rat tissue data, may not be appropriate to reflect chemistry-dependent outcome variability in humans. By comparison, our modified model performed comparably to published models, but without the amplified variance observed for the Mathew model or the constrained flexibility associated with the calibrated Pearce model.

Choice of PBPK mathematical abstraction may frustrate attempts to reduce prediction uncertainty by improving QSAR parameter predictions

Our sensitivity analysis clarified how structural assumptions shape the emergence of variance and pointed to inputs whose refinements would most improve prediction robustness. Variability in V_{ss} was driven primarily by $logD$ and pKa, though their influence was expressed differently across models. For Mathew, variance arose almost exclusively from parameter interactions – indicating that prediction confidence cannot be improved by reducing uncertainty in any single input parameter. Clearer avenues for improvement appear for our model and that of Pearce, as reducing error in inputs such as fu or $logD$ would directly reduce variance for all PK outcomes. This analysis highlights priorities for improving property prediction models if ML-informed PBPK workflows are to provide informative assessments in virtual screening.

Limitations

Our analyses included relatively large molecules, which, along with those that are not hepatically metabolized, would violate key model assumptions. Molecular mass did not, however, consistently map to poor V_{ss} predictions. In the Obach dataset, of the 101 compounds with RMM greater than 600, only 16 had prediction errors greater than 10-fold in our model's predictions. Of the 62 total molecules that did have greater than 10-fold prediction errors (26 underpredicted and 36 overpredicted), many are chemotherapeutic compounds that intercalate with DNA (effectively, a target-mediated disposition

factor), antibiotics and natural products that are not metabolized, or compounds that are metabolized by intracellular processes that are not specific to the liver. Others partition extensively into blood cells in a way that is not captured by the partitioning phenomena addressed in our generic model – such as binding to cytosolic carbonic anhydrase[93]. A previously reported example of this latter issue, identified by Rodgers and Rowland[94], is chlorthalidone, for which our model underpredicts Vss by ~8-fold. This differs from the substantial overprediction that would result had we used the commonly employed approach of inferring K_{APL} from blood cell partitioning. In that case, we would have incorrectly attributed the high level of chlorthalidone association with blood cells to the presence of acidic phospholipids. Propagating that error through to Kpu predictions for all tissues would lead to >10-fold overestimation for Vss.

We must bear in mind that the fidelity evaluation herein employs point estimates for both the empirical data and our model predictions. We note that, for some drugs, there are discrepancies between the Vss metrics reported in Obach $et\ al.$[68] and findings elsewhere in the literature – cases where the alternative empirical value is within 10-fold of our model prediction.

Finally, the conclusions presented herein are particular to the IV route of administration simulated. Similar analyses would be required to determine model performance under uncertainty for other routes of administration. For example, CL_{int} is expected to strongly influence $Cmax,u$ for an oral dose, due to first-pass elimination and interplay of absorption kinetics with tissue disposition.

Methods

Table 1 summarizes the distinguishing features of the three *Kp* models. These features and relevant governing equations are elaborated upon in the section that follows.

Table 1. Distinguishing features of the models considered in this work. *MA*: membrane affinity. *Dow*: octanol-water distribution coefficient. *logD*: distribution coefficient. *logP*: partition coefficient. AP: acidic phospholipid. R&R: Rogers & Rowland. *fu*: fraction unbound.

	This model	Pearce (2017)	Mathew (2021)
fu adjustment	Modified Lukacova equation to use D_{ow} for neutral lipid partitioning, membrane affinity (MA) for neutral phospholipids, and to include plasma acidic phospholipids.	Addresses lipid partitioning in plasma using an effective volume fraction of neutral lipid that includes 30% of the neutral phospholipid volume.	Weights lipid partitioning by the fraction of molecules in a neutral state. Adjusts protein binding term to reflect neutral and charged fractions binding to plasma lipoprotein and albumin, respectively.
intracellular pH	Accounts for pH differences in intracellular compartments of different tissues.	Accounts for pH differences in intracellular compartments of different tissues.	pH=7 in intracellular compartment of all tissues.
neutral lipid partitioning	Describes partitioning into neutral lipids using *logD* at relevant pH.	Describes partitioning into neutral lipids using *logD* at relevant pH.	Assumes only neutral drugs partition into neutral lipids with partitioning described by logP.
neutral phospholipid partitioning	Partitioning into neutral phospholipids described using the logMA (membrane affinity) correlation of Yun and Edginton. Employs *logD* in this correlation, to reflect impact of drug ionization.	Partitioning into neutral phospholipids described using the *logMA* (membrane affinity) correlation of Yun and Edginton. Employs *logP* in this correlation.	Only neutral drugs partition into neutral phospholipids, with partition coefficient dependent on *logP* (*viz.* 0.3*P*+0.7).
acidic phospholipid partitioning	Assumes all molecules associate with acidic phospholipids, with coefficient dependent on ionization, as described by Schmitt. Zwitterionic fractions grouped with the cationic fraction in AP partitioning (following the assumption of R&R, which treats protonated sites as dominating AP interactions for zwitterions).	Assumes all molecules associate with acidic phospholipids, with coefficient dependent on ionization, as described by Schmitt. Zwitterionic fraction treated as part of the neutral fraction in AP partitioning.	Assumes drug fractions with cationic sites bind to acidic phospholipids, regardless of overall molecular charge.

interstitial protein binding	Assumes anionic and zwitterionic fractions bind to albumin and neutral drug fractions bind to lipoproteins. Derives corresponding protein binding affinity from fu. Interstitial protein content varies with tissue albumin and lipoprotein levels.	Assumes nonspecific binding to proteins. Derives corresponding protein binding affinity from fu. Interstitial protein content is invariant across tissues.	Assumes all ionized drug fractions bind to albumin and neutral fractions bind to lipoproteins. Derives protein binding affinity from fu.
intracellular protein binding	Assumes nonspecific binding to proteins using an empirical relationship linking protein binding to phospholipid partitioning.	Assumes nonspecific binding to proteins using an empirical relationship linking protein binding to phospholipid partitioning.	No intracellular protein binding, except in blood cells. In blood cells, non-cations assumed to bind to an unspecified protein, with affinity equal to that of cation binding to acidic phospholipids.
adjusted lipid partitioning for adipose	Replaces octanol-water partition coefficient with the vegetable oil-water partition coefficient.	N/A	Replaces octanol-water partition coefficient with the vegetable oil-water partition coefficient.

Model implementation

General equation for R&R-Lukacova models

Equation 1 captures the decomposition of PBPK partition coefficients into contributions from tissue
water, interstitial proteins, and lipid components of each tissue.

$$Kpu = f^{EW} + Ka_{PRp}[PROT,IS]_{eff} + {f_{neutral,p}}/{f_{neutral,IW}}\,(f^{IW} + K_{NL}f^{NL} + K_{NPL}f^{NPL} + K_{APL}f^{APL})$$
[1]

Here, f^i indicates the fractional volume of tissue that is occupied by component i, and f_j indicates the
micro-ionization fraction of drug in charge state j. Plasma is denoted by the subscript p, and Ka_{PRp}
denotes the binding affinity for plasma proteins. We denote tissue components as: extracellular water
(EW), intracellular water (IW), neutral lipids (NL), neutral phospholipids (NPL), acidic phospholipids
(APL), lipoproteins (LPP), albumin (ALB), and intracellular proteins (PROT). Binding to interstitial
proteins is expressed as a function of the effective concentrations of those proteins relative to the protein
content of blood plasma, $[PROT,IS]_{eff}$. We denote the association coefficient for each tissue component as
K_i.

Neutral lipids (plasma and intracellular)

The R&R-Lukacova approach typically employs lipophilicity, quantified by the octanol-water
partition coefficient, $logP_{ow}$, to predict drug partitioning into neutral lipids. These models capture the
consequences of ionization by weighting this term by the fraction of drug molecules expected to remain in
the neutral state. In our model, we instead follow the example of Schmitt, and use the pH-specific
distribution coefficient, $logD_{ow}$, to represent neutral lipid partitioning.

$$K_{NL} \approx D_{ow}$$ [2]

We address the difference in plasma pH and reported differences in intracellular pH across tissues by
adjusting D_{ow} to reflect local states of micro-ionization. Using the Henderson-Hasselbalch equation, we
predict the fraction of molecules in neutral and charged states at the relevant local pH. The neutral
fraction then informs our distribution coefficient estimates, as indicated in **Equation 3**, where α=0.001 [95].

$$D_{ow} \approx P_{ow}[f_{neutral}(1 - \alpha) + \alpha]$$ [3]

Neutral phospholipids

We also apply this local distribution coefficient to predict neutral phospholipid association of
ionizable species. Our approach follows that of Pearce et $al.$[60], who reported improved Kp predictions
when using the membrane affinity correlation of Yun and Edginton[96] to approximate neutral phospholipid
partitioning (**Equation 4**). Herein, we have assessed the further impact of replacing $logP$ with the local
distribution coefficient, $logD$, as a potential approach to account for charge effects.

$$logK_{NPL} \approx logMA \approx 1.294 + 0.304\,logD_{ow}$$ [4]
34

1   where *MA* is the membrane affinity constant. Finally, we note the adjustment made to account for the
2   triglyceride rich composition of neutral lipids in adipose tissue. As with prior models, we replace the
3   octanol-water partition coefficient, P_{ow}, with the vegetable oil-water partition coefficient, P_{vow}, when
4   modeling tissue partitioning for adipose tissue – estimating P_{vow} from P_{ow} as per **Equation 5**.[97] Converting
5    $logP_{vow}$ to $logD_{vow}$, using **Equation 3**, provides the required value for K_{NL} in adipose tissue.

6
$$logP_{vow} = 1.115 * logP_{ow} - 1.35 \tag{5}$$

7   Intracellular proteins

8   The R&R-Lukacova framework does not typically reflect intracellular protein binding[62,63,98]. For our
9   model we add this component to the general equation as:

10
$$\left. f_{neutral,p} \middle/ f_{neutral,IW} \right. K_{prot} f^{PROT} \tag{6}$$

11   with the first term denoting the charge-dependent difference in concentration between unbound drug in
12   the plasma and intracellular water. Following the example of Schmitt[95], we approximate the non-specific
13   protein binding coefficient as a function of the phospholipid partitioning coefficient (**Equation 7**).

14
$$K_{prot} \approx 0.163 + 0.0221 K_{NPL} \tag{7}$$

Acidic phospholipids (plasma and intracellular)

The R&R-Lukacova framework infers acidic phospholipid partition coefficients from empirically
assessed blood-to-plasma ratio and treats the association with acidic phospholipids as a site-specific, 1:1
binding event. Here, we instead adopt the approach of Schmitt and project the affinity for acidic
phospholipids as a perturbation of the affinity for neutral phospholipids. Due to electrostatic effects, the
cationic fraction of drug is expected to augment the apparent membrane association, and the anionic
fraction is expected to reduce that affinity. Based on the correlation proposed by Schmitt[95], Pearce *et al.*[60]
report the corresponding estimate for K_{APL} as:

23
$$K_{APL} \approx K_{NPL}\left(f_{neutral,IW} + 0.05\,f_{anionic,IW} + 20\,f_{cationic,IW}\right) \tag{8}$$

For zwitterions, we adopt the assumption stated by Rodgers and Rowland – namely, that interactions
between protonated sites and acidic phospholipids dominate[62]. Thus, we add the zwitterionic fraction,
$f_{zwitterionic,IW}$ to the cationic term in **Equation 8**. Note that this differs from the approach implemented
in the high-throughput toxicokinetic model Httk, as part of the framework originally reported by Pearce[60].
There, the zwitterionic fraction was combined with the neutral fraction.

Binding to plasma and interstitial proteins

As in the work of Rodgers and Rowland, we use the apparent binding coefficient between plasma
proteins and drug to approximate binding to tissue interstitial proteins. If plasma protein binding is
empirically quantified via the measurement of unbound drug in plasma samples (*fu*), the apparent binding
coefficient can be calculated using **Equation 9**.

34
$$K_{aPRp} \approx \frac{f^{PW}}{1/fu^{-1}} \tag{9}$$

1 where PW indicates plasma water. This equation assumes that the drug sequestration observed in plasma
2 binding assays is dominated by protein interactions.

3 As is common practice, we adjust this value to account for the potential impact of drug partitioning into
4 plasma lipid components. Here, we use the adjusted form suggested by Lukacova *et al.*[98], modified to
5 account separately for neutral lipid, neutral phospholipid, and acidic phospholipid components of plasma.

6
$$fu_{adj} = \frac{f^{PW}}{f^{PW}+K_{aPRp}+K_{NL}f^{NL,p}+K_{NPL}f^{NPL,p}+K_{APL}f^{APL,p}}$$
[10]

Finally, we capture binding to interstitial albumin and lipoproteins by weighting the apparent ratio of
those proteins' contribution against the estimated binding propensity of albumin and lipoproteins in
plasma. As neutral compounds are expected to associate with lipoproteins and anionic components are
expected to interact with albumin[99,100], each protein's contribution to interstitial binding is weighted by
the relevant micro-ionization fraction (**Equation 11**). Here, we have assumed that zwitterions follow the
behavior of anions, interacting with albumin.

13
$$[PROT,IS]_{eff} = f_{neutral,EC}[LPP] + (f_{anionic,EW} + f_{zwitterionic,EW})[ALB]$$
[11]

Prior Models

Pearce model

Our implementation of the model reported by Pearce *et al.* closely follows the approach described in
their 2017 publication and laid out in the Httk toxicokinetic R package.[60,66] The general form of their
tissue partitioning equation is captured in **Equations 12-15**, and key distinguishing assumptions of the
model are detailed in Table 1. The *Kpu* equation has 2 broad components, one term each for interstitial
and intracellular partitioning.

21
$$Kpu = f^{int} K_{int} + \kappa_{IW:PW} f^{cell}K_{cell}$$
[12]

22 where *int* indicates tissue interstitial space, f^{int} and f^{cell} are the fractional volumes of the interstitial and
23 cellular spaces, and K_{int} and K_{cell} are the net partition coefficients for each of these compartments (with
24 respect to plasma water). $\kappa_{IW:PW}$ captures the impact of the ionization gradient that arises from the
25 difference between intra- and extra-cellular pH, as determined by **Equation 13**.

26
$$\kappa_{IW:PW} = \frac{(f_{neutral}+f_{zwitterion}+\alpha(f_{cationic}+f_{anionic}))|_p}{(f_{neutral}+f_{zwitterion}+\alpha(f_{cationic}+f_{anionic}))|_{IW}}$$
[13]

27 The interstitial partition coefficient, K_{int}, reflects drug dissolved in interstitial water and non-specific
28 protein binding as follows.

29
$$K_{int} = f^{w,int} + {f^{prot,int}}\Big/{f^{prot,p}}\left(1\Big/{fu_{adj}} - f^{w,p}\right)$$
[14]

30 where *w* indicates water, $f^{w,int}$ and $f^{prot,int}$ are the fractional volume of water and protein in the interstitial
31 space, and $f^{w,p}$ and $f^{prot,p}$ are the corresponding values for blood plasma. The intracellular partition
32 coefficient, K_{cell}, reflects drug dissolved in intracellular water, along with lipid and protein binding.

$$K_{cell} = f^{w,cell} + K_{NL}f^{NL,cell} + K_{NPL}f^{NPL,cell} + K_{APL}f^{APL,cell} + K_{prot}f^{prot,cell} \tag{15}$$

All superscripts with 'cell' are fractional volumes expressed relative to cell volume. K_{NL} and K_{prot} follow the same assumptions captured in **Equations 2 and 4**, which our model has adopted from that of Pearce. Note that, for all drugs, we used the correlation of Yun and Edginton as a surrogate for predicting membrane affinity, which serves, in turn, as a surrogate for neutral phospholipid partitioning. This differs from the approach reported by Pearce, which employed empirical values for *logMA*, where available. Finally, this model includes a correction to adjust the *fu* measured *in vitro* to account for the role that lipid partitioning is expected to play in augmenting the apparent bound fraction (**Equation 16**).

$$fu_{adj} = \frac{f^{PW}}{K_{NL}\,f^{NL,p}+1/fu} \tag{16}$$

Mathew model

Our implementation of the model reported by Mathew *et al.* follows the governing equations made available in the VBA code included with the supplement to that publication[59]. However, we made two key modifications. Firstly, we corrected the equation used to adjust the unbound fraction of drug in plasma, as per Equation 17.

$$fu_{adj} = \frac{f^{PW}}{f^{PW}+K_a+f_{neutral,p}(K_{NL}f^{NL,p}+K_{NPL}f^{NPL,p})}\,;\,K_a = 1/fu - f^{PW} \tag{17}$$

This addresses a sign error that adjusted *fu* upward instead of decreasing its value, as is necessary to account for postulated lipid partitioning. This change obviated the need for the upper limit on *fu_adj* that the authors imposed. Secondly, we sought to remove the model's reliance on blood-to-plasma ratio as an input parameter. Our motivation for this is two-fold: (1) the molecule datasets used in this study do not include empirical values for blood-to-plasma concentration ratio (*B2P*); and (2) this parameter proves challenging to predict via QSAR models[101], and its uncertainty has pronounced first-order impacts on the prediction of human volume of distribution[102]. In lieu of using *B2P* to infer association coefficients for acidic phospholipids, we adopted the Schmitt-Pearce approach and predicted K_{APL} from the charge-based modification of K_{NPL}. In Table 1, we further highlight distinguishing assumptions of the Mathew model.

PBPK simulations

Our 14-compartment PBPK models (**Figure S9**) assume perfusion-limited disposition following a 15 mg/kg IV dose simulated over 1 minute. We simulated disposition dynamics over a time span of 48 hours. For elimination, the models assume hepatic metabolism (via the well-stirred approximation) and passive renal filtration. As such, the models neglect active transport and permeability limitations. They also do not account for target-mediated disposition or non-linear PK phenomena. For example, by using first order kinetics to describe hepatic clearance, the models neglect any saturation of metabolic capacity.

The dynamic PK statistics selected for analysis were the area under the unbound concentration-time curve ($AUCu_{0-\infty}$), maximum unbound concentration (*Cmax,u*), and time for which unbound concentration exceeds a threshold of 0.1 μM ($T_{0.1\mu M}$). In all cases, the analysis focused on the predicted unbound concentration in the muscle intracellular space.

Definition of ion classes

When classifying molecules, we defined zwitterions as any compound possessing both protonatable
and de-protonatable sites, irrespective of the ionization state at physiological pH. Bases and acids were
those with only an acceptor site and those with only a donor site, respectively. Compounds without
ionizable sites were classified as neutral. For our synthetic dataset of pseudomolecules, we defined sites
to be ionizable if $pKa_{donor} < 9.4$ or $pKa_{acceptor} > 5.4$ – each of which maps to 1% ionization at pH of 7.4.

Evaluation of model performance

To assess the prediction fidelity, relative performance, and uncertainty associated with each model,
we predicted Kp and Vss quantities for which empirical data have been reported. Kp represents the ratio of
tissue concentration to that of blood plasma, and is related to Kpu by:

$$Kp = Kpu\ fu_{adj} \qquad [18]$$

Vss, the steady-state volume of distribution, reflects the extent to which a drug is expected to distribute
beyond the blood plasma into tissue matrices. Estimation of Vss based on mechanistic predictions of Kpu
is carried out using **Equation 19.**

$$V_{ss} = V_p + K_{blood\ cells}V_{blood\ cells} + \sum_{i=1}^{N} Kp_iV_i \qquad [19]$$

where V_i represents tissue volume, and the sum runs over all tissues.

17
We used the following statistics to assess model fidelity to data:

• fraction of drugs with PK predictions within 2-fold, 3-fold, and 10-fold error

$$fold\ error = 10^{|log(pred_i/obs_i)|} \qquad [20]$$

• root mean squared error of log-transformed values

$$RMSLE = \sqrt{\frac{1}{N}\sum_i \left(log\left(\frac{pred_i+1}{obs_i+1} \right) \right)^2} \qquad [21]$$

• average fold error

$$AFE = 10^{\frac{1}{N}\sum_i log(pred_i/obs_i)} \qquad [22]$$

• concordance correlation coefficient (of log-transformed values).

$$CCC = \frac{2\sigma_{pred,obs}}{\sigma_{pred}^2+\sigma_{obs}^2+(\mu_{pred}-\mu_{obs})^2} \qquad [23]$$

Monte Carlo simulations

To evaluate the local variance of model predictions subject to parameter uncertainty, we generated
pseudomolecules from the Obach dataset by correlated sampling of the key biophysicochemical input
parameters RMM, fu, $logD$, pKa, and CL_{int}. To achieve this, we first assumed a uniform input distribution

for each parameter and then imposed correlation by applying a Gaussian copula derived from the property
distributions reflected in the Obach dataset (using MATLAB function *copulafit*). We thereby generated a
dataset of 10^4 pseudomolecules, each defined by *RMM*, *fu*, *logD*, *pKa*, and *CL$_{int}$*.

4   For each pseudomolecule, we then perturbed its properties by sampling from a distribution with mean
5 equal to that of the molecule and mean absolute error matched to the predictive accuracy of typical QSAR
6 models[56–58]. **Table 2** reports the corresponding distribution types and typical errors.

Table 2. Distribution types and mean absolute errors used for Monte Carlo sampling of input parameters.

	fu	*logD*	*pKa*	*log(CL$_{int}$)*
sampling distribution	beta	normal	beta (scaled to [0, 14])	normal
mean absolute error	0.075	0.467	0.42	0.25

7   To evaluate the effects of parameter uncertainty, we simulated 1,000 Monte Carlo realizations for
each of 10^4 pseudomolecules via each of the four PBPK models of interest – giving a total of 4×10^7
simulations. Each realization predicted four PK outcomes: steady-state volume of distribution (*Vss*),
maximum unbound concentration (*Cmax,u*), area under the unbound concentration-time curve (*AUCu*),
and the duration of time for which concentration exceeded 0.1 μM ($T_{0.1\mu M}$). The latter three were
generated for intracellular concentrations in the tissue of interest.

Pseudocode for generation of synthetic pseudomolecules and parameter perturbations for
Monte Carlo simulations

Algorithm 1 Generation of Synthetic Pseudomolecules

1: **Goal:** Define the synthetic dataset pseudomolecules with comparable properties to the joint distribution $\boldsymbol{\theta} = [\mathrm{p}K_a^{\mathrm{acceptor}}, \mathrm{p}K_a^{\mathrm{donor}}, f_u, \log D, \log(\mathrm{CL_{int}}), \mathrm{RMM}]$ of 862 drugs in the Obach dataset.

2: Determine correlation coefficient ρ_{ij} between model parameters θ_i and θ_j.

3: Sample $\mathcal{X} = \{x_1, x_2, \ldots, x_N\}$ with $N = 10^4$ from the Gaussian copula $\mathcal{C}_{\rho_{ij}}$ for the parameters $\boldsymbol{\theta}$.

4: Define typical errors ε for each parameter.

5: **for** each $x_i \in \mathcal{X}$ **do**

6:  For $\log D$ and $\log(\mathrm{CL_{int}})$, find σ^2 of the Gaussian distribution such that the expected absolute deviation matches QSAR model fidelity for x_i.

7:  For f_u, $\mathrm{p}K_a^{\mathrm{acceptor}}$, and $\mathrm{p}K_a^{\mathrm{donor}}$, find α, β of the Beta distribution such that the expected absolute deviation matches QSAR model fidelity for x_i.

8:  Draw $R = 10^3$ samples from:

  • $f_u, \mathrm{p}K_a^{\mathrm{acceptor}}, \mathrm{p}K_a^{\mathrm{donor}} \sim \mathrm{Beta}(\alpha, \beta)$

  • $\log D, \log(\mathrm{CL_{int}}) \sim \mathcal{N}(x_i, \sigma^2)$

9: **end for**

10: Resulting in a total of $R = 10^3$ samples for each of the N pseudomolecules.

Model agreement analysis

The Monte Carlo simulations yielded distributions of the four predicted PK outcomes for each
molecule and each PBPK model. We used these distributions to assess the extent to which models
generate predictions that deviate under parameter uncertainty. To this end, we quantified the model-to-
model Wasserstein distance between the predicted distributions for a given PK outcome. For each
molecule ($N_{pseudomolecules} = 10^4$), given four models ($N_{models} = 4$) and four PK outcomes ($N_{PKoutcomes} = 4$), this
yielded 24 Wasserstein distance features (as per **Equation 24** and **Figure S3**).

8
$$N_{features} = \binom{N_{models}}{2} N_{PK\ outcomes} \tag{24}$$

We then used these 24 features to group molecules into consensus clusters based on model agreement
patterns. The PK preprocessing applied before distance calculation and the feature transformation applied
prior to clustering are described in detail in **Algorithm 2**. Also detailed is our approach to defining robust
cluster assignments. We performed 1000 K-means clustering attempts, noting with each attempt which
pairs of molecules were assigned to the same group. By tallying co-clustering events, we were able to
identify groups of molecules that clustered together in at least 95% of the clustering attempts. This
approach removes any molecule that does not meet the 95% frequency threshold for stably clustering with
other molecules. For this dataset, however, we found that all molecules could robustly be assigned to a
cluster.

In addition to our whitened/PCA-transformed analysis, we evaluated clusters defined by omitting the
preprocessing step (*i.e.*, calculating Wasserstein distances based on raw PK data) and/or by applying min-
max or principal component feature transformations. The final number of clusters (2 or 3) – and their size
– varied with the choice of pre-processing and feature transformation approach (see adjusted Rand
indices, **Table S5**). We arrived at the decision to employ whitening and PCA transformation by
examining the silhouette statistic distributions resulting from each approach (**Table S5**). We also
considered the potential of whitening decorrelation to partially segregate global and local influences on
tissue exposure (*i.e.*, bulk changes in exposure to all body tissues that might drive increases in *Vss* vs.
local changes in muscle intracellular exposure that are driven by the interplay of muscle perfusion and
tissue composition).

Pseudocode for whitening transformation and K-means clustering

---

Algorithm 2 Model agreement analysis: PK preprocessing and feature scaling

1: **Goal:** Quantify PBPK model agreement under uncertainty and cluster molecules by agreement patterns.
2: **Input:** Monte Carlo predictions $\mathcal{M} \in \mathbb{R}^{R \times M \times N \times O}$ where $R = 10^3$ realizations, $M = 4$ models, $N = 10^4$ pseudomolecules, $O = 4$ PK outcomes; Nattempts $= 1000$ (clustering attempts); $K_{\max} = 6$ (maximum number of clusters evaluated); $F = 0.95$ (co-assignment frequency cutoff); fixed seed list (one seed per attempt).
3: **Output:** Consensus cluster labels and co-assignment matrix $\mathbf{A}$.
4: **PK preprocessing:** For each PK outcome, apply a base-10 log transform to all Monte Carlo realizations in $\mathcal{M}$, then apply whitening using *pooled covariance* across the M models.
5: Compute 1-Wasserstein distances between all $\binom{M}{2}$ model pairs for each outcome and molecule.
6: Concatenate distances to form the agreement-feature matrix $\mathbf{D} \in \mathbb{R}^{N \times 24}$ (6 model pairs $\times$ 4 outcomes).
7: **Feature transformation:** Apply a feature-wise PCA transformation to $\mathbf{D}$ to obtain $\mathbf{X}_{\text{scaled}} \in \mathbb{R}^{N \times 24}$.
8: **for** $q = 1, \ldots, \text{Nattempts}$ **do**
9:      Set RNG using the q-th seed from the fixed seed list.
10:      Select K_q via silhouette evaluation over $K = 2, \ldots, K_{\max}$.
11:      Run k-means on $\mathbf{X}_{\text{scaled}}$ with K_q clusters and store labels $\mathbf{c}^{(q)}$.
12: **end for**
13: Compute $\mathbf{A} \in [0, 1]^{N \times N}$, where for $i \neq j$,

$$\mathbf{A}_{ij} = \frac{1}{\text{Nattempts}} \sum_{q=1}^{\text{Nattempts}} \mathbb{I}\left[c_i^{(q)} = c_j^{(q)}\right].$$

14: Define robust groups using $\mathbf{A}_{ij} \geq F$ and assign consensus cluster IDs.

Obach molecule discordance analysis

To connect the cluster patterns from the pseudomolecule analysis to real drug space and distinguish
structure-driven disagreement from the effects of parameter uncertainty, we analyzed 862 molecules from
the Obach dataset. We performed Monte Carlo simulations for these molecules and computed the 24-
element Wasserstein distance features required to map these real molecules to the feature-space defined
by our synthetic dataset. This required whitening and PCA transformations using the same transformation
matrices obtained in our pseudomolecule analysis. We then assigned molecules to clusters by identifying
which cluster centroid was closest to each Obach molecule's feature vector (as defined by Euclidean
distance).

Statistical analysis of sub-group properties

After mapping molecules to clusters, we used two-sample Kolmogorov-Smirnov (KS) tests to assess
whether these groups corresponded to distinct sub-regions of chemical space. We applied this analysis to
the six biophysicochemical properties of interest across the two clusters, using a Bonferroni correction to
adjust the chosen confidence level of $\alpha = 0.05$ for these multiple comparisons. For all comparisons, we
also quantified the Cohen's d effect size associated with property differences between groups. A similar
workflow was used to determine any differences between cluster sub-groups corresponding to $\leq$ 2-fold
and >2-fold differences in *Vss* values predicted by our model and that of Mathew.

<u>Sobol sensitivity analysis</u>

We performed variance-based global sensitivity analyses using the approach described by
Kucherenko *et al.* for correlated input parameters[89]. The statistically significant correlation coefficients
for molecules in our chosen subset of 862 molecules from the Obach dataset are presented as **Equation**
25, with the range of drug property values indicated in **Table 3**.

$$\rho = \begin{bmatrix} RMM & pKa,d & pKa,a & fu & log(CLint) & logD \\ 1 & -0.15 & 0 & -0.20 & 0 & 0.16 \\ -0.15 & 1 & 0.25 & 0 & 0.29 & 0.49 \\ 0 & 0.25 & 1 & 0.10 & 0.099 & 0 \\ -0.20 & 0 & 0.10 & 1 & -0.33 & -0.56 \\ 0 & 0.29 & 0.099 & -0.33 & 1 & 0.47 \\ 0.16 & 0.49 & 0 & -0.56 & 0.47 & 1 \end{bmatrix} \qquad [25]$$

Table 3. 95% interval of drug property values for selected molecules in the Obach dataset

	RMM	pKa, donor	pKa, acceptor	fu	log(CLint [L/kg])	logD
lower bound	153	0.713	0	0.002	-2.81	-2.20
upper bound	1160	14	11.0	1	3	5.03

Convergence analysis

For the Sobol' analysis, we assumed a uniform input distribution for each parameter and imposed
correlation by applying a Gaussian copula derived from the property distributions reflected in the Obach
dataset (using MATLAB function *copulafit*). We confirmed convergence of the Monte Carlo estimator by
assessing the 95% interval of sensitivity indices estimated from 100 bootstrapped samples. As a
convergence statistic[103,104], we monitored the largest 95% estimation interval characterizing predicted PK
statistic *j*, observed over all *n* parameters, as per **Equation 26**.

$$convergence\ statistic, CI_j \ = \ \max_{1 \leq i \leq n} \left(P_{0.95}(S_{ij}) \right) \qquad [26]$$

In most cases, we evaluated sufficient samples to reduce this interval to << 0.1 (**Figures S10 & S11**). A
notable exception is the case of *Vss* prediction by the Mathew model, where the Monte Carlo estimator
had not converged after running $\sim 2.6 \times 10^7$ samples (requiring $> 10^8$ simulations).

To validate that apparently influential parameters could be distinguished statistically from those of
negligible influence, we implemented a dummy model output to act as a control. This output's value was

set equal to a uniform random input parameter (*i.e.*, a parameter that did not feature in the PBPK model

and was not correlated with any other parameter). When defined this way, the dummy output acted as a

positive control for true, first order sensitivity to one parameter ($S_i^{TC} = S_i^{TU} = 1$) and a negative control

reflecting true insensitivity to all other parameters ($S_{\sim i}^{TC} = S_{\sim i}^{TU} = 0$). A parameter was then considered to

be influential if the median of its bootstrapped index estimates differed from that of the negative control

(*i.e.*, medians were determined to be statistically distinct by a Wilcoxon rank-sum test at confidence level

of $\alpha = 0.05$).

Programming languages and code availability

We performed all simulations and analyses in MATLAB (The MathWorks Inc., Natick,

Massachusetts). Differences between distributions were computed using the Wasserstein distance metric,

as implemented in the *ws_distance* function[105], and covariance normalization was performed using ZCA

whitening as implemented in the *whiten* function[106]. Differences in clustering outputs across

preprocessing and feature-standardization approaches were quantified using an adjusted Rand index,

computed using function *randindex*[107]. Seeds for random number generation are detailed in **Table S9**.

Details of our computing environments appear in **Table 4**. Data and code used to perform the analyses for

this paper are hosted at https://github.com/MADBioLab/PBPKPredictionVariability.

Table 4. Software and computing environment details

Operating system	Laptops: Windows 11 Version 25H2. Build 26200.8037	Compute nodes on UTK ISAAC-NG HPC cluster: RedHat Enterprise Linux 8.10
MATLAB version	9.14.0.2206163 (R2023a) 25.1.0.2973910 (R2025a)	9.14.0.2306882 (R2023a)
CPU	12th Gen Intel® Core™ i7-12700H @2.30 GHz ×14 cores.	AMD Bergamo EPYC™ 9734 @2.2 GHz ×112 cores.
RAM	32GB RAM	1TB RAM

Funding statement

18   This study was partially funded by the Laboratory Directed Research and Development Program of

19   Oak Ridge National Laboratory, managed by UT-Battelle, LLC, for the U. S. Department of Energy

20   (Award 10493). BSA and MF acknowledge Institutional Start-up funding from the University of

21   Tennessee. The funders played no role in study design, data collection, analysis and interpretation of data,

22   or the writing of this manuscript.

Acknowledgements

24   A portion of the computation for this work was performed on the University of Tennessee

25   Infrastructure for Scientific Applications and Advanced Computing (ISAAC) computational resources.

26   BSA acknowledges the Accelerating Therapeutics for Opportunities in Medicine (ATOM) Consortium.

27   The authors would like to thank MADBio Lab member Ghizelle Abarro for her comments on the

28   manuscript.

Author Contributions (CRediT Taxonomy).

Investigation: BSA, MF

Conceptualization: BSA
Funding Acquisition: BSA
Writing-original draft: BSA, MF, ZRF
Writing-review and editing: BSA, MF, DTF, ZRF
Methodology: BSA, MF, DTF
Software: BSA, MF, DTF
Formal Analysis: BSA, MF, ZRF
8
9 **Competing interests.** The authors have no competing interest to declare.
10
11 **Materials & Correspondence:** Belinda S Akpa

References

1. Qureshi, R. *et al.* AI in drug discovery and its clinical relevance. *Heliyon* **9**, e17575 (2023).

2. Narayanan, R. R., Durga, N. & Nagalakshmi, S. Impact of Artificial Intelligence (AI) on Drug Discovery and Product Development. *Indian J. Pharm. Educ. Res.* **56**, s387–s397 (2022).

3. Dara, S., Dhamercherla, S., Jadav, S. S., Babu, C. M. & Ahsan, M. J. Machine Learning in Drug Discovery: A Review. *Artif. Intell. Rev.* **55**, 1947–1999 (2022).

4. Schneider, P. *et al.* Rethinking drug design in the artificial intelligence era. *Nat. Rev. Drug Discov.* **19**, 353–364 (2020).

5. Dermawan, D. & Alotaiq, N. From Lab to Clinic: How Artificial Intelligence (AI) Is Reshaping Drug Discovery Timelines and Industry Outcomes. *Pharmaceuticals* **18**, 981 (2025).

6. Vamathevan, J. *et al.* Applications of machine learning in drug discovery and development. *Nat. Rev. Drug Discov.* **18**, 463–477 (2019).

7. Dash, D. K., Pattnaik, S. & Namdeo, A. The role of artificial intelligence in drug development: enhancing pharmaceutical chemistry through machine learning and predictive modeling. *Drug Dev. Ind. Pharm.* **51**, 1430–1438 (2025).

8. Staszak, M. *et al.* Machine learning in drug design: Use of artificial intelligence to explore the chemical structure–biological activity relationship. *WIREs Comput. Mol. Sci.* **12**, e1568 (2022).

9. Moshkov, N. *et al.* Predicting compound activity from phenotypic profiles and chemical structures. *Nat. Commun.* **14**, 1967 (2023).

10. Rao, M. S. *et al.* Novel Computational Approach to Predict Off-Target Interactions for Small Molecules. *Front. Big Data* **2**, (2019).

11. Rao, M., McDuffie, E. & Sachs, C. Artificial Intelligence/Machine Learning-Driven Small Molecule Repurposing via Off-Target Prediction and Transcriptomics. *Toxics* **11**, 875 (2023).

12. Liu, J. *et al.* In silico off-target profiling for enhanced drug safety assessment. *Acta Pharm. Sin. B* **14**, 2927–2941 (2024).

13. Basile, A. O., Yahi, A. & Tatonetti, N. P. Artificial Intelligence for Drug Toxicity and Safety. *Trends Pharmacol. Sci.* **40**, 624–635 (2019).

14. Rao, M. *et al.* AI/ML Models to Predict the Severity of Drug-Induced Liver Injury for Small Molecules. *Chem. Res. Toxicol.* **36**, 1129–1139 (2023).

15. Yang, K. *et al.* Analyzing Learned Molecular Representations for Property Prediction. *J. Chem. Inf. Model.* **59**, 3370–3388 (2019).

16. Fooladi, H., Vu, T. N. L., Mathea, M. & Kirchmair, J. Evaluating Machine Learning Models for Molecular Property Prediction: Performance and Robustness on Out-of-Distribution Data. *J. Chem. Inf. Model.* **65**, 9871–9891 (2025).

17. Chen, E. P., Dutta, S., Ho, M.-H. & DeMartino, M. P. Model-Based Virtual PK/PD Exploration and Machine Learning Approach to Define PK Drivers in Early Drug Discovery. *J. Med. Chem.* **67**, 3727–3740 (2024).

18. Pähler, A. *et al.* The "five-star matrix" for patient-centric drug discovery and development in neuroscience. *Neuron* **113**, 2885–2909 (2025).

19. McGinnity, D. F., Meneyrol, J., Boldron, C. & Johnstone, C. Every Compound a Candidate: experience-led risk-taking approaches to accelerate small-molecule drug discovery. *Drug Discov. Today* **30**, 104354 (2025).

20. Liu, G. *et al.* GraphDTI: A robust deep learning predictor of drug-target interactions from multiple heterogeneous data. *J. Cheminformatics* **13**, 58 (2021).

21. Zhou, Y. *et al.* ToxMPNN: A deep learning model for small molecule toxicity prediction. *J. Appl. Toxicol.* **44**, 953–964 (2024).

22. Francoeur, P. G. *et al.* Three-Dimensional Convolutional Neural Networks and a Cross-Docked Data Set for Structure-Based Drug Design. *J. Chem. Inf. Model.* **60**, 4200–4215 (2020).

23. McNutt, A. T. *et al.* GNINA 1.0: molecular docking with deep learning. *J. Cheminformatics* **13**, 43 (2021).

24. Chen, E. P. *et al.* Applications of Model-Based Target Pharmacology Assessment in Defining Drug Design and DMPK Strategies: GSK Experiences. *J. Med. Chem.* **65**, 6926–6939 (2022).

25. Chen, E. P., Bondi, R. W. & Michalski, P. J. Model-based Target Pharmacology Assessment (mTPA): An Approach Using PBPK/PD Modeling and Machine Learning to Design Medicinal Chemistry and DMPK Strategies in Early Drug Discovery. *J. Med. Chem.* **64**, 3185–3196 (2021).

26. Kuepfer, L. *et al.* Applied Concepts in PBPK Modeling: How to Build a PBPK/PD Model. *CPT Pharmacomet. Syst. Pharmacol.* **5**, 516–531 (2016).

27. Poulin, P. *et al.* PHRMA CPCDC initiative on predictive models of human pharmacokinetics, part 5: prediction of plasma concentration-time profiles in human by using the physiologically-based pharmacokinetic modeling approach. *J. Pharm. Sci.* **100**, 4127–4157 (2011).

28. Rowland, M., Peck, C. & Tucker, G. Physiologically-based pharmacokinetics in drug development and regulatory science. *Annu. Rev. Pharmacol. Toxicol.* **51**, 45–73 (2011).

29. Jones, H. & Rowland-Yeo, K. Basic Concepts in Physiologically Based Pharmacokinetic Modeling in Drug Discovery and Development. *CPT Pharmacomet. Syst. Pharmacol.* **2**, 63 (2013).

30. Zhuang, X. & Lu, C. PBPK modeling and simulation in drug research and development. *Acta Pharm. Sin. B* **6**, 430–440 (2016).

31. Edginton, A. N., Theil, F.-P., Schmitt, W. & Willmann, S. Whole body physiologically-based pharmacokinetic models: their use in clinical drug development. *Expert Opin. Drug Metab. Toxicol.* **4**, 1143–1152 (2008).

32. Shardlow, C. E. *et al.* Impact of physiologically based pharmacokinetic modeling and simulation in drug development. *Drug Metab. Dispos. Biol. Fate Chem.* **41**, 1994–2003 (2013).

33. Basic Concepts in Physiologically Based Pharmacokinetic Modeling in Drug Discovery and Development - PMC. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3828005/.

34. Rowland Yeo, K., Gil Berglund, E. & Chen, Y. Dose Optimization Informed by PBPK Modeling: State-of-the Art and Future. *Clin. Pharmacol. Ther.* **116**, 563–576 (2024).

35. Zhang, X. *et al.* Application of PBPK Modeling and Simulation for Regulatory Decision Making and Its Impact on US Prescribing Information: An Update on the 2018-2019 Submissions to the US FDA's Office of Clinical Pharmacology. *J. Clin. Pharmacol.* **60**, S160–S178 (2020).

36. Perry, C., Davis, G., Conner, T. M. & Zhang, T. Utilization of Physiologically Based Pharmacokinetic Modeling in Clinical Pharmacology and Therapeutics: an Overview. *Curr. Pharmacol. Rep.* **6**, 71–84 (2020).

37. Franchetti, Y. & Nolin, T. D. Dose Optimization in Kidney Disease: Opportunities for PBPK Modeling and Simulation. *J. Clin. Pharmacol.* **60**, S36–S51 (2020).

38. Templeton, I. E., Jones, N. S. & Musib, L. Pediatric Dose Selection and Utility of PBPK in Determining Dose. *AAPS J.* **20**, 31 (2018).

39. Moj, D. *et al.* Clarithromycin, Midazolam, and Digoxin: Application of PBPK Modeling to Gain New Insights into Drug–Drug Interactions and Co-medication Regimens. *AAPS J.* **19**, 298–312 (2017).

40. Chou, W.-C. & Lin, Z. Machine learning and artificial intelligence in physiologically based pharmacokinetic modeling. *Toxicol. Sci. Off. J. Soc. Toxicol.* **191**, 1–14 (2023).

41. Habiballah, S. & Reisfeld, B. Adapting physiologically-based pharmacokinetic models for machine learning applications. *Sci. Rep.* **13**, 14934 (2023).

42. Gruber, A. *et al.* Prediction of Human Pharmacokinetics From Chemical Structure: Combining Mechanistic Modeling with Machine Learning. *J. Pharm. Sci.* **113**, 55–63 (2024).

43. Mavroudis, P. D., Teutonico, D., Abos, A. & Pillai, N. Application of machine learning in combination with mechanistic modeling to predict plasma exposure of small molecules. *Front. Syst. Biol.* **3**, (2023).

44. Wu, K. *et al.* Predicting pharmacodynamic effects through early drug discovery with artificial intelligence-physiologically based pharmacokinetic (AI-PBPK) modelling. *Front. Pharmacol.* **15**, (2024).

45. Wang, W. *et al.* An Integrated AI-PBPK Platform for Predicting Drug In Vivo Fate and Tissue Distribution in Human and Inter-Species Extrapolation. *Clin. Pharmacol. Ther.* **n/a**,.

46. Li, Y. *et al.* A Combination of Machine Learning and PBPK Modeling Approach for Pharmacokinetics Prediction of Small Molecules in Humans. *Pharm. Res.* **41**, 1369–1379 (2024).

47. Naga, D., Parrott, N., Ecker, G. F. & Olivares-Morales, A. Evaluation of the Success of High-Throughput Physiologically Based Pharmacokinetic (HT-PBPK) Modeling Predictions to Inform Early Drug Discovery. *Mol. Pharm.* **19**, 2203–2216 (2022).

48. Parrott, N., Paquereau, N., Coassolo, P. & Lavé, T. An Evaluation of the Utility of Physiologically Based Models of Pharmacokinetics in Early Drug Discovery. *J. Pharm. Sci.* **94**, 2327–2343 (2005).

49. Deepika, D. & Kumar, V. The Role of 'Physiologically Based Pharmacokinetic Model (PBPK)' New Approach Methodology (NAM) in Pharmaceuticals and Environmental Chemical Risk Assessment. *Int. J. Environ. Res. Public. Health* **20**, 3473 (2023).

50. Ni, M. *et al.* IVIVE-PBPK based new approach methodology for addressing early life toxicity induced by Bisphenol A. *Environ. Res.* **240**, 117343 (2024).

51. Korzekwa, K. & Nagar, S. On the Nature of Physiologically-Based Pharmacokinetic Models -A Priori or A Posteriori? Mechanistic or Empirical? *Pharm. Res.* **34**, 529–534 (2017).

52. Miller, N. A., Reddy, M. B., Heikkinen, A. T., Lukacova, V. & Parrott, N. Physiologically Based Pharmacokinetic Modelling for First-In-Human Predictions: An Updated Model Building Strategy Illustrated with Challenging Industry Case Studies. *Clin. Pharmacokinet.* **58**, 727–746 (2019).

53. Korzekwa, K., Radice, C. & Nagar, S. A permeability- and perfusion-based PBPK model for improved prediction of concentration-time profiles. *Clin. Transl. Sci.* **15**, 2035–2052 (2022).

54. Cherkasov, A. *et al.* QSAR Modeling: Where Have You Been? Where Are You Going To? *J. Med. Chem.* **57**, 4977–5010 (2014).

55. Neves, B. J. *et al.* QSAR-Based Virtual Screening: Advances and Applications in Drug Discovery. *Front. Pharmacol.* **9**, (2018).

56. Johnston, R. C. *et al.* Epik: pKa and Protonation State Prediction through Machine Learning. *J. Chem. Theory Comput.* **19**, 2380–2388 (2023).

57. Rodríguez-Pérez, R., Trunzer, M., Schneider, N., Faller, B. & Gerebtzoff, G. Multispecies Machine Learning Predictions of In Vitro Intrinsic Clearance with Uncertainty Quantification Analyses. *Mol. Pharm.* **20**, 383–394 (2023).

58. Huang, K. *et al.* Therapeutics Data Commons: Machine Learning Datasets and Tasks for Drug Discovery and Development. *Proc. Neural Inf. Process. Syst. Track Datasets Benchmarks* **1**, (2021).

59. Mathew, S. *et al.* Evaluation of Prediction Accuracy for Volume of Distribution in Rat and Human Using In Vitro, In Vivo, PBPK and QSAR Methods. *J. Pharm. Sci.* **110**, 1799–1823 (2021).

60. Pearce, R. G., Setzer, R. W., Davis, J. L. & Wambaugh, J. F. Evaluation and calibration of high-throughput predictions of chemical distribution to tissues. *J. Pharmacokinet. Pharmacodyn.* **44**, 549–565 (2017).

61. Poulin, P. & Theil, F.-P. Prediction of pharmacokinetics prior to in vivo studies. 1. Mechanism-based prediction of volume of distribution. *J. Pharm. Sci.* **91**, 129–156 (2002).

62. Rodgers, T. & Rowland, M. Physiologically based pharmacokinetic modelling 2: predicting the tissue distribution of acids, very weak bases, neutrals and zwitterions. *J. Pharm. Sci.* **95**, 1238–1257 (2006).

63. Rodgers, T., Leahy, D. & Rowland, M. Physiologically based pharmacokinetic modeling 1: predicting the tissue distribution of moderate-to-strong bases. *J. Pharm. Sci.* **94**, 1259–1276 (2005).

64. Berezhkovskiy, L. M. Volume of distribution at steady state for a linear pharmacokinetic system with peripheral elimination. *J. Pharm. Sci.* **93**, 1628–1640 (2004).

65. Peyret, T., Poulin, P. & Krishnan, K. A unified algorithm for predicting partition coefficients for PBPK modeling of drugs and environmental chemicals. *Toxicol. Appl. Pharmacol.* **249**, 197–207 (2010).

66. Pearce, R. G., Setzer, R. W., Strope, C. L., Wambaugh, J. F. & Sipes, N. S. httk: R Package for High-Throughput Toxicokinetics. *J. Stat. Softw.* **79**, 1–26 (2017).

67. Edginton, A. N., Schmitt, W. & Willmann, S. Development and evaluation of a generic physiologically based pharmacokinetic model for children. *Clin. Pharmacokinet.* **45**, 1013–1034 (2006).

68. Lombardo, F., Berellini, G. & Obach, R. S. Trend Analysis of a Database of Intravenous Pharmacokinetic Parameters in Humans for 1352 Drug Compounds. *Drug Metab. Dispos. Biol. Fate Chem.* **46**, 1466–1477 (2018).

69. Sato, S., Matsumiya, K., Tohyama, K. & Kosugi, Y. Translational CNS Steady-State Drug Disposition Model in Rats, Monkeys, and Humans for Quantitative Prediction of Brain-to-Plasma and Cerebrospinal Fluid-to-Plasma Unbound Concentration Ratios. *AAPS J.* **23**, 81 (2021).

70. Pfeifer, N. D., Harris, K. B., Yan, G. Z. & Brouwer, K. L. R. Determination of Intracellular Unbound Concentrations and Subcellular Localization of Drugs in Rat Sandwich-Cultured Hepatocytes Compared with Liver Tissue. *Drug Metab. Dispos.* **41**, 1949–1956 (2013).

71. Chu, X. *et al.* Intracellular Drug Concentrations and Transporters: Measurement, Modeling, and Implications for the Liver. *Clin. Pharmacol. Ther.* **94**, 126–141 (2013).

72. Riede, J., Camenisch, G., Huwyler, J. & Poller, B. Current In Vitro Methods to Determine Hepatic Kp uu : A Comparison of Their Usefulness and Limitations. *J. Pharm. Sci.* **106**, 2805–2814 (2017).

73. Brown, R. P., Delp, M. D., Lindstedt, S. L., Rhomberg, L. R. & Beliles, R. P. Physiological parameter values for physiologically based pharmacokinetic models. *Toxicol. Ind. Health* **13**, 407–484 (1997).

74. Holt, K., Nagar, S. & Korzekwa, K. Methods to Predict Volume of Distribution. *Curr. Pharmacol. Rep.* **5**, 391–399 (2019).

75. Björkman, S. Prediction of the volume of distribution of a drug: which tissue-plasma partition coefficients are needed? *J. Pharm. Pharmacol.* **54**, 1237–1245 (2002).

76. Yun, Y. E., Tornero-Velez, R., Purucker, S. T., Chang, D. T. & Edginton, A. N. Evaluation of quantitative structure property relationship algorithms for predicting plasma protein binding in humans. *Comput. Toxicol.* **17**, 100142 (2021).

77. Al-Qassabi, J. *et al.* Facing the Facts of Altered Plasma Protein Binding: Do Current Models Correctly Predict Changes in Fraction Unbound in Special Populations? *J. Pharm. Sci.* **113**, 1664–1673 (2024).

78. Wood, F. L., Houston, J. B. & Hallifax, D. Clearance Prediction Methodology Needs Fundamental Improvement: Trends Common to Rat and Human Hepatocytes/Microsomes and Implications for Experimental Methodology. *Drug Metab. Dispos. Biol. Fate Chem.* **45**, 1178–1188 (2017).

79. Lombardo, F., Bentzien, J., Berellini, G. & Muegge, I. Prediction of Human Clearance Using In Silico Models with Reduced Bias. *Mol. Pharm.* **21**, 1192–1203 (2024).

80. Ring, B. J. *et al.* PhRMA CPCDC initiative on predictive models of human pharmacokinetics, part 3: Comparative assessement of prediction methods of human clearance. *J. Pharm. Sci.* **100**, 4090–4110 (2011).

81. Banerjee, S., Bhattacharya, A., Dasgupta, I., Gayen, S. & Amin, S. A. Exploring molecular fragments for fraction unbound in human plasma of chemicals: a fragment-based cheminformatics approach. *SAR QSAR Environ. Res.* **35**, 817–836 (2024).

82. Shen, J. & Nicolaou, C. A. Molecular property prediction: recent trends in the era of artificial intelligence. *Drug Discov. Today Technol.* **32–33**, 29–36 (2019).

83. Li, Z., Jiang, M., Wang, S. & Zhang, S. Deep learning methods for molecular representation and property prediction. *Drug Discov. Today* **27**, 103373 (2022).

84. Handa, K., Hirano, M., Kageyama, M. & Bender, A. Computational approaches to DMPK: A realistic assessment of current methods and their practical impact. Part I: Physicochemical and in vitro properties. *Drug Discov. Today* **30**, 104422 (2025).

85. Deng, J. *et al.* A systematic study of key elements underlying molecular property prediction. *Nat. Commun.* **14**, 6395 (2023).

86. Kantorovich, L. V. On the Translocation of Masses. *J. Math. Sci.* **133**, 1381–1382 (2006).

87. Carrillo, J. a. & Toscani, G. Wasserstein metric and large?time asymptotics of nonlinear diffusion equations. in *New Trends in Mathematical Physics* 234–244 (WORLD SCIENTIFIC, 2005). doi:10.1142/9789812702319_0022.

88. Kessy, A., Lewin, A. & Strimmer, K. Optimal Whitening and Decorrelation. *Am. Stat.* **72**, 309–314 (2018).

89. Kucherenko, S., Tarantola, S. & Annoni, P. Estimation of global sensitivity indices for models with dependent variables. *Comput. Phys. Commun.* **183**, 937–946 (2012).

90. Hilhorst, P. L. J., Quicken, S., van de Vosse, F. N. & Huberts, W. Efficient sensitivity analysis for biomechanical models with correlated inputs. *Int. J. Numer. Methods Biomed. Eng.* **40**, e3797 (2024).

91. Li, L., Lu, Z., Zhang, K. & Gao, Q. General validation and decomposition of the variance-based measures for models with correlated inputs. *Aerosp. Sci. Technol.* **62**, 75–86 (2017).

92. Wambaugh, J. F. *et al.* Evaluating In Vitro-In Vivo Extrapolation of Toxicokinetics. *Toxicol. Sci. Off. J. Soc. Toxicol.* **163**, 152–169 (2018).

93. Hinderling, P. H. Red Blood Cells: A Neglected Compartment in Pharmacokinetics and Pharmacodynamics. *Pharmacol. Rev.* **49**, 279–295 (1997).

94. Rodgers, T. & Rowland, M. Mechanistic Approaches to Volume of Distribution Predictions: Understanding the Processes. *Pharm. Res.* **24**, 918–933 (2007).

95. Schmitt, W. General approach for the calculation of tissue to plasma partition coefficients. *Toxicol. Vitro Int. J. Publ. Assoc. BIBRA* **22**, 457–467 (2008).

96. Yun, Y. E. & Edginton, A. N. Correlation-based prediction of tissue-to-plasma partition coefficients using readily available input parameters. *Xenobiotica* **43**, 839–852 (2013).

97. Poulin, P., Schoenlein, K. & Theil, F. Prediction of adipose tissue: Plasma partition coefficients for structurally unrelated drugs. *J. Pharm. Sci.* **90**, 436–447 (2001).

98. Lukacova, V. *et al.* Role of Fraction Unbound in Plasma in Calculations of Tissue:Plasma Partition Coefficients - Simulations Plus. (2008).

99. Zhivkova, Z. D., Mandova, T. & Doytchinova, I. Quantitative Structure - Pharmacokinetics Relationships Analysis of Basic Drugs: Volume of Distribution. *J. Pharm. Pharm. Sci. Publ. Can. Soc. Pharm. Sci. Soc. Can. Sci. Pharm.* **18**, 515–527 (2015).

100. Smith, D. A., Beaumont, K., Maurer, T. S. & Di, L. Volume of Distribution in Drug Design. *J. Med. Chem.* **58**, 5691–5698 (2015).

101. Murad, N. *et al.* Predicting Volume of Distribution in Humans: Performance of In Silico Methods for a Large Set of Structurally Diverse Clinical Compounds. *Drug Metab. Dispos.* **49**, 169 (2021).

102. Yau, E. *et al.* Global Sensitivity Analysis of the Rodgers and Rowland Model for Prediction of Tissue: Plasma Partitioning Coefficients: Assessment of the Key Physiological and Physicochemical Factors That Determine Small-Molecule Tissue Distribution. *AAPS J.* **22**, 41 (2020).

103. Sarrazin, F., Pianosi, F. & Wagener, T. Global Sensitivity Analysis of environmental models: Convergence and validation. *Environ. Model. Softw.* **79**, 135–152 (2016).

104. Awad, M., Senga Kiesse, T., Assaghir, Z. & Ventura, A. Convergence of sensitivity analysis methods for evaluating combined influences of model inputs. *Reliab. Eng. Syst. Saf.* **189**, 109–122 (2019).

105. Kolbe, N. nklb/wasserstein-distance: v1.0. Zenodo https://doi.org/10.5281/zenodo.10912241 (2024).

106. Reed, C. Data matrix whitening (https://www.mathworks.com/matlabcentral/fileexchange/34471-data-matrix-whitening), MATLAB Central File Exchange. (2012).

107. qqffssxx. Rand and Adjusted Rand Index Calculator for Cluster Analysis https://www.mathworks.com/matlabcentral/fileexchange/130779-rand-and-adjusted-rand-index-calculator-for-cluster-analysis, MATLAB Central File Exchange. (2023).