Early-Life Commensal Lung Bacteria Modulate Immune Responses and

Susceptibility to RSV Infection in Neonatal Mice

Quentin Marquant[§1,2,3], Claire Chottin[§1], Cécile Ferret[1,3], Eloi Poucet[1], Vinciane Saint-Criq[4], Daphné

Laubreton[1], Carole Drajac[1], Edwige Bouguyon[1], Aude Remot[5], Samuel Constant[6], Song Huang[6],

Edouard Sage[1,2,3], Isabelle Schwartz-Cornil[1,3], Muriel Thomas[4], Sabine Riffault[1,3], and Delphyne

Descamps[¶1,3]

[§] Quentin Marquant and Claire Chottin should be considered joint first authors.

[¶]Corresponding author

[1]Université Paris-Saclay, INRAE, UVSQ, VIM, 78350, Jouy-en-Josas, France

[2]Hôpital Foch, Suresnes, France

[3]ERL U1369, INSERM, V2I, 78350, Jouy-en-Josas, France

[4]Université Paris-Saclay, INRAE, Micalis, 78350, Jouy-en-Josas, France

[5]Université de Tours, INRAE, ISP, 37380, Nouzilly, France

[6]Epithelix, 18, Chemin des Aulx, CH-1228 Plan-les-Ouates, Geneva, Switzerland

Correspondence to: Delphyne Descamps, Tel: (33) 01 34 65 26 10, Email:

delphyne.descamps@inrae.fr. Virologie et Immunologie Moléculaires, INRAE, UVSQ, INSERM

Domaine de Vilvert, 78350 Jouy-en-Josas, France.

Abstract

The lung mucosa at birth and shortly after is particularly vulnerable to respiratory syncytial virus (RSV) infection. Respiratory disease severity is strongly influenced by the age at first exposure, which may also affect the trajectory of airway function. Early-life represents a critical window for lung development and initial microbiota colonization, both of which shape mucosal immune responses to RSV. The impact of the initial establishment of the lung microbiota by its primo-colonizing strains on susceptibility to RSV infection in neonatal mice remains poorly described. In the present study, we showed that early-life primo-colonizing bacterial strains in the mouse lung differentially induce innate immune responses and influence RSV susceptibility in *ex vivo* models using lung explants and alveolar macrophages (AMs). We identified a specific bacterial strain (strain 17) whose prior exposure enhances type I interferon (IFN-I) responses in AMs upon RSV infection and reduces viral replication both *ex vivo* and *in vivo* in lung tissues. Intranasal administration of this strain during early life prevented the development of immunopathological responses upon RSV reinfection in adult mice. Finally, using a translational human airway epithelium model, we demonstrated that pre-exposure to strain 17 restricts RSV spread without cytotoxicity, likely *via* enhanced β-defensin 2 production. These findings highlight the potential of early-life microbiota modulation as a promising intervention for preventing RSV disease and its long-term respiratory consequences.

Keywords

Respiratory Syncytial Virus (RSV), Early-life, Lung microbiota, Primo-colonizing bacteria, Mucosal immunity, Immune imprinting**,** Interferon, Alveolar macrophages, lung explants

1 Introduction

The respiratory syncytial virus (RSV), known as the primary cause of bronchiolitis in infants, is a major contributor to lower respiratory tract infections (LRTIs), and remains the leading cause of child mortality worldwide [1]. Moreover, severe RSV infection in early-life has been associated with long-term respiratory problems, such as repeated LRTI, wheezing, asthma, and reduced lung function, which can continue into adulthood in the form of chronic respiratory disease [2]. New passive immunisation strategies, such as maternal vaccination or administration of monoclonal antibodies, have been approved to reduce the burden of acute RSV infection in high-risk infants. However, some complications have been reported, including an increase in preterm births [3]. Direct vaccination in children remains a challenge, particularly after the FDA recently suspended all RSV vaccine trials in infants following reports of increased cases of severe illness in one trial [4]. The mechanisms underlying pulmonary hypersensitivity to RSV in early-life and its long-term health effects remain to be elucidated in order to develop effective prophylactic strategies. An increasing number of studies on RSV-infected neonatal mice (less than 7 days old at initial infection) have validated this model as relevant for understanding RSV infection in human infants [5]. It has been shown that the age at first infection plays a key role in shaping subsequent immune responses, notably by inducing epigenetic modifications [6]. This may increase the severity of later RSV infections [7], potentially by driving a type 2-biased immunity [8], which promotes allergic responses later in life [6]. Moreover, the early-life in mice is characterized by an immediate permissiveness to RSV replication, which gradually decreases with age as the ability to mount type I interferon (IFN-I) responses improves [9]. IFN-I are rarely detected in nasal washes from RSV-infected children [10]. However, IFN-α and IFN-β control viral pathogenesis by inducing interferon-stimulated genes that provide an early antiviral defense and shape inflammatory and adaptive immune responses [11]. IFN-I responses appear to play a key role in the outcome of RSV infection [11], as treatments that enhance IFN-I levels in infected neonatal mice have been shown to reduce viral replication and long-term RSV-induced lung effects [12, 13]. Thus, modulation of pulmonary innate immune defenses during the perinatal period can affect susceptibility to RSV infection, and determine long-term respiratory outcomes.

Early-life is indeed a crucial period for lung mucosa development, during which environmental factors influence the maturation of three interconnected components, namely, the airway epithelium, immune cells, and the resident microbiota. When the tuning between these three components is inadequate, it can ultimately affect susceptibility to infections or predispose to chronic respiratory diseases [14]. There is increasing evidence that postnatal lung mucosa is biased toward a type 2 immune response, as indicated by the spontaneous accumulation of type 2 innate lymphoid

cells (ILC2), eosinophils, and basophils, along with the production of cytokines such as IL-33 and

IL-13 [15], which are typically involved in anti-inflammatory responses and tissue repair. Such a

tissue environment is prone to the early polarization of tissue-resident alveolar macrophages (AMs)

toward a tolerogenic, anti-inflammatory M2 phenotype [16]. Moreover, the lung microbiota

establishes rapidly after birth and it shaped by factors linked to the host and the environment [17-19].

A recent prospective pediatric cohort study showed that between birth and 3 months of age, rapid

changes in the airway bacterial microbiota are accompanied by a major shift in the JAK/STAT and

interferon immune pathways, revealing a link between bacterial communities and early-life

respiratory immunity [20]. In addition, the lung microbiota contributes to immune tolerance in early-

life according to an asthma model [17] and attenuates innate immune reactivity to PAMPs [21].

The incidence of respiratory infections and the occurrence of dysbiosis in the lung microbiota

are closely related [22]. The severity of RSV infection, [23, 24] and the persistence of symptoms in

children may be influenced by the bacterial ecosystem [25]. Conversely, the presence of *Lactobacillus*

in the nasopharynx is associated with a reduced risk of childhood wheezing illnesses at 2 years of age

[26]. In mice, viable commensal bacteria are detected in the lungs as early as 3 days of age, with their

number and diversity increasing over time [17] and reaching a plateau by weaning at around 21 days

of age [19]. A collection of primo-colonizing bacteria was established *via* their isolation from lung

homogenates of neonatal mice using aerobic or anaerobic culture conditions [19]. Intranasal

administration of certain strains of this collection during early life influenced the outcome of an

allergic asthma model [19], while instillation of other strains in adult mice attenuated pulmonary

inflammation following *Mycobacterium tuberculosis* infection [27]. These findings suggest that lung

primo-colonizing commensal bacteria can modulate pulmonary immunity and influence

susceptibility to respiratory diseases. However, the mechanisms by which early bacterial colonization

of the lungs modulates immune responses and determines both immediate and long-term respiratory

outcomes, including during early-life RSV infection, remain largely unexplored [28]. In the present

study, we demonstrated that early-life commensal lung bacteria modulate immune responses and RSV

susceptibility in neonatal mice. We established a proof of concept demonstrating that it is possible to

select a single strain from a panel of lung primo-colonizing commensal bacteria based on its

immunomodulatory and antiviral properties on the neonatal lungs. One bacterial strain in particular,

hereafter referred to as strain 17, promoted IFN-I antiviral pathways and reduced RSV replication

both *ex vivo* and *in vivo* in lung tissues. Finally, its intranasal administration to neonatal mice

prevented early-life immunopathological priming, which became apparent after a second exposure to

RSV in adulthood.

2 Materials and methods

2.1 Bacterial strains' identification and growth

Bacterial strains (Supplemental Table 1) previously isolated from 3-day-old mouse lungs were stored at −80°C [19]. Strains were revived in 10 mL yeast-heminBHI or MRS medium and subcultured (100 µL/10 mL). Growth was monitored hourly for 8h by spectrophotometry to determine exponential and stationary phases. Bacterial counts were obtained using a BD Accuri™ C6 flow cytometer, and strain identity was confirmed by 16S PCR (J4/J7 primers) and MALDI-TOF mass spectrometry (CIRM, INRAE, Nouzilly, France).

2.2 Virus

Human RSV strain A2 and recombinant RSV-Long expressing mCherry (RSV-mCherry) or luciferase (RSV-Luc) were produced, titrated on HEp-2 cells, and stored at -80°C [29].

2.3 *Ex vivo* lung explants, stimulation and infection

Pregnant BALB/cJrj mice were purchased from Janvier (Le Genest, France). Lungs from 6-day-old neonatal mice were filled *via* tracheotomy with RPMI containing 1.5% low-melting agarose (37°C). After solidification, lungs were placed in a Krumdieck tissue slicer (MD6000) chamber containing PBS with 1× antibiotic-antimycotic and cut into 150 µm slices to obtain precision-cut lung slices (PCLS) [19, 21]. PCLS were stimulated for 16h at 37 °C, 5% CO_2 with strains (50 CFU/well in RPMI with 5% FBS and L-glutamine). Inoculations were verified by CFU counts on agar plates. Three hours later, PCLS were infected with RSV-Luc (7.19×10^4 PFU/well) or Mock control. Supernatants were collected. PCLS were lysated in lysis buffer (5 mM EDTA, 150 mM NaCl, 50 mM Tris-HCl, 1% Triton X100, pH 7,4) or Passive Lysis Buffer (30 mM Tris, 10 mM $MgCl_2$, 1% Triton 100X, 2% Glycerol) using a Precellys 24-bead grinder homogenizer (Bertin Technologies) for 2×15s at 6,000 x g. Protein levels were quantified with DC™ or RC-DC™ Protein Assays (Bio-Rad).

2.4 Experiments with differentiated human primary bronchial epithelial cells

MucilAir® from healthy donors were purchased from Epithelix. Cells ($\approx 5 \times 10^5$ cells/transwell) were stimulated with bacterium 17 (10^3 CFU/transwell) in 20 µL in apical side. After 3h, cells were infected with RSV-mCherry (MOI = 0.1) or Mock control during 3h. The inoculum was removed and medium replaced by strain 17 containing fresh medium. RSV-mCherry replication was visualized using the ZOE Fluorescent Cell Imager (Bio-Rad) and quantified with a fluorescence plate reader (Tecan Infinite 200 Pro). Supernatants and cell lysates were then collected.

2.5 *Ex vivo* AM culture and infection

Bronchoalveolar lavages (BALs) were performed on 6-days-old mice to isolate AM, as previously described [9, 30, 31]. Cells (1×10^5) were plated, 24h after seeding, incubated with bacterial strains

(50 CFU/well) for 3h before exposure to RSV-Long (MOI = 5) or a mock control. Supernatants and

cells were collected 24h post-infection.

2.6 Cell cytotoxicity

Cytotoxicity was measured by lactate deshydrogenase (LDH) activity, using the Non-Radioactive

Cytotoxicity Assay kit (Promega).

2.7 Cytokine, IFN, and RSV-specific Ab assays

Cytokine production was measured using the 15-plex or IL-6/TNF-α/IFN-α/IFN-β custom 4-Plex

mouse ProcartaPlex™ immunoassay (ThermoFisher Scientific). Human IL-6, TNF-α or IP-10 were

measured by Luminex and human β-defensin-2 by ELISA (NB-50-1634-1000, NeoBiotech). Data

were read on MagPix instrument (Luminex), and analyzed using Bio-Plex Manager software (Bio-

Rad). HEK-Blue™ IFN-λ (InvivoGen) cells were used for the detection of bioactive human IFN-λ.

Individual mouse sera were assayed for specific Ab (IgG1, IgG2a) against RSV lysates by ELISA

[13].

2.8 *In vivo* bacterial strains inoculation and RSV infection

Neonate mice (age 3 and 5 days) received 10 µL of strain 17 intranasally (i.n.) route. At 6 days old,

mice were infected with RSV-Luc (7.187 x 10^6 PFU/mL, 10 µL, i.n) or a Mock control. In other

experiments, neonate mice were first infected by RSV-A2 (2.3 × 10^6 PFU/mL, 10 µL, i.n.), and were

kept until 8 weeks of age when they were all infected with RSV-A2 (100 µL, i.n.) [13]. BAL and lung

tissues were collected.

2.9 Bioluminescence measurements

Photon emission was measured in lysates of the nasal cavity or lung tissues and PCLS using the IVIS

system (Xenogen Biosciences) and Living Image software (version 4.0, Caliper Life Sciences),

following a previously described method [9], and was normalized to protein content.

2.10 Cell infliltration

Cells were collected from BALs and centrifugated at 400 x g for 10min at 4°C. May-Grunwald-

Giemsa coloration was performed to assess the differential cell counts using light microscopy. Two

hundred cells were counted to determine relative and absolute cell numbers.

2.11 RT-qPCR

Total RNA was extracted from lung lysates using a NucleoSpin® RNA midi column (Macherey

Nagel) and reverse transcribed using random primers and SuperScript II (Invitrogen), according to

the manufacturer's instructions. The primers (Sigma–Aldrich) used are listed (Supp. Table 2). RT-

qPCR was performed in duplicate for each gene using the CFX Connect™ Real-Time PCR Detection

System (Bio-rad) and SYBRGreen PCR Master Mix (Eurogenetec). Data were analyzed using CFX

Manager™ Software (Bio-rad) to determine the quantification cycle (Cq) values. Results were

determined using the formula $2^{-\Delta Cq}$, with $\Delta CQ = Cq_{gene} - Cq_{(mean\ (bActine, Gapdh,\ Hprt))}$.

2.12 Data and statistical analysis

Data are expressed as mean ± SEM. Statistical significance was set at *$P < 0.05$, **$P < 0.01$, ***$P <$

0.001, and ****$P < 0.0001$ using GraphPad Prism 8 (GraphPad Software Inc.). Data normality was

assessed using the D'Agostino–Pearson and Shapiro–Wilk tests. Depending on the distribution,

differences between groups were analyzed using a parametric *t*-test or a nonparametric Mann–

Whitney test (two groups), or by one-way ANOVA/Kruskal–Wallis or two-way ANOVA with

Tukey's multiple comparisons test.

3 Results

3.1 Primo-colonizing bacteria exert distinct immunostimulatory effects on neonatal lung mucosa *ex vivo* model.

The goal was to test the concept that a commensal strain can promote a protective, antiviral type 1 immunity in the lungs during early-life. To evaluate the immunostimulatory effects of primo-colonizing bacterial strains (Supplemental Table 1) on the neonatal lung, we established a PCLS-based screening assay using lung tissue from 6-day-old mice, an *ex vivo* model that recapitulates pulmonary cytokine production [32]. PCLS were exposed to each strain at MOI=50 (Figure 1A). Among 20 strains tested, two (strains 1 and 11) caused significant cytotoxicity after 16h and were excluded, while most showed cytotoxicity levels comparable to the control medium (mean = 21.3% $\pm$ 1.1%), indicating no harmful effects (Supplemental Figure S1A). Cytokines were quantified using a 15-plex assay and grouped by functional category (pro-inflammatory, type 1 immunity, type 2 immunity, Th17/Th22/Th9-responses, and lymphocyte proliferation, Supplemental Figure S1B-F). Both the nature and magnitude of cytokine production by PCLS varied across bacterial strains. Strains 2, 19 and 16 induced high levels of pro-inflammatory cytokines such as TNF-α and IL-6, together with robust a type 2 immune response, including IL-10 (Supplemental Figures S1B and S1D). In contrast, most other strains induced low cytokine production. A bacterial strain, the strain 17, stood out from the others. Strain 17, identified as *Escherichia coli* by 16S sequencing and mass spectrometry (Supplemental Table 1), also inducing type 2 cytokines (Supplemental Figure S1D) but was unique in triggering a strong type 1 profile, characterized by a high production of IFN-γ and IL-12p70 (Figure 1B), and increased IL-22 and IL-9 secretion (Supplemental Figure S1E). To highlight strain-specific immune signatures, cytokine levels were expressed as fold-change relative to control PCLS and visualized as heat map (Figure 1C), with hierarchical clustering revealing groups of strains with similar immunostimulatory patterns. Overall, primo-colonizing lung bacteria displayed distinct immunostimulatory capacities in PCLS from neonatal mice, with strain 17 standing out for its strongly type 1-skewed immune profile and lack of cytotoxicity.

3.2 Strain 17 exerts *ex vivo* antiviral activity against RSV by limiting viral replication and enhancing the innate immune response.

We next focused on the early events following RSV infection to explore how strain 17 affects early innate antiviral response. To assess whether strain 17 influences early-life RSV susceptibility, we performed co-exposure experiments using PCLS from 6-day-old mice. PCLS were first exposed to strain 17 (50 CFU/well as previously used [19]) to elicit its immunostimulatory effects, and then infected with luciferase-expressing RSV (RSV-Luc) or mock-treated for 16h to evaluate viral

replication (Figure 2A). Strains 16, 24, and 25, displaying distinct cytokine profiles (see Supplemental

Figure S1), were included as comparators. Cytotoxicity induced by these strains, RSV-Luc, or their

combination did not exceed 21% and was comparable to that of the control medium. Thus, co-

exposure did not affect PCLS viability and supporting the safe use of these bacteria during RSV

infection (Figure 2B). Viral replication was quantified in PCLS lysates using a luciferase activity

assay (Figure 2C). PCLS exposed to strain 25 showed a slight increase in luciferase activity compared

to the control condition (not statistically different), suggesting a potential enhancement of RSV

replication associated with the presence of strain 25. Pre-exposure to strains 16 or 24 resulted in only

a slight reduction in RSV replication (non stastistically significant). In contrast, exposure to strain 17

resulted in a strong significant reduction in luciferase activity compared to both control and PCLS

exposed to strain 16 or 24 conditions. Altogether, these results suggest that pre-exposure to lung

primo-colonizing commensal bacteria can modulate RSV replication in PCLS from 6-day-old mice,

with strain 17 emerging as the most effective strain in reducing RSV replication.

We also investigated the effect of strain 17 on AMs, the main pulmonary resident immune

cells, involved in IFN-I production during RSV infection [33]. AMs isolated from BAL of 6-day-old

mice were first exposed to strain 17 (50 CFU/well as previously used [19]) or control medium,

followed by exposure to RSV (Figure 3A). After 16h of incubation, exposure of AMs to strain 17

alone induce a slight but unsignificant production of IL-6, TNF-α, and IFN-I in the supernatants

compared to basal control conditions (no strain, no RSV, Figures 3B-C). When AMs exposed to strain

17 were subsequently infected with RSV, they produced markedly more inflammatory (IL-6 and

TNF-α, Figure 3B) and antiviral (IFN-α and IFN-β, Figure 3C) cytokines than RSV-infected AMs

without bacterial pre-exposure. Notably, RSV-infected AMs produced 3-fold higher levels of IFN-I

when exposed to strain 17. Thus, neonatal AMs exposure to strain 17 improved their ability to mount

an innate antiviral immune response.

3.3 Pulmonary exposure to strain 17 decreases the susceptibility of neonatal mice to RSV

infection.

Collectively, these *in vitro* results identify strain 17 as non cytotoxic strain capable of enhancing IFN-

I production and limiting RSV replication. We therefore evaluated whether pulmonary exposure to

strain 17 confers protection against RSV infection in neonatal mice. To assess the *in vivo* relevance

of its effects, we first investigated whether intranasal (i.n.) administration of this commensal strain

(10^6 CFU; dose and method previously shown to deliver sufficient bacteria to the lungs [19]) was

non-pathogenic in neonatal mice. No mortality was observed in neonatal mice following

administration of strain 17 (data not shown). A modest increase in neutrophil counts was detected in

BALs, although total cell numbers remained unchanged (Supplemental Figure S2). Next, we

performed a dose-response study to determine the minimal amount of strain 17 required to achieve its *in vivo* immunomodulatory effect. Neonatal mice received three different amounts of strain 17 or PBS diluent intranasally at 3- and 5-days-old to modulate the neonatal lung mucosa before infection with RSV-Luc at day 6 (Figure 4A). Mice were euthanized 1 or 2 d.p.i. to assess viral replication in the upper and lower respiratory tracts, as well as the associated immune response in lung tissues. As previously shown [9, 13], i.n. RSV-Luc infection of neonatal mice led to rapid viral replication in the nasal cavity, reaching $1.42 \times 10^5 \pm 0.38 \times 10^5$ and $0.89 \times 10^5 \pm 0.28 \times 10^5$ photons/mg at 1 and 2 d.p.i., respectively (Figure 4B). In PBS-treated neonates, lung replication was detectable at 1 d.p.i. $(1.02 \times 10^4 \pm 0.38 \times 10^4$ photons/mg) and increased by 2 d.p.i. $(2.09 \times 10^4 \pm 0.32 \times 10^4$ photons/mg), confirmed by RSV *N gene* expression (Figures 4C-D). Strain 17 significantly reduced nasal viral replication at 1 d.p.i. at all doses (3-4-fold, Figure 4B), whereas only the 10^6 CFU dose significantly lowered lung replication $(0.098 \times 10^4 \pm 0.047 \times 10^4$ photons/mg, Figure 4C). The 10^5 CFU dose showed delayed lung protection at 2 d.p.i., while the 10^3 CFU dose had no effect, indicating that upper airway protection requires lower doses, whereas lung protection needs higher or delayed dosing (Figures 4B-D). RSV infection in neonatal mouse lungs induced low *IFN-I* gene expression at 1 d.p.i. (Figure 5A), and led to a slight, non-significant, increase in BAL cellularity between 1 and 2 d.p.i. (Figure 5B), primarily due to neutrophil recruitment (Figure 5D), as previously shown [9, 12, 13]. In neonates treated with high doses of strain 17, *IFN-I gene* expressions, and notably IFN-α, were increased at 1 d.p.i. (Figure 5A), while total BAL cells (Figure 5B) and macrophages (Figure 5C) remained unchanged. In contrast, neutrophil recruitment was significantly enhanced from 1 to 2 d.p.i., with an ~8-fold increase at 2 d.p.i. in RSV-infected mice pre-treated with the highest doses of strain 17 (Figure 5D). Overall, these results demonstrate that neonatal exposure to strain 17 enhances early innate immune responses, leading to reduced RSV replication, dose-dependent protection of the lungs and a strong protective effect in the nasal cavity.

3.4 Strain 17 treatment of neonates attenuates the long-term pulmonary consequences of early-life RSV infection upon rechallenge in adulthood.

In vitro, bacterial strain 17 demonstrated a pronounced enhancement of type 1 immune responses. Thus, we employed a previously described model of adult RSV re-exposure to investigate the ability of strain 17 to promote protective immunity and counteract the immunopathological lung priming induced by early-life RSV infection [8]. Three-day-old mice were treated intranasally with strain 17 or PBS vehicle at days 3, 5, 8, and 10 days of age. At 6-days-old, mice were infected intranasally with RSV, followed by a second exposure in adulthood (Figure 6A). Weight change was monitored (Figure 6B), and antibody responses (Figures 6C-D) and cell infiltration in BALs were assessed (Supplemental Figures S3B-E). As expected, mice initially infected as pups and re-exposed to RSV

in adulthood (PBS/RSV/RSV group) lost weight for over the 14 days post-infection (d.p.i), whereas mice treated with strain 17 and infected with RSV as neonates (strain 17/RSV/RSV group) lost little weight and regained their initial weight by 8 d.p.i., consistently showing a better although not stastically significant, maintenance of body weight compared with controls (Figure 6B). We next investigated whether neonatal strain administration would impact the anti-RSV IgG1 response indicative of type 2 immunity and expected for adults upon RSV reexposure [8]. As expected, the re-exposure to RSV in adulhood induced a significant production of anti-RSV-IgG1 at 14 d.p.i. in the PBS/RSV/RSV group (Figure 6C). In contrast, strain 17-treated neonates showed an early induction of anti-RSV-IgG2a antibody production as soon as 7 d.p.i. (Figure 6D) and no significant increase in IgG1 at 14 d.p.i. (Figure 6C). Finally, we monitored cellular inflammation. At 7 d.p.i., total BAL cell numbers were significantly higher in mice first infected as neonates and re-exposed as adults (PBS/RSV/RSV, $2.51 \times 10^5 \pm 0.38 \times 10^5$ cells) than in mice infected only as adults (PBS/Mock/RSV, $1.26 \times 10^5 \pm 0.33 \times 10^5$ cells ; Supplemental Figure S3B, $p < 0.05$). BAL cell counts in strain 17/RSV/RSV-treated mice did not differ significantly from the PBS/RSV/RSV group (Supplemental Figure S3B). Macrophage and neutrophil numbers were similar across groups at 7 and 9 d.p.i. (Supplemental Figures S3C–D), while the overall leukocyte increase was mainly due to lymphocyte recruitment ($0.98 \times 10^5 \pm 0.19 \times 10^5$ cells in PBS/RSV/RSV *vs.* $0.81 \times 10^5 \pm 0.24 \times 10^5$ cells in strain 17/RSV/RSV; Supplemental Figure S3E), which persisted until 9 d.p.i. despite declining over time. Significant eosinophil recruitment was observed in the BAL of PBS/RSV/RSV mice at 7 d.p.i. ($0.08 \times 10^5 \pm 0.012 \times 10^5$ cells) and persisted at 9 d.p.i. ($0.065 \times 10^5 \pm 0.019 \times 10^5$ cells), but was absent in PBS/Mock/RSV mice, reflecting type 2 immune priming from early-life RSV infection (Figures 6E–F). Notably, eosinophil infiltration was nearly abolished in mice treated with strain 17 prior to the first RSV infection ($0.006 \times 10^5 \pm 0.004 \times 10^5$ cells at 7 d.p.i.). Accordingly, the strain 17/RSV/RSV group had 9.6- and 3.4-fold fewer eosinophils in the BAL compared to PBS/RSV/RSV animals at 7 and 9 d.p.i., respectively. Neonatal treatment with strain 17 did not affect overall leukocyte infiltration upon adult RSV re-exposure but prevented immunopathologic eosinophilia. Altogether, these data indicate that the early-life presence of strain 17 during the first RSV infection of mice promoted a shift in the RSV-specific antibody toward a type 1 profile (higher IgG2a, lower IgG1), associated with minimal eosinophil recruitment and improved disease outcomes (trend to maintain their weight).

3.5 Strain 17 limits RSV spread in a translational human airway model.

Finally, to validate the beneficial anti-RSV effect of strain 17 in a translational model, human primary airway epithelial cells cultured at the air-liquid interface were pre-exposed to strain 17 and subsequently infected with mCherry-expressing RSV to visualize cellular RSV replication (Figure 7A). Without bacterial pre-exposure, RSV replication was detectable at 48h post-infection, as shown

by mCherry fluorescence (Figure 7B). In contrast, strain 17 exposure markedly reduced mCherry

activity compared to the control (Figures 7B-C). No cytotoxicity was detected (Figure 7D), and

neither RSV exposure nor treatment with strain 17 induced the production of cytokines (IL-6, TNF-

α or IP-10, Supplemental Figure S4) in these experimental conditions. Type III interferons (IFN-λ)

are produced by airway epithelial cells infected with influenza virus earlier than IFN-I, helping to

restrict viral spread without provoking inflammation [34]. As expected, RSV infection induced IFN-

λ production in epithelial cells (Figure 7E). Strain 17 alone did not induce IFN-λ and did not enhance

the IFN-λ response to RSV infection; if anything, it tended to reduce secretion. Overall, the effect of

strain 17 is therefore unlikely to be explained by an increase in IFN-λ. We then measured another

antiviral factor, the antimicrobial peptide human β-defensin 2, in epithelial cells (Figure 7F). This

cationic molecule, secreted by lung epithelial cells, blocks RSV entry by destabilizing its envelope

[35]. Strain 17 strongly induced β-defensin 2 (2040 ± 571 pg/mL *vs* 65 ± 19 pg/mL in controls),

whereas RSV alone triggered low levels (120 ± 32 pg/mL). Thus, pre-exposure to strain 17 markedly

reduced RSV propagation in a human airway epithelium model without cytotoxicity, likely *via*

enhanced β-defensin 2 secretion.

4 Discussion

The lung microbiota and its influence on RSV infection severity have recently been identified as critical research areas in the fight against bronchiolitis in young children [23, 24] [21]. While modulation of the lung microbiota or metabolites appears to be a promising strategy for strengthening immune defenses in adults [36, 37], the impact during the critical period of lung mucosal development and susceptibility to RSV infection is yet unknown [28]. In this study, we demonstrated that the immune responses of neonatal mice to RSV infection, as well as their long-term consequences, could be favorably modulated by lung primo-colonizing bacterial strains given shortly after birth. In particular, using complementary cellular and *in vivo* approaches, we identified a bacterial strain 17 (*E. coli*), that promoted protective type 1 immune responses and limited RSV replication in both mouse neonatal models and human epithelial cells.

We employed a PCLS-based screening platform from neonatal mice to assess cytokine responses to a collection of primo-colonizing bacterial strains isolated from mouse neonatal lungs. This strain collection encompasses major bacterial families, ranging from *Staphylococcus* to *Lactobacillus*. While most *Lactobacillus* strains exhibited low immunostimulatory activity on neonatal PCLS, the majority of *Staphylococcus* strains induced a pro-inflammatory cytokine profile. In contrast, strain 17, identified as *Escherichia coli*, uniquely induced a strong polarized type-1 cytokine response, particularly IFN-γ and IL-12p70, without causing cytotoxicity, suggesting its potential to prime antiviral immunity in the neonatal lung. However, strain 17 also induced the production of IL-9, a cytokine linked to asthma and detected in airway aspirates from patients with severe bronchiolitis [38]. Its role in early-life pulmonary mucosa and RSV pathology remains poorly understood [39]. Interestingly, strain 17 also induced the secretion of IL-22, which is known to promote epithelial barrier regeneration [40], and has a beneficial inhibitory effect on RSV infection has been demonstrated in a neonatal mouse model [41]. Although cytokine accumulation during 24-hour PCLS-based screening prevents the evaluation of secretion kinetics or autocrine effects, this approach nevertheless reveals the broad potential of early-life lung primo-colonizing bacteria to induce immune stimulation and polarization. Strain 17 differed markedly from the other lung primo-colonizing strains tested due to its unique ability to limit RSV replication in both PCLS and a human airway epithelium model, underscoring its translational relevance. It is interesting to note that strains taken individually may display disease-specific immunobeneficial effects. Indeed, strain 24, previously shown to be protective in a juvenile asthma model [19], had neither a protective nor a detrimental effect on RSV replication in this study. These findings underscore the diverse immunomodulatory potential of bacteria within the pulmonary microbiota. Functionally, the antiviral activity of strain 17 in human epithelial cells appeared to be independent of IFN-λ and was associated with increased production of

β-defensin 2, a known host defense peptide capable of destabilizing the viral envelope, thereby
indirectly interfering with RSV entry and/or propagation [35]. Our study also demonstrated that strain
17 can prime anti-RSV immunity in neonatal AMs by enhancing their production of IFN-I. Thus,
strain 17 confers innate protective immunity to the lung mucosa in early-life.

Our *in vivo* studies confirmed the ability of strain 17 to reduce RSV replication in the upper and
lower respiratory tracts of neonatal mice. The degree of protection was dose-dependent and associated
with an induction of IFN-I expression, as well as increased neutrophil recruitment. The role of the
neutrophils recruited in strain 17-treated mice, with or without RSV infection, remains unclear,
particularly regarding their protective *versus* pathological effects in neonates [42]. Nevertheless,
early-life exposure to strain 17 mitigated long-term immunopathology upon re-exposure to RSV in
adulthood. Mice treated with strain 17 exhibited a relatively stable body weight curve with no major
alterations following viral reinfection, a shift toward type 1 immunity (increased IgG2a, decreased
IgG1), and reduced eosinophilic infiltration. These data indicate that early-life administration of strain
17 induces protective immunomodulation in the lung mucosa, mitigating the long-term consequences
of type 2 immune imprinting established by neonatal RSV infection. Although colonization was not
assessed, our study highlights the importance of dosing relative to the targeted respiratory region. For
instance, the instillation of 10^6 CFU into the lungs of a neonatal mouse represents a substantial
bacterial load compared to the naturally sparse neonatal lung microbiota [19]. Higher doses were
required to confer early protection in the lungs, whereas lower doses sufficed for the upper airways.
This likely reflects the larger tissue surface of the lungs that must be exposed to the bacteria, rather
than differences in microbiota density, which is already low in the pulmonary mucosa [43]. However,
before proposing this bacterial strain as a therapeutic candidate for early immunomodulation to
prevent acute and chronic RSV-related diseases, it is essential to elucidate the molecular pathways
underlying its immunostimulatory effects in both epithelial and immune cells. Future studies should
investigate whether the beneficial effects of strain 17 require live microorganisms, which colonize
tissues and are likely more effective than inert components to produce metabolites. Testing
conditioned supernatants could then clarify the contribution of bacterial metabolic activity, such as
acetate production *in vitro* (data not shown), a metabolite known to enhance antiviral responses in
AMs during RSV infection [44, 45]. Indeed, in an adult context, airway administration of
Lactobacillus or *Corynebacterium* strains [46, 47], or a standardized bacterial lysate derived from
human respiratory strains [48] is associated with improved resistance of mice to RSV challenge.
Morever, higher acetate levels boosts the IFN-β response of AMs and epithelial ISG expression,
reducing viral load and pulmonary inflammation in RSV-infected adult mice [44, 45]. It has been
shown that *E. coli* probiotic strains, can have both deleterious and beneficial effects on digestive

health [49]. So, before exploiting a such promising strain as a probiotic to enhance respiratory

defenses, a global safety assessement is required.

5 Conclusion

Overall, this study reveals that the immune responses of neonatal mice to RSV infection, as well

as their long-term consequences, can be favorably modulated by early-life lung-colonizing bacterial

strains. These results reinforce the notion that the neonatal lung microbiota plays a key role in shaping

pulmonary immune responses, and that its disruption, for example by antibiotics, can have profound

and lasting effects on host immunity and respiratory vulnerability [50]. A recent study in pediatric

RSV cohorts indicates that persistent respiratory symptoms during the convalescence phase are

associated with disruptions to the respiratory microbiota [25]. It would be interesting to explore

further the potential benefits of primo-colonizing strains during recovery.

Author Contibutions

Q.M. and C.C.: Methodology, Validation, Investigation, Data curation, Visualization, Writing – review & editing. C.F.: Validation, Data curation, Resources. E.P. and E.B.: Validation, Data curation. V.S-C. and D.L.: Methodology, Writing – review & editing. C.D.: Methodology, Investigation. A.R.: Methology, Conceptualization. S.C., S.H. and E.S.: Resources. I.S-C., M.T. and S.R.: Resources, Conceptualization, Writing – review & editing. D.D.: Conceptualization, Methodology, Investigation, Validation, Formal analysis, , Writing – Original Draft, Writing – review & editing, Supervision, Visualization, Project administration, Funding acquisition.

Acknowledgements

We greatly acknowledge M. Prost and A. Mialet for their assistances with experiments. We are grateful to M-A. Rameix-Welti (INSERM, U1173, UVSQ) and J-F Eléouët (INRAE, UVSQ, VIM) for providing recombinant RSV-Cherry. We are grateful to the IERP unit, INRAE (Infectiology of fishes and rodent facility, doi: 10.15454/1.5572427140471238E12). We thank the Emerg'in platform for access to IVIS-Spectrum that was financed by the Region Ile De France (DIM One Health). We thank E Helloin and C Darrigo (CIRM, INRAE, Nouzilly) for bacterial strains identification by mass spectrometry.

Fundings

C. Drajac. and Q. Marquant were the recipient of a Ph.D. and post-doctoral fellowships of Région Ile-de-France (DIM-Malinf or DIM-OneHealth, respectively). C. Chottin was the recipient of a Ph.D. fellowship from the French research ministry. D. Laubreton and experiments were supported by a grant of the French national research agency "Agence Nationale de la Recherche" (ANR-13-BSV3-0016 and ANR-24-NEOMIS). This project was supported by the Fondation Air Liquide, Vaincre la Mucoviscidose and by Association Grégory Lemarchal.

Ethics Statement

All experiments were approved by the ethics committee COMETHEA (Ethical Committee for Animal Experimentation, authorization number: APAFIS#14654-2018032116445169).

Conflict of Interest statement

The authors declare no competing financial interests. The authors declare that all the other data supporting the findings of this study are available within the article and its supplementary information files and from the corresponding author upon request.

Data Availability Statement

Data are available from the corresponding author upon reasonable request.

References.

1.   Liu, L., et al., *Global, regional, and national causes of under-5 mortality in 2000-15: an updated systematic analysis with implications for the Sustainable Development Goals.* Lancet, 2016. **388**(10063): p. 3027–3035.

2.   Zar, H.J., et al., *Early-life respiratory syncytial virus disease and long-term respiratory health.* Lancet Respir Med, 2024. **12**(10): p. 810–821.

3.   Dieussaert, I., et al., *RSV Prefusion F Protein-Based Maternal Vaccine - Preterm Birth and Other Outcomes.* N Engl J Med, 2024. **390**(11): p. 1009–1021.

4.   Mahase, E., *FDA pauses all infant RSV vaccine trials after rise in severe illnesses.* BMJ, 2024. **387**: p. q2852.

5.   Cormier, S.A., D. You, and S. Honnegowda, *The use of a neonatal mouse model to study respiratory syncytial virus infections.* Expert Rev Anti Infect Ther, 2010. **8**(12): p. 1371–80.

6.   Morris, S.B., et al., *Long-term alterations in lung epithelial cells after EL-RSV infection exacerbate allergic responses through IL-1beta-induced pathways.* Mucosal Immunol, 2024. **17**(5): p. 1072–1088.

7.   Malinczak, C.A., et al., *Early-life pulmonary viral infection leads to long-term functional and lower airway structural changes in the lungs.* Am J Physiol Lung Cell Mol Physiol, 2024. **326**(3): p. L280–L291.

8.   Culley, F.J., J. Pollott, and P.J. Openshaw, *Age at first viral infection determines the pattern of T cell-mediated disease during reinfection in adulthood.* J Exp Med, 2002. **196**(10): p. 1381–6.

9.   Laubreton, D., et al., *Regulatory B Lymphocytes Colonize the Respiratory Tract of Neonatal Mice and Modulate Immune Responses of Alveolar Macrophages to RSV Infection in IL-10-dependant Manner* Viruses, 2020. **12**(822).

10.   Hall, C.B., et al., *Interferon production in children with respiratory syncytial, influenza, and parainfluenza virus infections.* J Pediatr, 1978. **93**(1): p. 28–32.

11.   Hijano, D.R., et al., *Role of Type I Interferon (IFN) in the Respiratory Syncytial Virus (RSV) Immune Response and Disease Severity.* Front Immunol, 2019. **10**: p. 566.

12.   Cormier, S.A., et al., *Limited type I interferons and plasmacytoid dendritic cells during neonatal respiratory syncytial virus infection permit immunopathogenesis upon reinfection.* J Virol, 2014. **88**(16): p. 9350–60.

13.   Remot, A., et al., *Flt3 ligand improves the innate response to respiratory syncytial virus and limits lung disease upon RSV reexposure in neonate mice.* Eur J Immunol, 2016. **46**(4): p. 874–84.

14.   Torow, N., et al., *Neonatal mucosal immunology.* Mucosal Immunol, 2017. **10**(1): p. 5–17.

15.   de Kleer, I.M., et al., *Perinatal Activation of the Interleukin-33 Pathway Promotes Type 2 Immunity in the Developing Lung.* Immunity, 2016. **45**(6): p. 1285–1298.

16.   Saluzzo, S., et al., *First-Breath-Induced Type 2 Pathways Shape the Lung Immune Environment.* Cell Rep, 2017. **18**(8): p. 1893–1905.

17.   Gollwitzer, E.S., et al., *Lung microbiota promotes tolerance to allergens in neonates via PD-L1.* Nat Med, 2014. **20**(6): p. 642–7.

18.   Pattaroni, C., B.J. Marsland, and N.L. Harris, *Early-Life Host-Microbial Interactions and Asthma Development: A Lifelong Impact?* Immunol Rev, 2025. **330**(1): p. e70019.

19.   Remot, A., et al., *Bacteria isolated from lung modulate asthma susceptibility in mice.* ISME J, 2017. **11**(5): p. 1061–1074.

20.   Pattaroni, C., et al., *Bacterial communities co-develop with respiratory immunity early in life, linking dysbiosis to systemic monocyte signature and wheezing.* Sci Adv, 2025. **11**(42): p. eadw1410.

21.   Marquant, Q., et al., *The microbiota plays a critical role in the reactivity of lung immune components to innate ligands.* FASEB J, 2021. **35**(4): p. e21348.

22.   de Steenhuijsen Piters, W.A.A., et al., *Early-life viral infections are associated with disadvantageous immune and microbiota profiles and recurrent respiratory infections.* Nat Microbiol, 2022. **7**(2): p. 224–237.

23.   Schippa, S., et al., *Nasal Microbiota in RSV Bronchiolitis.* Microorganisms, 2020. **8**(5).

24.   Yagi, K., et al., *Early-Life Lung and Gut Microbiota Development and Respiratory Syncytial Virus Infection.* Front Immunol, 2022. **13**: p. 877771.

25.   Kristensen, M., et al., *The respiratory microbiome is linked to the severity of RSV infections*
and the persistence of symptoms in children. Cell Rep Med, 2024. **5**(12): p. 101836.
26.   Rosas-Salazar, C., et al., *Nasopharyngeal Lactobacillus is associated with a reduced risk of*
childhood wheezing illnesses following acute respiratory syncytial virus infection in infancy. J
Allergy Clin Immunol, 2018. **142**(5): p. 1447–1456 e9.
27.   Bernard-Raichon, L., et al., *A Pulmonary Lactobacillus murinus Strain Induces Th17 and*
RORgammat(+) Regulatory T Cells and Reduces Lung Inflammation in Tuberculosis. J Immunol,
2021. **207**(7): p. 1857–1870.
28.   de Steenhuijsen Piters, W.A.A., J. Binkowska, and D. Bogaert, *Early Life Microbiota and*
Respiratory Tract Infections. Cell Host Microbe, 2020. **28**(2): p. 223–232.
29.   Rameix-Welti, M.A., et al., *Visualizing the replication of respiratory syncytial virus in cells*
and in living mice. Nat Commun, 2014. **5**: p. 5104.
30.   Descamps, D., et al., *Toll-like receptor 5 (TLR5), IL-1beta secretion, and asparagine*
endopeptidase are critical factors for alveolar macrophage phagocytosis and bacterial killing. Proc
Natl Acad Sci U S A, 2012. **109**(5): p. 1619–24.
31.   Drajac, C., et al., *Control of IFN-I responses by the aminopeptidase IRAP in neonatal*
C57BL/6 alveolar macrophages during RSV infection. Mucosal Immunol, 2021. **14**(4): p. 949–962.
32.   Liu, G., et al., *Use of precision cut lung slices as a translational model for the study of lung*
biology. Respir Res, 2019. **20**(1): p. 162.
33.   Makris, S., et al., *Alveolar Macrophages Can Control Respiratory Syncytial Virus Infection*
in the Absence of Type I Interferons. J Innate Immun, 2016. **8**(5): p. 452–63.
34.   Galani, I.E., et al., *Interferon-lambda Mediates Non-redundant Front-Line Antiviral*
Protection against Influenza Virus Infection without Compromising Host Fitness. Immunity, 2017.
46(5): p. 875–890 e6.
35.   Kota, S., et al., *Role of human beta-defensin-2 during tumor necrosis factor-alpha/NF-*
kappaB-mediated innate antiviral response against human respiratory syncytial virus. J Biol Chem,
2008. **283**(33): p. 22417–29.
36.   Drajac, C., et al., *Pulmonary Susceptibility of Neonates to Respiratory Syncytial Virus*
Infection: A Problem of Innate Immunity? J Immunol Res, 2017. **2017**: p. 8734504.
37.   Fonseca, W., N.W. Lukacs, and C. Ptaschinski, *Factors Affecting the Immunity to*
Respiratory Syncytial Virus: From Epigenetics to Microbiome. Front Immunol, 2018. **9**: p. 226.
38.   McNamara, P.S., et al., *Interleukin 9 production in the lungs of infants with severe*
respiratory syncytial virus bronchiolitis. Lancet, 2004. **363**(9414): p. 1031–7.
39.   Dodd, J.S., et al., *IL-9 regulates pathology during primary and memory responses to*
respiratory syncytial virus infection. J Immunol, 2009. **183**(11): p. 7006–13.
40.   Dean, L.S., et al., *I don't know about you, but I'm feeling IL-22.* Cytokine Growth Factor
Rev, 2024. **80**: p. 1–11.
41.   Das, S., et al., *Interleukin-22 Inhibits Respiratory Syncytial Virus Production by Blocking*
Virus-Mediated Subversion of Cellular Autophagy. iScience, 2020. **23**(7): p. 101256.
42.   Tahamtan, A., et al., *Neutrophils in respiratory syncytial virus infection: From harmful*
effects to therapeutic opportunities. Br J Pharmacol, 2021. **178**(3): p. 515–530.
43.   Ozcam, M. and S.V. Lynch, *The gut-airway microbiome axis in health and respiratory*
diseases. Nat Rev Microbiol, 2024. **22**(8): p. 492–506.
44.   Antunes, K.H., et al., *Microbiota-derived acetate protects against respiratory syncytial virus*
infection through a GPR43-type 1 interferon response. Nat Commun, 2019. **10**(1): p. 3273.
45.   Ji, J.J., et al., *Probiotics protect against RSV infection by modulating the microbiota-*
alveolar-macrophage axis. Acta Pharmacol Sin, 2021. **42**(10): p. 1630–1641.
46.   Villena, J., et al., *Orally administered Lactobacillus rhamnosus modulates the respiratory*
immune response triggered by the viral pathogen-associated molecular pattern poly(I:C). BMC
Immunol, 2012. **13**: p. 53.
47.   Kanmani, P., et al., *Respiratory Commensal Bacteria Corynebacterium*
pseudodiphtheriticum Improves Resistance of Infant Mice to Respiratory Syncytial Virus and
Streptococcus pneumoniae Superinfection. Front Microbiol, 2017. **8**: p. 1613.
48.   Antunes, K.H., et al., *Airway Administration of Bacterial Lysate OM-85 Protects Mice*
Against Respiratory Syncytial Virus Infection. Front Immunol, 2022. **13**: p. 867022.

49.   Nougayrede, J.P., et al., *A Toxic Friend: Genotoxic and Mutagenic Activity of the Probiotic*
Strain Escherichia coli Nissle 1917. mSphere, 2021. **6**(4): p. e0062421.
50.   Perdijk, O., et al., *Antibiotic-driven dysbiosis in early life disrupts indole-3-propionic acid*
production and exacerbates allergic airway inflammation in adulthood. Immunity, 2024. **57**(8): p.
1939–1954 e7.

Figure legends

Figure. 1: Cytokine production by precision-cut lung slices (PCLS) from neonatal mice varies in response to exposure to lung primo-colonizing bacteria. **A**. Experimental design. The lungs from 6-day-old mice were isolated and cut with a Krumdieck tissue slicer to obtain PCLS. They were next exposed to bacteria (50 CFU/well) or cultured in medium control. Supernatants were collected 16h after stimulation with bacterial strains and PCLS were lysed. **B**. Secretion of cytokines associated with type 1 immune response (IFN-β, IL-12p70). Results are expressed as mean ± SEM and represent a pool of five independent experiments (n = 2-5 biological replicate samples/condition). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ and ****$P < 0.0001$ between the control medium and the strain-exposed condition (ANOVA, Kruskal-Wallis test followed by Dunn's post-hoc test). **C**. The heat map illustrates the fold-change (FC) of cytokine production induced after bacterial exposure. For each strain and cytokine secretion, the FC were calculated using the following formula: FC = (average of the stimulated conditions)/(average of the control conditions). FC were entered into Morpheus online software (Broad Institute) to obtain the heatmaps.

Figure. 2: RSV replication in neonatal PCLS is differentially affected by primo-colonizing commensal bacteria. **A**. Experimental design. The lungs from 6-day-old mice were isolated and cut with a Krumdieck tissue slicer to obtain PCLS. They were next exposed to strains 16, 17, 24 or 25 (50 CFU/well) or cultured in control medium. PCLS were infected with RSV expressing luciferase (RSV-Luc, 7.187×10^4 PFU/well) or mock control (-) 3h after bacterial stimulation. Supernatants were collected 16h after bacterial stimulation and PCLS were lysed. **B**. Cytotoxicity observed in PCLS was measured by LDH activity assay. Results are expressed as the mean ± SEM and represent a pool of four independent experiments (n = 5-18 per group in simplicate). **C**. RSV-Luc replication was assessed by measuring luciferase activity in lysates of PCLS exposed or unexposed to strains. Results are expressed as mean ± SEM and represent a pool of four independent experiments (n = 6-21 per group in duplicate). *$P < 0.05$ and ****$P < 0.0001$ between the control medium or the strain-exposed condition (ANOVA, Kruskal-Wallis test followed by Dunn's post-hoc test).

Figure. 3: Cytokines and type I interferon (IFN-I) production by RSV-infected alveolar macrophages (AMs) from neonatal mice is enhanced in response to exposure to strain 17. **A**. Experimental design. AMs isolated from BAL of neonatal mice were exposed to strain 17 (50 CFU/well) or cultured in medium control. AMs were infected with rRSV-long (MOI = 5) 3h after bacterial stimulation. Supernatants were collected 16h after bacterial stimulation and AMs were lysed. The production of pro-inflammatory cytokines (IL-6 and TNF-α, **B**) and IFN-I (IFN-α and IFN-β, **C**) was measured in

supernatants using multiplex immunoassay. Data are representative of two independent experiments.

Means ± SEM are represented. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ and ****$P < 0.0001$ (2way

ANOVA with Tukey's multiple comparisons test).

Figure. 4: Early-life airway administration of strain 17 restricts RSV replication in neonatal mice. **A**.

Experimental design. Neonatal mice were treated with different quantities of strain 17 (10^3, 10^5 or

10^6 CFU) or PBS vehicle at 3 and 5-days old (i.n., 10 µL). At 6-days old, mice were infected with

RSV-Luc (7.2 x 10^4 PFU, 10 µL, i.n) and were euthanized at 1 d.p.i or 2 d.p.i. to assess viral

replication by luciferase activity assay in the lysats of nasal cavities (**B**) and lungs (**C**). **D**. The viral

replication was evaluated by quantification of the RSV *N gene* expression. The level of expression

was calculated by the formula $2^{-\Delta Cq}$ with $\Delta CQ = Cq_{N gene} - Cq_{reference genes}$. Results are expressed as mean

± SEM. Data are representative of two independent experiment (n = 5-8 per group). *$P < 0.05$, **P

< 0.01, ***$P < 0.001$, Mann-Whitney test.

Figure. 5: Early-life airway administration of strain 17 enhances neonatal pulmonary immunity

against RSV. Neonatal mice were treated with different quantities of strain 17 (10^3, 10^5 or 10^6 CFU)

or PBS vehicle at 3 and 5-days old (i.n., 10 µL). At 6 days old, mice were infected with RSV-Luc

(7.2 x 10^4 PFU, 10 µL, i.n) and were euthanized at 1 or 2 d.p.i. **A**. *IFN-I gene* expression was

measured by qRT-PCR in lung lysates at 1 d.p.i.. Gene expression levels were calculated using the 2^{-}

$^{\Delta Cq}$ method, with Cq values normalized to housekeeping genes. **B**. Cell counts were performed on

BAL samples at 1 or 2 d.p.i.. Cell infiltration in BALs was assessed by May-Grünwald-Giemsa

staining to identify and differentiate macrophages (**C**) and polynuclear neutrophils (**D**) at 1 or 2 d.p.i..

Results are expressed as mean ± SEM. Data are representative of two independent experiments (n =

5-8 per group). *$P < 0.05$, **$P < 0.01$, Mann-Whitney test.

Figure. 6 : Early-life airway administration of strain 17 attenuates the long-term pulmonary

consequences of neonatal RSV infection during reinfection in adulthood. **A**. Experimental design.

Three-day-old mice were treated intranasally with bacterium 17 (10^6 CFU, i.n., 10 µL) or PBS vehicle

at days 3, 5, 8, and 10 days of age. At 6 days old, mice were infected intranasally with RSV-A2 (2.3

$\times 10^4$ PFU in 10 µL), followed by a second infection with RSV-A2 (2.3×10^5 PFU in 100 µL, i.n.) at

8 weeks of age. Mice were euthanized at 7, 9 or 14 days post-reinfection (d.p.i.) to assess cellular

infiltration in the BALs and blood was collected. **B**. Weight evolution after the second infection.

Results are expressed as mean ± SEM and presented as a percentage of the starting weight (n = 4 per

group). **C-D**. Anti-RSV antibody assays. Serum was used to quantify anti-RSV IgG1 (**C**) or IgG2a

(**D**) antibodies by ELISA. Results are expressed as the mean ± SEM (n = 4-5 per group). **P < 0.05*,

P < 0.01, Mann-Whitney test. **E-F. The infiltration of eosinophils in BALs was assessed by May-

Grünwald-Giemsa staining at 7 or 9 d.p.i.. Results are expressed as mean ± SEM. Data are

representative of two independent experiments (n = 5-9 per group). **P < 0.05*, ***P < 0.01*, Mann-

Whitney test.

Figure. 7: RSV replication in human primary airway epithelial cells is reduced by the pre-exposure

to strain 17. **A**. Experimental design. Human primary bronchial epithelial cells cultured at the air

liquid interface (MucilAir™) were first exposed to strain 17 (10^3 CFU/well) for 3h, followed by RSV

expressing mCherry (RSV-mCherry, MOI = 0.1) infection or an incubation with mock for a further

3h. Supernatants and cell lysates were collected 48h after incubation. RSV replication was evaluated

by detecting mCherry using microscopy (**B**, three images representative of six inserts) or a

fluorescence plate reader (**C**). Results are expressed as the mean fluorescence intensity (MFI)

observed in lysats of RSV-mCherry-infected cells, after subtraction of the background MFI from

mock-infected controls (± SEM), and represent a pool of two independent experiments (n = 3 inserts

per group per experiment, ***P < 0.01*, Mann-Whitney test). **D**. Cytotoxicity observed in MucilAir™

was measured by LDH activity assay. Results are expressed as mean ± SEM. **E**. Bioactive IFN-λ was

evaluated by measuring the response of HEK-Blue™ IFN-λ cells to the supernatants. SEAP activity

was assessed using QUANTI-Blue™ Solution. The optical density (OD) at 630 nm is shown as mean

± SEM. **F**. β-defensin 2 secretion in supernatants was evaluated by ELISA. **E-F**. Data represent a

pool of two independent experiments (n = 9 inserts per group, ****P < 0.001*, *****P < 0.0001*, 2way

ANOVA with Tukey's multiple comparisons test).

Early-Life Commensal Lung Bacteria Modulate Immune Responses and Susceptibility ito RSV Infection in Neonatal Mice

Figures

Figure 1

Figure 2

A

BALB/c neonates
(6 days of life)

Lung extraction

Krumdieck tissue slicer

Precision cut lung slices (PCLS)
2 per wells

Agarose 1,5%

Medium or Strain (16, 17, 24 or 25)
(50 CFU/well)

-24h   0h   3h   16h

Mock or RSV-Luc

- Cytotoxicity (LDH assay)
- RSV replication (Luciferase activity)

B

Cytotoxicity (%)

RSV-Luc   - +   - +   - +   - +   - +
Strain   -   16   17   24   25

C

Viral replication
(luciferase activity (radiance) normalized to PCLS protein content (mg))

****   ****
*   *

RSV-Luc   +   +   +   +   +
Strain   -   16   17   24   25

Figure 3

A

B

IL-6

Concentration in supernatant (pg/mL)

RSV   −   +   −   +
       Medium   Strain 17

ns
ns
****
****

TNF-α

Concentration in supernatant (pg/mL)

RSV   −   +   −   +
       Medium   Strain 17

ns
ns
****
****

C

IFN-α

Concentration in supernatant (pg/mL)

RSV   −   +   −   +
       Medium   Strain 17

*
ns
***
****

IFN-β

Concentration in supernatant (pg/mL)

RSV   −   +   −   +
       Medium   Strain 17

**
ns
****
****

Figure 4

A

PBS or Strain 17
(10^3, 10^5, 10^6 CFU/well, 10 µL, i.n)

BALB/c neonates
(3 days of life)

D3 — D5 — D6 — D7 — D8

1 d.p.i.   *2 d.p.i.*

Mock or RSV-Luc

- **RSV replication** (Luciferase activity, qRT-PCR)
- **IFN-I response** (qRT-PCR)
- **Cellular infiltration** (MGG coloration)

B

Nasal cavity (1 d.p.i.)

Viral replication (Avg Radiance/mg of tissues)

| RSV-Luc | - | + | + | + | + |
| Strain 17 | - | - | 10^3 | 10^5 | 10^6 |

Nasal cavity (2 d.p.i.)

Viral replication (Avg Radiance/mg of tissues)

| RSV-Luc | - | + | + | + | + |
| Strain 17 | - | - | 10^3 | 10^5 | 10^6 |

C

Lung (1 d.p.i.)

Viral replication (Avg Radiance/mg of tissues)

| RSV-Luc | - | + | + | + | + |
| Strain 17 | - | - | 10^3 | 10^5 | 10^6 |

Lung (2 d.p.i.)

RSV replication (Avg Radiance/mg of tissues)

| RSV-Luc | - | + | + | + | + |
| Strain 17 | - | - | 10^3 | 10^5 | 10^6 |

D

Lung (1 d.p.i.)

Viral replication (N gene expression ($2^{-\Delta Cq}$))

| RSV-Luc | - | + | + | + | + |
| Strain 17 | - | - | 10^3 | 10^5 | 10^6 |

Lung (2 d.p.i.)

Viral replication (N gene expression ($2^{-\Delta Cq}$))

| RSV-Luc | - | + | + | + | + |
| Strain 17 | - | - | 10^3 | 10^5 | 10^6 |

Figure 5

A

Lung lysates
IFN-α (1 d.p.i.)

Lung lysates
IFN-β (1 d.p.i.)

B

BAL (1 d.p.i.)

BAL (2 d.p.i.)

C

BAL (1 d.p.i.)

BAL (2 d.p.i.)

D

BAL (1 d.p.i.)

BAL (2 d.p.i.)

Figure 6

A

B

C

D

E

F

Figure 7

A

Medium or Strain 17

Human bronchial epithelial cells cultured at the air liquid interface

0h   3h   48h

Mock or RSV-mCherry

- **Cytotoxicity** (LDH assay)
- **RSV replication** (fluorescence)
- **Innate immune response** (Immunoassays and multiPlex)

B

Medium   Strain 17

24h

48h

C

mCherry

Fluorescence intensity (MFI)

5×10^4
4×10^4
3×10^4
2×10^4
1×10^4
0

**

RSV-Cherry   +   +
              Medium   Strain 17

D

Cytotoxicity (%)

RSV-Cherry   -   +   -   +
              Medium   Strain 17

E

Bioactive IFN-λ

OD (630 nm)

1.50
1.25
1.00
0.75
0.15
0.10
0.05
0.00

****   ****
        ****

RSV-Cherry   -   +   -   +
              Medium   Strain 17

F

β-defensin 2

Concentration (pg/mL)

6000
4000
2000
0

***
$p = 0.0561$
$p = 0.4193$
ns

RSV-Cherry   -   +   -   +
              Medium   Strain 17