Mitophagy-related prohibitin 2 is an orthoflavivirus restriction factor targeted for degradation during infection

Imanol Rodrigo,[a,b,c,*] María José Romero de Ávila,[a,b,c,*] Belén García-Navarro,[a,b,c] Martín Arellano,[a,b,c] Laura Albentosa-González,[a,b,c,d] Pilar Clemente-Casares,[a,b,c,d] Edward Emmott,[e,#] Armando Arias,[a,b,c,d,f,#]

Affiliations:

[a]Unidad de Medicina Molecular, Instituto de Biomedicina de UCLM (IB-UCLM), Universidad de Castilla-La Mancha (UCLM), Albacete, Spain.

[b]Unidad de Biomedicina UCLM-CSIC, Albacete, Spain.

[c]Instituto de Investigación Sanitaria de Castilla-La Mancha (IDISCAM), Spain.

[d]Facultad de Farmacia, UCLM, Albacete, Spain.

[e]Centre for Proteome Research, Department of Biochemistry, Cell & Systems Biology, Institute of Systems, Molecular & Integrative Biology, Biosciences Building, Crown Street, University of Liverpool, Liverpool, L69 7ZB, United Kingdom.

[f]Escuela Técnica Superior de Ingenieros Agrónomos y de Montes y Biotecnología (ETSIAMB), UCLM, Albacete, Spain.

Running title: Prohibitin 2 restricts orthoflaviviral replication

23   # Address correspondence to:

24   Armando Arias, armando.arias@uclm.es; Edward Emmott, e.emmott@liverpool.ac.uk

25   * These authors equally contribute to this work

29   Keywords: Zika, Usutu, host-virus interactions, NS5 polymerase, NS2B/NS3 protease,

30   myoferlin, restriction factors, proviral factors.

1. ABSTRACT

Orthoflaviviruses (OFVs) are the primary cause of arboviral disease worldwide, leading to large numbers of hospitalisations, sequelae and deaths associated. Viral genome replication is catalysed by non-structural protein 5 (NS5), the largest and most conserved protein in these viruses. Here, we used quantitative proteomics to identify host cell interactors of Zika virus (ZIKV), Usutu virus (USUV), and West Nile virus (WNV) NS5 proteins. A total of 141 proteins were detected in the analysis, being 26 of them associated with all three NS5s. Downregulation of myoferlin and LGALS3BP led to decreases in the virus yields, indicating that their presence is needed for efficient virus replication. Conversely, silencing of mitophagy-related prohibitin 2 (PHB2) resulted in significant increases in ZIKV and USUV titres and genomic RNA. Supporting a connection between this antiviral behaviour and mitophagy, an inhibitor of this pathway also led to larger virus titres and viral RNA copies. During infection, PHB2 protein levels gradually decreased, becoming transiently undetectable, and suggesting that it is specifically targeted by the virus for degradation. Ectopic expression of NS5 alone did not affect PHB2 intracellular abundance, however a significant decrease was observed in cells expressing the viral protease NS2B/NS3. Its elimination, was more pronounced in cells where both viral enzymes were co-expressed, supporting that NS5 could be assisting PHB2 recognition by NS2B/NS3. Overall, our results support that PHB2 is an orthoflaviviral restriction factor which is targeted for degradation to favour viral replication.

2. IMPORTANCE

Orthoflaviviruses (OFVs) are one of the main causes of disease transmitted by mosquitoes, with millions of infections occurring annually, and large numbers of deaths associated. Some important members include viruses causing dengue, yellow fever, and Zika-associated disease (e.g. microcephaly). OFVs contain a genome which is copied by a highly conserved viral protein termed NS5. Prohibitin 2 (PHB2) is a host cellular protein involved in different cellular processes including mitophagy, a mechanism to remove damaged mitochondria. Here, we show that human PHB2 interacts with orthoflaviviral NS5. Depletion of PHB2 results in increased viral yields, suggesting that this protein restricts orthoflaviviral infection. We found that PHB2 is targeted for degradation by the orthoflaviviral protease in a process aided by NS5, supporting that several viral proteins act concertedly to eliminate PHB2, counteracting its antiviral activity, and thus favouring infection.

3. INTRODUCTION

Over 50 per cent of the entire human population is at risk of infection by members of the *Orthoflavivirus* genus (1–3). Global warming and a broader geographic dissemination of competent vectors are factors contributing to a steady increase in the incidence and expansion of orthoflaviviral disease (4). Most human orthoflaviviruses (OFVs) are transmitted by mosquitoes, with estimates suggesting ~100 million infections occur annually, leading to over 100,000 associated deaths (4, 5). The economic impact caused by OFVs is sizeable, with data supporting that Zika virus (ZIKV) and dengue virus (DENV) infections combined lead to >3 billion USD losses annually (6). ZIKV was declared a human pathogen of international concern in 2016, following large epidemic outbreaks that primarily affected the Americas. Now, the virus is present in six continents and more than 90 countries (7–9). Sequelae associated with infection during pregnancy can lead to severe congenital malformations, being microcephaly the most common syndrome (10). Although generally asymptomatic, infection in adults can cause clinical manifestations of different degree of severity, including meningitis, encephalitis and the Guillain-Barré syndrome (11, 12).

OFVs infecting birds as their natural vertebrate hosts are an additional source of public health concern (13). Even though humans are typically dead-end hosts for avian OFVs, neuroinvasive infection occur in a significant number of cases. West Nile virus (WNV) is the most relevant avian OFV from a public health perspective, with thousands of cases of severe disease and hundreds of fatalities annually (14). Despite that it generally leads to milder outcomes, a gradual increase in the number of neurological disorders associated with Usutu virus (USUV) has also been documented (15–18). WNV and USUV are genetically similar and exhibit serological cross-reactivity, which compromises their correct diagnosis in areas where cocirculation occur. Some lines of evidence imply that USUV

infection in humans has been historically misdiagnosed as WNV, and its true impact

underestimated (17, 19).

The OFV genome is a positive-stranded RNA molecule which is directly translated into a

single polyprotein. This polypeptide is proteolytically processed by both cellular activities

and the viral NS2B/NS3 protease, resulting in the release of different structural and non-

structural proteins (20, 21). Non-structural protein 5 (NS5) contains a C-terminal RNA-

dependent RNA polymerase (RdRp) domain, for the synthesis of new viral genomes, and

an N-terminal methyltransferase (MTase) domain for the 5' end capping of the viral

genome (22). OFV replication factories are assembled inside partially secluded

invaginations of endoplasmic reticulum (ER) membranes where viral RNA is isolated from

host antiviral sensors while allowing the recruitment of replication factors (23–25). In

addition to viral proteins, forty-six cellular proteins have been identified inside these

vesicles, most of them displaying proviral activities (23).

The use of drugs targeting host factors required for viral replication has gained momentum

in the antiviral research field (26, 27). In addition to viral replication, NS5 has been

revealed as a multifaceted protein, engaging in diverse processes that include suppression

of innate immunity, transcription reprogramming, and interference with the cell cycle

progression (28–30). Several NS5 cellular partners have been identified, including proviral

(23, 27, 31–34) and antiviral activities (35–37). Recent data have revealed that ZIKV NS5

binds to host protein Ajuba, preventing mitophagy activation (38). Mitophagy is a type of

selective autophagy to remove irreparably damaged or depolarised mitochondria and

promote cell survival. Since damaged mitochondria can activate innate immunity and

proinflammatory responses, many viruses modulate the activity of this pathway (39). Ajuba

knockdown *in vivo* resulted in enhanced inflammatory responses, leading to larger ZIKV

dissemination in mouse tissues (38). These results convey that mitophagy suppression by

NS5 favours viral spread in the host.

During mitophagy activation, PINK1 is accumulated in the outer mitochondrial membrane

(OMM), leading to the recruitment of E3-ubiquitin ligase Parkin (40). Parkin ubiquitylates

different OMM proteins, attracting other factors needed to initiate mitophagy (41).

Prohibitin 2 (PHB2), a multitasking protein found in several subcellular locations, has been

recently identified as a mitophagy receptor (40, 42). PHB2 interacts with SQSTM1 and

LC3 in the inner mitochondrial membrane to drive damaged mitochondria to the

autophagosome (43). Besides mitophagy, mitochondrial PHB2 in complex with prohibitin 1

(PHB1) has been proposed as major contributor to the MAVS-mediated antiviral signal,

leading to the activation of type I interferon (IFN-I) responses (44). PHB2 is also found in

the nucleus where it acts as a repressor of diverse transcription factors such as the

oestrogen receptor (ESR), MyoD and MEF2 (45–48).

In the present study, we identified 141 host factors that interact with the NS5 proteins of

ZIKV, WNV, and USUV. We found that 26 of these factors were common interactors for all

three NS5s, including PHB2. PHB2 gene silencing led to increases in both virus titres and

genomic RNA, which is suggestive of a viral restriction activity. During both ZIKV and

USUV infection, PHB2 abundance gradually decreased, becoming transiently

undetectable, and supporting that OFVs specifically target PHB2 to promote viral

replication. Ectopic expression of NS2B/NS3 protease but not NS5 led to PHB2

degradation, although its elimination was more efficient when both viral proteins were co-

expressed. In the study, we identified other NS5 interactors that are necessary for efficient

virus infection, such as myoferlin (MYOF) and LGALS3BP, encouraging the use of specific

drugs against them.

4. MATERIALS AND METHODS

4.1. Cells, viruses, drugs and cell viability assays

The origin of ZIKV and USUV strains, and protocols for infection have been detailed in previous manuscripts (32, 49–51). Cell culture infections and virus titrations by 50% tissue culture infectious dose (TCID$_{50}$) assays have been performed following protocols described before (31, 32, 49). The antiviral efficacy of different host-targeting drugs has been tested as described in previous manuscripts (32, 51). To these assays, WJ460 [3-(3-(2-methoxyphenyl)-4-oxo-2-thiazolidinyl)-N-(4-phenylbutyl) benzamide] and mdivi-1 [3-(2,4-dichloro-5-methoxyphenyl)-2-mercaptoquinazolin-4(3H)-one] were purchased from MedChemExpress. The number of metabolically live cells after exposure to these compounds relative to mock-treated cultures, or after transfection with different host-targeting DsiRNA molecules relative to negative control DsiRNAs (not recognising human sequences) were obtained with the *CellTiterBlue* kit (Promega), as described before (31, 32).

4.2. Gene silencing with Dicer-substrate small-interfering RNAs (DsiRNAs)

Host-specific DsiRNA duplexes were obtained from Integrated DNA Technologies (IDT), with Information regarding their target sequences provided as supplemental material (Table S1). As a negative control, a non-specific DsiRNA (no target in the human genome), namely NonSP, was used. For gene silencing, 5 pmol of each DsiRNA were incubated during 20 min at room temperature in the presence of 0.25 µl of lipofectamine RNAiMAX (Invitrogen) and 50 µl of opti-MEM (Gibco). The mix was transferred into 96-plate wells, and then 1.5×10^4 A549 cells were added on top. Twenty-four hours after transfection, the supernatants were removed, and virus was inoculated at an MOI of 0.1 or

1 $TCID_{50}$/cell. Following adsorption, the inoculum was removed, cells washed, and fresh media containing 1% foetal bovine serum (FBS) added.

4.3. Viral RNA extraction and quantitative RT-PCR (RT-qPCR)

RNA was extracted from whole cell monolayers with the *NZY Total RNA Isolation* kit (NZYtech). ZIKV RNA was detected with a FAM-TAMRA probe targeting the E gene (1214 to 1244), and primers spanning positions 1192 to 1208 (forward) and 1269 to 1246 (reverse), while USUV RNA was detected using a FAM-TAMRA probe targeting the NS1 gene (3322 to 3345), and sense and antisense primers spanning positions 3288 to 3309 and 3372 to 3351, respectively. To the amplification of viral RNA, we used the *NZYSupreme One-step RT-qPCR Probe Master Mix (2x), ROX* kit (NZYtech), and RT-PCR conditions suggested by the manufacturer (15 min at 50ºC, 3 min at 95°C, followed by 40 cycles of 5s at 95°C and 45s at 60°C). To obtain a standard curve with known amounts of RNA, we produced *in vitro* transcripts of the target genes. Briefly, 4µl of whole RNA extracted from infected cells were reverse transcribed using *SuperScript III Reverse Transcriptase* (Invitrogen). Then, 3µl of cDNA were PCR-amplified using *Phire Hot Start II DNA Polymerase* (ThermoFisher), and primers spanning residues 949 to 967 (forward) and 1934 to 1915 (antisense) for ZIKV, and 2809 to 2829 (forward) and 3808 to 3787 (antisense) for USUV (both forward primers contained a T7 promoter. PCR amplicons were purified (*NZY GelPure* kit, NZYtech), and ligated into the PCR Blunt plasmid (*Zero Blunt PCR cloning* kit, ThermoFisher), and the linearised plasmids or directly the PCR amplicons were *in vitro* transcribed with the *mMESSAGE mMACHINE T7 Transcription* kit (Ambion). After DNAse treatment, the RNA transcript was purified and quantitated, and the standards were prepared with known amounts of each transcript in 100 ng/µl yeast RNA.

4.4. Relative quantification of host mRNAs and poly I:C treatment

To confirm host gene knockdown after DsiRNA transfection, host gene mRNA levels were

determined with *One-step NZYSpeedy RT-qPCR Green kit, ROX* kit (NZYtech) in a CFX

Duet Real Time PCR instrument (BioRad) and specific primer pairs, all provided by IDT

[reference numbers (Hs.PT.58.): 27059869, 19355804, 40826680, 2219540, 3098270,

40209489, 456578.g, 3838519.g, 38465792, 22859503, 22973334, 27422678.g,

40506149, 40261049]. As a a housekeeping gene we used primers amplifying the GAPDH

transcripts (Hs.PT.39a.22214836). The efficiency of downregulation for each transcript

was calculated following the $2^{-\Delta\Delta CT}$ method (32, 52).

To stimulate IFN-I expression, A549 cells were transfected with *Poly (I:C) (LMW)*

(Invivogen), using Lipofectamine 3000 in the absence of the P3000 reagent (Invitrogen).

Twenty-four hours prior poly I:C stimulation, cells were transfected with DsiRNAs (either

targeting PHB2 or a non-specific DsiRNA), as explained above. Total RNA was extracted

from poly I:C-stimulated cells to analyse IFN-β gene transcription (relative IFNB1 mRNA; a

primer pair purchased from IDT, reference Hs.PT.58.39481063.g).

4.5. Vectors and mutagenesis

Constructs expressing HA-tagged versions of full-length NS5 (HA-NS5) or the RdRp

domain alone (HA-RdRp) from ZIKV, USUV, and WNV were used as documented in

previous manuscripts (31, 32, 53). To obtain the ZIKV NS2B/NS3 protease vector (pZIKV-

NS2B3), we used a pcDNA3 clone expressing ZIKV RdRp fused to an HA tag and the

EYFP (pZIKV-HA-RdRp-EYFP) as a backbone. We swapped the insert in this vector by

the corresponding NS2B3-coding region, using the *In-Fusion HD EcoDry Cloning Kit*

(Takara Bio), using specific reverse (5'-AGCGTAATCTGGAACATCGTATGGGTA-3') and

forward (5'-GATAACTCGAGCATGCATCTAGAG-3') primers, leading to the amplification

of the vector devoid of the RdRp-EYFP insert. To the amplification of the NS2B3 coding

region we used ZIKV primers spanning genomic positions 4224 to 4244 (forward) and

6464 to 6444 (antisense). Both primers contained 15-nt tails overlapping with the linear

vector, enabling the ligation of the insert by recombination.

4.6. Co-immunoprecipitation and mass spectrometry

For coimmunoprecipitation assays, HA-NS5 or HA-RdRp plasmids were transfected in

A549 or 293T cells with either the *Lipofectamine 3000* kit (Invitrogen) or the *FuGENE 4K*

reagent (Promega). At 24 to 48 h post-transfection, cells were scrapped on ice-cold PBS,

washed in PBS twice, spun at 300x *g* for 5 min, and the pellet resuspended at 4°C in 450

µl of RIPA buffer (10 mM Tris-HCl pH 7.5; 150 mM NaCl; 0.5 mM EDTA; 1% SDS; 1%

Triton X-100; 1% deoxycholate) in the presence of 1x *Complete Mini EDTA-free protease*

inhibitor cocktail (Merck) and 250 U/ml benzonase (Sigma). To remove the cellular debris,

the lysate was spun during 10 mins at 15000x *g*. Protein concentration in the supernatants

was determined with the *BCA Protein Assay Kit* (Pierce), and all samples normalised to

the same concentration in a final volume of 400 µl. The samples were then equilibrated in

1.5 volumes (600 µl) of dilution buffer (10 mM Tris-HCl pH 7.5; 150 mM NaCl; 0.5 mM

EDTA), and then mixed with either agarose or magnetic anti-HA beads (Pierce), and

incubated in a rotary shaker overnight at 4ºC. The beads were then either magnetically

separated (magnetic) or pelleted by centrifugation (agarose), the supernatant eliminated,

and washed four times in 1x dilution buffer containing 0.05% Nonidet P40 substitute (Fluka

BioChemika). Finally, the co-immunoprecipitated proteins were released from the beads

by heating at 90°C for 5 min in 1x *Bolt LDS Sample Buffer* (Life Technologies).

For the identification of host cell interactors by mass spectrometry, HA-NS5 proteins

expressed in A549 cells were pulled down with anti-HA magnetic beads, as described

above. The pulldowns were SP3-precipitated (54), trypsin digested and labelled with

TMTpro. LC-MS/MS was performed on a Dionex U3000 UHPLC, in-line with a Q-Exactive-

HF mass spectrometer as described previously (55). Data were analysed with MaxQuant

1.6.7.0 (56). N-terminal protein acetylation and methionine were set as variable

modifications, enzyme specificity was set to trypsin with up to two missed cleavages. Data

were searched against fasta files containing the Swissprot human proteome entries

(20,300 entries, downloaded 2020/03/12) and a custom fasta file containing the HA-tagged

NS5 sequences. For each NS5, four independent transfections and co-IPs were

performed, one control was removed at the data analysis stage as an outlier (Fig. S1). To

identify those factors that were significantly enriched in the IPs, four independent plates

were mock-transfected in parallel with an empty vector (pcDNA3). NS5-interacting factors

were identified using the *Perseus software* version 2.0.6.0., as previously described (57).

A list with all host factors that are significantly enriched in ZIKV, USUV and WNV HA-NS5

co-IPs followed by TMT-based proteomics are provided as supporting material (Tables S2

– S4). The mass spectrometry data, including .fasta files has been submitted to

proteomeXchange via the PRIDE repository and has accession number PXD071636.

4.7.Western blot analysis

To the detection of host and virus proteins, we followed protocols available in previous

studies (32). Antibodies against the HA epitope (BioLegend, reference 901501), the FLAG

epitope (Proteintech, reference 20543-1-AP), PHB2 (Invitrogen, PA5-21417), STAT2 (Cell

Signaling, 72604S), ZIKV NS3 (Invitrogen, PA5-143436) and ZIKV NS5 (Genetex,

GTX133328) were used. As loading controls, we purchased HRP-conjugated antibodies

against GAPDH (Genetex, GTX627408-01) and tubulin (Cell Signaling, 12351S).

4.8. Gene ontology analysis

Gene Ontology (GO) enrichment analysis was performed on differentially abundant
proteins (FDR < 0.05) using g:Profiler with the Benjamini-Hochberg correction. Redundant
GO Biological Process (BP) terms were reduced using rrvgo (v1.10.0) with a Wang
semantic similarity threshold of 0.7. GO Cellular Component (CC) terms with <5 genes
were excluded. Remaining terms were grouped only when sub-compartments shared a
parent GO category (e.g., mitochondrial compartments) and the combined gene count
remained statistically robust (≥5 genes). A summary of all enrichment analysis to the
identification of BP and CC terms are provided as supporting information (Tables S5 and
S6).

5. RESULTS

5.1. Identification of host factors associated with orthoflaviviral NS5s

Full-length versions of NS5 proteins were expressed in A549 cells, and their interacting
host factors isolated as described in the Methods section. Mass spectrometry analyses
resulted in the identification of 72, 63 and 77 cellular proteins significantly enriched in
pulldowns of ZIKV, USUV and WNV NS5s, respectively (Fig. 1, Tables S1 – S4). The GO
Biological Process enrichment analysis revealed that most of them were involved in RNA
metabolism (splicing and processing), ribosome biogenesis and protein translation,
chromatin assembly and chromosome organisation, or respiration and proton transport in
the mitochondria (Fig. 1 and 2, and Fig. S2). The intersection of the three sets revealed
that 26 proteins were common interactors for ZIKV, WNV and USUV NS5s (Fig. 1A and
Fig. S2). To the identification of those factors that could be more biologically relevant, we
re-examined the intersection, but only using the top 50 significantly enriched proteins in
each pulldown. This resulted in 14 host factors shared by all three sets (Fig. 2, Tables S2

297 – S4). Most of them are mitochondrial proteins (ATP5B, PHB2, PGAM5, FH and SFXN1)

298 or participate in ribosome biogenesis and/or protein translation (RSL1D1, RPL21, RPL24,

299 RPL24).

5.2. Silencing of NS5-host partners identifies proviral and restriction activities

To elucidate the biological impact of these 14 host factors to OFVs, we examined the

effect of their silencing during infection (Fig. 3). All the DsiRNAs used in this study elicited

efficient gene downregulation, with over 70% reduction in mRNA levels compared to

controls (Fig. S3). Except for ribosomal proteins RPL21, RPL24 and RPL34, gene

silencing had no effect or caused only a modest decrease in cell viability (>75% live cells)

at 48 h post-transfection (Fig. S4). Downregulation of either PHB2 or ribosomal L1

domain-containing protein 1 (RSL1D1) resulted in larger virus titres in the supernatant of

both ZIKV- and USUV-infected cells, suggesting these two proteins are restriction factors

for the OFVs (Fig. 3). Downregulation of RPL21, PGAM5, SFXN1 and SMARCA5 also led

to significant increases in USUV but not in ZIKV titres, while DDX21 silencing resulted in

increased infectivity for ZIKV, but not for USUV.

Silencing of LGALS3BP and MYOF negatively affected both ZIKV and USUV, with

pronounced losses observed in the virus titres (Fig. 3). Additional analyses at different time

points and MOIs further confirmed this observation, conveying that these two are proviral

proteins (Fig. 4 and Fig. S5). Viral replication was significantly impaired when MYOF was

silenced, with 1,000- to 10,000-fold decreases in virus titres at 72 h post-infection (hpi).

These findings enticed us to evaluate whether specific drugs against MYOF, such as

WJ460, exhibit an antiviral behaviour. Although WJ460 specifically inhibits all known

MYOF cellular functions (58), we observed only a modest reduction in ZIKV titres, whilst

no effect on USUV was found (Fig. 4E – 4G). This apparent lack of antiviral effect was observed even at concentrations causing significant cell toxicity (40 to 160 µM), in contrast to data obtained with DsiRNAs.

5.3. PHB2 downregulation stimulates orthoflaviviral replication

Since the preliminary DsiRNA-based screening identified PHB2 and RSL1D1 as putative restriction factors (Fig. 3), we further examined how their silencing affected virus replication (Fig. 5 and Fig. S6). In agreement with our previous observations, downregulation of these genes resulted in significantly larger ZIKV and USUV titres at different time points and MOIs tested. Owing to precedent studies reporting that ZIKV NS5 interferes with mitophagy to promote viral dissemination (38), and that PHB2 is a mitophagy receptor, we aimed at further characterising this protein during the virus life cycle. In addition to enhanced virus titres, PHB2 silencing also led to significant increases in intracellular viral RNA (Figs 5G – 5J) that was also accompanied with increases in viral protein NS5 expression (Fig. 5B and Fig. S7), further supporting that PHB2 restricts viral replication. Treatment of infected cells with mdivi-1, an inhibitor of mitochondrial fission-related protein DRP1 which represses mitophagy (59), also resulted in both larger virus titres and genomic RNA (Fig. 6 and Fig. S8). Altogether, these results are pointing towards mitophagy and PHB2, a key receptor in this pathway, as restriction factors of the orthoflaviviral life cycle.

To examine whether larger titres and RNA levels could be related to an increase in virus particle release, we compared extracellular to intracellular viral RNA in PHB2-depleted and unspecific DsiRNA-treated cells (Fig. S9), but no significant differences were found. This observation imply that larger virus titres and genomic RNA in PHB2-depleted cells are likely the result of enhanced genome replication, and hence PHB2 is an OFV antagonist.

5.4. PHB2 is transiently eliminated during infection

Owing its apparent influence on viral replication, we decided to examine PHB2 protein

abundance dynamics during infection (Fig. 7). We found that PHB2 intensity gradually

decreased to become barely detectable at 48 hpi (< 3% relative to mock-infected cells) in

both ZIKV- and USUV-infected A549 cells (Fig. 7). At later time points, PHB2 intensity was

detected at a similar level than in control uninfected cells (101% and 58% at 60 hpi, in

ZIKV- and USUV-infected cells, respectively). PHB2 signal was also transiently reduced in

293T cells, supporting that is either targeted for degradation or its expression repressed

(Fig. 7H). Since mock- and virus-infected cells exhibit no significant differences in the

amount of PHB2 mRNA (Fig. S10), reduced protein levels are possibly connected to

degradation.

5.5. PHB2 depletion correlates with larger interferon expression in infected cells

Since PHB2 has been linked to MAVS-signalling, we considered that its degradation may

be interfering with IFN-I expression. To assess this possibility, we analysed IFN-β gene

(IFNB1) transcription in cells previously knocked down for PHB2. In both ZIKV- and USUV-

infected cultures, we found larger IFNB1 mRNA levels when PHB2 was silenced than in

controls transfected with a non-specific DsiRNA (Figs 8A – 8D), in contrast with previous

observations (44). A reasonable scenario is that enhanced IFNB1 mRNA transcription

could be incidental rather than directly prompted by PHB2 knockdown, given that more

viral RNA genomes are detected in PHB2-depleted cells (Figs. 5G and 5I). Thus, we

quantitated the relative amount of IFNB1 mRNA in cells stimulated with poly I:C, a non-

replicating double-stranded RNA. Although less pronounced than in infected cells, PHB2

silencing also resulted in increased IFNB1 transcription (Figs 8E – 8H). To further explore

whether IFN-I activation was connected to mitophagy inhibition, we analysed transcription

levels in ZIKV-infected cells treated with mdivi-1, but in this case, IFNB1 mRNA levels

were lower (Fig. 6D).

5.6. NS5 stimulates PHB2 degradation by the viral protease NS2B/NS3

Since both ZIKV and DENV NS5 directly promote host STAT2 degradation by the

proteasome (35, 37), and that NS5 specifically interacts with PHB2 (Fig. 2), direct PHB2

degradation by NS5 was a plausible scenario. Further supporting this possibility, ZIKV

NS5 and host PHB2 protein levels coincidentally decrease at 48 hpi in A549 cells when

compared to earlier time points (Fig. 7). We first confirmed that PHB2 and NS5 interact in

co-IP assays followed by Western blot (Fig. 9A), as inferred from mass spectrometry data

(Tables S2 – S4). PHB2 was also pulled down when the RdRp alone was used as a bait,

supporting that the interaction occurs through the polymerase domain (Fig. 9B). To assess

whether NS5 expression and PHB2 degradation were linked, we used increasing amounts

of all three NS5 constructs, but no significant change in PHB2 was detected (Fig. 9C and

Fig. S11). As expected, the expression of recombinant ZIKV NS5 significantly affected

STAT2 protein levels detected (Fig. 9C and Fig. S11), as previously documented (37).

Since NS5 is not sufficient for PHB2 degradation, we anticipated that other viral factors

might be involved. In addition to viral polyprotein processing, NS2B/NS3 also participates

in the proteolytic cleavage of different cellular proteins (60–62). We confirmed that the

ectopic expression of NS2B/NS3 precursor (NS2B3) resulted in decreased PHB2 levels

(Fig. 9D). To examine a possible synergistic activity of both viral proteins, we analysed the

effect of NS5 coexpression in cells transfected with a suboptimal amount of pZIKV-NS2B3.

When both NS2B/NS3confirming that, when both proteins are present, a lower amount of

PHB2 is detected (Figs 9E and 9F).

6. DISCUSSION

With the aim advancing our knowledge on the complex virus-host networks controlling genome replication, we characterised the cellular proteins interacting with three different NS5s. As a result of this analysis, we have obtained evidence supporting that PHB2, a mitophagy receptor, is a restriction factor for OFV replication. Previous studies have established a relationship between the OFVs and mitophagy (63). While its activation is necessary for Japanese encephalitis virus replication in a hepatoma cell line (64), mitophagy is silenced during DENV infection (65). For ZIKV, mitophagy activation has been associated with both proviral and antiviral activities (63). Ponia and colleagues demonstrated that ZIKV supresses mitophagy in astrocytes, in a process where NS5 interferes with host protein Ajuba (38). By contrast, Lee and Shin found that mitophagy inhibition, through RNAi-mediated silencing of PINK1, repressed viral replication, suggesting that mitophagy is distinctly regulated by different OFVs and/or cellular contexts (63, 66).

In our study, PHB2 depletion resulted in increased viral titres, genomic RNA and protein translation, sustaining that this protein is a restriction factor (Fig. 5). In addition to PHB2, we have identified other mitochondrial proteins as interactors, such as phosphoglycerate mutase family member 5 (PGAM5), which also plays an important role during mitophagy (40, 67), several subunits of the mitochondrial ATP synthase (e.g. ATP5F1B), fumarate hydratase (FH) and sideroflexin-1 (SFXN1). These findings are suggestive of mitochondria dynamics playing a crucial role during infection, and of NS5 as a factor subverting some of the processes controlled by these organelles. Despite this previous connection between ZIKV NS5 and mitophagy, it is conceivable, however, that viral restriction is linked to other cellular processes controlled by PHB2. A tantalising possibility is that PHB2 antiviral activity is related to its nuclear role as an ESR repressor (68–70). Supporting this

possibility, inhibition of the oestrogen-dependent signal by tamoxifen also affects orthoflaviviral replication (70). Additional studies on ESR and other transcription factors regulated by PHB2 are necessary to fully understand this antiviral activity.

PHB2 is transiently eliminated during infection in different human cell contexts (Fig. 7). Since both ZIKV and DENV NS5 proteins target STAT2 for degradation in a ubiquitin-dependent manner we initially considered this possibility for PHB2. In this process, orthoflaviviral NS5s act as a bridge between STAT2 and an E3 ubiquitin ligase (35, 37). We have recently documented that OFV NS5s also interact with the main E1 ubiquitin-activating enzyme UBA1 (32). However, we found that the ectopic expression of NS5 does not affect PHB2 levels, being required the viral protease NS2B/NS3 to its degradation (Fig. 9D). When NS2B/NS3 is expressed at suboptimal levels, NS5 is needed for PHB2 elimination, suggesting that NS5 operates as an adaptor between the viral protease and the host factor (Figs 9E and 9F). A physical interaction between NS5 and NS3 has been previously documented, and hence it is also expected to take place between NS5 and the NS2B/NS3 complex (71, 72). An attractive possibility is that NS5 could be acting as a protein bond between NS2B/NS3 and various host restriction factors, in addition to PHB2.

Previous studies have established a connection between PHB2 and MAVS antiviral signalling (44). Therefore, a forthcoming explanation would be that OFVs targeting PHB2 to degradation might help uncouple MAVS from IFN-I expression (44). Our data suggest however that PHB2 knockdown activates IFN-I transcription, with larger IFNB1 mRNA levels in either infected or poly I:C–stimulated cells (Fig. 8). Larger IFNB1 transcription levels observed in PHB2-silenced cells seems counterintuitive to the antiviral role proposed for this host protein (Fig. 8). This could be partially explained in the context of PHB2 fluctuating levels in the infected cells (Fig. 7). We hypothesise that transient PHB2

degradation at 48 hpi is necessary for efficient orthoflaviviral genome replication. However, subsequent PHB2 restoration to normal levels at 60 hpi is predicted to reactivate mitophagy, and thereby mitigate the ensuing inflammation and MAVS-mediated IFN-I responses (73–76).

Gene downregulation assays have identified other NS5-interacting host partners affecting orthoflaviviral replication. Larger virus titres were recovered in RSL1D1-depleted cells, also supporting an antiviral activity associated with this cellular factor (Fig. S6). RSL1D1 is an RNA-binding protein participating in multiple cellular processes, including antagonising senescence, proliferation and apoptosis, and acting as an autophagy suppressor in different types of cancer (77, 78). Conversely, MYOF and LGALS3BP downregulation led to significant decreases in both ZIKV and USUV titres, indicating that they may be contributing to critical steps of the life cycle (Fig. 4 and Fig. S4). MYOF is a multifunctional Ca2+ – binding protein which regulates vesicle trafficking and the cellular membrane repair (79, 80). Since OFVs assemble their replication sites on ER invaginations, requiring extensive membrane remodelling, membrane-repair activities might be relevant to their formation (81). WJ460, a promising anticancer drug targeting MYOF Ca2+ – binding C2D domain (79, 82), showed only a minor effect on virus titres, in contrast with DsiRNA-mediated silencing (Fig. 4). Recent studies suggest that MYOF might not be the primary host target of WJ460, and thus the activities observed for this drug not being totally associated with this host factor (83). LGALS3BP is a secreted glycoprotein that has been related to immune modulation, tumour progression, and extracellular matrix organization. Mice infected with WNV show upregulated LGALS3BP expression in the CNS (84). However, while KO mice for this gene exhibit reduced neuroinflammation and alterations in T cell populations, no significant differences in viral titres were observed (84). This

evidence suggests that LGALS3BP is not as necessary for WNV replication, as it appears to be for ZIKV and USUV, according to our study.

In addition to advance our understanding on the biological processes that control the orthoflaviviral life cycle, we deem these data presented here open new avenues for their therapeutic control. Approaches based on stimulating PHB2 antiviral activity or, alternatively, molecules triggering mitophagy activation, which are expected to antagonise viral replication, could be considered in future studies. The identification of the interacting epitopes between NS5 and other cellular partners can be of use to the rational design of host-targeting antivirals.

7. ACKNOWLEDGEMENTS

We thank José Joaquín Lorenzo for his technical assistance to this work.

8. FUNDING INFORMATION

This research was supported by grant SBPLY/24/180225/000240, from Junta de Comunidades de Castilla-La Mancha (JCCM), awarded to AA and LAG. AA is also the recipient of grants PID2022-137974OB-I00 and CNS2022-135258 from Ministerio de Ciencia, Innovación y Universidades (MICIU/AEI/10.13039/501100011033) and FEDER (EU), and grant SBPLY/21/180501/000076 from Junta de Comunidades de Castilla-La Mancha (JCCM). Work in AA's lab is also supported by core funding from UCLM (grant numbers 2025-GRIN-38349 and 2022-GRIN-34150). IR and MJRdA are, respectively, the recipients of fellowships 2020-PREDUCLM-16723 and 2024-INVGO-12348, both granted by UCLM. Research in EE's lab is supported by an AMS Springboard Award (SBF006\1008) funded by the Academy of Medical Sciences (AMS), the Wellcome Trust, the Government Department of Business, Energy and Industrial Strategy (BEIS), the

British Heart Foundation and Diabetes UK, an MRC NIRG (MR/X000885/1) and Wellcome

Career Development Award (227831/Z/23/Z).

9. FIGURE LEGENDS

Fig. 1. Summary of orthoflaviviral NS5-interacting network. (A) Overlap of host

proteins interacting with different orthoflaviviral NS5 proteins (ZIKV, USUV and WNV).

BioVenn was used to the graphic representation of the diagram (85). (B) Gene Ontology

analysis of cellular components (GO-CC). Related sub-compartments, i.e. mitochondrial

inner membrane (GO:0005743) and mitochondrial nucleoid (GO:0042645) were grouped

as 'Mitochondria' to reflect organelle-level localization. The pie chart displays the

proportional distribution of significantly enriched terms (adjusted $p < 0.05$) among all

factors identified in NS5-interacting datasets. (C) Heatmap of significantly enriched Gene

Ontology Biological Process (GO-BP) terms in co-IPs of cells transfected with either ZIKV,

USUV, or WNV HA-tagged NS5 proteins (adjusted $p < 0.05$). Each GO-BP is represented

in a colour-based gradient for statistical significance [$\sqrt{}$(–$\log_{10}$ adjusted p-value]. GO terms

were clustered hierarchically using semantic similarity (rrvgo; similarity threshold = 0.7).

See Tables S5 and S6 for full terms.

Fig. 2. Orthoflavivirus NS5 interaction network. Quantitative proteomics was used as

described in the text to identify host factors interacting with HA-tagged versions of ZIKV,

USUV or WNV NS5 proteins. Only the top 50 most significantly enriched factors in each

data set are used to build the network and to identify overlaps between data sets. The

protein-protein interaction network was generated using the STRING database (86). Data

clustering was generated in Cystoscape (87), using the MCL algorithm. Those factors

identified in two datasets are coloured in pale yellow, and those in all three datasets in

coral. Each grey-shaded area represents a cluster positively associated with a biological

process (i.e. ribosome biogenesis, cytoplasmic translation, mRNA processing, chromatin

organisation, proton transmembrane transport/aerobic respiration). Mitochondrial proteins

are encircled by a dotted line.

Fig. 3. DsiRNA-mediated gene silencing identifies putative proviral and restriction

host factors for OFV replication. Virus titres in the cellular supernatant of infected cells

($n \geq 4$ biologically independent samples), previously transfected with a DsiRNA targeting

an NS5-interacting factor. At 24 h post-transfection, A549 cells were infected with either

ZIKV (black columns) or USUV (grey columns) at an MOI of 0.1 $TCID_{50}$/cell. Control

infections were carried out in parallel experiments in cells transfected with a non-human

target DsiRNA (NonSP). The bars represent relative changes in the viral titres of gene-

silenced cultures compared to NonSP controls, both at 48 hpi. To improve clarity, the

effect of host factor downregulation on viral infectivity is represented in 3 different graphs:

ribosome biogenesis and translation (A), mitochondrial (B), and other ungrouped host

factors (C). Significant differences observed between unspecific (NonSP) and specific

host-targeting DsiRNA-transfected cells are indicated (Two-way ANOVA, Dunnett's

multiple comparisons test; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$). Bars are representing

the standard error of the mean (SEM) for each data point.

Fig. 4. Reduced virus titres in MYOF-depleted cells. (A – D) Effect of MYOF gene

silencing on viral replication. (A) Relative amount of metabolically active cells after

transfection with a specific (DsiMYOF) or an unspecific DsiRNA (DsiNonSP). Cell viability

is determined at 48 and 72 h post-transfection as suggested in the Methods section.

Values are obtained as a percentage relative to mock untransfected cells. (B) Relative

MYOF mRNA expression was examined by RT-qPCR in A549 cells transfected with either

a MYOF-targeting (DsiMYOF) or a non-human target DsiRNA (DsiNonSP). MYOF mRNA

levels were normalised to GAPDH mRNA levels. Values obtained are the average of at

least 4 independent biological replicas (n ≥ 4), and relative to untransfected cells carried

out in parallel. (C, D) Virus infectivity was examined by $TCID_{50}$ assays of samples

collected at different time points from cells previously transfected with each DsiRNA.

Twenty-four hours after transfection, cells were infected with ZIKV (C) or USUV (D) at an

MOI of 1 $TCID_{50}$/cell. Each value in the graph is the average of ≥ 4 independent biological

replicas. (E – G) Effect of MYOF-targeting drug WJ460 on ZIKV and USUV replication. (E)

The relative amount of metabolically active cells after treatment during 48 h with increasing

concentrations of WJ460 is calculated as indicated in (A). Values are represented as a

percentage relative to mock-treated cells. (F, G) ZIKV (F) or USUV (G) titres detected in

cells treated with WJ460 at 48 h postinfection. In each panel, significant differences are

indicated (Two-way ANOVA, Sidak's multiple comparisons test; *, $p<0.05$; ***, $p < 0.001$).

The standard error of the mean (SEM) is represented for each data point.

Fig. 5. Increased viral titres, RNA and protein levels in PHB2-depleted cells. (A)

Relative PHB2 mRNA expression in A549 cells previously transfected with either a PHB2-

targeting (DsiPHB2) or a non-human target DsiRNA (DsiNonSP). PHB2 expression was

normalised to GAPDH mRNA levels. Values obtained are the average of at least 4

independent biological replicas (n ≥ 4), and relative to untransfected cells carried out in

parallel. (B) PHB2 downregulation leads to increased ZIKV NS5 protein abundance in

infected cells. Representative WB of ZIKV NS5 and PHB2 protein levels in A549 cells

previously transfected with each DsiRNA and infected at 24 h post-transfection with ZIKV.

As a loading reference, an anti-GAPDH antibody was used. (C – F) Virus infectivity was

examined in the supernatants of cells previously transfected with each DsiRNA. Twenty-

four hours after DsiRNA transfection, cells were infected with ZIKV (C, D) or USUV (E, F)

at an MOI of 0.1 (C, E) or 1 $TCID_{50}$/cell (D, F). Each value in the graph is the average of

four independent biological replicas (n = 4). (G – J) Viral RNA genome equivalents

detected in samples collected at 48 hpi from cells previously transfected with each

DsiRNA. Twenty-four hours after DsiRNA transfection, cells were infected at an MOI of 0.1

or 1 $TCID_{50}$/cell with ZIKV (G, H) or USUV (I, J). Each value in the graph represents the

average of viral RNA detected in four independent biological replicas (n = 4) extracted

from infected cells (intracellular, graphs G and I) or the supernatant of these cultures

(extracellular, graphs H and J). The statistical significance in each panel was determined

by Two-way ANOVA and Sidak's multiple comparison test (C – J) or t (A) tests (*, p<0.05;

**, p<0.01; ***, p < 0.001). Bars are representing the standard error of the mean (SEM).

Fig. 6. Increased ZIKV titres and RNA levels in cells treated with a mitophagy

inhibitor. (A) Relative number of metabolically active A549 cells after 48 hours in the

presence of increasing concentrations of mdivi-1. Values are relative to untreated cells (0

μM). (B) ZIKV titres at 48 hpi in the supernatant of cells treated with increasing

concentrations of mdivi-1. Cells were previously inoculated with ZIKV at an MOI of 0.1

$TCID_{50}$/cell. (C) Genomic RNA equivalents in cells infected with ZIKV, and treated with

mdivi-1. D, Relative IFNB1 mRNA expression in ZIKV-infected cells treated with mdivi-1.

Fig. 7. PHB2 is transiently eliminated in ZIKV- and USUV-infected cells. (A, E)

Representative Western blot of PHB2 kinetics in cells infected with either ZIKV (A) or

USUV (E). (B, C, F, G) PHB2 protein levels in mock- (B, F), ZIKV- (C) or USUV-infected

(G) A549 cell lysates at different time points. PHB2 to tubulin band intensity ratios are

relative to those obtained at 48 h in Mock-infected cultures. (D) ZIKV NS5 protein levels

detected in ZIKV-infected A549 cell lysates collected at different time points. NS5 to

tubulin band intensity ratios for each time point are relative to 72 hpi. Each value in the

graphs is the average of three independent biological replicas (n = 3). Bars are

representing the standard error of the mean (SEM) for each data point. Significant differences between mock and infected samples are indicated (One-way ANOVA; *, p < 0.05; **, p < 0.01). (H) Representative blot of PHB2 protein levels in 293T cells collected at different time post-infection in cells infected with ZIKV (top) or USUV (bottom). An anti-GAPDH antibody was used to calibrate total protein loading.

Fig. 8. Increased IFN-I transcription in PHB2-downregulated cells. (A − D), Relative IFNB1 mRNA expression in A549 cells previously transfected with either a PHB2-targeting (DsiPHB2) or a non-human target (DsiNonSP) DsiRNA, and then infected with ZIKV (A, B) or USUV (C, D). IFNB1 expression was normalised to GAPDH mRNA levels. (E − H) Relative IFNB1 mRNA expression in A549 cells transfected with either a PHB2-targeting (DsiPHB2) or a non-human target (DsiNonSP) DsiRNA, and then stimulated with different amounts of poly I:C. IFNB1 expression was normalised to GAPDH mRNA levels. Values obtained are the average of at least 4 independent biological replicas, and relative to control cells (no DsiRNA). (E, G) IFNB1 mRNA levels in poly I:C-stimulated cells compared to control unstimulated cells. (F, H) Relative IFNB1 mRNA expression in PHB2-depleted cells compared to control cells (DsiNonSP) after poly I:C stimulation.

Fig. 9. Ectopic co-expression of NS2B3 and NS5 mediates efficient PHB2 elimination. (A, B) PHB2 is co-immunoprecipitated with ZIKV, USUV and WNV NS5 proteins (A) or their RdRp domains (B). To this, 293T cells were transfected with constructs expressing HA-tagged versions of each NS5 (A), or the RdRp domain alone (B). As controls, there were included mock-transfected cells or constructs expressing EYFP (pEYFP-N1, shown as pEYFP), HA-tagged LacZ (pCruz HA-LacZ, shown as HA-LacZ) or an empty vector (pcDNA3). Cell lysates were incubated with anti-HA agarose beads binding to NS5 and RdRp, and the presence of PHB2 in the co-IPs was examined

with a specific antibody. (C) ZIKV NS5 overexpression leads to reduced STAT2 but not PHB2 levels. 293T cells were transfected with increasing amounts (400, 800, 1600, 3200 ng) of either a construct expressing HA-tagged ZIKV NS5 or an empty vector (pcDNA3). (D) Ectopic expression of NS2B/NS3 precursor (NS2B3) leads to reduced PHB2 levels in 293T cells. To this, 5×10^5 293T cells were seeded on a 35-mm plate and transfected with either 2 or 4 µg of pZIKV-NS2B3 or an empty vector (pcDNA3). (E) PHB2 elimination by NS2B3 is enhanced in the presence of NS5. In this assay, 5×10^5 293T cells were seeded on a 35-mm plate and co-transfected with 2 µg of pZIKV-NS2B3 and 2 µg of pZIKV-HA-NS5 or an empty vector (pcDNA3). (F) Larger amounts of NS5 lead to reduced PHB2 levels in cells expressing a suboptimal amount of NS2B/NS3. Transfections were carried out upon 5×10^6 293T cells on 100-mm plates with 2 µg of pZIKV-NS2B3 and different amounts of pZIKV-HA-NS5 (gradually increasing from 0.5 to 6 µg).

10. REFERENCES

1. Lim A, Shearer FM, Sewalk K, Pigott DM, Clarke J, Ghouse A, Judge C, Kang H, Messina JP, Kraemer MUG, Gaythorpe KAM, de Souza WM, Nsoesie EO, Celone M, Faria N, Ryan SJ, Rabe IB, Rojas DP, Hay SI, Brownstein JS, Golding N, Brady OJ. 2025. The overlapping global distribution of dengue, chikungunya, Zika and yellow fever. Nat Commun 16:3418.

2. Xu Y, Zhou J, Liu T, Liu P, Wu Y, Lai Z, Gu J, Chen X. 2022. Assessing the risk of spread of Zika virus under current and future climate scenarios. Biosaf Health 4:193–204.

3. Malavige GN, Sjö P, Singh K, Piedagnel J-M, Mowbray C, Estani S, Lim SCL, Siquierra AM, Ogg GS, Fraisse L, Ribeiro I. 2023. Facing the escalating burden of dengue: Challenges and perspectives. PLOS Global Public Health 3:e0002598.

4. Pierson TC, Diamond MS. 2020. The continued threat of emerging flaviviruses. Nat Microbiol 5:796–812.

5. de Souza WM, Weaver SC. 2024. Effects of climate change and human activities on vector-borne
diseases. Nat Rev Microbiol 22:476–491.

6. Roiz D, Pontifes PA, Jourdain F, Diagne C, Leroy B, Vaissière A-C, Tolsá-García MJ, Salles J-M, Simard
F, Courchamp F. 2024. The rising global economic costs of invasive Aedes mosquitoes and Aedes-
borne diseases. Sci Total Environ 933:173054.

7. Moore SM, Oidtman RJ, James Soda K, Siraj AS, Reiner RC, Barker CM, Alex Perkins T. 2020.
Leveraging multiple data types to estimate the size of the Zika epidemic in the Americas. PLoS Negl
Trop Dis 14:e0008640.

8. Musso D, Ko AI, Baud D. 2019. Zika Virus Infection — After the Pandemic. New England Journal of
Medicine 381:1444–1457.

9. Rabe IB, Hills SL, Haussig JM, Walker AT, dos Santos T, San Martin JL, Gutierrez G, Mendez-Rico J,
Rodriguez JC, Elizondo-Lopez D, Gonzalez-Escobar G, Chanda E, Al Eryani SM, Kodama C, Yajima A,
Kakkar M, Kato M, Wijesinghe PR, Samaraweera S, Brindle H, Tissera H, Kelley J, Lackritz E, Rojas DP.
2025. A Review of the Recent Epidemiology of Zika Virus Infection. Am J Trop Med Hyg
https://doi.org/10.4269/ajtmh.24-0420.

10. Miner JJ, Diamond MS. 2017. Zika Virus Pathogenesis and Tissue Tropism. Cell Host Microbe 21:134–
142.

11. Ferraris P, Yssel H, Missé D. 2019. Zika virus infection: an update. Microbes Infect 21:353–360.

12. Santana LMB, de Moura IA, Mouzinho Ramos Tanaka Y, França RF de O. 2025. Zika Virus: A Review
of Biology, Clinical Impacts, and Coinfections. Viruses 17:637.

13. Scaramozzino P, Carvelli A, Bruni G, Cappiello G, Censi F, Magliano A, Manna G, Ricci I, Rombolà P,
Romiti F, Rosone F, Sala MG, Scicluna MT, Vaglio S, De Liberato C. 2021. West Nile and Usutu viruses
co-circulation in central Italy: outcomes of the 2018 integrated surveillance. Parasit Vectors 14:243.

14. Brüssow H, Figuerola J. 2025. The Spread of the Mosquito-Transmitted West Nile Virus in North
America and Europe. Microb Biotechnol 18.

15. Simonin Y. 2022. Un second cas autochtone d'infection par le virus Usutu détecté en France : que
faut-il savoir ? The Conversation. https://theconversation.com/un-second-cas-autochtone-

dinfection-par-le-virus-usutu-detecte-en-france-que-faut-il-savoir-97944. Retrieved 16 January
2023.

16.   Clé M, Beck C, Salinas S, Lecollinet S, Gutierrez S, Van de Perre P, Baldet T, Foulongne V, Simonin Y.
2019. Usutu virus: A new threat? Epidemiol Infect 147:e232.

17.   Simonin Y. 2024. Circulation of West Nile Virus and Usutu Virus in Europe: Overview and Challenges.
Viruses 16:599.

18.   Cadar D, Maier P, Müller S, Kress J, Chudy M, Bialonski A, Schlaphof A, Jansen S, Jöst H, Tannich E,
Runkel S, Hitzler WE, Hutschenreuter G, Wessiepe M, Schmidt-Chanasit J. 2017. Blood donor
screening for West Nile virus (WNV) revealed acute Usutu virus (USUV) infection, Germany,
September 2016. Eurosurveillance 22:30501.

19.   Bakonyi T, Jungbauer C, Aberle SW, Kolodziejek J, Dimmel K, Stiasny K, Allerberger F, Nowotny N.
2017. Usutu virus infections among blood donors, Austria, July and August 2017 – Raising awareness
for diagnostic challenges. Eurosurveillance 22.

20.   Madere FS, Andrade da Silva AV, Okeze E, Tilley E, Grinev A, Konduru K, García M, Rios M. 2025.
Flavivirus infections and diagnostic challenges for dengue, West Nile and Zika Viruses. npj Viruses
3:36.

21.   Wahaab A, Mustafa BE, Hameed M, Stevenson NJ, Anwar MN, Liu K, Wei J, Qiu Y, Ma Z. 2021.
Potential Role of Flavivirus NS2B-NS3 Proteases in Viral Pathogenesis and Anti-flavivirus Drug
Discovery Employing Animal Cells and Models: A Review. Viruses 14:44.

22.   Ferrero DS, Ruiz-Arroyo VM, Soler N, Usón I, Guarné A, Verdaguer N. 2019. Supramolecular
arrangement of the full-length Zika virus NS5. PLoS Pathog 15:e1007656.

23.   Denolly S, Stukalov A, Barayeu U, Rosinski AN, Kritsiligkou P, Joecks S, Dick TP, Pichlmair A,
Bartenschlager R. 2023. Zika virus remodelled ER membranes contain proviral factors involved in
redox and methylation pathways. Nat Commun 14:8045.

24.   Welsch S, Miller S, Romero-Brey I, Merz A, Bleck CKE, Walther P, Fuller SD, Antony C, Krijnse-Locker
J, Bartenschlager R. 2009. Composition and three-dimensional architecture of the dengue virus
replication and assembly sites. Cell Host Microbe 5:365–75.

25. Ci Y, Shi L. 2021. Compartmentalized replication organelle of flavivirus at the ER and the factors
involved. Cellular and Molecular Life Sciences 2021 78:11 78:4939–4954.

26. Scaturro P, Stukalov A, Haas DA, Cortese M, Draganova K, Płaszczyca A, Bartenschlager R, Götz M,
Pichlmair A. 2018. An orthogonal proteomic survey uncovers novel Zika virus host factors. Nature
561:253–257.

27. Schneider WM, Hoffmann HH. 2022. Flavivirus-host interactions: an expanding network of proviral
and antiviral factors. Curr Opin Virol 52:71–77.

28. Li P, Wu J, Liu S, Lu R, Jiang H, Wang N, Luo M, Guo L, Xiao J, Bu L, Liu L, Xing F, Peng H, Li C, Ma L,
Zhao B, Zhou Z, Guo D. 2022. The RNA polymerase of cytoplasmically replicating Zika virus binds
with chromatin DNA in nuclei and regulates host gene transcription. Proc Natl Acad Sci U S A
119:e2205013119.

29. Lebeau G, Hoareau M, Rivière S, El Safadi D, Da Silva CR, Krejbich-Trotot P, Viranaicken W. 2024. Cell
cycle and mitosis progression during ZIKA virus infection: The viral non-structural protein NS5 as a
master regulator of the APC/cyclosome? Biochimie 221:75–80.

30. Best SM. 2017. The Many Faces of the Flavivirus NS5 Protein in Antagonism of Type I Interferon
Signaling. J Virol 91.

31. Albentosa-González L, Jimenez de Oya N, Arias A, Clemente-Casares P, Martin-Acebes MÁ, Saiz JC,
Sabariegos R, Mas A. 2021. Akt Kinase Intervenes in Flavivirus Replication by Interacting with Viral
Protein NS5. Viruses 13:896.

32. Rodrigo I, Albentosa-González L, Romero de Ávila MJ, Bassi MR, Sempere RN, Clemente-Casares P,
Arias A. 2025. Ubiquitin-like modifier-activating enzyme 1 interacts with Zika virus NS5 and
promotes viral replication in the infected cell. J Gen Virol 106:002063.

33. Saade M, Ferrero DS, Blanco-Ameijeiras J, Gonzalez-Gobartt E, Flores-Mendez M, Ruiz-Arroyo VM,
Martínez-Sáez E, Ramón y Cajal S, Akizu N, Verdaguer N, Martí E. 2020. Multimerization of Zika
Virus-NS5 Causes Ciliopathy and Forces Premature Neurogenesis. Cell Stem Cell 27:920-936.e8.

34. Deb A, Nagpal S, Yadav RK, Thakur H, Nair D, Krishnan V, Vrati S. 2024. Japanese encephalitis virus
NS5 protein interacts with nucleolin to enhance the virus replication. J Virol 98:e0085824.

35. Grant A, Ponia SS, Tripathi S, Balasubramaniam V, Miorin L, Sourisseau M, Schwarz MC, Sánchez-
Seco MP, Evans MJ, Best SM, García-Sastre A. 2016. Zika Virus Targets Human STAT2 to Inhibit Type i
Interferon Signaling. Cell Host Microbe 19:882–890.

36. Gonzalez Lopez Ledesma MM, Costa Navarro G, Pallares HM, Paletta A, De Maio F, Iglesias NG,
Gebhard L, Oviedo Rouco S, Ojeda DS, de Borba L, Giraldo M, Rajsbaum R, Ceballos A, Krogan NJ,
Shah PS, Gamarnik A V. 2023. Dengue virus NS5 degrades ERC1 during infection to antagonize NF-kB
activation. Proceedings of the National Academy of Sciences 120.

37. Ren W, Fu C, Zhang Y, Ju X, Jiang X, Song J, Gong M, Li Z, Fan W, Yao J, Ding Q. 2024. Zika virus NS5
protein inhibits type I interferon signaling via CRL3 E3 ubiquitin ligase-mediated degradation of
STAT2. Proc Natl Acad Sci U S A 121:e2403235121.

38. Ponia SS, Robertson SJ, McNally KL, Subramanian G, Sturdevant GL, Lewis M, Jessop F, Kendall C,
Gallegos D, Hay A, Schwartz C, Rosenke R, Saturday G, Bosio CM, Martens C, Best SM. 2021.
Mitophagy antagonism by ZIKV reveals Ajuba as a regulator of PINK1 signaling, PKR-dependent
inflammation, and viral invasion of tissues. Cell Rep 37:109888.

39. Liu Y, Zhou T, Hu J, Jin S, Wu J, Guan X, Wu Y, Cui J. 2022. Targeting Selective Autophagy as a
Therapeutic Strategy for Viral Infectious Diseases. Front Microbiol 13.

40. Yan C, Gong L, Chen L, Xu M, Abou-Hamdan H, Tang M, Désaubry L, Song Z. 2020. PHB2 (prohibitin
2) promotes PINK1-PRKN/Parkin-dependent mitophagy by the PARL-PGAM5-PINK1 axis. Autophagy
16:419–434.

41. Jin SM, Youle RJ. 2012. PINK1- and Parkin-mediated mitophagy at a glance. J Cell Sci 125:795–799.

42. Cheng M, Lin N, Dong D, Ma J, Su J, Sun L. 2021. PGAM5: A crucial role in mitochondrial dynamics
and programmed cell death. Eur J Cell Biol 100:151144.

43. Xiao Y, Zhou Y, Lu Y, Zhou K, Cai W. 2018. PHB2 interacts with LC3 and SQSTM1 is required for bile
acids-induced mitophagy in cholestatic liver. Cell Death Dis 9:160.

44. Yoshinaka T, Kosako H, Yoshizumi T, Furukawa R, Hirano Y, Kuge O, Tamada T, Koshiba T. 2019.
Structural Basis of Mitochondrial Scaffolds by Prohibitin Complexes: Insight into a Role of the Coiled-
Coil Region. iScience 19:1065.

45.   Sun L, Liu L, Yang X-J, Wu Z. 2004. Akt binds prohibitin 2 and relieves its repression of MyoD and
812   muscle differentiation. J Cell Sci 117:3021–3029.

814   46.   Ding T, Xu H, Zhang X, Yang F, Zhang J, Shi Y, Bai Y, Yang J, Chen C, Zhu C, Zhang H. 2024. Prohibitin 2
815   orchestrates long noncoding RNA and gene transcription to accelerate tumorigenesis. Nat Commun
816   15:8385.

818   47.   Kasashima K, Ohta E, Kagawa Y, Endo H. 2006. Mitochondrial Functions and Estrogen Receptor-
819   dependent Nuclear Translocation of Pleiotropic Human Prohibitin 2. Journal of Biological Chemistry
820   281:36401–36410.

822   48.   Kim J, Akiyama M, Park J, Lin M, Shimo A, Ueki T, Daigo Y, Tsunoda T, Nishidate T, Nakamura Y,
823   Katagiri T. 2009. Activation of an estrogen/estrogen receptor signaling by BIG3 through its inhibitory
824   effect on nuclear transport of PHB2/REA in breast cancer. Cancer Sci 100:1468–1478.

826   49.   Bassi MR, Sempere RN, Meyn P, Polacek C, Arias A. 2018. Extinction of Zika virus and Usutu virus by
827   lethal mutagenesis reveals different patterns of sensitivity to three mutagenic drugs. Antimicrob
828   Agents Chemother 62:e00380-18.

830   50.   Sempere RN, Arias A. 2019. Establishment of a Cell Culture Model of Persistent Flaviviral Infection:
831   Usutu Virus Shows Sustained Replication during Passages and Resistance to Extinction by Antiviral
832   Nucleosides. Viruses 11:560.

834   51.   Rodrigo I, Ballesta C, Nunes EB, Pérez P, García-Arriaza J, Arias A. 2022. Eeyarestatin I, an inhibitor of
835   the valosin-containing protein, exhibits potent virucidal activity against the flaviviruses. Antiviral Res
836   207:105416.

838   52.   Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative
839   PCR and the 2-ΔΔCT method. Methods 25:402–408.

841   53.   Albentosa-González L, Sabariegos R, Arias A, Clemente-Casares P, Mas A. 2021. Akt interacts with
842   usutu virus polymerase, and its activity modulates viral replication. Pathogens 10:1–15.

844   54.   Hughes CS, Moggridge S, Müller T, Sorensen PH, Morin GB, Krijgsveld J. 2019. Single-pot, solid-
845   phase-enhanced sample preparation for proteomics experiments. Nat Protoc 14:68–85.

847   55.   Meyer B, Chiaravalli J, Gellenoncourt S, Brownridge P, Bryne DP, Daly LA, Grauslys A, Walter M,
848   Agou F, Chakrabarti LA, Craik CS, Eyers CE, Eyers PA, Gambin Y, Jones AR, Sierecki E, Verdin E,

Vignuzzi M, Emmott E. 2021. Characterising proteolysis during SARS-CoV-2 infection identifies viral
cleavage sites and cellular targets with therapeutic potential. Nature Communications 2021 12:1
12:1–16.

56.   Cox J, Mann M. 2008. MaxQuant enables high peptide identification rates, individualized p.p.b.-
range mass accuracies and proteome-wide protein quantification. Nat Biotechnol 26:1367–1372.

57.   Nguyen SK, Anders N, Holmes S, Barrow HG, Lukhovitskaya N, Jahun AS, Georgana I, Caller LG, Edgar
JR, Graham SC, Emmott E, Firth AE, Stewart H. 2025. Encephalomyocarditis virus protein 2B*
interacts with 14-3-3 proteins through a phosphorylated C-terminal binding motif. mBio 16.

58.   Zhang T, Li J, He Y, Yang F, Hao Y, Jin W, Wu J, Sun Z, Li Y, Chen Y, Yi Z, Liu M. 2018. A small molecule
targeting myoferlin exerts promising anti-tumor effects on breast cancer. Nat Commun 9:3726.

59.   Bordt EA, Clerc P, Roelofs BA, Saladino AJ, Tretter L, Adam-Vizi V, Cherok E, Khalil A, Yadava N, Ge
SX, Francis TC, Kennedy NW, Picton LK, Kumar T, Uppuluri S, Miller AM, Itoh K, Karbowski M, Sesaki
H, Hill RB, Polster BM. 2017. The Putative Drp1 Inhibitor mdivi-1 Is a Reversible Mitochondrial
Complex I Inhibitor that Modulates Reactive Oxygen Species. Dev Cell 40:583-594.e6.

60.   Zhou P, Zhang Q, Yang Y, Wu W, Chen D, Zheng Z, Jongkaewwattana A, Jin H, Zhou H, Luo R. 2024.
Cleavage of SQSTM1/p62 by the Zika virus protease NS2B3 prevents autophagic degradation of viral
NS3 and NS5 proteins. Autophagy 20:2769–2784.

61.   Ding Q, Gaska JM, Douam F, Wei L, Kim D, Balev M, Heller B, Ploss A. 2018. Species-specific
disruption of STING-dependent antiviral cellular defenses by the Zika virus NS2B3 protease.
Proceedings of the National Academy of Sciences 115.

62.   Lennemann NJ, Coyne CB. 2017. Dengue and Zika viruses subvert reticulophagy by NS2B3-mediated
cleavage of FAM134B. Autophagy 13:322–332.

63.   Boytz R, Keita K, Pawlak JB, Laurent-Rolle M. 2024. Flaviviruses manipulate mitochondrial processes
to evade the innate immune response. npj Viruses 2:47.

64.   Agarwal A, Alam MF, Basu B, Pattanayak S, Asthana S, Syed GH, Kalia M, Vrati S. 2022. Japanese
Encephalitis Virus NS4A Protein Interacts with PTEN-Induced Kinase 1 (PINK1) and Promotes
Mitophagy in Infected Cells. Microbiol Spectr 10:e0083022.

65. Singh B, Avula K, Sufi SA, Parwin N, Das S, Alam MF, Samantaray S, Bankapalli L, Rani A, Poornima K,
Prusty B, Mallick TP, Shaw SK, Dodia H, Kabi S, Pagad TT, Mohanty S, Syed GH. 2022. Defective
Mitochondrial Quality Control during Dengue Infection Contributes to Disease Pathogenesis. J Virol
96:e0082822.

66. Lee JK, Shin OS. 2023. Zika virus modulates mitochondrial dynamics, mitophagy, and mitochondria-
derived vesicles to facilitate viral replication in trophoblast cells. Front Immunol 14.

67. Qi Y, Rajbanshi B, Hao R, Dang Y, Xu C, Lu W, Dai L, Zhang B, Zhang X. 2025. The dual role of PGAM5
in inflammation. Exp Mol Med 57:298–311.

68. Oyang L, Li J, Jiang X, Lin J, Xia L, Yang L, Tan S, Wu N, Han Y, Yang Y, Luo X, Li J, Liao Q, Shi Y, Zhou Y.
2022. The function of prohibitins in mitochondria and the clinical potentials. Cancer Cell Int 22:343.

69. Eyre NS, Kirby EN, Anfiteatro DR, Bracho G, Russo AG, White PA, Aloia AL, Beard MR. 2020.
Identification of Estrogen Receptor Modulators as Inhibitors of Flavivirus Infection. Antimicrob
Agents Chemother 64:e00289-20.

70. Grady SF, Pinto AK, Hassert M, D'Angelo JA, Brien JD, Arnatt CK. 2021. Selective estrogen receptor
modulator, tamoxifen, inhibits Zika virus infection. J Med Virol 93:6155–6162.

71. Celegato M, Sturlese M, Vasconcelos Costa V, Trevisan M, Lallo Dias AS, Souza Passos IB, Queiroz-
Junior CM, Messa L, Favaro A, Moro S, Teixeira MM, Loregian A, Mercorelli B. 2023. Small-Molecule
Inhibitor of Flaviviral NS3-NS5 Interaction with Broad-Spectrum Activity and Efficacy *In Vivo*. mBio
14.

72. Tay MYF, Saw WG, Zhao Y, Chan KWK, Singh D, Chong Y, Forwood JK, Ooi EE, Grüber G, Lescar J, Luo
D, Vasudevan SG. 2015. The C-terminal 50 Amino Acid Residues of Dengue NS3 Protein Are
Important for NS3-NS5 Interaction and Viral Replication. Journal of Biological Chemistry 290:2379–
2394.

73. Wang L, Zhang X, Huang X, Sha X, Li X, Zheng J, Li S, Wei Z, Wu F. 2024. Homoplantaginin alleviates
high glucose-induced vascular endothelial senescence by inhibiting mtDNA–cGAS–STING pathway
via blunting DRP1–mitochondrial fission–VDAC1 axis. The FASEB Journal 38.

74. Jiménez-Loygorri JI, Villarejo-Zori B, Viedma-Poyatos Á, Zapata-Muñoz J, Benítez-Fernández R,
Frutos-Lisón MD, Tomás-Barberán FA, Espín JC, Area-Gómez E, Gomez-Duran A, Boya P. 2024.
Mitophagy curtails cytosolic mtDNA-dependent activation of cGAS/STING inflammation during
aging. Nat Commun 15:830.

75.   Gkikas I, Palikaras K, Tavernarakis N. 2018. The Role of Mitophagy in Innate Immunity. Front
927   Immunol 9:1283.

929   76.   Yuk J-M, Silwal P, Jo E-K. 2020. Inflammasome and Mitophagy Connection in Health and Disease. Int
930   J Mol Sci 21:4714.

932   77.   Jin Y, Zhao L, Wang S, Zhang X, Quan J, Lin Z, Piao J. 2023. *RSL1D1* knockdown induces ferroptosis
933   and mediates ferrous iron accumulation in senescent cells by inhibiting *FTH1* mRNA stability.
934   Carcinogenesis 44:129–142.

936   78.   Liu X, Chen J, Long X, Lan J, Liu X, Zhou M, Zhang S, Zhou J. 2022. RSL1D1 promotes the progression
937   of colorectal cancer through RAN-mediated autophagy suppression. Cell Death Dis 13:43.

939   79.   Gupta S, Yano J, Mercier V, Htwe HH, Shin HR, Rademaker G, Cakir Z, Ituarte T, Wen KW, Kim GE,
940   Zoncu R, Roux A, Dawson DW, Perera RM. 2021. Lysosomal retargeting of Myoferlin mitigates
941   membrane stress to enable pancreatic cancer growth. Nat Cell Biol 23:232–242.

943   80.   Bernatchez PN, Sharma A, Kodaman P, Sessa WC. 2009. Myoferlin is critical for endocytosis in
944   endothelial cells. American Journal of Physiology-Cell Physiology 297:C484–C492.

946   81.   Morita E, Suzuki Y. 2021. Membrane-Associated Flavivirus Replication Complex—Its Organization
947   and Regulation. Viruses 13:1060.

949   82.   Cretu C, Chernev A, Kibédi Szabó CZ, Pena V, Urlaub H, Moser T, Preobraschenski J. 2025. Structural
950   insights into lipid membrane binding by human ferlins. EMBO J https://doi.org/10.1038/s44318-025-
951   00463-8.

953   83.   He Y, Kan W, Li Y, Hao Y, Huang A, Gu H, Wang M, Wang Q, Chen J, Sun Z, Liu M, Chen Y, Yi Z. 2021. A
954   potent and selective small molecule inhibitor of myoferlin attenuates colorectal cancer progression.
955   Clin Transl Med 11.

957   84.   Arutyunov A, Durán-Laforet V, Ai S, Ferrari L, Murphy R, Schafer DP, Klein RS. 2024. West Nile Virus-
958   Induced Expression of Senescent Gene Lgals3bp Regulates Microglial Phenotype within Cerebral
959   Cortex. Biomolecules 14:808.

961   85.   Hulsen T, de Vlieg J, Alkema W. 2008. BioVenn – a web application for the comparison and
962   visualization of biological lists using area-proportional Venn diagrams. BMC Genomics 9:488.

86. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, Gable AL, Fang T, Doncheva NT,
Pyysalo S, Bork P, Jensen LJ, von Mering C. 2023. The STRING database in 2023: protein–protein
association networks and functional enrichment analyses for any sequenced genome of interest.
Nucleic Acids Res 51:D638–D646.

87. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T.
2003. Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction
Networks. Genome Res 13:2498–2504.

A Ribosome and translation

■ ZIKV
■ USUV

Change in virus titre (Log$_{10}$ TCID$_{50}$/ml)

DsiRNA target gene

NonSP, RPL21, RPL24, RPL34, RSL1D1

B Mitochondrial proteins

■ ZIKV
■ USUV

Change in virus titre (Log$_{10}$ TCID$_{50}$/ml)

DsiRNA target gene

NonSP, ATP5F1B, FH, PGAM5, PHB2, SFXN1

C Other proteins

■ ZIKV
■ USUV

Change in virus titre (Log$_{10}$ TCID$_{50}$/ml)

DsiRNA target gene

NonSP, DDX21, HSD17B12, LGALS3BP, MYOF, SMARCA5

A Cell viability (RNAi)

B Silencing efficacy

C ZIKV - DsiRNA

D USUV - DsiRNA

E Cell viability - WJ460

F ZIKV - WJ460

G USUV - WJ460

A Cell viability

B ZIKV titres

C ZIKV gRNA

D IFN (ZIKV infection)