Evaluating the roles of testosterone and sex-linked genes in territorial aggression of sex-reversed XY females in the African pygmy mouse

Louise D. Heitzmann[1], Julie Perez[1], Romann Charbonnier[1], Marc Fichter[1], Xavier Hautecoeur[2], Theo Deremarque[1], Agnes O. Martin[3] & Frederic Veyrunes[1]

[1] Institut des Sciences de l'Evolution de Montpellier, University of Montpellier, CNRS, IRD, UMR 5554, Montpellier, France

[2] Institut de Génétique Humaine, University of Montpellier, CNRS, UMR 9002, Montpellier, France

[3] Institut de Génétique Fonctionnelle, University of Montpellier, CNRS, INSERM, UMR 5203, Montpellier, France

Sex differences in social aggression are widespread across the animal kingdom, with males typically displaying greater territoriality. While this dimorphism has traditionally been attributed to sex hormones, sex chromosomes can also contribute to it independently of hormonal influence. In the African pygmy mouse *Mus minutoides*, naturally occurring sex-reversed XY females (named X*Y due to a mutation on the X chromosome) are highly territorial in comparison to the other female genotypes present in the population (XX and XX*). However, the molecular basis of this phenotype remains unknown. Here, we evaluate molecular factors – known to correlate with aggressiveness – following a standardized behavioural assessment of aggression. We focus on i) the androgen pathway by quantifying testosterone serum levels and expression of its receptor in the brain; ii) the brain dopaminergic system through expression of dopamine-regulating genes, including the sex-determining *Sry* gene; and iii) neuroendocrine circuits, via vasopressin and oxytocin expression. These systems, although distinct, converge on hypothalamic-limbic circuits that regulate social behaviour and threat responses, providing an integrated framework for the modulation of aggression. Contrary to earlier reports, we found reduced level of aggression in X*Y females, likely to reflect a breeding facility effect. We also observe no correlations between aggressiveness and androgen levels or gene expression of the tested factors. However, our results support a stimulation of the dopaminergic system and of the oxytocin pathway following the agonistic assay suggesting their potential involvement in aggression-related responses. This further supports the idea that aggression is multifactorial. It is shaped by the interaction of several neuroendocrine and neurotransmitter pathways rather than a single determinant.

Introduction

Aggression is a complex social behaviour widespread across the animal kingdom. In many species, social aggression is highly sexually dimorphic, prevalent in males competing for territories and mates, whereas female aggression is often associated with maternal defense (Darwin, 1871; Pandolfi et al. 2021). This sexual dimorphism is mostly known to arise from sex differences in steroid hormones levels, especially testosterone which can promote aggression either directly or through its metabolites (e.g. aromatization of testosterone to estradiol) (Rose et al. 1971; Book et al. 2001; Han & De Vries, 2003; Giammanco et al. 2005; Phoenix, 2009; Phoenix et al. 1959; Wu et al. 2009; Zha et al. 2025).

Sex chromosomes and imbalance in sex-linked genes can also contribute to sex differences in social aggression, factors that have received relatively little attention (Gatewood et al. 2006; Lovell-Badge, 2005; Saunders et al. 2016; Sluyter et al. 1996). For instance, levels of dopamine – a well-characterized neurotransmitter associated with both social and maternal aggression – can be directly influenced by the expression of sex-linked genes (Czech et al. 2012; Godar et al. 2016; Korzan et al. 2006; Kudryavtseva et al. 2017; Mahadevia et al. 2021; van Erp & Miczek, 2000; Yamaguchi & Lin, 2018). The sex-determining Y-linked gene *Sry* is expressed in specific brain regions and upregulates the rate-limiting enzyme required for dopamine production: tyrosine hydroxylase (*Th*). Elevated level of *Th,* and associated dopamine levels, is directly correlated to increased agonistic behaviours (Mahadevia et al. 2021). Conversely, the gene encoding the enzyme responsible for dopamine degradation, monoamine oxydase A enzyme (*MaoA;* Shih et al. 1999), is located on the X chromosome. Consistent with the relationship between dopamine levels and aggression, knock-out or reduced expression of *MaoA* have been related to enhanced aggressiveness in mice (Cases et al. 1995 ; Scott et al. 2008; Kudryavtseva et al. 2017) and humans (Godar et al. 2016;). There is also evidence that the neuropeptide vasopressin (AVP), which can promote or inhibit aggression (depending on species), is influenced by X chromosome number (Gatewood et al. 2006; Bester-Meredith & Marler, 1999; de Moura Oliveira et al. 2021; Everts et al. 1997; Cox et al. 2015). These examples highlight that sex-linked genes (and the autosomal genes they regulate) can shape the neural circuits underpinning aggressive behaviours, but their role is often difficult to separate from that of steroid hormones as chromosomal sex (XX or XY) usually aligns with gonadal sex (ovaries or testes). Nonetheless, there are a few mammalian exceptions in which these two factors are dissociated (Saunders & Veyrunes, 2021).

The African pygmy mouse is a mammal species showing a natural dissociation between gonadal and chromosomal sexes, with the presence of sex-reversed females. About 1 million years ago, there was a mutation on the X chromosome resulting in a third sex chromosome, called X*, and induced sex reversal in X*Y individuals (Veyrunes et al. 2010a). Three female genotypes are found in natural populations: XX, XX* and X*Y, while all males are XY. In addition, the three sex chromosomes are fused to autosomes, so called neo-sex chromosomes, which are subject to sex-specific recombination and inheritance patterns as well (Veyrunes et al. 2010b; Veyrunes et al. 2006; Baudat et al. 2019; Saunders et al. unpublished data). Although X*Y individuals are fully fertile females (Rahmoun et al. 2014; Saunders et al. 2014), they have different behavioural phenotypes compared to XX and XX* females, including an enhanced territoriality assessed by aggressive behaviours in a resident-intruder assay (Saunders et al. 2016). Yet, whether sex chromosomes influence directly (by regulating the expression of sex-linked and/or autosomal genes; e.g., dopamine pathway) or indirectly (through their control on hormone levels and notably testosterone) social aggression is unknown.

Here, we reproduce and conduct the agonistic assay from Saunders et al. (2016) to investigate candidates for the molecular basis of territorial aggression in X*Y females: testosterone and sex chromosomes/sex-linked genes. We evaluate agonistic behaviours from which we define a probability of offensiveness as well as an aggressiveness level (i.e., intensity of aggression) in individuals of the three female genotypes. We then correlated the aggressiveness level with markers of the androgen (hormonal output and receptor expression), dopaminergic (expression of sex-linked modulators) and neuropeptide (expression of *Avp*) pathways. Oxytocin was included in the neuropeptide pathway due to its established role in modulating territorial aggression as well (Tan et al. 2019; Goodson et al. 2015; Love, 2014). Our results suggest that dopamine and oxytocin pathways are stimulated following resident-intruder assays in X*Y females, but do not necessarily trigger aggressive behaviours.

Material & Methods

Animals

Individuals were reared in a room held at 23°C with a 15:9 light-dark cycle in our breeding facility at the University of Montpellier (CECEMA). Virgin females were kept in groups of 5-6 individuals in environmentally enriched cages (38 x 26 x 24 cm) filled with woodchips' bedding, with constant food and water supply. Males were grouped by two (brothers) or isolated in small cages (32 x 45 cm) to prevent agonistic behaviour. We genotyped females by PCR of *Sry* (Veyrunes et al. 2010a)

and using X/X* markers with the following primers: SC328.2.R 5'-
AAGCCTTGACCCAGCCTTCT-3' and SC328.2L1 5'-TGAAGGACAGGACAGGAACACT-3'.
The primers amplified one band for XX females 553 bp, two bands for XX* females (553 and 595
bp) and one band for X*Y females (595 bp). Animal experimentations were performed with the
approval of the French Ethical Committee on animal care and use (No. APAFIS#23541), and
following European guidelines.
108
Resident-intruder assay
Tests were conducted following the same procedures as in Saunders et al. (2016). A virgin female
aged between 7-10 months (table 1) is isolated in a large plexiglass cage (40 L x 30 W x 30 H cm)
with its own bedding and enrichment for 48 hours, in order to establish the resident status. We used
females in the estrus stage (determined by vaginal smears with the presence of anucleate squamous-
type epithelial cell; see Veryunes et al. 2024 for details) to make sure they would not be sexually
receptive by the time of experimentations two days later, which is likely to influence aggression
(Hyde & Sawyer, 1977). Prior the assay, male is placed in a side cage (10 x 10 x 10 cm) with a trap
door leading to the experimental environment. Males are tested once or twice at most, with a
minimal gap of seven days between trials. An assay started once the male entered the female
environment and female-male interactions were then recorded for 10 minutes. Each experiment was
recorded on camera (using phone camera) and videos were analysed manually by one rater blind to
females' genotype. We measured the latency of the first offensive attack, the frequency of attacks
(i.e., number of bite attempts corrected by the overall number of female-male interactions to
account for differences in female-male encounters between assays), the type of attacks (defensive or
offensive) as well as the percent of time females spend chasing the male during an experiment. At
the end of each assay, females were anaesthetized and then decapitated. Blood and brain were
instantaneously collected for hormonal dosage and candidate gene RNA quantifications. Overall,
we tested 19 X*Y females, 11 XX and 8 XX* females. XX and XX* females were considered as
one control group with low aggression level (Saunders et al. 2016).
129
Testosterone dosage assay
The whole blood collected after female euthanasia was left to clot for approximately 45-60 minutes.
Samples were then centrifuged for 10 minutes at 4500 rpm and serum were collected into new tubes
and stored at -20°C until processing. Testosterone levels were quantified by immuno-enzymatic
assay using IBL International ELISA kit (#RE52151) and following manufacturer's instructions.
Females' sera were not diluted except for two females which did not have enough material (i.e., 25

µL). These samples were treated with caution as dilution can impede detection and measures due to
already low hormonal levels (Veyrunes et al. 2024). Most samples were measured only once due to
limited amount of blood collected but some samples with large serum volume were duplicated and
values of testosterone were then averaged across replicates. Measurement for the 38 females were
spread over three ELISA assays. One female (X*Y) did not pass detection threshold and the females
in the last assay (2 XX, 3 XX* and 3 X*Y) showed unusually low concentrations, indicative of
unstable testosterone in serum, and likely to reflect a storage issue with prolonged thawing. We thus
applied a correction factor for these values specifically: when possible (enough serum left), we
repeated the ELISA test on females from the 1st and 2nd ELISA assays. We then calculated the
difference between the first measure (used in the study) and the new ones (i.e., following thawing
issue) and applied the averaged difference as a correction factor to be multiplied by the values from
the 3rd assay.

RNA extraction , reverse transcription and RT-qPCR of Th, MaoA, Drd2, Sry, Avp, Oxt and Ar
Brains collected were immediately placed in RNAlater®, maintained at 4°C for 48 hours and then
stored at -80 °C until further use. Whole brain RNA was extracted using Rneasy® Plus Mini Kit
(Qiagen). Tissues were first grounded in liquid nitrogen using a mortar and a pestle under sterile
environment. We added a lysis buffer to homogenize materials and because we used the whole
tissue, we adjusted volumes to four times volumes recommendation (4 x 6 µL of β-mercapto and 4
x 600 µL of RLT buffer). This gave us four tubes per sample. RNA was purified for two tubes
following manufacturer's protocol and the remaining two were stored at -80°C. Extracted RNA was
eluted in 50 µL of RNase free water and stored at -80°C until further use. Before real-time PCR
quantifications (RT-qPCR) assays, samples were treated with DNase (TurboDNA-free™,
Invitrogen) followed by reverse transcription of 1 µg of RNA to convert into cDNA using
SuperScript IV Reverse Transcriptase kit (Invitrogen), with random hexamers and following
provided protocol. We then diluted cDNA samples to 1:20 prior quantifications. However, *Sry* is
poorly expressed, with variable haplotypes of different lengths (Veyrunes et al. 2013), as well as in
very few regions of the brain (Lahr et al. 1995; Dewing et al. 2006); its signal might be diluted and
difficult to capture using whole-brain RNA and low concentrations. Therefore, for *Sry* assay
specifically, we concentrated DNase-treated RNA samples (i.e.,speed vacuum centrifugation for
~10 min at room temperature) prior the reverse transcription and diluted cDNA to 1:5 to increase
RNA/cDNA yields (up to 2.5 µg of RNA) as well as qPCR performance, All RT-qPCR assays were
performed on a LightCycler 480 (Roche) and using SensiFAST™ SYBR® No-ROX kit (Meridian).
We used *Rps29* as a reference gene, which had previously been used (see Rahmoun et al. 2014).

Expression levels of *Sry, Th, MaoA, Drd2, Ar, Avp* and *Oxt* were measured in triplicates and genes

were distributed across three distinct assays. Primers and qPCR conditions are shown in table 2.

Relative expression levels were measured on LightCycler 480 software v 1.5 (Roche). We included

12 females (4 XX, 2 XX* and 6 X*Y) that did not perform behavioural assays to control for the

treatment effect on expression levels (i.e., quantification of baseline expression levels).

Statistical analyses

All statistical analyses were performed on R v4.5.2 (R core Team 2025). XX/XX* females represent

the control group with low aggression level and are therefore used as the reference group for

downstream analyses. We modelled the probability of offensive aggression between the two

genotypic group (i.e. X*Y *vs* XX+XX* females) with a generalized linear model, fitting a binomial

distribution. We then cumulated aggressive behaviours to determine an aggressiveness level: we

added the frequency of attacks to the percent of time females spent chasing the male (i.e., chasing

duration over the experiment duration), and to the inverse function of the latency of first attack (i.e.,

+1/latency). We log(x+1) transformed the aggressiveness level to control for heteroscedasticity and

compared it between females using a linear model with a Gaussian distribution. We did not include

the age nor the weight of individuals as random effects due to computational limitations (i.e.,

singular fit). Testosterone and gene expression levels were compared between genotypic group with

linear models, fitting a Gaussian distribution. Looking at gene expression specifically, we included

a treatment variable (i.e. whether females performed the behavioural assays or not: 'post-assay' *vs*

'baseline' categories, respectively) as a fixed effect in interaction with the genotypic group.

Pairwise post-hoc comparisons were made using *emmeans* package, and p-values were adjusted for

multiple testing. We then performed a Kendall rank correlation test to assess the relationship

between aggressiveness level and testosterone / gene expression levels and applied a Benjamini-

Hochberg correction on p-values for multiple testing. We acknowledge that the small sample size

might bias our estimations (e.g low power to detect small effects) and accounted for this limitation

in the discussion section accordingly.

Results

Agonistic behaviours

Despite a tendency for a greater aggressiveness in X*Y females (Fig. 1a-e), there is no significant

difference in the probability of being offensive between X*Y females and XX/XX* females (Fig.

1d, GLM, df = 36, estimate = 0.86 ± 0.66 SE, p=0.2), nor there is a significant difference in

aggressiveness level (combined agonistic behaviours, see Fig. 1, table 1) between females (Fig. 1e,

LM, df = 36, estimate = 0.54 ± 0.34 SE, p=0.12).

Androgen pathway: testosterone and its receptor

There is no significant difference in average testosterone levels between X*Y females and XX/XX*

females following the agonistic assay (Fig 2a, LM, df = 35, estimate = 0.16 ± 0.18 SE, p=0.38).

Similarly, there are no differences in *Ar* expression levels between females, regardless of whether

they were tested (Fig 2b, post-hoc Sidak, df = 46, post assay ratio = 2.03 ± 1 SE, p = 0.5,) or not

(post-hoc Sidak, df = 46, baseline ratio = 1.28 ± 0.36 SE, p = 0.5). Baseline *Ar* expression levels are

not significantly different from post-assay expression levels, despite a tendency for higher post-

assay levels in X*Y females (post-hoc Sidak, df = 46, X*Y ratio = 0.41 ± 0.17 SE, p=0.12;

XX/XX* ratio = 0.66 ± 0.26 SE, p=0.76). Finally, there is no significant correlation between

androgens and aggressiveness levels in X*Y (Fig. 2c-d, Kendall test, testosterone: τ = -0.0.7, p=0.7;

Ar : τ =-0.04, p=0.94) nor in XX/XX* females (Kendall test, testosterone, τ =0.31, p=0.1; *Ar* : τ =-

0.12, p=0.64).

Sry and the dopaminergic pathway: Th, MaoA and *Drd2*

The basal expression level of *Sry* in the whole brain of X*Y females is extremely low, with an

average at 2.2e-5 [95% CI = 1.38e-5 - 3.06e-5], and with no significant difference between basal

and post-assay expression levels (Fig. 3a, LM, df = 23, estimate = -3.2e-6 ± 4.6e-6 SE, p=0.5).

Baseline and post-assay *Th* expression levels are not significantly different between X*Y and

XX/XX* females, despite a tendency for a greater baseline expression level in XX/XX* females

(Fig. 3b, post-hoc Sidak, df = 46, baseline ratio = 2.86 ± 1.23 SE, p=0.07; post-assay ratio = 1.13 ±

0.27 SE, p=0.98). However, only X*Y females show higher post-assay *Th* expression levels

compared to baseline expression levels (post-hoc Sidak, df = 46, X*Y ratio = 0.32 ± 0.11 SE,

p=0.008; XX/XX* ratio= 0.81 ± 0.28 SE, p=0.96). Baseline and post-assay *MaoA* expression levels

are lower in X*Y females (Fig. 3c, post-hoc Sidak, df = 46, baseline ratio =7.05 ± 2.27 SE,

p<0.0001; post-assay ratio = 4.26 ± 0.77 SE, p<0.0001), even though X*Y females show a greater

post-assay expression level compared to baseline (post-hoc Sidak, df = 46, ratio = 0.42 ± 0.11 SE,

p=0.006). This difference between post-assay and baseline expression levels is not observed in

XX/XX* females (post-hoc Sidak, df = 46, ratio= 0.69 ± 0.18 SE, p=0.5). *Drd2* expression levels

are not significantly different between females, regardless of whether they were tested (Fig. 3d,

post-hoc Sidak, df = 46, post-assay ratio = 1.17 ± 0.28 SE, p=0.93) or not (post-hoc Sidak, df = 46,

baseline ratio = 2.03 ± 0.89 SE, p =0.31). There is also no significant differences between post-

assay and baseline levels for each genotypic group, despite a tendency for a greater post-assay

expression level in X*Y females (post-hoc Sidak, df = 46, X*Y ratio= 0.43 ± 0.15 SE, p=0.07;

XX/XX* ratio = 0.77 ± 0.27 SE, p= 0.91). Finally, none of these genes have an expression level that

significantly correlates with aggression in X*Y (Fig 3e-h, Kendall test, *Sry* : τ=0.16, p =0.84; *Th*:

τ= 0.03, p=0.94; *MaoA*: τ=0.01, p =0.94; *Drd2*: τ= 0.07, p=0.94) or XX/XX* females (Kendall test,

Th: τ= 0.14, p=0.64; *MaoA*: τ=0.18, p =0.64; *Drd2*: τ= 0.16, p=0.64).

Neuropeptides: Oxt and Avp

Avp expression levels are overall not significantly different between X*Y and XX/XX* females

(Fig. 4a, post-hoc Sidak, df = 46, baseline ratio= 0.88 ± 0.34 SE, p=0.99; post-assay ratio= 0.86 ±

0.19 SE, p=0.93). Moreover, *Avp* expression are very low, with an average relative post-assay level

at 0.0003 [95% CI = 0.0002-0.0004] for instance, whereas the post-assay *Oxt* level is at 0.02 [95%

CI= 0.016-0.035] in average for both genotypic group. There is also no significant differences

comparing baseline and post-assay expression levels for X*Y (post-hoc Sidak, df = 46, ratio= 0.9 ±

0.28 SE, p=0.99) and XX/XX* (post-hoc Sidak, df = 46, ratio= 0.92 ± 0.29 SE, p=0.99) females,

nor there is a significant correlation between aggressiveness and *Avp* levels (Fig. 4b) in each

genotypic group (Kendall test, X*Y, τ = 0.22, p = 0.75; XX/XX*, τ=-0.22, p=0.64). Baseline *Oxt*

expression levels are significantly greater in XX/XX* than in X*Y females (Fig 4c, Post-hoc Sidak,

df = 46, ratio=3.66 ± 1.78 SE, p=0.04) but there is no difference between females looking at post-

assay expression levels (Post-hoc Sidak, df = 46, ratio=1.05 ± 0.29 SE, p=0.99). This is consistent

with post-assay *Oxt* expression levels which are greater than baseline expression levels in X*Y

females only (Post-hoc Sidak, df = 46, X*Y ratio=0.3 ± 0.12 SE, p=0.01; XX/XX* ratio=1.04 ±

0.41 SE, p=1). Finally, we did not find a significant correlation between the level of aggressiveness

and *Oxt* expression levels in X*Y (Fig 4d, Kendall test, τ = 0.22, p=0.75), nor in XX/XX* (τ=0.09,

p=0.64) females.

Discussion

In the African pygmy mouse, sex-reversed X*Y females have been shown to be highly territorial

but the molecular basis of this behaviour is unknown. By correlating behaviours with testosterone

and candidate gene expression levels, we propose that the territoriality of the X*Y females is more

likely to be driven by the influence of sex chromosomes on dopamine and/or neuropeptide levels

than by the androgen pathway.

Tendencies for enhanced aggressiveness in X*Y females

In a previous study, Saunders et al. (2016) highlighted that X*Y females were more aggressive than both XX and XX* females during a resident-intruder assay with the opposite sex (i.e. males): X*Y females had a shorter latency to first attack and attack approximately three times as much as the other females. Here, our results reveal a reduced level of aggressiveness in X*Y females in comparison to this primary observation. We did not find significant differences in aggressiveness level between X*Y and XX/XX* females, although X*Y females tend to be more offensive and XX/XX* seem less likely to be aggressive (Fig. 1). In addition, our results show that the effect size of aggressive behaviours in X*Y females is not as strong as the one previously reported (in Saunders et al. 2016; see Fig. 1), with a greater variance in aggressiveness level. Because we used individuals from the same colony, it is likely that these differences reflect an artificial selection due to breeding conditions (i.e., 'breeding facility' effect). This phenomenon could be common in animals that are held in captivity. In birds for instance, it has been shown that one generation of captive breeding is sufficient for the loss of anti-predatory behaviours (Carrete & Tella, 2015). In our case, there has been a gap of about nine years between experimentations representing 18 generations (given that there are approximately two generations per year in our breeding colony). Considering that highly aggressive X*Y females can be difficult to pair – some were even excluded from the breeding program if they have killed a male (Heitzmann et al. 2023) –, it is likely that highest levels of aggressiveness were artificially selected against in our breeding facility. Consequently, the lack of significance may reflect a combination of lower aggressiveness (i.e., lower effect size), and of a lower statistical power because of a sample size too low to detect smaller effects. These finding highlights that although captivity provide great advantages by enhancing research accessibility, it also has its limitations and can be subject to artificial selection which can bias phenotypic studies. Including animals from the wild would greatly benefits future studies investigating territorial aggression in the African pygmy mouse. Nonetheless, we still observed aggressive females, enabling the establishment of a relationship between aggressiveness and hormonal or gene expression levels.

No causal link between androgens and territoriality

Gonadal hormones and sex chromosomes interact to shape brain neural circuits underlying behaviours. The role of steroid hormones and especially testosterone have been widely investigated in relation to territorial aggression (e.g. Giammanco et al. 2005; Wu et al. 2009; Xu et al. 2012).

Testosterone is a key factor for the masculinization of brain neural pathway, either directly or through the aromatization of testosterone to estrogen (Han & De Vries, 2003; Phoenix, 2009; Phoenix et al. 1959; Wu et al. 2009). More precisely, testosterone has been shown to regulate social aggression in males, but also in females as illustrated in some bird species, rats and humans (Cantarero et al. 2015; Denson et al. 2018; Giammanco et al. 2005; Gill et al. 2007; Rose et al. 1971; Wu et al. 2009). Here, testosterone levels are comparable between females, as previously observed when examining basal testosterone levels (see Veyrunes et al. 2024), albeit concentrations seem overall greater in the present study, which could be explained by the post-behavioural assay context. We did not find a correlation between aggressiveness and testosterone levels, with a slight tendency for a positive relationship in XX/XX* females only, a find that is mainly driven by one outlier individual (Fig. 2c). Because testosterone action depends on its receptor, the androgen receptor (AR) (see Cunningham et al. 2012), we also evaluated the expression of *Ar* and included females that did not perform the test to control for baseline expression levels. It is also noteworthy that the *Ar* gene is on the X chromosome and in the region that does not recombine with the X* chromosome (Saunders et al. unpublished data). Its expression is thus likely to be influenced by differences in the sex chromosome complement between females (e.g., X dosage, divergence between X and X* copies). There are no differences in *Ar* expression levels between females but most importantly, basal and post-assay levels remain within the same expression range, suggesting that resident-intruder encounter does not induce an increase in *Ar* expression. Moreover, there is no apparent correlation between *Ar* expression and aggression. Taken together, our results are not sufficient to support an involvement of androgens in females' territorial aggression in the African pygmy mouse.

It has been shown that the ratio of corticosterone over testosterone could be a more reliable measure to correlate with aggression than testosterone itself (Montoya et al. 2012; Terburg et al. 2009). Interestingly, X*Y females exhibit lower basal corticosterone than XX and XX* females, but this difference disappears in stressful conditions (Veyrunes et al. 2024), which are presumed to occur here. Given the greater fold-change in corticosterone levels between basal and stressful conditions in X*Y females, it is possible that these females display a different sensitivity to corticosterone. Nevertheless, this suggests that the ratio of testosterone/corticosterone is unlikely to be a better predictor of aggressiveness than testosterone alone in *M. minutoides*. Secondly, it has been shown that the intrauterine hormonal environment could also influence aggressiveness, where females developing between two males are more aggressive than females developing with adjacent females (Ryan & Vandenbergh, 2002). However, in our case, due to a complex meiotic drive (Saunders et al.

2022), the least aggressive XX females have mostly brothers, while the most aggressive X*Y
females have mainly sisters, which does not support an intrauterine effect of androgens on the
aggressive behaviours of adult females. Finally, estrogens have also been shown to regulate female
aggression in mice and humans (Denson et al. 2018; Hashikawa et al. 2017; Woodley & Moore,
1999). While we did not include estrogens here, a recent investigation highlighted no differences in
circulating estradiol concentrations between female genotypes of the pygmy mouse (Veyrunes et al.
2024). Nonetheless, future studies looking at estrogen receptors in specific regions of the brain are
mandatory to completely rule out the role of estrogens and more generally, of hormones.

Resident-intruder assays stimulate the dopaminergic pathway in X*Y females only
Dopamine is a catecholamine neurotransmitter that regulates many social behaviours and elevated
levels of dopamine have been associated with aggressive behaviours (Czech et al. 2012; Korzan et
al. 2006; Mahadevia et al. 2021; Scott et al. 2008). In *M. minutoides*, dopamine levels are likely to
vary according to the sex chromosome complement of females (XX, XX* or X*Y). First, the sex-
determining *Sry* gene which can promote dopamine synthesis (Dewing et al. 2006) has been shown
to be expressed in the brain of X*Y females (Rahmoun et al. 2014). Then, the X-linked *MaoA* gene
responsible for dopamine degradation is located in the non-recombining region of X and X*
chromosomes, and in multiple copies on the X chromosome only (unpublished data). Finally,
dopamine precursor (*Th*), and dopamine receptor D2 (*Drd2*) – which inhibits dopamine release and
synthesis in an auto-negative feedback loop –, are also sex-linked in *M. minutoides. Th* and *Drd2*
are located in the recombining regions of two distinct neo-sex chromosomes but because their
inheritance pattern differs between sexual genotype, their expression pattern may vary accordingly
(Saunders et al. unpublished data; Heitzmann et al. 2025). Among these four genes, *Th* and *MaoA*
had previously been identified as potential candidates underlying female genotype divergences in
social behaviours (Heitzmann et al. 2023; 2025). A study on the whole-brain transcriptome revealed
greater *Th* and lower *MaoA* expression levels in X*Y females compared to the other female
genotypes (Heitzmann et al. 2025). Consistently, neuroanatomical investigations of the maternal
behaviours circuitry also suggested greater Th levels in X*Y females, although these results were
not statistically significant (Heitzmann et al. 2023). In the present study, we confirm the overall
lower expression levels of *MaoA* in X*Y females compared to XX and XX* females, but we also
observed a lower basal *Th* expression level in X*Y females, contrary to previous findings. Female
differences in *Th* expression disappear post-assay whereas *MaoA* expression levels remain
significantly lower in X*Y females. Taken together, these results suggest elevated post-assay
dopamine levels in X*Y females compared to XX/XX* females. In fact, only X*Y females show

greater post-assay expression of *Th*, *MaoA* and presumably *Drd2* when compared to basal expression. These results are in line with a stimulation of dopamine metabolism in X*Y females only, where encounters with unfamiliar males induce dopamine production, which in turn triggers compensatory upregulation of its receptor and degradation pathways to restore homeostasis.

Given the absence of differences in *Sry* expression -which was relatively low compared to previous reports (Rahmoun et al. 2014)- it is unlikely that *Sry* enhances dopamine activity via *Th* upregulation (Dewing et al. 2006). However, since *Sry* expression is restricted to specific brain regions (e.g. Lahr et al. 1995), its potential involvement cannot be entirely ruled out, as its signal may be diluted in whole-brain analyses. Taken together, our results suggest that the dopaminergic pathway is stimulated following an agonistic assay in X*Y females only. Nevertheless, because there is no apparent correlation between aggressiveness and gene expression levels, this highlights that aggression is a complex and multifactorial behaviour that cannot be reduced to a few genes. While dopamine may be permissive for aggression, it does not necessarily lead to it; the behavioural output and onset of aggression seem to vary according to other factors. In the future, multifactorial investigations as well as quantification of dopamine levels in brain regions of interest (e.g., ventral tegmental area, nucleus accumbens; Mahadevia et al. 2021) should help unravel the proximate causes of territorial aggression in the African pygmy mouse.

Vasopressin and oxytocin

AVP and OXT are two neuropeptides involved in the regulation of social behaviours including aggression (e.g. Gatewood et al. 2006; Moura Oliveira et al. 2021). While both are known to stimulate maternal aggression (see Bosch & Neumann, 2012), their influence on territorial aggression is rather species-dependant due to controversial results with both stimulating and inhibitory effects (e.g. Bester-Meredith & Marler; de Moura Oliveira et al. 2021; Gatewood et al. 2006; Tan et al. 2019; Everts et al. 1997). For instance, AVP immunoreactive density of fibers correlates positively with aggression in the California mouse, *Peromyscus californicus*, while *Avp* inhibits aggression in rats (Bester-Meredith & Marler; de Moura Oliveira et al. 2021; Everts et al. 1997). Here, our results do not support an increase in Avp expression following the resident-intruder assay, which is moreover very low compared to Oxt levels for instance (Fig 4a). We also did not find a correlation between *Avp* and aggression levels in females (Fig4b), despite a steeper negative slope in XX/XX* females. Given the low levels and narrow range of expression, we assume it to reflect an artefact rather than a true but undetected effect. It is thus unlikely that vasopressin

influences territorial aggression in X*Y females or XX/XX* females. Conversely, our data support
an increase in *Oxt* expression following the resident-intruder assay in X*Y females only, as
observed for dopamine. Given the well-documented interactions between oxytocin and dopamine
systems in shaping social behaviours (Baskerville & Douglas, 2010; Melis et al., 2007; Scott et al.,
2015; Peris et al. 2017), these findings suggest that their interplay may underlie the unique
behavioural phenotype of X*Y females. An interesting possibility is that *Oxt* increase reflects how
females cope with the stress induced by the resident-intruder experiment. In the violet-eared waxbill
for instance, oxytocin has been hypothesized to enable (and not trigger) aggression due to its
anxiolytic effect, as aggressive individuals tend to be less responsive to stress (Goodson et al.
2015). Therefore, one might suggest that O*xt* and dopamine interact in stressful conditions to shape
(at least partly) X*Y female behaviours during female-male encounters, which remains to be tested.
418

Conclusion

X*Y females of the African pygmy mouse have previously been shown to be highly territorial
compared to other females. We highlight that the dopaminergic and Avp/oxytocin pathways, rather
than androgens, are more likely to drive this trait in *M. minutoides*. Because the increase in
dopamine/*Oxt* levels does not correlate with agonistic behaviours, the motivation to be aggressive
may thus depend on other interacting factors. In addition, because changes in gene expression
mostly occur in X*Y females, it is likely that X*Y and XX/XX* females have different proximate
mechanisms promoting aggression. Future investigations such as *in situ* hybridization in specific
regions of the brain (e.g. ventral tegmental area) that include wild animals to remove breeding
effects and other biological factors should help unravel the proximate causes of territorial
aggression. Nonetheless, our findings strongly suggest differences in activation of the dopaminergic
pathway between X*Y and XX/XX* females. Given that this pathway is tightly linked to sex and
neo-sex chromosomes in *M. minutoides*, our findings support an influence of sex reversal on the
dopaminergic system and therefore a direct influence of sex chromosomes on the sexual
differentiation of the brain. This further defies the paradigm that has dominated for decades, of the
unique role of hormones (see Breedlove et al. 1999 ; Arnold, 2009).
435

Acknowledgements

We thank the CECEMA for the housing of animals and Marie Challe for her assistance on the
maintenance of the breeding colony. We also thank the qPCR Platform for allowing us to perform
RT-qPCR and Philippe Clair for his assistance on RT-qPCR as well as on data analysis on

LightCycler 480 software. We thank Emmanuel Valjent and Francis Poulat for constructive

comments and discussions on the topic. We kindly thank Phillip C. Boan for proofreading the paper.

This study was supported by the ANR grant SEXREV (no. 18-CE02-0018-01).

References

Albert, D. J., Jonik, R. H., Walsh, M. L., & Petrovic, D. M. (1989). Testosterone supports hormone-dependent aggression in female rats. *Physiology & Behavior*, *46*(2), 185–189. https://doi.org/10.1016/0031-9384(89)90253-9

Arnold, A. P. (2009). The organizational-activational hypothesis as the foundation for a unified theory of sexual differentiation of all mammalian tissues. *Hormones and Behavior*, *55*(5), 570–578. https://doi.org/10.1016/j.yhbeh.2009.03.011

Baskerville, T. A., & Douglas, A. J. (2010). Dopamine and Oxytocin Interactions Underlying Behaviors: Potential Contributions to Behavioral Disorders. *CNS Neuroscience & Therapeutics*, 16(3), e92. https://doi.org/10.1111/J.1755-5949.2010.00154.X

Baudat F, de Massy B, Veyrunes F. Sex chromosome quadrivalents in oocytes of the African pygmy mouse Mus minutoides that harbors non-conventional sex chromosomes. *Chromosoma* 2019;128:397–411. https://doi.org/10.1007/s00412-019-00699-4.

Bester-Meredith, J. K., Young, L. J., & Marler, C. A. (1999). Species differences in paternal behavior and aggression in peromyscus and their associations with vasopressin immunoreactivity and receptors. *Hormones and Behavior*, 36(1), 25–38. https://doi.org/10.1006/HBEH.1999.1522

Book, A. S., Starzyk, K. B., & Quinsey, V. L. (2001). The relationship between testosterone and aggression: A meta-analysis. *Aggression and Violent Behavior*, *6*(6), 579–599. https://doi.org/10.1016/S1359-1789(00)00032-X

Breedlove, S. M., Cooke, B. M., & Jordan, C. L. (1999). The Orthodox View of Brain Sexual Differentiation. *Brain, Behavior and Evolution*, *54*(1), 8–14. https://doi.org/10.1159/000006607

Cantarero, A., Laaksonen, T., Järvistö, P. E., Gil, D., López-Arrabé, J., Redondo, A. J., & Moreno, J. (2015). Nest Defence Behaviour and Testosterone Levels in Female Pied Flycatchers. *Ethology*, *121*(10), 946–957. https://doi.org/10.1111/eth.12407

Carrete, M., & Tella, J. L. (2015). Rapid loss of antipredatory behaviour in captive-bred birds is linked to current avian invasions. *Scientific Reports*, *5*(1), 18274. https://doi.org/10.1038/srep18274

Cases, O., Seif, I., Grimsby, J., Gaspar, P., Chen, K., Pournin, S., Müller, U., Aguet, M., Babinet, C., Shih, J. C., & De Maeyer, E. (1995). Aggressive Behavior and Altered Amounts of Brain Serotonin and Norepinephrine in Mice Lacking MAOA. *Science*, *268*(5218), 1763–1766. https://doi.org/10.1126/science.7792602

Cox KH, Quinnies KM, Eschendroeder A, Didrick PM, Eugster EA, Rissman EF, Rissman EF. 2015 Number of X-chromosome genes influences social behavior and vasopressin gene expression in mice. Psychoneuroendocrinology 51, 271–281. https://doi.org/10.1016/j.psyneuen.2014.10.010

Cunningham, R. L., Lumia, A. R., & McGinnis, M. Y. (2012). Androgen Receptors, Sex Behavior, and Aggression. *Neuroendocrinology*, *96*(2), 131–140. https://doi.org/10.1159/000337663

Czech, D. P., Lee, J., Sim, H., Parish, C. L., Vilain, E., & Harley, V. R. (2012). The human testis-determining factor SRY localizes in midbrain dopamine neurons and regulates multiple components of catecholamine synthesis and metabolism. *Journal of Neurochemistry*, *122*(2), 260–271. https://doi.org/10.1111/j.1471-4159.2012.07782.x

Darwin, C. (1871). *The Descent of Man and Selection in Relationship to Sex*. London, UK: John Murray

De Moura Oliveira, V. E., Lukas, M., Wolf, H. N., Durante, E., Lorenz, A., Mayer, A.-L., Bludau, A., Bosch, O. J., Grinevich, V., Egger, V., De Jong, T. R., & Neumann, I. D. (2021). Oxytocin and vasopressin within the ventral and dorsal lateral septum modulate aggression in female rats. *Nature Communications*, 12, 2900. https://doi.org/10.1038/s41467-021-23064-5

Denson, T. F., O'Dean, S. M., Blake, K. R., & Beames, J. R. (2018). Aggression in Women: Behavior, Brain and Hormones. *Frontiers in Behavioral Neuroscience*, *12*, 81. https://doi.org/10.3389/fnbeh.2018.00081

Dewing, P., Chiang, C. W. K., Sinchak, K., Sim, H., Fernagut, P. O., Kelly, S., Chesselet, M. F., Micevych, P. E., Albrecht, K. H., Harley, V. R., & Vilain, E. (2006). Direct regulation of adult brain function by the male-specific factor SRY. *Current Biology*, *16*(4), 415–420. https://doi.org/10.1016/j.cub.2006.01.017

Everts, H. G. J., De Ruiter, A. J. H., & Koolhaas, J. M. (1997). Differential Lateral Septal Vasopressin in Wild-type Rats: Correlation with Aggression. *Hormones and Behavior*, *31*(2), 136–144. https://doi.org/10.1006/HBEH.1997.1375

Gatewood, J. D., Wills, A., Shetty, S., Xu, J., Arnold, A. P., Burgoyne, P. S., & Rissman, E. F. (2006). Sex chromosome complement and gonadal sex influence aggressive and parental behaviors in mice. *Journal of Neuroscience*, *26*(8), 2335–2342. https://doi.org/10.1523/JNEUROSCI.3743-05.2006

Giammanco, M., Tabacchi, G., Giammanco, S., Di Majo, D., & La Guardia, M. (2005). Testosterone and aggressiveness. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research*, *11*(4), RA136-145.

Gill, S. A., Alfson, E. D., & Hau, M. (2007). Context matters: Female aggression and testosterone in a year-round territorial neotropical songbird (Thryothorus leucotis). *Proceedings of the Royal Society B: Biological Sciences*, *274*, 2187–2194. https://doi.org/10.1098/rspb.2007.0457

Godar, S. C., Fite, P. J., Mcfarlin, K. M., & Bortolato, M. (2016). The role of monoamine oxidase A in aggression: Current translational developments and future challenges HHS Public Access. *Prog Neuropsychopharmacol Biol Psychiatry*, *69*, 90–100. https://doi.org/10.1016/j.pnpbp

Goodson, J. L., Schrock, S. E., & Kingsbury, M. A. (2015). Oxytocin mechanisms of stress response and aggression in a territorial finch. *Physiology & Behavior*, 141, 154–163. https://doi.org/10.1016/J.PHYSBEH.2015.01.016

Han, T. M., & De Vries, G. J. (2003). Organizational effects of testosterone, estradiol, and dihydrotestosterone on vasopressin mRNA expression in the bed nucleus of the stria terminalis. *Journal of Neurobiology*, *54*(3), 502–510. https://doi.org/10.1002/neu.10157

Hashikawa, K., Hashikawa, Y., Tremblay, R., Zhang, J., Feng, J. E., Sabol, A., Piper, W. T., Lee, H., Rudy, B., & Lin, D. (2017). Esr1+ cells in the ventromedial hypothalamus control female aggression. *Nature Neuroscience*, *20*(11), 1580–1590. https://doi.org/10.1038/nn.4644

Heitzmann LD, Challe M, Perez J, Castell L, Galibert E, Martin AO, Valjent E, Veyrunes F. (2023). Genotypic sex shapes maternal care in the African pygmy mouse, Mus minutoides. *Proceedings of the Royal Society B: Biological Sciences* . 290, 20231224. https://doi.org/10.1098/rspb.2023.1224

Heitzmann LD, Saunders PA, Perez J, Boursot P, Veyrunes F. Influence of Gonadal and Chromosomal Sex on the Brain Transcriptome in a Mouse Species with Natural Sex Reversal. *Molecular Biology and Evolution,* 2025;42:1–14. https://doi.org/10.1093/molbev/msaf208.

Hyde, J. S., & Sawyer, T. F. (1977). Estrous cycle fluctuations in aggressiveness of house mice. *Hormones and Behavior*, *9*(3), 290–295. https://doi.org/10.1016/0018-506X(77)90064-2

Korzan, W. J., Forster, G. L., Watt, M. J., & Summers, C. H. (2006). Dopaminergic activity modulation via aggression, status, and a visual social signal. *Behavioral Neuroscience*, *120*(1), 93–102. https://doi.org/10.1037/0735-7044.120.1.93

Kudryavtseva, N. N., Smagin, D. A., Kovalenko, I. L., Galyamina, A. G., Vishnivetskaya, G. B., Babenko, V. N., & Orlov, Y. L. (2017). Serotonergic genes in the development of anxiety/depression-like state and pathology of aggressive behavior in male mice: RNA-seq data. *Molecular Biology*, *51*(2), 251–262. https://doi.org/10.1134/S0026893317020133

Lahr G, Maxson SC, Mayer A, Just W, Pilgrim C, Reisert I. 1995 Transcription of the Y chromosomal gene, Sry, in adult mouse brain. Mol. Brain Res. 33, 179–182. https://doi.org/10.1016/0169-328X(95)00136-G

Love, T. M. (2014). Oxytocin, motivation and the role of dopamine. In Pharmacology Biochemistry and Behavior (Vol. 119, pp. 49–60). Elsevier Inc. https://doi.org/10.1016/j.pbb.2013.06.011

Lovell-Badge, R. (2005). Aggressive behaviour: Contributions from genes on the Y chromosome. *Novartis Foundation Symposium*, *268*, 20–33. https://doi.org/10.1002/0470010703.ch3

Mahadevia, D., Saha, R., Manganaro, A., Chuhma, N., Ziolkowski-Blake, A., Morgan, A. A., Dumitriu, D., Rayport, S., & Ansorge, M. S. (2021). Dopamine promotes aggression in mice via ventral tegmental area to lateral septum projections. *Nature Communications, 12*(1). https://doi.org/10.1038/S41467-021-27092-Z

Melis, M. R., Melis, T., Cocco, C., Succu, S., Sanna, F., Pillolla, G., Boi, A., Ferri, G. L., & Argiolas, A. (2007). Oxytocin injected into the ventral tegmental area induces penile erection and increases extracellular dopamine in the nucleus accumbens and paraventricular nucleus of the hypothalamus of male rats. *The European Journal of Neuroscience*, 26(4), 1026–1035. https://doi.org/10.1111/J.1460-9568.2007.05721.X

Montoya, E. R., Terburg, D., Bos, P. A., & van Honk, J. (2012). Testosterone, cortisol, and serotonin as key regulators of social aggression: A review and theoretical perspective. *Motivation and Emotion, 36*(1), 65–73. https://doi.org/10.1007/s11031-011-9264-3

Pandolfi, M., Scaia, M. F., & Paz, M. F. (2021). Sexual Dimorphism in Aggression: Sex-Specific Fighting Strategies Across Species. *Frontiers in Behavioural Neuroscience, 15,* 659615. https://doi.org/10.3389/fnbeh.2021.659615

Peris, J., MacFadyen, K., Smith, J. A., de Kloet, A. D., Wang, L., & Krause, E. G. (2017). Oxytocin Receptors Are Expressed on Dopamine and Glutamate Neurons in the Mouse Ventral Tegmental Area That Project to Nucleus Accumbens and Other Mesolimbic Targets. *The Journal of Comparative Neurology*, 525(5), 1094. https://doi.org/10.1002/CNE.24116

Phoenix, C. H. (2009). Organizing action of prenatally administered testosterone propionate on the Tissues mediating mating behavior in the female guinea pig. *Hormones and Behavior, 55*(5), 566–566. https://doi.org/10.1016/j.yhbeh.2009.01.004

Phoenix, C. H., Goy, R. W., Gerall, A. A., & Young, W. C. (1959). Organizing action of prenatally administered testosterone propionate on the tissues mediating mating behavior in the female guinea pig. *Endocrinology, 65*, 369–382. https://doi.org/10.1210/endo-65-3-369

Rahmoun, M., Perez, J., Saunders, P. A., Boizet-Bonhoure, B., Wilhelm, D., Poulat, F., & Veyrunes, F. (2014). Anatomical and molecular analyses of XY ovaries from the african pygmy mouse Mus minutoides. *Sexual Development, 8*(6), 356–363. https://doi.org/10.1159/000368664

R core Team. (2025). *R: A Language and Environment for Statistical Computing.* R Foundation for Statistical Computing. https://www.r-project.org/

Rose, R. M., Holaday, J. W., & Bernstein, I. S. (1971). Plasma Testosterone, Dominance Rank and Aggressive Behaviour in Male Rhesus Monkeys. *Nature, 231*(5302), 366–368. https://doi.org/10.1038/231366a0

Ryan, B. C., & Vandenbergh, J. G. (2002). Intrauterine position effects. *Neuroscience & Biobehavioral Reviews, 26*(6), 665-678. https://doi.org/10.1016/S0149-7634(02)00038-6

Saunders, P. A., Franco, T., Sottas, C., Maurice, T., Ganem, G., & Veyrunes, F. (2016). Masculinised Behaviour of XY Females in a Mammal with Naturally Occuring Sex Reversal. *Scientific Reports, 6*, 1–9. https://doi.org/10.1038/srep22881

Saunders, P. A., Perez, J., Ronce, O., & Veyrunes, F. (2022). Multiple sex chromosome drivers in a mammal with three sex chromosomes. *Current Biology*, *32*(9), 2001-2010.e3. https://doi.org/10.1016/j.cub.2022.03.029

Saunders, P. A., & Veyrunes, F. (2021). Unusual Mammalian Sex Determination Systems: A Cabinet of Curiosities. *Genes*, *12*(11), 1770. https://doi.org/10.3390/genes12111770

Scott, A. L., Bortolato, M., Chen, K., & Shih, J. C. (2008). Novel monoamine oxidase A knock out mice with human-like spontaneous mutation. *Neuroreport*, *19*(7), 739–739. https://doi.org/10.1097/WNR.0B013E3282FD6E88

Scott, N., Prigge, M., Yizhar, O., & Kimchi, T. (2015). A sexually dimorphic hypothalamic circuit controls maternal care and oxytocin secretion. *Nature*, 525(7570), 519–522. https://doi.org/10.1038/nature15378

Shih, J. C., Chen, K., & Ridd, M. J. (1999). Role of MAO A and B in neurotransmitter metabolism and behavior. *Polish Journal of Pharmacology*, *51*(1), 25–29.

Sluyter, F., Van Oortmerssen, G. A., De Ruiter, A. J. H., & Koolhaas, J. M. (1996). Aggression in wild house mice: Current state of affairs. *Behavior Genetics*, *26*(5), 489–496. https://doi.org/10.1007/BF02359753

Tan, O., Musullulu, H., Raymond, J. S., Wilson, B., Langguth, M., & Bowen, M. T. (2019). Oxytocin and vasopressin inhibit hyper-aggressive behaviour in socially isolated mice. *Neuropharmacology*, 156. https://doi.org/10.1016/J.NEUROPHARM.2019.03.016

Terburg, D., Morgan, B., & van Honk, J. (2009). The testosterone–cortisol ratio: A hormonal marker for proneness to social aggression. *International Journal of Law and Psychiatry*, *32*(4), 216–223. https://doi.org/10.1016/J.IJLP.2009.04.008

van Erp, A. M. M., & Miczek, K. A. (2000). Aggressive Behavior, Increased Accumbal Dopamine, and Decreased Cortical Serotonin in Rats. *The Journal of Neuroscience*, *20*(24), 9320–9325. https://doi.org/10.1523/JNEUROSCI.20-24-09320.2000

Veyrunes, F., Chevret, P., Catalan, J., Castiglia, R., Watson, J., Dobigny, G., Robinson, T. J., & Britton-Davidian, J. (2010a). A novel sex determination system in a close relative of the house mouse. *Proceedings of the Royal Society B: Biological Sciences*, *277*(1684), 1049–1056. https://doi.org/10.1098/rspb.2009.1925

Veyrunes, F., Catalan, J., Tatard, C., Cellier-Holzem, E., Watson, J., Chevret, P., Robinson, T. J., & Britton-Davidian, J. (2010b). Mitochondrial and chromosomal insights into karyotypic evolution of the pygmy mouse, Mus minutoides, in South Africa. *Chromosome Research*, *18*(5), 563–574. https://doi.org/10.1007/s10577-010-9144-9

Veyrunes, F., Dobigny, G., Yang, F., O'Brien, P. C. M., Catalan, J., Robinson, T. J., & Britton-Davidian, J. (2006). Phylogenomics of the genus Mus (Rodentia; Muridae): Extensive genome repatterning is not restricted to the house mouse. *Proceedings of the Royal Society B: Biological Sciences*, *273*(1604), 2925–2934. https://doi.org/10.1098/rspb.2006.3670

Veyrunes F, Perez J, Heitzmann LD, Saunders PA, Givalois L. (2024) Hormone profiles of the African pygmy mouse *Mus minutoides* , a species with XY female sex reversal. *Journal of Experimental Zoology Part A: Ecological and Integrative Physiology. 341, 130–137.* https://doi.org/10.1002/jez.2767

Woodley, S. K., & Moore, M. C. (1999). Female territorial aggression and steroid hormones in mountain spiny lizards. *Animal Behaviour, 57*(5), 1083–1089. https://doi.org/10.1006/anbe.1998.1080

Wu, M. V., Manoli, D. S., Fraser, E. J., Coats, J. K., Tollkuhn, J., Honda, S. I., Harada, N., & Shah, N. M. (2009). Estrogen Masculinizes Neural Pathways and Sex-Specific Behaviors. *Cell, 139*(1), 61–72. https://doi.org/10.1016/j.cell.2009.07.036

Xu, X., Coats, J. K., Yang, C. F., Wang, A., Ahmed, O. M., Alvarado, M., Izumi, T., & Shah, N. M. (2012). Modular genetic control of sexually dimorphic behaviors. *Cell, 148*(3), 596–607. https://doi.org/10.1016/j.cell.2011.12.018

Yamaguchi, T., & Lin, D. (2018). Functions of medial hypothalamic and mesolimbic dopamine circuitries in aggression. *Current Opinion in Behavioral Sciences, 24*, 104–112. https://doi.org/10.1016/J.COBEHA.2018.06.011

Zha X et al. 2025 Estrogen signaling in the ventromedial hypothalamus is required for the development of aggression circuitry in male mice. Curr. Biol. 35, 2684-2696.e6. https://doi.org/10.1016/j.cub.2025.04.076

Figure 1. Tendency for enhanced aggressiveness in X*Y females. XX and XX* females formed the control group with low aggressiveness (Saunders et al. 2016). **a-b)** Attack Frequency (i.e. number of attacks standardized by the number of encounters) and percent of time females spent chasing males (i.e. cumulated chasing duration over the 10 min experiment), respectively. Boxplots represent median (central line), 25^{th} and 75^{th} percentile of values (box bounds). **c)** Latency of first attack (s). Values are mean ± 95 % confidence intervals. **d,** Probabilities of offensive aggression. Values are estimated average probabilities ± 95% CI. **e,** Aggressiveness level defined by the sum of agonistic behaviours (attack frequency, percent of time spent chasing and +1 for offensive behaviours). Values are mean ± 95 % confidence intervals. Dots represent raw data. $n_{XX+XX*}=19$ and $n_{X*Y}=19$.

Figure 2. Androgen levels and correlation with aggression in X*Y and XX/XX* females. **a, c)** serum testosterone concentration in X*Y and XX/XX* females following the resident-intruder test and correlation between testosterone levels and the level of aggressiveness, respectively. $n_{XX+XX*}=19$ and $n_{X*Y}=18$. **b,** relative basal and post-assay expression of the androgen receptor (*Ar*) and **d)** correlation between post-assay *Ar* expression and the level of aggressiveness. Baseline, $n_{XX+XX*}=6$ and $n_{X*Y}=6$; Post-assay, $n_{XX+XX*}=19$ and $n_{X*Y}=19$. XX and XX* females formed the control group with low aggressiveness (Saunders et al. 2016). Bars are mean ± 95% confidence intervals and dots/triangles represent raw data. Regression lines are for visual purpose only.

Figure 3. Expression of *Sry* and sex-linked genes from the dopaminergic pathway in the whole brain. Relative baseline and post-assay expression of *Sry* (**a**) *Th* (**b**) *MaoA* (**c**) and *Drd2* (**d**), and correlation between expression levels and aggression (**e**,**f**,**g**, **h** for *Sry, Th, MaoA,* and *Drd2,* respectively) in X*Y and XX/XX* females. Bars are mean ± 95% confidence intervals and dots/triangles represent raw data. XX and XX* females formed the control group with low

555   aggressiveness (Saunders et al. 2016). Baseline, $n_{XX+XX*}=6$ and $n_{X*Y}=6$; Post-assay, $n_{XX+XX*}=19$ and

556   $n_{X*Y}=19$. Regression lines are for visual purpose only. ***$p \leq 0.001$, **$p \leq 0.01$, *$p \leq 0.05$, ·$p \leq 0.1$

Figure 4. Expression of vasopressin (*Avp*) and oxytocin (*Oxt*) in the brain of X*Y and XX/XX* females. Relative baseline and post-assay expression of *Avp* (**a**) and *Oxt* (**c**) and correlations with aggression type (**b**, *Avp*, and **d**, *Oxt*). Bars are mean ± 95% confidence intervals and dots/triangles represent raw data. XX and XX* females formed the control group with low aggressiveness (Saunders et al. 2016). Baseline, n_{XX+XX*}=6 and n_{X*Y} =6; Post-assay, n_{XX+XX*}=19 and n_{X*Y} =19. Regression lines are for visual purpose only. ***p≤0.001, **p≤0.01, *p≤0.05, ·p ≤0.1

Genotype	N	Average female age (months)	Average female weight (g) ± SD	Average intruder age (months)	Average intruder weight (g) ± SD	Mean Latency of 1st attack (s) ± SD	Mean Number of attacks ± SD	Mean Chasing duration (s) ± SD	Mean Number of interactions ± SD
X*Y	19	9.2 ±1.3	8.05 ± 0.9	8.7 ± 2.4	8.8 ± 1.5	93.5 ± 67.18	7.8 ± 9.4	7.4 ± 12.4	14 ± 11.5
XX+XX*	19	9.6 ± 2.4	7.2 ± 1.6	8.7 ± 2.5	8.9 ± 2.2	234.7 ± 153.76	6.7 ± 10	9.4 ± 20.6	16.9 ± 18

596 **Table 1**. Summary statistics of females (residents) and males (intruders) during agonistic assays

Gene name		Primer sequence (5'-3')	Annealing T°C	Primer concentration (μM)	Product size (bp)
Sry	F	AAAAGCCTTACAGAAGACGAAAAA	60°C	0.4	71
	R	TCTCTGTGTAGGGTCTTCAGTCTC			
Th	F	AGTGCACACAGTACATCCGTCATGC	60°C	0.4	128
	R	AATGTCCTGGGAGAACTGGGCAAATG			
MaoA	F	GGGCTGTCATCAAGTGCATG	60°C	0.4	101
	R	TGAAATTGGAGCCTCCTCATCA			
Drd2	F	GCTCACCCTCCTCATCTTCA	60°C	0.4	178
	R	CTCACCCACCACCTCCAG			
Ar	F	CTTCTGTTTCCCTTCAGCGG	60°C	0.4	209
	R	TGGGACCTTGGATGGAGAAC			
Avp	F	ATTTTCCATGCTGTAGGGGC	60°C	0.4	154
	R	GGAACCAAGCAACGCCAC			
Oxt	F	TCAGACTGAGCACCATCGC	60°C	0.4	148
	R	GGGAGACACTTGCGCATATC			

600 **Table 2**. Sequences of primers and conditions for RT-qPCR.

605