1 Experimental evolution of cellular miniaturization reveals a

2 putative mechanism for cell size evolution

3 Garoña Ana.[1], Lemos Victoria Morgane[1], Giometto Andrea[2], Fumasoni Marco[1#].

4 [1] Gulbenkian Institute for Molecular Medicine (GIMM), Oeiras, Portugal

5 [2] Cornell University, School of Civil and Environmental Engineering, Ithaca, USA

6 [#] To whom correspondence may be addressed: marco.fumasoni@gimm.pt

8 ABSTRACT

Cell volume, a key determinant of physiology, is maintained by cell size homeostasis.

Large deviations from typical size are often harmful, yet cell sizes have diverged

drastically in evolution. How size homeostasis evolves to support such diversity without

impairing cellular function remains unclear. To address this question, we used

experimental evolution to select progressively smaller *Saccharomyces cerevisiae* cells.

Over 1,500 generations, we achieved a four-fold modal volume reduction compared to

wild type. Despite this reduction, evolved populations maintained robust size homeostasis

and relatively high fitness, revealing that substantial decreases in cell volume can occur

without severe growth defects. Whole-genome sequencing and genetic perturbations

identify mutations in the G1 cyclin Cln3 and the TOR pathway effector Sch9 as major

contributors to cell size reduction, with the Greatwall kinase Rim15 signaling cascade

further modulating this phenotype. Manipulating these pathways produced a six-fold

range in cell size without changes in ploidy, with engineered mutations recapitulating the

volumes of evolved lines and loss-of-function mutations yielding enlarged cells. Our

results demonstrate the evolutionary plasticity of cell size homeostasis and reveal a

putative mechanism for eukaryotic cell size evolution.

INTRODUCTION

Cells are the fundamental unit of life, and their size is a key determinant of cell and organismal physiology (1). In microbial systems, variation in cell size can shape nutrient acquisition, metabolic scaling, and growth rate, making it a key target of evolutionary and ecological pressures (2). Through its effects on biosynthesis and proliferation, cell size constrains tissue architecture and regulates functional specialization (3–5).

Not surprisingly, cell size varies greatly across the tree of life. Among prokaryotes, cell diameters range from ~0.2 µm in *Mycoplasma* species (6) to nearly 1 cm for *Thiomargarita magnifica* (7). Eukaryotes exhibit a similarly broad distribution, ranging from the smallest known free-living species *Ostreococcus tauri,* with a 1–2 µm diameter (8), to 15 cm long ostrich eggs (9). Even within multicellular organisms, such as humans, cells range from micron-scale lymphocytes to meter-long neurons (10).

Despite this large variability, specific cell types typically maintain their size within a narrow range through a process known as cell size homeostasis (11, 12). At its core, cell size homeostasis relies on the precise coordination of cell growth and division. The most direct evidence of such control on cell size is that cells born small typically grow more before dividing, and vice versa (13). While size control could in theory be performed throughout the cell cycle (14), most organisms exert it mainly prior to DNA synthesis, such as in budding yeast (15) and humans (16), or before nuclear division, such as in fission yeast (17). The general principles followed by cells to regulate their size, as well as the molecular mechanisms that execute this control, have been studied and debated for decades. Cell size control is best understood in the budding yeast *S. cerevisiae*, where cells are thought to commit to division once they reach a critical size (15) or after adding a constant volume each cycle (18). The mechanism that determines the transition from G1 to S phase has been proposed to depend mainly on titration of cell cycle

activators (e.g., Cln3), inhibitors (e.g., Whi5), transcription factors (e.g., SBF), or their

combination (for detailed mechanisms, see refs. (19–26)).

Strong evidence for the importance of cell size homeostasis is represented by the severe

defects experienced by cells as they leave their narrow range of stereotypical sizes.

Excessive cell size has been shown to cause cytoplasm dilution and genetic instability,

ultimately increasing cells' propensity to senescence (27–34). On the other hand,

mutations causing the largest decrease in cell size impair the Target of Rapamycin (TOR)

signaling and ribosome biogenesis, imposing severe reductions in cellular biosynthesis

and fitness (35–38). Notably, altered cell size control has been observed in cancers (39–

42).

It is therefore unclear how evolution has achieved changes in cell size of many orders of

magnitude, when even slight increases or decreases are associated with severe cellular

defects. These detrimental consequences could severely constrain the evolution of the

cell size trait, posing an apparent paradox between the broad evolutionary diversity of cell

sizes and the limited tolerance observed within species. To address this paradox, we

sought to observe in real time the evolution of a large decrease in cell size in the model

eukaryote *S. cerevisiae.* To this goal, we resorted to an experimental evolution regimen

(43–45), simultaneously imposing direct selection on cellular size by Fluorescent-

Activated Cell Sorting (FACS), followed by selection for competitive fitness. Applying this

regimen over 1,500 generations produced miniaturized cells with modal volumes reduced

four-fold relative to wild type (WT), with only marginal fitness tradeoffs. Miniaturized cells

maintained their phenotype in the absence of selection and had size homeostasis

comparable to their ancestor. Whole-genome sequencing revealed adaptive mutations in

key players of the G1 cyclins and TOR pathway, with additional contributions from

components of the Greatwall kinase cascade. Reconstruction of these mutations showed

that their combined effects generate markedly smaller cells while maintaining near-wild-

type fitness, thereby decoupling cell size from growth rate. Targeted engineering of these

pathways substantially altered cell size control and cell cycle dynamics, yielding a six-fold

range in cell volume. Our work demonstrates the plasticity of the mechanisms governing

cell size homeostasis and suggests a putative mechanism for cell size evolution.

RESULTS

To investigate how cell size can evolve under sustained proliferative capacity, we

developed an experimental system in *S. cerevisiae* that alternates two selective regimens

targeting cell size and reproductive fitness, respectively. Cell populations were analyzed

by FACS, using the light scattered by cells (forward scatter area, FSC-A), a widely used

proxy for their size (46). Only the smallest 7% of cells were collected into separate tubes

and subjected to the second selection step (Figure 1A; see SI Appendix and Figure S1A

for details). The second selection consisted of the unconstrained growth of the sorted

population for at least 11 generations, during which cells competed for resources (Figure

1A). After reaching marked turbidity, typical of cells reaching stationary phase, the three

parallel small (S1-3) populations were subjected again to both selection regimens.

Repeating this cycle daily resulted in approximately 1,500 generations of evolution

(Figure 1B). Under these conditions, selection favors mutations that reduce cell size

without compromising proliferation. To speed the evolution, the starting haploid

population (ancestor) carried a mutator allele in the catalytic subunit of DNA polymerase

delta (*pol3-L523D* (47), Table S1). In addition, we engineered the ancestor strain to

express a cytoplasmic fluorescent protein (ymCitrine) to prevent subcellular debris from

being sorted.

Over 1,500 generations, the evolving populations (S1–3) showed a gradual decrease in

FCS values as measured by flow cytometry (Figure S1B). Microscopy and Coulter

counter analyses confirmed that this reduction was accompanied by a marked decrease

in cell volume (Figure 1C–E). Literature values from Coulter counter measurements are

typically reported from exponential-phase cultures, when cells divide rapidly in the

presence of abundant nutrients. In contrast, in our experimental protocol, size selection

occurred as cultures approached the stationary phase, when nutrients become limiting

and cell division progressively slows. To determine whether the evolved reduction in cell

size was maintained across growth phases, we measured cell volumes in both

exponentially growing and saturated cultures (Figure 1C-D and Figure S1C–E). In

exponential phase, WT and Ancestor cells exhibited unimodal size distributions with

modes of 54.7 ± 1.6 fL (mean ± SD) and 61.2 ± 7.29 fL (mean ± SD), respectively. By

generation 1,500, the S-populations displayed an approximately four-fold reduction in

modal size over WT (12.8–13.8 fL; Figure 1C-D), accompanied by ~2.2-fold and ~2.7-fold

decreases in mean and median volumes, respectively (means ~30–31 fL; medians ~22–

23 fL), approaching volumes closer to those of the bacterium *B. subtilis* than to their own

ancestor. Comparatively, perturbations of the TOR signaling pathway (*sch9Δ* and *sfp1Δ*),

which produce some of the smallest previously described yeast cells (35, 48), only

reduced modal volumes by ~2.5-fold (22.8 ± 2.9 fL and 22.6 ± 1.9 fL mean ± SD,

respectively, Figure 1C). Upon entry into stationary phase, WT populations displayed a

broader, and frequently bimodal volume distribution, with peaks corresponding to small

daughter cells (~22 fL) and larger mother cells (~80 fL) (Figure S1C-D). In contrast, the

evolved populations remained unimodal, with a single peak around ~11 fL. Because the

WT stationary-phase distribution reflects a mixture of cell types, we compared median cell

volumes, which revealed that the evolved lines remained approximately 2.5-fold smaller

than WT cells, indicating a robust size reduction also at the end of the growth curve

(Figure S1E). Unless otherwise noted, all cell size measurements reported below were

obtained from exponentially growing cultures.

Together, these results indicate that, to the best of our knowledge, this selection

experiment produced the smallest *S. cerevisiae* cells reported to date under standard

laboratory growth conditions. Modal cell sizes as low as 16 fL have previously been

reported for mutants affecting the G1/S transition when grown in poor carbon sources

such as glycerol; however, these conditions markedly slow cell growth and division (20,

35, 49, 50) (Figure S1F). Comparable size reductions in *S. pombe* have been reported to

cause mitotic catastrophe (51, 52), highlighting the severe constraints on extreme size

reduction. Strikingly, the evolved cells reached volumes similar to those of the smallest

species within the *Saccharomycotina* subphylum (Figure S1G), spanning nearly half of

the natural size diversity observed across the phylum, despite over 400 million years of

divergence (53).

Figure 1. Experimental evolution of cellular miniaturization. (A and B) Schematic representation of size selection and

evolution regimens. A haploid *pol3-L523D* mutator ancestor population was sorted into three small (S) subpopulations and

subjected to daily size selection. Cells in the lowest 7% of the forward-scatter distribution were sorted (100,000 cells) and

cultured in 10 mL rich medium (YPD). Populations were serially grown and sorted for 116 cycles (~1,500 generations), with

samples collected and stored every 24 h. (C) Cell volume distributions of exponential-phase cultures. Smooth density curves

represent kernel density estimates fitted to histograms of cell volumes measured with a Coulter counter. For each strain,

densities were computed per replicate and averaged to obtain a mean density curve. Shaded areas represent the standard

deviation across replicates. In large cell size strains, the first peak corresponds to debris rather than intact cells, a known

Coulter counter artifact that is captured as an artificial peak in the density estimate. (D) Modal cell volume from measurements

of exponential-phase cultures (Coulter counter) across generations. Data from the three S-populations are plotted together

("evolved"), showing progressive evolution toward smaller size, reaching up to a four-fold reduction in volume. Grey lines

indicate wild-type and final evolved sizes. Box plots show medians, interquartile ranges, whiskers (1.5× IQR). (E) Light

micrographs of wild-type and evolved small-cell populations at different cell-cycle stages (G1, S, G2/M). Scale bar, 5 µm.

Gradual cell size decrease reshapes cell cycle dynamics, cellular morphology, and

nutrient-dependent size control

The dramatic changes in cell volumes observed during our evolution experiment

prompted us to examine potential alterations in cellular architecture. The cell cycle is a

major determinant of cell morphology in budding yeast (54). To determine whether

changes in cell size were associated with alterations in the cell cycle, we analyzed DNA

content in exponentially growing populations, where signal intensity distinguishes cell

cycle phases (1C for G1, 2C for G2/M, and 1C < DNA < 2C for S phase). Strikingly, by

generation 1,500, evolved small cells displayed a near absence of G1 compared with WT

(Figure 2A and S2A). Notably, cell cycle changes during evolution occurred in two steps:

during the first ~150 generations, the proportion of cells in G1 increased, followed by a

sharp shift around generation 700 that resulted in the near disappearance of G1 (Figure

2B and S2B). Calculating the absolute time spent in each phase confirmed this trend,

showing that the reduction of G1 duration was primarily compensated by a corresponding

extension of S/G2/M phases (Figure 2B-C and S2B).

We next focused on intracellular organization, reasoning that reduced volume might occur

at the expense of specific organelles. To obtain a general view of subcellular structures,

we performed transmission electron microscopy (55) on exponentially growing WT,

ancestral, and evolved S-populations. Buds smaller than mother cells, consistent with

asymmetric cell division (15), were commonly observed. Interestingly, major organelles

displayed their typical morphology, though with altered proportions, suggesting a potential

differential miniaturization of organelles (Figure 2D and Figure S2C). Despite the

presence of petite clones (cells lacking functional mitochondrial respiration) within the

populations, most evolved isolates, including the smallest, retained growth on non-

fermentable carbon sources, indicating preserved mitochondrial function. Petite clones

did not differ in cell size from non-petites (Figure S2D).

Given preserved respiratory capacity, we tested whether evolved cells maintain the

canonical relationship between nutrient quality and cell size. Typically, cells decrease in

volume in lower-quality media (e.g., non-fermentable carbon sources) relative to high-

quality media such as glucose. When cultured in the non-fermentable carbon source

glycerol, WT and ancestor cells reduced their modal cell size from 55 and 60 fL to 37 and

34 fL, respectively, corresponding to a ~1.5–2-fold decrease in modal cell volume. In

contrast, evolved populations maintained almost identical modal cell sizes in glucose and

glycerol (13.22 vs. 12.12 fL; Figure 2E-F and Figure S2E). This result suggests either that

evolved cells have reached a minimal attainable cell volume, or that the molecular

pathway linking nutrient quality to cell size has been altered to mimic poor-nutrient

conditions even in high-quality media like glucose.

Together, these observations indicate that cellular miniaturization in evolved populations

is accompanied by extensive remodeling of cell cycle progression, cellular organization,

and nutrient-dependent size control.

Figure 2. Physiology of cell miniaturization. (A) DNA-content profiles of ancestral and representative evolved (S1) populations during exponential growth across 1,500 generations, measured by flow cytometry. Lines show means; shaded areas, SD (n = 3). G2 peak height was normalized to 1. Arrows indicate time points of cell cycle profile changes (see main text). (B) Percentage of doubling time spent in G1 and S/G2/M phases for the S1 population at generation 1,500. Bars show mean ± SD (n = 3). (C) Absolute time spent in each cell cycle phase, calculated as the fraction of the doubling time allocated to each phase (56). (D) Transmission electron micrograph of evolved *S. cerevisiae* cells during exponential growth in YPD. (E) Modal cell volume measurements (Coulter counter) grown in glucose versus glycerol as carbon sources. Data from the three S-populations are plotted together ("evolved"). Box plots show medians, interquartile ranges, and whiskers (1.5× IQR). Pairwise comparisons between carbon sources (glucose vs glycerol) were performed using the Mann–Whitney U test; p-values are: WT (p = 0.012), Ancestor (p = 0.045), evolved-Gen1500 (p = 0.051), and *sch9Δ* (p = 1.000). (F) Cell volume distributions of exponential-phase cultures grown in glucose or glycerol as a carbon source.

Evolved cells decouple miniaturized size from pronounced slow growth

Marked decreases in cell size, whether caused by mutations, poor nutrients, or a

combination of both, are typically associated with pronounced slow growth. We therefore

wondered how changes in cell size affected the evolved populations' growth rates.

Growth curves of WT, ancestral, and S-populations revealed a marked increase in

maximum population density in the evolved lines (Figure 3A). Unexpectedly, their growth

rates were only marginally reduced relative to ancestors, which themselves grew slower

than WT, potentially due to the mutator phenotype (Figure 3B and Figure S3A-C). To

precisely estimate this defect, we resorted to pairwise competition assays, which

measure relative fitness, a measure of reproductive success of organisms (57). All three

evolved S-populations had equivalent relative fitness (–7,79% ± 1.4, mean ± SD vs.

Ancestor), whereas WT showed a ~5% advantage relative to the ancestor (Figure 3C). In

contrast, deletion of *SCH9*, which produces some of the smallest cells in rich media,

caused a severe fitness defect (~24% vs. Ancestor & ~28% vs. WT) despite reducing

modal cell volume only 2.5-fold in exponential phase. Across all populations, fitness

followed a non-uniform trajectory: (i) declining during the first 200 generations, (ii)

recovering by generation ~250, (iii) dropping sharply until ~500, and (iv) stabilizing with

minor decreases until 1,500 (Figure 3D, S3D-G). Fitness was more strongly associated

with cell-cycle dynamics than with cell volume (Pearson $r = 0.67$, $p < 0.001$ for G1

fraction; $r = 0.30$, $p = 0.11$ for cell volume), indicating that changes in fitness primarily

reflect alterations in cell-cycle progression (Figure 2B and 3D). Importantly, these results

highlight how evolved lines fall outside of an inverse linear relationship between cell size

and doubling time observable across a wide range of mutants (Figure 3E).

The most pronounced decreases in cell size have previously been associated with

nutrient limitation or mutations in nutrient-sensing pathways. These conditions were

proposed to modulate cell size through reduced ribosome biogenesis (48). However,

numerous microbial physiology studies have linked reduced ribosome biogenesis with

decreased growth rates (58, 59). This raises the question: how do evolved cells maintain

a miniaturized cell size while sustaining growth rate and fitness? To address this, we

examined the biosynthetic capacity of the evolved lines. At saturation, the dry mass

produced per gram of dextrose by evolved lines was comparable to that of WT, whereas

sch9Δ cells produced approximately one-third less (Figure 3F). We next estimated the

ribosomal mass fraction of the proteome from the RNA/protein ratio, a commonly used

proxy for ribosome biogenesis in microbial physiology, also applicable to yeast (59). As

expected, *sch9Δ* cells showed a marked reduction in ribosomal mass fraction, whereas

the evolved lines maintained ribosome biogenesis at levels closer to WT (Figure 3G).

Altogether, these results suggest that evolved lines maintain efficient growth despite a

markedly reduced cell volume, by preserving ribosome biogenesis and biomass

production.

Figure 3. Fitness and biomass production. (A) Growth curves of selected strains measured as cell density (cells/mL) versus

time (h). (B) Maximum growth rates of reference and evolved strains (generation 1,500) in YPD. Box plots show medians,

interquartile ranges, and whiskers (1.5× IQR). (C) Relative fitness of wild-type, ancestral, and evolved populations (S1–S3) at

generation 1,500. Relative fitness is the ratio of reproductive success of a strain to that of a reference strain under identical

conditions. Fitness of evolved and *sch9Δ* strains differed significantly from the ancestor and wild type (adjusted $p < 0.01$), with

no significant differences among evolved lines. Box plots as in (B). Full pairwise statistics are provided in Supplementary file:

RawData-Figures. (D) Fitness and cell-size trajectories across 1,500 generations for the S1 population. Lines show mean

values; shaded areas show SD. (E) Relationship between fold decrease relative to WT median cell size and doubling time in *S.*

cerevisiae. Circles represent data from (35, 49, 50). Diamonds represent strains measured in this study. Error bars represent

propagated measurement uncertainty. The solid black line shows a linear regression computed using only mutant strains, with

261 the shaded area representing the 95% confidence interval. Genes of interest are annotated. (F) Biomass derived from dry mass

262 per cell, converted to yield relative to dextrose content in the medium. (G) Scaling of the ribosomal fraction of the proteome with

263 the population's growth rate. Data from (60) (circles) are compared with experimental measurements obtained in this study

264 (diamonds). A linear regression (dashed line) was fitted to the ribosomal fraction values in (60). The total RNA-to-protein ratio

265 (RNA/protein) is plotted on a secondary y-axis as a proxy for the ribosomal fraction of the proteome.

New evolved size homeostasis is stable and robust

How stable is the evolved miniaturized size? Measurements performed right after the last sorting round may reflect a transient physiological state, rather than an evolutionarily stable phenotype. To address this possibility, clones were isolated from two independent populations (S1Cl2 and S2Cl4) and passaged for 150 generations in the absence of any size selection. Interestingly, S-clones stably maintained their size across the experiment, resulting in a final fold change similar to that observed in control WT lines (Figure 4A and Figure S4A-D). In contrast, *sch9Δ* lines consistently reverted to WT sizes between generation 50 to 100 despite starting with a larger volume than S-clones (Figure 4A and Figure S4A-D). A key question is whether miniaturized cells actively regulate their size, or whether they merely preserve the volumes attained at the end of the experiment. To address this, we transiently arrested cells at the G1/S transition using the pheromone α-factor, which permits growth while blocking division (61) (Figure S4E-F). While the arrest was not fully efficient in S-clones, it nevertheless led to a ~1.8-fold increase in cell volume in both WT and evolved cells. The increased volume, however, was progressively reversed upon release into fresh medium, indicating that cells actively return to their characteristic small size (Figure S4E, S4G).

The ability of miniaturized cells to maintain their size across many generations and under transient perturbations reflects efficient size homeostasis, which is expected to manifest as a narrow distribution of cell sizes around the population mean. To test this prediction, we compared the width of cell size distributions across WT, ancestral strains, and evolved populations at successive generations (Fig. 4A). To avoid spurious contributions from cell debris and non-cellular particles at low volumes, which bias conventional statistics such as

the mean and standard deviation, we quantified distribution width as the mean excess above

the modal volume $E[v - v_{mode}|v \geq v_{mode}]$, normalized by the modal volume (SI Appendix).

This dimensionless metric captures the average relative width of the right tail of the size

distribution independently of debris at low volumes (Figure 4B). Over the course of evolution,

populations initially reduced both their modal volume and their normalized mean excess,

converging toward values below those of the ancestor and comparable to WT (Figure 4C).

Once modal volumes fell below ~30 fL, however, normalized mean excess increased again,

returning to values comparable to the ancestor despite a dramatically reduced characteristic

cell size. Overall, miniaturized S-clones thus maintain a degree of size homeostasis similar

to the ancestor, demonstrating that the reduction in cell size was not accompanied by a

proportional loss of size control.

Together, these results demonstrate that miniaturized cells have evolved a new cell size

homeostasis that is evolutionarily stable, robust to transient perturbations, and capable of

maintaining population-level size control as tightly as their own ancestor.

Figure 4. Evolution of cell size homeostasis. (A) Stability of cell size during unconstrained experimental evolution. Left: cell-size measurements of WT, Ancestor, *sch9Δ*, and evolved clones at 1,500 generations from populations evolved for 150 generations without size selection. Box plots show medians, interquartile ranges, and whiskers (1.5× IQR). Pairwise comparisons between strains were performed using the Mann–Whitney U test: WT vs WTGen150 (p = 0.328), Ancestor vs AncestorGen150 (p = 0.082), S1Cl2 vs S1Cl2Gen150 (p = 0.011), S2Cl4 vs S2Cl4Gen150 (p = 0.008) and *sch9Δ* vs *sch9Δ*Gen150 (p = 0.015). Right: fold change in cell size relative to time 0. (B) Our metric of cell size distribution width calculates the mean excess above the mode, which is a generalization of the half-width at half maximum that accounts for the distribution width across all density values, rather than just the half maximum (SI Appendix). We focus on the right tail of the distribution to avoid spurious effects due to the presence of cell debris at small cell volumes, highlighted here by logarithmic cell volume binning. Shown here is the distribution of WT (W303). (C) Width of cell size distributions, quantified as the mean excess above the mode (SI Appendix), normalized by modal volume. The Ancestor (blue, ~62.5 fL modal volume, normalized mean excess ~2.8) had a broad right tail relative to WT, leading to a large normalized mean excess. Evolved populations progressively reduced both modal volume and normalized mean excess over generations, with early-generation populations (Gen100–Gen200) converging near WT (W303) values (~50-55 fL, normalized mean excess ~1.2). At modal volumes below ~30 fL, the normalized mean excess rose again to ~2.5, approaching the ancestral value.

Genetic landscape of cellular miniaturization

We sought to determine the genetic basis of the evolved phenotype. Evolved cells

retained their characteristic size when co-cultured with wild-type strains, indicating that

miniaturization was not driven by secreted or diffusible factors (Figure S5A). The

phenotype's persistence over 150 generations in the absence of selection (Figure 4A)

strongly suggested a genetic basis. Supporting this idea, crosses between evolved clones

and WT produced diploids of intermediate size between haploid and diploid WT,

consistent with the presence of both recessive and dominant mutations. Furthermore,

crosses between evolved clones and mating-type–switched clones yielded diploids that

retained a large reduction in cell volume relative to WT diploids (Figure S5B), suggesting

that the reduced cell size is independent of ploidy.

To identify causative mutations, we performed whole-genome, whole-population

sequencing across multiple evolutionary timepoints. As expected from the mutator allele

present in the ancestral population, we detected 869 new unique mutations that arose

during evolution in coding and regulatory regions across the three populations (Figure 5A,

Table S2). By the end of the experiment, an average of 267 single-nucleotide

polymorphisms (SNPs) and small insertions/deletions (indels) persisted per population

(285 in S1, 301 in S2 and 216 in S3, Figure S5C). No aneuploidies, segmental

amplifications, or large-scale changes in DNA copy number were observed, thus

confirming the presence of a full haploid genome (Figure S5D). The evolved populations

exhibited distinct clonal compositions (Figure S5E), however, several alleles were shared

across populations. These mutations were absent from the ancestral strain and, in most

cases, involved identical nucleotide substitutions. The probability that a large number of

identical mutations arose independently across populations is therefore extremely low,

pointing towards a likely cross-contamination during the FACS selection steps.

Accordingly, the evolved lines were treated as a single evolving population for

subsequent analyses rather than as independent replicates (Figure 5A).

To identify putative adaptive mutations associated with the evolved cell size, we

employed four non-exhaustive, but complementary approaches: (i) tracking lineages that

rose to fixation in any of the three sub-populations, identifying cohorts of mutations under

positive selection (Table S2); (ii) searching for genes previously implicated in cell size

regulation (Table S3); (iii) using statistical inference to identify genes mutated more

frequently than expected given their length and the observed mutation rates (parallel

evolution; Table S4); and (iv) performing two consecutive rounds of backcrossing and

selection for small cell size to pinpoint mutations segregating with the phenotype (bulk

segregant analysis; Table S5; see S1 Appendix for details). A summary of the genes

identified by these methods, as well as their reported interactions, is portrayed in Figure

5B-C (Table S6). Despite several genetic and physical interactions among the identified

genes, no gene ontology term was found to be statistically enriched.

Our analysis indicates that the miniaturized phenotype is heritable and genetically

encoded, but the underlying adaptive landscape is complex. The large number of

candidate mutations likely includes false positives arising from hitchhiking or linkage, as

well as variants under indirect selection. Some of these may reflect selection on traits

influencing the flow-cytometry size proxy (FSC-A) rather than cell size itself, or

compensatory adaptations to physiological changes associated with miniaturization (62).

Figure 5. Genomic landscape of the evolved phenotype. (A) Allele-frequency trajectories of mutations detected in the combined population (S1–S3). A 300% frequency indicates fixation in all three populations. Line thickness reflects the final frequency of the mutation at generation 1,500. (B) List of putative adaptive gene mutations ranked by the number of satisfied criteria outlined in the main text and affected by non-synonymous mutations located within coding sequences (CDS). Only genes appearing in at least three categories are included (full list in Table S6). (C) Interaction network of mutations detected in S1–S3 evolved populations curated in Cytoscape (52). Grey lines indicate known genetic and physical interactions (STRING database (53)). Node diameter reflects the number of satisfied criteria we used to detect putative adaptive mutations. Only genes appearing in at least two categories are included. Nodes are color-coded: dark yellow for non-synonymous mutations within CDS fixed at generation 1,500, light yellow for mutations in the regulatory regions fixed at generation 1,500, and grey for non-fixed mutations. Bold outlines mark genes linked to cell size changes in the literature (either decrease or increase).

Combined mutations in the G1 cyclin and Greatwall signalling cascades cause

large deviations in cell size

We set out to identify the mutations primarily responsible for the change in cell size among those identified as putatively adaptive. *CIM1*, *CLN3*, and *SCH9* ranked highest in our analysis and were therefore selected for further investigation. *CIM1* encodes for a mitochondrial HMG-box protein that limits mitochondrial DNA copy numbers (63), and independently accumulated up to six mutations in the protein over the course of the experiment, including three frameshifts resulting in premature stop codons (Figure S6A). Cln3, a G1 cyclin that partners with Cdk1 to drive the G1/S transition by activating phase-specific gene expression, has been shown to reduce cell size when truncated at its C-terminus (e.g. *cln3-1*), due to increased protein stability (49, 64, 65). Our evolved strains carried a similar truncation, *cln3-S410**, just downstream of the *cln3-1* allele (Figure 6A). Sch9 is a key effector of the TORC1 pathway, which integrates nutrient sensing with anabolic and catabolic processes. We identified a mutation (*sch9*-L343S) within the conserved C2 domain (residues 184–402), a calcium-dependent membrane-targeting module that may also inhibit the kinase domain (residues 403–738; Figure 6A, (66)).

To test the causality of these mutations, we reconstructed the *cim1*, *cln3*, and *sch9* alleles in a WT background. Deletion of *CIM1*, used to mimic the observed frameshift mutations, did not alter cell size (Figure S6A), suggesting a potential adaptive role in other cellular features or fitness. In contrast, mutations in *CLN3* and *SCH9* individually caused a modest but significant decrease in cell volume (Figure 6C). Combining the two mutations produced synergistic effects, with the double mutant exhibiting a three-fold reduction in modal volume (28.5 fL; Figure 6C).

Truncations of the C-terminal region of Cln3 confer a gain-of-function phenotype by increasing protein stability and cyclin abundance (65). Loss-of-function mutations in *SCH9*

have likewise been associated with reduced cell size, typically attributed to its role in

ribosome biogenesis (48). However, the elevated ribosome biogenesis observed in evolved

lines (Figure 3G), together with their sustained fitness (Figure 3B–C), indicates that *sch9-*

L343S is a separation-of-function allele that affects cell size through a mechanism distinct

from ribosome biogenesis.

Recent work suggests that Sch9 can also regulate cell size via the Greatwall kinase Rim15,

albeit under conditions of low CDK activity (67). This pathway is highly conserved among

eukaryotes and controls cell cycle progression in metazoans through endosulfine-mediated

inhibition of the phosphatase PP2A (68, 69). While it links nutrient sensing to cell size in *S.*

pombe (70), in *S. cerevisiae* it has primarily been implicated in the regulation of quiescence

(71). We identified mutations across multiple components of this pathway, including a fixed

mutation in *RIM15*, as well as lower-frequency mutations in the promoter of the endosulfine

IGO1 and in *CDC55*, which encodes a regulatory subunit of PP2A (72). Reconstruction of

the evolved *RIM15* allele (*rim15-A132T*) resulted in a modest, non-significant reduction in

cell volume (Figure 6C). However, the distribution of mutations across the pathway (*sch9-*

L343S, *rim15-A132T*, *igo1-T-75G*, and *cdc55-L159P*; Figure S6B) suggests that the

Greatwall pathway activity was modulated during evolution. To test this hypothesis, we

engineered a copper-inducible tunable promoter (*prCUP1*) to control expression of *RIM15*

and *rim15-A132T*. Copper titration revealed a clear dosage-dependent effect: basal

expression reduced cell size, and overexpression induced further shrinkage in WT

backgrounds (Figure S6B). Notably, overexpression of *rim15-A132T* in the *sch9-L343S cln3-*

S410* background further reduced modal cell volume, matching that of the evolved lines (16

± 0.1 fL, mean ± SD; Figures 6C and S6C–F). Consistent with increased Greatwall pathway

activity, deletion of *RIM15* increased exponential-phase cell volume in evolved clones but

had little effect in WT (Figure 6D and Figure S7A). In stationary phase, however, *RIM15*

deletion increased cell size in both backgrounds, with a stronger effect in evolved cells

(Figure S7B-D). Furthermore, combined deletion of *RIM15* and *cln3Δ* showed synergistic

effects on increasing WT cell size in stationary phase (Figure S7E-J).

Importantly, despite their reduced size, all reconstructed strains carrying evolved alleles

retained near-WT fitness (Figure 6C). Although enforced expression of *rim15-A132T*

phenocopied the size of evolved clones in the *sch9-L343S cln3-S410** background, it

imposed a fitness penalty. This penalty is unlikely to arise from reduced cell size itself, as

fitness improved while size decreased further at higher copper concentrations (Figure 6C),

and instead points to deleterious consequences of constitutive Rim15 activation across the

cell cycle. These observations suggest that, rather than simply increasing pathway output,

evolution selected for combinations of mutations that tune Greatwall signaling in a context-

dependent manner to minimize fitness trade-offs.

These results demonstrate that cell size in the evolved lines is dependent on Rim15 dosage

and support a model in which, together with a hyperactive G1 cyclin Cln3, increased

Greatwall pathway activity contributes to reduced cell volume, although the precise

molecular mechanisms remain to be defined.

Figure 6. Genetic basis of the cell size phenotype. (A) Schematic of cell size–regulating genes. Protein domains are indicated, and mutations identified during evolution are marked: PK, kinase catalytic domain; D, consensus disorder prediction; PAS, N-terminal PAS domain; C2HC, CCHC-type zinc-finger domains; REC, C-terminal receiver domain. (B) Allele-frequency trajectories of size-associated mutations. Only variants reaching ≥80% frequency by generation 1,500 are shown; causative mutations are highlighted. The right axis shows the S1 population volume (circles) over time. (C) Relationship between relative fitness and modal cell volume. Each point represents a strain (wild type, ancestor, evolved populations, or engineered mutants). Fitness (x-axis) and cell volume (y-axis) are shown; error bars indicate SD. (D) Modal cell volume following deletion of *RIM15* in the wild-type background and in final evolved clones from exponential-phase cultures. Box plots show medians, interquartile ranges, whiskers (1.5× IQR).

DISCUSSION

How can evolution produce large shifts in cell size without incurring the detrimental effects associated with even small deviations from the norm? In development, abrupt changes often arise through ploidy increase or reduction (73). Linear scaling relationships between DNA content and cell volume are also observed across diverse taxa (74). Ploidy increases are important in plant evolution and likely account for part of the variation in cell

size among related species (75). Yet, genome duplications alone cannot explain the wide

diversity of cell sizes, as they also create barriers to sexual reproduction and would leave

more pervasive genomic signatures if they were the main driver. Cells must therefore be

able to modify size independently of DNA content, at least initially, especially under

gradual selection for divergent sizes.

Decades of studies on eukaryotic cell cycle regulation have identified the main principles

allowing cells to regulate their size, as well as many molecular players involved (reviewed

in (11, 76). Mutations in many such players have been demonstrated to produce altered

sizes (35, 36, 77). It is therefore intuitive to imagine that evolution could modulate cell

size by fine-tuning these molecular players. Yet most such mutations reduce fitness, and

it remains unclear which, if any, combinations of the dozens of known regulators could

provide viable evolutionary paths to new cell sizes.

We addressed this question experimentally by imposing selection on *S. cerevisiae* cell

size and fitness. Over 1,500 generations, cells shrank four-fold in modal volume

compared to WT, without changes in ploidy and with minimal fitness costs. We show that

miniaturized cells maintain an efficient and evolutionarily stable cell size homeostasis,

despite a significantly altered cell cycle progression characterized by the near absence of

a G1 phase. Whole genome sequencing revealed a complex genomic landscape, due to

the mutator allele employed, and likely linked to the many direct and indirect changes

cells experienced during the miniaturization. Nevertheless, re-engineering putative

adaptive alleles showed that simultaneous perturbations of the G1 cyclin Cln3 and the

TOR effector Sch9 recapitulate much of the evolved phenotype with minimal fitness cost

(Figure 6C). Although consistent with a gain-of-function effect from C-terminal truncation

of Cln3, our data do not support a primary role for Sch9 in size control via ribosome

biogenesis (Figure 3G). We argue instead for its involvement as an inhibitor of Rim15.

Together, manipulations in the G1 cyclin and Greatwall kinase signaling cascades

produced a six-fold range within an otherwise constant genome, capturing up to 70%

(221 / 315) of the natural variation in size seen across the *Saccharomycotina* over 400

million years of evolution (Figure 6C, S7E-J and Figure 7A).

Three questions arise. First, how can our evolved cells reach sizes smaller than those

achieved previously through gene deletions? We propose that evolution selected alleles

with modest individual effects, but that act synergistically on cell size by working on

multiple mechanisms simultaneously. We speculate the following model to account for the

phenotype observed (Figure 7B): Truncations in *CLN3* stabilize the protein, increasing

CDK1–Cln3 activity and phosphorylation of the transcriptional inhibitor Whi5 (78). In

parallel, loss of Sch9 inhibition and gain-of-function mutations in the Greatwall kinase

cascade inhibit the phosphatase PP2A–Cdc55, further driving Whi5 phosphorylation and

nuclear exclusion (67). But why would this lead to a volume smaller than *whi5Δ*, where

the inhibitor is absent? First, CDK1-Cln3 levels could simultaneously hyper-phosphorylate

the transcriptional regulator Stb1 (20, 79, 80) and accelerate the degradation of the CDK1

inhibitor Sic1 (81–83), eliminating the G1 phase. Furthermore, *RIM15* activity in mitosis

has been reported to destabilize the CDK1 inhibitor and size/morphogenesis checkpoint

Swe1 (84), offering an additional route to reduced volume. To test this model, we deleted

WHI5 and *SWE1*, inhibitors of the G1/S and G2/M transitions, respectively, in evolved

clones (Figure 7C and Figure S8A). *WHI5* deletion increased cell size in evolved clones

(19.3fL), whereas *SWE1* deletion caused a stronger size reduction than in WT (9% vs

33% decrease), producing cells with a modal volume of ~9 fL. These results indicate that

whi5Δ no longer contributes to size decrease in evolved lines, while Swe1 plays a more

prominent role, suggestive of cell size regulation occurring in G2/M rather than G1/S.

The second question regards what allows these miniaturized cells to retain high fitness.

Eukaryotic cells physiologically shrink upon nutrient scarcity (49). Under these conditions,

cells may find it advantageous to increase population-wide survival by dividing at a smaller

volume while making a parsimonious use of the resources left by reducing ribosome

biogenesis. The deletion of *SCH9* hard-wires this response, resulting in small cells with

severe fitness defects due to strongly decreased protein production. Here, we show that our

evolved lines do not shrink further in poor-quality nutrients. Because we achieved even

smaller cells through the deletion of Swe1, we conclude that the evolved lines have

genetically altered the mechanism linking nutrient quality to cell size, rather than reaching a

minimal size limit. Although such a mechanism has been proposed to operate through the

modulation of ribosome biogenesis, our results show that ribosome biogenesis remains

efficient in our evolved lines. This indicates that the effect of nutrients on cell size has

become uncoupled from their effect on biomass production. This decoupling explains the

high fitness of these evolved miniaturized cells and challenges current models of nutrient

control of cell size.

Finally, have the evolved cells reached the lower size limit for *S. cerevisiae*? Gain-of-function

mutants of the G1 cyclin Cln3, when grown in glycerol, approach the size of our evolved

lines (16 fL, (53)). This observation supports our model in which the evolved lines combine

size effects from the G1 cyclin pathway with those mediated by nutrient signaling. However,

we argue that this combined effect does not represent the lower physical limit, as additional

deletion of *SWE1* further reduces the modal cell volume to ~9 fL. The long-term continuation

of this evolutionary experiment will help determine whether, and how, a physical lower size

limit can be reached experimentally.

In summary, our work shows that large deviations in cell volume can be achieved while

maintaining cellular fitness and size homeostasis. We show that this phenomenon is

accompanied by changes in cell cycle regulation and in the nutrient-dependent control of cell

size. To achieve this phenotype, we propose a mechanism based on the simultaneous gain

of function of the G1 cyclin and Greatwall kinase signaling pathways. In this model,

increased Greatwall pathway activity arises, at least in part, from the relief of Sch9-mediated

inhibition, thereby linking nutrient signaling to cell cycle control. We speculate that this model

identifies a simple yet powerful route for evolution to achieve large deviations in volume

without altering ploidy. Future studies should test the generality of the principles we identified

and determine how broadly it underpins size diversity across the tree of life. The range of

cell sizes generated within a single species, while maintaining high fitness and stable ploidy,

also offers a framework to study subcellular scaling and the limits of cell size.

Figure 7. Speculative model of the effect of the mutations identified on cell cycle regulation. (A) Distribution of cell

volumes across the *Saccharomycotina* subphylum (derived from (83)). Highlighted are the triple mutant (*sch9-L343S cln3-*

S410 prCUP1-rim15-A132T*) under a copper-inducible promoter and the *CLN3* and *RIM15* double deletions, showing shifts

in size relative to natural variation. (B) Modal cell volume of deletion of *WHI5* and *SWE1* in the wild-type background and in

final evolved clones from exponential-phase cultures. Box plots show medians, interquartile ranges, and whiskers (1.5×

IQR). (C) In wild-type (WT) cells growing under nutrient-rich conditions, TOR is active, promoting ribosome biogenesis and

biomass accumulation (right, green rectangle) while delaying cell cycle transitions (left, orange rectangle). This results in

fast-growing, large cells. Under nutrient limitation or in TOR-related mutants, Sch9 repression of Rim15 is relieved, leading

to earlier cell cycle transitions. However, the accompanying reduction in biomass production limits growth and fitness. We

propose that the combined effect of the identified mutations accelerates cell cycle transitions through truncations in Cln3,

loss of Sch9 inhibition, and gain-of-function mutations in the Greatwall kinase cascade, while maintaining ribosome

biogenesis and biomass production. As a result, cells grow rapidly but, by spending less time in interphase, remain smaller

than their WT counterparts.

MATERIAL AND METHODS

All strains used in this study were W303 derivatives. Yeast cultures were propagated using

standard techniques (85). Unless otherwise stated, experiments were performed in standard

rich medium (YPD). Experimental evolution followed the procedure described in (86), with

the addition of the size-selection step by sorting detailed in the SI Appendix. Growth-curve

measurements and fitness assays were performed as described in (87). Samples for cell-

cycle analyses were processed as in (87) and analyzed as in (86). Whole-genome

sequencing and analysis followed procedures described in (88). A full description of all

methods is provided in the SI Appendix, Methods.

Data Availability

The genomic datasets used within this publication were deposited at European Nucleotide

Archive (PRJEB101156). Scripts used for data analysis are available at the GitHub repository

(https://github.com/FumaLab/Garona2025). The raw data corresponding to each figure are

provided in the Supplementary file Raw Data–Figures. Expanded view data, supplementary

information, and appendices are available for this paper. Raw data from Coulter Counter

measurements is available at Zenodo (DOI: 10.5281/zenodo.19254220).

Authors contribution

M.F. conceived the project. A.Ga. and M.F. designed the research, performed the

experiments, and analyzed the data. A.Gi. contributed data analysis for Figure 4B–C. M.V.L.

contributed experimental work for Figure 5C, E. A.Ga. and M.F. wrote the original

manuscript draft. All authors edited and commented on the final version of the manuscript.

Acknowledgments

We thank Mina Abrantes, Joana Boas, Filipa Cantanhede, Olga Carreira, João Martins, and

Ana Quendera for experimental assistance; members of the Genome Maintenance and

Evolution lab for support and discussions; and Gabriel Neurohr, Andrew W. Murray, Paula

Gonçalves, Partha Pratim Chakraborty, and Mariana Natalino for manuscript review. We

acknowledge the Flow Cytometry, Electron Microscopy, Bioimaging, and Genomics Facility

of the Gulbenkian Institute for Molecular Medicine for technical support, and Beatriz Tomaz,

Erin Tranfield, and Muriel Christel Mari for assistance with establishing electron microscopy

sample preparation. We also thank Giorgio Tallarico and Marco Cosentino Lagomarsino for

advice on quantifying the ribosomal fraction of the proteome. This work was supported by

the Gulbenkian Foundation and the Gulbenkian Institute for Molecular Medicine (A.Ga.,

M.V.L., M.F.); Human Frontier Science Program (HFSP, RGEC28/2023; M.F., A.Gi.);

European Molecular Biology Organization (EMBO, IG 5349-2023; M.F.); Fundação para a

Ciência e a Tecnologia (FCT, 2023.09068.CEECIND; M.F.); and the National Institute of

General Medical Sciences (NIGMS), National Institutes of Health (NIH), USA

(1R35GM147493; A.Gi.).

BIBLIOGRAPHY

1. M. B. Ginzberg, R. Kafri, M. Kirschner, Cell biology. On being the right (cell) size.
Science **348**, 1245075 (2015).

2. F. Vasseur, L. Mahaut, B. J. Enquist, C. Violle, From organism traits to ecosystem
processes: Why size is so important. *Annu. Rev. Ecol. Evol. Syst.* **56**, 145–170
(2025).

3. C. Cadart, R. Heald, Scaling of biosynthesis and metabolism with cell size. *Mol. Biol.*
Cell **33** (2022).

4. T. P. Miettinen, M. J. Caldez, P. Kaldis, M. Björklund, Cell size control - a mechanism
for maintaining fitness and function. *Bioessays* **39**, 1700058 (2017).

5.   T. L. Orr-Weaver, When bigger is better: the role of polyploidy in organogenesis.
Trends Genet. **31**, 307–315 (2015).

6.   M. G. Gabridge, Current status of the mycoplasmas: The smallest living cells. *Lab.*
Med. **9**, 36–40 (1978).

7.   J.-M. Volland, *et al.*, A centimeter-long bacterium with DNA contained in
metabolically active, membrane-bound organelles. *Science* **376**, 1453–1458 (2022).

8.   M.-J. Chrétiennot-Dinet, *et al.*, A new marine picoeucaryote: Ostreococcus tauri gen.
et sp. nov. (Chlorophyta, Prasinophyceae). *Phycologia* **34**, 285–292 (1995).

9.   C. Di Meo, G. Stanco, M. I. Cutrignelli, S. Castaldo, A. Nizza, Physical and chemical
quality of ostrich eggs during the laying season. *Br. Poult. Sci.* **44**, 386–390 (2003).

10.   R. Milo, R. Philips, Cell Biology by the Numbers. Available at:
https://book.bionumbers.org/ [Accessed 23 October 2025].

11.   A. A. Amodeo, J. M. Skotheim, Cell-Size Control. *Cold Spring Harb. Perspect. Biol.*
8, a019083 (2016).

12.   N. Rhind, Cell-size control. *Curr. Biol.* **31**, R1414–R1420 (2021).

13.   G. C. Johnston, J. R. Pringle, L. H. Hartwell, Coordination of growth with cell division
in the yeast Saccharomyces cerevisiae. *Exp. Cell Res.* **105**, 79–98 (1977).

14.   X. Liu, J. Yan, M. W. Kirschner, Cell size homeostasis is tightly controlled throughout
the cell cycle. *PLoS Biol.* **22**, e3002453 (2024).

15.   L. H. Hartwell, M. W. Unger, Unequal division in Saccharomyces cerevisiae and its
implications for the control of cell division. *J. Cell Biol.* **75**, 422–435 (1977).

16.   D. Killander, A. Zetterberg, Quantitative cytochemical studies on interphase growth. I.
Determination of DNA, RNA and mass content of age determined mouse fibroblasts
in vitro and of intercellular variation in generation time. *Exp. Cell Res.* **38**, 272–284
(1965).

17.   P. Nurse, Genetic control of cell size at cell division in yeast. *Nature* **256**, 547–551
(1975).

18.   I. Soifer, L. Robert, A. Amir, Single-Cell Analysis of Growth in Budding Yeast and
Bacteria Reveals a Common Size Regulation Strategy. *Curr. Biol.* **26**, 356–361
(2016).

19.   E. Vergés, N. Colomina, E. Garí, C. Gallego, M. Aldea, Cyclin Cln3 is retained at the
ER and released by the J chaperone Ydj1 in late G1 to trigger cell cycle entry. *Mol.*
Cell **26**, 649–662 (2007).

20.   H. Wang, L. B. Carey, Y. Cai, H. Wijnen, B. Futcher, Recruitment of Cln3 cyclin to
promoters controls cell cycle entry via histone deacetylase and other targets. *PLoS*
Biol. **7**, e1000189 (2009).

21.   K. M. Schmoller, J. J. Turner, M. Kõivomägi, J. M. Skotheim, Dilution of the cell cycle
inhibitor Whi5 controls budding-yeast cell size. *Nature* **526**, 268–272 (2015).

22.   X. Liu, *et al.*, Reliable cell cycle commitment in budding yeast is ensured by signal
integration. *Elife* **4** (2015).

23. S. Dorsey, *et al.*, G1/S transcription factor copy number is a growth-dependent
determinant of cell cycle commitment in yeast. *Cell Syst.* **6**, 539-554.e11 (2018).

24. F. S. Heldt, R. Lunstone, J. J. Tyson, B. Novák, Dilution and titration of cell-cycle
regulators may control cell size in budding yeast. *PLoS Comput. Biol.* **14**, e1006548
(2018).

25. A. Litsios, *et al.*, Differential scaling between G1 protein production and cell size
dynamics promotes commitment to the cell division cycle in budding yeast. *Nat. Cell
Biol.* **21**, 1382–1392 (2019).

26. Y. Chen, G. Zhao, J. Zahumensky, S. Honey, B. Futcher, Differential Scaling of Gene
Expression with Cell Size May Explain Size Control in Budding Yeast. *Mol. Cell* **78**,
359-370.e6 (2020).

27. Z. N. Demidenko, M. V. Blagosklonny, Growth stimulation leads to cellular
senescence when the cell cycle is blocked. *Cell Cycle* **7**, 3355–3361 (2008).

28. G. E. Neurohr, *et al.*, Excessive Cell Growth Causes Cytoplasm Dilution And
Contributes to Senescence. *Cell* **176**, 1083-1097.e18 (2019).

29. J. Lengefeld, *et al.*, Cell size is a determinant of stem cell potential during aging. *Sci
Adv* **7**, eabk0271 (2021).

30. M. C. Lanz, *et al.*, Increasing cell size remodels the proteome and promotes
senescence. *Mol. Cell* **82**, 3255-3269.e8 (2022).

31. S. Manohar, *et al.*, Genome homeostasis defects drive enlarged cells into
senescence. *Mol. Cell* **83**, 4032-4046.e6 (2023).

32. L. Crozier, *et al.*, CDK4/6 inhibitor-mediated cell overgrowth triggers osmotic and
replication stress to promote senescence. *Mol. Cell* **83**, 4062-4077.e5 (2023).

33. R. Foy, *et al.*, Oncogenic signals prime cancer cells for toxic cell overgrowth during a
G1 cell cycle arrest. *Mol. Cell* **83**, 4047-4061.e6 (2023).

34. S. Manohar, G. E. Neurohr, Too big not to fail: emerging evidence for size-induced
senescence. *FEBS J.* **291**, 2291–2305 (2024).

35. P. Jorgensen, J. L. Nishikawa, B.-J. Breitkreutz, M. Tyers, Systematic Identification of
Pathways That Couple Cell Growth and Division in Yeast. *Science* **297**, 395–400
(2002).

36. J. Zhang, *et al.*, Genomic Scale Mutant Hunt Identifies Cell Size Homeostasis Genes
in S. cerevisiae. *Curr. Biol.* **12**, 1992–2001 (2002).

37. M. Ohanna, *et al.*, Atrophy of S6K1(-/-) skeletal muscle cells reveals distinct mTOR
effectors for cell cycle and size control. *Nat. Cell Biol.* **7**, 286–294 (2005).

38. S. M. I. Goorden, *et al.*, Rheb is essential for murine development. *Mol. Cell. Biol.* **31**,
1672–1678 (2011).

39. C. D. Bell, E. Waizbard, Variability of cell size in primary and metastatic human
breast carcinoma. *Invasion Metastasis* **6**, 11–20 (1986).

40. B. M. Iritani, R. N. Eisenman, c-Myc enhances protein synthesis and cell size during
B lymphocyte development. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 13180–13185 (1999).

41. I. Jones, *et al.*, Characterization of proteome-size scaling by integrative omics
reveals mechanisms of proliferation control in cancer. *Sci. Adv.* **9**, eadd0636 (2023).

42. Asadullah, *et al.*, Combined heterogeneity in cell size and deformability promotes
cancer invasiveness. *J. Cell Sci.* **134**, jcs250225 (2021).

43. T. J. Kawecki, *et al.*, Experimental evolution. *Trends Ecol. Evol.* **27**, 547–560 (2012).

44. M. Natalino, M. Fumasoni, Experimental approaches to study evolutionary cell
biology using yeasts. *Yeast* **40**, 123–133 (2023).

45. J. Helsen, G. Sherlock, G. Dey, Experimental evolution for cell biology. *Trends Cell*
Biol. **33**, 903–912 (2023).

46. Practical Flow Cytometry Shapiro, 4th Edition.

47. Y. H. Jin, *et al.*, The multiple biological roles of the 3'-->5' exonuclease of
Saccharomyces cerevisiae DNA polymerase delta require switching between the
polymerase and exonuclease domains. *Mol. Cell. Biol.* **25**, 461–471 (2005).

48. P. Jorgensen, *et al.*, A dynamic transcriptional network communicates growth
potential to ribosome synthesis and critical cell size. *Genes Dev.* **18**, 2491–2505
(2004).

49. R. Nash, G. Tokiwa, S. Anand, K. Erickson, A. B. Futcher, The WHI1+ gene of
Saccharomyces cerevisiae tethers cell division to cell size and is a cyclin homolog.
EMBO J. **7**, 4335–4346 (1988).

50. R. S. Nash, T. Volpe, B. Futcher, Isolation and characterization of WHI3, a size-
control gene of Saccharomyces cerevisiae. *Genetics* **157**, 1469–1480 (2001).

51. P. Russell, P. Nurse, cdc25+ functions as an inducer in the mitotic control of fission
yeast. *Cell* **45**, 145–153 (1986).

52. K. Lundgren, *et al.*, Mik1 and wee1 cooperate in the inhibitory tyrosine
phosphorylation of cdc2. *Cell* **64**, 1111–1122 (1991).

53. C. M. Chavez, *et al.*, The cell morphological diversity of Saccharomycotina yeasts.
FEMS Yeast Res. **24**, foad055 (2024).

54. D. H. Williamson, A. W. Scopes, The behaviour of nucleic acids in synchronously
dividing cultures of Saccharomyces cerevisiae. *Exp. Cell Res.* **20**, 338–349 (1960).

55. M. Osumi, Visualization of yeast cells by electron microscopy. *J. Electron Microsc.*
61, 343–365 (2012).

56. A. Alsina, M. Fumasoni, P. Sartori, Model-based inference of cell cycle dynamics
captures alterations of the DNA replication programme. *PLoS Comput. Biol.* **21**,
e1013570 (2025).

57. R. E. Lenski, M. R. Rose, S. C. Simpson, S. C. Tadler, Long-Term Experimental
Evolution in Escherichia coli . I . Adaptation and Divergence During. *Am. Nat.* **138**,
1315–1341 (1991).

58. M. Scott, T. Hwa, Bacterial growth laws and their applications. *Curr. Opin.*
Biotechnol. **22**, 559–565 (2011).

59. R. Droghetti, *et al.*, Hands-on growth laws theory cookbook. *PRX Life* **4** (2026).

60. E. Metzl-Raz, *et al.*, Principles of cellular resource allocation revealed by condition-
dependent proteome profiling. *Elife* **6** (2017).

61. A. I. Goranov, *et al.*, The rate of cell growth is governed by cell cycle stage. *Genes*
Dev. **23**, 1408–1422 (2009).

62. T. LaBar, Y. Y. Phoebe Hsieh, M. Fumasoni, A. W. Murray, Evolutionary Repair
Experiments as a Window to the Molecular Diversity of Life. *Curr. Biol.* **30**, R565–
R574 (2020).

63. S. Schrott, C. Osman, Two mitochondrial HMG-box proteins, Cim1 and Abf2,
antagonistically regulate mtDNA copy number in Saccharomyces cerevisiae. *Nucleic*
Acids Res. **51**, 11813–11835 (2023).

64. B. L. A. Carter, P. E. Sudbery, SMALL-SIZED MUTANTS OF SACCHAROMYCES
CEREVISIAE. *Genetics* **96**, 561–566 (1980).

65. M. Tyers, G. Tokiwa, R. Nash, B. Futcher, The Cln3-Cdc28 kinase complex of S.
cerevisiae is regulated by proteolysis and phosphorylation. *EMBO J.* **11**, 1773–1784
(1992).

66. M. Caligaris, *et al.*, Snf1/AMPK fine-tunes TORC1 signaling in response to glucose
starvation. *Elife* **12** (2023).

67. N. Talarek, E. Gueydon, E. Schwob, Homeostatic control of START through negative
feedback between Cln3-Cdk1 and Rim15/Greatwall kinase in budding yeast. *Elife* **6**
(2017).

68. S. Mochida, S. L. Maslen, M. Skehel, T. Hunt, Greatwall phosphorylates an inhibitor
of protein phosphatase 2A that is essential for mitosis. *Science* **330**, 1670–1673
(2010).

69. A. Gharbi-Ayachi, *et al.*, The substrate of Greatwall kinase, Arpp19, controls mitosis
by inhibiting protein phosphatase 2A. *Science* **330**, 1673–1677 (2010).

70. N. Chica, *et al.*, Nutritional control of cell size by the greatwall-endosulfine-PP2A·B55
pathway. *Curr. Biol.* **26**, 319–330 (2016).

71. S. Bontron, *et al.*, Yeast endosulfines control entry into quiescence and chronological
life span by inhibiting protein phosphatase 2A. *Cell Rep.* **3**, 16–22 (2013).

72. A. M. Healy, *et al.*, CDC55, a Saccharomyces cerevisiae gene involved in cellular
morphogenesis: identification, characterization, and homology to the B subunit of
mammalian type 2A protein phosphatase. *Mol. Cell. Biol.* **11**, 5767–5780 (1991).

73. G. S. Darmasaputra, L. M. van Rijnberk, M. Galli, Functional consequences of
somatic polyploidy in development. *Development* **151**, dev202392 (2024).

74. T. R. Gregory, Coincidence, coevolution, or causation? DNA content, cell size, and
the C-value enigma. *Biol. Rev. Camb. Philos. Soc.* **76**, 65–101 (2001).

75. J. S. P. Heslop-Harrison, T. Schwarzacher, Q. Liu, Polyploidy: its consequences and
enabling role in plant diversification and evolution. *Ann. Bot.* **131**, 1–10 (2023).

76. Y. Chadha, A. Khurana, K. M. Schmoller, Eukaryotic cell size regulation and its
implications for cellular function and dysfunction. *Physiol. Rev.* **104**, 1679–1717
(2024).

77. I. Soifer, N. Barkai, Systematic identification of cell size regulators in budding yeast.
Mol. Syst. Biol. **10**, 761 (2014).

78. R. A. M. de Bruin, W. H. McDonald, T. I. Kalashnikova, J. Yates 3rd, C. Wittenberg,
Cln3 activates G1-specific transcription via phosphorylation of the SBF bound
repressor Whi5. *Cell* **117**, 887–898 (2004).

79. Y. Ho, M. Costanzo, L. Moore, R. Kobayashi, B. J. Andrews, Regulation of
transcription at the Saccharomyces cerevisiae start transition by Stb1, a Swi6-
binding protein. *Mol. Cell. Biol.* **19**, 5267–5278 (1999).

80. R. A. M. de Bruin, T. I. Kalashnikova, C. Wittenberg, Stb1 collaborates with other
regulators to modulate the G_1-specific transcriptional circuit. *Mol. Cell. Biol.* **28**,
6919–6928 (2008).

81. R. Verma, *et al.*, Phosphorylation of Sic1p by G1 Cdk required for its degradation
and entry into S phase. *Science* **278**, 455–460 (1997).

82. M. Costanzo, *et al.*, CDK activity antagonizes Whi5, an inhibitor of G1/S transcription
in yeast. *Cell* **117**, 899–913 (2004).

83. F. R. Cross, L. Schroeder, J. M. Bean, Phosphorylation of the Sic1 inhibitor of B-type
cyclins in Saccharomyces cerevisiae is not essential but contributes to cell cycle
robustness. *Genetics* **176**, 1541–1555 (2007).

84. M. A. Juanes, *et al.*, Budding yeast greatwall and endosulfines control activity and
spatial regulation of PP2A(Cdc55) for timely mitotic progression. *PLoS Genet.* **9**,
e1003575 (2013).

85. *Guide to yeast genetics and molecular biology* (Academic Press, 1990).

86. M. Fumasoni, A. W. Murray, The evolutionary plasticity of chromosome metabolism
allows adaptation to constitutive DNA replication stress. *Elife* **9**, 1–28 (2020).

87. M. Natalino, M. Fumasoni, Compensatory evolution to DNA replication stress is
robust to nutrient availability. *Mol. Syst. Biol.* (2025). https://doi.org/10.1038/s44320-
025-00127-z.

88. M. Fumasoni, A. W. Murray, Ploidy and recombination proficiency shape the
evolutionary adaptation to constitutive DNA replication stress. *PLoS Genet.* **17**,
e1009875 (2021).