Experimental evolution of cellular miniaturization reveals a mechanism for cell size evolution

Garoña A.[1], Lemos V. M.[1], Giometto A.[2], Fumasoni M[1#].

[1] Gulbenkian Institute for Molecular Medicine (GIMM), Oeiras, Portugal

[2] Cornell University, School of Civil and Environmental Engineering, Ithaca, USA

[#] To whom correspondence may be addressed: marco.fumasoni@gimm.pt

ABSTRACT

Cell volume, a key determinant of physiology, is maintained by cell size homeostasis. Large deviations from typical size are often harmful, yet cell sizes have diverged drastically in evolution. How does size homeostasis evolve to support such diversity without impairing physiology? To address this, we used experimental evolution to select progressively smaller *Saccharomyces cerevisiae* cells. Over 1,500 generations, we achieved a six-fold volume reduction compared to wild type. Size homeostasis remained robust and evolutionary stable, with cells maintaining competitive fitness, indicating that dramatic volume changes need not compromise physiology. We investigated the genetic basis of cellular miniaturization through whole-genome sequencing of the evolving populations. We show that genetic manipulations of the G1 cyclin *CLN3* and the Greatwall kinase *RIM15* signaling cascades produce a fourteen-fold change in cell size, with adaptive mutations recapitulating the evolved volumes and loss-of-function mutations yielding enlarged cells. Our results demonstrate the evolutionary plasticity of cell size homeostasis and reveal a mechanism for eukaryotic cell size evolution.

Significance statement

Cell size affects nearly every aspect of cell function and varies enormously among eukaryotes.

How such diversity arises while cells remain stable and healthy is not well understood. Using

experimental evolution in budding yeast, we selected for progressively smaller cells and

identified mutations in conserved cellular pathways that enable stable size reduction without

major fitness costs. These results show that gradual evolutionary tuning of core regulatory

networks can support large changes in cell size while preserving normal function, providing an

experimental framework to explore how fundamental cellular traits evolve.

INTRODUCTION

Cells are the fundamental unit of life, and their size is a key determinant of cell and organismal

physiology (1). Through its effects on biosynthesis and proliferation, cell size constrains tissue

architecture and regulates functional specialization (2–4).

Not surprisingly, cell size varies greatly across the tree of life. Among prokaryotes, cell

diameters range from ~0.2 μm in *Mycoplasma* species (5) to nearly 1 cm in *Thiomargarita*

magnifica (6). Eukaryotes exhibit a similarly broad distribution, from the smallest known free-

living species *Ostreococcus tauri,* with a 1–2 μm diameter (7), to 15 cm long ostrich eggs (8).

Even within multicellular organisms, as in humans, cells range from lymphocytes (<10 μm) to

meter-long neurons (9).

Despite this large variability, specific cell types typically maintain their size within a narrow

range through a process defined as cell size homeostasis (10, 11). At its core, cell size

homeostasis relies on the precise coordination of cell growth and division. The most direct

evidence of such control on cell size is that cells born small typically grow more before

dividing, and vice versa (12). While, in theory, size control could be performed throughout the

cell cycle (13), most organisms exert it mainly prior to DNA synthesis, such as in budding yeast

(14) and humans (15), or before nuclear division, such as in fission yeast (16). The general

principles followed by cells to regulate their size, as well as the molecular mechanisms that

execute this control, have been studied and debated for decades. These are best understood

in the budding yeast *S. cerevisiae*, where cells are thought to commit to division once they

reach a critical size (14) or after adding a constant volume each cycle (17). The mechanism

that determines the transition from G1 to S phase has been proposed to depend mainly on

titration of cell cycle activators (e.g. Cln3), inhibitors (e.g. Whi5), transcription factors (e.g.

SBF), or their combination (for detailed mechanisms, see refs. (18–25)).

Strong evidence for the importance of cell size homeostasis is represented by the severe

defects experienced by cells as they leave their narrow range of stereotypical sizes. Excessive

cell growth has been shown to cause cytoplasm dilution and genetic instability, ultimately

increasing cells' propensity to senescence (26–33). On the other hand, mutations causing the

largest decrease in cell size impair the Target of Rapamycin (TOR) signaling and ribosome

biogenesis, imposing severe reductions in cellular biosynthesis and fitness (34–37). Notably,

altered cell size control has been observed in cancers (38–41).

It is therefore unclear how evolution has achieved changes in cell size of many orders of

magnitude, when even slight increases or decreases are associated with severe cellular

defects. These detrimental consequences could severely constrain the evolution of the cell

size trait, posing an apparent paradox between the broad evolutionary diversity of cell sizes

and the limited tolerance observed within species. To address this paradox, we sought to

observe in real time the evolution of a large decrease in cell size in the model eukaryote *S.*

cerevisiae. To this goal, we resorted to an experimental evolution regime (42–44),

simultaneously imposing direct selection of cellular size by Fluorescent-Activated Cell

Sorting (FACS), followed by selection for competitive fitness. Applying this regime over 1,500

generations produced miniaturized cells with six-fold reduction over wild type (WT) volumes

and bearing only marginal fitness tradeoffs. Miniaturized cells maintained their phenotype in

the absence of selection and had tight control over size distribution at the population level.

Whole-genome sequencing revealed adaptive mutations in the *CLN3*, *SCH9,* and *RIM15*

genes. Collectively, these mutations decouple the tight link between growth rate and cell size

by altering the G1 cyclin and Greatwall signaling pathway. Targeted engineering of these

pathways rewired cell size control and cell cycle dynamics, yielding a ten-fold range in cell

volume. Our work demonstrates the plasticity of the mechanisms governing cell size

homeostasis and suggests an effective mechanism for cell size evolution.

RESULTS

To investigate how cell size can evolve under sustained proliferative capacity, we developed

an experimental system in *S. cerevisiae* alternating two selective regimes targeting cell size

and reproductive fitness, respectively. Cell populations were analyzed by Fluorescence

Activated Cell Sorting (FACS), using the light scattered by cells (forward scatter area, FSC-A),

a widely used proxy for their size (45). Only the smaller 7% of cells were collected into separate

tubes and subjected to a second selection step (Figure 1A, see Materials and Methods and

Figure S1A for details). The second selection consisted of the unconstrained growth of the

sorted population for at least 11 generations, during which cells competed for resources

(Figure 1A). Three parallel small (S1-3) populations were subjected daily to the two selection

regimes, for a total of 1,500 generations (Figure 1B). Under these conditions, selection favors

mutations that reduce cell size without compromising proliferation. To speed the evolution,

the starting haploid population (ancestor) carried a mutator allele in the catalytic subunit of

DNA polymerase delta (*pol3-L523D*, (46) Table S1). In addition, we engineered the ancestor

strain to express a cytoplasmic fluorescent protein (ymCitrine) to prevent subcellular debris

from being sorted.

Over 1,500 generations, the evolving populations (S1–3) showed a gradual decrease in FCS

values as measured by flow cytometry (Figure S1B). Microscopy and Coulter counter analyses

confirmed that this reduction was accompanied by a marked decrease in cell volume (Figure

1C–D). Literature values from Coulter counter measurements are often reported from log-

phase cultures, where the prevalence of daughter cells skews the volume distribution toward

smaller sizes. To avoid biases from growth phase and cell cycle differences, we measured

volumes at saturation (48 h), which normalizes these effects and yields consistently higher

values (Figure S1C). Relative to WT cells (101 fL ± 17 (mean ± SD)), S-populations at 1,500

generations displayed on average a six-fold decrease, reaching 14 fL ± 1.5 (mean ± SD, Figure

1C), a volume closer to the bacterium *B. subtilis* than their own ancestor. Comparatively,

perturbations of the TOR signaling pathway (*sch9Δ* and *sfp1Δ*), which produced the smallest

previously known yeast cells (34, 47), only reduced volumes by threefold (30 fL ± 1.7 and 28 fL

± 2.8, respectively (mean ± SD)) (Figure 1C). Our selection experiment therefore produced the

smallest *S. cerevisiae* cells reported to date, representing, to our knowledge, the largest

reduction in cell volume observed in proliferating cells, excluding cases associated with

developmental programs and ploidy changes. Comparable reductions in *S. pombe* have been

reported to cause mitotic catastrophe (48, 49). Strikingly, the evolved cells reached volumes

similar to those of the smallest species within the *Saccharomycotina* subphylum (Figure

S1D), spanning nearly half of the natural size diversity observed across the phylum despite

over 400 million years of divergence (50).

Figure 1. Experimental evolution of cellular miniaturization. (A and B) Schematic representation of size selection and

evolution regimes. A haploid *pol3-L523D* mutator ancestor population was sorted into three small (S) subpopulations and

subjected to daily size selection. Cells in the lowest 7% of the forward-scatter distribution were sorted (100,000 cells) and cultured

in 10 mL rich medium (YPD). Populations were serially grown and sorted for 116 cycles (~1,500 generations), with samples

collected and stored every 24 h. (C) Cell volume measurements (Coulter counter) across generations. Data from the three S-

populations are plotted together, showing progressive evolution toward smaller size, reaching up to a sixfold reduction in volume.

Grey lines indicate wild-type and final evolved sizes. Box plots show medians, interquartile ranges, whiskers (1.5× IQR), and

outliers. (D) Light micrographs of wild-type and evolved small-cell populations at different cell-cycle stages (G1, S, G2/M). Scale

bar, 5 µm.

Gradual cell size decrease leads to altered morphology and cell cycle dynamics, with marginal fitness tradeoffs.

The dramatic changes in cell volumes observed during our evolution prompted us to examine

potential alterations in cellular architecture. First, we focused on intracellular organization,

reasoning that reduced volume might occur at the expense of specific organelles. To obtain a

general view of subcellular structures, we performed transmission electron microscopy (51)

on exponentially growing WT, ancestral, and evolved S-populations. Buds smaller than

mother cells, consistent with asymmetric cell division (14), were commonly observed.

Interestingly, major organelles displayed their typical morphology, though with altered

proportions, suggesting a potential differential miniaturization of organelles (Figure 2A and

Figure S2A). Despite the presence of petite mutants (cells lacking functional mitochondrial

respiration), most evolved isolates, including the smallest, retained growth on non-

fermentable carbon sources, indicating preserved mitochondrial function (Figure S2B).

Cell cycle is a major determinant of cell morphology in budding yeast (52). To determine

whether changes in cell size were associated with alterations in the cell cycle, we analyzed

DNA content in exponentially growing populations, where signal intensity distinguishes cell

cycle phases (1C for G1, 2C for G2/M, and 1C < DNA < 2C for S phase). Strikingly, by generation

1,500, evolved small cells displayed a near absence of G1 compared with WT (Figure 2B and

S2C). Notably, cell cycle changes during evolution occurred in two steps: during the first ~150

generations, the proportion of cells in G1 increased, followed by a sharp shift around

generation 700 that resulted in the near disappearance of G1 (Figure 2C and S2D). Calculating

the absolute time spent in each phase confirmed this trend, showing that the reduction of G1

duration was primarily compensated by a corresponding extension of G2/M (Figure 2C-D and

S2D).

We next examined how changes in cell size and cycle affected population growth. Growth

curves of WT, ancestral, and S-populations revealed a marked increase in maximum

population density of the evolved lines (Figure 2E). Their growth rates were only marginally

reduced relative to ancestors, which themselves grew slower than WT, potentially due to the

mutator phenotype (Figure 2F). To precisely estimate this defect, we resorted to pairwise

competition assays, which measure relative fitness, a measure of the reproductive success

of organisms (53). All three evolved S-populations had equivalent relative fitness (ω = –7,79%

± 1.4, mean ± SD vs. ancestor), whereas WT showed a ~5% advantage relative to the ancestor

(Figure 2G). In contrast, deletion of *SCH9*, which produces the smallest cells previously

reported, caused a severe fitness defect (–28.76% ± 1.1, mean ± SD) despite reducing cell

volume only threefold. Across all populations, fitness followed a non-uniform trajectory: (i)

declining during the first 200 generations, (ii) recovering by ~250, (iii) dropping sharply until

~500, and (iv) stabilizing with minor decreases until 1,500 (Figure 2H, S2E). Fitness was more

strongly associated with cell-cycle dynamics than with cell volume (Pearson r = 0.67, p < 0.001

for G1 fraction; r = 0.30, p = 0.11 for cell volume), indicating that changes in fitness primarily

reflect alterations in cell-cycle progression (Figure 2I, S2F).

Thus, cellular miniaturization was accompanied by significant changes in cell cycle dynamics

while maintaining a substantially higher fitness than previously known small-size mutants.

Figure 2. Physiology of cell miniaturization. (A) Transmission electron micrograph of evolved *S. cerevisiae* cells during

exponential growth in YPD. Scale bar, 1 μm. (B) DNA-content profiles of ancestral and representative evolved (S1) populations

during exponential growth across 1,500 generations, measured by flow cytometry. Lines show means; shaded areas, SD (n = 3).

G2 peak height was normalized to 1. Arrows indicate time points of cell-cycle profile changes (see main text). (C) Percentage of

doubling time spent in G1 and S/G2/M phases for the S1 population at generation 1,500. Bars show mean ± SD (n = 3). (D)

Absolute time spent in each cell cycle phase, calculated as the fraction of the doubling time allocated to each phase. (E) Growth

curves of selected strains measured as cell density (cells/mL). (F) Maximum growth rates of reference and evolved strains

(generation 1,500) in YPD. Box plots show medians, interquartile ranges, whiskers (1.5× IQR), and outliers. Statistical significance

assessed by the Mann–Whitney–Wilcoxon test; p-values indicated. Differences among evolved lines (S1–S3) were not significant

(adjusted p = 0.082–0.143), whereas growth rates differed significantly between wild type and evolved populations (p = 0.006).

The ancestor did not differ from S1 (p = 0.65) but differed from S2 and S3 (p = 0.006; n = 4 each). Full pairwise statistics in

RawData-Figures Table Fig2F. (G) Relative fitness of wild-type, ancestral, and evolved populations (S1–S3) at generation 1,500.

Relative fitness is the ratio of reproductive success of a strain to a reference strain under identical conditions. Fitness of evolved

and *sch9Δ* strains differed significantly from the ancestor and wild type (adjusted p < 0.01), with no significant differences among

evolved lines. Box plots as in (F). Full pairwise statistics in RawData-Figures Table Fig2G. (H and I) Fitness and cell-size

trajectories across 1,500 generations for the S1 population. Lines show mean values; shaded areas, SD.

New evolved size homeostasis is stable and robust

How stable is the evolved miniaturized size? Measurements performed right after the last sorting

round may reflect a transient physiological state, rather than an evolutionarily stable phenotype.

To address this possibility, clones were isolated from two independent populations (S1Cl2 and

S2Cl4) and passaged for 150 generations in the absence of any size selection. Interestingly, S-

clones stably maintained their size across the experiment, resulting in a smaller final fold change

than the one observed in control WT lines (Figure 3A). In contrast, *sch9Δ* lines, consistently

reverted to WT sizes between 50 and 100 generations despite starting with a larger volume than

S-clones (Figure 3A). An important question is whether miniaturized cells actively regulate their

size, or whether they merely preserve the volumes attained at the end of the experiment. To

address this, we transiently arrested cells at the G1/S transition using the pheromone α-factor,

which permits growth while blocking division (54) (Figure S3A and S3B). While the arrest was not

fully efficient in S-clones, it nevertheless led to a ~1.8-fold increase in cell volume in both WT and

evolved cells. The increased volume, however, was progressively reversed upon release into

fresh medium, indicating that cells actively return to their characteristic size (Figure S3A, S3C).

The ability of miniaturized cells to maintain their size across many generations and under

transient perturbations reflects efficient size homeostasis, which is expected to manifest as a

narrow distribution of cell sizes around the population mean. To test this prediction, we

compared the size distributions of WT, ancestral strains, S-clones, and *sch9Δ* controls. Cell size

distributions are described by a universal statistical structure (55), by which the distribution of

rescaled cell sizes $V = v/\langle v \rangle$ is invariant across strains and species. Here, $\langle v \rangle$ denotes the mean

size of a strain. This invariance leads to a "collapse" of rescaled cell size distributions onto a

single curve (Figure 3B and 3C) and implies that the coefficient of variation (CV) of the original

(non-rescaled) distributions is constant. Such rescaling enables meaningful comparison of size

homeostasis across strains with markedly different sizes. *sch9Δ* controls showed a poor

collapse of the rescaled distributions compared to WT (Figure 3C), accompanied by an increased

CV ($p < 0.001$, Figure 3D), consistent with greater variability than expected for their mean size.

By contrast, S-clones exhibited collapse and CV values close to WT (SI Appendix), leading to

overlapping rescaled distributions with WT and ancestral strains (Figures 3B–D).

Together, these results demonstrate that miniaturized cells have evolved a new cell size

homeostasis that is evolutionarily stable, robust to transient perturbations, and capable of

maintaining population-level size control as tightly as WT.

Figure 3. Evolution of cell size homeostasis. (A) Stability of cell size during experimental evolution. Left: cell-size measurements of WT, ancestor, *sch9Δ*, and evolved clones at 1,500 generations from populations evolved without size selection. No significant differences were detected between S1Cl2 and S2Cl4 replicates (Mann–Whitney U test, p > 0.1), indicating stable size maintenance. Right: fold change in cell size relative to time 0. (B) Distributions of cell volumes for the analyzed strains before rescaling. Leftmost peaks correspond to debris. (C) Collapse of rescaled cell-volume distributions around their mean, estimated using truncated log-normal fits applied to data above the mode to exclude debris. The dashed black curve shows the best log-normal fit to the rescaled, debris-corrected WT distribution (see Materials and Methods). (D) Coefficient of variation of the collapsed volume distributions, quantifying size variability after rescaling and revealing differences in size homeostasis among genotypes.

Genetic landscape of cellular miniaturization

We sought to determine the genetic basis of the evolved phenotype. Evolved cells retained their characteristic size when co-cultured with wild-type strains, indicating that miniaturization was not driven by secreted or diffusible factors (Figure S4A). The phenotype's persistence over 150 generations in the absence of selection (Figure 3A) strongly suggested a genetic basis. Supporting this idea, crosses between evolved clones and WT produced

diploids of intermediate size between haploid and diploid WT, consistent with the presence

of both recessive and dominant mutations. Furthermore, crosses of evolved clones with

derivatives whose mating type had been switched yielded diploids that maintained a 6-fold

volume reduction relative to WT diploids (Figure S4B), suggesting that the reduction of cell size

is independent of ploidy state.

To identify causative mutations, we performed whole-genome, whole-population sequencing

across multiple evolutionary timepoints. As expected from the mutator allele present in the

ancestral population, we detected 869 unique mutations that arose during evolution in coding

and regulatory regions across the three populations that were not present in the ancestor

(Figure 4A, Table S2). By the end of the experiment, an average of 267 single-nucleotide

polymorphisms (SNPs) and small insertions/deletions (indels) persisted per population (285

in S1, 301 in S2 and 216 in S3). No aneuploidies, segmental amplifications, or large-scale

changes in DNA copy number were observed, thus confirming the presence of a full haploid

genome (Figure S4D). The evolved populations exhibited distinct clonal compositions (Figure

S4C–D). However, several alleles were shared across populations, indicating that cross-

contamination likely occurred during the FACS selection steps. Consequently, the evolved

lines were treated as a single, large evolving population for subsequent analyses (Figure 4A).

To identify putative adaptive mutations associated with the evolved cell size, we employed

four non exhaustive, but complementary approaches: (i) tracking lineages that rose to fixation

in any of the three sub-populations, identifying cohorts of mutations under positive selection

(Table S2); (ii) searching for genes previously implicated in cell size regulation (Table S3); (iii)

using statistical inference to identify genes mutated more frequently than expected given their

size and the observed mutation rates (parallel evolution; Table S4); and (iv) performing two

consecutive rounds of backcrossing and selection for small cell size to pinpoint mutations

segregating with the phenotype (bulk segregant analysis; Table S5; see Materials and Methods

for details). A summary of the genes identified by these methods, as well as their reported

interactions, is portrayed in Figure 4B-C (Table S6). Despite several genetic and physical

interactions among the identified genes, no gene ontology term was found to be statistically

enriched.

Our analysis indicates that the miniaturized phenotype is heritable and genetically encoded,

but the underlying adaptive landscape is complex. The large number of candidate mutations

likely includes false positives arising from hitchhiking or linkage, as well as variants under

indirect selection. Some of these may reflect selection on traits influencing the flow-

cytometry size proxy (FSC-A) rather than cell size itself, or compensatory adaptations to

physiological changes associated with miniaturization (56).

Figure 4. Genomic landscape of the evolved phenotype. (A) Allele-frequency trajectories of mutations detected in the combined population (S1–S3). A 300% frequency indicates fixation in all three populations. Line thickness reflects the final frequency of the mutation at generation 1,500. (B) List of putative adaptive gene mutations ranked by the number of satisfied criteria outlined in the main text and affected by non-synonymous mutations located within coding sequences (CDS). Only genes appearing in at least three categories are included (full list in Table S6). (C) Interaction network of mutations detected in S1–S3 evolved populations curated in Cytoscape (52). Grey lines indicate known genetic and physical interactions (STRING database (53)). Node diameter reflects the number of satisfied criteria we used to detect putative adaptive mutations. Only genes appearing in at least two categories are included. Nodes are color-coded: dark yellow for non-synonymous mutations within CDS fixed at generation 1,500, light yellow for mutations in the regulatory regions fixed at generation 1,500, and grey for non-fixed mutations. Bold outlines mark genes linked to cell size changes in the literature (either decrease or increase).

Combined mutations in the G1 cyclin and Greatwall signalling cascades cause large deviations in cell size

We set out to identify the mutations primarily responsible for the change in cell size among those identified as putatively adaptive. *CIM1*, *CLN3*, and *SCH9* ranked highest in our analysis and were therefore selected for further investigation. *CIM1* encodes for a mitochondrial HMG-box protein that limits mitochondrial DNA copy numbers (57), and independently accumulated up to 6 mutations in the protein over the course of the experiment, including three frameshifts resulting in premature stop codons (Figure S5A). Cln3, a G1 cyclin that partners with Cdk1 to drive the G1/S transition by activating phase-specific gene expression, has been shown to reduce cell size when truncated at its C-terminus (e.g. *cln3-1*), due to increased protein stability (58, 59). Our evolved strains carried a similar truncation, *cln3-S410**, just downstream of the *cln3-1* allele (Figure 5A). Sch9 is a key effector of the TORC1 pathway, which integrates nutrient sensing with anabolic and catabolic processes. We identified a mutation (*sch9*-L343S) within the conserved C2 domain (residues 184–402), a calcium-dependent membrane-targeting module that may also inhibit the kinase domain (residues 403–738; Figure 5A, (60)). The influence of Sch9 on cell size has traditionally been attributed to its role in ribosome biogenesis (47). However, recent work suggests an additional function in regulating cell size through the Greatwall kinase Rim15, albeit under conditions of low CDK activity (61). The Greatwall kinase cascade is highly conserved among eukaryotes and regulates the cell cycle in humans and *Xenopus* via endosulfines-mediated inhibition of the phosphatase PP2A (62, 63). Notably, whereas in *S. pombe* this pathway links nutrient sensing to cell size (64), in *S. cerevisiae* it is thought to function mainly in the regulation of quiescence (65). Interestingly, we also identified a fixed mutation in *RIM15* (*rim15-A132T*) and another in *CDC55* (*cdc55-L159P*), which encodes a regulatory subunit of PP2A (66), though at a lower frequency.

We tested the causality of the *cim1*, *cln3*, *sch9* and *rim15* mutant alleles by reconstructing

them in a WT background. We mimicked frameshift mutations in *CIM1* with a gene deletion,

which did not cause any change in cell size (Figure S5A), perhaps reflective of an adaptive role

in other cellular features or fitness. Mutations in *CLN3*, *SCH9*, and *RIM15* individually caused

a marginal but significant decrease in volumes compared to WT (Figure 5C-D). Combinations

of the three mutations lead to additive and synergistic effects, with the triple mutant

producing a three-fold reduction in volume (28.5 fL, Figure 5C). Although *sch9-L343S cln3-*

*S410** double mutant already manifested a small cell size, the additional *rim15-A132T* allele

marginally decreased the volume (p= 3.2587e-03, Figure 5C and S5B). To further test the

involvement of Rim15 in the final evolved size, we engineered a tunable promoter (*prCUP1*)

driving the *RIM15* and *rim15-A132T* alleles. Copper titration confirmed Rim15 dosage

sensitivity: basal expression reduced size, and overexpression caused further shrinkage in

both WT and *cln3-S410** backgrounds (Figure S5C). Notably, overexpressing *rim15-A132T* in

the *sch9-L343S cln3-S410** background decreased size even further, matching that of the final

evolved clones (16 fL ± 0.1 (mean ± SD), Figure 5C and S5C). These results suggest that, in

addition to a gain of function of the G1 cyclin Cln3, a gain of function of the Greatwall signaling

cascade may be responsible for the miniaturized volume. Consistent with this, deleting *RIM15*

in evolved clones restored larger volumes (Figure S5D). Importantly, despite their small size,

all strains reconstructed with evolved alleles retained near-WT fitness (Figure 5C). The

engineered *prCUP1-rim15-A132T* in the *sch9-L343S cln3-S410** background resulted in lower

fitness than the evolved clones but higher than *sch9Δ*. However, these fitness levels likely

reflect the toxic effects of deregulated *Rim15* expression throughout the cell cycle rather than

reduced cell size, as fitness remained constant while size decreased at higher copper

concentrations (Figure 5C).

Finally, following the hypothesis that the combined effect of the reconstructed mutations

results in a gain of function in the G1 cyclin and Greatwall signaling cascades, we asked

whether removing Cln3 and Rim15 would enlarge cell size. Indeed, simultaneous deletion of

CLN3 and *RIM15* produced a synergistic ~2.3-fold volume increase (Figure 5E). Altogether,

combining alleles that modulate G1 cyclin activity and Greatwall signalling generated a

fourteen-fold size range, comparable to 82.86% of the variation in sizes observed across the

entire *Saccharomycotina* subphylum (Figure 5E).

Figure 5. Genetic basis of cell size phenotype. (A) Schematic of cell size–regulating genes. Protein domains are indicated, and mutations identified during evolution are marked: PK, kinase catalytic domain; D, consensus disorder prediction; PAS, N-terminal PAS; C2HC, CCHC-type zinc-finger domains; REC, C-terminal receiver domain. (B) Allele-frequency trajectories of size-associated mutations. Only variants reaching ≥80% frequency by generation 1,500 are shown; causative mutations are highlighted. The right axis shows the S1 population volume (× symbols) over time. (C) Relationship between relative fitness and modal cell volume. Each point represents a strain (WT, ancestor, evolved populations, or engineered mutants). Fitness (x-axis) and cell volume (y-axis); error bars indicate SD. (D) Coefficient of variation of rescaled volume distributions, quantifying size variability and differences in size homeostasis among genotypes. (E) Distribution of cell volumes across the *Saccharomycotina* subphylum (derived from (50)). Highlighted are the triple mutant (*sch9-L343S cln3-S410* prCUP1-rim15-A132T*) under a copper-inducible promoter and the *CLN3* and *RIM15* double deletions, showing shifts in size relative to natural variation.

DISCUSSION

How can evolution produce large shifts in cell size without incurring in the detrimental effects associated with even small deviations from the norm? In development, abrupt changes often arise through ploidy increase or reduction, and cell volume scales linearly with DNA content across taxa (67). Ploidy increases are important in plant evolution and likely account for part of the variation in cell size among related species (68). Yet, genome duplications alone cannot explain the wide diversity of cell sizes, as they also create barriers to sexual reproduction and would leave more pervasive genomic signatures if they were the main driver. Cells must therefore be able to modify size independently of DNA content, at least initially, especially under gradual selection for divergent sizes.

Decades of studies on eukaryotic cell cycle regulation have identified the main principles allowing cells to regulate their size, as well as many molecular players involved (reviewed in (10, 69). Mutations in many such players have been demonstrated to produce altered sizes (34, 35, 70). It is therefore intuitive to imagine that evolution could modulate cell size by fine-tuning these molecular players. Yet most such mutations reduce fitness, and it remains unclear which, if any, combinations of the dozens of known regulators could provide viable evolutionary paths to new cell sizes.

We addressed this question experimentally by imposing selection on *S. cerevisiae* cell size

and fitness. Over 1,500 generations, cells shrank six-fold in volume without changes in ploidy

and with minimal fitness costs. We show that miniaturized cells maintain an efficient and

evolutionarily stable cell size homeostasis, despite a significantly altered cell cycle

progression characterized by the near absence of a G1 phase. Whole genome sequencing

revealed a complex genomic landscape, due to the mutator allele employed, and likely linked

to the many direct and indirect changes cells experienced during the miniaturization.

Nevertheless, re-engineering putative adaptive alleles showed that simultaneous

perturbations of the G1 cyclin *CLN3* and the Greatwall kinase *RIM15* pathways recapitulate

the miniaturized phenotype, whereas loss-of-function in the same genes drives cell

enlargement. Together, manipulations in these signaling cascades produced a fourteen-fold

size range within a constant genome, capturing up to 82.86% (261 / 315) of the natural

variation in size seen across the *Saccharomycotina* over 400 million years of evolution.

Three questions arise. First, how can our evolved cells reach sizes smaller than those

achieved previously through gene deletions? We propose that evolution selected alleles with

modest individual effects, but that act synergistically on cell size by working on multiple

mechanisms simultaneously. We speculate the following model to account for the phenotype

observed (Figure 6): Truncations in *CLN3* stabilize the protein, increasing *CDK1–Cln3* activity

and phosphorylation of the transcriptional inhibitor Whi5 (71). In parallel, gain-of-function

mutations in the Greatwall kinase Rim15 enhance Igo1/2-mediated inhibition of the

phosphatase PP2A–Cdc55, further driving Whi5 phosphorylation and nuclear exclusion (61).

But why would this lead to a volume smaller than *whi5Δ* (Figure S5E), where the inhibitor is

absent? First, CDK1-Cln3 levels could simultaneously hyper-phosphorylate the

transcriptional regulator Stb1 (19, 72, 73) and accelerate the degradation of the CDK1

inhibitor Sic1 (74–76), eliminating the G1 phase. Furthermore, *RIM15* activity in mitosis has

been reported to destabilize the CDK1 inhibitor and size/morphogenesis checkpoint Swe1

(77), offering an additional route to reduced volume.

The second question regards what allows these miniaturized cells to retain high fitness.

Eukaryotic cells physiologically shrink upon nutrient scarcity (12), where cells may find it

advantageous to increase population-wide survival by dividing at a smaller volume while making

a parsimonious use of the resources left by reducing ribosome biogenesis. We propose that the

deletion of *SCH9* hard-wires this response, resulting in severe fitness defects due to strongly

decreased protein production. We speculate that the allele we obtained is a separation-of-

function allele, which interferes with Sch9's ability to inhibit the Greatwall kinase, but allows the

signaling that promotes ribosome biogenesis.

Finally, our miniaturized cells spend virtually no time in G1, yet they maintain robust size

homeostasis at the population level. The G1 duration is thought to buffer stochastic differences

in birth size (14, 17). Does the G1/S transition still perform this function on such short timescales,

or has size control shifted to a cryptic checkpoint in another phase, such as G2/M, as observed

in *S. pombe?*

In summary, our work shows how cell cycle regulation alone can achieve large deviations in cell

size while maintaining homeostasis. It also identifies a simple yet powerful route for evolution to

simultaneously modulate the G1 cyclin and Greatwall kinase pathways synergistically to control

the time spent in G1/G2, and thus cells' ability to grow in interphase. Finally, by coordinating

multiple points of size control while preserving biosynthetic capacity, cells can achieve large

deviations in volume without altering ploidy. Future studies should test the generality of the

principles we identified and determine how broadly it underpins size diversity across the tree of

life.

Overall, our work illustrates how long-term evolution, quantitative physiology, and molecular

genetics, together, can inform the mechanisms underlying the evolution of cellular traits (78).

The generation of an unprecedented range of cell sizes within one species, with proliferating

cells and stable ploidy, provides a useful framework for future work on subcellular scaling and

the limits of cell size.

Figure 6. Speculative model of the effect of the mutations identified on cell cycle regulation. In wild-type (WT) cells

growing under nutrient-rich conditions, TOR is active, promoting ribosome biogenesis and biomass accumulation (right, green

rectangle) while delaying cell cycle transitions (left, orange rectangle). This results in fast-growing, large cells. Under nutrient

limitation or in TOR-related mutants, Sch9 repression of Rim15 is relieved, leading to earlier cell cycle transitions. However,

the accompanying reduction in biomass production limits growth and fitness. We propose that the combined effect of the

identified mutations accelerates cell cycle transitions while maintaining ribosome biogenesis and biomass production. As a

result, cells grow rapidly but, by spending less time in interphase, remain smaller than their WT counterparts.

MATERIAL AND METHODS

All strains used in this study were W303 derivatives. Yeast cultures were propagated using

standard techniques (79). Unless otherwise stated, experiments were performed in standard rich

medium (YPD). Experimental evolution followed the procedure described in (80), with the

addition of the size-selection step by sorting detailed in the SI Appendix. Growth-curve

measurements and fitness assays were performed as described in (81). Samples for cell-cycle

analyses were processed as (81) and analyzed as in (82). Whole-genome sequencing and

analysis followed procedures described in (83). A full description of all methods is provided in

the SI Appendix, Methods.

Data Availability

The genomic datasets used within this publication were deposited at European Nucleotide

Archive (PRJEB101156). Scripts used for data analysis are available at the GitHub repository

(https://github.com/FumaLab/Garona2025). The raw data corresponding to each figure are

provided in the Raw Data–Figures Table. Expanded view data, supplementary information, and

appendices are available for this paper.

Authors contribution

M.F. conceived the project. A.Ga. and M.F. designed the research, performed the experiments,

and analyzed the data. A.Gi. contributed data analysis for Figure 3B–D. M.V.L. contributed

experimental work for Figure 5C, E. A.Ga. and M.F. wrote the original manuscript draft. All authors

edited and commented on the final version of the manuscript.

Acknowledgments

We thank Mina Abrantes, Joana Boas, Filipa Cantanhede, Olga Carreira, and João Martins for

their assistance in the experimental work; all the members of the Genome Maintenance and

Evolution lab for daily support helpful discussions; Gabriel Neurohr, Andrew W. Murray, Paula

Gonçalves, Partha Pratim Chakraborty and Mariana Natalino for critical reading of the

manuscript. The Flow Cytometry, Electron Microscopy, Bioimaging, and Genomics Facility of the

Gulbenkian Institute for Molecular Medicine for their technical support. We also thank Beatriz

Tomaz, Erin Tranfield and Muriel Christel Mari for their help in the establishment of the electron

microscopy sample preparation. The authors acknowledge support from the Gulbenkian

Foundation and the Gulbenkian Institute for Molecular Medicine (A.Ga., M.V.L and M.F), the

Human Frontier Science Program (HFSP) -RGEC28/2023 (M.F and A.Gi), European Molecular

Biology Organization (EMBO) - IG 5349-2023 (M.F), Fundação para a Ciência e a Tecnologia (FCT)

- 2023.09068.CEECIND (M.F) and the National Institute of General Medical Sciences (NIGMS) of

the National Institutes of Health (NIH), United States of America, 1R35GM147493 (A.Gi).

BIBLIOGRAPHY

1. M. B. Ginzberg, R. Kafri, M. Kirschner, Cell biology. On being the right (cell) size. *Science*
348, 1245075 (2015).

2. C. Cadart, R. Heald, Scaling of biosynthesis and metabolism with cell size. *Mol. Biol.*
Cell **33** (2022).

3. T. P. Miettinen, M. J. Caldez, P. Kaldis, M. Björklund, Cell size control - a mechanism for
maintaining fitness and function. *Bioessays* **39**, 1700058 (2017).

4. T. L. Orr-Weaver, When bigger is better: the role of polyploidy in organogenesis. *Trends*
Genet. **31**, 307–315 (2015).

5. M. G. Gabridge, Current status of the mycoplasmas: The smallest living cells. *Lab. Med.*
9, 36–40 (1978).

6. J.-M. Volland, *et al.*, A centimeter-long bacterium with DNA contained in metabolically
active, membrane-bound organelles. *Science* **376**, 1453–1458 (2022).

7. M.-J. Chrétiennot-Dinet, *et al.*, A new marine picoeucaryote: Ostreococcus tauri gen. et
sp. nov. (Chlorophyta, Prasinophyceae). *Phycologia* **34**, 285–292 (1995).

8. C. Di Meo, G. Stanco, M. I. Cutrignelli, S. Castaldo, A. Nizza, Physical and chemical
quality of ostrich eggs during the laying season. *Br. Poult. Sci.* **44**, 386–390 (2003).

9. R. Milo, R. Philips, Cell Biology by the Numbers. Available at:
https://book.bionumbers.org/ [Accessed 23 October 2025].

10. A. A. Amodeo, J. M. Skotheim, Cell-Size Control. *Cold Spring Harb. Perspect. Biol.* **8**,
a019083 (2016).

11. N. Rhind, Cell-size control. *Curr. Biol.* **31**, R1414–R1420 (2021).

12. G. C. Johnston, J. R. Pringle, L. H. Hartwell, Coordination of growth with cell division in
the yeast Saccharomyces cerevisiae. *Exp. Cell Res.* **105**, 79–98 (1977).

13. X. Liu, J. Yan, M. W. Kirschner, Cell size homeostasis is tightly controlled throughout the
cell cycle. *PLoS Biol.* **22**, e3002453 (2024).

14. L. H. Hartwell, M. W. Unger, Unequal division in Saccharomyces cerevisiae and its
implications for the control of cell division. *J. Cell Biol.* **75**, 422–435 (1977).

15. D. Killander, A. Zetterberg, Quantitative cytochemical studies on interphase growth. I.
Determination of DNA, RNA and mass content of age determined mouse fibroblasts in
vitro and of intercellular variation in generation time. *Exp. Cell Res.* **38**, 272–284 (1965).

16. P. Nurse, Genetic control of cell size at cell division in yeast. *Nature* **256**, 547–551
(1975).

17. I. Soifer, L. Robert, A. Amir, Single-Cell Analysis of Growth in Budding Yeast and Bacteria
Reveals a Common Size Regulation Strategy. *Curr. Biol.* **26**, 356–361 (2016).

18. E. Vergés, N. Colomina, E. Garí, C. Gallego, M. Aldea, Cyclin Cln3 is retained at the ER
and released by the J chaperone Ydj1 in late G1 to trigger cell cycle entry. *Mol. Cell* **26**,
649–662 (2007).

19. H. Wang, L. B. Carey, Y. Cai, H. Wijnen, B. Futcher, Recruitment of Cln3 cyclin to
promoters controls cell cycle entry via histone deacetylase and other targets. *PLoS Biol.*
7, e1000189 (2009).

20. K. M. Schmoller, J. J. Turner, M. Kõivomägi, J. M. Skotheim, Dilution of the cell cycle
inhibitor Whi5 controls budding-yeast cell size. *Nature* **526**, 268–272 (2015).

21. X. Liu, *et al.*, Reliable cell cycle commitment in budding yeast is ensured by signal
integration. *Elife* **4** (2015).

22. S. Dorsey, *et al.*, G1/S transcription factor copy number is a growth-dependent
determinant of cell cycle commitment in yeast. *Cell Syst.* **6**, 539-554.e11 (2018).

23. F. S. Heldt, R. Lunstone, J. J. Tyson, B. Novák, Dilution and titration of cell-cycle
regulators may control cell size in budding yeast. *PLoS Comput. Biol.* **14**, e1006548
(2018).

24. A. Litsios, *et al.*, Differential scaling between G1 protein production and cell size
dynamics promotes commitment to the cell division cycle in budding yeast. *Nat. Cell
Biol.* **21**, 1382–1392 (2019).

25. Y. Chen, G. Zhao, J. Zahumensky, S. Honey, B. Futcher, Differential Scaling of Gene
Expression with Cell Size May Explain Size Control in Budding Yeast. *Mol. Cell* **78**, 359-
370.e6 (2020).

26. Z. N. Demidenko, M. V. Blagosklonny, Growth stimulation leads to cellular senescence
when the cell cycle is blocked. *Cell Cycle* **7**, 3355–3361 (2008).

27. G. E. Neurohr, *et al.*, Excessive Cell Growth Causes Cytoplasm Dilution And
Contributes to Senescence. *Cell* **176**, 1083-1097.e18 (2019).

28. J. Lengefeld, *et al.*, Cell size is a determinant of stem cell potential during aging. *Sci Adv*
7, eabk0271 (2021).

29. M. C. Lanz, *et al.*, Increasing cell size remodels the proteome and promotes
senescence. *Mol. Cell* **82**, 3255-3269.e8 (2022).

30. S. Manohar, *et al.*, Genome homeostasis defects drive enlarged cells into senescence.
Mol. Cell **83**, 4032-4046.e6 (2023).

31. L. Crozier, *et al.*, CDK4/6 inhibitor-mediated cell overgrowth triggers osmotic and
replication stress to promote senescence. *Mol. Cell* **83**, 4062-4077.e5 (2023).

32. R. Foy, *et al.*, Oncogenic signals prime cancer cells for toxic cell overgrowth during a G1
cell cycle arrest. *Mol. Cell* **83**, 4047-4061.e6 (2023).

33. S. Manohar, G. E. Neurohr, Too big not to fail: emerging evidence for size-induced
senescence. *FEBS J.* **291**, 2291–2305 (2024).

34. P. Jorgensen, J. L. Nishikawa, B.-J. Breitkreutz, M. Tyers, Systematic Identification of
Pathways That Couple Cell Growth and Division in Yeast. *Science* **297**, 395–400 (2002).

35. J. Zhang, *et al.*, Genomic Scale Mutant Hunt Identifies Cell Size Homeostasis Genes in
S. cerevisiae. *Curr. Biol.* **12**, 1992–2001 (2002).

36. M. Ohanna, *et al.*, Atrophy of S6K1(-/-) skeletal muscle cells reveals distinct mTOR
effectors for cell cycle and size control. *Nat. Cell Biol.* **7**, 286–294 (2005).

37. S. M. I. Goorden, *et al.*, Rheb is essential for murine development. *Mol. Cell. Biol.* **31**,
1672–1678 (2011).

38. C. D. Bell, E. Waizbard, Variability of cell size in primary and metastatic human breast
carcinoma. *Invasion Metastasis* **6**, 11–20 (1986).

39. B. M. Iritani, R. N. Eisenman, c-Myc enhances protein synthesis and cell size during B
lymphocyte development. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 13180–13185 (1999).

40. I. Jones, *et al.*, Characterization of proteome-size scaling by integrative omics reveals
mechanisms of proliferation control in cancer. *Sci. Adv.* **9**, eadd0636 (2023).

41. Asadullah, *et al.*, Combined heterogeneity in cell size and deformability promotes
cancer invasiveness. *J. Cell Sci.* **134**, jcs250225 (2021).

42. T. J. Kawecki, *et al.*, Experimental evolution. *Trends Ecol. Evol.* **27**, 547–560 (2012).

43. M. Natalino, M. Fumasoni, Experimental approaches to study evolutionary cell biology
using yeasts. *Yeast* **40**, 123–133 (2023).

44. J. Helsen, G. Sherlock, G. Dey, Experimental evolution for cell biology. *Trends Cell Biol.*
33, 903–912 (2023).

45. Practical Flow Cytometry Shapiro, 4th Edition.

46. Y. H. Jin, *et al.*, The multiple biological roles of the 3'-->5' exonuclease of
Saccharomyces cerevisiae DNA polymerase delta require switching between the
polymerase and exonuclease domains. *Mol. Cell. Biol.* **25**, 461–471 (2005).

47. P. Jorgensen, *et al.*, A dynamic transcriptional network communicates growth potential
to ribosome synthesis and critical cell size. *Genes Dev.* **18**, 2491–2505 (2004).

48. P. Russell, P. Nurse, cdc25+ functions as an inducer in the mitotic control of fission
yeast. *Cell* **45**, 145–153 (1986).

49. K. Lundgren, *et al.*, Mik1 and wee1 cooperate in the inhibitory tyrosine phosphorylation
of cdc2. *Cell* **64**, 1111–1122 (1991).

50. C. M. Chavez, *et al.*, The cell morphological diversity of Saccharomycotina yeasts. *FEMS*
Yeast Res. **24**, foad055 (2024).

51. M. Osumi, Visualization of yeast cells by electron microscopy. *J. Electron Microsc.* **61**,
343–365 (2012).

52. D. H. Williamson, A. W. Scopes, The behaviour of nucleic acids in synchronously
dividing cultures of Saccharomyces cerevisiae. *Exp. Cell Res.* **20**, 338–349 (1960).

53. R. E. Lenski, M. R. Rose, S. C. Simpson, S. C. Tadler, Long-Term Experimental Evolution
in Escherichia coli . I . Adaptation and Divergence During. *Am. Nat.* **138**, 1315–1341
(1991).

54. A. I. Goranov, *et al.*, The rate of cell growth is governed by cell cycle stage. *Genes Dev.*
23, 1408–1422 (2009).

55. A. Giometto, F. Altermatt, F. Carrara, A. Maritan, A. Rinaldo, Scaling body size
fluctuations. *Proceedings of the National Academy of Sciences* **110**, 4646–4650 (2013).

56. T. LaBar, Y. Y. Phoebe Hsieh, M. Fumasoni, A. W. Murray, Evolutionary Repair
Experiments as a Window to the Molecular Diversity of Life. *Curr. Biol.* **30**, R565–R574
(2020).

57. S. Schrott, C. Osman, Two mitochondrial HMG-box proteins, Cim1 and Abf2,
antagonistically regulate mtDNA copy number in Saccharomyces cerevisiae. *Nucleic*
Acids Res. **51**, 11813–11835 (2023).

58. B. L. A. Carter, P. E. Sudbery, SMALL-SIZED MUTANTS OF SACCHAROMYCES
CEREVISIAE. *Genetics* **96**, 561–566 (1980).

59. M. Tyers, G. Tokiwa, R. Nash, B. Futcher, The Cln3-Cdc28 kinase complex of S.
cerevisiae is regulated by proteolysis and phosphorylation. *EMBO J.* **11**, 1773–1784
(1992).

60. M. Caligaris, *et al.*, Snf1/AMPK fine-tunes TORC1 signaling in response to glucose
starvation. *Elife* **12** (2023).

61. N. Talarek, E. Gueydon, E. Schwob, Homeostatic control of START through negative
feedback between Cln3-Cdk1 and Rim15/Greatwall kinase in budding yeast. *Elife* **6**
(2017).

62. S. Mochida, S. L. Maslen, M. Skehel, T. Hunt, Greatwall phosphorylates an inhibitor of
protein phosphatase 2A that is essential for mitosis. *Science* **330**, 1670–1673 (2010).

63. A. Gharbi-Ayachi, *et al.*, The substrate of Greatwall kinase, Arpp19, controls mitosis by
inhibiting protein phosphatase 2A. *Science* **330**, 1673–1677 (2010).

64. N. Chica, *et al.*, Nutritional control of cell size by the greatwall-endosulfine-PP2A·B55
pathway. *Curr. Biol.* **26**, 319–330 (2016).

65. S. Bontron, *et al.*, Yeast endosulfines control entry into quiescence and chronological
life span by inhibiting protein phosphatase 2A. *Cell Rep.* **3**, 16–22 (2013).

66. A. M. Healy, *et al.*, CDC55, a Saccharomyces cerevisiae gene involved in cellular
morphogenesis: identification, characterization, and homology to the B subunit of
mammalian type 2A protein phosphatase. *Mol. Cell. Biol.* **11**, 5767–5780 (1991).

67. T. R. Gregory, Coincidence, coevolution, or causation? DNA content, cell size, and the
C-value enigma. *Biol. Rev. Camb. Philos. Soc.* **76**, 65–101 (2001).

68. J. S. P. Heslop-Harrison, T. Schwarzacher, Q. Liu, Polyploidy: its consequences and
enabling role in plant diversification and evolution. *Ann. Bot.* **131**, 1–10 (2023).

69. Y. Chadha, A. Khurana, K. M. Schmoller, Eukaryotic cell size regulation and its
implications for cellular function and dysfunction. *Physiol. Rev.* **104**, 1679–1717 (2024).

70. I. Soifer, N. Barkai, Systematic identification of cell size regulators in budding yeast.
Mol. Syst. Biol. **10**, 761 (2014).

71. R. A. M. de Bruin, W. H. McDonald, T. I. Kalashnikova, J. Yates 3rd, C. Wittenberg, Cln3
activates G1-specific transcription via phosphorylation of the SBF bound repressor
Whi5. *Cell* **117**, 887–898 (2004).

72. Y. Ho, M. Costanzo, L. Moore, R. Kobayashi, B. J. Andrews, Regulation of transcription at
the Saccharomyces cerevisiae start transition by Stb1, a Swi6-binding protein. *Mol.*
Cell. Biol. **19**, 5267–5278 (1999).

73. R. A. M. de Bruin, T. I. Kalashnikova, C. Wittenberg, Stb1 collaborates with other
regulators to modulate the G_1-specific transcriptional circuit. *Mol. Cell. Biol.* **28**, 6919–
6928 (2008).

74. R. Verma, *et al.*, Phosphorylation of Sic1p by G1 Cdk required for its degradation and
entry into S phase. *Science* **278**, 455–460 (1997).

75. M. Costanzo, *et al.*, CDK activity antagonizes Whi5, an inhibitor of G1/S transcription in
yeast. *Cell* **117**, 899–913 (2004).

76. F. R. Cross, L. Schroeder, J. M. Bean, Phosphorylation of the Sic1 inhibitor of B-type
cyclins in Saccharomyces cerevisiae is not essential but contributes to cell cycle
robustness. *Genetics* **176**, 1541–1555 (2007).

77. M. A. Juanes, *et al.*, Budding yeast greatwall and endosulfines control activity and
spatial regulation of PP2A(Cdc55) for timely mitotic progression. *PLoS Genet.* **9**,
e1003575 (2013).

78. M. Lynch, *et al.*, Evolutionary cell biology: two origins, one objective. *Proc. Natl. Acad.*
Sci. U. S. A. **111**, 16990–16994 (2014).

79. *Guide to yeast genetics and molecular biology* (Academic Press, 1990).

80.   M. Fumasoni, A. W. Murray, The evolutionary plasticity of chromosome metabolism
allows adaptation to constitutive DNA replication stress. *Elife* **9**, 1–28 (2020).

81.   M. Natalino, M. Fumasoni, Compensatory evolution to DNA replication stress is robust
to nutrient availability. *Mol. Syst. Biol.* (2025). https://doi.org/10.1038/s44320-025-
00127-z.

82.   A. Alsina, M. Fumasoni, P. Sartori, Model-based inference of cell cycle dynamics
captures alterations of the DNA replication programme. *PLoS Comput. Biol.* **21**,
e1013570 (2025).

83.   M. Fumasoni, A. W. Murray, Ploidy and recombination proficiency shape the
evolutionary adaptation to constitutive DNA replication stress. *PLoS Genet.* **17**,
e1009875 (2021).