Examples of adaptive peak tracking

as found in the fossil record

Rolf Ergon

rolf.ergon@usn.no

University of South-Eastern Norway

January 12, 2026

Abstract

Species that have persisted over millions of years have done so because they have been able to track peaks in an adaptive landscape well enough to survive and reproduce. Such optima are defined by the mean phenotypic values that maximize mean fitness, and they are predominantly functions of the environment, for example the sea temperature. The mean phenotypic values over time will thus predominantly be determined by the environment over time, and the trait history may be found in the fossil record. Here I use fossil data from four cases found in the literature, and show that adaptive peak tracking models give better results than alternative weighted least squares and directional random walk models. The model performances are compared by use of weighted mean squared errors and Akaike information criterion results.

Keywords: adaptive landscape, adaptive peak tracking, AIC, fitness landscape, fossil record, moving average smoothing, WMSE

1 Introduction

Adaptive peak tracking is known from various engineering applications, but tracking models are also relevant in evolutionary biology settings. They are then based on an adaptive landscape metaphor, which assumes that the mean population fitness in a population is a function of possibly multivariate mean phenotypic traits, with an optimum (adaptive peak) for a specific set of mean traits. See Pigliucci (2013) for historical background and clarifying discussion. Below, I also make use of an individual fitness landscape spanned by axes for individual phenotypic values, for example body size or number of fin rays. Here, individual fitness is a measure of reproductive success, i.e., to which degree genes are passed on to future generations. For species with sexual reproduction the phenotypic values are midparent values, i.e., the means of male and female traits.

It has for a long time been known that populations and species of larger size are found in colder environments, while populations and species of smaller size are found in warmer regions. This is called Bergman's rule after the German biologist Carl Bergman, who described the pattern in 1847. Although there are exceptions from this rule, it is today referred to also in cases where a gradual change in temperature leads to gradual changes in body size, and it is then at least implicitly assumed that the body size tracks a peak in an adaptive landscape. As shown by the last real data example in Section 3, such tracking of environmental change is not restricted to body size, although it indirectly may be so.

Adaptive peak tracking in evolutionary biology can be described by the block diagram in Fig. 1. From a possibly multivariate environment $u(t)$ follows a peak $\theta_i(t)$ in the individual fitness landscape, i.e., individual phenotypic values that maximize fitness, and when $u(t)$ changes the position of this peak will change accordingly. This will again result in a moving adaptive peak $\theta_p(t)$ in the adaptive landscape, which with a non-symmetrical individual fitness function will have a different position than $\theta_i(t)$. As discussed in Ergon (2025), the two peaks may also have

different locations owing to constraints on the phenotypic values, and conceivably also for

other reasons. The essential feature of the model in Fig. 1 is that the tracking process seeks to

keep the mean phenotypic trait $\bar{y}(t)$ equal to the adaptive peak value $\theta_p(t)$, and from a

feedback control point of view it will do so with integral action. In each of the real data cases in

Section 3, we will find that $\theta_p(t)$ is a nearly linear function of a dominating environmental driver,

which means that also $\bar{y}(t)$ is a linear function of $u(t)$. As will be seen in these cases, the

environmental variables may be quite noisy, and I will therefore use moving average smoothed

signals as $u(t)$. This is also a way to handle errors in the estimated age of the fossils.

Figure 1. Block diagram for adaptive peak tracking system in evolutionary biology.

The adaptive peak tracking model in Fig. 1 can be applied on all types of evolutionary

systems, but in the following I will focus on systems involving fossil data. One example is found

in Liow et al. (2024), with fossil data over 2.3 million years on the size of feeding zooids

(autozooids or AZ) in encrusting cheilostome bryozoans. The data were collected from 985

fossil colonies as shown in Fig. 2, with a large total number of individual bryozoans. Results in

the form of nine *log AZ area mean* values with standard errors were analyzed in Ergon (2025),

see panel B in Fig. 3, with present time as zero point. Two samples were omitted for reasons as

discussed in Ergon (2025).

In addition to mean trait standard errors, as shown by error bars in Fig. 3, there are

considerable errors in the nine time points, i.e., in the estimated age. This is because mean trait

values are found as mean values of mean values for various numbers of colonies found in nine

different geological formations with various time spans, and the nine time points are estimated

midpoints for these formations. The dominating environmental driver is the sea temperature,

with oxygen isotope $\partial^{18}O(t)$ values used as proxy (Lisiecki and Raymo, 2005). The $\partial^{18}O(t)$

values increase when the temperature decreases, and they can be converted to temperature by

linear transformations (Evans et al., 2024). Because of the potentially large age errors in the

mean phenotype data, Ergon (2025) used $u(t) = \partial^{18}O_{smooth}(t)$ as a centered moving average

smoothed version of $\partial^{18}O(t)$. Using the moving average window size as a tuning variable, the

nine mean phenotypic values fell quite nicely on a straight line (Fig. 3, panel C), and from

weighted least squares (WLS) parameter estimation thus followed continuous predictions of

the mean trait values (Fig. 3, panel B). Again, note that a high $\partial^{18}O(t)$ value means low

temperature, and that the AZ area thus in accordance with Bergman's rule increases when the

temperature decreases. See Section 3 for detailed results regarding the ovicell (OV) trait as

shown in Fig. 2.

Figure 2. Scanning electron micrograph showing part of a *Microporella agonistes*

colony, with perimeters of autozooid (AZ) and ovicell (OV) areas outlined. The length of

the scale bar is 500 μm. From Liow et al. (2024).

Figure 3. Results for the bryozoan species *Microporella agonistes*, as shown in Fig. 2.

Panel A shows raw $\partial^{18}O(t)$ data (green line) and a centered moving average smoothed

version $\partial^{18}O_{smooth}(t)$ with window size 100 samples (red line). For the four last

samples, the time window size is 0.1 million years, while it is 0.2 million years for

samples 4 and 5, and 0.25 million years for samples 1 to 3. Panel B shows *log AZ area*

mean with error bars, as found from Fig. S2 in Liow at al. (2024) (circles), as well as

predictions (red line). Panel C shows the estimated adaptive peak versus environment

function (red line) computed by the weighted least squares method. Panel B also shows

predictions with window size 800 samples (dashed red line).

The adaptive peak tracking model in Fig. 1 is useful only if the following error $e(t)$ is kept

sufficiently small, such that the mean phenotypic value stays in the vicinity of the adaptive

peak. As referred to in Ergon (2025), there is a lot of support for this assumption. A persistent

system excitation in the form of short-term environmental fluctuations as exemplified in Fig. 3,

panel A, makes it unlikely that a tracking system gets stuck in a local optimum or in a valley in

the adaptive landscape. As illustrated in simulations in Ergon (2025), it is more likely that the

mean phenotypic values fluctuate around global optima.

Under the assumption that the following error $e(t)$ in Fig. 1 is kept sufficiently small, such that

$\bar{y}(t) \approx \theta_p(t)$, the adaptive peak function $\theta_p(t) = f\big(u(t)\big)$ can be identified from $u(t)$ and $\bar{y}(t)$

data, and that is especially simple when this function is linear. As illustrated in Fig. 3 it is then

possible to find the adaptive peak function from few and irregularly spaced samples, and thus

also to predict the mean trait values $\bar{y}(t)$. Note that this can be done without detailed

knowledge of the tracking process in Fig. 1.

Section 2 describes the methods used in the article. As also used in Liow et al. (2024), a well-

established method for analyses of fossil time series data is to find optimal biased or unbiased

random walk models (Hunt, 2006). Biased or general random walk (GRW) models are also

known as directional evolution models (Hunt, 2012). An alternative model for directional

evolution is to use WLS directly on the $\bar{y}(t)$ data. In the real data cases in Section 3 I will thus

compare three models, (i) tracking with use of WLS on the adaptive peak function $\bar{y}(u)$, (ii) WLS

directly on $\bar{y}(t)$, and (iii) GRW. The performances of these models will be compared by use of

weighted mean squared errors (WMSE). The models are also compared by use of the Akaike

Information Criterion (AIC), which is often used for comparison of different random walk

models of fossil data (Hunt, 2006).

In Section 3, the different methods are applied on four real case data sets in the fossil record.

The first case is from the already introduced field study of the bryozoan species *Microporella*

agonistes, with results for *log AZ area mean* as shown in Fig. 3. The second case is from the

same bryozoan study, but with results for *log OV area mean*. The third case is a study of the

ostracod species *Poseidonamics major* in Hunt and Roy (2006), while the fourth is a stickleback

fish study in Bell et al. (1985), which shows that tracking of environmental change is not

restricted to body size traits (although it indirectly may be so). Finally follows a summary and

discussion in Section 4.

2 Theory and Methods

2.1 Essential elements of the tracking process

As discussed in the introduction, it is not necessary to know the details of the tracking process in Fig. 1, but it must have two essential elements. First, there must be a mean fitness function such that $\overline{W}(t) \to \overline{W}_{max}$ when $\bar{y}(t) \to \theta_p(u,t)$. This function defines the adaptive peak, and in the univariate case it can for example be (as in Ergon, 2025)

$$\overline{W}(t) = W_{max}\sqrt{\frac{\omega^2}{\omega^2 + \sigma_y^2(t)}}\exp\left(-\frac{\left(\bar{y}(t) - \theta_p(u,t)\right)^2}{2\left(\omega^2 + \sigma_y^2(t)\right)}\right), \tag{1}$$

where $\sigma_y^2(t)$ is the variance of $y(t)$, while ω^2 is the width of the underlying individual fitness function. Second, there must be a selection mechanism, such that individuals with high fitness get the best chance for reproduction (Lande, 1979). A simple example of such a set of selection equations is given in Ergon (2025). This will in most cases result in slow evolution, but owing to plasticity there may still be rapid changes in the mean trait $\bar{y}(t)$ (Ergon, 2025). In addition, mutations may give sudden and positive jumps in the individual fitness (Ch. 7, Walsh and Lynch, 2018).

2.1 Weighted least squares estimation for tracking models

In the tracking method, least squares predictions are found as $\hat{\bar{y}}_{track}(i) = \hat{a}_{track} + \hat{b}_{track}(u(i) - u(1))$, where $u(i)$ is a moving average smoothed version of the $\partial^{18}O(i)$ measure at the N sampling points. Here, the parameters are found as

$$\begin{bmatrix} \hat{a}_{track} \\ \hat{b}_{track} \end{bmatrix} = (\boldsymbol{X}^T\boldsymbol{W}\boldsymbol{X})^{-1}\boldsymbol{X}^T\boldsymbol{W}\boldsymbol{y}, \tag{2}$$

where $\boldsymbol{X} = \begin{bmatrix} 1 & 1 & \cdots & 1 \\ u(1) - u(1) & u(2) - u(1) & \cdots & u(N) - u(1) \end{bmatrix}^T$ and $\boldsymbol{y} = [\bar{y}(1) \quad \bar{y}(2) \quad \cdots \quad \bar{y}(N)]^T$,

while $\boldsymbol{W}$ is a diagonal matrix with weights $w_i = \sigma_i^{-2}$ that are the reciprocals of the measurement variances. From this follows continuous predictions

$$\hat{\bar{y}}_{track}(t) = \hat{a}_{track} + \hat{b}_{track}(u(t) - u(1)), \qquad (3)$$

144 such that $\hat{\bar{y}}_{track}(t)$ becomes a smoothed and scaled version of the environmental variable.

2.2 Weighted least squares estimation for direct WLS models

146 In the direct WLS method, least squares predictions are found as $\hat{\bar{y}}_{track}(t) = \hat{a}_{WLS} + \hat{b}_{WLS}t$.

147 Here, the parameters are found as in Eq. (2), except that $X = \begin{bmatrix} 1 & 1 & \cdots & 1 \\ t_1 & t_2 & \cdots & t_N \end{bmatrix}^T$, where t_i are

148 the sampling time points.

2.3 Random walk models

Hunt (2006) developed what he called a general random walk (GRW) model for analyses of

fossil time series, later referred to as a model for directional evolution (Hunt, 2012). This model

assumes a random walk process, where at each timestep an increment of evolutionary change

in a mean trait value is drawn at random from a distribution of evolutionary steps. It is also

assumed that the incremental change is normally distributed with mean value μ_{step} and

variance σ_{step}^2. For an observed evolutionary mean trait change ΔY over T time steps, the log-

likelihood function for a GRW process is given by (Hunt, 2006)

$$\ell\left(\mu_{step}, \sigma_{step}^2\right) = -\frac{1}{2}\ln(2\pi) - \frac{1}{2}\ln\left(T\sigma_{step}^2 + \frac{V_{p_A}}{n_A} + \frac{V_{p_D}}{n_D}\right) - \frac{\left(\Delta Y - T\mu_{step}\right)^2}{2\left(T\sigma_{step}^2 + \frac{V_{p_A}}{n_A} + \frac{V_{p_D}}{n_D}\right)}, \qquad (4)$$

where n_A and n_D are the numbers of observed ancestors and descendants, respectively, while

V_{p_A} and V_{p_D} are the corresponding population phenotypic variances. With N irregular and sparse

samples of mean trait values, multiple ancestor-descendant trait differences over $N - 1$

evolutionary transitions may be used jointly to estimate μ_{step} and σ_{step}^2 by summing the log-

likelihoods according to Eq. (4) over the transitions, i.e., by use of

$\ell_{total}\left(\mu_{step}, \sigma_{step}^2\right) = \sum_{i=1}^{N-1} \ell_i\left(\mu_{step}, \sigma_{step}^2\right)$. From this expression, step parameters can be found

by means of maximum likelihood estimation, and the estimated step size $\hat{\mu}_{step}$ will then be a

measure of directional change over time. In the real data examples in Section 3, the optimal

GRW parameters are found by use of the MATLAB function *fmincon*, with initial parameter

values sat to zero.

The GRW model has an obvious weakness in that it is difficult to estimate σ_{step}^2 for small

values of T and large values of V_{p_A}/n_A and V_{p_D}/n_D. In the three examples in Section 3 the result

is $\hat{\sigma}_{step}^2 < 0$, such that the step variance must be set to $\hat{\sigma}_{step}^2 = 0$.

Eq. (4) is not intended for predictions, but once $\hat{\mu}_{step}$ is determined with $\hat{\sigma}_{step}^2 = 0$, a

prediction model

$$\hat{\bar{y}}_{GRW}(t) = \hat{a}_{GRW} + \hat{b}_{GRW}t \tag{5}$$

172   is found by use of $\hat{b}_{GRW} = c\hat{\mu}_{step}$, and by fitting a parameter a_{GWR} to the data by a least squares

173   approach. Here, c is a scaling constant that I consistently will set to $c = 1$.

2.4 Weighted mean squared errors

175   Comparisons of responses $\hat{\bar{y}}_{track}(t)$, $\hat{\bar{y}}_{WLS}(t)$ and $\hat{\bar{y}}_{GRW}(t)$ can be done by use of weighted

176   mean squared errors

$$WMSE = (\bar{\boldsymbol{y}} - \hat{\bar{\boldsymbol{y}}})\boldsymbol{W}(\bar{\boldsymbol{y}} - \hat{\bar{\boldsymbol{y}}})/\sum w_i, \tag{6}$$

177   where $\bar{\boldsymbol{y}}$ and $\hat{\bar{\boldsymbol{y}}}$ are $N \times 1$ data vectors, and where $\boldsymbol{W}$ is a diagonal matrix with the weights w_i.

178   Note that we in the special ordinary least squares (OLS) case have $\boldsymbol{W} = \boldsymbol{I}$ and $\sum w_i = N$. I thus

179   assume independent and normally distributed random variables $\bar{y}(t)$. Note that when the

180   weights are the reciprocals of the measurement variances, as done here, the prediction slope

181   parameters $\hat{b}_{track}$ and $\hat{b}_{WLS}$ are the best linear unbiased estimates (BLUE).

2.4 Theory for the Akaike Information Criterion

183   Model performances may be compared by use of the Akaike information criterion (AIC), and the

184   model with the lowest AIC value should then be chosen (Hunt, 2006). For GRW, the AIC value

185   with compensation for short data is given by (Hunt, 2006),

$$AIC_{c,GRW} = 2k - 2\ell_{total}(\hat{\mu}_{step}, \hat{\sigma}_{step}^2) + 2k(k+1)/(N-k-2), \qquad (7)$$

where $k = 2$ is the number of estimated parameters, while N is the number of samples (the number of transitions is $N - 1$). Since in all the real data examples in Section 3 the estimated variance $\hat{\sigma}_{step}^2$ is negative, the realistic value of $AIC_{c,GRW}$ is found by finding $\hat{\mu}_{step}$ with $\sigma_{step}^2 = 0$.

For comparisons with tracking and WLS models, Eq. (7) must be replaced by (Banks and Joyner, 2017)

$$AIC_c = 2k - 2\ln\left(\mathcal{L}(\hat{\Theta}_{MLE}|\overline{y})\right) + 2k(k+1)/(N-k-1), \qquad (8)$$

where $\mathcal{L}(\hat{\Theta}_{MLE}|\overline{y})$ is the likelihood of the estimated parameters using maximum likelihood estimation, given the vector $\overline{y}$ of observed data. Banks and Joyner (2017) further assume a statistical model $Y_i = f(t_i, \boldsymbol{q}_0) + \omega_i \mathcal{E}_i$, where $\boldsymbol{q}_0$ is a parameter vector, while ω_i are known weights. Here, ε_i for $i = 1, 2, \dots, N$ are i.i.d. $\mathcal{N}(0, \sigma^2)$, such that σ^2 must be estimated together with $\boldsymbol{q}_0$. In the present setting the weights are $\omega_i = \sqrt{1/w_i}$, and the final AIC expression in Banks and Joyner (2017) thus becomes

$$AIC = 2k + Nln(2\pi) + N + 2\sum_{i=1}^{N} \ln\left(\sqrt{1/w_i}\right) + Nln\left(\frac{\sum_{i=1}^{N} w_i(\overline{y}_i - \hat{\overline{y}}_i)^2}{N}\right),$$
$$(9)$$

from which $AIC_{c,track}$ and $AIC_{c,WLS}$ follow by adding $2k(k+1)/(N-k-1)$. Here, $k = 3$, since also the underlying parameter σ^2 must be estimated in addition to a_{WLS} and b_{WLS}. Note that $\hat{\overline{y}}_i$ in Eq. (9) is the result of either a tracking model or WLS, and that Eq. (9) therefore is not valid with use of $\hat{\overline{y}}_{i,GRW}$ as found from Eq. (5).

Two models with known AIC values can be compared by use of Akaike weights (Wagenmakers and Farrell, 2004). Assuming that that the tracking model is best, as we will find in the real data examples in Section 3, we first compute

$$\triangle_{WLS} = AIC_{WLS} - AIC_{track}, \qquad (10a)$$

and

$$\triangle_{GRW} = AIC_{GRW} - AIC_{track}. \qquad (10b)$$

205   We then find the Akaike weights

$$A_{w,track} = \frac{1}{1 + exp\left(-\frac{\triangle_{WLS}}{2}\right) + exp\left(-\frac{\triangle_{GRW}}{2}\right)}, \qquad (11a)$$

$$A_{w,WLS} = \frac{exp\left(-\frac{\triangle_{WLS}}{2}\right)}{1 + exp\left(-\frac{\triangle_{WLS}}{2}\right) + exp\left(-\frac{\triangle_{GRW}}{2}\right)}, \qquad (11b)$$

206   and

$$A_{w,GRW} = \frac{exp\left(-\frac{\triangle_{GRW}}{2}\right)}{1 + exp\left(-\frac{\triangle_{WLS}}{2}\right) + exp\left(-\frac{\triangle_{GRW}}{2}\right)}, \qquad (11c)$$

207   The sum of these weights is one, and the weights can be directly interpreted as conditional

208   probabilities for the models (Wagenmakers and Farrell, 2004).

3 Real data cases

3.1 Bryozoan AZ case

As already introduced, Liow et al. (2024) presented results from a field study of the bryozoan

species *Microporella agonistes*, based on a fossil record over 2.3 million years. The scanning

electron micrograph in Fig. 2 shows perimeters of autozooid (AZ) and ovicell (OV) areas, and

Fig. 3 shows predictions of *log AZ area mean* as found in Ergon (2025). I used the environmental

driver $\partial^{18}O(t)$ as found in Lisiecki and Raymo (2005), and the optimal moving average window

size was 100. In these data the time resolution for $t > -0.6$ million years is 0.001, such that the

moving time window size is 0.1 million years. For $-1.5 < t < -0.6$ the time window size is 0.2

million years, while it for $t < -1.5$ is 0.25 million years. With σ^2_{step} as a free variable, the term

$T\sigma^2_{step} + \frac{V_{p_A}}{n_A} + \frac{V_{p_D}}{n_D}$ in Eq. (4) for one or several of the transitions becomes negative in the course

of the parameter search, such that the log-likelihood becomes a complex number. The search

is therefore limited to $\sigma^2_{step} \geq 0$, resulting in $\hat{\sigma}^2_{step} = 0$. See Table A1 in Appendix A for data, and

Fig. 4 and Table 1 for results. Note that present time is used as zero point. Also note that there

are some unfortunate errors in the AIC results given in Ergon (2025).

Figure 4. Prediction results for Bryozoan AZ case, with mean trait observations (circles

and error bars), tracking predictions (solid blue line), WLS predictions (red line), and

GRW predictions with $\hat{\sigma}^2_{step} = 0$ (dashed red line).

Table 1. Prediction slope, WMSE and AIC results for Bryozoan AZ case.

	Tracking model	WLS model	GRW model
Prediction slope	–	0.1519	0.1441
WMSE	0.0010	0.0019	0.0019
AIC_c	-23	-18	-10
Akaike weights	0.9372	0.0613	0.0015

3.2 Bryozoan OV case

Fig. 5 shows results corresponding to Fig. 4, but now for *log OV area mean*. The optimal moving

window size was here 700 (0.7 to 1.75 million years). The parameter search was also in this

case limited to $\sigma_{step}^2 \geq 0$, resulting in $\hat{\sigma}_{step}^2 = 0$. See Table A2 in Appendix A for data, and Fig. 5

and Table 2 for results.

Figure 5. Prediction results for Bryozoan case 1, with mean trait observations (circles

with standard error bars), tracking predictions (solid blue line), WLS predictions (red

line), and GRW predictions with $\hat{\sigma}_{step}^2 = 0$ (dashed red line).

Table 2. Prediction slope, WMSE and AIC results for Bryozoan OV case.

	Tracking model	WLS model	GRW model
Prediction slope	—	0.1227	0.1805
WMSE	0.0020	0.0021	0.0038
AIC_c	−9	−8	−4
Akaike weights	0.52	0.43	0.05

3.3 Ostracod case

Body size data for the deep-sea ostracod (seed shrimp) species *Poseidonamicus major* are found in Hunt and Roy (2006), Table 4, see Table A3 in Appendix A for details. The body size varies from $y = 707$ to $y = 876\ \mu m$, with the variance in the measurements estimated to be $467/n\ (\mu m)^2$, where n is the number of samples. The standard error is thus $SE = \sqrt{467/n}\ \mu m$, which on a log scale translates to $SE_{log} = \sqrt{467/n}\ \log(y)\,/y$.

   The data in Hunt and Roy (2006) include Mg/Ca temperatures, but a tracking model based on this information is not any better than the WLS model. Instead, I used the environmental driver $\partial^{18}O(t)$ as found in Lisiecki and Raymo (2005), and the optimal moving average window size was here 80 samples (0.08 to 0.2 million years). The parameter search was also in this case limited to $\sigma^2_{step} \geq 0$, resulting in $\hat{\sigma}^2_{step} = 0$. See Table A3 in Appendix A for data, and Fig. 6 and Table 3 for results.

Figure 6. Prediction results for Ostracod case, with mean trait observations (circles with standard error bars), tracking predictions (solid blue line), WLS predictions (red line), and GRW predictions with $\hat{\sigma}^2_{step} = 0$ (dashed red line).

257 **Table 3.** Prediction slope, WMSE and AIC results for Ostracod case.

	Tracking model	WLS model	GRW model
Prediction slope	—	0.0402	0.0316
WMSE	0.000108	0.000241	0.000319
AIC_c	−51	−43	−23
Akaike weights	0.9826	0.0174	$9 \cdot 10^{-7}$

3.4 Stickleback fish case

In this case I use data that show how the dorsal fin ray number evolved in *Gasterosteus*

doryssus, an extinct species of a small freshwater stickleback fish that inhabited inland

freshwater habitats of the North American Great Basin during the Miocene (Bell et al., 1985).

See Table A4 in Appendix A for data. In the data analysis the mean trait values are log-

transformed, with the measurement errors transformed accordingly, i.e.,

$e_{i,log} = e_i \cdot \bar{y}_{i,log}/\bar{y}_i$, where e_i is found from the given standard deviations SD as $e_i = \sqrt{SD_i^2/n_i}$.

The original and log-transformed samples thus have equal coefficients of variation.

For a tracking model, the available environmental data was $\partial^{18}O(t)$ as found in Westerhold et

al. (2020), Table S34, column marked ISOBENbinned_d18Ointerp. This is obviously not the

direct driver of evolution in what is now Nevada, but there is an association between deep sea

temperatures and local Nevada temperatures.

The zero time point not given in Bell et al. (1985), but since $t = 0$ in the data is approximately

10 million years ago (Ma) (personal information) it was possible to align the data in two different

ways such that the prediction results had local optima, and we thus have two stickleback fish

cases. In both cases optimal moving average window sizes of n samples were also found by a

manual search using raw data from 9.8 to 10.3 Ma, and since the time resolution is 0.002

million years, the moving time window sizes were $n \cdot 0.002$ million years. In addition to the

underlying parameter σ^2 and the elevation and slope parameters there are thus two additional

unknown parameters, such that the number of estimated parameters is 5.

- Case 1: The best tracking result was obtained when $t = 0$ in the data corresponds to
  $t_0 = -9.994$ million years with present time as zero point (see Figures 7 and 8, and Table 4). Optimal moving average time window size was 0.192 million years.

- Case 2: The zero point $t_0 = -10.034$ gave almost the same good tracking result (see Figure 9, and Table 4). Optimal moving average time window size was 0.198 million years.

284   The time windows are in both cases very wide compared to the time span of 0.1 million years

285   for the data, such that the effect of smoothing mainly is feature extraction, see Section 4 for

286   discussion and Appendix B for details. The parameter search was also in these cases limited to

287    $\sigma^2_{step} \geq 0$, resulting in $\hat{\sigma}^2_{step} = 0$.

Figure 7. Plots for the Stickleback fish case 1, with upper panel showing $\partial^{18}O(t)$ (blue line) and $u(t) = \partial^{18}O_{smooth}(t)$ with window size 96 (red line), while the lower panel shows data points $\bar{y}(i)$ and $u(i) = \partial^{18}O_{smooth}(i)$ with standard error bars, and the estimated linear adaptive peak function (red line).

Figure 8. Prediction results for Stickleback fish case 1, with mean trait observations

(circles with standard error bars), tracking predictions (blue line), WLS predictions (red

line), and GRW predictions with $\hat{\sigma}_{step}^2 = 0$ (dashed red line, almost identical to the solid

red line).

Figure 9. Prediction results for Stickleback fish case 2, with mean trait observations

(circles with standard error bars), tracking predictions (blue line), WLS predictions (red

line), and GRW predictions with $\hat{\sigma}_{step}^2 = 0$ (dashed red line, almost identical to the solid

red line).

303   **Table 4.** Prediction slope, WMSE and AIC results for Stickleback fish case.

	Tracking model 1	Tracking model 2	WLS model	GRW model
Prediction slope	−	−	1.2605	1.2531
Zero time point	−9.994	-10.034	−	−
Time window size	192	196	−	−
WMSE	0.000216	0.000259	0.000401	0.000401
AIC_c	−131	−126	−121	−104
Akaike weights	0.9941	−	0.0059	$1.8 \cdot 10^{-6}$
Akaike weights	−	0.9405	0.0595	$18 \cdot 10^{-6}$
Akaike weights	0.91	0.09	−	−

4 Summary and discussion

In this article I use fossil data from four real data cases. I show that in cases with directional

evolution, adaptive peak tracking models give better results than alternative weighted least

squares (WLS) and general random walk (GRW) models. The model performances are

compared by use of weighted mean squared errors (WMSE) and Akaike information criterion

(AIC) results.

In all the studied cases, the evolution is driven by a dominating environmental component,

but the tracking model in Fig. 1 is in general multivariate. In multivariate cases, the influences

from several environmental factors on the adaptive peak movements must be found by causal

modeling, which is far beyond the scope of this paper. The adaptive peak may also be

influenced by non-symmetrical fitness functions and developmental constraints as discussed

in Ergon (2025). In the four real data examples, the positions of the adaptive peaks are

approximately linear functions of temperature.

The methodological background is given in Section 2, including the theory for GRW models,

and a description of how AIC values are found from WLS based WMSE results. The results for

the various models are summarized in Figures 4, 5, 6 and 8, and Tables 1, 2, 3 and 4. Note that

the $AIC_{c,GRW}$ values do not take the estimation of a_{GRW} in Eq. (5) into account.

In the tracking models I use oxygen isotope data for the environmental drivers. These data

were found in deep sea drilling projects, and they can be converted to global ground

temperatures through linear transformations (Evans et al., 2024). It is especially interesting that

the last case is a study of fossils of a freshwater fish, which verifies that deep sea oxygen

isotope data contains information on global temperature. This is perhaps the most convincing

case when it comes to tracking model performance (Fig. 8), but it also raises questions as

discussed below.

The reason for use of moving average smoothing in the two Bryozoan cases is that the

individual data are collected from geological formations with time spans from 0.019 to 0.203

million years, and that there is a lot of variation in the raw $\partial^{18}O(t)$ data within each time

window (Ergon, 2025). As seen in Fig. 3, panel B, smoothing is also a feature extraction method

(Herff and Krusienski, 2018), in that the larger window size of 800 samples only captures the

overall trend, while the shorter and optimal window size 100 samples reveals more details.

Such feature extraction is the main purpose in the Stickleback fish case, where possible errors

in the age data are much smaller than in the Bryozoan cases (see Appendix B for details). Note,

however, that the raw $\partial^{18}O(t)$ data as seen in Fig. 7, upper panel, are found from deep-water

drilling samples, and that there may be a complex relationship involving various delays,

feedback mechanisms and oscillations between these data and the local environment at the

field site in what is now Nevada. It is far beyond the aim of this article to consider such complex

relationships, and it is also unclear exactly how the local environment affects mean traits. I

therefore simply assume that the smoothed deep-sea $\partial^{18}O(t)$ data captures an essential part

of the environmental influence on the mean trait, and that an optimal performance can be

found by use of the moving window size and time scale zero point as tuning parameters. This

assumption is supported by the prediction results in Figures 8 and 9, and the good AIC results in

Table 4. It is also supported by the observation that $\partial^{18}O_{smooth}(t)$ appears to be cyclic with an

approximate period of 41,000 years, which can be explained by variations in obliquity, i.e., in

the Earth's axial tilt. The relationship between the local obliquity cycles, as experienced by the

stickleback fish, and the cycles in the deep-sea drilling samples may as pointed out above be

complex, but the essential points are that these cycles have equal frequency but different

amplitude and phase, and that these differences are accounted for by the tuning and regression

procedures as described.

As shown in Appendix B, replacement of the $\partial^{18}O_{smooth}(t)$ input signal by a non-linear

function with a sinus term, may in the stickleback fish case give a reduction in WMSE value.

However, owing to the increased number of parameters the AIC result will be poorer than for

the tracking model. This result supports the evidence that temperature drove the change in

dorsal fin ray number.

One may ask whether the obliquity cycles as seen in Figures 8 and 9 are real, or artifacts

caused by random noise. The reality of the cycles is seen in the results in Appendix B, Fig. B1,

where it, from smoothed and detrended data from 9.2 to 10.8 Ma, is easy to find the period

41,000 years from the total time for 11 cycles. It turns out that that it is more difficult to find the

period in a periodogram based on spectral analysis (Fig. B4). The reality of the cycles is also

supported by the results in Tian et al. (2013), where obliquity and long eccentricity pacing of the

Middle Miocene climate transition is studied.

The stickleback fish case also illustrates an inherent difficulty with the directional evolution

concept. The trait evolution as shown in Fig. 8 has an overall positive directional trend, but in

several shorter time periods the trend is negative. If the fossils had been 200,000 years younger,

it follows from Fig. B1 that the trait evolution instead would have had an overall negative trend,

but that the effect of the obliquity cycles would have been the same. This shows that directional

evolution is not a system property, but a property of the environment. The essential system

property is thus tracking, as illustrated in Fig. 1. In the linear mean trait versus environment

cases studied here, the mean trait tracking predictions are simply smoothed and scaled

versions of the dominating environmental variable.

In the Bryozoan AZ case, the tracking model was clearly best, while the WLS and GRW models

gave very similar results (Fig. 4 and Table 1). The Bryozoan OV case is special in the sense that

tracking and WLS models give quite similar results (Fig. 5 and Table 2). This is not surprising

since the optimal moving window is quite wide (700 samples). In the Ostracod case, the

tracking model was again clearly the best, while the GWR model gave the poorest results (Fig. 6

and Table 3). In the two Stickleback fish cases, the WLS and GRW models gave very similar

results (Figures 8 and 9, and Table 4), while the tracking model was much better.

It would, finally, be interesting to find more linear cases in the literature. It would be even

more interesting to find cases with non-linear mean trait versus environment functions, as

discussed in Ergon (2025). The tracking methodology used here can easily be extended to

handle such cases. It would also be interesting to find more cases with obliquity cycles as vital

parts of the environmental driver.

386 Acknowledgements

387 I thank Mike Bell for information regarding the time span of the stickleback fish data and for the

388 observation of possible obliquity cycles. I also thank University of South-Eastern Norway for

389 support and funding.

390 Competing interests

391 There are no competing interests.

Data Availability Statement

Oxygen isotope data for Figures 4, 5 and 6 are archived as *Raw data for simulations* on bioRxiv,

https://doi.org/10.1101/2024.10.30.621046

Oxygen isotope data for Figures 7, 8 and 9 are archived as *Data for Stickleback fish case* on

bioRxiv, doi: https://doi.org/10.64898/2025.12.07.692820

References

Banks, H.T., and Joyner, M.L. (2017). AIC under the framework of least squares estimation. Applied Mathematics Letters. http://dx.doi.org/10.1016/j.aml.2017.05.005

Bell, M. A., Baumgartner, J. V., & Olson, E. C. (1985). Patterns of temporal change in single morphological characters of a Miocene stickleback fish. Paleobiology 11(3): 258-271.

Ergon, R. (2025). Adaptive Peak Tracking as Explanation of Sparse Fossil Data Across Fluctuating Ancient Environments. Ecology and Evolution. First published: 03 July 2025 https://doi.org/10.1002/ece3.71705

Evans, D., Brugger, J., Inglis, G.N., and Valdes, P. (2024). The Temperature of the Deep Ocean is a Robust Proxy for Global Mean Surface Temperature During the Cenozoic. Paleoceanography and Paleoclimatology. doi: 10.1029/2023PA004788

Herff, C., and Krusienski, D.J. (2018). Extracting Features from Time Series, Chapter 7 in Fundamentals of Clinical Data Science [Internet]. Kubben P, Dumontier M, Dekker A, editors. Cham (CH): Springer; 2019.

Hunt, G. (2006). Fitting and comparing models of phyletic evolution: random walks and beyond. Paleobiology 32: 578–601.

Hunt, G., and Roy, K. (2006). Climate change, body size evolution, and Cope's Rule in deep-sea ostracodes. PNAS, vol. 103/5: 1347–1352.

415 www.pnas.org/cgi/doi/10.1073/pnas.0510550103

416 Hunt, G. (2012). Measuring rates of phenotypic evolution and the inseparability of tempo and

417 Mode. Paleobiology, 38(3), 2012, pp. 351–373. paleo_Hunt_2012_Pb_rates.pdf

418 Hunt, G., Hopkins, M.J., and Lidgard, S. (2015). Simple versus complex models of trait evolution

419 and stasis as a response to environmental change. PNAS, vol. 112 /16: 4885–4890.

420 Lande R (1979) Quantitative genetic analysis of multivariate evolution, applied to

421 brain:body size allometry. Evolution 33:402-416

422 www.pnas.org/cgi/doi/10.1073/pnas.1403662111

423 Lisiecki, L. E., and Raymo, M. E. (2005). A Pliocene–Pleistocene stack of 57 globally distributed

424 benthic δ18O records. Paleoceanography 20, A1003. doi:10.1029/ 2004PA001071

425 Liow, L.H., Porto, A., and Di Martino, E. (2024). Trait-Fitness Associations via Fecundity and

426 Competition in a Two-Million-Year-Long Fossil Record. Am. Nat. 204/3.

427 https://doi.org/10.1086/73133

428 Pigliucci, M. (2013). Landscapes, Surfaces, and Morphospaces: What Are They Good For? In

429 The Adaptive Landscape in Evolutionary Biology, Eds. Svensson, E. and R. Calsbeek. Oxford

430 University Press. https://doi.org/10.1093/acprof:oso/9780199595372.001.0001

431 Tian, J., Yang, M., Lyle, M.W., Wilkens, R., Shackford, J.K., (2013). Obliquity and long eccentricity

432 pacing of the Middle Miocene climate transition. Geochemistry, Geophysics, Geosystems,

433 doi:10.1002/ggge.20108

434 Wagenmakers, E-J., and Farrell, S. (2004). AIC model selection using Akaike weights.

435 Psychonomic Bulletin & Review, 11 (1), 192-196

436 Walsh, B., and Lynch, M. (2018). Evolution and selection of quantitative traits.

437 Oxford University Press

438 Westerhold et al. (2020). An astronomically dated record of Earth's climate and its

439   predictability over the last 66 million years. Science 369, 1383–1387.

440   DOI: 10.1126/science.aba6853

Appendix A Data for real data cases

442 **Table A1.** Data for AZ bryozoan case

Age Ma	$\partial^{18}O(t)$ temperature	log AZ area mean	SE value
2.1895	3.6518	11.3930	0.0500
1.9660	3.7409	11.4230	0.0500
1.8505	3.7487	11.4550	0.1200
0.9265	4.0233	11.6150	0.0500
0.6485	4.2053	11.6600	0.0400
0.5480	4.0763	11.5550	0.0450
0.5055	4.1545	11.6550	0.0400
0.4510	4.1775	11.7500	0.0850
0.3990	4.1735	11.7000	0.1200

444 **Table A2.** Data for OV bryozoan case

Age Ma	$\partial^{18}O(t)$ temperature	log AZ area mean	SE value
2.1895	3.6110	10.2600	0.0650
1.9660	3.6934	10.1700	0.0650
0.9265	4.0426	10.4300	0.0950
0.6485	4.0779	10.3900	0.0550
0.5480	4.0935	10.3600	0.0700
0.5055	4.1210	10.4300	0.0600
0.4510	4.1357	10.5000	0.1350

446 **Table A3**. Data for Ostracod case

Age Ma	Mg/Ca temperature	n	Mean valve length μm
3.33	2.54	13	707
2.60	2.23	4	744
2.40	2.15	21	742
2.19	2.07	23	743
1.90	1.96	25	745
1.68	1.88	22	763

0.89	1.55	28	812
0.14	1.19	24	797
0.12	1.18	3	793
0.00	1.12	31	812

448 **Table A4.** Data for Stickleback fish case with $\bar{y} = dorsal\ fin\ ray\ number$

Time	N	$\bar{y}$	SD
0	30	8.8300	0.7470
0.0053	33	8.7600	0.6630
0.0100	30	8.8700	0.6810
0.0152	36	9.0600	0.7150
0.0193	55	8.5800	0.7620
0.0256	40	8.7500	0.8990
0.0305	75	8.5600	0.7750
0.0340	52	8.7700	0.7030
0.0366	22	9.1800	0.5010
0.0381	28	9.1800	0.7720
0.0410	54	8.9400	0.6270
0.0506	45	9.2000	0.5880
0.0552	35	9.1400	0.6480
0.0603	37	9.0500	0.6640
0.0653	41	9.1000	0.5390
0.0701	25	9.2000	0.7070
0.0751	30	9.1300	0.7300
0.0811	58	9.3100	0.7540
0.0865	32	9.7500	0.7180
0.0920	84	9.6300	0.8030
0.0934	82	10.0200	0.8460
0.0967	89	9.8000	0.8280
0.0994	95	9.6200	0.7320
0.1013	57	9.6700	0.7150
0.1070	45	9.8000	0.6940
0.1077	64	9.8400	0.7180

Appendix B Obliquity cycles in Stickleback fish case

B1. Data smoothing for feature extraction

452 For an elementary theoretical analysis, assume the model

$$u(t) = A sin(2\pi f t) + u_s(t) + e(t),\qquad\qquad\text{(B1)}$$

where $u_s(t)$ has a structure that is important for prediction of the mean traits in the block

diagram in Fig. 1, while $e(t)$ is unimportant high frequence noise. Also assume a sampling

frequency $f_s \gg f$, moving average smoothing with window sample size n, and n large enough to

make the smoothed high frequency noise $e_{smooth}(t) \approx 0$.

For an integral number of periods in the moving average time window, the average of the

sinusoidal component in Eq. (B1) is zero. For a non-integral number of periods we thus find

$\sum_{\tau=t-0.5n/f_s}^{\tau=t+0.5n/f_s} sin(2\pi f\tau) = \sum_{\tau=0}^{n_p/f_s} sin(2\pi f\tau)$, where $0 < n_p < f_s/f$ (where f_s/f is the number of

samples per period). We thus find

$$u_{smooth}(t) = \frac{1}{n} A \sum_{\tau=o}^{n_p/f_s} sin(2\pi f\tau) + \frac{1}{n} \sum_{\tau=t-0.5n/f_s}^{\tau=t+0.5n/f_s} u_s(\tau). \tag{B2}$$

For a non-integral number of periods in the moving time window, the sinusoidal component will

thus be seen in $u_{smooth}(t)$ with the frequency f, but with an amplitude depending on n and n_p

according to Eq. (B2), while the structured information will be smoothed. The sampling window

must thus be wide enough to filter out the noise component $e(t)$, but small enough to retain the

information on the sinusoidal component and the essential features in the structured

information in $u_s(t)$. And the window should not be equal to an integer number of periods. As

shown in Fig. 7, an optimal solution can be found by use of the window size as a tuning

parameter.

As an example, Fig. B1, upper panel, shows the same smoothed mean trait data as in Fig. 7,

upper panel, but detrended and over a longer time span. The mean trait apparently includes

cycles with an approximate period of 41,000 years, i.e., with $f = 24\ \mu Hz$, which indicates

variations caused by alterations in the Earth's axial tilt (obliquity). Since $f_s = 500\ \mu Hz$, the

number of samples per period is $f_s/f \approx 20$. From the theory above follows that the obliquity

cycles should vanish with an integer number of cycles in the sampling window, for example for

a window size of 164,000 years, as shown in Fig. B1, mid-panel. When the window size is further

reduced, the obliquity cycles reappear, now with a larger amplitude owing to the reduced

number of samples.

Figure B1. Obliquity cycles with an approximate period of 41,000 years, as seen with

window size 192,000 years (upper panel). The cycles vanish with window size 164,000

years (mid-panel), and reappear with window size 140,000 years (lower panel).

Fig. B2 shows the smoothed mean trait data for different window sizes without detrending. With

short window sizes the obliquity cycles are hidden in noise, with medium window sizes the

cycles stand out, and with wide windows they are so much reduced that they again tend to be

lost.

Figure B2. Smoothed $\partial^{18}O(t)$ data for different time-window sizes.

B2. Periodogram

In an attempt to find the obliquity cycles by spectral density analysis, the smoothed and

detrended data $u(t)$ in Fig. B1, upper panel, was used to find a periodogram, i.e., estimates of

the power spectral density as function of frequency. First, a further smoothed signal $u_{smooth}(t)$

was found using a moving average time window of 80,000 years. Second, a residual

$u_{res}(t) = u(t) - u_{smooth}$ was computed as shown in Fig. B3. This residual signal was finally

used in the MATLAB function *periodogram*, with results as shown in Fig. B4, upper panel. Here,

the peak at 27 cycles per million years (period 37,000 years) indicates the obliquity cycles. As

shown in Fig. B1, upper panel, a more correct result is found by simply counting peaks over

time. The peak at 51 cycles per million years (period 20,000 years) could possibly be explained

by precession (axial wobble) with periods of 23,000 years ([Milankovitch cycles - Wikipedia](#)). The

error in the estimated periods of the obliquity and precession cycles may thus be caused by

interference between these cycles, as also indicated in Fig. B3 (red line), and possibly also

other Milankovitch cycles. The periodogram with window size 164,000 years, such that the

obliquity cycles in theory should vanish, is shown in Fig. B4, mid-panel. Here, the peak at 27

cycles per million years is very much reduced, but instead, and for unknown reasons, strong

peaks at 21 and 32 cycles per million years are dominating features. It is interesting to note that

the peak at 51 cycles per million years also is very much reduced with window size 164,000

years, and that 164,000 is close to the integer number 7 times the precession period of 23,000

years.

Figure B3. Detrended signal $u(t)$ as in Fig, B1, upper panel, smoothed signal $u_{smooth}(t)$

(dashed blue line), and residual $u_{res}(t) = u(t) - u_{smooth}(t)$ (red line).

Figure B4. Periodograms corresponding to the time series in Fig. B1, as found from

$u_{res}(t)$ as in Fig. B3 (red line) by use of the MATLAB function *periodogram*. Note the

different scaling in the lower panel.

B3. Sinusoidal model

For comparison with the tracking response in Fig. (8) we may use a non-linear prediction model

with a sinus function,

$$\bar{y}_{non-lin} = a + b_1(t - t_0) + b_2(t - t_0)^2 + A sin(2\pi f(t - t_0) + \varphi), \qquad (B3)$$

where $f = \frac{1}{0.041} = 24.4$ periods pr. million years. A numerical parameter search for minimum

WMSE using the function *fmincon* in MATLAB resulted in $a = 2.16$, $b_1 = 0.27$, $b_2 = 8.64$,

$A = 0.0176$ and $\varphi = 0.52$. With $b_2 = 0$ this was changed into $a = 2.14$, $b_1 = 1.28$, $b_2 = 0$,

$A = 0.0177$ and $\varphi = 0.38$. See Fig. B3 and Table B for results. Note that the model in Eq. (B3)

with b_2 as a free variable is only slightly better than the tracking model, while $b_2 = 0$ gives a

somewhat poorer model.

Figure B4. Tracking response as in Fig. 8 (blue line) and by use of Eq. (B3) with b_2 as free

variable (red line). The response with $b_2 = 0$ is shown by dashed red line.

528   **Table B.** Prediction slope, WMSE and AIC results for Stickleback fish case.

	Tracking model	Model in Eq. (B1)	Model in Eq. (B1) with $b_2 = 0$
WMSE	0.000216	0.000201	0.000258
AIC_c	-131	-133	-126
Akaike weights	0.28	0.72	$-$
Akaike weights	0.91	$-$	0.09

Appendix C MATLAB code for Stickleback fish case

```
531    clear
532    load Data_9800_to_10300.mat
533    Age=flipud(Age);
534    d18O=flipud(d18O);
535    tdata=-Age';        % Step size 0.002 My
536    rawdata=d18O';
537    window=96;
538    u=movmean(rawdata,window);
539    t0=-10+0.0060;
540
541    %% Sample data
542    tlabel1=[0 5317 9950 15182 19318 25551 30541 34030 36631 38052 41008 50629 55246];
543    tlabel2=[60289 65312 70119 75075 81150 86454 92006 93445 96659 99377 101306 106969 107667];
544    tlabel=[tlabel1 tlabel2]/1000000;
545    tlabel=t0+tlabel;
546    n=[30 33 30 36 55 40 75 52 22 28 54 45 35 37 41 25 30 58 32 84 82 89 95 57 45 64];
547    y1=0.01*[883 876 887 906 858 875 856 877 918 918 894 920 914 ];
548    y2=0.01*[905 910 920 913 931 975 963 1002 980 962 967 980 984];
549    y=[y1 y2];
550    sd1=0.001*[747 663 681 715 762 899 775 703 501 772 627 588 648  ];
551    sd2=0.001*[664 539 707 730 754 718 803 846 828 732 715 694 718];
552    sd=[sd1 sd2];
553    var=sd.^2;
554    err0=sqrt(var./n);
```

```matlab
555    ylabell=log(y);
556    for i=1:26
557      err(i)=err0(i)*(ylabell(i)/y(i))^1;
558    end
559
560    %%% Tracking model
561    tplot=tdata;
562    ulabel=zeros(1,length(tlabel));
563    tlab=tlabel;
564    for i=1:length(tlabel)
565      for t=1:length(tplot)
566        if tplot(t)>min(tlab)
567          ulabel(1,i)=u(1,t);
568          tlab(1,i)=0;
569          break
570        end
571      end
572    end
573    w=err.^-2;
574    W=diag(w);
575    N=26;
576    k=3;
577    X=[ones(length(tlabel),1) ulabel'-2.9559*ones(length(tlabel),1)];
578    bls1=inv(X'*W*X)*X'*W*ylabell'
579    a1=bls1(1);
580    b1=bls1(2);
581    yhat1=a1+b1*(ulabel-2.9559*ones(1,26));
582    for t=1:length(u)
583      yhatplot(t)=a1+b1*(u(t)-2.9559);
584    end
585    WMSE_Tracking=sum((ylabell-yhat1)*W*(ylabell-yhat1)')/sum(diag(W))
586    for j=1:N
587      lnw(j)=log(sqrt(1/w(j)));
588      z(j)=(ylabell(j)-yhat1(j)).^2*w(j);
589    end
590    AICc_Tracking=2*k+N*log(2*pi)+N+2*sum(lnw)+N*log(sum(z)/N)+2*k*(k+1)/(N-k-1);
591
```

```matlab
%% WLS model
X=[ones(26,1) tlabel'];
bls=inv(X'*W*X)*X'*W*ylabell';
a2=bls(1);
b2=bls(2);
for i=1:26
    yhat2(i)=a2+b2*tlabel(i);
end
WMSE_WLS=(ylabell-yhat2)*W*(ylabell-yhat2)'/sum(diag(W))
for j=1:N
    lnw(j)=log(sqrt(1/w(j)));
    z(j)=(ylabell(j)-yhat2(j)).^2*w(j);
end
AICc_WLS=2*k+N*log(2*pi)+N+2*sum(lnw)+N*log(sum(z)/N)+2*k*(k+1)/(N-k-1);

%% GRW model
k=2;
N=25;
n_samples=ones(1,26);
var_samples=err.^2;
Tsample=tlabel;
for i=1:25
    dT(i)=-Tsample(i)+Tsample(i+1);
    dX(i)=ylabell(i+1)-ylabell(i);
    nA(i)=n_samples(i);
    nD(i)=n_samples(i+1);
    varA(i)=var_samples(i);
    varD(i)=var_samples(i+1);
end

% Contraints and
mustep_min=-10; mustep_max=10;
varstep_min=0; varstep_max=1;
par_lb=[mustep_min varstep_min];
par_ub=[mustep_max varstep_max];
par_guess=[0 0];
Aineq=[]; Bineq=[]; Aeq=[]; Beq=[];
```

```matlab
629    fun_objective_handle=...
630       @(par)fun_objective(par,dT,dX,varA,nA,varD,nD);
631    [par_opt,fval,exitflag,output,lambda,grad,hessian] =...
632    fmincon(fun_objective_handle,par_guess,Aineq,Bineq,Aeq,Beq,par_lb,par_ub);
633    mustep=par_opt(1);
634    varstep=par_opt(2);
635    for i=1:25
636       varterm(i)=dT(i)*varstep+varA(i)/nA(i)+varD(i)/nD(i);
637    end
638    for i=1:25
639       Li(i)=-log(2*pi)/2-log(varterm(i))/2-(dX(i)-dT(i)*mustep)^2/(2*varterm(i));
640    end
641    L=sum(Li);
642    AICc_GRW=2*k-2*L+2*k*(k+1)/(N-k-1);
643    X=ones(26,1);
644    b3=mustep;
645    a3=inv(X'*W*X)*X'*W*(ylabell'-b3*tlabel');
646    for i=1:26
647       yhat3(i)=a3+b3*tlabel(i);
648    end
649    WMSE_GRW=sum((ylabell-yhat3)*W*(ylabell-yhat3)')/sum(diag(W));
650
651    % Akaike weights
652     D_Tracking=0;
653     D_WLS=AICc_WLS-AICc_Tracking;
654     D_GRW=AICc_GRW-AICc_Tracking;
655     L_WLS=exp(-0.5*D_WLS);
656     L_GRW=exp(-0.5*D_GRW);
657     Aweight_Tracking=1/(1+L_WLS+L_GRW);
658     Aweight_WLS=L_WLS/(1+L_WLS+L_GRW);
659     Aweight_GRW=L_GRW/(1+L_WLS+L_GRW);
660
661
662    yplot=a1+b1*(ulabel'-2.95*ones(26,1));
663
664    figure(7)
665    subplot(2,1,1)
```

```matlab
666    plot(tplot,rawdata,'b','LineWidth',1.0), hold on
667    plot(tplot,u,'m','LineWidth',1.0), hold off, grid
668    axis([-10 -9.88 2.5 3.3])
669    xlabel('Million years')
670    ylabel('d^1^8O and d^1^8O_s_m_o_o_t_h')
671    subplot(2,1,2)
672    errorbar(ulabel,ylabell,err,'bo','LineWidth',1.0), hold on, grid
673    plot(ulabel,yplot,'m','LineWidth',1.0), hold off
674    axis([2.95 3.03 2.12 2.33])
675    xlabel('d^1^8O_s_m_o_o_t_h')
676    ylabel('log mean dorsal fin ray number')
677
678    figure(8)
679    errorbar(tlabel,ylabell,err,'bo','LineWidth',1.0), hold on
680    plot(tplot,yhatplot,'b','LineWidth',1.0)
681    plot(tlabel,yhat2,'m');
682    plot(tlabel,yhat3,'m--','LineWidth',1.0)
683    hold off, grid
684    axis([-10 -9.88 2.12 2.33])
685    xlabel('Million years')
686    ylabel('log mean dorsal fin ray number')
687
688    b2
689    b3
690    WMSE_Tracking
691    AICc_Tracking
692    WMSE_WLS
693    AICc_WLS
694    WMSE_GRW
695    AICc_GRW
696    Aweight_Tracking
697    Aweight_WLS
698    Aweight_GRW
699
700
701    %%% Parameter search
702    function f = fun_objective(par,dT,dX,varA,nA,varD,nD)
```

```
703    mustep=par(1);
704    varstep=par(2);
705    for i=1:25
706       varterm=dT(i)*varstep+varA(i)/nA(i)+varD(i)/nD(i);
707       Li(i)=-log(2*pi)/2-log(varterm)/2-(dX(i)-dT(i)*mustep)^2/(2*varterm);
708    end
709    L=sum(Li);
710    f=-L;
711    end
```