Examples of adaptive peak tracking

as found in the fossil record

Rolf Ergon

rolf.ergon@usn.no

University of South-Eastern Norway

December 10, 2025

Abstract

Species that have persisted over millions of years have done so because they have been able to track peaks in an adaptive landscape well enough to survive and reproduce. Such optima are defined by the mean phenotypic values that maximize mean fitness, and they are predominantly functions of the environment, for example the sea temperature. The mean phenotypic values over time will thus predominantly be determined by the environment over time, and the trait history may be found in the fossil record. Here I use fossil data from three cases found in the literature, and I show that adaptive peak tracking models give better results than alternative weighted least squares and directional random walk models. The model performances are compared by use of weighted mean squared errors and Akaike information criterion results.

Keywords: adaptive landscape, adaptive peak tracking, AIC, fitness landscape, fossil record, moving average smoothing, WMSE

1 Introduction

Adaptive peak tracking is known from various engineering applications, but tracking models are also relevant in evolutionary biology settings. They are then based on an adaptive landscape metaphor, which assumes that the mean population fitness in a population is a function of possibly multivariate mean phenotypic traits, with an optimum (adaptive peak) for a specific set of mean traits. See Pigliucci (2013) for historical background and clarifying discussion. Below, I also make use of an individual fitness landscape spanned by axes for individual phenotypic values, for example body size or breeding time. Here, individual fitness is a measure of reproductive success, i.e., to which degree genes are passed on to future generations. For species with sexual reproduction the phenotypic values are midparent values, i.e., the means of male and female traits.

It has for a long time been known that populations and species of larger size are found in colder environments, while populations and species of smaller size are found in warmer regions. This is called Bergman's rule after the German biologist Carl Bergman, who described the pattern in 1847. Although there are exceptions from this rule, it is today referred to also in cases where a gradual change in temperature leads to gradual changes in body size, and it is then at least implicitly assumed that the body size tracks a peak in an adaptive landscape. As shown by the first real data example in Section 3, such tracking of environmental change is not restricted to body size, although it indirectly may be so.

Adaptive peak tracking in evolutionary biology can be described by the block diagram in Fig. 1. From a possibly multivariate environment $u(t)$ follows a peak $\theta_i(t)$ in the individual fitness landscape, i.e., individual phenotypic values that maximize fitness, and when $u(t)$ changes the position of this peak will change accordingly. This will again result in a moving adaptive peak $\theta_p(t)$ in the adaptive landscape, which with a non-symmetrical individual fitness function will have a different position than $\theta_i(t)$. As discussed in Ergon (2025), the two peaks may also have

different locations owing to constraints on the phenotypic values, and conceivably also for

other reasons. The essential feature of the model in Fig. 1 is that the tracking process seeks to

keep the mean phenotypic trait $\bar{y}(t)$ equal to the adaptive peak value $\theta_p(t)$, and it will do so

with integral action. In the real data cases in Section 3 we will find that $\theta_p(t)$ are nearly linear

functions of a dominating environmental driver, which means that also $\bar{y}(t)$ are linear functions

of $u(t)$. As will be seen in the cases in Section 3, the environmental variables may be quite

noisy, and I will therefore use moving average smoothed signals as $u(t)$. This is also a way to

handle errors in the estimated age of the fossils.

Figure 1. Block diagram for adaptive peak tracking system in evolutionary biology.

The adaptive peak tracking model in Fig. 1 can be applied on all types of evolutionary

systems, but in the following I will focus on systems involving fossil data. One example is found

in Liow et al. (2024), with fossil data over 2.3 million years on the size of feeding zooids

(autozooids or AZ) in encrusting cheilostome bryozoans. The data were collected from 985

fossil colonies as shown in Fig. 2, with a large total number of individual bryozoans. Results in

the form of nine *log AZ area mean* values with standard errors were analyzed in Ergon (2025),

see panel B in Fig. 3, with present time as zero point. Two samples were omitted for reasons as

discussed in Ergon (2025).

In addition to mean trait standard errors, as shown by error bars in Fig. 3, there are

considerable errors in the nine time points, i.e., in the estimated age. This is because mean trait

values are found as mean values of mean values for various numbers of colonies found in nine

different geological formations with various time spans, and the nine time points are estimated

midpoints for these formations. The dominating environmental driver is the sea temperature,

with oxygen isotope $\partial^{18}O(t)$ values used as proxy (Lisiecki and Raymo, 2005). The $\partial^{18}O(t)$

values increase when the temperature decreases, and they can be converted to temperature by

linear transformations (Evans et al., 2024). Because of the potentially large age errors in the

mean phenotype data, Ergon (2025) used $u(t) = \partial^{18}O_{smooth}(t)$ as a centered moving average

smoothed version of $\partial^{18}O(t)$. Using the moving average window size as a tuning variable, the

nine mean phenotypic values fell quite nicely on a straight line (Fig. 3, panel C), and from

weighted least squares (WLS) parameter estimation thus followed continuous predictions of

the mean trait values (Fig. 3, panel B). Again, note that a high $\partial^{18}O(t)$ value means low

temperature, and that the AZ area thus in accordance with Bergman's rule increases when the

temperature decreases. See Section 3 for detailed results regarding the ovicell (OV) trait as

shown in Fig. 2.

Figure 2. Scanning electron micrograph showing part of a *Microporella agonistes*

colony, with perimeters of autozooid (AZ) and ovicell (OV) areas outlined. The length of

the scale bar is 500 μm. From Liow et al. (2024).

Figure 3. Results for the bryozoan species *Microporella agonistes*, as shown in Fig. 2.

Panel A shows raw $\partial^{18}O(t)$ data (green line) and a centered moving average smoothed

version $\partial^{18}O_{smooth}(t)$ with window size 100 samples (red line). For the four last

samples, the time window size is 0.1 million years, while it is 0.2 million years for

samples 4 and 5, and 0.25 million years for samples 1 to 3. Panel B shows *log AZ area*

mean with error bars, as found from Fig. S2 in Liow at al. (2024) (circles), as well as

predictions (red line). Panel C shows the estimated adaptive peak versus environment

function (red line) computed by the weighted least squares method. Panel B also shows

predictions with window size 800 samples (dashed red line).

The adaptive peak tracking model in Fig. 1 is useful only if the following error $e(t)$ is kept

sufficiently small, such that the mean phenotypic value stays in the vicinity of the adaptive

peak. As referred to in Ergon (2025), there is a lot of support for this assumption. A persistent

system excitation in the form of short-term environmental fluctuations as exemplified in Fig. 3,

panel A, makes it unlikely that a tracking system gets stuck in a local optimum or in a valley in

the adaptive landscape. As illustrated in simulations in Ergon (2025), it is more likely that the

mean phenotypic values fluctuate around global optima.

Under the assumption that the following error $e(t)$ in Fig. 1 is kept sufficiently small, such that

$\bar{y}(t) \approx \theta_p(t)$, the adaptive peak function $\theta_p(t) = f\big(u(t)\big)$ can be identified from $u(t)$ and $\bar{y}(t)$

data, and that is especially simple when this function is linear. As illustrated in Fig. 3 it is then

possible to find the adaptive peak function from few and irregularly spaced samples, and thus

also to predict the mean trait values $\bar{y}(t)$. Note that this can be done without detailed

knowledge of the tracking process in Fig. 1.

Section 2 describes the methods used in the article. As also used in Liow et al. (2024), a well-

established method for analyses of fossil time series data is to find optimal biased or unbiased

random walk models (Hunt, 2006). Biased or general random walk (GRW) models are also

known as directional evolution models (Hunt, 2012). An alternative model for directional

evolution is to use WLS directly on the $\bar{y}(t)$ data. In the real data cases in Section 3 I will thus

compare three models, (i) tracking with use of WLS on the adaptive peak function $\bar{y}(u)$, (ii) WLS

directly on $\bar{y}(t)$, and (iii) GRW. The performance of these models will be compared by use of

weighted mean squared errors (WMSE). The models are also compared by use of the Akaike

Information Criterion (AIC), which is often used for comparison of different random walk

models of fossil data (Hunt, 2006).

In Section 3, the different methods are applied on three real case data sets in the fossil

record. The first is a stickleback fish study in Bell et al. (1985), which shows that tracking of

environmental change is not restricted to body size traits (although it indirectly may be so). The

second is a study of the ostracod species *Poseidonamics major* in Hunt and Roy (2006), while

the third is the already introduced field study of the bryozoan species *Microporella agonistes*,

with results as in Fig. 3, but now with use of data for the ovicell (OV) area as shown in Fig. 2.

Finally follows a summary and discussion in Section 4.

2 Theory and Methods

2.1 Essential elements of the tracking process

It is not necessary to know the details of the tracking process in Fig. 1, but it must have two essential elements. First, there must be a mean fitness function such that $\overline{W}(t) \to \overline{W}_{max}$ when $\overline{y}(t) \to \theta_p(u, t)$. This function defines the adaptive peak, and in the univariate case it can for example be (as in Ergon, 2025)

$$\overline{W}(t) = W_{max}\sqrt{\frac{\omega^2}{\omega^2 + \sigma_y^2(t)}} \exp\left(-\frac{\left(\overline{y}(t) - \theta_p(u, t)\right)^2}{2\left(\omega^2 + \sigma_y^2(t)\right)}\right), \tag{1}$$

where $\sigma_y^2(t)$ is the variance of $y(t)$, while ω^2 is the width of the underlying individual fitness function. Second, there must be a selection mechanism, such that individuals with high fitness get the best chance for reproduction (Lande, 1979). A simple example of such a set of selection equations is given in Ergon (2025). This will in most cases result in slow evolution with small changes in $\overline{y}(t)$ from generation to generation, but in addition there may be mutations that give sudden and positive jumps in the individual fitness (Ch. 7, Walsh and Lynch, 2018).

2.1 Weighted least squares estimation for tracking models

In the tracking method, least squares predictions are found as $\hat{\overline{y}}_{track}(i) = \hat{a}_{track} + \hat{b}_{track}(u(i) - u(1))$, where $u(i)$ is a moving average smoothed version of the $\partial^{18}O(i)$ measure at the N sampling points. Here, the parameters are found as

$$\begin{bmatrix} \hat{a}_{track} \\ \hat{b}_{track} \end{bmatrix} = (X^TWX)^{-1}X^TWy, \tag{2}$$

where $X = \begin{bmatrix} 1 & 1 & \cdots & 1 \\ u(1) - u(1) & u(2) - u(1) & \cdots & u(N) - u(1) \end{bmatrix}^T$ and $y = [\overline{y}(1) \quad \overline{y}(2) \quad \cdots \quad \overline{y}(N)]^T$, while W is a diagonal matrix with weights $w_i = \sigma_i^{-2}$ that are the reciprocals of the measurement variances. From this follows continuous predictions

$$\hat{\bar{y}}_{track}(t) = \hat{a}_{track} + \hat{b}_{track}\big(u(t) - u(1)\big), \tag{3}$$

143 such that $\hat{\bar{y}}_{track}(t)$ becomes a smoothed and scaled version of the environmental variable.

2.2 Weighted least squares estimation for direct WLS models

145 In the direct WLS method, least squares predictions are found as $\hat{\bar{y}}_{track}(t) = \hat{a}_{WLS} + \hat{b}_{WLS}t$.

146 Here, the parameters are found as in Eq. (2), except that $\boldsymbol{X} = \begin{bmatrix} 1 & 1 & \cdots & 1 \\ t_1 & t_2 & \cdots & t_N \end{bmatrix}^{\boldsymbol{T}}$, where t_i are

147 the sample time points.

2.3 Random walk models

Hunt (2006) developed what he called a general random walk (GRW) model for analyses of

fossil time series, later referred to as a model for directional evolution (Hunt, 2012). This model

assumes a random walk process where at each timestep an increment of evolutionary change

in a mean trait value is drawn at random from a distribution of evolutionary steps, and that this

incremental change is normally distributed with mean value μ_{step} and variance σ_{step}^2. For an

observed evolutionary mean trait change ΔY over T time steps, the log-likelihood function for a

GRW process is given by (Hunt, 2006)

$$\ell\big(\mu_{step}, \sigma_{step}^2\big) = -\frac{1}{2}\ln(2\pi) - \frac{1}{2}\ln\left(T\sigma_{step}^2 + \frac{V_{p_A}}{n_A} + \frac{V_{p_D}}{n_D}\right) - \frac{\big(\Delta Y - T\mu_{step}\big)^2}{2\left(T\sigma_{step}^2 + \frac{V_{p_A}}{n_A} + \frac{V_{p_D}}{n_D}\right)}, \tag{4}$$

where n_A and n_D are the numbers of observed ancestors and descendants, respectively, while

V_{p_A} and V_{p_D} are the corresponding population phenotypic variances. With N irregular and sparse

samples of mean trait values, multiple ancestor-descendant trait differences over $N-1$

evolutionary transitions may be used jointly to estimate μ_{step} and σ_{step}^2 by summing the log-

likelihoods according to Eq. (4) over the transitions, i.e., by use of

$\ell_{total}\big(\mu_{step}, \sigma_{step}^2\big) = \sum_{i=1}^{N-1} \ell_i\big(\mu_{step}, \sigma_{step}^2\big)$. From this expression, step parameters can be found

by means of maximum likelihood estimation, and the estimated step size $\hat{\mu}_{step}$ will then be a

measure of directional change over time. In the real data examples in Section 3, the optimal

GRW parameters are found by use of the MATLAB function *fmincon*, with initial parameter

values sat to zero.

The GRW model has an obvious weakness in that it is difficult to estimate σ_{step}^2 for small

values of T and large values of V_{p_A}/n_A and V_{p_D}/n_D. In the three examples in Section 3 the result

is $\hat{\sigma}_{step}^2 < 0$, such that the step variance must be set to $\hat{\sigma}_{step}^2 = 0$.

Eq. (4) is not intended for predictions, but once $\hat{\mu}_{step}$ is determined with $\hat{\sigma}_{step}^2 = 0$, a

prediction model

$$\hat{\bar{y}}_{GRW}(t) = \hat{a}_{GRW} + \hat{b}_{GRW}t \tag{5}$$

171   is found by use of $\hat{b}_{GRW} = c\hat{\mu}_{step}$, and by fitting a parameter a_{GWR} to the data by a least squares

172   approach. Here, c is a scaling constant that I consistently will set to $c = 1$.

2.4 Weighted mean squared errors

174   Comparisons of responses $\hat{\bar{y}}_{track}(t)$, $\hat{\bar{y}}_{WLS}(t)$ and $\hat{\bar{y}}_{GRW}(t)$ can be done by use of weighted

175   mean squared errors

$$WMSE = (\bar{y} - \hat{\bar{y}})W(\bar{y} - \hat{\bar{y}})/\sum w_i, \tag{6}$$

176   where $\bar{y}$ and $\hat{\bar{y}}$ are $N \times 1$ data vectors, and where W is a diagonal matrix with the weights w_i.

177   Note that we in the special ordinary least squares (OLS) case have $W = I$ and $\sum w_i = N$. I thus

178   assume independent and normally distributed random variables $\bar{y}(t)$. Note that when the

179   weights are the reciprocals of the variances of the measurements, as done here, the prediction

180   slope parameters $\hat{b}_{track}$ and $\hat{b}_{WLS}$ are the best linear unbiased estimates (BLUE).

2.4 Theory for the Akaike Information Criterion

182   Model performances may be compared by use of the Akaike information criterion (AIC), and the

183   model with the lowest AIC value should then be chosen (Hunt, 2006). For GRW, the AIC value

184   with compensation for short data is given by (Hunt, 2006),

$$AIC_{c,GRW} = 2k - 2\ell_{total}\left(\hat{\mu}_{step}, \hat{\sigma}_{step}^2\right) + 2k(k+1)/(N-k-2), \tag{7}$$

185 where $k = 2$ is the number of estimated parameters, while N is the number of samples (the

186 number of transitions is $N - 1$). Since in all the real data examples in Section 3 the estimated

187 variance $\hat{\sigma}_{step}^2$ is negative, the realistic value of $AIC_{c,GRW}$ is found by finding $\hat{\mu}_{step}$ with $\sigma_{step}^2 = 0$.

188   For comparisons with tracking and WLS models, Eq. (7) must be replaced by (Banks and

189 Joyner, 2017)

$$AIC_c = 2k - 2\ln\left(\mathcal{L}\left(\hat{\Theta}_{MLE}|\overline{\mathbf{y}}\right)\right) + 2k(k+1)/(N-k-1), \tag{8}$$

where $\mathcal{L}(\hat{\Theta}_{MLE}|\overline{\mathbf{y}})$ is the likelihood of the estimated parameters using maximum likelihood

estimation, given the vector $\overline{\mathbf{y}}$ of observed data. Banks and Joyner (2017) further assume a

statistical model $Y_i = f(t_i, \mathbf{q}_0) + \omega_i \mathcal{E}_i$, where $\mathbf{q}_0$ is a parameter vector, while ω_i are known

weights. Here, ε_i for $i = 1, 2, \dots, N$ are i.i.d. $\mathcal{N}(0, \sigma^2)$, such that σ^2 must be estimated together

with $\mathbf{q}_0$. In the present setting the weights are $\omega_i = \sqrt{1/w_i}$, and the final AIC expression in

Banks and Joyner (2017) thus becomes

$$AIC = 2k + N\ln(2\pi) + N + 2\sum_{i=1}^{N} \ln\left(\sqrt{1/w_i}\right) + N\ln\left(\frac{\sum_{i=1}^{N} w_i(\overline{y}_i - \hat{\overline{y}}_i)^2}{N}\right),$$
$$\tag{9}$$

from which $AIC_{c,track}$ and $AIC_{c,WLS}$ follow by adding $2k(k+1)/(N-k-1)$. Here, $k = 3$, since

also the underlying parameter σ^2 must be estimated in addition to a_{WLS} and b_{WLS}. Note that $\hat{\overline{y}}_i$

in Eq. (9) is the result of either a tracking model or WLS, and that Eq. (9) therefore is not valid

with use of $\hat{\overline{y}}_{i,GRW}$ as found from Eq. (5).

Two models with known AIC values can be compared by use of Akaike weights (Wagenmakers

and Farrell, 2004). Assuming that that the tracking model is best, as we will find in the real data

examples in Section 3, we first compute

$$\triangle_{WLS} = AIC_{WLS} - AIC_{track}. \tag{10a}$$

203 and

$$\triangle_{GRW} = AIC_{GRW} - AIC_{track}. \tag{10b}$$

204   We then find the Akaike weights

$$A_{w,track} = \frac{1}{1 + exp\left(-\frac{\triangle_{WLS}}{2}\right) + exp\left(-\frac{\triangle_{GRW}}{2}\right)}, \tag{11a}$$

$$A_{w,WLS} = \frac{exp\left(-\frac{\triangle_{WLS}}{2}\right)}{11 + exp\left(-\frac{\triangle_{WLS}}{2}\right) + exp\left(-\frac{\triangle_{GRW}}{2}\right)}, \tag{11b}$$

205   and

$$A_{w,GRW} = \frac{exp\left(-\frac{\triangle_{GRW}}{2}\right)}{1 + exp\left(-\frac{\triangle_{WLS}}{2}\right) + exp\left(-\frac{\triangle_{GRW}}{2}\right)}, \tag{11c}$$

206   The sum of these weights is one, and the weights can be directly interpreted as conditional

207   probabilities for the models (Wagenmakers and Farrell, 2004).

3 Real data cases

3.1 Stickleback fish case

In this case I use data that show how the dorsal fin ray number evolved in *Gasterosteus*

doryssus, an extinct species of a small freshwater stickleback fish that inhabited inland

freshwater habitats of the North American Great Basin during the Miocene (Bell et al., 1985).

See Table A1 in Appendix A for data. In the data analysis the mean trait values are log-

transformed, with the measurement errors transformed accordingly, i.e., $e_{i,log} = e_i \cdot \frac{\bar{y}_{i,log}}{\bar{y}_i}$,

where e_i is found from the given standard deviations SD as $e_i = \sqrt{SD_i^2/n_i}$. The original and log-

transformed samples thus have equal coefficients of variation.

In the tracking model, the environmental driver was $\partial^{18}O(t)$ as found in Westerhold et al.

(2020), Table S34, column marked ISOBENbinned_d18Ointerp. The exact sampling time points

are not given in Bell et al. (1985), but since $t = 0$ in the data is approximately 10 million years

ago (personal information) it was possible to align the data such that the prediction results

were optimized. The result was that $t = 0$ in the data corresponds to $t = -9.994$ million years

with present time as zero point. An optimal window size 96 was also found by a manual search

using raw data from 9.8 to 10.3 million years ago. Since the time resolution is 0.002 million

years, the moving time window size is 0.192 million years.

With σ_{step}^2 as a free variable, the term $T\sigma_{step}^2 + \frac{V_{p_A}}{n_A} + \frac{V_{p_D}}{n_D}$ in Eq. (4) for one or several of the

transitions becomes negative in the course of the parameter search, such that the log-

likelihood becomes a complex number. The search is therefore limited to $\sigma_{step}^2 \geq 0$, resulting in

$\hat{\sigma}_{step}^2 = 0$. See results in Figures 4 and 5, with present time as zero point, and in Table 1.

Figure 4. Plots for the Stickleback fish case. Upper panel shows $\partial^{18}O(t)$ (blue line) and

$u(t) = \partial^{18}O_{smooth}(t)$ with window size 96 (red line). Lower panel shows data points $\bar{y}(i)$

and $u(i) = \partial^{18}O_{smooth}(i)$ with standard error bars, and the estimated linear adaptive

peak function (red line).

Figure 5. Prediction results for the Stickleback fish case, with mean trait observations (circles with standard error bars), tracking predictions (blue line), WLS predictions (red line), and GRW predictions with $\sigma^2_{step} \geq 0$ in the search for the optimum (dashed red line, almost identical to the solid red line).

Table 1. Prediction slope, WMSE and AIC results for predictions in Fig. 5.

	Tracking model	WLS model	GRW model
Prediction slope	—	1.2605	1.2531
WMSE	0.000216	0.000401	0.000401
AIC_c	−137	−121	−104
Akaike weights	0.9997	$3 \cdot 10^{-4}$	$9 \cdot 10^{-8}$

3.2 Ostracod case

Body size data for the deep-sea ostracod (seed shrimp) species *Poseidonamicus major* are found in Hunt and Roy (2006), Table 4, see Table A2 in Appendix A for details. The body size varies from $y = 707$ to $y = 876\ \mu m$, with the variance in the measurements estimated to be

$467/n\ (\mu m)^2$, where n is the number of samples. The standard error is thus $SE = \sqrt{467/n}\ \mu m$,

which on a log scale translates to $SE_{log} = \sqrt{467/n}\ \log(y)\ /y$.

The data in Hunt and Roy (2006) include Mg/Ca temperatures, but a tracking model based on

this information is not any better than the WLS model. Instead, I used the environmental driver

$\partial^{18}O(t)$ as found in Lisiecki and Raymo (2005), and the optimal moving avearage window size

was here 80. In these data the time resolution for $t > -0.6$ million years is 0.001, such that the

moving time window size is 0.08 million years. For $-1.5 < t < -0.6$ the time window size is 0.16

million years, while it for $t < -1.5$ is 0.2 million years. The parameter search was also in this

case limited to $\sigma^2_{step} \geq 0$, resulting in $\hat{\sigma}^2_{step} = 0$. See results in Fig. 6 and Table 2.

Figure 6. Prediction results for Ostracod case, with mean trait observations (circles with

standard error bars), tracking predictions (solid blue line), WLS predictions (red line),

and GRW predictions with $\sigma^2_{step} \geq 0$ in the search for the optimum (dashed red line).

Table 2. Prediction slope, WMSE and AIC results for predictions in Fig. 6.

	Tracking model	WLS model	GRW model
Prediction slope	−	0.0402	0.0316
WMSE	0.000108	0.000241	0.000319
AIC_c	−51	−43	−23
Akaike weights	0.9826	0.0174	$9 \cdot 10^{-7}$

3.3 Bryozoan OV case

Liow et al. (2024) presented results from a field study of the bryozoan species *Microporella*

agonistes, based on a fossil record over 2.3 million years. The scanning electron micrograph in

Fig. 2 shows perimeters of autozooid (AZ) and ovicell (OV) areas, and Fig. 3 shows predictions of

log AZ area mean as found in Ergon (2025). Fig. 7 shows the corresponding results for *log OV*

area mean. As for the results in Fig. 6, the environmental driver was $\partial^{18}O(t)$ as found in Lisiecki

and Raymo (2005), and the optimal moving window size was here 700 (0.7 to 1.75 million years).

The parameter search was also in this case limited to $\sigma_{step}^2 \geq 0$, resulting in $\hat{\sigma}_{step}^2 = 0$. See Table

3 for results.

Figure 7. Prediction results for Bryozoan case 1, with mean trait observations (circles

with standard error bars), tracking predictions (solid blue line), WLS predictions (red

line), and GRW predictions with $\sigma^2_{step} \geq 0$ in the search for the optimum (dashed red

line).

Table 3. Prediction slope, WMSE and AIC results for predictions in Fig. 7.

	Tracking model	WLS model	GRW model
Prediction slope	−	0.1227	0.1805
WMSE	0.0020	0.0021	0.0038
AIC_c	−9	−8	−4
Akaike weights	0.52	0.43	0.05

3.4 Bryozoan AZ case

Finally, and for completeness, I include results from Ergon (2025), as already shown in Fig. 3,

panel B. The moving window size is here 100 (0.1 to 0.25 million years). See results in Fig. 8 and

Table 4. Note that there are some unfortunate errors in the AIC results given in Ergon (2025).

Figure 8. Prediction results for Bryozoan case 2, with mean trait observations (circles

and error bars), tracking predictions (solid blue line), WLS predictions (red line), and

GRW predictions with $\sigma^2_{step} \geq 0$ in the search for the optimum (dashed red line).

Table 4. Prediction slope, WMSE and AIC results for predictions in Fig. 8.

	Tracking model	WLS model	GRW model
Prediction slope	−	0.1519	0.1441
WMSE	0.0010	0.0019	0.0019
AIC_c	−23	−18	−10
Akaike weights	0.9372	0.0613	0.0015

4 Summary and discussion

In this article I use fossil data from three real data cases. I show that in cases with directional

evolution, the adaptive peak tracking models give better results than alternative weighted least

squares (WLS) and general random walk (GRW) models. The model performances are

compared by use of weighted mean squared errors (WMSE) and Akaike information criterion

(AIC) results.

The tracking model in Fig. 1 is in principle multivariate, although the influences from several

environmental factors on the adaptive peak movements in general must be found by causal

modeling that is far beyond the scope of this paper. The adaptive peak may also be influenced

by non-symmetrical fitness functions and developmental constraints as discussed in Ergon

(2025). In the three real data examples, positions of the adaptive peaks are approximately linear

functions of temperature.

The methodological background is given in Section 2, including the theory for GRW models,

and a description of how AIC values are found from WLS based WMSE results. The results for

the various models are summarized in Figures 5, 6, 7 and 8, and Tables 1, 2, 3 and 4. Note that

the $AIC_{c,GRW}$ values do not take the estimation of a_{GRW} in Eq. (5) into account.

In the tracking models I use oxygen isotope data for the environmental drivers. These data

were found in deep sea drilling projects, and they can be converted to global ground

temperatures through linear transformations (Evans et al., 2024). It is especially interesting that

the first case is a study of fossils of a freshwater fish, which verifies that deep sea oxygen

isotope data can be used as proxy for global temperature. This is perhaps the most convincing

case when it comes to tracking model performance (Fig. 5), but it also raises questions as

discussed below.

The reason for use of moving average smoothing in the Bryozoan cases is that the individual

data are collected from geological formations with time spans from 0.019 to 0.203 million

years, and that there is a lot of variation in the raw $\partial^{18}O(t)$ data within each time window

(Ergon, 2025). This is less clear in the Stickleback fish case, where possible errors in the age

data are much smaller. Note, however, that the raw $\partial^{18}O(t)$ data as seen in Fig. 4, upper panel,

are found from deep water drilling samples, and that the local temperature at the field site in

what is now Nevada might have had rapid variations that are not captured in these data.

The stickleback fish case also illustrates an inherent difficulty with the directional evolution

concept. The evolution as shown in Fig. 5 has an overall positive directional trend, but in several

shorter time periods the evolution has a negative trend. This shows that directional evolution is

not a system property, but a property of the environment. The essential system property is thus

tracking, as illustrated in Fig. 1. In the linear mean trait versus environment cases studied here,

the mean trait tracking predictions are simply smoothed and scaled versions of the dominating

environmental variable.

In the Stickleback fish case, the WLS and GRW models gave very similar results (Fig. 5 and

Table 1), while the tracking model was much better. In the Ostracod case, the tracking model

was again clearly the best, while the GWR model gave the poorest results (Fig. 6 and Table 2).

The Bryozoan OV case is special in the sense that tracking and WLS models give quite similar

results (Fig. 7 and Table 3). This is not surprising since the optimal moving window is quite wide.

When instead the autozooid (AZ) area was used, the tracking model was clearly best, while the

WLS and GRW models gave very similar results (Fig. 8 and Table 4).

It would, finally, be interesting to find more linear cases in the literature. It would be even

more interesting to find cases with non-linear mean trait versus environment functions, as

discussed in Ergon (2025). The tracking methodology used here can easily be extended to

handle such cases.

Acknowledgements

336   I thank Mike Bell for information regarding the time span of the stickleback fish data, and

337   University of South-Eastern Norway for support and funding.

Competing interests

339   There are no competing interests.

Data Availability Statement

341   Oxygen isotope data for Figures 4 and 5 are archived as *Data for Stickleback fish case* on

342   bioRxiv, doi: https://doi.org/10.64898/2025.12.07.692820

343   Oxygen isotope data for Figures 6, 7 and 8 are archived as *Raw data for simulations* on bioRxiv,

344   https://doi.org/10.1101/2024.10.30.621046

References

Banks, H.T., and Joyner, M.L. (2017). AIC under the framework of least squares estimation.

Applied Mathematics Letters. http://dx.doi.org/10.1016/j.aml.2017.05.005

Bell, M. A., Baumgartner, J. V., & Olson, E. C. (1985). Patterns of temporal change in single

morphological characters of a Miocene stickleback fish. Paleobiology 11(3): 258-271.

Ergon, R. (2025). Adaptive Peak Tracking as Explanation of Sparse Fossil Data Across

Fluctuating Ancient Environments. Ecology and Evolution. First published: 03 July 2025

https://doi.org/10.1002/ece3.71705

Evans, D., Brugger, J., Inglis, G.N., and Valdes, P. (2024). The Temperature of the Deep Ocean is

a Robust Proxy for Global Mean Surface Temperature During the Cenozoic.

Paleoceanography and Paleoclimatology. doi: 10.1029/2023PA004788

Hunt, G. (2006). Fitting and comparing models of phyletic evolution: random walks and

beyond. Paleobiology 32: 578–601.

Hunt, G. and Roy, K. (2006). Climate change, body size evolution, and Cope's Rule in

deep-sea ostracodes. PNAS, vol. 103/5: 1347–1352.

www.pnas.org/cgi/doi/10.1073/pnas.0510550103

Hunt, G. (2012). Measuring rates of phenotypic evolution and the inseparability of tempo and

Mode. Paleobiology, 38(3), 2012, pp. 351–373. paleo_Hunt_2012_Pb_rates.pdf

Hunt, G., Hopkins, M.J., and Lidgard, S. (2015). Simple versus complex models of trait evolution

and stasis as a response to environmental change. PNAS, vol. 112 /16: 4885–4890.

Lande R (1979) Quantitative genetic analysis of multivariate evolution, applied to

brain:body size allometry. Evolution 33:402-416

www.pnas.org/cgi/doi/10.1073/pnas.1403662111

Lisiecki, L. E., and Raymo, M. E. (2005). A Pliocene–Pleistocene stack of 57 globally distributed

benthic δ18O records. Paleoceanography 20, A1003. doi:10.1029/ 2004PA001071

Liow, L.H., Porto, A., and Di Martino, E. (2024). Trait-Fitness Associations via Fecundity and

Competition in a Two-Million-Year-Long Fossil Record. Am. Nat. 204/3.

https://doi.org/10.1086/73133

Pigliucci, M. (2013). Landscapes, Surfaces, and Morphospaces: What Are They Good For? In

The Adaptive Landscape in Evolutionary Biology, Eds. Svensson, E. and R. Calsbeek. Oxford

University Press. https://doi.org/10.1093/acprof:oso/9780199595372.001.0001

Wagenmakers, E-J., and Farrell, S. (2004). AIC model selection using Akaike weights.

Psychonomic Bulletin & Review, 11 (1), 192-196

Walsh, B., and Lynch, M. (2018). Evolution and selection of quantitative traits.

Oxford University Press

Westerhold et al. (2020). An astronomically dated record of Earth's climate and its

predictability over the last 66 million years. Science 369, 1383–1387.

DOI: 10.1126/science.aba6853

Appendix A Data for real data cases

Table A1. Data for Stickleback fish case with $\bar{y} = dorsal\ fin\ ray\ number$

Time	n	$\bar{y}$	SD
0	30	8.8300	0.7470
0.0053	33	8.7600	0.6630
0.0100	30	8.8700	0.6810
0.0152	36	9.0600	0.7150
0.0193	55	8.5800	0.7620
0.0256	40	8.7500	0.8990
0.0305	75	8.5600	0.7750
0.0340	52	8.7700	0.7030
0.0366	22	9.1800	0.5010
0.0381	28	9.1800	0.7720
0.0410	54	8.9400	0.6270
0.0506	45	9.2000	0.5880
0.0552	35	9.1400	0.6480
0.0603	37	9.0500	0.6640
0.0653	41	9.1000	0.5390
0.0701	25	9.2000	0.7070
0.0751	30	9.1300	0.7300
0.0811	58	9.3100	0.7540
0.0865	32	9.7500	0.7180
0.0920	84	9.6300	0.8030
0.0934	82	10.0200	0.8460
0.0967	89	9.8000	0.8280
0.0994	95	9.6200	0.7320
0.1013	57	9.6700	0.7150
0.1070	45	9.8000	0.6940
0.1077	64	9.8400	0.7180

Table A2. Data for Ostracod case

Age Ma	Mg/Ca temperature	n	Mean valve length μm
3.33	2.54	13	707
2.60	2.23	4	744
2.40	2.15	21	742
2.19	2.07	23	743
1.90	1.96	25	745
1.68	1.88	22	763
0.89	1.55	28	812
0.14	1.19	24	797
0.12	1.18	3	793
0.00	1.12	31	812

389   **Table A3.** Data for OV bryozoan case

Age Ma	$\partial^{18}O(t)$ temperature	log AZ area mean	SE value
2.1895	3.6110	10.2600	0.0650
1.9660	3.6934	10.1700	0.0650
0.9265	4.0426	10.4300	0.0950
0.6485	4.0779	10.3900	0.0550
0.5480	4.0935	10.3600	0.0700
0.5055	4.1210	10.4300	0.0600
0.4510	4.1357	10.5000	0.1350

391   **Table A4.** Data for AZ bryozoan case

Age Ma	$\partial^{18}O(t)$ temperature	log AZ area mean	SE value
2.1895	3.6518	11.3930	0.0500
1.9660	3.7409	11.4230	0.0500
1.8505	3.7487	11.4550	0.1200
0.9265	4.0233	11.6150	0.0500
0.6485	4.2053	11.6600	0.0400
0.5480	4.0763	11.5550	0.0450
0.5055	4.1545	11.6550	0.0400
0.4510	4.1775	11.7500	0.0850
0.3990	4.1735	11.7000	0.1200

Appendix B MATLAB code for Stickleback fish case

```matlab
clear
load Data_9800_to_10300.mat
Age=flipud(Age);
d18O=flipud(d18O);
tdata=-Age';        % Step size 0.002 My
rawdata=d18O';
window=96;
u=movmean(rawdata,window);
t0=-10+0.0060;

%% Sample data
tlabel1=[0 5317 9950 15182 19318 25551 30541 34030 36631 38052 41008 50629 55246];
tlabel2=[60289 65312 70119 75075 81150 86454 92006 93445 96659 99377 101306 106969 107667];
tlabel=[tlabel1 tlabel2]/1000000;
tlabel=t0+tlabel;
n=[30 33 30 36 55 40 75 52 22 28 54 45 35 37 41 25 30 58 32 84 82 89 95 57 45 64];
y1=0.01*[883 876 887 906 858 875 856 877 918 918 894 920 914 ];
y2=0.01*[905 910 920 913 931 975 963 1002 980 962 967 980 984];
y=[y1 y2];
sd1=0.001*[747 663 681 715 762 899 775 703 501 772 627 588 648  ];
sd2=0.001*[664 539 707 730 754 718 803 846 828 732 715 694 718];
sd=[sd1 sd2];
var=sd.^2;
err0=sqrt(var./n);
ylabell=log(y);
for i=1:26
  err(i)=err0(i)*(ylabell(i)/y(i))^1;
end

%% Tracking model
tplot=tdata;
ulabel=zeros(1,length(tlabel));
tlab=tlabel;
for i=1:length(tlabel)
```

```matlab
429      for t=1:length(tplot)
430        if tplot(t)>min(tlab)
431          ulabel(1,i)=u(1,t);
432          tlab(1,i)=0;
433          break
434        end
435      end
436    end
437    w=err.^-2;
438    W=diag(w);
439    N=26;
440    k=3;
441    X=[ones(length(tlabel),1) ulabel'-2.9559*ones(length(tlabel),1)];
442    bls1=inv(X'*W*X)*X'*W*ylabell'
443    a1=bls1(1);
444    b1=bls1(2);
445    yhat1=a1+b1*(ulabel-2.9559*ones(1,26));
446    for t=1:length(u)
447      yhatplot(t)=a1+b1*(u(t)-2.9559);
448    end
449    WMSE_Tracking=sum((ylabell-yhat1)*W*(ylabell-yhat1)')/sum(diag(W))
450    for j=1:N
451      lnw(j)=log(sqrt(1/w(j)));
452      z(j)=(ylabell(j)-yhat1(j)).^2*w(j);
453    end
454    AICc_Tracking=2*k+N*log(2*pi)+N+2*sum(lnw)+N*log(sum(z)/N)+2*k*(k+1)/(N-k-1);
455
456    %%% WLS model
457    X=[ones(26,1) tlabel'];
458    bls=inv(X'*W*X)*X'*W*ylabell';
459    a2=bls(1);
460    b2=bls(2);
461    for i=1:26
462      yhat2(i)=a2+b2*tlabel(i);
463    end
464    WMSE_WLS=(ylabell-yhat2)*W*(ylabell-yhat2)'/sum(diag(W))
465    for j=1:N
```

```matlab
466         lnw(j)=log(sqrt(1/w(j)));
467          z(j)=(ylabell(j)-yhat2(j)).^2*w(j);
468     end
469     AICc_WLS=2*k+N*log(2*pi)+N+2*sum(lnw)+N*log(sum(z)/N)+2*k*(k+1)/(N-k-1);
470
471     %%% GRW model
472     k=2;
473     N=25;
474     n_samples=ones(1,26);
475     var_samples=err.^2;
476     Tsample=tlabel;
477     for i=1:25
478        dT(i)=-Tsample(i)+Tsample(i+1);
479        dX(i)=ylabell(i+1)-ylabell(i);
480        nA(i)=n_samples(i);
481        nD(i)=n_samples(i+1);
482        varA(i)=var_samples(i);
483        varD(i)=var_samples(i+1);
484     end
485
486     % Contraints and
487     mustep_min=-10; mustep_max=10;
488     varstep_min=0; varstep_max=1;
489     par_lb=[mustep_min varstep_min];
490     par_ub=[mustep_max varstep_max];
491     par_guess=[0 0];
492     Aineq=[]; Bineq=[]; Aeq=[]; Beq=[];
493     fun_objective_handle=...
494        @(par)fun_objective(par,dT,dX,varA,nA,varD,nD);
495     [par_opt,fval,exitflag,output,lambda,grad,hessian] =...
496     fmincon(fun_objective_handle,par_guess,Aineq,Bineq,Aeq,Beq,par_lb,par_ub);
497     mustep=par_opt(1);
498     varstep=par_opt(2);
499     for i=1:25
500        varterm(i)=dT(i)*varstep+varA(i)/nA(i)+varD(i)/nD(i);
501     end
502     for i=1:25
```

```
503    Li(i)=-log(2*pi)/2-log(varterm(i))/2-(dX(i)-dT(i)*mustep)^2/(2*varterm(i));

504    end

505    L=sum(Li);

506    AICc_GRW=2*k-2*L+2*k*(k+1)/(N-k-1);

507    X=ones(26,1);

508    b3=mustep;

509    a3=inv(X'*W*X)*X'*W*(ylabell'-b3*tlabel');

510    for i=1:26

511      yhat3(i)=a3+b3*tlabel(i);

512    end

513    WMSE_GRW=sum((ylabell-yhat3)*W*(ylabell-yhat3)')/sum(diag(W));

514

515    % Akaike weights

516     D_Tracking=0;

517     D_WLS=AICc_WLS-AICc_Tracking;

518     D_GRW=AICc_GRW-AICc_Tracking;

519     L_WLS=exp(-0.5*D_WLS);

520     L_GRW=exp(-0.5*D_GRW);

521     Aweight_Tracking=1/(1+L_WLS+L_GRW);

522     Aweight_WLS=L_WLS/(1+L_WLS+L_GRW);

523     Aweight_GRW=L_GRW/(1+L_WLS+L_GRW);

524

525

526    yplot=a1+b1*(ulabel'-2.95*ones(26,1));

527

528    figure(4)

529    subplot(2,1,1)

530    plot(tplot,rawdata,'b','LineWidth',1.0), hold on

531    plot(tplot,u,'m','LineWidth',1.0), hold off, grid

532    axis([-10 -9.88 2.5 3.3])

533    xlabel('Million years')

534    ylabel('d^1^8O and d^1^8O_s_m_o_o_t_h')

535    subplot(2,1,2)

536    errorbar(ulabel,ylabell,err,'bo','LineWidth',1.0), hold on, grid

537    plot(ulabel,yplot,'m','LineWidth',1.0), hold off

538    axis([2.95 3.03 2.12 2.33])

539    xlabel('d^1^8O_s_m_o_o_t_h')
```

```matlab
540    ylabel('log mean dorsal fin ray number')
541
542    figure(5)
543    errorbar(tlabel,ylabell,err,'bo','LineWidth',1.0), hold on
544    plot(tplot,yhatplot,'b','LineWidth',1.0)
545    plot(tlabel,yhat2,'m');
546    plot(tlabel,yhat3,'m--','LineWidth',1.0)
547    hold off, grid
548    axis([-10 -9.88 2.12 2.33])
549    xlabel('Million years')
550    ylabel('log mean dorsal fin ray number')
551
552    b2
553    b3
554    WMSE_Tracking
555    AICc_Tracking
556    WMSE_WLS
557    AICc_WLS
558    WMSE_GRW
559    AICc_GRW
560    Aweight_Tracking
561     Aweight_WLS
562     Aweight_GRW
563
564
565    %%% Parameter search
566    function f = fun_objective(par,dT,dX,varA,nA,varD,nD)
567    mustep=par(1);
568    varstep=par(2);
569    for i=1:25
570      varterm=dT(i)*varstep+varA(i)/nA(i)+varD(i)/nD(i);
571      Li(i)=-log(2*pi)/2-log(varterm)/2-(dX(i)-dT(i)*mustep)^2/(2*varterm);
572    end
573    L=sum(Li);
574    f=-L;
575    end
576
```