Deferred mortality: cyclic thermal stress during pupation triggers irreversible carry-

over effects in a key pollinator

Maximilian M. Mandlinger[1] and Christoph Kurze[1]*

[1] Evolutionary Ecology, University of Regensburg, Germany

* Corresponding author, email: Christoph.Kurze@ur.de

Abstract

Extreme heatwaves threaten global biodiversity, yet the sublethal carry-over effects of

developmental thermal stress on adult fitness remain poorly understood. We addressed

this knowledge gap for bumblebees, key pollinators that face alarming population

declines, using a controlled *in vitro* approach. We exposed *Bombus terrestris* pupae to

cyclic thermal stress to evaluate emergence success, adult longevity, and morphological

traits. Cyclic thermal stress caused significant acute pupal mortality (up to 30% reduced

emergence) and, crucially, deferred mortality, significantly reducing adult longevity

(hazard ratio, HR = 2.00) in both workers and males. Rearing *in vitro* at 34°C showed no

significant difference compared to individuals that developed naturally in their natal colony

(HR = 0.95). Antennal and wing deformations served as powerful hazard indicators (HR =

2.38) for premature adult mortality. Our findings reveal that cyclic thermal stress during

pupal development imposes irreversible developmental damage, undermining adult

physiological resilience.

Keywords: climate change, metamorphosis, lifespan, deformations, malformations, body

size, morphometrics, bees

1. Introduction

Extreme weather events, such as heatwaves, are increasing in frequency and intensity

due to climate change, leading to distinct and widespread implications for biodiversity,

including insects [1, 2]. While thermal stress is known to have lethal and sublethal effects

across a wide range of insects, most studies have focused on acute effects in adults, such

as mortality at critical thermal maxima, metabolic disruption, and impaired flight

performance and cognition [3-5]. Little is known about how thermal stress during an earlier

life stage impose sublethal, yet long-lasting, carry-over effects on adult fitness [6-9].

Gaining a better understanding of such sublethal carry-over effects, impacting colony

fitness, is essential for accurately forecasting population declines in keystone insect taxa

such as bumblebees (*Bombus* spp., Hymenoptera: Apidae) [10].

Bumblebees, as a cold-adapted taxon, have experienced population declines,

which are partially attributed to climate change [11-16]. This is a concerning trend, given

that they provide crucial pollination services in ecosystems and agriculture [17, 18]. While

ground-nesting species can mitigate extreme ambient temperatures to some extent, other

species, such as *B. hypnorum* and *B. pascuorum* that typically nest above the ground

[19], or commercially used species in greenhouses such as *B. terrestris* [20], face a higher

risk of thermal stress. Colony-level studies have demonstrated that constant prolonged

elevated temperatures negatively impact colony energetics, growth and reproductive

fitness [21-24], and such thermal stress has been linked to altered offspring morphology

[25-27]. As behavioural responses vary widely between colonies, it is difficult to

disentangle indirect effects arising from worker thermoregulation [28, 29], from direct

physiological effects [30]. For example, if workers need to spend more time fanning to

cool the nest or incubating the brood to maintain optimal brood temperatures (generally

above ambient nest temperatures) rather than foraging and nursing [28, 29], this can

affect larval development and ultimately colony fitness. To gain a precise mechanistic

understanding of how thermal stress affects specific larval stages, *in vitro* rearing offers

the required control at the cost of losing natural realism [30, 31].

Our previous *in vitro* studies have already revealed stage-specific acute responses

to constant thermal stress in *B. terrestris* larvae and pupae [30, 31]. While smaller fourth-

instar (L4) larvae may have a selective advantage under thermal stress due to their ability

to more easily reach the critical body mass for pupation [31], thermal stress at the pupal

stage appears to select for individuals with increased relative lipid reserves [30]. A

constant *in vitro* rearing temperature of 38°C during L4 larval and pupal development was

identified as a critical thermal threshold, resulting in an increased direct mortality and

higher a incidence of wing deformations [30, 31]. While these studies illustrate impaired

larval and pupal development at constant thermal stress, the magnitude of the

downstream impact on adult survival and the effect of fluctuating thermal stress scenarios

remains unknown.

To address this fundamental knowledge gap, we used highly controlled *in vitro*

rearing of *B. terrestris* pupae to isolate the effect of cyclic thermal stress scenarios on

individual life history. Our main aim was to assess the carry-over effects under fluctuating

thermal extremes, an approach more ecologically relevant than constant temperature

regimes [21, 30, 32, 33]. We focused on the pupal stage as metamorphosis in

holometabolous insects, such as bees, represents the most extreme transition between

phenotypes [34] and is likely the most vulnerable to environmental stress [30, 35]. We

quantified both the direct effect on emergence success and the subsequent long-term

adult survival, morphology, and energy reserves. We hypothesised that increasing cyclic

thermal stress during pupal development (metamorphosis) would affect adult morphology

(specifically wing morphology [30, 35]) and trigger deferred mortality, significantly reducing

adult longevity. Furthermore, we expected that the impact of thermal stress would differ

between workers and males.

2. Materials and methods

2.1 Colony husbandry

We maintained ten queen-right source colonies of *B. terrestris* (Natupol Research Hives, Koppert B.V., Netherlands) under standardized laboratory conditions [36]. Colonies were housed in standard nest boxes (23 x 25 x 15 cm), which were connected to foraging arenas (59 x 39 x 26 cm). We maintained all colonies at 25 ± 1°C and 30-50% RH, under a 14:10 h light:dark regime. Bumblebees had *ad libitum* access to a 66% w/w sucrose solution (Natupol Bee Happy, Koppert B.V., Netherlands). We supplied each colony with 6-11 g of pollen candy (67% w/w organic pollen, 25% w/w sucrose, 8% w/w water) daily.

2.2 Larvae collection and *in vitro* rearing

We collected 4th instar (L4) larvae from fully developed source colonies (for sample size see Table 1; Fig. S1*a*). Collecting L4 larvae was crucial to ensure that we could start all thermal regimes consistently at the beginning of the pupal phase. Larvae were reared *in vitro* following previous protocols [for details see 30, 31, 37].

Briefly, L4 larvae were carefully transferred into 3D-printed polylactide (PLA) artificial brood cells (Fig. S1*b*), weighed (initial larval weight; d = 0.1 mg, Sartorius M-Pact AX224, Sartorius GmbH, Germany), then placed inside 24- or 48-well clear flat bottom plates (Fig. S1*c*; Falcon/Corning, USA) and put within a humidity-controlled plastic container (18.5 x 18.5 x 11.5 cm) kept at a constant temperature of 34°C (KB115, BINDER GmbH, Germany). While 34°C is at the upper range of the typical nest temperature in *B. terrestris* [19], it aligns with brood temperatures, which are typically 2°C warmer than ambient nest temperatures [28] and likely represent the optimal temperature for larval development [29]. Moreover, 34°C has been used in previous *in vitro* rearing protocols for *B. terrestris* [30, 31, 37], making it comparable to previous *in vitro* studies. To maintain a relative humidity (RH) of 65 ± 10%, a 120 mL cup of saturated sodium chloride solution was also placed within the container.

Larvae were fed a pollen medium (Fig. S1*b*), consisting of 50% w/v sucrose

solution, 40% organic pollen, 10% yeast extract (Bacto™, BD, USA), and 1% casein

sodium salt from bovine milk (Sigma-Aldrich, Germany) [37], twice daily to satiation [31].

Feeding was stopped with the beginning of the pupal phase. Our setup allowed us to

measure pupal weight non-invasively and facilitated the easy removal of any dead

individuals.

2.3 Experimental procedure

At the beginning of pupation (metamorphosis), each individual was weighed (initial pupal

weight; Sartorius M-Pact AX224) and randomly assigned to one of three experimental

treatment groups (for sample sizes see Table 1). Pupae were transferred to new,

humidity-controlled containers within one of three programmed incubators. The control

group was maintained at a constant temperature of 34°C (34°C, constant *in vitro* control).

The two cyclic thermal stress treatment groups experienced a 12:12 h temperature cycle

between a baseline of 34°C (night phase) and a thermal maximum (day phase) of either

37°C (37°C cyclic) or 38°C (38°C cyclic) until adult emergence (i.e. 9-10 days for males,

8-9 days for workers; Fig. 1*c*). Given that bumblebee species regulate their core nest

temperatures between 30-35°C for optimal brood development [19], our two cyclic

temperature treatments simulate a thermal stress scenario when colony thermoregulation

fails during the day. Pupal survival and adult emergence were monitored blindly twice

daily. In addition to our *in vitro* treatments, newly emerged adults (< 1 day old) were

collected directly from their natal colonies to serve as an additional control group (colony-

reared) for the subsequent longevity assay.

2.4 Adult survival assay

Newly emerged adult bees from each thermal treatment (i.e. those that experienced as

pupae *in vitro*) and the additional colony-reared group (for sample sizes, see Table 1)

were weighed (emergence weight; Sartorius M-Pact AX224) and then individually housed

in modified 50 mL tubes (Fig. S1*d*; Greiner Bio-One GmbH, Germany). These were stored

in a wooden rack within the insectary room housing their natal colonies at 25 ± 1°C and

30-50% RH (Fig. S1*e*). Each tube contained a 3D-printed PLA mesh insert for sanitary

reasons. The lids were modified to include three 3 mm ventilation holes and one 1 cm

feeding port, through which a 1.5 mL perforated tube provided *ad libitum* access to a 60%

w/v sucrose solution. Pollen candy was offered separately in the caps of 2 mL reaction

tubes and was replaced weekly or as necessary. Housing tubes and meshes were

replaced as needed to ensure sanitary conditions. Adult survival was monitored around

noon daily. Dead individuals were recorded and immediately frozen at -21°C until further

analysis.

Table 1. Overview of the experimental design. This table details the rearing conditions (regime and

temperature) of pupae for all treatment groups, including individually developed naturally in their natal colony.

Sample sizes (n) reflect the total number of pupae per treatment at start of *in vitro* rearing experiment

(indicated by §) and emerged adults at the start of the survival assay (indicated by $^\#$). N indicates the number

of colonies used in each treatment group.

Treatment	Regime§	Temperature§	Sample size (n)	N
colony-reared	developed naturally	25 ± 1°C*	$^\#$115	10
34°C	constant	34°C	§132/ $^\#$102	6
37°C cyclic	12:12 h cycle	37°C:34°C	§125/ $^\#$82	5
38°C cyclic	12:12 h cycle	38°C:34°C	§146/ $^\#$78	5

* colonies kept in the insectary
§ pupae / $^\#$ adults

2.5 Morphological and physiological analysis

We conducted the morphological and physiological analyses on adult workers and males,

which were exposed to different thermal scenarios as pupae (Table 1), following our

previous protocols [30, 31, 36]. In addition to their emergence weight, the intertegular

distance (ITD) and head width, used as proxies for adult body size, were measured on

dead bees from the survival assay using a digital microscope (VHX-500F, Keyence

GmbH, Germany). The sex of each adult bee was determined by counting flagellomeres

(workers = 10; males = 11). Furthermore, any deformations of their antennae were

recorded (Fig. S2*a*). To assess wing integrity, forewings were carefully detached,

embedded side-by-side on a microscope slide (Roti®Mount, Carl Roth GmbH + Co. KG,

Germany) and photographed at 20x magnification. Wings were subsequently classified as

deformed (figure S2*b*) when visually apparent malformations were present.

The remaining bee bodies were processed to determine dry mass and relative lipid

content following our previous protocol. Specimens were first cut open from the stinger to

the fourth sternite and then dried at 60°C for three days (U40, Memmert GmbH & Co. KG,

Germany). Subsequently, their initial dry mass, including body lipids, was measured

(Sartorius M-Pact AX224), after which lipids were extracted using a petroleum ether

extraction method [30, 38]. For this, dried specimens were placed in 5 mL petroleum ether

for five days. Afterwards, the ether containing lipids was discarded and specimens were

rinsed with fresh ether. The specimens were re-dried for another three days and weighed

again post-extraction, allowing us to calculate the difference between the initial and post-

extraction dry mass. The relative lipid content, serving as a proxy for fat body size and

energy reserves, was calculated as absolute lipid content divided by initial dry mass.

2.6 Statistics and reproducibility

All statistical analyses and data visualizations were performed in R version 4.5.1 [39]. For

full transparency and reproducibility, the complete code as including model selection

procedures (provided as R Markdown output) and the dataset are available in the Zenodo

Repository (https://doi.org/10.5281/zenodo.18298098). Experimental reproducibility was

ensured by balancing initial sample sizes (pupae at the start of the experiment) across

treatments and colonies (Table 1). Specifically, we used individuals from 5-6 colonies per

treatment group, while naturally developed individuals were taken from 10 sources

colonies (including four additional colonies), defining colonies as the primary unit of

biological replication. We accounted for colony variability in all statistical models described

in detail below.

To analyse the effect of cyclic thermal stress, while accounting for the effects of in

vitro rearing by including data from colony-reared individuals where possible, and

additional explanatory variables (e.g. sex, initial pupal weight) and their interactions on

adult emergence rates, adult morphological traits (antennae and wing deformation,

emergence weight, and head width), and relative lipid content, we fitted generalized linear

mixed effect models (GLMMs) using the *glmmTMB* package [40]. Emergence rates and

deformation status were fitted with binomial distribution (logit link), while the continuous

variables proportional pupal weight loss, emergence weight, head width, and relative lipid

content were fitted with a gamma distribution (log link). To analyse the duration of pupal

development (number of days from the start of pupation to adult emergence), we applied

an ordinal cumulative link mixed model (CLMM) with the Laplace approximation using the

ordinal package [41], as assumptions for fitting a GLMM were not met (described below).

While the CLMM included only colony ID as a random factor, GLMMs utilising only data

from the *in vitro* rearing also included collection date as a random factor to account for

colony and temporal-specific variability.

For each response variable, a set of candidate models was compared. The most

parsimonious and best-fitting model was selected based on the Akaike information

criterion (AIC), with lower AIC values indicating a better fit. The best-fitting models were

compared to a simpler nested model using likelihood ratio tests (LRTs). When added

terms did not improve model fit significantly, the simpler model was chosen and compared

against its respective null-model using LRT. Model assumptions and dispersion of the

data were checked using the *DHARMa* package [42]. When model assumptions were

met, statistical significance (p < 0.05) was determined using the *Anova* function. Pairwise

comparisons between treatment groups were conducted using the *emmeans* package [43]

with Holm-Bonferroni corrections. Smooth predictions of emergence rate across the

observed range of larval weights were generated using a prediction dataset with 100

evenly spaced values.

Differences in survival probability between treatment groups were initially

determined for workers and males separately using Kaplan-Meier analyses via the

survival and *ggsurvfit* packages [44], following to previous research [45]. To further

analyse the main factors affecting adult survival, we fitted a set of candidate Cox

Proportional Hazards (CoxPH) models using the *coxme* package [44]. The full model

contained treatment (including data from individuals that developed naturally), sex (and

their interactions), adult dry mass, relative lipid content, and deformations (antennae and

wing deformation combined). The colony ID was included as a random factor. Model

selection followed the AIC and LRT procedure as described above. The final best-fitting

model to predict mortality included the explanatory variables treatment, sex, and their

interactions, alongside deformation status. As proportional hazards (PH) assumptions

were not entirely met (based on diagnostics using the *ggcoxdiagnostics* function of the

surmiver package [46]), the final model was stratified by sex. Pairwise comparisons

between treatment groups were performed using the *glht* function with Holm-Bonferroni

correction.

3. Results

3.1 Emergence success and developmental duration

Cyclic thermal stress and pupal weight significantly affected the proportion of pupae

reaching adulthood (pupal survival; GLMM: x^2 = 35.187, df = 3, p < 0.0001; Fig. 1*a,b*).

This effect was primarily driven by the increase in thermal stress (x^2 = 22.276, df = 2, p <

0.0001; Fig. 1*a*). Pairwise comparisons revealed that pupae exposed to 37°C and 38°C

cyclic thermal regimes had a 15% and 30% lower adult emergence rate, respectively,

compared to pupae reared at constant 34°C (37°C vs 34°C: p < 0.05; 38°C vs 34°C: p <

0.0001). The difference between the 38°C and 37°C cyclic treatments was also significant

(37°C vs 38°C: p < 0.05).

Emergence probability was significantly higher with an increase in initial pupal

weight (x^2 = 11.065, df = 1, p < 0.001; figure 1*b*). Despite the marginally significant effect

of sex on proportional pupal weight loss (GLMM: x^2 = 3.777, df = 1, p = 0.052; Fig. 1*d,*

absolute pupal weight loss data are provided in figure S3*b*), including treatment did not

improve model fit (LRT: x^2 = 0.904, df = 2, p = 0.904). Additionally, there was a strong

positive relationship between adult emergence weight and the initial pupal weight (GLMM:

x^2 = 402.03, df = 1, p < 0.0001; Pearson's correlation: r = 0.88, p < 0.0001; Fig. S3*c*) with

a small but significant effect of sex (x^2 = 4.03, df = 1, p = 0.045), whereas the interaction

between initial pupal weight and sex had no effect (x^2 = 2.169, df = 1, p = 0.141).

Including treatment did not significantly improve model fit (LRT: x^2 = 6.205, df = 3, p =

0.102)

While we observed a significant overall effect on pupation duration (CLMM: x^2 =

43.958, df = 3, p < 0.0001; Fig. 1*c*), this effect was not related to the thermal stress

treatment (x^2 = 1.6121, df = 2, p = 0.447), but was sex-dependent (x^2 = 34.311, df = 1, p <

0.0001). Pairwise comparisons revealed that workers across all treatment groups were

significantly different from males of all treatment groups (p < 0.001).

Figure 1. Direct effects of cyclic thermal stress during pupation on emergence success and development duration in *B. terrestris*. (a) Proportions of pupae emerging as adult bees observed per colony (grey points; n ≥ 5 per colony) and GLMM means ± 95% CI (bars). Sample sizes (n) across all colonies are provided below for each treatment group. **(b)** Predicted probability of emergence (± 95% CI, shaded areas) in relation to pupal weight and treatment. **(c)** Pupation duration until emergence for males (open violins) and workers (filled violins). **(d)** Proportional pupal weight loss from the beginning of pupation until emergence as adult males (open boxes) and workers (filled boxes). Colours **(a-d)** indicate different *in vitro* rearing conditions for pupae reared at constant 34°C (blue), and under cyclic thermal stress regimes of 37°C (orange) and 38°C (reddish-purple). Individual data points in **(b-d)** are shown as triangles (males, m) and circles (workers, w). Sex-specific samples sizes for 34°C ($n_m = 67$, $n_w = 35$), 37°C cyclic ($n_m = 58$, $n_w = 24$), and 38°C cyclic ($n_m = 60$, $n_w = 18$). Significant differences ($p < 0.05$), based on post-hoc comparisons from GLMMs, are indicated by different letters above the bars and next to the violin plots (groups sharing a letter are not significantly different).

3.2 Carry-over effects on adult longevity

Cyclic thermal stress exposure during pupation resulted in reduced adult survival

(longevity) for both workers (Log-rank test: χ^2 = 8.9, df = 2, p = 0.01; Kaplan-Meier

survival curves in Fig. 2a) and males (χ^2 = 11.1, df = 2, p < 0.01; Fig. 2b). The presence of

carry-over effects on longevity was strongly supported by the CoxPH model stratified by

sex (χ^2 = 517.34, df = 4, p < 0.0001, model concordance = 0.64 ± 0.01 SE; Fig. 2e).

Pupae exposed to 38°C cyclic treatment had a significantly higher mortality risk as adults

than pupae reared at 34°C (HR = 2.00, 95% CI = 1.61-2.48, p < 0.0001, Fig. 2e). The

37°C cyclic treatment during pupal development posed a marginally significant higher

mortality risk during adulthood (HR = 1.39, 95% CI = 1.024-1.89, p = 0.035, Fig. 2e). We

also found that individuals with morphological deformations (combining deformed

antennae and wings) had a significantly higher mortality risk (HR = 2.38, 95% CI = 1.94-

2.92, p < 0.0001, Fig. 2e). Importantly, we found no evidence that mortality risk during

adulthood significantly differed between individuals reared *in vitro* as L4 larvae and pupae

at 34°C and those developed naturally in their natal colony until adult emergence (HR =

0.95, 95% CI = 0.70-1.30; p = 0.756, Fig. 2e). This was consistent with the survival

analyses in both adult workers (Log-rank test: χ^2 = 0, df = 1, p = 0.9, Fig. 2c) and males

(χ^2 = 0.8, df = 1, p = 0.4, Fig. 2d).

Figure 2. Carry-over effects of cyclic thermal stress during pupation on adult longevity in *B. terrestris*.
Kaplan-Meier survival curves (± 95% CI, shaded areas) showing the carry-over (indirect) effect of thermal treatments during pupation *in vitro* on the longevity in adult workers (***a***) and males (***b***). The group kept at a constant 34°C (blue) is compared to the cyclic thermal stress treatments (37°C cyclic, orange; 38°C cyclic, reddish-purple). Comparison of *in vitro* rearing of L4 larvae and pupae at 34°C (blue) to those naturally developed in their natal colony (bluish green) until emergence on adult survival in workers (***c***) and males (***d***), respectively. Sample sizes (n) and overall statistical significance (p) from the Kaplan-Meier analyses are provided in each plot. **(e)** Forest plot of the hazard ratios (HR) ± 95% CI based on the stratified CoxPH model for treatment (cyclic thermal stress during pupal development) and morphological deformations (combined antennae and wings deformations; Fig. S2) on adult longevity. HR values > 1 indicate significantly increased hazard. Asterisks denote significance (* p < 0.05; *** p < 0.001) relative to the 34°C group (dashed blue line).

3.3 Morphology and energy reserves

We found evidence that cyclic thermal stress during pupal development increased the

incidence of deformed wings in adults (GLMM: x^2 = 20.383, df = 3, p = 0.0001; Fig. 3*a,*

filled bars; Fig. S2*b*). Pupae exposed to cyclic 38°C developed deformed wings more

frequently than pupae kept at 34°C (38°C vs 34°C: 3.6-fold increase, p < 0.01) and those

reared inside their natal colony (38°C vs colony: 4.5-fold increase, p < 0.001), while there

was no significant difference compared to pupae exposed to cyclic 37°C (38°C vs 37°C: p

= 0.155). Nonetheless, there was no significant difference between pupae exposed to

cyclic 37°C compared to pupae kept at 34°C (37°C vs 34°C: p = 0.294) or those reared

inside their natal colony (37°C vs colony: p = 0.169). There was no significant difference in

the incidence of deformed wings between pupae reared *in vitro* at 34°C compared to

colony-reared individuals (34°C vs colony: p = 0.636). Furthermore, there was no

significant effect of thermal treatment during pupal development on the incidence of adult

bees with deformed antennae (x^2 = 6.241, df = 3, p = 0.100; Fig. 3*a,* open bars; Fig. S2*a*).

We found a significant effect on adult emergence weight (GLMM: x^2 = 158.96, df =

4, p < 0.0001; Fig.3*b*), which was both treatment-dependent (x^2 = 82.129, df = 3, p <

0.0001) and sex-dependent (x^2 = 94.806, df = 1, p < 0.0001) without an interaction effect.

Pairwise comparisons revealed that the emergence weight of adults reared *in vitro* during

larval and pupal development was lighter compared to adults that emerged in their natal

colony (colony vs all *in vitro* treatments: p < 0.0001), and males were significantly heavier

than workers across all treatment groups (males vs workers: p < 0.0001).

Further morphometric analyses revealed that head width strongly increased with

emergence weight (GLMM: x^2 = 1403.577, df = 1, p < 0.0001; Fig. 3*c*). This relationship

was also both treatment-dependent (x^2 = 50.95, df = 3, p < 0.0001) and sex-dependent

(x^2 = 40.242, df = 1, p < 0.0001) with a weak three-way interaction (weight*treatment*sex:

x^2 = 15.966, df = 3, p < 0.001), but no significant two-way interactions (weight*treatment:

x^2 = 2.102, df = 3, p = 0.552, weight*sex: x^2 = 0.417, df = 1, p = 0.518, treatment*sex: x^2 =

0.861, df = 3, p = 0.835). However, the scaling relationship between emergence weight

and head width remained largely consistent across treatments and sexes, with slopes

ranging from 0.17 to 0.27 and no significant differences between slopes (all interactions p

> 0.05, detailed comparisons in Table S1; Fig. 3*c*).

There was also a strong morphometric relationship between head width and

intertegular distance (ITD) measured in adults (GLMM: χ^2 = 567.94, p < 0.0001; Pearson's

correlation: r = 0.87, Fig. S3*d*), with similar scaling relationships between sexes (both

slopes = 0.353) despite males being larger than workers (χ^2 = 5.43, p < 0.05). Including

treatment did not significantly improve the model fit (LRT: χ^2 = 6.205, p = 0.102).

Analysis of relative lipid content (as a proxy for energy reserves) at the end of

adult life revealed a significant effect of sex (GLMM: χ^2 = 4.27, df = 1, p = 0.039, Fig. 3*d*),

which was independent of treatment, as including treatment did not improve the model fit

(LRT: χ^2 = 0.923, df = 2, p = 0.631).

Figure 3. Effects of cyclic thermal stress during pupation on morphological and physiological traits in adult _B. terrestris_. (a) Proportion of individuals per colony (n ≥ 5 per colony) with deformed antennae (open circles; Fig. S2*a*) and wings (filled grey circles; figure S2*b*), and the GLMM means ± 95% CI (bars). Sample sizes (n) are provided below for each treatment group. Colours (*a-d*) indicate different rearing conditions *in vitro* (treatments during pupal development: blue = 34°C; orange = 37°C cyclic; reddish-purple = 38°C cyclic) and inside their natal colony (bluish green). **(b)** Box plots showing the weight of newly emergence adults (males = open boxes; workers = filled boxes) for each treatment. **(c)** Scaling of adult head width relative to emergence weights for adults from each treatment group. Lines (dashed = males; dotted = workers) show fitted trends ± 95% CI from GLMs (ignoring colony effects) with log-transformed emergence weight. **(d)** Box plots showing the relative lipid content (proxy for remaining energy reserves) in adults at the end of their lifespan for each treatment group. Individual data points in **(b-d)** are shown as triangles (males, m) and circles (workers, w). Sex-specific samples sizes for 34°C (n_m = 67, n_w = 35), 37°C cyclic (n_m = 58, n_w = 24), 38°C cyclic (n_m = 60, n_w = 18), and colony-reared (n_m = 74, n_w = 41). Significant differences (p < 0.05), based on post-hoc comparisons from GLMMs, are indicated by different letters above the bars and boxes (groups sharing a letter are not significantly different).

4. Discussion

Our study addresses the knowledge gap concerning the lethal and sublethal, long-term carry-over effects of thermal extremes in insect pollinators [3, 4, 10]. We provide empirical evidence that thermal stress during pupal development leads to deferred mortality at the adult life-stage in bumblebees. This finding reveals a mechanism of deferred mortality that may contribute significantly to pollinator decline. Specifically, cyclic thermal stress not only caused acute pupal mortality (i.e. reduced emergence; Fig. 1*a*) but also resulted in significantly reduced adult longevity (HR = 2.00) in both workers and males of *B. terrestris* (Fig. 2*a,b,e*). This finding suggests that thermal stress during pupation imposes irreversible developmental damage that undermines adult physiological resilience.

Developmental bottlenecks and irreversible damage

The high sensitivity observed during the pupal phase underscores its role as a critical developmental bottleneck. Reductions in emergence success by 15% and 30% under cyclic thermal stress of 37°C and 38°C respectively (Fig. 1*a*) align with patterns observed under constant thermal stress [30], though to a lesser extent. Noteworthy, average emergence rates of about 80% at 34°C were slightly higher than in our previous *in vitro* study under constant thermal stress [30] and for *B. impatiens* colonies exposed to 25°C, 30°C, and 35°C [23], likely reflecting temporal, colony, and species variation. Interestingly, despite high emergence rates reported for those *B. impatiens* colonies, there was also a drastic reduction in the number of offspring for colonies housed at 35°C, indicating colony failure which could be linked to the increase in adult workers abandoning their nest [23]. This observation may indicate insufficient nursing (feeding) of larvae at 35°C. Indeed, our observation that heavier pupae (strongly dependent on L4 larvae weight, Fig. S3*a*) were more likely to emerge as adults (Fig. 1*b*) supports the idea that successful pupa-to-adult transition might be resource-dependent [30]. This suggests that resource limitations within a colony could trigger negative synergistic interactions, amplifying the detrimental effects of thermal stress.

We found no significant effect on the duration of metamorphosis (pupation)

between the cyclic thermal stress treatments and incubating pupae at a constant 34°C

(Fig. 1*c*). Regardless, pupation duration differed between workers (8-9 days) and males

(9-10 days), aligning with data reported for pupae reared in *B. impatiens* colonies [47] and

B. terrestris reared *in vitro* [30, 37]. In contrast to how constant thermal stress of 38°C

may slightly prolong pupal development, exposure to 36°C appears to shorten

developmental time compared to 34°C [30].

Most notably, 38°C cyclic thermal stress led to a higher incidence of deformed

wings (Fig. 3*a*), but this effect was much less pronounced compared to pupae exposed to

a constant temperature of 38°C for five days during pupation [30]. Interestingly, cold

stress during pupal development, mimicking a spring cold snap, has been shown to cause

wing deformities and reduced flight ability in the alfalfa leafcutting bee *Megachile*

rotundata (Hymenoptera: Apiformes: Megachilidae) [33]. While wing morphology is a

known indicator of irreparable developmental damage [30, 48, 49], our data provide clear

empirical evidence that deformations (antennae and wings) can serve as a powerful

hazard indicator (HR = 2.38) that predicts adult longevity regardless of treatment (Fig. 2*e*).

Although survival was reduced in adults that were exposed to the cyclic 38°C treatment

(Fig. 2*a,b,e*), there was a steep decline in adult survival during the first two weeks. We

speculate that this may stem from a lack of social interactions [50]. Supporting this and

validating our *in vitro* rearing approach, no significant difference was detected in the

mortality risk during adulthood between individuals developed naturally in their natal

colonies until emergence and those reared *in vitro* at 34°C as L4 larvae and pupae (HR =

0.95; Fig. 2*c,d*).

Beyond individual mortality, workers and males with severely deformed wings are

unable to fly, which consequently reduces colony fitness. In fact, it has already been

shown that thermal stress during development can reduce adult fitness across diverse

insect taxa [6, 7, 9]. Adding to the consequences of deformed wings, deformed antennae

likely impair the ability to detect scents of nestmates, mating partners (important for

males), and floral resources (important for workers). This idea is supported by a study

demonstrating heat stress-induced olfactory impairment in adult bumblebee workers [51].

Decoupling fitness costs from adult morphometrics and fat body

Despite the difference in functional longevity, there was no evidence that deferred

mortality was associated with easily measurable morphometric traits (i.e. pupal weight

loss, emergence weight, head width, and ITD) between *in vitro* treatments (Fig. 3*b-d* and

S3*b-d*). While this result suggests that selection pressure from short-term thermal stress

for body size might be more pronounced at earlier larval stages [30, 31], this is in contrast

with some previous colony-level findings. For example, a smaller male wing size was

reported for *B. terrestris* micro-colonies housed at 33°C compared to 26°C [26].

Furthermore, worker offspring were shown to have a smaller ITD in colonies housed at

33°C compared to 25°C, whereas males appeared to be unaffected [25, 27]. It is worth

noting that our colony-reared individuals, housed at 25°C, also emerged slightly heavier

than *in vitro* reared bees (Fig. 3*b*). Although this difference could indicate a temperature-

related ceiling effect, with pupae reared at 34°C *in vitro* being already heat-stressed, our

chosen baseline aligns with brood temperatures measured in *B. terrestris* nests [19, 28,

29].

An alternative explanation for our observed weight difference would be the

possibility of initial food consumption by newly emerged bees inside the nest, unlike bees

emerging from the *in vitro* rearing. However, as the median proportional pupal weight loss

was almost identical between treatments (Fig. 1*d* and S3*b*) with a strong positive

relationship between initial pupal weight and emergence weight (Fig. S3*c*) and the scaling

between emergence weight and head width was also very similar across all treatment

groups (Fig. 3*c*), we think that our observed pattern reflects the limitations of *in vitro*

rearing in general compared to naturally reared larvae inside the colony. It is very likely

that colony-reared L4 larvae received more food than our *in vitro* reared L4 larvae, as

workers can sense hungry larvae and provide food whenever needed [52].

We would even speculate that the body size effects observed in thermally stressed

colonies of the aforementioned studies [25-27] play out at an earlier larval stage [30, 31]

and are an indirect result of altered worker nursing and foraging behaviour [23, 24] rather

than a direct physiological consequence of the heat itself. It could well be that either

resources are much more limited or larvae are simply insufficiently fed by thermally

stressed adult workers. This highlights the importance of using an *in vitro* approach to

isolate direct physiological effects from comprehensive colony dynamics [30, 31].

While increased lipid content is advantageous for constantly heat-stressed pupae

to successfully emerge [30], we found no significant difference in relative lipid content at

the end of their adult life (Fig. 3d), aligning with previous findings of energy homeostasis in

ageing bumblebee workers [53]. This indicates the capacity of surviving adults to maintain

homeostasis of their fat body, which is the primary site for heat shock protein (HSP)

synthesis and storage of energy reserves [48, 54, 55], by resource allocation or

compensatory foraging. Hence, any sublethal damage carrying over to adulthood might be

genetically, structurally, and/or physiologically mediated rather than simply due to

depleted energy reserves. This aligns with findings where developmental thermal stress

indirectly affected insect adult physiology, such as flight performance and behaviour [56-

58].

Conclusion

These findings highlight that sublethal thermal stress during insect development may lead

to profound fitness costs. Our study provides compelling empirical evidence that cyclic

thermal stress during pupal development significantly reduces adult longevity in *B.*

terrestris. Furthermore, our findings confirm that 38°C is a critical threshold for

developmental disruption in *B. terrestris* [30, 31]. Although our study oversimplifies

realistic thermal stress scenarios, such as those experienced during heatwaves, and *B.*

terrestris only serves as a model system, we are convinced that these findings provide

valuable insights into the consequences of thermal stress during pupal development in

bumblebees. Species that are less heat-tolerant than *B. terrestris*, such as *B. lapidarius*

and *B. alpinus* [13], or typically nest above ground, such as *B. hypnorum* and *B.*

pascuorum [19] or are used in greenhouses for pollination services, would be expected to

be especially at risk [30]. As sublethal carry-over effects are rarely considered, current

prediction models may underestimate the long-term impact of environmental stressors on

crucial pollinator populations. To generate a more holistic picture of environmental stress

impact, gaining a nuanced mechanistic understanding of carry-over effects across diverse

insect species and life stages, incorporating both individual-level, and for social insects,

also colony-level physiological and life-history data, remains crucial.

Data accessibility statement:

483 Dataset and R script used for statistical analysis and producing all Fig. are available from

484 the Zenodo Repository: https://doi.org/10.5281/zenodo.18298098. Electronic supplementary

485 material is available online.

Statement of authorship:

488 M.M.M.: methodology, investigation, data curation. C.K.: conceptualization, methodology,

489 investigation, data curation, resources, validation, formal analysis, writing – original draft

490 preparation, writing – review and editing, visualization, supervision, project administration.

Acknowledgments

493 We thank Sandra Laußer, Laura Wögler, Ramona Schöpf, and Michael Ott for their

494 assistance in the laboratory. We are grateful to Prof. Dr. Tomer J. Czaczkes for

495 sponsoring us bumblebee colonies. We have used AI for proofreading.

Ethics statement

498 This study was conducted in accordance with the ethical regulations of the German

499 Animal Welfare Act (TierSchG) for conducting experiments with insects.

Funding

This study was carried out without any support of third-party funding.

Competing interests

The authors have no competing interests.

5. Reference

1.   Garcia RA, Cabeza M, Rahbek C, Araújo MB. Multiple dimensions of climate
change and their implications for biodiversity. Science. 2014;344(6183):1247579.
2.   Meehl GA, Tebaldi C. More intense, more frequent, and longer lasting heat waves
in the 21st century. Science. 2004;305(5686):994-7.
3.   Ma C-S, Ma G, Pincebourde S. Survive a warming climate: insect responses to
extreme high temperatures. Annual Review of Entomology. 2021;66:163-84.
4.   González-Tokman D, Córdoba-Aguilar A, Dáttilo W, Lira-Noriega A,
Sánchez-Guillén RA, Villalobos F. Insect responses to heat: physiological mechanisms,
evolution and ecological implications in a warming world. Biological Reviews.
2020;95(3):802-21.
5.   Jørgensen LB, Ørsted M, Malte H, Wang T, Overgaard J. Extreme escalation of
heat failure rates in ectotherms with global warming. Nature. 2022;611(7934):93-8.
PubMed PMID: Jørgensen2022.
6.   Zhang W, Rudolf VHW, Ma C-S. Stage-specific heat effects: timing and duration of
heat waves alter demographic rates of a global insect pest. Oecologia. 2015;179(4):947-
57. PubMed PMID: Zhang2015.
7.   Padde JR, Zhou Y, Chen Y, Zhu Y, Yang Y, Hou M, et al. Adaptation and carry
over effects of extreme sporadic heat stress in *Culex* mosquitoes. 2024;260:107417.
8.   Klockmann M, Kleinschmidt F, Fischer K. Carried over: Heat stress in the egg
stage reduces subsequent performance in a butterfly. PLoS one. 2017;12(7):e0180968.
9.   Sales K, Vasudeva R, Gage MJG. Fertility and mortality impacts of thermal stress
from experimental heatwaves on different life stages and their recovery in a model insect.
Royal Society Open Science. 2021;8(3):201717. doi: doi:10.1098/rsos.201717.
10.   White SA, Dillon ME. Climate warming and bumble bee declines: the need to
consider sub-lethal heat, carry-over effects, and colony compensation. Frontiers in
Physiology. 2023;14. doi: 10.3389/fphys.2023.1251235.
11.   Ghisbain G, Thiery W, Massonnet F, Erazo D, Rasmont P, Michez D, et al.
Projected decline in European bumblebee populations in the twenty-first century. Nature.
2024;628(8007):337-41.

12. Soroye P, Newbold T, Kerr J. Climate change contributes to widespread declines
among bumble bees across continents. Science. 2020;367(6478):685-8.

13. Martinet B, Dellicour S, Ghisbain G, Przybyla K, Zambra E, Lecocq T, et al. Global
effects of extreme temperatures on wild bumblebees. Conservation Biology.
2021;35(5):1507-18.

14. Cameron SA, Lozier JD, Strange JP, Koch JB, Cordes N, Solter LF, et al. Patterns
of widespread decline in North American bumble bees. Proceedings of the National
Academy of Sciences. 2011;108(2):662-7.

15. Kerr JT, Pindar A, Galpern P, Packer L, Potts SG, Roberts SM, et al. Climate
change impacts on bumblebees converge across continents. Science.
2015;349(6244):177-80.

16. White SA, Dillon ME. Climate warming and bumble bee declines: the need to
consider sub-lethal heat, carry-over effects, and colony compensation. Frontiers in
Physiology. 2023;Volume 14 - 2023. doi: 10.3389/fphys.2023.1251235.

17. Klein A-M, Vaissiere BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen
C, et al. Importance of pollinators in changing landscapes for world crops. P Roy Soc B.
2007;274:303-13. doi: 10.1098/rspb.2006.3721. PubMed PMID: WOS:000243354100001.

18. Goulson D, Lye GC, Darvill B. Decline and conservation of bumble bees. Annu
Rev Entomol. 2008;53(1):191-208.

19. Kevan PG, Rasmont P, Martinet B. Thermodynamics, thermal performance and
climate change: temperature regimes for bumblebee (Bombus spp.) colonies as examples
of superorganisms. Frontiers in Bee Science. 2024;Volume 2 - 2024. doi:
10.3389/frbee.2024.1351616.

20. Potts SG, Imperatriz-Fonseca V, Ngo HT, Aizen MA, Biesmeijer JC, Breeze TD, et
al. Safeguarding pollinators and their values to human well-being. Nature. 2016;540:220-
9.

21. Theodorou P, Kühn O, Baltz LM, Wild C, Rasti SL, Bucksch CR, et al. Bumble bee
colony health and performance vary widely across the urban ecosystem. Journal of
Animal Ecology. 2022;91(10):2135-48. doi: https://doi.org/10.1111/1365-2656.13797.

22. Bretzlaff T, Kerr JT, Darveau C-A. High temperature sensitivity of bumblebee
castes and the colony-level costs of thermoregulation in Bombus impatiens. Journal of
Thermal Biology. 2023;117:103710.

23. Bretzlaff T, Kerr JT, Darveau C-A. Handling heatwaves: balancing
thermoregulation, foraging and bumblebee colony success. Conservation Physiology.
2024;12(1). doi: 10.1093/conphys/coae006.

24. Sepúlveda Y, Nicholls E, Schuett W, Goulson D. Heatwave-like events affect
drone production and brood-care behaviour in bumblebees. PeerJ. 2024;12:e17135.

25. Guiraud M, Cariou B, Henrion M, Baird E, Gérard M. Higher developmental
temperature increases queen production and decreases worker body size in the
bumblebee *Bombus terrestris*. Journal of Hymenoptera Research. 2021;88:39-49.

26. Gérard M, Michez D, Debat V, Fullgrabe L, Meeus I, Piot N, et al. Stressful
conditions reveal decrease in size, modification of shape but relatively stable asymmetry
in bumblebee wings. Sci Rep. 2018;8(1):15169. PubMed PMID: Gerard2018.

27. Gérard M, Guiraud M, Cariou B, Henrion M, Baird E. Elevated developmental
temperatures impact the size and allometry of morphological traits of the bumblebee
Bombus terrestris. Journal of Experimental Biology. 2023;226(8):jeb245728.

28. Weidenmüller A, Kleineidam C, Tautz J. Collective control of nest climate
parameters in bumblebee colonies. Animal Behaviour. 2002;63(6):1065-71.

29. Wynants E, Lenaerts N, Wäckers F, Van Oystaeyen A. Thermoregulation
dynamics in commercially reared colonies of the bumblebee Bombus terrestris.
Physiological Entomology. 2021;46(1):110-8. doi: https://doi.org/10.1111/phen.12350.

30. Laußer S, Kurze C. Impact of thermal stress during pupal development in a key
pollinator. Proceedings B. 2025;292(2061):20252029.

31. Wögler L, Kurze C. Experimental short-term heatwaves negatively impact body
weight gain and survival during larval development in *Bombus terrestris* L. (Hymenoptera:
Apidae). Biology Open. 2025;14(4). doi: 10.1242/bio.061781.

32. Radmacher S, Strohm E. Effects of constant and fluctuating temperatures on the
development of the solitary bee *Osmia bicornis* (Hymenoptera: Megachilidae). Apidologie.
2011;42:711-20.

33. Earls KN, Porter MS, Rinehart JP, Greenlee KJ. Thermal history of alfalfa
leafcutting bees affects nesting and diapause incidence. Journal of Experimental Biology.
2021;224(22). doi: 10.1242/jeb.243242.

34. Rolff J, Johnston PR, Reynolds S. Complete metamorphosis of insects.
Philosophical Transactions of the Royal Society B: Biological Sciences.
2019;374(1783):20190063. doi: doi:10.1098/rstb.2019.0063.

35. Lausser S, Kurze C. Heatwave length and severity drive development disruption in
a key pollinator. bioRxiv. 2025:2025.03.26.645459. doi: 10.1101/2025.03.26.645459.

36. Gilgenreiner M, Kurze C. Age dominates flight distance and duration, while body
size shapes flight speed in *Bombus terrestris* L.(Hymenoptera: Apidae). Proceedings B.
2024;291(2027):20241001. doi: https://doi.org/10.1098/rspb.2024.1001.

37. Kato Y, Kikuta S, Barribeau SM, Inoue MN. In vitro larval rearing method of
eusocial bumblebee *Bombus terrestris* for toxicity test. Scientific Reports.
2022;12(1):15783. PubMed PMID: Kato2022.

38.   Brown MJF, Loosli R, Schmid-Hempel P. Condition-dependent expression of
virulence in a trypanosome infecting bumblebees. Oikos. 2000;91(3):421-7. doi:
https://doi.org/10.1034/j.1600-0706.2000.910302.x.

39.   R_Core_Team. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/;
2024.

40.   Brooks ME, Kristensen K, Van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et
al. glmmTMB balances speed and flexibility among packages for zero-inflated generalized
linear mixed modeling. The R journal. 2017;9(2):378-400.

41.   Christensen RHB. ordinal—regression models for ordinal data. R package version.
2019;10(2019):54.

42.   Hartig F. DHARMa: residual diagnostics for hierarchical (multi-level/mixed)
regression models. R package version 03. 2020;3(5).

43.   Lenth R, Lenth MR. Package 'lsmeans'. The American Statistician.
2018;34(4):216-21.

44.   Therneau TM, Grambsch PM. Modeling survival data : extending the Cox model.
New York: Springer; 2000.

45.   Kurze C, Jenkins NE, Hughes DP. Evaluation of direct and indirect transmission of
fungal spores in ants. Journal of Invertebrate Pathology. 2020;172. doi:
10.1016/j.jip.2020.107351. PubMed PMID: WOS:000525892100006.

46.   Kassambara A, Kosinski M, Biecek P. survminer: Drawing Survival Curves
using'ggplot2'. CRAN: Contributed Packages. 2016.

47.   Cnaani J, Schmid-Hempel R, Schmidt J. Colony development, larval development
and worker reproduction in Bombus impatiens Cresson. Insectes Sociaux.
2002;49(2):164-70.

48.   Vanderplanck M, Breeze T, Cini E, Potts SG, Senapathi D. Monitoring bee health
in European agroecosystems using wing morphology and fat bodies. One Ecosystem.
2021;6.

49.   Gérard M, Michez D, Debat V, Fullgrabe L, Meeus I, Piot N, et al. Stressful
conditions reveal decrease in size, modification of shape but relatively stable asymmetry
in bumblebee wings. Scientific Reports. 2018;8(1):15169. PubMed PMID: Gerard2018.

50.   Wang ZY, McKenzie-Smith GC, Liu W, Cho HJ, Pereira T, Dhanerawala Z, et al.
Isolation disrupts social interactions and destabilizes brain development in bumblebees.
Current Biology. 2022;32(12):2754-64.e5.

51.   Nooten SS, Korten H, Schmitt T, Kárpáti Z. The heat is on: reduced detection of
floral scents after heatwaves in bumblebees. Proceedings of the Royal Society B.
2024;291(2029):20240352.

52. Boer SPAd, Duchateau MJHM. A larval hunger signal in the bumblebee Bombus
terrestris. 2006;53(3):369-73. PubMed PMID: Boer2006.

53. Schulte H, Bäuml C, Strohm E, Kurze C. Age-dependent shift from high lipid
reserves to high haemolymphs sugars supply underlies worker maturation in *Bombus*
terrestris (Hymenoptera: Apidae). bioRxiv. 2026:10.64898/2026.01.17.700087. doi:
https://www.biorxiv.org/content/10.64898/2026.01.17.700087v1.

54. Arrese EL, Soulages JL. Insect fat body: energy, metabolism, and regulation.
Annual review of entomology. 2010;55:207-25.

55. King AM, MacRae TH. Insect heat shock proteins during stress and diapause.
Annual review of entomology. 2015;60(1):59-75.

56. Gérard M, Cariou B, Henrion M, Descamps C, Baird E. Exposure to elevated
temperature during development affects bumblebee foraging behavior. Behavioral
Ecology. 2022;33(4):816-24.

57. Galarza JA, Dhaygude K, Ghaedi B, Suisto K, Valkonen J, Mappes J. Evaluating
responses to temperature during pre-metamorphosis and carry-over effects at post-
metamorphosis in the wood tiger moth (*Arctia plantaginis*). Philosophical Transactions of
the Royal Society B: Biological Sciences. 2019;374(1783):20190295. doi:
doi:10.1098/rstb.2019.0295.

58. Arambourou H, Sanmartín-Villar I, Stoks R. Wing shape-mediated carry-over
effects of a heat wave during the larval stage on post-metamorphic locomotor ability.
2017;184(1):279-91. PubMed PMID: Arambourou2017.

667