Deferred mortality: cyclic thermal stress during pupation triggers irreversible carry-

over costs in a key pollinator

Maximilian M. Mandlinger[1] and Christoph Kurze[1]*

[1] Evolutionary Ecology, University of Regensburg, Germany

* Corresponding author, email: Christoph.Kurze@ur.de

Abstract

Extreme weather events, such as heatwaves, pose a concerning threat to global

biodiversity, particularly to terrestrial ectotherms like insects. While lethal effects of

thermal stress are widely studied, very little is known about sublethal carry-over effects of

heatwaves experienced during development on adult fitness. Addressing this fundamental

knowledge gap is crucial for key pollinators such as bumblebees, which face alarming

population declines. Therefore, we used a highly controlled *in vitro* approach to expose

Bombus terrestris pupae to ecologically relevant cyclic thermal stress, and studied their

emergence success, subsequent adult longevity, and morphological traits. We found that

cyclic thermal stress caused significant acute pupal mortality (up to 30% reduction in

emergence). Most notably, this developmental stress resulted in deferred mortality,

significantly reducing adult longevity (HR = 1.81) in both workers and males. Furthermore,

we identified antennae and wing deformations as a powerful hazard indicator (HR = 2.50)

that strongly predicts premature adult mortality. Our findings reveal that cyclic thermal

stress during pupation imposes irreversible developmental damage that undermines adult

physiological resilience. We argue that life-stage specific carry-over effects should be

considered more in the future to avoid underestimating the long-term impact of climate

change on crucial insect pollinator populations.

Keywords: climate change, heatwave, metamorphosis, mortality, survival, lifespan,

deformations, malformations, social insects, bees

1. Introduction

Extreme weather events, such as heatwaves, are increasing in frequency and intensity due to climate change, leading to distinct and widespread implications for biodiversity, including insects (Meehl & Tebaldi 2004; Garcia *et al.* 2014). While thermal stress is known to have lethal and sublethal effects across a wide range of insects (González-Tokman *et al.* 2020; Ma, Ma & Pincebourde 2021; Jørgensen *et al.* 2022), most studies focus on the acute effects in adults. Little is known about how thermal stress during an earlier life-stage impose sublethal, yet long-lasting, carry-over effects on adult fitness (Zhang, Rudolf & Ma 2015; Klockmann, Kleinschmidt & Fischer 2017; Sales, Vasudeva & Gage 2021; Padde *et al.* 2024). Gaining a better understanding of such sublethal carry-over effects is essential for accurately forecasting population declines in keystone insect taxa such as bumblebees (*Bombus* sp., Hymenoptera: Apidae) (White & Dillon 2023b).

Bumblebees, as a cold-adapted taxon, have experienced populations declines, which are partially attributed to climate change (Cameron *et al.* 2011; Kerr *et al.* 2015; Soroye, Newbold & Kerr 2020; Martinet *et al.* 2021; White & Dillon 2023a; Ghisbain *et al.* 2024). This is a concerning trend, given that they provide crucial pollination services in ecosystems and agriculture (Klein *et al.* 2007; Goulson, Lye & Darvill 2008). While ground-nesting species can mitigate extreme ambient temperatures to some extent, other species, such as *B. hypnorum* and *B. pascuorum* that nest typically above the ground (Kevan, Rasmont & Martinet 2024) or commercially used species in greenhouses such as *B. terrestris* (Potts *et al.* 2016), face a higher risk of thermal stress. Colony-level studies have demonstrated that constant prolonged elevated temperatures negatively impact colony energetics, growth and reproductive fitness (Theodorou *et al.* 2022; Bretzlaff, Kerr & Darveau 2023; Bretzlaff, Kerr & Darveau 2024; Sepúlveda *et al.* 2024), and such thermal stress have been linked to altered offspring morphology (Gérard *et al.* 2018; Guiraud *et al.* 2021; Perl *et al.* 2022; Gérard *et al.* 2023). However, behavioural responses vary vastly between colonies and make it difficult to disentangle indirect effects (e.g.

fanning workers spend less time for nursing or foraging) from the direct physiological

effects. To gain a precise mechanistic understanding of how thermal stress affects

specific larval stages, *in vitro* rearing offers the required control at the cost of losing

natural realism (Laußer & Kurze 2025; Wögler & Kurze 2025).

Our previous *in vitro* studies revealed stage-specific responses to thermal stress in

B. terrestris (Laußer & Kurze 2025; Wögler & Kurze 2025). While smaller fourth-instar

larvae may have a selective advantage under thermal stress due to their ability of more

easily reaching critical body mass for pupation (Wögler & Kurze 2025), thermal stress at

the pupal stage appears to select for individuals with increased relative lipid reserves

(Laußer & Kurze 2025). Importantly, we found strong evidence for a reduction in adult

emergence and ab increase of wing deformations in thermally stressed larvae and pupae

(Laußer & Kurze 2025; Wögler & Kurze 2025). While these studies illustrate impaired

larval and pupal development, the magnitude of the downstream impact on adult survival

remains unknown.

To address this fundamental knowledge gap, we used highly controlled *in vitro*

rearing of *B. terrestris* pupae to isolate the effect of cyclic thermal stress scenarios

(simulating heatwaves) on individual life history. Our main aim was to assess the carry-

over risk under fluctuating thermal extremes, an approach more ecologically relevant than

constant temperature regimes (Radmacher & Strohm 2011). We focused on the pupal

stage as metamorphosis in holometabolous insects, such as bees, represents the most

extreme transition between phenotypes (Rolff, Johnston & Reynolds 2019) and is likely

the most vulnerable to environmental stress (Laußer & Kurze 2025). We quantified both

the direct effect on emergence success and the subsequent long-term adult survival,

morphology, and energy reserves. We hypothesised that increasing cyclic thermal stress

during pupation would affect adult morphology (specifically wing morphology (Laußer &

Kurze 2025)) and trigger deferred mortality, significantly reducing adult longevity.

Furthermore, we expected that the impact of thermal stress would differ between workers

and males.

2. Materials and methods

2.1 Colony husbandry

We maintained ten queen-right source colonies of *B. terrestris* (Natupol Research Hives, Koppert B.V., Netherlands) under standardized laboratory conditions (Gilgenreiner & Kurze 2024; Laußer & Kurze 2025; Wögler & Kurze 2025). Colonies were housed in standard nest boxes (23 x 25 x 15 cm), which connected to foraging arenas (59 x 39 x 26 cm). We maintained all colonies at 25 ± 1°C and 30-50% RH, under a 14:10 h light:dark regime. Bumblebees had *ad libitum* access to a 66% w/w sucrose solution (Natupol Bee Happy, Koppert B.V., Netherlands). We supplied each colony with 6-11 g of pollen candy (67% w/w organic pollen, 25% w/w sucrose, 8% w/w water) daily.

2.2 Larvae collection and *in vitro* rearing

We collected 691 fourth-instar larvae (L4) from fully developed source colonies (figure S1*a*). Collecting L4 was crucial to ensure that we could start all thermal regimes consistently at the beginning of the pupal phase. Larvae were reared *in vitro* following our established protocol (for details see Laußer & Kurze 2025; Wögler & Kurze 2025).

Briefly, L4 were carefully transferred into 3D-printed polylactide (PLA) artificial brood cells (figure S1*b*), placed inside 24- or 48-well clear flat bottom plates (figure S1*c*; Falcon/Corning, USA), and put within a humidity-controlled plastic container (18.5 x 18.5 x 11.5 cm), kept at a constant temperature of 34°C (KB115, BINDER GmbH, Germany). To maintain a relative humidity (RH) of 65 ± 10%, a 120 mL cup of saturated sodium chloride solution was also placed within the container.

Larvae were fed a pollen medium (figure S1*b*), consisting of 50% w/v sucrose solution, 40% organic pollen, 10% yeast extract (Bacto™, BD, USA), and 1% casein sodium salt from bovine milk (Sigma-Aldrich, Germany) (Kato *et al.* 2022), twice daily to satiation (Wögler & Kurze 2025). Our setup allowed us to measure larval and pupal weight non-invasively and facilitated the easy removal of any dead individuals. Feeding was stopped with the beginning of the pupal phase.

2.3 Experimental procedure

At the beginning of pupation, each individual was weighed (d = 0.1 mg, Sartorius M-Pact

AX224, Sartorius GmbH, Germany) and randomly assigned to one of three experimental

thermal treatment groups (Table 1). Pupae were transferred to new, humidity-controlled

containers within one of three programmed incubators. The control group was maintained

at a constant 34°C (34°C ctrl). The two cyclic treatment groups experienced a 12:12 h

temperature cycle between the base of 34°C (night phase) and a thermal maximum (day

phase) of either 37°C (37°C cyclic) or 38°C (38°C cyclic). We monitored pupal survival

and emergence success twice daily. In addition to the *in vitro* treatments, newly emerged

adults (< 1 day old) were collected directly from their natal colonies to serve as an external

colony control group (colony ctrl) for the subsequent longevity assay.

2.4 Adult survival assay

Newly emerged bees from the *in vitro* treatments and the additional colony control group

(Table 1) were weighed and then individually housed in modified 50 mL tubes (figure S1*d*;

Greiner Bio-one GmbH, Germany). These were stored in a wooden rack within the

insectary room of their natal colonies at 25 ± 1°C and 30-50% RH (figure S1*e*). Each tube

contained a 3D-printed PLA mesh insert for sanitary reasons. The lids were modified to

include three 3 mm ventilation holes and one 1 cm feeding port, through which a 1.5 mL

perforated tube provided *ad libitum* access to a 60% w/v sucrose solution. Pollen candy

was offered separately in the caps of 2 mL reaction tubes and was replaced weekly or as

necessary. Housing tubes and meshes were replaced as needed to ensure sanitary

conditions. Adult survival was monitored around noon daily. Dead individuals were

recorded and immediately frozen at -21°C until further analysis.

140 **Table 1. Overview of the experimental design.** This table details the rearing conditions (regime and

141 temperature) of pupae for all experimental groups and an additional colony control. Sample sizes (n) reflect

142 the total number of pupae per treatment at start of *in vitro* rearing experiment (indicated by [§]) and emerged

143 adults at the start of the survival assay (indicated by [#]).

Treatment	Regime[§]	Temperature[§]	Sample size (n)
colony ctrl	colony reared	25 ± 1°C*	[#]115
34°C ctrl	constant	34°C	[§]132/ [#]102
37°C cyclic	12:12 h cycle	37°C:34°C	[§]125/ [#]82
38°C cyclic	12:12 h cycle	38°C:34°C	[§]146/ [#]78

* colonies kept in the insectary
[§] pupae / [#] adults

2.5 Morphological and physiological analysis

We contacted the morphological and physiological analysis following established

protocols (Gilgenreiner & Kurze 2024; Laußer & Kurze 2025; Wögler & Kurze 2025). The

intertegular distance (ITD), used as proxy for body size, was measured using a digital

microscope (VHX-500F, Keyence GmbH, Germany). The sex of each bee was

determined by counting flagellomeres (workers = 10; males = 11). Furthermore, any

deformations of their antennae were recorded (figure S2*a*).

To assess wing integrity, forewings were carefully detached, embedded side-by-

side on a microscope slide (Roti®Mount, Carl Roth GmbH + Co. KG, Germany) and

photographed at 20x magnification. Wings were subsequently classified as deformed

(figure S2*b*) when visually apparent malformations were present.

The remaining bee bodies were processed to determine dry mass and relative lipid

content. Specimens were first drying at 60°C for three days (U40, Memmert GmbH & Co.

KG, Germany) and immediately weighed to obtain their dry mass.

The absolute body lipid content was determined by using a petroleum ether

extraction method. The specimens were re-drying and weighted again post-extraction to

allowing us to calculate the difference between the initial and post-extraction dry masses.

The relative lipid content, which served as a proxy for the fat body size and energy

reserves, was calculated as the absolute lipid content divided by the initial dry mass.

2.6 Statistical analyses

All statistical analyses and data visualizations were performed in R version 4.5.1 (R Core Team 2024), similar to our previous work (Kurze, Jenkins & Hughes 2020; Laußer & Kurze 2025). The complete code, including model selection procedures (R Markdown output), and datasets are provided in the Dryad Digital Repository (XXX).

To analyse the effect of cyclic thermal stress and additional explanatory variables (e.g. sex, pupal weight) and their interactions on emergence rates, ITD, dry mass, relative lipid content, as well as antennae and wing deformation, we fitted generalized linear mixed effect models (GLMMs) using the *glmmTMB* package (Brooks *et al.* 2017). Emergence rates and deformation status were fitted with binomial distribution (logit link), while the continuous variables ITD, dry mass, and relative lipid content were fitted with a gamma distribution (log link). For analysing the effects on pupation duration (days to emergence), we applied an ordinal cumulative link mixed model (CLMM) with the Laplace approximation using the *ordinal* package (Christensen 2019), as assumption for fitting a GLMM were not met (described below). While the CLMM included only colony ID as a random factor, all GLMMs also included collection date as random factor to account for colony and temporal-specific variability.

For each response variable, a set of candidate models was compared. The most parsimonious and best-fitting model based on the Akaike information criterion (AIC), with lower AIC values indicating a better fit. The best-fitting models were compared to a simpler nested model using likelihood ratio tests (LRTs). When added terms did not improve model fit significantly, the simpler model was chosen and compared against its respective null-model using LRT. Model assumptions and dispersion of the data were checked using the *DHARMa* package (Hartig 2020). When model assumptions were met, statistical significance ($p < 0.05$) was determined using the *Anova* function. Pairwise comparisons between treatment groups were conducted using the *emmeans* package (Lenth & Lenth 2018) with Holm-Bonferroni corrections. Smooth predictions of emergence

rate across the observed range of larval weights were generated using a prediction

dataset with 100 evenly spaced values.

Differences in survival probability between treatment groups were initially

determined for workers and males separately based on Kaplan-Meier analyses using the

survival and *ggsurvfit* packages (Therneau & Grambsch 2000). To further analyse the

main factors affecting adult survival, we fitted a set of candidate Cox Proportional Hazards

(CoxPH) models the using *coxme* packges (Therneau & Grambsch 2000). The full model

contained treatment and sex (and their interactions), adult dry mass, relative lipid content,

and deformations (antennae and wing deformation combined). The colony ID was

included as a random factor. Model selection followed the AIC and LRT procedure as

described above. The final best-fitting model to predict mortality included the explanatory

variables treatment, sex, and their interactions, alongside deformations status. As

proportional hazards (PH) assumptions were not entirely met (as diagnosed model

diagnostics using *ggcoxdiagnostics* function of the *surmiver* package (Kassambara,

Kosinski & Biecek 2016)), the final model was stratified by sex. Pairwise comparisons

between treatment groups were performed using the *glht* function with Holm-Bonferroni

correction.

3. Results

3.1 Emergence success and developmental duration

Cyclic thermal stress and pupal weight significantly affected the proportion of pupae

reaching adulthood (GLMM: χ^2 = 35.187, df = 3, p < 0.0001; figures 1*a,b*). This effect was

primarily driven by the increase in thermal stress (χ^2 = 22.276, df = 2, p < 0.0001; figure

1*a*). Pairwise comparisons revealed that pupae exposed to the 38°C cyclic treatment

resulted in a 30% lower likelihood of emergence compared to the controls (p < 0.0001),

while the 37°C cyclic treatment had a 15% reduction in emerged adults (p < 0.05). The

difference between the 38°C and 37°C cyclic treatments was also significant (p < 0.05).

Furthermore, emergence probability was significantly higher with an increase in initial

pupal weight (χ^2 = 11.065, df = 1, p < 0.001; figure 1*b*).

While we observed a significant overall effect on pupation duration (CLMM: χ^2 =

43.958, df = 3, p < 0.0001; figure 1*c*), this effect was not related to the thermal stress

treatment (χ^2 = 1.6121, df = 2, p > 0.05), but was sex-dependent (χ^2 = 34.311, df = 1, p <

0.0001). Pairwise comparisons revealed that workers of all treatment groups were

significantly different from males of all treatment groups (p < 0.001).

3.2 Carry-over effects on adult longevity

Adult survival was significantly reduced with increasing cyclic thermal stress exposure

during pupation for both workers and males (Log-rank test for workers: χ^2 = 8.9, df =

2, p = 0.01; for males: χ^2 = 11.1, df = 2, p < 0.01; Kaplan-Meier survival curves in figure

2*a,b*). The presence of carry-over effects on longevity was strongly supported by the

CoxPH model stratified by sex (χ^2 = 349.78, df = 3, p < 0.0001, model concordance =

0.64; figure 2*e*). Adults of 38°C cyclic treatment had a significantly higher mortality risk

than the controls (HR = 1.81, 95% CI = 1.56-2.13, p < 0.0001). The 37°C cyclic treatment

posed a marginally significant increase in mortality risk (HR = 1.30, 95% CI = 0.98-1.73, p

= 0.07). We also found that individuals with morphological deformations (combining

deformed antennae and wings) had a significantly higher mortality risk (HR = 2.50, 95% CI

= 2.02-3.10, p < 0.0001).

Importantly, we found no significant difference in adult survival between *in vitro*

reared controls and naturally colony reared controls (workers: $\chi^2 = 0$, df = 1, p > 0.05,

figure 2*c*; males: $\chi^2 = 0.8$, df = 1, p > 0.05, figure 2*d*).

Figure 1. Direct effects of cyclic thermal stress during pupation on emergence success and

development duration in *B. terrestris*. Panel *(a)* illustrates the proportion of pupae emerging as adult bees,

and *(b)* shows the predicted probability of emergence in relation to pupal weight and treatment. Panel *(c)*

displays the pupation duration until emergence for male (open violins) and workers (filled violins). Controls

(34°C ctrl, reared at constant 34°C, blue) were compared to individuals exposed to cyclic thermal stress

regimes of 37°C (37°C cyclic, orange) and 38°C (38°C cyclic, reddish-purple). Individual data points in (*b*) and

(*c*) are shown as circles. In panels (*a*) and (*b*), sample sizes (n) are provided in plots. Significant differences (p

< 0.05), based on post-hoc comparisons from GLMMs, are indicated by different letters above the bars and

next to the violin plots (groups sharing a letter are not significantly different).

Figure 2. Carry-over effects of cyclic thermal stress during pupation on adult longevity in *B. terrestris*.

Kaplan-Meier survival curves (± 95% confidence intervals, CI, shaded areas) showing the effect of in vitro

thermal treatments on worker (*a*) and male (*b*) adult longevity. The constant 34°C control group (34°C ctrl,

blue) is compared to the cyclic thermal stress treatments (37°C cyclic, orange; 38°C cyclic, reddish-purple).

Panel (*c*) and (*d*) compare the survival of *in vitro*-reared controls (34°C ctrl) to naturally reared colony controls

(colony ctrl, bluish green) for worker and male, respectively. Sample sizes (n) and overall statistical

significance (p) from the Kaplan-Meier analyses are provided on each plot. Panel (*e*) shows a forest plot of the

hazard ratios (HR) ± 95% CI based on the stratified CoxPH model for treatment and morphological

deformations (combined antennae and wings deformations; figure S2) on longevity. HR values > 1 indicate

significantly increased hazard. Asterisks denote significance (*** p < 0.001) relative to the 34°C ctrl baseline

group (dashed blue line).

3.3 Morphology

We found evidence that thermal stress treatment during pupal development increased the

likelihood of adults having deformed wings (GLMM: x^2 = 12.791, df = 2, p < 0.01; figures

3a, filled bars). Pupae of the 38°C cyclic treatment developed 3.6-fold more deformed

wings compared to the controls (p < 0.01). This difference was only marginally significant

compared to individuals of the 37°C cyclic treatment (p = 0.08). Individuals of the 37°C

cyclic treatment developed non-significantly more deformed wings compared to the

controls (p > 0.05). We found also no evidence that thermal stress developed significantly

more deformed antennae (x^2 = 2.298, df = 2, p > 0.05; figures 3a, open bars).

While we found a significant effect on adult body size metrics ITD (x^2 = 32.155, df

= 3, p < 0.0001; figure 3b) and dry mass (x^2 = 32.155, df = 3, p < 0.0001; figure 3c), this

effect was primarily sex-dependent (ITD: x^2 = 37.343, df = 1, p < 0.0001; dry mass: x^2 =

32.155, df = 3, p < 0.0001). Although treatment had no effect on the ITD (x^2 = 1.097, df =

2, p > 0.05), there was a weak interaction effect between treatment and sex on adult dry

mass (x^2 = 8.827, df = 2, p < 0.05). Pairwise comparisons revealed strongest effects

between workers of the 34°C control group compared to males of all treatment groups

(34°C ctrl: p < 0.0001; 37°C cyclic: p < 0.0001; 38°C cyclic: p < 0.001). Males of the

control group were also significantly different from workers in the cyclic treatments (37°C

cyclic: p < 0.01; 38°C cyclic: p < 0.01).

3.4 Energy reserves

For the analysis of the effects on the relative lipid content at the end of their adult life, our

model including treatment and sex was not significant (x^2 = 5.16, df = 3, p > 0.05, figure

3d). Within this model, we found a small but significant effect of sex on the relative lipid

content (x^2 = 4.412, df = 1, p < 0.05), whereas treatment had no effect (x^2 = 0.921, df =

2, p > 0.05).

Figure 3. Effects of cyclic thermal stress during pupation on morphological and physiological traits in

adult *B. terrestris*. Panel *(a)* illustrates the proportion of individuals with deformed antennae (open bars;

figure S2*a*) and wings (filled bars; figure S2*b*), both separately analysed. Sample sizes (n) are provided below

for each treatment group. Box plots (*b-d*) show the treatment effects (constant 34°C ctrl, blue; 37°C cyclic,

orange; 38°C cyclic, reddish-purple) on body size, quantified by intertegular distance (ITD) and dry mass, and

relative lipid content as metric for energy reserves, in males (open boxes) and workers (filled boxes).

Individual data points are displayed as circles. Significant differences (p < 0.05), based on post-hoc

comparisons from GLMMs, are indicated by different letters above the bars and boxes (groups sharing a letter

are not significantly different).

4. Discussion

Our study addresses a fundamental knowledge gap concerning the sublethal, long-term

carry-over effects of thermal extremes in insect pollinators (González-Tokman *et al.* 2020;

Ma, Ma & Pincebourde 2021; White & Dillon 2023b). To our knowledge, we provide the

first evidence that thermal stress during pupal development leads to deferred mortality at

the adult life-stage in bumblebees. This finding reveals a novel and previously

unrecognized mechanism that may contribute significantly to pollinator decline.

Specifically, cyclic thermal stress not only caused acute pupal mortality (i.e. reduced

emergence; figure 1*a*) but also resulted in significantly reduced adult longevity (HR = 1.81)

in both workers and males of *B. terrestris* (figures 2*a,b,e*). This finding suggests that

thermal stress during pupation imposes irreversible developmental damage that

undermines adult physiological resilience.

Developmental bottlenecks and irreversible damage

The high sensitivity observed during the pupal phase underscores its role as a critical

developmental bottleneck. The reductions in emergence success from cyclic thermal

stress (15% at 37°C and 30% at 38°C cyclic stress; figure 1*a*) confirm patterns previously

observed under constant stress (Laußer & Kurze 2025). Our observation that heavier

pupae were more likely to emerge (figure 1*b*) supports the idea that the pupae-to-adult

transition might be resource-dependent (Laußer & Kurze 2025). Critically, this suggests

that resource limitations within a colony could unleash negative synergistic interactions,

amplifying the detrimental effects of heatwaves.

Most notably, 38°C cyclic thermal stress led to a higher incidence of deformed

wings (figure 3*a*). While wing morphology is a known indicator of irreparable

developmental damage (Vanderplanck *et al.* 2021; Laußer & Kurze 2025), here we

demonstrate that antennae and wing deformations together can serve as a powerful

hazard indicator (HR = 2.50) that predicts adult longevity (figures 2*e*). Beyond individual

mortality, workers and males with severely deformed wings are unable to fly, which

consequently would reduce colony fitness. In fact, it has already been shown that thermal

stress during development can reduce adult fitness across diverse insect taxa (Zhang,

Rudolf & Ma 2015; Sales, Vasudeva & Gage 2021; Padde *et al.* 2024). Adding to the

consequences of deformed wings, deformed antennae likely impair their ability to detect

scents of nest mates, mating partners (important for males), and floral resources

(important for workers). This idea would be supported by a study demonstrating heat

stress-induced olfactory impairment in adult bumblebee workers (Nooten *et al.* 2024).

Importantly, the observed survival curves showed no significant difference between *in*

vitro reared controls and colony-reared controls, validating the application of our controlled

in vitro rearing approach to natural contexts.

Decoupling fitness costs from adult body size and fat body

Our study provides evidence that deferred mortality was not associated with easily

measurable morphological traits. Despite the direct impact on emergence, we observed

no treatment-dependent effects on body size proxies (ITD or dry mass; figures 3*b,c*), in

contrast to some colony-level studies (Gérard *et al.* 2018; Guiraud *et al.* 2021; Perl *et al.*

2022; Gérard *et al.* 2023). This result suggests that selection pressure from short-term

thermal stress for body size might be more pronounced at earlier larval stages (Wögler &

Kurze 2025) or is insufficient (Laußer & Kurze 2025). Alternatively, it would be plausible

that body size effects observed in stressed colonies are an indirect result of altered worker

nursing and foraging behaviour (Bretzlaff, Kerr & Darveau 2024; Sepúlveda *et al.* 2024),

rather than a direct physiological consequence of the heat itself. This highlights the

importance of our *in vitro* approach isolating the direct physiological effects from

comprehensive colony dynamics (Laußer & Kurze 2025; Wögler & Kurze 2025).

While increased lipid content is advantageous for thermally stressed pupae to

successfully emerge (Laußer & Kurze 2025), we found no significant difference in relative

lipid content at the end of their adult life (figure 3d). This indicates the capacity of surviving

adults to maintain homeostasis of their fat body, which is the primary site for heat shock

protein (HSP) synthesis and storage of energy reserves (Arrese & Soulages 2010; King &

MacRae 2015; Vanderplanck *et al.* 2021), by resource allocation or compensatory

foraging. Hence, any sublethal damage carrying over to adulthood might be genetically,

structurally, and/or physiologically mediated rather than simply due depleted energy

reserves. This aligns with findings where developmental thermal stress indirectly affected

insect adult physiology, such as flight performance and behaviour (Arambourou,

Sanmartín-Villar & Stoks 2017; Galarza *et al.* 2019; Gérard *et al.* 2022; Perl *et al.* 2022).

Conclusion

These findings highlight that sublethal thermal stress during insect development may lead

to profound fitness costs. Our study provides compelling empirical evidence that cyclic

thermal stress during pupation significantly reduces adult longevity. Furthermore, our

findings confirm that 38°C is a critical threshold for development disruption in *B. terrestris*

(Laußer & Kurze 2025; Wögler & Kurze 2025). Although our study simplifies realistic

thermal stress scenarios, heatwaves with peak temperatures exceeding 38°C for several

days are becoming more common in Europe (Barriopedro *et al.* 2011; Rita *et al.* 2020;

Lhotka & Kyselý 2024). Bumblebee species which are less heat-tolerant than *B. terrestris*,

such as *B. lapidarius* and *B. alpinus* (Martinet *et al.* 2021), and colonies nesting above

ground would be expected to be especially at risk. As sublethal carry-over effects are

rarely considered, current prediction models may underestimate the long-term impact of

heatwaves on crucial pollinator populations. To generate a more holistic picture of

environmental stress impact, gaining a nuanced mechanistic understanding of carry-over

effects across diverse insect and life stages, incorporating both individual-level, and for

social insects, also colony-level physiological and life-history data, remains crucial.

Statement of authorship:

M.M.M.: methodology, investigation, data curation. C.K.: conceptualization, methodology,

investigation, data curation, resources, validation, formal analysis, writing – original draft

preparation, writing – review and editing, visualization, supervision, project administration.

Data accessibility statement:

Dataset and R script used for statistical analysis and producing all figures are available from the Dryad Digital Repository: XXX. Electronic supplementary material is available online.

Acknowledgments

We thank Sandra Laußer, Laura Wögler, Ramona Schöpf, and Michael Ott for their assistance in the laboratory. We are grateful to Prof. Dr. Tomer J. Czaczkes for sponsoring us bumblebee colonies. We have used AI for proof-reading.

Funding

This study was carried out without any support of third-party funding.

Competing interests

The authors have no competing interests.

5. References

Arambourou, H., Sanmartín-Villar, I. & Stoks, R. (2017) Wing shape-mediated carry-over effects of a heat wave during the larval stage on post-metamorphic locomotor ability. **184,** 279-291.

Arrese, E.L. & Soulages, J.L. (2010) Insect fat body: energy, metabolism, and regulation. *Annual review of entomology,* **55,** 207-225.

Barriopedro, D., Fischer, E.M., Luterbacher, J., Trigo, R.M. & García-Herrera, R. (2011) The hot summer of 2010: redrawing the temperature record map of Europe. *Science,* **332,** 220-224.

Bretzlaff, T., Kerr, J.T. & Darveau, C.-A. (2023) High temperature sensitivity of bumblebee castes and the colony-level costs of thermoregulation in Bombus impatiens. *Journal of Thermal Biology,* **117,** 103710.

Bretzlaff, T., Kerr, J.T. & Darveau, C.-A. (2024) Handling heatwaves: balancing
thermoregulation, foraging and bumblebee colony success. *Conservation*
Physiology, **12**.

Brooks, M.E., Kristensen, K., Van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A.,
Skaug, H.J., Machler, M. & Bolker, B.M. (2017) glmmTMB balances speed and
flexibility among packages for zero-inflated generalized linear mixed modeling. *The*
R journal, **9**, 378-400.

Cameron, S.A., Lozier, J.D., Strange, J.P., Koch, J.B., Cordes, N., Solter, L.F. & Griswold,
T.L. (2011) Patterns of widespread decline in North American bumble bees.
Proceedings of the National Academy of Sciences, **108**, 662-667.

Christensen, R.H.B. (2019) ordinal—regression models for ordinal data. *R package*
version, **10**, 54.

Galarza, J.A., Dhaygude, K., Ghaedi, B., Suisto, K., Valkonen, J. & Mappes, J. (2019)
Evaluating responses to temperature during pre-metamorphosis and carry-over
effects at post-metamorphosis in the wood tiger moth (*Arctia plantaginis*).
Philosophical Transactions of the Royal Society B: Biological Sciences, **374**,
20190295.

Garcia, R.A., Cabeza, M., Rahbek, C. & Araújo, M.B. (2014) Multiple dimensions of
climate change and their implications for biodiversity. *Science,* **344**, 1247579.

Ghisbain, G., Thiery, W., Massonnet, F., Erazo, D., Rasmont, P., Michez, D. & Dellicour,
S. (2024) Projected decline in European bumblebee populations in the twenty-first
century. *Nature,* **628**, 337-341.

Gilgenreiner, M. & Kurze, C. (2024) Age dominates flight distance and duration, while
body size shapes flight speed in *Bombus terrestris* L.(Hymenoptera: Apidae).
Proceedings of the Royal Society B, **291**, 20241001.

González-Tokman, D., Córdoba-Aguilar, A., Dáttilo, W., Lira-Noriega, A.,
Sánchez-Guillén, R.A. & Villalobos, F. (2020) Insect responses to heat:
physiological mechanisms, evolution and ecological implications in a warming
world. *Biological Reviews,* **95**, 802-821.

Goulson, D., Lye, G.C. & Darvill, B. (2008) Decline and conservation of bumble bees.
Annu. Rev. Entomol., **53**, 191-208.

Guiraud, M., Cariou, B., Henrion, M., Baird, E. & Gérard, M. (2021) Higher developmental
temperature increases queen production and decreases worker body size in the
bumblebee *Bombus terrestris*. *Journal of Hymenoptera Research,* **88**, 39-49.

Gérard, M., Cariou, B., Henrion, M., Descamps, C. & Baird, E. (2022) Exposure to
elevated temperature during development affects bumblebee foraging behavior.
Behavioral Ecology, **33**, 816-824.

Gérard, M., Guiraud, M., Cariou, B., Henrion, M. & Baird, E. (2023) Elevated developmental temperatures impact the size and allometry of morphological traits of the bumblebee Bombus terrestris. *Journal of Experimental Biology,* **226,** jeb245728.

Gérard, M., Michez, D., Debat, V., Fullgrabe, L., Meeus, I., Piot, N., Sculfort, O., Vastrade, M., Smagghe, G. & Vanderplanck, M. (2018) Stressful conditions reveal decrease in size, modification of shape but relatively stable asymmetry in bumblebee wings. *Sci. Rep.,* **8,** 15169.

Hartig, F. (2020) DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression models. *R package version 0.3,* **3**.

Jørgensen, L.B., Ørsted, M., Malte, H., Wang, T. & Overgaard, J. (2022) Extreme escalation of heat failure rates in ectotherms with global warming. *Nature,* **611,** 93-98.

Kassambara, A., Kosinski, M. & Biecek, P. (2016) survminer: Drawing Survival Curves using'ggplot2'. *CRAN: Contributed Packages*.

Kato, Y., Kikuta, S., Barribeau, S.M. & Inoue, M.N. (2022) In vitro larval rearing method of eusocial bumblebee *Bombus terrestris* for toxicity test. *Scientific Reports,* **12,** 15783.

Kerr, J.T., Pindar, A., Galpern, P., Packer, L., Potts, S.G., Roberts, S.M., Rasmont, P., Schweiger, O., Colla, S.R. & Richardson, L.L. (2015) Climate change impacts on bumblebees converge across continents. *Science,* **349,** 177-180.

Kevan, P.G., Rasmont, P. & Martinet, B. (2024) Thermodynamics, thermal performance and climate change: temperature regimes for bumblebee (Bombus spp.) colonies as examples of superorganisms. *Frontiers in Bee Science,* **Volume 2 - 2024**.

King, A.M. & MacRae, T.H. (2015) Insect heat shock proteins during stress and diapause. *Annual review of entomology,* **60,** 59-75.

Klein, A.-M., Vaissiere, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A., Kremen, C. & Tscharntke, T. (2007) Importance of pollinators in changing landscapes for world crops. *P. Roy. Soc. B.,* **274,** 303-313.

Klockmann, M., Kleinschmidt, F. & Fischer, K. (2017) Carried over: Heat stress in the egg stage reduces subsequent performance in a butterfly. *PLoS one,* **12,** e0180968.

Kurze, C., Jenkins, N.E. & Hughes, D.P. (2020) Evaluation of direct and indirect transmission of fungal spores in ants. *Journal of Invertebrate Pathology,* **172**.

Laußer, S. & Kurze, C. (2025) Impact of thermal stress during pupal development in a key pollinator. *Proceedings of the Royal Society B,* **292,** 20252029. *in press* https://doi.org/10.1098/rspb.2025.2029

Lenth, R. & Lenth, M.R. (2018) Package 'lsmeans'. *The American Statistician,* **34,** 216-221.

Lhotka, O. & Kyselý, J. (2024) Three-dimensional analysis reveals diverse heat wave
types in Europe. *Communications Earth & Environment,* **5,** 323.

Ma, C.-S., Ma, G. & Pincebourde, S. (2021) Survive a warming climate: insect responses
to extreme high temperatures. *Annual Review of Entomology,* **66,** 163-184.

Martinet, B., Dellicour, S., Ghisbain, G., Przybyla, K., Zambra, E., Lecocq, T., Boustani,
M., Baghirov, R., Michez, D. & Rasmont, P. (2021) Global effects of extreme
temperatures on wild bumblebees. *Conservation Biology,* **35,** 1507-1518.

Meehl, G.A. & Tebaldi, C. (2004) More intense, more frequent, and longer lasting heat
waves in the 21st century. *Science,* **305,** 994-997.

Nooten, S.S., Korten, H., Schmitt, T. & Kárpáti, Z. (2024) The heat is on: reduced
detection of floral scents after heatwaves in bumblebees. *Proceedings of the Royal*
Society B, **291,** 20240352.

Padde, J.R., Zhou, Y., Chen, Y., Zhu, Y., Yang, Y., Hou, M., Chen, L., Xu, Z., Zhang, D.,
Chen, L. & Ji, M. (2024) Adaptation and carry over effects of extreme sporadic
heat stress in *Culex* mosquitoes. **260,** 107417.

Perl, C.D., Johansen, Z.B., Moradinour, Z., Guiraud, M., Restrepo, C.E., Wen Jie, V.,
Miettinen, A. & Baird, E. (2022) Heatwave-like events during development are
sufficient to impair bumblebee worker responses to sensory stimuli. *Frontiers in*
Ecology and Evolution, **9**.

Potts, S.G., Imperatriz-Fonseca, V., Ngo, H.T., Aizen, M.A., Biesmeijer, J.C., Breeze,
T.D., Dicks, L.V., Garibaldi, L.A., Hill, R. & Settele, J. (2016) Safeguarding
pollinators and their values to human well-being. *Nature,* **540,** 220-229.

Radmacher, S. & Strohm, E. (2011) Effects of constant and fluctuating temperatures on
the development of the solitary bee *Osmia bicornis* (Hymenoptera: Megachilidae).
Apidologie, **42,** 711-720.

Rita, A., Camarero, J.J., Nolè, A., Borghetti, M., Brunetti, M., Pergola, N., Serio, C.,
Vicente-Serrano, S.M., Tramutoli, V. & Ripullone, F. (2020) The impact of drought
spells on forests depends on site conditions: The case of 2017 summer heat wave
in southern Europe. *Global change biology,* **26,** 851-863.

Rolff, J., Johnston, P.R. & Reynolds, S. (2019) Complete metamorphosis of insects.
Philosophical Transactions of the Royal Society B: Biological Sciences, **374,**
20190063.

R Core Team (2024) R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-
project.org/.

Sales, K., Vasudeva, R. & Gage, M.J.G. (2021) Fertility and mortality impacts of thermal
stress from experimental heatwaves on different life stages and their recovery in a
model insect. *Royal Society Open Science,* **8,** 201717.

Sepúlveda, Y., Nicholls, E., Schuett, W. & Goulson, D. (2024) Heatwave-like events affect
drone production and brood-care behaviour in bumblebees. *PeerJ,* **12,** e17135.

Soroye, P., Newbold, T. & Kerr, J. (2020) Climate change contributes to widespread
declines among bumble bees across continents. *Science,* **367,** 685-688.

Theodorou, P., Kühn, O., Baltz, L.M., Wild, C., Rasti, S.L., Bucksch, C.R., Strohm, E.,
Paxton, R.J. & Kurze, C. (2022) Bumble bee colony health and performance vary
widely across the urban ecosystem. *Journal of Animal Ecology,* **91,** 2135-2148.

Therneau, T.M. & Grambsch, P.M. (2000) *Modeling survival data : extending the Cox
model*. Springer, New York.

Vanderplanck, M., Breeze, T., Cini, E., Potts, S.G. & Senapathi, D. (2021) Monitoring bee
health in European agroecosystems using wing morphology and fat bodies. *One
Ecosystem,* **6**.

White, S.A. & Dillon, M.E. (2023a) Climate warming and bumble bee declines: the need to
consider sub-lethal heat, carry-over effects, and colony compensation. *Frontiers in
Physiology,* **Volume 14 - 2023**.

White, S.A. & Dillon, M.E. (2023b) Climate warming and bumble bee declines: the need to
consider sub-lethal heat, carry-over effects, and colony compensation. *Frontiers in
Physiology,* **14**.

Wögler, L. & Kurze, C. (2025) Experimental short-term heatwaves negatively impact body
weight gain and survival during larval development in *Bombus terrestris* L.
(Hymenoptera: Apidae). *Biology Open,* **14**.

Zhang, W., Rudolf, V.H.W. & Ma, C.-S. (2015) Stage-specific heat effects: timing and
duration of heat waves alter demographic rates of a global insect pest. *Oecologia,*
179, 947-957.

559