Strain-dependent differences in the capacity of peste des petits ruminants virus to infect antigen-presenting cells

Aadel Bouziane[a,b]#, Vincent Lasserre[a,b]*, Roger-Junior Eloiflin[a,b], Manon Chambon[a,b], Camille Piro-Mégy[a,b], Samia Guendouz[a,b], Philippe Holzmuller[a,b]*, Philippe Totté[a,b], Arnaud Bataille[a,b]#

[a] CIRAD, UMR ASTRE, F-34398 Montpellier, France

[b] ASTRE, University of Montpellier, CIRAD, INRAE, Montpellier, France

Running head: APC Infection Differences Among Two PPRV Strains

#Address correspondence to Aadel Bouziane (ORCID: https://orcid.org/0009-0007-1090-9110), aadel.bouziane@cirad.fr; address correspondence to Arnaud Bataille (ORCID: https://orcid.org/0000-0002-3508-2144), arnaud.bataille@cirad.fr

*Present address: V. Lasserre, Laboratoire Ecologie et Biologie des Interactions, équipe MHE, UMR CNRS 7267, Université de Poitiers, 5 rue Albert Turpin, 86073 Poitiers Cedex 9, France ; R.-J. Eloiflin, Institut de Génétique Moléculaire de Montpellier (IGMM), Université de Montpellier, Montpellier, France; P. Holzmuller, CIRAD, UMR Selmet, Montpellier, France

The order of authors reflects the relative contributions of the individuals listed up to S. Guendouz. The last three authors, P. Holzmuller, P. Totté, and A. Bataille, contributed to the study in supervisory roles.

Word count: Abstract, 243 words; Importance, 153 words; Introduction, 718 words; Results, 1152 words; Discussion, 1538 words; Materials and Methods, 1279 words.

KEYWORDS: Morbillivirus, monocyte-derived macrophage, monocyte-derived dendritic cell, virulence, host susceptibility

ABSTRACT

The ability of morbilliviruses to infect antigen-presenting cells (APCs) plays a major role in

infection outcome, because these cells transport peste des petits ruminants virus (PPRV) from

the respiratory tract to the lymph nodes before spreading to lymphoid and epithelial tissues.

This study used two cell models, — monocyte-derived macrophages (MoMs) and monocyte-

derived dendritic cells (MoDCs), — from three host species (goat, sheep and cattle) to evaluate

the differential ability of APCs to be infected by two strains of PPRV with contrasting virulence

in goats. Infection was analysed using flow cytometry, infectious titers ($TCID_{50}$/mL), RT-qPCR

and immunofluorescence. Goat and sheep MoDCs proved highly permissive to both strains,

with viral titers at 48 hpi in the range of 10^4–10^5 $TCID_{50}$/mL and a high percentage of infected

cells. Cattle MoDCs were weakly infected and had no detectable viral titer, as expected for this

dead-end host. In comparison, MoMs showed reduced permissiveness to PPRV in goats but not

in sheep. Post-infection supernatants in goat MoMs infected with the low-virulent strain showed

low or undetectable $TCID_{50}$/mL, despite persistence of detectable viral genome up to 96 hpi.

Immunofluorescence revealed double-stranded RNA but no nucleoprotein of the low-virulent

strain in goat MoMs, suggesting a partial engagement of the viral cycle, resulting in a restricted

or semi-permissive infection. Overall, these results suggest that this combined in vitro model

of MoDC and MoM infections would be useful in anticipating host susceptibility and PPRV

strain virulence, providing complementary tools for PPR control efforts.

IMPORTANCE

Peste des petits ruminants (PPR) disease affects goats, sheep and multiple other hosts across

the globe, with major impacts on economies, food security and biodiversity. The sensitivity of

host species to PPRV infection varies considerably, with clinical manifestations and disease

progression being shaped by the viral strain, host species and breed infected. In vitro models

focusing on immune cells, which are the primary target of PPRV could provide a relevant tool

for studying the mechanisms of viral adaptation to the host and cellular antiviral responses,

while contributing to surveillance and control strategies. The results obtained here suggest that

a dual-cell model focusing on macrophages and dendritic cells could be useful for pre-screening

studies aimed at assessing the susceptibility of a given host species to a PPRV strain and

estimating its relative virulence. Testing this in vitro approach in additional host species and a

wider range of strains is required to confirm its relevance.

INTRODUCTION

Peste des petits ruminants virus (PPRV) is a member of the Paramyxoviridae family, belonging

to the *Morbillivirus* genus, which includes the measles virus (MeV), rinderpest virus (RPV) and

canine distemper virus (CDV). PPRV has a linear, non-segmented, single-stranded RNA

genome with a negative polarity [1]. First identified in the Ivory Coast in 1942, PPRV has since

spread extensively, becoming endemic in large regions of Africa, the Middle East, and Asia [1,2],

and has recently emerged in several European countries [3,4]. The virus primarily affects domestic

small ruminants, such as goats and sheep, but it can also infect suids and a variety of wild

species within the order Artiodactyla, including antelopes and gazelles [5].

PPRV is transmitted primarily through direct contact with infected animals, via respiratory

secretions, feces, and contaminated feed or water sources. PPR can cause severe clinical

manifestations in sheep and goats, with high morbidity and mortality, leading to significant

economic losses for farmers. Given its high transmissibility and significant economic impact,

PPR is classified as a notifiable disease by the World Organisation for Animal Health (WOAH,

formerly OIE) [6], and has been the focus of a control programme jointly led by WOAH and the

Food and Agriculture Organization of the United Nations (FAO) since 2015, aiming for its

eradication by 2030 [7]. However, the sensitivity of the host species to PPRV infection varies

considerably, with clinical manifestations and disease progression being shaped by the viral

strain, host species and breed infected[8,9]. A better understanding of PPRV virulence and host

susceptibility is considered a research priority for disease control [10].

The main targets of PPRV and other morbilliviruses are immune cells via the SLAMF1

(Signalling Lymphocytic Activation Molecule Family 1) receptor, also known as CD150 [11,12],

and epithelial cells via the Nectin-4 receptor [13]. At the early stage of infection, morbilliviruses

infect antigen-presenting cells (APCs), such as macrophages and dendritic cells, which

transport the viral particles to lymphoid organs, in particular lymph nodes and tonsils, where

lymphocytes reside [1]. Infection of epithelial cells occurs once viraemia is established, with the

Nectin-4 receptor playing an essential role in viral dissemination and excretion [11,13].

For PPRV, direct infection of APCs in vivo has not yet been demonstrated. However, draining

lymphoid tissues such as the tonsils and lymph nodes, are the first major sites of replication [14].

Recent studies have shown that monocyte-derived dendritic cells are permissive to PPRV in

vitro [15]. Permissiveness is defined as the ability of the virus to replicate and produce infectious

virions. We know that for other morbilliviruses permissiveness varies depending on the cell

type and the virus infecting these cells [16,17]. Due to the importance of APCs in morbillivirus

infection, it is possible that the observed variability in PPRV virulence and host susceptibility

can be related to differences in the viral replication capacity in APCs. Among APCs, dendritic

cells and macrophages are essential for detecting viral infections and initiating innate

immunity[18,19].

So, the study of infection of monocyte-derived macrophages (MoMs) and monocyte-derived

dendritic cells (MoDCs) could be a relevant and reproducible model for comparing the

permissiveness of host species to PPRV. Here, we will focus on the infection of MoDCs and

MoMs from goats, sheep and cattle (the latter being identified as dead-end hosts for PPRV [20])

with two strains of PPRV. The two strains used in this study, Morocco 2008 (MA08) and Ivory

Coast 1989 (IC89), demonstrated different levels of virulence in certain goat breeds [21]. Strain

MA08 belongs to lineage IV of PPRV and was isolated from an Alpine goat that was severely

affected during the 2008 outbreak in Morocco [22]. The IC89 strain belongs to lineage I. It was

isolated from a goat in Ivory Coast in 1989 and is considered to be a mildly virulent strain,

inducing a mild disease in various goat breeds [21,23,24], although it is used as a virulent strain for

the study of immune response in sheep [25].

Based on representative known virulence to goats of the two strains of PPRV, low for IC89 and

high for MA08, the aim of this study was to assess the comparative in vitro replication of these

PPRV strains in both MoMs and MoDCs of goats, sheep and cattle. The research question was

to evaluate whether any difference could be observed between these APCs in relation to PPRV

strain virulence.

RESULTS

1. PPRV infections of MoMs and MoDCs

To determine the infectivity of the two PPRV strains, IC89 and MA08, for macrophages and

dendritic cells, circulating monocytes from the blood of the different animal species studied

were differentiated into dendritic cells (MoDCs) and macrophages (MoMs). These two cell

types were infected with each of the viral strains studied, IC89 and MA08, for 48 hours at a

multiplicity of infection (MOI) of 0.1. Infection was assessed by flow cytometry after

intracellular labelling with an antibody targeting the virus nucleoprotein (anti-NPPRV) coupled

to a fluorochrome.

In goat MoMs, an average of 20% of the cells were positive for the N protein of PPRV strain

IC89, compared with an average of 32.6% for those infected with strain MA08. However, no

significant difference was observed (Figure 1A). In addition, for goat MoDCs, no significant

difference was observed between the two viral strains and in most cases the cells were positive

for NPPRV with a range of 40 to 73% (Figure 1B). Intercellular comparison (i.e. MoMs versus

MoDCs) revealed a significant difference in the rate of NPPRV-positive cells, with MoDCs

being twice as infected as MoMs. This was observed for both PPRV strains (Fig. 2A and 2B).

For sheep MoMs, there was no significant difference in the rate of NPPRV-positive cells

between the two PPRV strains, and the variability of responses was high, with an average rate

of 14.7% for IC89 and 21% for MA08 (Fig. 1A). Similarly, for sheep MoDCs, there was no

significant difference between these two viral strains in terms of percentage of NPPRV-positive

cells. In both cases, the average was 30%, with comparable extreme values (Fig. 1B). A

significant difference was observed between sheep MoMs and MoDCs infected with IC89,

again MoDCs being about twice as infected as MoMs (P<0.0001) (Fig. 2A). In the case of

PPRV strain MA08, the infection rate of MoDCs was significantly higher than in MoMs

(P<0.001), with an average of 30% compared to 22% for MoMs (Fig. 2B).

MoMs and MoDCs from cattle were also infected with IC89 and MA08 strains using the same

parameters. Flow cytometry analysis after intracellular staining of the NPPRV protein revealed

very low levels of NPPRV-positive cells, no matter which cell type and viral strain was used

(Fig. 1A, 1B)

2. Replication of PPRV in MoMs and MoDCs

To study viral replication in MoMs and MoDCs of different hosts, post-infection supernatants

were added to cultures of SLAM-expressing Vero cells (VDS) [26] at dilutions from 10^{-1} to 10^{-7}

with a minimum of four replicates per dilution. The cytopathic effect (CPE) was observed under

the microscope and viral titers were calculated as described in Materials and Methods.

In goats, post-infection supernatants from MoDCs had, on average for both PPRV strains MA08

and IC89, significantly higher titers than supernatants from MoMs (P<0.0001) (Fig. 3A and

3B). For MoMs, the viral titers of the post-infection supernatants for MA08 reached an average

$TCID_{50}$/mL of $3x10^3$ (Fig3B) whereas, for the IC89 strain, the viral titer was barely detectable

with an average $TCID_{50}$/mL of 34 (Fig 3A) including 9 out of 17 replicates where no titer was

detected (Circled in Fig 3A).

In sheep, post-infection supernatants from MoDCs infected with strain MA08 had, on average,

significantly higher titers than those from MoMs infected with MA08 (Fig. 3B) with a $TCID_{50}$

of around $1.2x10^4$ for MoDCs infected with MA08 compared with $9.8x10^2$ for MoMs infected

with MA08 (P<0.0001). There were no significant differences between MoMs and MoDCs

infected with the IC89 strain even though there was a trend towards higher titers for post-

infection supernatants from MoDCs. The IC89 strain yielded a significantly higher mean titer

than the MA08 strain in both MoDCs (P<0.001) and MoMs (P<0.0001) (Supplemental fig. S1).

The titration of 48hpi supernatants of the IC89 strain on sheep MoMs versus goat MoMs,

indicated a significant difference (P<0.0001), with sheep MoMs exhibiting high titers with an

average $TCID_{50}$ of $2x10^4$. In contrast, whereas goat MoMs infected with IC89 had titers that

are barely or not at all measurable (Fig. 3A). In cattle APCs, no viral titer was detected in the

post-infection supernatants (Fig. 3A and 3B).

3. Kinetics of infection of goat MoMs by PPRV IC89 strain

159   a. Infection and replication

After observing low or non-detectable viral titers in supernatants of goat MoMs infected for 48

hours with the IC89 strain, infection kinetics were studied to see if replication was somewhat

delayed for this low virulent strain. Goat MoMs were infected for 24 h, 48 h, 72 h and 96 h, at

an MOI of 0.1. As previously, the rate of NPPRV-positive cells was measured using a flow

cytometer and post-infection supernatants were collected and viral titers were determined.

The results show that the rate of infected cells is relatively stable over time, with an average

NPPRV positivity of around 30% regardless of the infection time. No significant increase in

the percentage of NPPRV-positive cells was observed between 24hpi and 96hpi (Fig. 4A).

The post-infection supernatants from IC89-infected goat MoMs were collected in triplicates per

animal throughout the kinetics, and titrated on VDS. No $TCID_{50}$/mL viral titers could be

detected for any time point during the kinetics, with values remaining below the test detection

threshold (Fig. 4B), and cytopathic effects (CPEs) sometimes observed in a single well of the

first dilution (10^{-1}). The titration plates used for this kinetic analysis were then labelled with the

monoclonal antibody 38.4 targeting the NPPRV coupled with tetramethylrhodamine

isothiocyanate (TRITC) in order to observe the presence or absence of the virus at the cellular

level. We observed the presence of the virus in VDS cells in the first dilution, and sometimes

up to the second and third dilutions (10^{-2} and 10^{-3}). However, this detection occurred in the

absence of a cytopathic effect (CPE), or with CPE limited to a single well in the first dilution,

which is insufficient to calculate a reliable viral titer (Supplemental fig. S2).

b. Detection of viral RNA

To determine the presence of viral genomic RNA, an RT-qPCR was performed on post-

infection supernatants throughout the kinetic. The cycle number (Cq) between 24 hpi and 48hpi

remained stable at around 20.5. A decrease in the Cq value was first observed at 72hpi, falling

to an average of 19.5, with an even more significant decrease at 96hpi, when the Cq averaged

17.8 (Fig. 4C), representing an increase in genetic material of the IC89 strain detected in

supernatants.

4.   Detection of dsRNA and NPPRV protein in goat MoMs and MoDCs

To investigate a possible blockage or slowdown in the replication cycle of the IC89 strain in

goat MoMs, immunofluorescence was used to detect separately double-stranded RNA

(dsRNA), an indicator of the accumulation of aberrant viral RNAs in morbilliviruses, and the

PPRV nucleoprotein (NPPRV), a marker of viral protein expression, in infected MoMs. The

profiles obtained were compared with those observed in goat MoDCs. A positive dsRNA signal

was observed in both cell types in the presence of IC89, although the dsRNA signal appeared

slightly less intense in MoMs (Fig. 5A). The dsRNA signal appeared stronger in goat MoMs

when infected by MA08 (Supplemental fig. S3). NPPRV is detected in goat MoDCs infected

with IC89, but not in goat MoMs (Fig. 5B).

DISCUSSION

PPRV infection can lead to a variety of outcomes ranging from acute symptoms and death of

the infected host to subclinical infection. Surveillance and control measures must be adapted to

provide an efficient response. However, we have limited capacity to predict disease outcomes

in different hosts and prepare accordingly. A better understanding of differences in sensitivity

of the host to viral strains, studied in vitro, would make it possible to anticipate disease impacts

and develop a predictive model that can demonstrate not only the sensitivity of a host to

different PPRV strains but also the differences in virulence expressed by the same strain

depending on the host [27,28]. To move towards this objective, we selected two strains of PPRV

with distinct virulence profiles in goats, mildly virulent IC89 and highly virulent MA08 [24,29,30].

Goat and sheep cells were used because these species are the main natural hosts of PPRV, while

bovine cells served as a relevant negative control, as cattle are considered a dead-end host [20,31].

This in vitro experimental model focused on a comparative overview of PPRV infection in

MoMs and MoDCs in order to describe possible differences in sensitivity and replication profile

between cell types, host species and depending on the viral strain.

Overall, our study shows that MoDCs from goats and sheep are more permissive to PPRV than

MoMs. On average, the rate of infected cells (or NPPRV-positive cells) and viral titers are

higher for MoDCs than for MoMs. In cattle, the absence of infection and of measurable

replication in MoDCs and MoMs confirms the dead-end host status of this species.

Despite significant inter-individual variability observed in the number of cells infected during

the experiment, the measurement of viral replication via titration provided a robust and

biologically meaningful reading for defining permissiveness. The data obtained 48 hours post-

infection at MOI of 0.1 likely reflect the combined effect of several infectious cycles,

incorporating both the efficiency of viral replication and the virus's capacity for secondary

spread in cell cultures. Experiments at lower MOI and at different time post infection may give

additional information on the PPRV primary infectious cycle in both cells, however our results

show that the conditions used in our in vitro experiments are suitable to detect differences of

interest for the overall goal of this study.

No significant differences in viral titers were observed between the IC89 and MA08 viral strains

in goat MoDCs. However, in sheep MoDCs, the mean titer of the IC89 strain was approximately

0.6 $\log_{10}$ higher that of the MA08 strain. MoDCs results concur with the multiple field and

experimental studies showing that cattle are not affected and cannot transmit PPRV, unlike the

main small ruminant hosts [20,32]. In small ruminants, MoDCs confirmed their susceptibility to

PPRV infection, but did not reveal any significant difference between goats and sheep. This

cell model could therefore potentially serve as an indicator of the ability of the virus to infect a

given species.

In contrast to MoDCs, MoMs are less permissive but more discriminating host cells for PPRV. In goats, strain MA08 infected MoMs more efficiently than strain IC89. Oppositely, IC89 produced detectable titers that were significantly higher than MA08 in sheep MoMs. Furthermore, the titers observed with IC89 in sheep MoMs were significantly higher than those obtained in goat MoMs. This suggests that the ability of this strain to replicate will depend on the host species and the type of cell infected. Based on this, and on knowledge of the virulence of these strains in vivo [29,21,24,30,33], we propose that this infection model with MoMs could serve as an indicator of strain-specific virulence in small ruminant breeds and may highlight host-specific signatures of viral adaptation.

For the remainder of the study, infection kinetics were performed on goat MoMs infected with IC89, since the discrepancy was most pronounced in this model (i.e. cellular infection was detected by flow cytometry, but the viral titer was undetectable). The objective was to determine whether the absence of a viral titer was due to a delay in replication or a persistent blockage of the viral cycle. Throughout the kinetics (24h to 96h infection), the rate of infected cells remained stable, averaging between 28 and 32% with no significant increase, suggesting continuous infection of a sub-population of macrophages present in the culture. Titrations of supernatants at all time points remained negative, indicating the absence of measurable infectious virion production. Nevertheless, RT-qPCR revealed a significant decrease in Cq between 48, 72 and 96 hpi, reflecting a gradual accumulation of extracellular viral genomic RNA. It should be noted that RT-qPCR does not distinguish between infectious and non-infectious viral particles. In addition, immunofluorescence microscopy of the titration plates showed that viral antigens were present in the cells because of the NPPR protein detection in VDS cells, at different levels but without CPE, which supports the viral titer determined in supernatants and the cytometry results.

Despite the absence of measurable infectious titer, the detection of viral RNA in the

supernatants suggests an incomplete or an abortive replication cycle. This could result in the

release of non-infectious or defective viral particles, which has already been observed with

several RNA viruses[34–36], including morbilliviruses[37–40].

In negative-sense single-stranded RNA viruses, such as PPRV, viral genome replication is

closely linked to nucleocapsid assembly, meaning that dsRNA is not considered a reliable

indicator of productive replication [41]. Nevertheless, the presence of dsRNA is more likely to be

an indicator of the accumulation of aberrant viral RNA, such as defective interfering RNA (DIs)

or defective viral genomes (DVG)[39]. Such DIs and DVGs can be produced during virus

production, but the contrasting levels of dsRNA observed in MoMs and MoDCs infected with

IC89 suggests differences in the accumulation of these particles during infection. Importantly,

no NPPRV signal was detected in MoMs infected with the IC89 strain, unlike in MoDCs. This

would suggest that the IC89 viral cycle in MoMs appears to be engaged until the accumulation

of aberrant viral RNA forms (DIs or DVGs), while nucleoprotein expression or accumulation

is severely limited or absent. Therefore, while partial expression of the viral genome is

detectable, this does not result in the production of infectious particles, indicating a restricted

or semi-permissive infection of goat MoMs by the IC89 strain.

Such a restriction may be associated with the activation of interferon-dependent antiviral

pathways known to limit viral protein synthesis[42,43].

Further work is needed to precisely understand the observations made within this promising

model in order to evaluate its usefulness. It would be interesting to move towards a multiomics

approach to understand the different mechanisms involved in PPRV infection of MoMs and

MoDCs, in particular those related to innate antiviral immune activity and the link between

different immune signalling pathways and the non-structural proteins C and V of

morbilliviruses. It would also be relevant to quantify the different production of cytokines in

the post-infection supernatants of MoMs and MoDCs, and to perform proteomic analysis of

these supernatants to detect and characterise viral proteins released during infection. Finally,

observing the state of the virion as it leaves the cells under electron microscope would provide

additional structural information on the viral particles produced and their infectious capacity.

In summary, here we have highlighted the difference in permissiveness to PPR virus strains

between MoDCs and MoMs. As demonstrated by the results we obtained with the IC89 and

MA08 strains in the two cell models obtained from three different hosts (goats, sheep and

cattle), in vitro infections of MoDCs may enable us to determine whether a given host is

susceptible to the PPR virus, and MoMs could serve as a cell model to determine the virulence

of a given strain among susceptible hosts. With the IC89 strain, an abortive infection is observed

in goat MoMs, while sheep MoMs are completely permissive. MoMs from goat or sheep also

remain permissive for the MA08 strain. The use of bovine MoMs and MoDCs in this study

constitutes a relevant negative control due to its status as a dead-end host. This combined model

of two cell types (MoDCs and MoMs) would be useful for pre-screening studies to assess the

sensitivity of species to a given strain of PPRV and thus estimate relative virulence. Testing

other PPRV strains with goat, sheep and cattle MoMs and MoDCs, and extending such tests to

other host species will confirm the utility of this model.

Since 2024, Europe has been facing the emergence of PPR, with small ruminants infected in

seven different countries and showing a wide variety of symptoms [44–47]. This recent dynamic

in the spread of PPRV shows that the epidemiological situation remains unstable and that the

virus still crosses borders, making it essential to strengthen risk assessment tools and early

detection capacities. The risk is also present in areas that interface with wildlife, with cases

observed in atypical hosts and endangered wild ruminants in contact with domestic animals [48–

52]. Furthermore, the impact of PPRV on wild species remains poorly understood, as does their

potential role in the transmission or maintenance of the virus and modulation of virulence. As

global eradication efforts led by FAO and WOAH intensify, having tools to anticipate host

susceptibility and strain virulence would be a major asset in guiding early detection and rapid

response measures.

MATERIALS AND METHODS

Cell lines and viral production

All viral strains were produced from Vero expressing the canine SLAM receptor (or Vero-Dog-

SLAM cells, VDS cells) [26]. VDS cell were cultured in complete Dulbecco's Modified Eagle's

Medium (DMEM, Gibco) supplemented with 10% foetal bovine serum (FBS, Dominique

Dutscher). Virus stocks were harvested from infected cell supernatants, clarified by

centrifugation, and stored at -80°C until use. The PPRV strains IC89 (lineage I) and MA08

(lineage IV) are from productions used in previous experimental studies [21]. Any risk of genetic

drift of PPRV strains during cell passages was assessed by sequencing the full genome of the

virus production used for the infection experiments using a protocol described elsewhere[53].

Genomic comparison showed that no mutations were detected following the passages required

for production.

Differentiation of macrophages and dendritic cells from CD14+ monocytes

Peripheral blood samples were obtained from clinically healthy goats (*Capra hircus*, N = 3),

sheep (*Ovis aries*, N = 3) and cattle (*Bos taurus*, N = 6) housed at the CIRAD animal facility

(Montpellier, France). Animal care and handling followed institutional ethical guidelines and

was approved by the Languedoc-Roussillon regional ethics committee (French CE-LR#36) in

an authorised project using animals for scientific purposes APAFIS#2628.

Peripheral blood mononuclear cells (PBMCs) were isolated from the whole blood of the

animals studied by Ficoll-Hypaque density gradient centrifugation using Histopaque-1077

(Sigma-Aldrich), following standard protocols [54]. After collection, viable PBMCs were counted

in a hemocytometer using the Trypan Blue exclusion method.

CD14+ monocytes were positively selected using CD14 MicroBeads UltraPure (Miltenyi

Biotec) and LS MACS columns, according to the manufacturer's instructions. After elution,

CD14+ monocytes were cultured in Roswell Park Memorial Institute medium and GlutaMAX^{TM}

(RPMI 1640 + GlutaMAX^{TM}, Gibco^{TM}) supplemented with 10% FBS and 1% antibiotics

(Penicillin-Streptomycin). A minimum purity of 80% CD14+ monocytes is required to ensure

homogeneous differentiation into MoMs or MoDCs (Supplemental fig. S4).

For differentiation into monocyte-derived dendritic cells (MoDCs), cultures were supplemented

with in-house bovine recombinant IL-4 and GM-CSF (50 ng/ml) From plasmid prepared with

EndoFree Plasmid Purification Maxi Kit Qiagen (reference 12362), cytokines were then

produced by transfection on HEK 293T cells. For the differentiation into monocyte-derived

macrophages (MoMs), no exogenous growth factors were added and monocytes were allowed

to spontaneously differentiate into macrophages under standard culture conditions. This

approach, although less controlled than cytokine-supplemented protocols, mimics a more

physiological environment and results in a heterogeneous macrophage population[55,56].

Differentiation was carried out for six to seven days with a medium change on day two or three

(Supplemental fig. S5). In goats, characterization based on CD205 marker expression was

performed to distinguish monocytes from cells differentiated into MoMs or MoDCs

(Supplemental fig. S6A). Similarly, the CD209 marker was used in cattle[57] (Supplemental fig.

S6B). To distinguish MoMs from MoDCs in goats and sheep, the CD172a marker was used

(Supplemental fig. S6C). In addition, SLAMF9 receptor (homolog of SLAMF1) labeling was

also performed on goat cells (Supplemental fig S6D). Viral infection assays were performed at

the end of the differentiation period. In addition to phenotyping, microscopic observation

(Supplemental fig. S5) confirmed that the cells had been differentiated in agreement with

previous work[15,57,58].

Viral infection

MoDCs and MoMs were infected with either MA08 or IC89 strains at a multiplicity of infection

(MOI) of 0.1. The cells were incubated with virus for one hour at 37°C in serum-free RPMI.

After incubation, cells were washed with PBS and cultured for 48 hours in RPMI supplemented

with 2% FBS. 48 hours post-infection, the culture supernatants were harvested, aliquoted, and

stored at –80°C for downstream analysis.

Flow cytometry

Following infection, adherent cells in culture were detached using TrypLE[TM] Express Enzyme

1X (Gibco) at 37°C for 10 minutes. Cells were harvested into PBS, centrifuged at 300 x g) for

5 minutes at 4°C, and fixed with 4% paraformaldehyde (or 4% PFA) for 10 minutes at 4°C.

For intracellular PPRV detection, cells were permeabilized with 0.3% saponin and stained with

FITC-conjugated monoclonal antibody (clone 38.4) targeting the N nucleoprotein of PPRV

developed by G. Libeau [59] and produced by ProteoGenix (Schiltigheim, France). No secondary

antibody was required due to conjugation of Mab 38.4 with FITC. Stained cells were analyzed

using a BD FACSCanto II flow-cytometer (BD Biosciences) and FlowJo_V10 software

(Supplemental fig. S7).

For phenotyping, the primary antibodies anti-CD14 (CAM36A), anti-CD205 (ILA114A), anti-

CD209 (209MD26A), anti-CD172a (DH59B) and anti-SLAMF9 CACT206A were provided

by WSU Monoclonal Antibody Center Veterinary Microbiology and Pathology (Washington State University – Pullman). The secondary antibodies used, rat anti-mouse IgG1, eBioscience™, PE-Cyanine7 Clone M1-14D12 (Reference: 24-4015-82), Invitrogen by Thermo Fisher Scientific and Alexa Fluor™ goat anti-mouse IgG2a (Reference: A21241), Invitrogen by Thermo Fisher Scientific.

Titration of PPRV

Viral production and cell culture supernatants were titrated on VDS cells seeded in 96-well plates using the Spearman-Kärber method [60–63]. Infectious titers were calculated as $TCID_{50}$/mL using the $TCID_{50}$ calculator developed by Marco Binder. (v2.1—20-01-2017_MB* by Marco Binder; adapted @ TWC https://www.klinikum.uni-heidelberg.de/fileadmin/inst_hygiene/molekulare_virologie/Downloads/TCID50_calculator_v2_17-01-20_MB.xlsx). The titration plates were marked with antibody 38:4 (anti-NPPRV) coupled with TRITC in order to observe the presence or absence of the virus (Supplemental fig. S2), according to the Eloiflin et al. [64] protocol.

RNA extraction from supernatant and real-time qPCR analyses

Viral RNA was extracted from supernatants using the IndiMag Pathogen Kit (Indical Biosciences), following the manufacturer's protocol. RNA isolation was performed on the IDEAL™ 32 automated extraction system (Innovative Diagnostics), based on magnetic bead technology.

RNA of PPRV present in the supernatants was quantified using real-time PCR by partial

amplification of the N protein gene using the ID GeneTM peste des petits ruminants Duplex kit

(Innovative Diagnostics) and then performed with LightCycler (Roche).

Immunofluorescence staining and inverted microscopy

Cells were fixed at 48 hpi using 4% paraformaldehyde (PFA) for 20 minutes at room

temperature. After fixation, permeabilization was performed with 0.1% Triton X-100 in 1X

PBS for 5 minutes at room temperature. The cells were then incubated in a buffer composed of

1X PBS, 10% FBS and 1 g BSA per 50 mL for 30 minutes. Primary staining was performed

with anti-dsRNA monoclonal antibody J2 (IgG2a, 1 mg/mL, SCICONS™, Jena Bioscience),

diluted 1:300, and incubated for 2 hours at room temperature. For detection, cells were

incubated with DAPI diluted 1:1000 (Sigma-Aldrich, MBD0015) and either a Donkey anti-

Mouse IgG secondary antibody (H+L) Alexa Fluor™ 555 secondary antibody (2 mg/mL)

applied at 4 µg/mL (Thermo Fisher Scientific, Cat# A-31570, RRID:AB_2536180), or with

TRITC-coupled anti-NPPRV (Mab 38.4) antibody diluted 1:250. In both cases, incubation took

place for 1 hour at room temperature, in the dark. The samples were then imaged using an

AXIOVERT A1 inverted microscope (Zeiss, France) with Archimed 6.1.4 software, and

analysed with ImageJ (version supplied with Java 1.8.0_172 64-bit).

Statistical analysis

Statistical analyses and histograms were performed using GraphPad Prism v.10.6.1 and R

software v.4.5.2. For the analysis of infection rates based on the number of cells infected by the

two strains studied on the different hosts, generalized linear mixed models (GLMM) with a

binomial distribution were used. Species, cell type, and strain were included as fixed effects,

and animal identify and/or experimental batches was included as a random effect to account for

repeated measurements within individuals.

Viral titers were analysed using generalized linear mixed-effects models with a negative

binomial distribution. Species, cell type and strain were included as fixed effects and animal

identify and/or experimental batches as a random effect.

For kinetic experiments, the analysis was performed using generalised mixed models with

binomial or beta-binomial distribution in cases of overdispersion, with post-infection time (hpi)

as a fixed effect and animal identify and/or experimental batches as a random effect.

RT-qPCR data from kinetics were analysed using linear mixed models (LMM) with the

restricted maximum likelihood (REML) method, with hpi as a fixed effect and animal identify

as a random effect to account for repeated measurements in the same animal.

Overdispersion and zero-inflation were evaluated where appropriate. Pairwise comparisons

between groups were performed using Tukey's multiple comparison test with adjustment for

multiple testing.

436
437
438 Figure legends
439
440 **Figure 1: Percentage of MoM (A) and MoDC (B) from cow, sheep and goat, positive for PPRV at 48hpi after in vitro infection with**
441 **IC89 or MA08 strain.** The MOCK group corresponds to uninfected control cells. Data are shown as individual biological replicates from a
442 minimum of three animals per species and per condition with mean $\pm$ SD; ns=non-significant; *P<0.05; Statistical analysis was performed
443 using a generalized linear mixed-effects model (GLMM). Post-hoc comparisons were conducted using estimated marginal means with Tukey
444 adjustment for multiple comparisons.

445
Figure 2: Percentage of MoM and MoDC from sheep and goat, positive for PPRV at 48hpi after in vitro infection with IC89 (A) or
MA08 (B) strain. The MOCK group corresponds to uninfected control cells. Data are shown as individual biological replicates from a
minimum of three animals per species and per condition with mean $\pm$ SD; ***P<0.001; ****P<0.0001; Statistical analysis was performed
using a generalized linear mixed-effects model (GLMM). Post-hoc comparisons were conducted using estimated marginal means with Tukey
adjustment for multiple comparisons.

Figure 3: Viral titers (TCID$_{50}$/mL) of IC89 (A) and MA08 (B) at 48 hpi in cow, sheep and goat MoMs and MoDCs. Nine of seventeen
replicates were below the limit of detection in goat MoMs infected with IC89 (circled in A). Data are shown as individual biological replicates
from a minimum of three animals per species and per condition with mean $\pm$ SD; ns=non-significant; ****P<0.0001. Replicates from goat
with no titer observed are highlighted with a circle. Statistical analysis was performed on viral titers using a generalized linear mixed-effects

model (GLMM) fitted with a negative binomial distribution. Post-hoc comparisons were performed using estimated marginal means with Tukey adjustment for multiple comparisons.

Figure 4: (A) Kinetics of infection of goat MoMs by strain IC89 of PPRV. Percentages of NPPRV+ cells were determined by flow cytometry at 24, 48, 72, and 96hpi. Data are shown as individual biological replicates with 3 goats and with minimum n=7 replicates, mean ± SD. A beta-binomial GLMM revealed no significant effect of time post-infection on infection rates (all P > 0.76). Post-hoc pairwise comparisons using Tukey-adjusted estimated marginal means confirmed the absence of significant differences between 24, 48, 72 and 96 hpi (all adjusted P > 0.99). **(B) Viral titration kinetics of MoMs supernatants from goats infected with strain IC89 at 24, 48, 72 and 96hpi.** Data are shown as individual biological replicates from 3 goats with n=9 replicates per time point. **(C) Kinetics of viral shedding in MoMs supernatant infected by IC89 strain at 24h, 48h, 72h and 96h.Viral shedding was monitored by RT-qPCR (expressed in Cq) detecting the presence of the PPRV N gene.** Data are shown as individual biological replicates from 3 goats with n ranging from 12 to 20 replicates per time point, mean ± SD; *P<0.05; P<0.0001. Statistical analysis was performed using a linear mixed-effects model (LMM) fitted by restricted maximum likelihood (REML). Pairwise comparisons were conducted using estimated marginal means with Tukey adjustment for multiple comparisons (Kenward–Roger approximation for degrees of freedom).

Figure 5: Immunofluorescent detection of viral double-stranded RNA (dsRNA) (A) and Nucleoprotein PPRV protein (NPPRV) (B) in goat MoMs and MoDCs infected with IC89. Infected and uninfected (MOCK) MoMs and MoDCs were stained with DAPI to identify cell nuclei, dsRNA (A) and NPPRV (B) are detected here in red, scale bar = 25 µm.

ACKNOWLEDGMENTS

A.Bo., A.Ba. and S.G. are supported by a grant (SI2.756606) from the European Commission Directorate General for Health and Food Safety awarded to the European Union Reference Laboratory for peste des petits ruminants (EURL-PPR, https://eurl-ppr.cirad.fr/).

The authors would like to thank all the staff at UMR ASTRE (CIRAD) for their technical support throughout this study, and in particular Olivier Kwiatek for his help in analysing the titration plates, and Hélène Jourdan for her valuable advice on statistical analyses. We also express our gratitude to the staff of the CIRAD animal facility in Baillarguet — Géraldine Bossard, Thierry Maccotta and Jade Fudaly — for their invaluable assistance. Finally, we would like to thank Aymeric Neyret for his expert advice on immunofluorescence microscopy analyses.

Conceptualisation: A.Bo., V.L., R.-J.E., M.C., P.H., P.T. and A.Ba.; data curation: A.Bo, V.L., M.C. and R.-J.E.; formal analysis: A.Bo., V.L., and P.T.; funding acquisition: A.Ba supervision: A.Ba, P.T. and P.H.; investigation: A.Bo., V.L., M.C. and R.-J.E.; methodology:

487 A.Bo, V.L., R.-J.E, P.T., C.P-G. and S.G.; validation: A.Ba., P.T. and P.H.; visualisation:

488 A.Bo., P.T., P.H. and A.Ba.; writing (original draft preparation): A.Bo.; writing (review and

489 editing): A.Bo, P.H., P.T. and A.Ba.

Declarations of interest

491 The authors declare no conflict of interest.
492

REFERENCES

(1) Kumar, N.; Maherchandani, S.; Kashyap, S. K.; Singh, S. V.; Sharma, S.; Chaubey, K.
K.; Ly, H. Peste Des Petits Ruminants Virus Infection of Small Ruminants: A
Comprehensive Review. *Viruses* **2014**, *6* (6), 2287–2327.
https://doi.org/10.3390/v6062287.

(2) Banyard, A. C.; Parida, S.; Batten, C.; Oura, C.; Kwiatek, O.; Libeau, G. Global
Distribution of Peste Des Petits Ruminants Virus and Prospects for Improved Diagnosis
and Control. *J. Gen. Virol.* **2010**, *91* (12), 2885–2897.
https://doi.org/10.1099/vir.0.025841-0.

(3) Guendouz, S.; Kwiatek, O.; Kirtzalidou, A.; Katsifa, A.; Gianniou, M.; Ancuceanu, C.;
Ghiță, M.; Mortasivu, C. L.; Zdravkova, A.; Kostov, I.; Ivanova, E.; Bărbuceanu, F.;
Tasioudi, K. E.; Bataille, A. Genomic Analysis of Peste Des Petits Ruminants Virus in
Europe: Common Origin for Emergence in Greece, Romania, and Bulgaria. *Infect.*
Genet. Evol. **2025**, *132*, 105774. https://doi.org/10.1016/j.meegid.2025.105774.

(4) Donduashvili, M.; Goginashvili, K.; Toklikishvili, N.; Tigilauri, T.; Gelashvili, L.; Avaliani,
L.; Khartskhia, N.; Loitsch, A.; Bataille, A.; Libeau, G.; Diallo, A.; Dundon, W. G.
Identification of Peste Des Petits Ruminants Virus, Georgia, 2016. *Emerg. Infect. Dis.*
2018, *24* (8), 1576–1578. https://doi.org/10.3201/eid2408.170334.

(5) Bataille, A.; Kwiatek, O.; Belfkhi, S.; Mounier, L.; Parida, S.; Mahapatra, M.; Caron, A.;
Chubwa, C. C.; Keyyu, J.; Kock, R.; Jones, B. A.; Libeau, G. Optimization and
Evaluation of a Non-Invasive Tool for Peste Des Petits Ruminants Surveillance and
Control. *Sci. Rep.* **2019**. https://doi.org/10.1038/s41598-019-41232-y.

(6) *Peste des petits ruminants - WOAH - World Organisation for Animal Health*.
https://www.woah.org/en/disease/peste-des-petits-ruminants/ (accessed 2025-02-10).

(7) Jones, B. A.; Rich, K. M.; Mariner, J. C.; Anderson, J.; Jeggo, M.; Thevasagayam, S.;
Cai, Y.; Peters, A. R.; Roeder, P. The Economic Impact of Eradicating Peste des Petits
Ruminants: A Benefit-Cost Analysis. *PLOS ONE* **2016**, *11* (2), e0149982.
https://doi.org/10.1371/journal.pone.0149982.

(8) Dou, Y.; Liang, Z.; Prajapati, M.; Zhang, R.; Li, Y.; Zhang, Z. Expanding Diversity of
Susceptible Hosts in Peste Des Petits Ruminants Virus Infection and Its Potential

Mechanism Beyond. *Front. Vet. Sci.* **2020**, *7*, 66.
https://doi.org/10.3389/fvets.2020.00066.

(9) Torsson, E. Peste des petits ruminants virus—in the field and in the host. *Acta Univ.*
Agric. Sueciae **2019**, No. 2019:80.

(10) Bataille, A.; Baron, M. D. Rinderpest and Peste Des Petits Ruminants: State of Play in
Disease Eradication Efforts. *Rev. Sci. Tech. Int. Off. Epizoot.* **2024**, *Special Edition*, 43–
52. https://doi.org/10.20506/rst.SE.3557.

(11) Parida, S.; Muniraju, M.; Mahapatra, M.; Muthuchelvan, D.; Buczkowski, H.; Banyard, A.
C. Peste Des Petits Ruminants. *Vet. Microbiol.* **2015**, *181* (1), 90–106.
https://doi.org/10.1016/j.vetmic.2015.08.009.

(12) Adombi, C. M.; Lelenta, M.; Lamien, C. E.; Shamaki, D.; Koffi, Y. M.; Traoré, A.; Silber,
R.; Couacy-Hymann, E.; Bodjo, S. C.; Djaman, J. A.; Luckins, A. G.; Diallo, A. Monkey
CV1 Cell Line Expressing the Sheep–Goat SLAM Protein: A Highly Sensitive Cell Line
for the Isolation of Peste Des Petits Ruminants Virus from Pathological Specimens. *J.*
Virol. Methods **2011**, *173* (2), 306–313. https://doi.org/10.1016/j.jviromet.2011.02.024.

(13) Birch, J.; Juleff, N.; Heaton, M. P.; Kalbfleisch, T.; Kijas, J.; Bailey, D. Characterization
of Ovine Nectin-4, a Novel Peste Des Petits Ruminants Virus Receptor. *J. Virol.* **2013**,
87 (8), 4756–4761. https://doi.org/10.1128/JVI.02792-12.

(14) Pope, R. A.; Parida, S.; Bailey, D.; Brownlie, J.; Barrett, T.; Banyard, A. C. Early Events
Following Experimental Infection with Peste-Des-Petits Ruminants Virus Suggest
Immune Cell Targeting. *PLOS ONE* **2013**, *8* (2), e55830.
https://doi.org/10.1371/journal.pone.0055830.

(15) Rodríguez-Martín, D.; García-García, I.; Martín, V.; Rojas, J. M.; Sevilla, N. A
Morbillivirus Infection Shifts DC Maturation Toward a Tolerogenic Phenotype to
Suppress T Cell Activation. *J. Virol.* **2022**, *96* (18), e01240-22.
https://doi.org/10.1128/jvi.01240-22.

(16) Pohl, C.; Shishkova, J.; Schneider-Schaulies, S. Viruses and Dendritic Cells: Enemy
Mine. *Cell. Microbiol.* **2007**, *9* (2), 279–289. https://doi.org/10.1111/j.1462-
5822.2006.00863.x.

(17) Hou, W.; Armstrong, N.; Obwolo, L. A.; Thomas, M.; Pang, X.; Jones, K. S.; Tang, Q.
Determination of the Cell Permissiveness Spectrum, Mode of RNA Replication, and
RNA-Protein Interaction of Zika Virus. *BMC Infect. Dis.* **2017**, *17* (1), 239.
https://doi.org/10.1186/s12879-017-2338-4.

(18) Roquilly, A.; Mintern, J. D.; Villadangos, J. A. Spatiotemporal Adaptations of
Macrophage and Dendritic Cell Development and Function. *Annu. Rev. Immunol.* **2022**,
40 (Volume 40, 2022), 525–557. https://doi.org/10.1146/annurev-immunol-101320-
031931.

(19) Mehrani, Y.; Morovati, S.; Keivan, F.; Tajik, T.; Forouzanpour, D.; Shojaei, S.; Bridle, B.
W.; Karimi, K. Dendritic Cells and Their Crucial Role in Modulating Innate Lymphoid
Cells for Treating and Preventing Infectious Diseases. *Pathogens* **2025**, *14* (8), 794.
https://doi.org/10.3390/pathogens14080794.

(20) Schulz, C.; Fast, C.; Wernery, U.; Kinne, J.; Joseph, S.; Schlottau, K.; Jenckel, M.;
Höper, D.; Patteril, N. A. G.; Syriac, G.; Hoffmann, B.; Beer, M. Camelids and Cattle Are

Dead-End Hosts for Peste-Des-Petits-Ruminants Virus. *Viruses* **2019**, *11* (12), 1133.
https://doi.org/10.3390/v11121133.

(21) Enchery, F.; Hamers, C.; Kwiatek, O.; Gaillardet, D.; Montange, C.; Brunel, H.;
Goutebroze, S.; Philippe-Reversat, C.; Libeau, G.; Hudelet, P.; Bataille, A. Development
of a PPRV Challenge Model in Goats and Its Use to Assess the Efficacy of a PPR
Vaccine. *Vaccine* **2019**, *37* (12), 1667–1673.
https://doi.org/10.1016/j.vaccine.2019.01.057.

(22) Muniraju, M.; El Harrak, M.; Bao, J.; Ramasamy Parthiban, A. B.; Banyard, A. C.;
Batten, C.; Parida, S. Complete Genome Sequence of a Peste Des Petits Ruminants
Virus Recovered from an Alpine Goat during an Outbreak in Morocco in 2008. *Genome*
Announc. **2013**, *1* (3), 10.1128/genomea.00096-13.
https://doi.org/10.1128/genomea.00096-13.

(23) Chard, L. S.; Bailey, D. S.; Dash, P.; Banyard, A. C.; Barrett, T. Full Genome
Sequences of Two Virulent Strains of *Peste-Des-Petits Ruminants Virus*, the Côte
d'Ivoire 1989 and Nigeria 1976 Strains. *Virus Res.* **2008**, *136* (1), 192–197.
https://doi.org/10.1016/j.virusres.2008.04.018.

(24) Eloiflin, R.-J.; Grau-Roma, L.; Python, S.; Mehinagic, K.; Godel, A.; Libeau, G.;
Summerfield, A.; Bataille, A.; García-Nicolás, O. Comparative Pathogenesis of Peste
Des Petits Ruminants Virus Strains of Difference Virulence. *Vet. Res.* **2022**, *53* (1), 57.
https://doi.org/10.1186/s13567-022-01073-6.

(25) Rojas, J. M.; Moreno, H.; Valcárcel, F.; Peña, L.; Sevilla, N.; Martín, V. Vaccination with
Recombinant Adenoviruses Expressing the Peste Des Petits Ruminants Virus F or H
Proteins Overcomes Viral Immunosuppression and Induces Protective Immunity against
PPRV Challenge in Sheep. *PLOS ONE* **2014**, *9* (7), e101226.
https://doi.org/10.1371/journal.pone.0101226.

(26) Seki, F.; Ono, N.; Yamaguchi, R.; Yanagi, Y. Efficient Isolation of Wild Strains of Canine
Distemper Virus in Vero Cells Expressing Canine SLAM (CD150) and Their Adaptability
to Marmoset B95a Cells. *J. Virol.* **2003**, *77* (18), 9943–9950.
https://doi.org/10.1128/jvi.77.18.9943-9950.2003.

(27) Couacy-Hymann, E.; Bodjo, C.; Danho, T.; Libeau, G.; Diallo, A. Evaluation of the
Virulence of Some Strains of Peste-Des-Petits-Ruminants Virus (PPRV) in
Experimentally Infected West African Dwarf Goats. *Vet. J.* **2007**, *173* (1), 178–183.
https://doi.org/10.1016/j.tvjl.2005.08.020.

(28) Diop, M.; Sarr, J.; Libeau, G. Evaluation of Novel Diagnostic Tools for Peste Des Petits
Ruminants Virus in Naturally Infected Goat Herds. *Epidemiol. Infect.* **2005**, *133* (4),
711–717. https://doi.org/10.1017/S0950268805003729.

(29) Eloiflin, R.-J.; Grau-Roma, L.; Lasserre, V.; Python, S.; Talker, S.; Totte, P.; García-
Nicolás, O.; Summerfield, A.; Bataille, A. Peste Des Petits Ruminants Virus Virulence Is
Associated with an Early Inflammatory Profile in the Tonsils and Cell Cycle Arrest in
Lymphoid Tissue. *Microbiol. Spectr.* **2025**, *13* (4), e03124-24.
https://doi.org/10.1128/spectrum.03124-24.

(30) Eloiflin, R.-J.; Auray, G.; Python, S.; Rodrigues, V.; Seveno, M.; Urbach, S.; El Koulali,
K.; Holzmuller, P.; Totte, P.; Libeau, G.; Bataille, A.; Summerfield, A. Identification of
Differential Responses of Goat PBMCs to PPRV Virulence Using a Multi-Omics

Approach. *Front. Immunol.* **2021**, *12*, 745315.
https://doi.org/10.3389/fimmu.2021.745315.

(31) Aklilu, F.; Ashenafi, H.; Kassa, T.; Chaka, H.; Sibhatu, D.; Shegu, D.; Mohammed, A.
A.; Belaineh, R.; Kidane, M.; Asgedom, H.; Chibssa, T.; Mekonnen, G.; Sirak, A.;
Gebredufe, S.; Schulz, C.; Herzog, C. M.; Kapur, V. Comparative Pathogenesis of
Ethiopia/Habru/2014 Lineage-IV Peste Des Petits Ruminants Virus in Goats and Cattle.
BMC Vet. Res. **2024**, *20* (1), 473. https://doi.org/10.1186/s12917-024-04313-3.

(32) Herzog, C. M.; Aklilu, F.; Sibhatu, D.; Shegu, D.; Belaineh, R.; Mohammed, A. A.;
Kidane, M.; Schulz, C.; Willett, B. J.; Cleaveland, S.; Bailey, D.; Peters, A. R.; Cattadori,
I. M.; Hudson, P. J.; Asgedom, H.; Buza, J.; Forza, M. S.; Chibssa, T. R.; Gebre, S.;
Juleff, N.; Bjørnstad, O. N.; Baron, M. D.; Kapur, V. Empirical and Model-Based
Evidence for a Negligible Role of Cattle in Peste Des Petits Ruminants Virus
Transmission and Eradication. *Commun. Biol.* **2024**, *7* (1), 937.
https://doi.org/10.1038/s42003-024-06619-2.

(33) Rojas, J. M.; Avia, M.; Pascual, E.; Sevilla, N.; Martín, V. Vaccination with Recombinant
Adenovirus Expressing Peste Des Petits Ruminants Virus-F or -H Proteins Elicits T Cell
Responses to Epitopes That Arises during PPRV Infection. *Vet. Res.* **2017**, *48* (1), 79.
https://doi.org/10.1186/s13567-017-0482-x.

(34) Dimmock, N. J.; Easton, A. J. Defective Interfering Influenza Virus RNAs: Time To
Reevaluate Their Clinical Potential as Broad-Spectrum Antivirals? *J. Virol.* **2014**, *88*
(10), 5217–5227. https://doi.org/10.1128/jvi.03193-13.

(35) Xu, J.; Sun, Y.; Li, Y.; Ruthel, G.; Weiss, S. R.; Raj, A.; Beiting, D.; López, C. B.
Replication Defective Viral Genomes Exploit a Cellular Pro-Survival Mechanism to
Establish Paramyxovirus Persistence. *Nat. Commun.* **2017**, *8* (1), 799.
https://doi.org/10.1038/s41467-017-00909-6.

(36) Vignuzzi, M.; López, C. B. Defective Viral Genomes Are Key Drivers of the Virus–Host
Interaction. *Nat. Microbiol.* **2019**, *4* (7), 1075–1087. https://doi.org/10.1038/s41564-019-
0465-y.

(37) Bora, M.; Yousuf, R. W.; Dhar, P.; Singh, R. P. An Overview of Process Intensification
and Thermo Stabilization for Upscaling of Peste Des Petits Ruminants Vaccines in View
of Global Control and Eradication. *VirusDisease* **2018**, *29* (3), 285–296.
https://doi.org/10.1007/s13337-018-0455-3.

(38) Mura, M.; Combredet, C.; Najburg, V.; Sanchez David, R. Y.; Tangy, F.; Komarova, A.
V. Nonencapsidated 5′ Copy-Back Defective Interfering Genomes Produced by
Recombinant Measles Viruses Are Recognized by RIG-I and LGP2 but Not MDA5. *J.*
Virol. **2017**, *91* (20), 10.1128/jvi.00643-17. https://doi.org/10.1128/jvi.00643-17.

(39) Pfaller, C. K.; Mastorakos, G. M.; Matchett, W. E.; Ma, X.; Samuel, C. E.; Cattaneo, R.
Measles Virus Defective Interfering RNAs Are Generated Frequently and Early in the
Absence of C Protein and Can Be Destabilized by Adenosine Deaminase Acting on
RNA-1-Like Hypermutations. *J. Virol.* **2015**, *89* (15), 7735–7747.
https://doi.org/10.1128/jvi.01017-15.

(40) Calain, P.; Roux, L. Generation of Measles Virus Defective Interfering Particles and
Their Presence in a Preparation of Attenuated Live-Virus Vaccine. *J. Virol.* **1988**, *62* (8),
2859–2866. https://doi.org/10.1128/JVI.62.8.2859-2866.1988.

(41) Weber, F.; Wagner, V.; Rasmussen, S. B.; Hartmann, R.; Paludan, S. R. Double-
Stranded RNA Is Produced by Positive-Strand RNA Viruses and DNA Viruses but Not
in Detectable Amounts by Negative-Strand RNA Viruses. *J. Virol.* **2006**, *80* (10), 5059–
5064. https://doi.org/10.1128/jvi.80.10.5059-5064.2006.

(42) Hsu, J. C.-C.; Laurent-Rolle, M.; Cresswell, P. Translational Regulation of Viral RNA in
the Type I Interferon Response. *Curr. Res. Virol. Sci.* **2021**, *2*, 100012.
https://doi.org/10.1016/j.crviro.2021.100012.

(43) Smart, A.; Gilmer, O.; Caliskan, N. Translation Inhibition Mediated by Interferon-
Stimulated Genes during Viral Infections. *Viruses* **2024**, *16* (7), 1097.
https://doi.org/10.3390/v16071097.

(44) Guendouz, S.; Kwiatek, O.; Kirtzalidou, A.; Katsifa, A.; Gianniou, M.; Ancuceanu, C.;
Ghiță, M.; Mortasivu, C. L.; Zdravkova, A.; Kostov, I.; Ivanova, E.; Bărbuceanu, F.;
Tasioudi, K. E.; Bataille, A. Genomic Analysis of Peste Des Petits Ruminants Virus in
Europe: Common Origin for Emergence in Greece, Romania, and Bulgaria. *Infect.*
Genet. Evol. **2025**, *132*, 105774. https://doi.org/10.1016/j.meegid.2025.105774.

(45) Iacobescu-Marițescu, R.; Morar, A.; Herman, V.; Tîrziu, E.; Dégi, J.; Imre, K. Field-
Based Characterization of Peste Des Petits Ruminants in Sheep in Romania: Clinical,
Pathological, and Diagnostic Perspectives. *Vet. Sci.* **2025**, *12* (7), 679.
https://doi.org/10.3390/vetsci12070679.

(46) Kirtzalidou, A.; Katsifa, A.; Komitas, G.; Xexaki, A.; Konstantinidis, A.; Vourvidis, D.;
Gianniou, M.; Kwiatek, O.; Guendouz, S.; Bataille, A.; Alexandropoulos, T.; Tasioudi, K.
E. Identification of Peste Des Petits Ruminants Virus, Greece, 2024. Social Science
Research Network: Rochester, NY June 6, 2025. https://doi.org/10.2139/ssrn.5279959.

(47) Moţ, D.; Tîrziu, E.; Olariu-Jurca, L. The Evolution of Peste Des Petits Ruminants (PPR)
in Romania and Neighbouring Countries. *Sci. Pap. Anim. Sci. Biotechnol.* **2025**, *58* (2),
301–308.

(48) Fine, A. E.; Pruvot, M.; Benfield, C. T. O.; Caron, A.; Cattoli, G.; Chardonnet, P.; Dioli,
M.; Dulu, T.; Gilbert, M.; Kock, R.; Lubroth, J.; Mariner, J. C.; Ostrowski, S.; Parida, S.;
Fereidouni, S.; Shiilegdamba, E.; Sleeman, J. M.; Schulz, C.; Soula, J.-J.; Van der
Stede, Y.; Tekola, B. G.; Walzer, C.; Zuther, S.; Njeumi, F.; Meeting Participants.
Eradication of Peste Des Petits Ruminants Virus and the Wildlife-Livestock Interface.
Front. Vet. Sci. **2020**, *7*. https://doi.org/10.3389/fvets.2020.00050.

(49) Pruvot, M.; Fine, A. E.; Hollinger, C.; Strindberg, S.; Damdinjav, B.; Buuveibaatar, B.;
Chimeddorj, B.; Bayandonoi, G.; Khishgee, B.; Sandag, B.; Narmandakh, J.;
Jargalsaikhan, T.; Bataa, B.; McAloose, D.; Shatar, M.; Basan, G.; Mahapatra, M.;
Selvaraj, M.; Parida, S.; Njeumi, F.; Kock, R.; Shiilegdamba, E. Outbreak of Peste Des
Petits Ruminants among Critically Endangered Mongolian Saiga and Other Wild
Ungulates, Mongolia, 2016–2017. *Emerg. Infect. Dis.* **2020**, *26* (1), 51–62.
https://doi.org/10.3201/eid2601.181998.

(50) Benfield, C. T. O.; Hill, S.; Shatar, M.; Shiilegdamba, E.; Damdinjav, B.; Fine, A.; Willett,
B.; Kock, R.; Bataille, A. Molecular Epidemiology of Peste Des Petits Ruminants Virus
Emergence in Critically Endangered Mongolian Saiga Antelope and Other Wild
Ungulates. *Virus Evol.* **2021**, *7* (2), veab062. https://doi.org/10.1093/ve/veab062.

(51) Saeed, I. K.; Haj, M. A.; Alhassan, S. M.; Mutwakil, S. M.; Mohammed, B. A.; Taha, K.
M.; Libeau, G.; Diallo, A.; Ali, Y. H.; Khalafalla, A. I. A Study on Transmission of Peste

Des Petits Ruminants Virus between Dromedary Camels and Small Ruminants. *J. Infect. Dev. Ctries.* **2022**, *16* (2), 374–382. https://doi.org/10.3855/jidc.14947.

(52) Jones, B. A.; Mahapatra, M.; Mdetele, D.; Keyyu, J.; Gakuya, F.; Eblate, E.; Lekolool, I.; Limo, C.; Ndiwa, J. N.; Hongo, P.; Wanda, J. S.; Shilinde, L.; Mdaki, M.; Benfield, C.; Parekh, K.; Mayora Neto, M.; Ndeereh, D.; Misinzo, G.; Makange, M. R.; Caron, A.; Bataille, A.; Libeau, G.; Guendouz, S.; Swai, E. S.; Nyasebwa, O.; Koyie, S. L.; Oyas, H.; Parida, S.; Kock, R. Peste Des Petits Ruminants Virus Infection at the Wildlife–Livestock Interface in the Greater Serengeti Ecosystem, 2015–2019. *Viruses* **2021**, *13* (5), 838. https://doi.org/10.3390/v13050838.

(53) *Comparative evolutionary analyses of peste des petits ruminants virus genetic lineages | Virus Evolution | Oxford Academic.* https://academic.oup.com/ve/article/10/1/veae012/7627268 (accessed 2026-02-14).

(54) Kaur, I.; Zulovich, J. M.; Gonzalez, M.; McGee, K. M.; Ponweera, N.; Thandi, D.; Alvarez, E. F.; Annandale, K.; Flagge, F.; Rezvani, K.; Shpall, E. Comparison of Two Methodologies for the Enrichment of Mononuclear Cells from Thawed Cord Blood Products: The Automated Sepax System versus the Manual Ficoll Method. *Cytotherapy* **2017**, *19* (3), 433–439. https://doi.org/10.1016/j.jcyt.2016.11.010.

(55) Rios, F. J.; Touyz, R. M.; Montezano, A. C. Isolation and Differentiation of Human Macrophages. In *Hypertension: Methods and Protocols*; Touyz, R. M., Schiffrin, E. L., Eds.; Springer: New York, NY, 2017; pp 311–320. https://doi.org/10.1007/978-1-4939-6625-7_24.

(56) Rigamonti, A.; Villar, J.; Segura, E. Monocyte Differentiation within Tissues: A Renewed Outlook. *Trends Immunol.* **2023**, *44* (12), 999–1013. https://doi.org/10.1016/j.it.2023.10.005.

(57) *Phenotype and Function of CD209+ Bovine Blood Dendritic Cells, Monocyte-Derived-Dendritic Cells and Monocyte-Derived Macrophages | PLOS One.* https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0165247 (accessed 2026-02-14).

(58) *Innate Immune Response to Rift Valley Fever Virus in Goats | PLOS Neglected Tropical Diseases.* https://journals.plos.org/plosntds/article?id=10.1371/journal.pntd.0001623 (accessed 2026-02-15).

(59) Libeau, G.; Lefevre, P. C. Comparison of Rinderpest and Peste Des Petits Ruminants Viruses Using Anti-Nucleoprotein Monoclonal Antibodies. *Vet. Microbiol.* **1990**, *25* (1), 1–16. https://doi.org/10.1016/0378-1135(90)90088-D.

(60) HAMILTON, M.; RC, R.; RV, T. TRIMMED SPEARMAN-KARBER METHOD FOR ESTIMATING MEDIAN LETHAL CONCENTRATIONS IN TOXICITY BIOASSAYS. *TRIMMED SPEARMAN-KARBER METHOD Estim. MEDIAN LETHAL Conc. Toxic. BIOASSAYS* **1977**.

(61) Wulff, N. H.; Tzatzaris, M.; Young, P. J. Monte Carlo Simulation of the Spearman-Kaerber TCID50. *J. Clin. Bioinforma.* **2012**, *2* (1), 5. https://doi.org/10.1186/2043-9113-2-5.

(62) Vieyres, G.; Pietschmann, T. Entry and Replication of Recombinant Hepatitis C Viruses in Cell Culture. *Methods* **2013**, *59* (2), 233–248. https://doi.org/10.1016/j.ymeth.2012.09.005.

743   (63) Wilham, J. M.; Orrú, C. D.; Bessen, R. A.; Atarashi, R.; Sano, K.; Race, B.; Meade-
744   White, K. D.; Taubner, L. M.; Timmes, A.; Caughey, B. Rapid End-Point Quantitation of
745   Prion Seeding Activity with Sensitivity Comparable to Bioassays. *PLOS Pathog.* **2010**, *6*
746   (12), e1001217. https://doi.org/10.1371/journal.ppat.1001217.

747   (64) Eloiflin, R.; Boyer, M.; Kwiatek, O.; Guendouz, S.; Loire, E.; Servan de Almeida, R.;
748   Libeau, G.; Bataille, A. Evolution of Attenuation and Risk of Reversal in Peste Des
749   Petits Ruminants Vaccine Strain Nigeria 75/1. *Viruses* **2019**, *11* (8), 724.
750   https://doi.org/10.3390/v11080724.

Strain-dependent differences in the capacity of peste des petits ruminants

virus to infect antigen presenting cells

Aadel Bouziane[a,b]#, Vincent Lasserre[a,b]*, Roger-Junior Eloiflin[a,b], Manon Chambon[a,b], Camille Piro-

Mégy[a,b], Samia Guendouz[a,b], Philippe Holzmuller[a,b]*, Philippe Totté[a,b], Arnaud Bataille[a,b]#

[a] CIRAD, UMR ASTRE, F-34398 Montpellier, France

[b] ASTRE, University of Montpellier, CIRAD, INRAE, Montpellier, France

Running Head: APC Infection Differences Among two PPRV Strains

#Address correspondence to Aadel Bouziane (ORCID: https://orcid.org/0009-0007-1090-9110),

aadel.bouziane@cirad.fr; address correspondence to Arnaud Bataille (ORCID: https://orcid.org/0000-

0002-3508-2144), arnaud.bataille@cirad.fr

*Present address: V. Lasserre, Laboratoire Ecologie et Biologie des Interactions, équipe MHE, UMR

CNRS 7267, Université de Poitiers, 5 rue Albert Turpin, 86073 Poitiers Cedex 9, France, Université

de Poitiers, Poitiers, France; R.-J. Eloiflin, Institut de Génétique Moléculaire de Montpellier (IGMM),

Université de Montpellier, Montpellier, France; P. Holzmuller, Cirad, UMR Selmet, Montpellier,

France

The order of authors reflects the relative contributions of the individuals listed up to S. Guendouz. The

last three authors, P. Holzmuller, P. Totté, and A. Bataille, contributed to the study in supervisory

roles.

KEYWORDS: Morbillivirus, monocyte-derived macrophage, monocyte-derived dendritic cell,

virulence, host susceptibility

Figure 1: Percentage of MoM (A) and MoDC (B) from cow, sheep and goat, positive for PPRV at 48hpi after in vitro infection with IC89 or MA08 strain. The MOCK group corresponds to uninfected control cells. Data are reported as individual biological replicates from minimum 3 animals per species and per condition with mean ± SD; ns=non-significant; *P<0.05; Statistical analysis was performed using a generalized linear mixed-effects model (GLMM). Post-hoc comparisons were conducted using estimated marginal means with Tukey adjustment for multiple comparisons.

45
46 **Figure 2: Percentage of MoM and MoDC from sheep and goat, positive for PPRV at 48hpi after in vitro infection with IC89 (A) or**
47 **MA08 (B) strain.** The MOCK group corresponds to uninfected control cells. Data are reported as individual biological replicates from
48 minimum 3 animals per species and per condition with mean ± SD; ***P<0.001; ****P<0.0001; Statistical analysis was performed using a
49 generalized linear mixed-effects model (GLMM). Post-hoc comparisons were conducted using estimated marginal means with Tukey
50 adjustment for multiple comparisons.
51

Figure 3: Viral titers (TCID$_{50}$/mL) of IC89 (A) and MA08 (B) at 48 hpi in cow, sheep and goat MoMs and MoDCs. Nine of seventeen replicates were below the limit of detection in goat MoMs infected with IC89 (circled in A). Data are shown as individual biological replicates from a minimum of three animals per species and per condition with mean ± SD; ns=non-significant; ****P<0.0001. Replicates from goat with no titer observed are highlighted with a circle. Statistical analysis was performed on viral titers using a generalized linear mixed-effects model (GLMM) fitted with a negative binomial distribution. Post-hoc comparisons were performed using estimated marginal means with Tukey adjustment for multiple comparisons

(A)

(B)   76

81   (C)

Figure 4: (A) Kinetics of infection of goat MoMs by strain IC89 of PPRV. Percentages of NPPRV+ cells were determined by flow cytometry at 24, 48, 72, and 96hpi. Data are shown as individual biological replicates with 3 goats and with minimum n=7 replicates, mean ± SD. A beta-binomial GLMM revealed no significant effect of time post-infection on infection rates (all P > 0.76). Post-hoc pairwise comparisons using Tukey-adjusted estimated marginal means confirmed the absence of significant differences between 24, 48, 72 and 96 hpi (all adjusted P > 0.99). **(B) Viral titration kinetics of MoMs supernatants from goats infected with strain IC89 at 24, 48, 72 and 96hpi.** Data are shown as individual biological replicates from 3 goats with n=9 replicates per time point. **(C) Kinetics of viral shedding in MoMs supernatant infected by IC89 strain at 24h, 48h, 72h and 96h.Viral shedding was monitored by RT-qPCR (expressed in Cq) detecting the presence of the PPRV N gene.** Data are shown as individual biological replicates from 3 goats with n ranging from 12 to 20 replicates per time point, mean ± SD; *P<0.05; P<0.0001. Statistical analysis was performed using a linear mixed-effects model (LMM) fitted by restricted maximum likelihood (REML). Pairwise comparisons were conducted using estimated marginal means with Tukey adjustment for multiple comparisons (Kenward–Roger approximation for degrees of freedom).

103
104
105

Figure 5: Immunofluorescent detection of viral double-stranded RNA (dsRNA) (A) and Nucleoprotein PPRV protein (NPPRV) (B) in goat MoMs and MoDCs infected with IC89. Infected and uninfected (MOCK) MoMs and MoDCs were stained with DAPI to identify cell nuclei, dsRNA (A) and NPPRV (B) are detected here in red, scale bar = 25 µm.