Apparent failure and cryptic success of disease control via intermediate host population
reduction

Authors: **Nadia Raytselis**[1], Maya Risin[1], Layla Steinbock[1], Logan T. Story[1], Jane Miller[1],
Ta'Nyia Heard[1], Kevin Xie[1], Ella Rintala[1], Inesha Gupta[1], Stephanie Gutierrez[1],
Alexander Strauss[2,3,4], David J. Civitello[1]

1. Department of Biology, Emory University, Atlanta, GA, 30322, USA
2. Odum School of Ecology, University of Georgia, Athens, GA, 30602, USA
3. Center for the Ecology of Infectious Diseases, University of Georgia, Athens, GA, 30602
4. River Basin Center, University of Georgia, Athens, GA, 30602

Keywords: population dynamics, negative density dependence, pest management, compensation,
overcompensation, disease control

Abstract

All populations experience density dependence, lest they grow infinitely. However,

elucidating what forms of density dependence act most strongly for a given species remains

incredibly challenging. Identifying the mechanisms that regulate population density is

particularly relevant to pest control, which often causes high mortality, reducing population

density but enabling recovery when survivors are freed from intraspecific competition and

density dependent vital rates (e.g., birth, death, and maturation rates). Understanding these

demographic responses is critical for effective pest management, especially when specific life

stages of pests differ in the harm they cause. Guinea worm disease (GWD) is a neglected tropical

disease with an obligate copepod intermediate host, and importantly, only large-bodied copepods

can be infected with GW parasites. GW disease has been the target of control for several decades,

with main management strategies including the use of a chemical larvicide, Abate™, which is

applied to water bodies to cull copepod populations. Despite the wide application of Abate in

GW endemic countries, little is known about long-term copepod population dynamics in

response to Abate™. Thus, we evaluated this mortality-based management of intermediate

copepod hosts to control GW transmission, combining mathematical models with a population

dynamics experiment designed to mimic Abate™ pesticide control methods across a range of

intensities. Despite initial reductions in both total and stage specific population densities,

copepod populations recovered so rapidly that control appeared to fail, enabled by extremely fast

maturation rates in low-density populations. Much to our surprise, model simulations of GW

transmission showed that although total and stage-specific densities rebounded, infectious adult

copepods —the proximate cause of infections—were strongly suppressed, indicating cryptic

success. This effect was enabled by the 15-day developmental period of GW larvae within

copepods, which blocks the accumulation of infectious copepods between interventions.

Ultimately, this study highlights the importance of understanding mechanisms of density

dependence when designing and optimizing pest control interventions, as well as interpreting

counterintuitive consequences of interventions.

Introduction

Elucidating the mechanisms that regulate populations is a fundamental goal of population

ecology. Negative feedbacks on population growth emerge because of competition for space

(Sebens 1982) and resources (Svanbäck and Bolnick 2007), degradation of the environment

(Calizza et al. 2017), or other mechanisms that reduce individual performance as density rises

(Lu and Wang 2025, Stockseth et al. 2025). Reductions in birth rates (Gotelli 2008) and

increases in death rates are ubiquitous inhibitors of population growth (Hixon and Jones 2005).

In populations with stage structure, negative density dependence can also arise through

maturation rates or cannibalism (Park et al. 1965, Dennis et al. 2001, Walker et al. 2021).

Despite the reality that all populations experience negative density dependence (lest they

increase infinitely), it remains challenging to identify which form(s) operate most strongly for

any species and few studies attempt to do so (Kaspersson et al. 2013).

The forms of density dependence are particularly relevant for pest control, because

interventions are designed to reduce pest density. This reduction is often followed by rapid

recovery, driven by relaxed density dependence (Karatayev et al. 2015) or potential acquisition

of resistance (Cao et al. 2025), limiting benefits and necessitating repeated interventions

(McIntire and Juliano 2018). The ensuing dynamics of repeated pest suppression and recovery

can be highly sensitive to density-dependent vital rates (Smith 1997). For example, strong

negative density dependence can cause compensation or overcompensation, the phenomena in

which control efforts actually increase the density or biomass of some or all life stages

(Karatayev et al. 2015, de Roos 2020, Grosholz et al. 2021). Even when interventions

successfully reduce density, populations can rebound faster than expected if the remaining

individuals are released from density-dependent reductions in maturation or birth rates

(Karatayev et al. 2015). Such responses may be strongest in stage-structured populations, where

removal of more competitive classes, or reductions in density overall, elicit stage-specific

compensatory responses (Karatayev et al. 2015). Thus, delineating the form(s) and magnitude of

density dependence can improve management of the ecological issues that pests cause, such as

disease transmission, crop losses, and biodiversity harm.

Here we combine an experiment manipulating pest control intensity with mechanistic

modeling to identify which density-regulating mechanisms affect copepods, the intermediate

hosts of the Guinea worm (GW) parasite, *Dracunculus medinensis*. Copepods exhibit stage-

structured populations (van den Bosch and Gabriel 1994) and experience repeated control

interventions due to their role in GW transmission (Gindola et al. 2022). Guinea worm disease

(GWD) is a globally impactful infectious disease, causing disability and negative economic

impacts in endemic areas (Pellegrino et al. 2022). Cyclopoid copepods are obligate intermediate

hosts that become infected through consumption of GW larvae and facilitate transmission to

humans and other definitive mammalian hosts (Simonetti et al. 2023). GWD has been targeted

for eradication for several decades (World Health Organization 1986), with few cases remaining

across five endemic countries (World Health Organization 2025). In GW-endemic regions,

application of the pesticide Abate™ (active compound: temephos) is a major component of

eradication efforts (Gindola et al. 2022). Abate™ can cause >99% immobilization of copepods

in laboratory studies (Grunert et al. 2022), although copepod species vary widely in their

sensitivity (Hanazato et al. 1989, Leboulanger et al. 2011). In Abate[TM]-treated sites, copepod

populations often exhibit immediate reductions in density, but routinely recover, necessitating

application at 28-day intervals to support eradication (Gindola et al. 2022). The density-

dependent demographic responses of these intermediate hosts – along with implications for

disease control – are unknown.

Copepod stage structure has implications for both their population dynamics and GW

transmission. Neonatal copepods (nauplii) are too small to consume GW larvae, but larger

juveniles (copepodites) and adults can consume parasites and become infected (McCullough

1983, Bimi 2007). Once ingested, parasites molt twice over 10-16 days within the copepod

before maturing into third stage larvae, the only stage infectious to definitive hosts that imbibe

infected copepods (Gonzalez Engelhard et al. 2021). Additionally, adults can cannibalize nauplii

(van den Bosch and Santer 1993). Finally, public health officials have shared anecdotally that

adults and copepodites may be more sensitive to Abate[TM] than nauplii. Critically, the

consequences of these stage-structured interactions for disease management are unknown.

To understand copepod demographic responses to episodic culling, and to predict

implications for disease control, we conducted a mesocosm experiment that mimicked the 28-

day treatment interval of Abate[TM] by physically removing copepods (either applied equally

across stages or biased toward larger individuals). We hypothesized that more intense

interventions (higher removal percentages) would cause greater reductions in population density,

but that relaxed density dependence might cause compensation or overcompensation between

interventions. Further, we speculated that these effects would be more pronounced in populations

where both size classes were removed. Finally, we hypothesized that even if copepods

rebounded rapidly ('apparent failure'), risk of parasite transmission might remain low ('cryptic

success') due to high adult turnover rate and the two-week period required for parasite

development from first stage to infectious third stage larvae.

Surprisingly, all interventions failed to reduce mean copepod density over the 14-week

intervention period. Copepods recovered extremely rapidly due to relaxed density dependence on

maturation rates. However, despite this apparent failure, model simulations indicate that

interventions may still reduce transmission risk by specifically suppressing the density of

infectious copepods. These results illustrate how density-dependent mechanisms interact with

transmission biology to drive cryptic success of control. Stage structure is a key feature of the

biology of many pest species, e.g., root-feeding crop pests (Johnson and Rasmann 2015) and

mosquito vectors of human diseases (Neale and Juliano 2019, Walker et al. 2021), and control

interventions often focus on specific life stages that are not always the proximate cause of harm,

e.g., larvicides to control pathogens transmitted by adult mosquitoes (Choi et al. 2019). Thus,

understanding the underlying mechanisms of stage-structured density dependence could enhance

our ability to address a diverse range of ecological challenges.

Methods

Copepod Maintenance and Propagation

We obtained *Eucyclops sp.* copepods isolated near Mandelia, Chad, an active focus of GW

transmission. We maintained copepods in 0.5x HHCOMBO artificial lake with 1X

micronutrients (hereafter COMBO; Kilham et al. 1998) in 1 L beakers at 26C with a 12:12

daylight cycle. We fed them 7.5 mg/L of powdered egg yolk (Judee's, Plain City, Ohio)

suspension every Monday, Wednesday, and Friday (M/W/F), and transferred copepods to new

media approximately every three weeks by sieving. Once densities reached ~500-1,000

copepods/L, we poured one 1 L culture into 42, 15 L propagation tanks with 4 L COMBO.

Mesocosm Setup, Sampling, and Experimental Design

On September 20[th], 2024, we filled 60 15 L tanks with 5 L COMBO and added 1.5 mg/L

powdered egg yolk on Monday/Wednesday/Friday to establish microbial communities. On

October 2[nd], 2024 (day 0), we introduced copepods to tanks by adding seven 1 L aliquots

collected randomly from the 42 propagation tanks after stirring to homogenize. We then added

COMBO to 15 L total and fed tanks 7.5 mg/L every Monday/Wednesday/Friday throughout the

experiment. We replaced water weekly to counteract evaporation (<1 L/tank/week).

We maintained the experiment for 126 days and sampled every Friday starting on day 2.

To sample, we homogenized tanks (stirring with a plastic paddle) and collected a 500 mL surface

grab sample with a beaker. We replaced this volume with COMBO. We intervened at 28-day

intervals (days 8, 36, and 64) to mimic the field use of Abate™ (Gindola et al. 2022). We

implemented mechanical removal, rather than Abate™, to enable consistent gradients of size-

biased and size-unbiased removal. We factorially crossed intervention type (equal vs. size-

biased) with the percent of the volume (i.e., copepod population) removed (0-90%: 0, 10, 20, 30,

40, 50, 60, 70, 80, or 90%). For size-biased removal, we sieved water through 100 μm nylon

screen, capturing adults, juveniles, and larger nauplii, and returned the water and smaller nauplii

to tanks. For size-unbiased (i.e., equal) removal, we replaced the volume removed with fresh

COMBO. In Appendix S7, we provide methodological details and results on the accuracy of

interventions relative to nominal treatment levels. After the first intervention, we converted the

10% and 20% treatments to 95% and 98% for the following interventions, because low initial

intensities caused negligible effects, and higher removal rates better match the stated efficacy of

the Abate™ (Grunert et al. 2022). On weeks with interventions, we sampled on Thursday and

intervened on Friday. We "skipped" the intervention scheduled for day 92 to observe responses

to treatment lapse, as can occur in the field due to inaccessibility from flooding.

Copepod Counting

We quantified copepods by sieving the 500 mL samples with 35 μm mesh (which captures all

stages) and preserving them in 10 mL ethanol. We gently resuspended the samples and examined

successive 1 mL aliquots at 10-25x magnification on a dissecting microscope until counting ≥ 50

copepods or screening the entire sample. We counted copepods in three morphologically

discernible categories (nauplii, juveniles and eggless adults, and adult females with eggs).

Statistical Analysis of Intervention Effects on Copepod Density

We conducted all analyses in the R Computing Language v4.3.2 (R Core Team 2023). We tested

for differences in mean population density over the first 14 weeks, when treatments were applied

"on schedule", using planned contrasts between all interventions versus controls (Dunnett's tests;

package emmeans (Lenth 2025). This analysis used a Generalized Linear Model (package

glmmTMB; Bolker 2017) with mean density as the response, removal proportion, removal type,

and their interaction as predictors, and the Gamma error distribution with a log link function.

State Space Modeling (SSM) of Copepod Population Dynamics
184
We built a stochastic population dynamic model and simultaneously fit it to all time series of

observations from the experiment using the Partially Observed Markov Process (POMP)

framework and the panelPomp package (Carles Bretó 2025). We connected likelihood estimates

generated by the function pfilter() to an adaptive Markov Chain Monte Carlo (MCMC) sampler

in the adaptMCMC package to conduct Bayesian inference, with priors informed by auxiliary

experiments (Appendix S2: Figure S2). POMP models are ideal for population dynamics data

(Newman et al. 2023) where state variables (the "true" abundances of nauplii, juveniles, and

adults) are partially observed as empirical subsample counts of nauplii (N_{obs}), juveniles/adult

males/adult females with no eggs (JOA_{obs} = "juveniles and other adults"), and adult females

(AF_{obs}) (Carles Bretó 2025).

We implemented the model by specifying ordinary differential equations that describe

deterministic rates of change in the densities of nauplii, N, juveniles, J, and adults, A. In this

model (Equations 1-6), nauplii are born, develop into juveniles, die from background causes, and

are cannibalized by adults. Juveniles mature from nauplii, develop into adults, and die of

background causes. Adults mature from juveniles, birth and cannibalize nauplii, and die of

background causes. Critically, birth rate (b), stage-specific death rates (d_i, with $i = N, J, or A$),

and stage-specific maturation rates (m_i, with $i = N or J$) depend exponentially on density, with

each stage potentially contributing differentially to density dependence via their impact on the

competitive environment (i.e., c_J and c_N represent the competitive effects of juveniles and nauplii

relative to adults). We allowed adults to cannibalize nauplii following a linear functional

response with attack rate α. On intervention days, we incorporated removal as additional

mortality on relevant stage(s). During all interventions, we assumed juveniles and adults were

removed according to treatment. We assumed nauplii were removed proportionally in size-

unbiased removal treatments and used a prior informed by an auxiliary experiment that showed

size-biased sieving removed approximately 45% of nauplii (Appendix S2: Methods S1).

$\frac{dN}{dt} = b(N,J,A)\frac{A}{2} - m_N(N,J,A)N - d_N(N,J,A)N - \alpha NA$ Eq. 1

$\frac{dJ}{dt} = m_N(N,J,A)N - m_J(N,J,A)J - d_J(N,J,A)J$ Eq. 2

$\frac{dA}{dt} = m_J(N,J,A)J - d_A(N,J,A)A$ Eq. 3

with density-dependent functions:

214
$$b(N, J, A) \; = \; b_M e^{\left(-\frac{c_b}{V}(c_N N + c_J J + A)\right)}$$
Eq. 4

215
$$m_i(N, J, A) = m_{i0} e^{\left(\frac{-c_m}{V}(C_N N + C_J J + A)\right)}$$
Eq. 5

216
$$d_i(N, J, A) = d_{i0} e^{\left(\frac{c_d}{V}(C_N N + C_J J + A)\right)}$$
Eq. 6

To simulate the dynamics of the latent (or "true and unobserved") state of the system we

assumed demographic stochasticity and implemented the model with a one-day time step.

Specifically, we tracked individuals transitioning through stages or dying using random

multinomial draws where the probability of three potential outcomes for any individual

(remaining in the same state, maturing to the next state, or dying) depends on the relative rates of

each process. We assumed births followed Poisson random draws depending on the total birth

rate. Our observations differentiate N_{obs}, AF_{obs}, and JOA_{obs}. We assumed that each observation

followed a negative binomial distribution with a common overdispersion parameter. For each

observation, we assumed the predicted count was determined by the product of the abundance of

individuals of that stage at time t, the proportion of the tank sampled (always $0.5/15$ L $= 3.3\%$),

and the proportion of the 10 mL subsample counted. To align true and observed categories, we

assumed a 50:50 sex ratio with female reproduction, with adult females releasing eggs every

third day and appearing eggless for one day in each three-day interval (personal observation

TDH). With this model specified, we used the pfilter() function in panelPomp to estimate model

likelihoods from the data. To maintain variance in log-likelihoods < 1 (Endo et al. 2019), we

used 20 particles per likelihood estimation, resulting in variances of approximately 0.5. We ran

32 MCMC chains, with dispersed initial parameter sets for 250,000 iterations each, permitting

adaptation during the first 50,000 iterations (burn-in), which we later discarded.

Model-derived estimates of density dependence

We used model-derived functions of density dependence, observed densities, and stage structure

from the experiment to visualize how the competitive environment shaped demographic rates.

We thinned chains to every 1,000[th] iteration, and calculated per-capita birth, death, cannibalism,

and maturation rates across a density gradient from 0 to 6,000 individuals/L for each parameter

set (the density range observed in the experiment). We fixed stage structure along this density

gradient (5.5% adults, 39% juveniles, and 55.5% nauplii, 1:7:10 ratio), based on the average

proportions in the experiment. We then generated the mean and 95% CI interval of posterior

predictions for each demographic rate along the gradient.

Compartmental (ODE) stage structured model with interventions and infection

Large-scale experiments with viable GW larvae are currently infeasible. Therefore, we extended

our model to gain applied insights into disease control. The extended model (Eq. S1-S11 in

Appendix S1: Methods and Equation S1) allows parasites to infect adult copepods. We

implemented a distributed developmental delay (Hurtado and Kirosingh 2019) for copepods

transitioning from exposed to infectious to definitive hosts (Appendix S1: Methods and Equation

S1). This approach uses the "linear chain trick" (Hurtado and Kirosingh 2019) to implement a

variable time lag, as we hypothesized this delay could be important for differentiating successful

population regulation from successful disease control. Once an adult copepod consumes the

parasite, it takes approximately 15 days to become infectious (Lindquist and Cross 2017). Thus,

we added a developmental delay of approximately 15 days by implementing a chain of 60

exposed stages that copepods transition through at a rate of 4.3 d^{-1}, yielding a 15.45 ± 1.82-day

delay from exposure to infectiousness.

Estimates of the force of infection for copepods are unavailable. Therefore, we assumed a

5% infection prevalence among adults, consistent with the limited field data available (Lyons

1972, Steib and Mayer 1988, Garrett et al. 2020), corresponding with $\lambda = 0.0034$/day. Further,

while parasites are likely introduced in discrete pulses (a single mature female can introduce 1

million L1 larvae (Greenwood 2012)), we implemented a constant force of infection to avoid

confounding results with day of introduction. Lastly, we assumed infected adults cannot

reproduce (Moorthy 1938).

Using this model, we evaluated the consequences of episodic mortality events on stage-

specific and total abundance, with emphasis on the density of infectious copepods, the proximate

hazard to definitive hosts. We plotted the temporal dynamics for the 90% mortality scenario, a

previously used metric for Abate^{TM} success (South Sudan Guinea Worm Eradication Program

2013). We plotted mean densities of all stages across the whole simulation period for scenarios

ranging from 0-90% mortality.

Results

Copepod population dynamics and mortality events

Regardless of removal intensity, we found no significant decreases in mean population density

over the first 14-weeks of the study for size-unbiased (light blue) or size-biased (dark blue)

removal (Fig. 1). This apparent failure was consistent for both the total population and when

broken down by stage (Fig. 1, diamond, triangle, circle; all Dunnett's test $P > 0.05$).

While we found no effect on mean density, the interventions did reduce densities

immediately after application (e.g., 95 and 98% removal, Fig. 2, Appendix S5: Figure S1A-B).

However, populations then rapidly rebounded. Populations in controls started at ~500-1,000

copepods/L, grew slowly for several weeks, and ended with a boom and bust (Fig. 2). In other

cases, density in culled tanks temporarily surpassed controls, indicating overcompensation. For

example, following the third intervention, densities in the 50% size-biased treatment were briefly

higher than controls. In some cases, population recovery occurred so rapidly that declines were

imperceptible one week later (e.g., after the second intervention with 80% removal). The best fit

of the model generally captured the dynamics in intervention tanks better than that of the control

treatment, which is partially expected, as the control treatment represented only 10% of the

experiment.

Size-biased Removal:

Relative to controls, the 30% and 60% size-biased treatments grew steadily throughout

the entire experiment (Fig. 2, Appendix S5: Figure S1A). Populations quickly recovered from

interventions, and observed dynamics generally fell within 95% credible intervals of the fitted

model predictions. The 40%, 50%, and 70% removal treatments more closely resembled the

controls, with large fluctuations in density at the end of the experiment. The 80%, 90%, and 95%

removal treatments had multiple extreme booms and busts. Finally, 98% removal treatments

fluctuated with low levels of population growth. These shifts sometimes fell outside the credible

interval of posterior predictions.

Size-unbiased Removal:

In contrast with controls, 30%, 70% and 80% size-unbiased removal treatments

maintained relatively constant abundances across all eighteen weeks, mostly within the credible

interval (Fig. 2, Appendix S5: Figure S1A). The 40% and 60% tanks resembled the controls. The

50% treatment also resembled the control; however, instead of final booms and busts,

populations declined steadily. The 90%, 95%, and 98% treatments had sustained peaks and

troughs, encompassed by the credible interval for most of the experiment. The last few weeks

had more extreme booms and busts, not captured within the credible interval.

Evaluating the Functional Forms of Density Dependence

The strength of negative density dependence varied significantly across processes of birth,

maturation, background death, and cannibalism. Density dependence was greatest for maturation

rates, with a >18,000-fold reduction across the observed density gradient (Fig. 3). Density

dependence on background death rate and cannibalism were also substantial, with a combined

4.4-fold increase over the density range (Fig. 3). However, we estimated essentially no effect on

birth rates, with <2% decrease over the density range.

Transmission Scenario Simulation Results

Simulated GW infection scenarios exhibited rapid, stage-specific recovery of copepods

following interventions (Fig. 4). With 90% intervention intensity, nauplii and non-infectious

adults rapidly rebounded to densities that surpassed those observed for untreated populations

(Fig. 4C, D), and juveniles reached similar, though lower densities until the next intervention

(Fig. 4B). Size-unbiased removal generally reduced copepod populations more strongly, but

these populations also showed slightly greater overcompensation (Fig. 4A, C). This transient

overcompensation reiterated a central result from the experiment (Fig. 1): that mean copepod

density appeared unaffected by interventions.

Despite this apparent failure, simulated interventions successfully reduced the density of

infectious copepods (Fig. 4A). Without intervention, control populations began with low

densities of infectious hosts, but these grew steadily (>75 infected copepods L^{-1} by the end of the

100-day simulation). However, both size-unbiased and size-biased suppressed infectious

copepod density below 10 L^{-1} (Fig. 4A). Evaluating mean densities of stages across an

intervention intensity gradient (Fig. 5), total population density and non-infected adults

(susceptible + exposed classes) remained relatively constant at all intensities, while the density of

infectious adults steadily declined across the gradient. Thus, despite observing that copepod

populations rebound rapidly – enabled by intense density dependence on maturation rates – the

density of infectious hosts, the proximate source of mammalian infections, is highly sensitive to

intervention.

Discussion

336   Uncovering the mechanisms regulating populations is a fundamental goal in population

ecology (Kaspersson et al. 2013, Karatayev et al. 2015). We combined a pest control experiment

and mechanistic modeling to partition the magnitude and importance of four general putative

mechanisms of density regulation. Then, we integrated these empirically-derived regulating

effects with a transmission model to evaluate how relaxation of density dependence via realistic

control methods shapes effectiveness of interventions for GW, a target for global parasite

eradication (Pellegrino et al. 2022).

343   Our experiment investigated how copepod populations – intermediate hosts of GWD –

responded to periodic culling. We designed treatments to mimic the treatment of water sources

with Abate™, an intervention commonly used to prevent GWD in endemic countries. We

hypothesized that more aggressive control efforts would better suppress copepod populations,

but that relaxed density dependence might lead to overcompensation. Strikingly, even when we

episodically removed up to 98% of the population, densities across were statistically

indistinguishable from controls (Fig. 1, Appendix S3: Table S1). Similar rebounds have been

observed in other copepod species, including *Mesocyclops leuckarti* which overshot pre-

treatment densities following application of Triphenyltin, a fungicide (Kulkarni et al. 2014). Our

interventions had failed to reduce copepod densities, an alarming prospect as other apparent pest

control failures have posed immense challenges for public health and conservation practitioners

(Rivero et al. 2010, Neale and Juliano 2019, Grosholz et al. 2021). For example, populations of

mosquitoes, sand flies, assassin bugs, and fleas, which are routinely treated to control diseases

such as malaria, leishmania, chagas, and plague, respectively, show evidence of insecticide

resistance, threatening vector control (Cao et al. 2025). Further, specific mosquito species

including *Aedes* and *Culex sp.*, vectors of diseases such as dengue, chikungunya, Zika virus

(*Aedes*) (Patterson et al. 2016), and West Nile Virus (*Culex*) (Hamer et al. 2008), show evidence

of culling-induced overcompensation, indicating that control strategies must be careful to not

impose mortality at levels that lead to overcompensatory releases from density dependence

(Neale and Juliano 2019). A striking example of a pest control program that backfired is the

effort to eliminate the invasive European green crab in an estuary off of California (Grosholz et

al. 2021). Culling of green crabs was an effective strategy for years until the population density

was reduced below a critical threshold, triggering stage-specific overcompensation in

recruitment (Grosholz et al. 2021). The result was a boom in juvenile crabs, a shift in size

structure, and exacerbation of negative impacts (Grosholz et al. 2021). Together, these examples

underscore the challenges of pest control – especially in light of stage structure.

To attempt to further explain the surprisingly fast recovery we observed, we investigated

the forms of density dependence regulating *Eucyclops sp.* populations. Few studies have parsed

specific drivers of density dependence in natural populations. In one example, stocking large

brown trout reduced the growth rate of smaller individuals (Kaspersson et al. 2013). In another,

dynamics of Swedish vole populations were consistent with density dependence in recruitment

alone (Lambin et al. 2025). Stage-specific density dependence in maturation rates has been

documented in some ecological systems (Karatayev et al. 2015, Walker et al. 2021), including

other copepods (Kulkarni et al. 2014), allowing for population persistence under significant

mortality (Karatayev et al. 2015). We determined that density dependence in *Eucyclops sp.* was

driven predominantly by maturation rates, with a >18,000-fold reduction along the density

gradient, with a modest ~4.4-fold increase in total mortality, and no change in birth rate (Fig. 3).

Thus, when interventions occur, surviving adults maintain birth rates, but nauplii mature

extremely quickly. Consequently, higher proportions of nauplii and juveniles survive to

development, enabling rapid recovery and temporary overcompensation. (Figs. 2 and 4).

For the goal of GWD management, the failure to reduce mean copepod population

densities seems alarming. However, not all copepod life stages host GW, (McCullough 1983,

Bimi 2007) and there is a significant delay between copepod exposure to GW and infectiousness

to mammals (~15 days; Gonzalez Engelhard et al. 2021). Thus, we speculated that if population

rebounds were driven by overcompensation in young stages that cannot transmit GW, then

apparent failures to control the overall population might actually conceal successes. We

evaluated this possibility by extending our model to include GW infection of adults with a

distributed delay from the exposed to infectious class. We tracked changes in the density and

stage structure of copepod populations across a gradient of intervention intensity.

When we simulated GW transmission scenarios, we found strong support for stage-

specific recovery dynamics, as documented in other zooplankton systems (Nilsson et al. 2010,

Kulkarni et al. 2014). Specifically, the densities of nauplii, juveniles, and non-infectious adults

all recovered rapidly to meet or exceed pre-intervention densities following each successive

mortality event (Fig. 4), indicating compensation or overcompensation (Abrams 2009, Kulkarni

et al. 2014, de Roos 2020, Grosholz et al. 2021). Critical to GW transmission, the density of

infectious adults remained extremely low. Thus, despite the apparent failure of these

interventions to control copepod populations, they appear highly effective at specifically

controlling the life stage that is the proximate cause of GW infections. We attribute this effect to

the approximately15-day developmental period from first stage to infectious third stage GW

larvae (Lindquist and Cross 2017, Gonzalez Engelhard et al. 2021), which is long enough to

prevent accumulation of infectious copepods before each successive intervention (Fig 4., Fig. 5).

Similar lags of parasite development in intermediate hosts and vectors are present in other

systems. For example, in Anopheline mosquitoes, a 2-3 week incubation period separates vector

exposure to *Plasmodium* and infectiousness to humans (Crutcher JM 1996). Similarly,

Trypanosoma maturation in tsetse flies requires approximately three weeks (Telleria et al. 2014).

Thus, reducing the overall density of intermediate hosts or vectors may not be essential for

success if control can cause high turnover in the stage(s) responsible for transmission.

Our results raise the importance of field surveillance targeted at infected copepods. At

present, diagnosing infections in copepods from field is infeasible due to GW rarity and the scale

of geographic surveillance (Pellegrino et al. 2022). Therefore, current eradication programs rely

on monitoring the total density of copepods before and after Abate[TM] applications. If copepod

communities in the field recover as rapidly as the *Eucyclops sp.* examined in our study, officials

might falsely conclude that control is ineffective, because population densities return to pre-

treatment levels every month. However, our experiment and model suggest that this observation

could conceal a cryptic success: reduction in the density of infectious copepods. Thus, we

recommend the development of tools to facilitate large-scale spatial and temporal diagnostics of

GW-infected copepods to evaluate the relevance of this counterintuitive finding.

We note some important caveats to translating these insights to the field. First, we

focused on a single copepod isolate. Greater understanding of copepod diversity in demographic

rates and sensitivity to Abate^{TM} would greatly benefit eradication programs. Second, we

implemented mechanical removal, which enabled a precise experimental design, but can differ

from chemical control. Experiments with Abate^{TM} are needed to evaluate chemical persistence,

delayed or sublethal effects, and the consequences of over- or underdosing. Lastly, our analysis

identified high importance for negative density dependence on maturation rates and mortality,

and no effect on birth rates. Generally, the model fit the data well for the first 70-80 days of the

experiment, albeit with some initial lags in population growth (Fig. 2). This imperfect fit could

be due to missing key processes in the model, such as an Allee effect, where mating failure

occurs at low population densities (seen for copepods (Kramer et al. 2008)). The inability of the

model to precisely recover empirical dynamics could also be due to nonconstant conditions in the

tanks (e.g., the accumulation of nutrients (Sahandi et al. 2023) and/or waste products or

differences in temperature or photoperiod over the course of the experiment). Nevertheless, the

POMP approach has major benefits, including representation of constraints in the observation

process and incorporation of process noise. Lastly, future studies of copepod control and

recovery could consider other functional forms for key demographic processes, including Type-

II or Type-III functional responses for cannibalism.

Chemical control of copepod populations represents a critical intervention for national

GW eradication programs (Gindola et al. 2022). This study highlights that nuance may be needed

to evaluate this intervention: population dynamics may suggest that control efforts are failing or

even backfiring, but direct analyses of infectious copepods, the proximate cause of human cases

and animal infections should be prioritized. Future laboratory studies should expand to greater

copepod diversity and directly evaluate pesticide formulations in use. Ultimately, a deeper

understanding of copepod control may improve the portfolio of GWD interventions and

contribute to the eradication of this devastating parasite.

More broadly, pest control is inherently designed to reduce population density (Ni et al.

2025), so understanding density dependence is critical to predicting responses (McIntire and

Juliano 2018). This work underscores the value of defining which life stages of pests are most

impactful, elucidating how stage-structured pest populations respond to interventions, and

developing a mechanistic understanding of density dependent mechanisms regulating

populations under control, especially in cases where focal populations experience drastic and

rapid shifts in density. Together, this information can be applied to anticipate and explain

potentially counterintuitive consequences of interventions. A thorough understanding of these

dynamics is critical for forecasting long-term population trajectories and their sensitivity to

density-dependent perturbations, paramount to predicting the influence of environmental and

ecological stressors on population dynamics.

Acknowledgements

This work was supported by funding from The Carter Center. We thank the Programme National

d'Eradication du Ver de Guinée - Tchad for collecting and providing the Chadian *Eucyclops*

sp. copepods used in our study, and the Yen lab at the Georgia Institute of Technology for

providing the culture to us. We thank The Carter Center and other Civitello lab members for

feedback on this manuscript. We also thank M. Hoogshagen and J. Raytselis for helping with

sampling over the holidays, and B. Lukubye for moral support.

Author Contributions

N.R., S.G., A.S., and D.J.C. developed and planned experiments. N.R., M.R., L.S., L.T.S, and J.M. conducted the experiment. N.R., M.R., L.S., L.T.S, J.M., T.H., E.R., and I.G., processed samples. N.R., K.X., and D.J.C. conducted analysis. A.S. and D.J.C. provided feedback to refine analysis. N.R. wrote manuscript and made figures with feedback from A.S. and D.J.C..

Conflict of Interest Statement

The authors report no potential conflicts of interest.

Literature Cited

Abrams, P. A. 2009. When does greater mortality increase population size? The long history and diverse mechanisms underlying the hydra effect. Ecology Letters **12**:462-474.

Bimi, L. 2007. Potential vector species of Guinea worm (Dracunculus medinensis) in Northern Ghana. Vector Borne Zoonotic Dis **7**:324-329.

Bolker, B. 2017. {glmmTMB} Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling.

Calizza, E., M. L. Costantini, G. Careddu, and L. Rossi. 2017. Effect of habitat degradation on competition, carrying capacity, and species assemblage stability. Ecology and Evolution **7**:5784-5796.

Cao, L. J., J. C. Chen, J. A. Thia, T. L. Schmidt, R. Ffrench-Constant, L. X. Zhang, Y. Yang, M. C. Yuan, J. Y. Zhang, X. Y. Zhang, Q. Yang, Y. J. Gong, H. Li, X. Chen, A. A. Hoffmann, and S. J. Wei. 2025. Recurrent mutations drive the rapid evolution of pesticide resistance in the two-spotted spider mite Tetranychus urticae. Elife **14**.

Carles Bretó, J. W., Aaron A. King, and Edward L. Ionides. 2025. panelPomp: Analysis of Panel Data via Partially Observed Markov Processes in R. The R Journal.

Choi, L., S. Majambere, and A. L. Wilson. 2019. Larviciding to prevent malaria transmission. Cochrane Database Syst Rev **8**:Cd012736.

Crutcher JM, H. S. 1996. Malaria, Medical Microbiology.

de Roos, A. 2020. The impact of population structure on population and community dynamics. Pages 53-73.

Dennis, B., R. A. Desharnais, J. M. Cushing, S. M. Henson, and R. F. Costantino. 2001. Estimating Chaos and Complex Dynamics in Insect Populations Ecological Monographs **71**:277-303.

Endo, A., E. van Leeuwen, and M. Baguelin. 2019. Introduction to particle Markov-chain Monte Carlo for disease dynamics modellers. Epidemics **29**:100363.

Garrett, K. B., E. K. Box, C. A. Cleveland, A. A. Majewska, and M. J. Yabsley. 2020. Dogs and the classic route of Guinea Worm transmission: an evaluation of copepod ingestion. Sci Rep **10**:1430.

Gindola, Y., D. Getahun, K. Mohammed, E. Kamau, B. Camara, M. Wossen, K. Demissie, S. Abdela, G. Gebrewolde, G. Hailu, M. Tegistu, A. Okugn, and G. Gikilo. 2022. Abate

application practices in the Guinea worm endemic region of Gambella, Ethiopia: identification of elimination gaps. The Journal of Infection in Developing Countries **16**:20S-25S.

Gonzalez Engelhard, C. A., A. P. Hodgkins, E. E. Pearl, P. K. Spears, J. Rychtář, and D. Taylor. 2021. A mathematical model of Guinea worm disease in Chad with fish as intermediate transport hosts. Journal of Theoretical Biology **521**:110683.

Gotelli, N. 2008. A Primer of Ecology Fourth edition.

Greenwood, D. 2012. 63 - Helminths: Intestinal worm infections; filariasis; schistosomiasis; hydatid disease. Pages 655-666 *in* D. Greenwood, M. Barer, R. Slack, and W. Irving, editors. Medical Microbiology (Eighteenth Edition). Churchill Livingstone, Edinburgh.

Grosholz, E., G. Ashton, M. Bradley, C. Brown, L. Ceballos-Osuna, A. Chang, C. de Rivera, J. Gonzalez, M. Heineke, M. Marraffini, L. McCann, E. Pollard, I. Pritchard, G. Ruiz, B. Turner, and C. Tepolt. 2021. Stage-specific overcompensation, the hydra effect, and the failure to eradicate an invasive predator. Proceedings of the National Academy of Sciences **118**:e2003955118.

Grunert, R., E. Box, K. Garrett, M. Yabsley, and C. Cleveland. 2022. Effects of Temephos (Abate®), Spinosad (Natular®), and Diflubenzuron on the Survival of Cyclopoid Copepods. Am J Trop Med Hyg **106**:818-822.

Hamer, G. L., U. D. Kitron, J. D. Brawn, S. R. Loss, M. O. Ruiz, T. L. Goldberg, and E. D. Walker. 2008. Culex pipiens (Diptera: Culicidae): A Bridge Vector of West Nile Virus to Humans. Journal of Medical Entomology **45**:125-128.

Hanazato, T., T. Iwakuma, M. Yasuno, and M. Sakamoto. 1989. Effects of temephos on zooplankton communities in enclosures in a shallow eutrophic lake. Environmental Pollution **59**:305-314.

Hixon, M. A., and G. P. Jones. 2005. Competition, Predation, and Density-Dependent Mortality in Demersal Marine Fishes Ecology **86**:2847-2859.

Hurtado, P. J., and A. S. Kirosingh. 2019. Generalizations of the 'Linear Chain Trick': incorporating more flexible dwell time distributions into mean field ODE models. J Math Biol **79**:1831-1883.

Johnson, S. N., and S. Rasmann. 2015. Root-feeding insects and their interactions with organisms in the rhizosphere. Annu Rev Entomol **60**:517-535.

Karatayev, V. A., C. E. Kraft, and E. F. Zipkin. 2015. Racing through life: maturation rate plasticity regulates overcompensation and increases persistence. Ecosphere **6**:art203.

Kaspersson, R., F. Sundström, T. Bohlin, and J. I. Johnsson. 2013. Modes of Competition: Adding and Removing Brown Trout in the Wild to Understand the Mechanisms of Density-Dependence. PLOS ONE **8**:e62517.

Kilham, S., D. Kreeger, S. Lynn, C. Goulden, and L. Herrera. 1998. COMBO: A defined freshwater culture medium for algae and zooplankton. Hydrobiologia **377**:147-159.

Kramer, A. M., O. Sarnelle, and R. A. Knapp. 2008. Allee Effect Limits Colonization Success of Sexually Reproducting Zooplankton. Ecology **89**:2760-2769.

Kulkarni, D., U. Hommen, A. Schäffer, and T. G. Preuss. 2014. Ecological interactions affecting population-level responses to chemical stress in Mesocyclops leuckarti. Chemosphere **112**:340-347.

Lambin, X., M. Begon, S. J. Burthe, I. M. Graham, J. L. MacKinnon, S. Telfer, and M. K. Oli. 2025. Density-dependent recruitment but not survival drives cyclic dynamics in a field vole population. Proceedings of the National Academy of Sciences **122**:e2509516122.

Leboulanger, C., C. Schwartz, P. Somville, A. O. Diallo, and M. Pagano. 2011. Sensitivity of Two Mesocyclops (Crustacea, Copepoda, Cyclopidae), from Tropical and Temperate Origins, to the Herbicides, Diuron and Paraquat, and the Insecticides, Temephos and Fenitrothion. Bulletin of Environmental Contamination and Toxicology **87**:487-493.

Lenth, R. V. 2025. emmeans: Estimated Marginal Means, aka Least-Squares Means.

Lindquist, H. D. A., and J. H. Cross. 2017. 195 - Helminths. Pages 1763-1779.e1761 *in* J. Cohen, W. G. Powderly, and S. M. Opal, editors. Infectious Diseases (Fourth Edition). Elsevier.

Lu, A., and L. Wang. 2025. Hygiene conditions explain larval density-dependent performance in Plutella xylostella with sufficient food. Journal of Asia-Pacific Entomology **28**:102444.

Lyons, G. R. 1972. Guineaworm infection in the Wa district of north-western Ghana. Bull World Health Organ **47**:601-610.

McCullough, F. S. 1983. Cyclopoid copepods: their role in the transmission and control of dracunculiasis. World Health Organization, Geneva, Ecology and Control of Vectors Unit, Division of Vector Biology and Control.

McIntire, K. M., and S. A. Juliano. 2018. How can mortality increase population size? A test of two mechanistic hypotheses. Ecology **99**:1660-1670.

Moorthy, V. N. 1938. Observations on the development of Dracunculus medinensis larvae in Cyclops. American Journal of Hygiene.

Neale, Z. R., and S. A. Juliano. 2019. Finding the sweet spot: What levels of larval mortality lead to compensation or overcompensation in adult production? Ecosphere **10**:e02855.

Newman, K., R. King, V. Elvira, P. de Valpine, Rachel S. McCrea, and B. J. T. Morgan. 2023. State-space models for ecological time-series data: Practical model-fitting. Methods in Ecology and Evolution **14**:26-42.

Ni, J., J. Wang, W. Zhang, E. Chen, Y. Gao, J. Sun, W. Huang, J. Xia, W. Zeng, J. Guo, and Z. Gong. 2025. Sustainable control and integrated management through a one health approach to mitigate vector-borne disease. One Health **20**:101018.

Nilsson, K. A., L. Persson, and T. Van Kooten. 2010. Complete compensation in Daphnia fecundity and stage-specific biomass in response to size-independent mortality. Journal of Animal Ecology **79**:871-878.

Park, T., D. B. Mertz, W. Grodzinski, and T. Prus. 1965. Cannibalistic Predation in Populations of Flour Beetles. Physiological Zoology **38**:289-321.

Patterson, J., M. Sammon, and M. Garg. 2016. Dengue, Zika and Chikungunya: Emerging Arboviruses in the New World. West J Emerg Med **17**:671-679.

Pellegrino, C., G. Patti, M. Camporeale, A. Belati, R. Novara, R. Papagni, L. Frallonardo, L. Diella, G. Guido, E. De Vita, V. Totaro, F. V. Segala, N. Veronese, S. Cotugno, D. F. Bavaro, G. Putoto, N. Bevilacqua, C. Castellani, E. Nicastri, A. Saracino, and F. Di Gennaro. 2022. Guinea Worm Disease: A Neglected Diseases on the Verge of Eradication. Trop Med Infect Dis **7**.

R Core Team. 2023. R: A Language and Environment for Statistical Computing.

Rivero, A., J. Vézilier, M. Weill, A. F. Read, and S. Gandon. 2010. Insecticide Control of Vector-Borne Diseases: When Is Insecticide Resistance a Problem? PLOS Pathogens **6**:e1001000.

Sahandi, J., P. Sorgeloos, H. Xiao, F. Mu, X. Wang, Z. Qi, Y. Zheng, and X. Tang. 2023. The use of microbes in culture of harpacticoid copepod (Tigriopus japonicus Mori) for suppression of Vibrio and enhancement of population growth, enzyme activity and microbial longevity. Aquaculture **563**:739008.

Sebens, K. P. 1982. Competition for Space: Growth Rate, Reproductive Output, and Escape in
Size. The American Naturalist **120**:189-197.
Simonetti, O., V. Zerbato, C. Maurel, L. Cosimi, S. Babich, F. Cavalli, S. Di Bella, D. Pavia, C.
Pesaresi, and R. Luzzati. 2023. The current state of knowledge on dracunculiasis: a
narrative review of a rare neglected disease. Infez Med **31**:500-508.
Smith, G. C. 1997. An analysis of the form of density dependence in a simulation model of a
seasonal breeder undergoing control. Ecological Modelling **95**:181-189.
South Sudan Guinea Worm Eradication Program. 2013. Management and Technical
Guidelines.*in* M. o. H.-R. o. S. Sudan, editor.
Steib, K., and P. Mayer. 1988. Epidemiology and vectors of Dracunculus medinensis in
northwest Burkina Faso, West Africa. Annals of Tropical Medicine & Parasitology
82:189-199.
Stockseth, L., Z. Neale, and V. H. W. Rudolf. 2025. Strengthening of negative density
dependence mediates population decline at high temperatures. Ecology **106**:e70030.
Svanbäck, R., and D. I. Bolnick. 2007. Intraspecific competition drives increased resource use
diversity within a natural population. Proceedings of the Royal Society B: Biological
Sciences **274**:839-844.
Telleria, E. L., J. B. Benoit, X. Zhao, A. F. Savage, S. Regmi, T. L. A. e Silva, M. O'Neill, and S.
Aksoy. 2014. Insights into the Trypanosome-Host Interactions Revealed through
Transcriptomic Analysis of Parasitized Tsetse Fly Salivary Glands. PLOS Neglected
Tropical Diseases **8**:e2649.
van den Bosch, F., and W. Gabriel. 1994. A model of growth and development in copepods.
Limnology and Oceanography **39**:1528-1542.
van den Bosch, F., and B. Santer. 1993. Cannibalism in Cyclops abyssorum. Oikos **67**:19-28.
Walker, M., M. A. Robert, and L. M. Childs. 2021. The importance of density dependence in
juvenile mosquito development and survival: A model-based investigation. Ecological
Modelling **440**:109357.
World Health Organization. 1986. Decisions and list of resolutions. World Health Assembly.
World Health Organization. 2025. Dracunculiasis (Guinea-worm disease).

Figure Captions

Figure 1. Mean observed copepod densities in the experiment over the 14 weeks in which the

interventions were applied on schedule. Points and error bars represent treatment means ± SEs

(log-transformed). Regardless of treatment type or removal proportion, there were no significant

reductions in total copepod density (squares), or stage-specific densities of egg-bearing adult

females (AF; diamonds), juveniles plus other adults (JOA; triangles), or nauplii (N; circles).

Similar dynamics were observed regardless of whether removal was selective for large-bodied

individuals (size-biased; dark blue) or equally applied across all stages (size-unbiased; light blue).

Figure 2. Observed copepod population dynamics (points and error bars represent means ± SE)

and best-fitting posterior predictions with 95% credible intervals (black lines and shaded grey

regions). Treatments that are representative of the major population dynamics are shown, with all

treatments can be found in the Appendix (Appendix S5: Figure S1A). Dynamics in control tanks

are plotted in green, size-biased removal tanks are plotted in light blue, and size-unbiased

removal tanks are in dark blue. Dashed vertical lines represent interventions. In general,

interventions caused immediate reductions in copepod abundance, but populations rebounded

rapidly. In some case, e.g., size-biased 50% and size-unbiased 90%, these rebounds surpassed

abundances observed in controls. For the 98% treatments, the first intervention was 20% removal,

explaining the small dip in the model fit.

Figure 3. Representation of the sources of negative density dependence affecting copepod

populations subject to control interventions. Lines and shaded regions represent mean and 95%

credible intervals for posterior model predictions for the effects of copepod population density

(with fixed stage structure) on key vital rates: birth rate (green), total death rate of nauplii (dark

blue; incorporating background deaths and cannibalism via adults), and naupliar maturation rate

(light blue). The rug indicates the observations of population density in the experiment. Adult

birth rate was insensitive to competition, declining only 0.9% over the observed density gradient

(indicated by the rug along the x-axis). In contrast, death rate increased 4.4-fold over the density

gradient. Most importantly, maturation rate was extremely sensitive, declining >18,000 fold over

the gradient.

Figure 4. Temporal dynamics from the modeled transmission scenarios incorporating 90%

mortality events at 28-day intervals. Dashed vertical lines represent interventions. Copepod

stages are shown in different colors. Stages are divided into infectious adults (I), juveniles (J),

nauplii (N), and non-infectious adults (NI = susceptible + exposed adults). Control tanks are

dashed lines, 90% size-unbiased removal are solid lines, and 90% size-biased removal are the

dotted lines. The force of infection parameter (λ =0.0034) corresponded with a ~5% infection

rate among adults(Lyons 1972, Steib and Mayer 1988, Garrett et al. 2020). Nauplii and non-

infected adults surpass densities of controls (green) with both forms of removal (size-biased

[dark blue] and size-unbiased [light blue]). Juveniles nearly recover to densities akin to controls.

Most importantly, infectious copepods are severely depressed by both removal types.

Figure 5: Mean total density of copepod populations (light blue), non-infectious adults (green),

and infectious adults (dark blue) in 120-day transmission scenarios modeled along a gradient of

intervention intensity for size-biased (solid line) and size-unbiased (dashed line) strategies. As in

the experiment, total density and the density of non-infectious adults (susceptible and exposed)

were largely insensitive to interventions, remaining relatively constant despite 90% population

reductions imposed every 28 days. In contrast, the mean density of infectious adults, the

proximate driver of mammalian hazard of infection, decreased almost 10-fold along the same

gradient. Thus, despite the rapid recovery of copepod populations and the appearance of failed

control, intervention can be highly effective at reducing the proximate driver of infection in

humans and other mammals.

704   **Figure 1**

713   **Figure 2**

722   **Figure 3**

731   **Figure 4**

739   **Figure 5**