Title:

Deep immune profiling pinpoints cellular and molecular drivers of lupus immunopathology

Authors:

Masahiro Nakano[1]†*, Michihiro Kono[1,2]†, Kenichiro Asahara[1,3], Takayuki Katsuyama[4], Satoshi Kubo[5], Eri

Katsuyama[4,6], Yuya Fujita[5], Takahiro Nishino[1,2], Hajime Inokuchi[1,2,7], Takahiro Arakawa[1], Tsugumi

Kawashima[1], Shohei Noma[1], Akiko Minowa[2], Reza Bagherzadeh[1,2], Yukiro Matsumoto[1,2,8], Jun Inamo[1,2],

Haruka Takahashi[1,2], Bunki Natsumoto[1], Xuejun Zhang[9,10,11], Sang-Cheol Bae[12,13,14], Akari Suzuki[15],

Hiroaki Hatano[1,2], Chikashi Terao[16,17,18], Yoshiya Tanaka[5,19], Yoshinori Matsumoto[4], Kazuhiko

Yamamoto[15], Kazuyoshi Ishigaki[1,2,20]*

Affiliations:

[1] Laboratory for Human Immunogenetics, RIKEN Center for Integrative Medical Sciences, Yokohama,

Japan

[2] Department of Microbiology and Immunology, Keio University School of Medicine, Tokyo, Japan

[3] Department of Ophthalmology, Graduate School of Medical Sciences, Kyushu University, Fukuoka,

Japan

[4] Department of Nephrology, Rheumatology, Endocrinology and Metabolism, Okayama University

Faculty of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan

[5] The First Department of Internal Medicine, University of Occupational and Environmental Health,

Kitakyushu, Japan

[6] Faculty of Health Science, Okayama University Medical School, Graduate School of Health Sciences,

Okayama University, Okayama, Japan

[7] Division of Rheumatology, Department of Internal Medicine, Keio University School of Medicine, Tokyo,

Japan

[8] Department of Dermatology, Keio University School of Medicine, Tokyo, Japan

[9] Department of Dermatology, the First Affiliated Hospital of Anhui Medical University, Hefei, Anhui,

China

[10] Institute of Dermatology, Anhui Medical University, Hefei, Anhui, China

[11] Key Laboratory of Dermatology (Anhui Medical University), Ministry of Education, Hefei, Anhui, China

[12] Department of Rheumatology, Hanyang University Hospital for Rheumatic Diseases, Seoul, Korea

[13] Hanyang Institute of Bioscience and Biotechnology, Seoul, Korea

[14] Hanyang University Institute for Rheumatology Research, Seoul, Korea

[15] Laboratory for Autoimmune Diseases, RIKEN Center for Integrative Medical Sciences, Yokohama,

Japan

[16] Laboratory for Statistical and Translational Genetics, RIKEN Center for Integrative Medical Sciences,

Yokohama, Japan

[17] Clinical Research Center, Shizuoka General Hospital, Shizuoka, Japan

[18] The Department of Applied Genetics, The School of Pharmaceutical Sciences, University of Shizuoka,

Shizuoka, Japan

[19] Department of Molecular Targeted Therapeutics, University of Occupational and Environmental Health,

Kitakyushu, Japan

[20] Keio University Human Biology-Microbiome-Quantum Research Center (WPI-Bio2Q)

† These authors contributed equally to this work.

*Correspondence:

Masahiro Nakano, M.D., Ph.D.

masahiro.nakano.fc@riken.jp

Kazuyoshi Ishigaki, M.D., Ph.D.

kazuyoshi.ishigaki@keio.jp

Abstract:

Systemic lupus erythematosus (SLE) is a complex autoimmune disease with an unknown etiology. To

pinpoint new disease-relevant cell states and their molecular profiles, we performed an in-depth

investigation of multimodal single-cell datasets comprising ~2.1 million peripheral blood mononuclear

cells from 346 donors. By resolving 123 fine-grained cell states across 27 cell types, we identified

previously uncharacterized populations distinctively associated with clinical severity and treatment

status, including $GZMK^+GZMH^+HLA\text{-}DR^+$ effector memory CD8$^+$ T cells (double-positive [DP] EMCD8)

and $FOXO1^+ARHGAP15^+$ T cells. Through extensive statistical frameworks and multimodal approaches,

we delineated their aberrant immune signaling networks, transcriptional regulators, key surface proteins,

T cell receptor repertoires, and genetic/epigenetic landscapes, underscoring them as candidate drivers

of SLE immunopathology. These findings provide new insights into therapeutic target discovery in SLE.

Main Text:

Introduction

Systemic lupus erythematosus (SLE) is a prototypic autoimmune disease with a broad spectrum of

clinical manifestations, triggered by genetic and environmental factors (*1*). Since the pathophysiology of

SLE involves various immune cell types, many bulk and single-cell transcriptome studies on peripheral

blood and affected organs have been conducted (*2–6*). While these efforts have begun to reveal key

gene signatures and core cell populations of SLE, the fine-resolution pathophysiology remains

unresolved, and the treatment options are still limited; to date, only two molecular-targeted drugs have

been approved for this disease (*7, 8*).

To understand the complex biology of SLE and identify therapeutic targets, it is crucial to

characterize disease-relevant cell states and their molecular signatures at high resolution. We previously

conducted a large-scale bulk transcriptome study across 27 immune cell types, identifying cell-type-

specific signatures linked to SLE development and exacerbation (*6*). However, as an inherent limitation

of bulk studies, we failed to identify more granular cell states within each cell type. While single-cell RNA

sequencing (scRNA-seq) can theoretically resolve hundreds of cell states, previous SLE studies have

not fully leveraged this potential, profiling only ~20 major cell types (*5*) or ~40 cell clusters (*3, 4*), thereby

lacking deeper mechanistic insights. Given that low-frequency cell populations often play pivotal roles in

various diseases (*9–13*)—such as peripheral helper T cells (Tph) and autoimmune-associated B cells

(ABCs) in autoimmunity—a comprehensive identification of disease-relevant cell states and their

orchestrated networks is essential to elucidating the complex pathophysiology and clinical heterogeneity

of SLE. Moreover, most single-cell studies have not formally distinguished between quantitative cellular

abundance shifts and qualitative transcriptional changes within specific cell types. Although recent tools

such as milo (*14*) and CNA (*15*) enable the detection of continuous quantitative shifts, they are not

designed to directly evaluate qualitative changes. Thus, it is important to extend these concepts and

construct a statistical framework capable of jointly dissecting these two scenarios. Finally, as previous

SLE studies have focused primarily on gene expression (*3–5*), integrating multi-layered information is

paramount to uncovering novel therapeutic targets within the complex molecular architecture of the

disease.

To address these gaps, we extensively investigated multimodal single-cell data of SLE with a

total of ~2.1 million cells from 346 donors (Lupus-derived Peripheral immune cell Integrated panels

[LuPIN]) to pinpoint granular disease-relevant cell states (**Fig. 1**). Using fine-grained clustering and

rigorous validation, we defined 123 cell states across 27 immune cell types. Notably, we identified

several previously uncharacterized cell states specifically tracking with severe clinical phenotypes and

treatment response. Our statistical framework, dissecting both quantitative and qualitative gene

signatures, indicated that key immunological pathways and SLE risk variants were closely linked to cell-

state compositional shifts within each cell type. Furthermore, these newly identified cell populations were

predicted to engage in intricate immune crosstalk with established pathogenic cell states, potentially

driving SLE pathogenesis. Finally, in-depth characterization delineated the multifaceted molecular

landscapes of these populations, including transcriptional regulators, epigenetic features, surface

proteins, and T-cell receptor (TCR) repertoires. Overall, our intensive approach elucidates the complex

immunopathology of SLE, providing a foundation for novel therapeutic strategies.

Results

Overview of this study

LuPIN comprises three datasets, each with a different purpose (**Fig. 1, left**). To robustly identify disease-

relevant cell states, we first used the largest, publicly available 3'-end scRNA-seq dataset for SLE (*5*)

(dataset 1; **table S1**). Our stringent multi-step quality control (QC) process yielded the final dataset

(1,456,041 peripheral blood mononuclear cells [PBMCs] from 97 healthy and 172 SLE samples [159

unique SLE donors: 11 donors have 2~3 timepoints]; **fig. S1**). Among 172 SLE samples, 21 exhibited

high disease activity (HDA), and 40 were clinically inactive (**Methods**). Following our previous study

scheme (*6*), we established two comparison axes: (i) disease-state comparison between healthy controls

and inactive SLE patients, reflecting the biology of disease development, and (ii) disease-activity

comparison between inactive vs. HDA SLE, reflecting the biology of disease exacerbation. For

replication analysis, we used another scRNA-seq dataset (*4*) (dataset 2, 344,235 PBMCs from 17

healthy and 41 SLE samples).

120   To overlay in-depth molecular information onto the fine-grained cell state map, we additionally

prepared a new multimodal 5'-end dataset using cellular indexing of transcriptomes and epitopes by

sequencing (CITE-seq)—a technology profiling gene expression and surface marker proteins from single

cells (*16*)—alongside TCR-seq and enhancer RNA profiling (*17*, *18*) (dataset 3, 268,783 PBMCs from 32

SLE samples). Through the efficient integration with dataset 1, we searched for key surface proteins,

TCR repertoires, and cis-regulatory elements specific to disease-relevant cell states.

Definition of 123 robust cell states using fine-grained clustering

To identify granular cell states of SLE, we employed a three-step intensive clustering approach in

dataset 1: lineage (layer 1), cell type (layer 2), and cell state (layer 3) (**Fig. 2**). First, Louvain clustering

categorized 1,456,041 PBMCs into seven cell lineages (layer 1: CD4$^+$ T cells [CD4 T], CD8$^+$ and

unconventional T cells [CD8 and other T], natural killer cells [NK], B cells, monocytes [Mono], dendritic

cells [DC], and hematopoietic stem and progenitor cells [HSPC]; **fig. S1D−F**). These lineages were

further subdivided into 27 coarse cell types (layer 2: **Fig. 2A−F**, **fig. S2A−F**, and **table S2**). Our analytical

pipelines (**table S3** and **Methods**) achieved higher annotation accuracy than the original study(*5*) (**fig.**

S2G).

To determine if these coarse cell-type-level annotations sufficiently explain disease-relevant

compositional changes, we performed co-varying neighborhood analysis (CNA) to identify single-cell-

resolution transcriptional neighborhoods associated with clinical phenotypes (*15*). We applied CNA to six

cell lineages (excluding HSPC) and examined the abundance of 26 coarse cell types (lineage-level CNA;

fig. S2H). The extent of expansion or depletion for each cell type was quantified at the single-cell level

as "neighborhood correlations". Using two comparison axes—disease-state (inactive SLE vs. healthy,

reflecting disease development) and disease-activity CNA (HDA vs. inactive SLE, reflecting disease

exacerbation) (*6*)—we recapitulated the depletion of NaiveCD4 and mucosal-associated invariant T cells

(MAIT) in disease-state comparison (*5*), while revealing that these patterns were remarkable in disease-

activity comparison. Similarly, regulatory T cells (Treg), proliferating (Prolif) CD4/CD8/NK cells, and $C1Q^+$

monocytes (Mono) expanded in disease-state and/or -activity CNA. Although neighborhoods in these cell

types exhibited universal expansion or depletion patterns, many other cell types showed heterogeneous

patterns in lineage-level CNA (**fig. S2H**), underscoring the need for a more fine-grained investigation.

To efficiently capture subtle gene-program heterogeneities within each cell type, we further

performed clustering on each cell type and defined 123 cell states with unique gene expression profiles

(layer 3: **Fig. 2A−F**, **fig. S3**, and **table S4**). As these cell-state definitions form the basis for downstream

analyses, we employed multiple rigorous approaches to ensure their robustness. First, a leave-one-

batch-out analysis, systematically excluding each of the 23 experimental batches from dataset 1,

confirmed the consistent preservation of all 123 cell states, albeit occasionally requiring slightly higher

resolution parameters (**Fig. 2G, left**, **fig. S4A**, and **table S5**). Moreover, 122 cell states (except for

gamma-delta T cells [gdT]-5: *ISG15*+*MX1*+) were replicated through independent clustering of datasets 2

and/or 3, demonstrating the cross-dataset reproducibility (**Fig. 2G, right**, and **fig. S4B−D**). Second, the

identity of each cluster was corroborated by an extensive literature review (*9–13*, *19–39*) (**table S6**).

Third, marker genes upregulated in each state exhibited minimal overlap within the same cell type (**fig.**

S4E), and pathway analysis using these markers revealed enrichment of cell-state-specific pathways

(**table S7**), supporting their non-redundancy and biological relevance. Finally, multimodal investigations

using dataset 3 identified cell-state-specific surface proteins and TCR features (**Fig. 6−7**), providing

orthogonal validation at the immune phenotypic level.

Our fine-grained annotations successfully captured multiple disease-relevant cell state

candidates (**Fig. 2**), most of which remained unidentified in the original study due to insufficient

resolution. We first recapitulated established pathogenic states including: (i) follicular and peripheral

helper T cells (Tfh and Tph) within central and effector memory CD4 T cells (CMCD4-3: *ICOS*+*PASK*+

and EMCD4-7: *PDCD1*+*TIGIT*+) and (ii) autoimmune-associated B cells (ABCs) in memory B cells

(MemB-6: *ITGAX*+*FCRL5*+), which are central to autoantibody production (*9*, *10*, *27*). We also confirmed

the presence of subpopulations highly activated by interferon (IFN) signaling (*ISG15*+*MX1*+ cell states)−a

hallmark signature of SLE (*1*, *26*)−across most cell types. The shared gene program across cell types

was not limited to IFN signaling; JUN^+CD69^+ cell states with upregulated AP-1 transcription factors (TFs;

e.g., *JUN, FOS*) and early activation markers (e.g., *CD69* and *NFKBIA*) (*21*) were identified in most

lymphoid cell types (**fig. S3A−D**). Importantly, we identified multiple previously uncharacterized cell

states with distinct gene programs, some of which were later found to expand especially in severe SLE

(e.g., EMCD8-5: $GZMK^+GZMH^+HLA\text{-}DRB1^+$, **Fig. 3**). Moreover, our approach captured very rare

subpopulations such as $ANXA2^+CTLA4^+$ double-negative T cells (dnT-1) (*32*), $RORA^+GATA3^+$ innate

lymphoid cells (ILC-1) (*33*), $MZB1^+PCNA^+$ plasmablasts (PlasmaB-1) (*35*), and $LAMP3^+CCR7^+$ DCs

(cDC-5) (*39*). Overall, our intensive clustering approach provided high-resolution cell state information for

SLE, serving as the foundation for subsequent analyses.

Mapping cell state compositional changes in SLE

To assess the cell-state compositional changes within each cell type, we next applied CNA to 18 major

cell types (>5,000 cells; cell-type-level CNA) in disease-state (inactive SLE [$n = 40$] vs. healthy [$n = 97$])

and disease-activity (HDA [$n = 21$] vs. inactive SLE [$n = 40$]) comparisons, accounting for age, sex, and

ancestry. Strikingly, we observed skewed neighborhood abundance in 34 out of 36 tests (18 cell types ×

2 comparisons, global $P < 0.05/36$), and 29 cell states showed evidence of remarkable expansion (**Fig.**

3A−B, **fig. S5A**, and **table S8**). These expanded cell states tended to be rare within each cell type,

supporting the importance of the fine-grained strategy (**Fig. 3C** and **fig. S5B**).

Among the 29 cell states exhibiting remarkable expansion,18 recapitulated known signals. All

IFN-activated ($ISG15^+MX1^+$) cell states within each cell type expanded in disease-state and -activity CNA

(**Fig. 3D**). While Tfh (CMCD4-3) expanded primarily in disease-state, Tph (EMCD4-7) expanded

specifically in disease-activity CNA. Notably, in memory B cells (MemB)—which harbor the highest

enrichment of SLE risk variants (*40*)—ABC (MemB-6) showed remarkable expansion in disease-activity

CNA but not in disease-state CNA (**Fig. 3A**), supporting their relevance in pathogenic autoantibody

production, particularly during disease activation (*10*). Furthermore, our ancestry-specific analysis

revealed the expansion of this population in Asian SLE compared to European (EUR) SLE (**fig. S5C**),

suggesting a potential need for stratified medicine using B-cell targeted therapies (*41*, *42*).

The remaining 11 cell states represented previously unreported expansions in SLE (*3–5*),

nominating them as new disease-relevant candidates. An example is $GZMK^+GZMH^+HLA\text{-}DRB1^+$ cells

(EMCD8-5), which showed expansion in both disease-state and -activity CNA (**Fig. 3A−B**). While other

EMCD8 cell states express either *GZMK* or *GZMH*, this population expresses both, along with *GZMB* to

a lesser extent (**fig. S3B**; hereafter "double-positive [DP] EMCD8"). Notably, this population exhibited

high expression of multiple T cell activation markers, including *HLA-DR* genes, *PDCD1*, and *TIGIT* (**fig.**

S3B). Another example, $FAM13A^+CDK6^+$ cells (NaiveCD4-4), further highlighted the necessity of fine-

grained analysis: although NaiveCD4 cells were, on average, depleted within the CD4 T lineage (**fig.**

S2H), this depletion pattern was not uniform across NaiveCD4 cell states. Lastly, $FOXO1^+ARHGAP15^+$

cell states expanded across most T-cell cell types in disease-state and -activity CNA (**Fig. 3A−B, D**).

These states were broadly shared across the majority of SLE samples, rather than being derived from a

limited number of donors (**fig. S5D**). Some cell states showed different patterns of abundance changes

between disease-state and activity CNA. While JUN^+CD69^+ cell states expanded in disease-state CNA,

they tended to deplete in disease-activity CNA across most cell types (**Fig. 3D**), suggesting distinct roles

of these states in SLE depending on the disease phase (*6*).

Our CNA results demonstrated high reproducibility in multiple approaches. First, cluster-based

abundance testing (MASC) (*11*) yielded concordant results with CNA (**fig. S5E**). Second, most CNA

signals were globally consistent between Asian and EUR ancestries (**fig. S5F**). Third, by independently

applying CNA to low (LDA) and moderate disease activity (MDA) cases—both of which were excluded

from the discovery set—we observed concordant signals with a gradual increase in disease-activity

effects, reflecting the continuous spectrum of SLE activity (**Fig. 3E**). Lastly, disease-state and activity

CNA using independent dataset 2 well replicated the signals detected in dataset 1 (**Fig. 3F**). Collectively,

these results robustly captured cell-state compositional changes linked to disease development and

exacerbation of SLE.

Statistical framework to dissect quantitative and qualitative changes within cell types

Having identified SLE-associated cellular compositional shifts within each cell type, we next introduced a

statistical framework to address the limitations of conventional cell-type-level analyses. Cell-type-level

analyses define genes driven by either of the following two distinct scenarios as disease-associated

differentially expressed genes (DEGs, **Fig. 4A**): (i) a gene is specifically expressed in a particular cell

state whose abundance shifts with phenotype (quantitative change) or (ii) a gene's per-cell expression is

directly dysregulated by the phenotype, regardless of cell-state abundance (qualitative change). To

dissect their respective pathogenic effects, our generalized linear mixed-effect model jointly regresses

each single-cell expression by (i) a phenotype-driven cell-state abundance term and (ii) a direct

phenotype term (e.g., inactive SLE vs. healthy) capturing per-cell gene dysregulation, while adjusting for

donor/batch effects and other covariates (**Fig. 4B**; **Methods**). Specifically, we incorporated single-cell-

level neighborhood correlations from cell-type-level CNA (**Fig. 3**) into the model as metrics for cell-state

abundance changes. By applying this model across 18 major cell types, we identified (i) "abundance

signature genes" (quantitative changes) and (ii) "dysregulated signature genes" (qualitative changes,

table S9). Notably, quantitative changes drove approximately 50% of the variation in the top 1,000

pseudobulk DEGs for each cell type (**fig. S6A**).

We tested the robustness of this model through multiple approaches. First, permutation tests

showed that this model is well calibrated (**fig. S6B**). Second, a leave-one-batch-out analysis,

systematically excluding each batch from dataset 1, confirmed the stability of model statistics across

genes (**fig. S6C**), demonstrating the robustness to technical confounders and donor-specific effects.

Finally, application of this model to dataset 2 showed good reproducibility of model statistics (**fig. S6D**).

For simplicity, we hereafter present disease-state comparison results in **Fig. 4C−F** and disease-activity

results in **fig. S6E−F**.

IFIH1 and *OAS1* are IFN response genes (IRGs) linked to the genetic loci in genome-wide

association studies of SLE (*43*) (SLE-GWAS; **table S10**). While cell-type-level pseudobulk analysis

identified both as DEGs in NaiveCD4 (false discovery rate [FDR] < 0.05), our model revealed different

behaviors of qualitative and quantitative changes for these genes (**Fig. 4C**). While *OAS1* expression

localized to NaiveCD4-7 (*ISG15$^+$MX1$^+$*), which expanded in SLE (abundance signature gene), *IFIH1* did

not show such localization and was globally overexpressed in SLE compared with healthy cells

(dysregulated signature gene). Hierarchical clustering of 100 IRGs (**table S11**) based on model statistics

across 18 cell types identified distinct modules representing abundance and dysregulated signature

genes, respectively, suggesting heterogeneous roles for IRGs in SLE pathophysiology (**Fig. 4D** and **fig.**

S6E).

Next, we performed pathway enrichment analyses on these signatures for each cell type (**Fig.**

4E). IFN signaling was enriched in both dysregulated and abundance signatures across almost all cell

types (FDR < 0.05; one-sided Fisher's exact test), demonstrating that IFN signaling exerts pathogenic

effects through both quantitative and qualitative changes. In contrast, other classical immunological

pathways were primarily enriched in abundance signature genes of relevant cell types (e.g., complement

signaling in myeloid cells; **Fig. 4E**, **fig. S6F**, and **table S12**), suggesting that key biological processes in

SLE are more closely linked to abundance signatures than dysregulated signatures.

Finally, we integrated these model statistics with the largest-scale SLE-GWASs in East Asian

(EAS) (*44*) and EUR (*45*). We evaluated the polygenic enrichment of SLE risk variants around signature

genes using stratified linkage disequilibrium score regression (S-LDSC) (*40*), followed by a meta-

analysis of the EAS and EUR results **(fig. S6G)**. While SLE risk variants were enriched around both

dysregulated and abundance signature genes across multiple cell types (FDR < 0.05), the enrichment

was more pronounced for abundance signatures ($P = 5.3 \times 10^{-4}$; paired Wilcoxon test; **Fig. 4F**). Notably,

this enrichment was specifically observed in abundance signature genes with positive statistics (**fig.**

S6H). Consistent with previous studies (*40*), strongest enrichment was identified in B cells. Although

these findings do not imply that SLE risk variants directly regulate cell-state compositional changes, our

framework suggests that SLE risk genes exert their effects primarily within these expanded cell states.

Characterization of aberrant immune networks between disease-relevant cell states

Motivated by our finding that key immunological pathways and SLE risk variants were predominantly

enriched in cell-state compositional changes (**Fig. 4**), we further investigated their clinical relevance.

Hierarchical clustering of all samples and cell states based on their proportions revealed that samples

broadly partitioned into SLE (sample group [SG]1) and healthy controls (SG2) (**Fig. 5A**). While a subset

of SLE patients clustered within SG2, SG1 was further stratified into four subgroups (SG1-1 to SG1-4).

Similarly, clustering of cell states identified six cell-state groups (CGs). Intriguingly, while all IFN-

activated ($ISG15^+MX1^+$) cell states correlated within CG3, other SLE-expanded states—including well-

known pathogenic (Tfh, Tph, and ABCs) and novel populations (e.g., DP EMCD8 and

$FOXO1^+ARHGAP15^+$ cells)—formed another distinct group (CG2). This suggests that these populations

are immunologically interconnected, partially independent of IFN signaling. Indeed, SG1-1 was

characterized by a remarkable expansion of CG3, while SG1-2 (and a subset of SG1-3) exhibited CG2

expansion, highlighting the cellular heterogeneity underlying SLE. Both SG1-1 and SG1-2 comprised a

higher proportion of HDA patients; conversely, SG1-4 and SG2 were enriched for inactive or LDA

patients (**Fig. 5B**). SG1-4 showed JUN^+CD69^+ (CG6) expansion (**Fig. 5A**), potentially marking a clinically

quiescent state. We found no significant associations between SGs and organ activity or treatment

agents (**fig. S7A**).

Leveraging longitudinal samples in dataset 1 (from "Flare" to "Treated" samples from 10

donors), we compared changes in cell-state proportions. Notably, while many CG2 cell states (e.g., Tph

and DP EMCD8) decreased following treatment, CG3 ($ISG15^+MX1^+$) remained stable or even increased

(**Fig. 5C** and **fig. S7B**), suggesting that CG2 might serve as a cellular biomarker reflecting disease

activity and treatment response.

297   Next, we inferred the inflammatory networks underlying these correlated cell states. By

298 evaluating 4,701 ligand−receptor (LR) interactions (*46*) across all 123 × 123 cell-state pairs, we screened

299 an extensive signaling space and identified 1,451,770 potentially active signals (LR score > 0.6;

300 **Methods**). To minimize redundancy, we prioritized the most upregulated cell-state pair as the

301 representative for each unique LR interaction within each of the 27 × 27 cell-type pairs. Among these

302 118,629 cell-state-specific interactions (of which 52,599 occurred within 18 × 18 major cell-type pairs),

303 9,428 were predicted to occur between states within the same CG. Intriguingly, within-CG2 interactions

304 accounted for more than one-third (3,386) of these signals, suggesting a prominent immune network

305 among SLE-expanded cell states (**Fig. 5D**). Within CG2, we delineated both established and novel

306 candidate disease-relevant networks (**Fig. 5E** and **table S13**). First, our analysis recapitulated enhanced

307 IL21-IL21R/IL2RG signaling from EMCD4-7 (Tph) to NaiveB-4 ($SOX5^+FCRL5^+$), a signaling axis

308 essential for their differentiation into MemB-6 (ABC) (*10*). Notably, we identified potential IFNG-IFNGR

309 signaling from EMCD8-5 (DP EMCD8) to NaiveB-4, which is also implicated in ABC differentiation (**Fig.**

310 **5E−F**), suggesting that Tph and DP EMCD8 may synergistically drive the maturation of pathogenic B

311 cells (*10*, *47*). Furthermore, Tph and DP EMCD8 were predicted to engage in bidirectional costimulatory

312 crosstalk (e.g., CD70-CD27 signaling) (*48*, *49*), mutually enhancing their effector functions (**Fig. 5E, G**).

313 Additionally, other CG2 cell states, including NaiveCD4-4 ($FAM13A^+CDK6^+$), NaiveCD4-5 and CMCD4-4

314 ($FOXO1^+ARHGAP15^+$), were predicted to amplify these pro-inflammatory cascades (*50–52*) (**Fig. 5E** and

fig. S7C−E). Collectively, high-resolution cell–cell interaction analysis provided a putative map of intricate

immune networks between established and newly identified pathogenic cell states, potentially driving the

complex immunopathology of SLE.

Multimodal molecular profiling across fine-grained cell states

To characterize multi-layered molecular profiles across fine-grained cell states, we prepared a new

dataset (dataset 3; 32 SLE donors) comprising CITE-seq (130 surface marker proteins), TCR-seq, and 5'

transcript-based enhancer RNA (eRNA) profiling. Reference mapping (*53*) successfully integrated

268,783 PBMCs from dataset 3 into the latent transcriptional space in dataset 1, extending our fine-

grained analysis across multimodal layers (**fig. S8A**). First, we comprehensively interrogated cell-state-

specific surface protein expression, identifying an average of nine upregulated proteins per cell state

(FDR < 0.05; **fig. S8B** and **table S14**). This analysis recapitulated well-established markers, such as (i)

CD278 (ICOS), CD279 (PD1), and TIGIT in Tph (EMCD4-7) (*9*), (ii) CD11c and CD19 in ABC (MemB-6)

(*10*), and (iii) CD38 and CD31 in $SOX4^+$ naive T cell states (NaiveCD4-3 and NaiveCD8-3) (*22*).

Furthermore, we identified several key proteins characterizing new disease-relevant cell states, which

facilitated our subsequent downstream investigations (**Fig. 6−7**).

We next examined TCR repertoire profiles. While we and others have shown that V gene usage

differs across cell types (*54, 55*) and represents a potential therapeutic target for autoimmunity (*56*),

systematic analysis at the cell-state resolution has been lacking. Principal component analysis (PCA) of

V-gene frequencies confirmed that cell-state embeddings globally clustered by cell type (**fig. S8C**).

Notably, among 94 V genes (45 α- and 49 β-chains), ~3.8 V genes per cell state exhibited skewed usage

(FDR < 0.05; two-sided Fisher's exact test; **fig. S8D** and **table S15**), highlighting distinct TCR signatures

unique to specific cell states. While most of them were observed in EMCD8 cell states, we identified

several cell-state-specific V genes in other cell types (e.g., NaiveCD4-5; **Fig. 6**).

Leveraging 5' transcript sequencing, we mapped transcription start sites (TSSs) at single-cell

resolution. Recent studies have shown that bidirectional transcription around TSSs marks enhancer

activity (*17*, *18*). Using this approach, we defined ~1,564 transcribed enhancers per cell type (**Methods**),

which were enriched for established PBMC enhancer marks (*57*) (**fig. S8E**). Among them, we identified

~68 cell-state-specific enhancers per cell type (**table S16**). Thus, the efficient integration of multimodal

data with large-scale scRNA-seq data provided a comprehensive molecular map of fine-grained cell

states.

FOXO1 **is a candidate driver of the pathogenic CD4$^+$ T cell state gene program**

We next investigated the molecular profiles of disease-relevant cell state candidates that exhibited

expansion in both disease-state and disease-activity CNA (**Fig. 3A−B**). Specifically, we interrogated the

molecular features of newly identified CD4$^+$ T cell states within CG2 (**Fig. 5**): (1) *FAM13A$^+$CDK6$^+$* cells

(NaiveCD4-4), and (2) *FOXO1$^+$ARHGAP15$^+$* cells (NaiveCD4-5 and CMCD4-4).

Pathway analysis of upregulated marker genes revealed an enrichment of JAK-STAT signaling

in *FAM13A$^+$CDK6$^+$* cells and TCR signaling in *FOXO1$^+$ARHGAP15$^+$* cells, reflecting their distinct

immunological roles (**Fig. 6A** and **table S7**). Cell-state-specific enhancers near the marker genes

suggested unique epigenetic regulations of these states (**fig. S9A** and **table S16**). Regarding ligand-

receptor signaling, both cell states were predicted to engage in pro-inflammatory networks within CG2,

including ADA–DPP4, CALM1–PTPRA, MIF−CD74/CD44, and COPA−CD74 axes (*50–52*), which

interact with Tph and DP EMCD8 cells (**Fig. 5D**, **fig. S7D−F, and table S13**). Beyond the CG2 network,

these states exhibited unique ligand-receptor expression profiles, potentially forming extensive crosstalk

with multiple cell types (**Fig. 6B**). Consistent with the upregulated TCR signaling pathway, NaiveCD4-5

and CMCD4-4 highly expressed *CD28*, *TNFSF8*, *PTPRA*, *CD44*, and *ITGA4*, suggesting an "effector-

primed" states triggered by persistent antigen stimulation (*49*, *50*, *58*); indeed, NaiveCD4-5 further

exhibited a unique TCR V gene usage pattern (**Fig. 6C** and **table S15**).

Surface protein profiling further characterized these states: *FAM13A$^+$CDK6$^+$* cells highly

expressed CD127 (IL-7Rα) (FDR < 0.05; **fig. S9B** and **table S14**), reflecting their distinct homeostatic

requirements (*59*). In *FOXO1$^+$ARHGAP15$^+$* cells, the upregulation of specific activation markers including

CD244, CD32, CD33, and CD38 was consistent with chronic antigen exposure via TCR signaling (*60*,

61) (**Fig. 6D**). Intriguingly, these upregulations were less pronounced at the mRNA level (**table S4**),

highlighting the significance of multi-layered investigation (*13*, *16*, *19*). We also examined the localization

of GWAS-prioritized genes: in *FAM13A*$^+$*CDK6*$^+$ cells, the transcription coactivator *ARID5B* was linked to

an SLE-GWAS locus (*43*) (**Fig. 6E** and **table S10**). In *FOXO1*$^+$*ARHGAP15*$^+$ cells, multiple genes

including *ARHGAP15* and *PRKCB* were associated with SLE-GWAS loci.

To identify transcriptional regulators driving the gene program of *FAM13A*$^+$*CDK6*$^+$ and

FOXO1$^+$*ARHGAP15*$^+$ cells, we integrated computational and experimental approaches, focusing on 22

candidate TFs upregulated in these states: (i) pathway and TF enrichment analyses, TF activity

inference based on (ii) decoupleR (*62*) and (iii) curated ChIP-seq data (*63*) limited to blood tissue, and

(iv) an arrayed CRISPR-KO experiment in primary CD4$^+$ T cells (**fig. S9C** and **table S17−19**). Notably,

FOXO1 consistently showed significant enrichment for NaiveCD4-5 and CMCD4-4 signature genes

across all approaches (Bonferroni-corrected *P* < 0.05, **Fig. 6F**). Within EMCD4 cells, *FOXO1* exhibited

its highest expression in Tph (EMCD4-7; **Fig. 6G** and **fig. S3A**). Trajectory analysis (*64*) on whole CD4$^+$

T cells also suggested a potential transcriptional continuity between *FOXO1*$^+$*ARHGAP15*$^+$ cells and Tph.

Recent studies have established that *FOXO1* is essential for maintaining T cell stemness and memory

programming (*24*, *25*). While persistent antigen exposure drives the T-cell activation program in

FOXO1$^+$*ARHGAP15*$^+$ cells, *FOXO1* may act as a differentiation gatekeeper, preventing terminal

exhaustion and maintaining a long-lived reservoir of pathogenic effectors. In contrast, *ARID5B* did not

robustly account for the *FAM13A*⁺*CDK6*⁺ gene program across our approaches (**fig. S9D**), although the

limitations of the in vitro context must be considered. Together, our multi-layered investigation pinpointed

key regulatory candidates in pathogenic CD4⁺ T cell states of SLE.

DP EMCD8 cells as a candidate pathogenic player in SLE

Among the new cell states within CG2, we next performed in-depth investigation of DP EMCD8 (EMCD8-

5; *GZMK*⁺*GZMH*⁺*HLA-DRB1*⁺). Notably, this population shared several transcriptional signatures with

Tph beyond cell lineages. Pathway analysis revealed an enrichment of oxidative phosphorylation

signaling in both populations, suggesting shared metabolomic reprogramming (**Fig. 7A** and **table S7**).

Furthermore, DP EMCD8 exhibited the highest "classical" exhaustion signature (e.g., *PDCD1* and *TIGIT*,

table S20) within EMCD8, mirroring the pattern of Tph within EMCD4. (**Fig. 7B**). Similarly, *RGS1*—an

SLE GWAS-prioritized gene and exhaustion marker—was upregulated in both (**fig. S10A**). While these

signatures were traditionally regarded as markers of T cell exhaustion, recent studies in autoimmunity

indicate that they reflect a highly activated state induced by chronic antigen exposure, rather than a

dysfunctional state (*9*, *65*). Consistently, DP EMCD8 showed evidence of TCR clonal expansion in

multiple SLE donors (**Fig. 7C**), with some expanded clonotypes potentially reactive to cytomegalovirus or

Epstein-Barr virus epitopes (**table S21**). Although many clones were shared with other EMCD8 states,

nine clonotypes and nine V genes were specific to DP EMCD8 (FDR < 0.05; two-sided Fisher's exact

test; **fig. S10B−C, table S15** and **21**).

As discussed above (**Fig. 5E−G**), DP EMCD8 cells were predicted to engage in pro-

inflammatory networks with other pathogenic populations. Specifically, DP EMCD8 and Tph may

synergistically drive the maturation of pathogenic B cells (ABCs) via *IFNG* and *IL21* signaling, while

potentially maintaining a feed-forward activation loop through bidirectional costimulatory crosstalk (e.g.,

CD70−CD27). Longitudinal analysis also showed that both DP EMCD8 and Tph decreased following

treatment (**Fig. 5C**), tracking with disease activity. Furthermore, DP EMCD8 exhibited high expression of

pro-inflammatory cytokines (*CCL5*, *IL16*, and *IL32*) and functional markers (*CD74*, *CD2*, and *ITGA4*; **Fig.**

7D), underscoring their activated effector function rather than exhaustion. Intriguingly, while DP EMCD8

cells expressed lower *GZMB* levels than classical cytotoxic states (EMCD8-2; **fig. S3B**), they showed

marked *GZMA* upregulation (**Fig. 7D**), suggesting that they might also mediate caspase-independent

cytotoxicity (*66*).

To identify key surface proteins, we integrated the multimodal dataset 3 into dataset 1,

identifying HLA-DR, CD27, CD38, and CD28 as the most upregulated proteins in DP EMCD8 (**Fig. 7E**

and **table S14**). Notably, CD45RA was downregulated, indicating that these cells are distinct from

terminally differentiated effector memory (TEMRA) cells (*67*). In-silico gating on HLA-

DR$^+$CD27$^+$CD38$^+$CD28$^+$ achieved >90% purity for DP EMCD8 (**fig. S10D**), enabling validation via flow

cytometry (FCM) and FACS. We sorted target EMCD8 cells (CCR7$^-$HLA-DR$^+$CD27$^+$CD38$^+$CD28$^+$) and

performed bulk RNAseq and ATAC-seq (**Fig. 7F**, **fig. S10E**, and **table S22**), confirming a gene

expression profile comparable to DP EMCD8 (**fig. S10F**). Motif analysis for differentially accessible

regions (DARs) in this population compared with whole EMCD8 (**table S23**) showed minimal TF

enrichment (**fig. S10G**). Combined with the few upregulated TFs in this population (**table S4**), this

suggests that the primary driver of DP EMCD8 may be extrinsic humoral signals rather than intrinsic

transcriptional reprogramming.

Since no DARs were initially detected near the *IFNG* locus in standard ATAC-seq, we employed

UNIChro-seq (*68*) to enrich tagmented DNA fragments in this region (**table S24**). A ~20-fold enrichment

compared with standard ATAC-seq identified a unique DAR (**Fig. 7G**). Consistently, this population

exhibited higher IFN-γ protein expression than whole EMCD8 upon in-vitro stimulation, alongside

elevated PD-1 and TIGIT levels (**Fig. 7H** and **fig. S10H**), suggesting dysregulation of the IFN-γ axis

across multimodal layers. Accumulating evidence suggests that IFN-γ (type II IFN), primarily produced by

CD8$^+$ T cells, plays a pivotal role in SLE, alongside type I IFN (*69–71*). Our strategy successfully

pinpointed DP EMCD8 as the cellular origin of this key cytokine with a pro-inflammatory phenotype,

underscoring its role as a candidate pathogenic player in SLE.

Discussion

In this study, we employed an in-depth investigation strategy to pinpoint disease-relevant cell states and

molecules from large-scale single-cell datasets of SLE. Fine-grained clustering resolved 123 cell states

across 27 cell types, including previously uncharacterized populations distinctively associated with

clinical severity and treatment status. Subsequent multimodal characterization delineated cell-state-

specific immune signaling networks, transcriptional regulators, epigenetic features, surface proteins, and

TCR repertoires.

While recent studies suggest that granular cell states within each cell type can exhibit

continuous plasticity (*72*), our results demonstrate that a fine-grained cell-state definition is essential for

a deeper understanding of disease pathophysiology, facilitating: (i) the interpretation of cluster-free cell

abundance testing (**Fig. 3**), (ii) patient stratification with high clinical relevance based on cell-state

abundance (**Fig. 5**), (iii) the delineation of aberrant immune crosstalk (**Fig. 5**), and (iv) the functional

characterization of disease-relevant populations and the identification of their potential therapeutic

targets (**Fig. 6−7**). Since disease-relevant cell states are often low-frequency populations easily obscured

by coarse clustering, our approach is consistent with studies that successfully identified rare pathogenic

subsets (*9*, *10*, *13*). Leveraging an extensive literature review (*9–13*, *19–39*) and rigorous validation, we

reconstructed 123 cell states with distinct immunological features from the largest publicly available

single-cell PBMC dataset (*5*) (~1.5 million cells). While recent studies on inflamed tissues in autoimmune

diseases have identified critical parenchymal subpopulations (*13*, *73*), the resolution of immune cells in

those studies remained limited (~60 immune cell states from 0.2~0.4 million cells). To benefit the

research community, we have released LuPIN, a high-resolution immune cell reference panel integrated

with a comprehensive atlas of immune signaling networks and multimodal features. Researchers can

map their datasets onto LuPIN for precise cell-state annotation and mechanistic inference, providing

opportunities to uncover new pathophysiology across multiple diseases.

We also proposed a statistical framework to dissect quantitative and qualitative changes linked

to disease within each cell type. Although methods like miloDE (*74*) infer differential expression unique to

transcriptional neighborhoods, they were not designed to jointly dissect abundance shifts and per-cell

gene dysregulation. Distinguishing these scenarios is a critical step in identifying therapeutic targets, as

each may play distinct roles in pathophysiology. The significant enrichment of SLE risk variants within

expanded cell states further suggests that these populations are pivotal for variant-to-function research

(*75*). While our model demonstrated reasonable robustness and reproducibility, its performance relies on

the latent space defined by current sparse single-cell gene expression. The distinction between

abundance and dysregulated signatures may be further refined as single-cell technologies advance.

The identification of cell-state-specific regulators and surface proteins has significant

implications for drug discovery. For instance, given that FOXO1 has been implicated in the efficacy of

cancer immunotherapy (*24, 25*), modulating FOXO1 could represent a novel treatment option for

autoimmunity by targeting aberrant "effector-primed" CD4[+] T cell states in SLE. Regarding surface

proteins, current therapeutic agents often target ubiquitous markers (e.g., CD19 and CD20 in B cells)

(*41*, *42*), which can deplete normal cells alongside autoreactive ones. Given the recent development of

bispecific antibodies (*76*), narrowing the target to disease-relevant cell states based on specific markers

could provide more precise and efficient treatment options.

This study has some limitations. First, we have yet to identify the optimal combination of surface

markers to purify $FOXO1^+ARHGAP15^+$ T cells; more comprehensive panels may address this in the

future. Second, extremely rare cell states within minor cell types could not be included in several

downstream analyses due to limited cell numbers (**Fig. 3−4**); improvements in throughput will likely

uncover their contribution to pathogenesis. Third, as this study focused on PBMCs, key tissue-resident

populations may have been overlooked; thus, our approach should be extended to inflamed tissues

across various diseases. Fourth, our study design lacked in-vivo functional validation of the identified cell

states. Collectively, our findings contribute to a deeper understanding of SLE pathophysiology and

facilitate the discovery of novel therapeutic targets.

Acknowledgments

490 The super-computing resource was provided by Human Genome Center, Institute of Medical Sciences,

491 The University of Tokyo. We appreciate the generosity of the donors for a RIKEN solicited donation

492 project, Support for SLE research and development. We appreciate the RIKEN-IMS Genome Platform

493 for its help in the sequencing experiments. We thank AIDA consortium and RIKEN-IMS Laboratory for

494 Gene Structure and Regulation for helpful feedback.

Funding

This study was supported by the Japan Agency for Medical Research and Development (AMED)

(JP22tm0424223, JP22ek0410099, JP23tm0524005, JP25ek0410139, JP22ama121015 and

JP223fa627010, JP22gm1810002, JP22gm1810005), JSPS Grants-in-Aid for Scientific Research

(JP21K20647, JP22H03114, JP23KJ2185, JP23K15361, JP25K19621), the Uehara Memorial

Foundation, the Nakajima Foundation, Japan Intractable Diseases Research Foundation, GSK Japan

Research Grant, Daiichi Sankyo Foundation of Life Science, Mochida Memorial Foundation for Medical

and Pharmaceutical Research, Takeda COCKPI-T Funding, Kanzawa Medical Research Foundation,

SENSHIN Medical Research Foundation, Japan Research Foundation for Clinical Pharmacology, the

Japan Rheumatism Foundation, the Ichiro Kanehara Foundation, Keio University Academic

Development Funds, and Keio University Program for the Advancement of Next Generation Research

Projects.

Authors contributions

M.N. and K.I. designed the study. M.N., K.I., and M.K. wrote the manuscript with critical inputs from T.N.,

Y.T., and K.Y. M.N. conducted analyses with support from K.I., M.K., K.A., H.I., T.N., J.I., H.T., and H.H.

M.K. M.N., and K.A. conducted experiments with support from T. A., T. K., S. N., R. B., Y.M., B.N., A.S.,

and K.Y. T.K., S.K., E.K., Y.F., Y.T., and Y.M. contributed to sample collection. X.Z., S.B. and C.T.

managed EAS SLE-GWAS data. All authors contributed to the final manuscript.

Competing interests

RIKEN has filed a patent application related to UNIChro-seq (WO/2024/242037). K.I, M.K, and T.A are

inventors on this patent. S.K. has received speaking fees from Eli Lilly, GlaxoSmithKline, Bristol-Myers,

Abbvie, Eisai, Pfizer, Astra-Zeneca, and research grants from Daiichi-Sankyo, Abbvie, Boehringer

Ingelheim, and Astellas.

Data, code, and materials availability

Single-cell count matrices of gene expression, surface protein, and TCR features in the CITE-seq

dataset will be available at the National Bioscience Database Center (NBDC) Human Database

(https://humandbs.dbcls.jp/en/) as of the date of publication. Public data used in this study are available

at https://cellxgene.cziscience.com/collections. Reference panels for each lineage and each cell type are

available on GitHub (https://github.com/MasahiroNakano-hub/LuPIN). Original codes used in this study

are available on GitHub (https://github.com/MasahiroNakano-hub/LuPIN).

Fig. 1. Overview of this study. (left) We extensively investigated multimodal single-cell data of SLE of ~2.1 million cells from 346 donors to pinpoint granular disease-relevant cell states. **(middle)** Using fine-grained clustering, we defined 123 cell states across 27 immune cell types, identifying previously uncharacterized cell states associated with clinical severity and treatment status. **(right)** Our statistical framework indicated that key immunological pathways and SLE risk variants were linked to cell-state compositional shifts. Furthermore, these newly identified cell populations were predicted to engage in immune crosstalk with established pathogenic cell states, alongside in-depth multimodal characterization of their molecular profiles. CITE-seq, cellular indexing of transcriptomes and epitopes by sequencing; TCR, T cell receptor; EMCD8, effector memory CD8⁺ T cells; TF, transcription factor; GWAS, genome-wide association studies; eRNA, enhancer RNA; UNIChro-seq, unique molecular identifier counting of regional chromatin accessibility with sequencing; FCM, flowcytometry; FACS, fluorescence-activated cell sorting.

Fig. 2. Definition of 123 robust cell states using fine-grained clustering. (A-F) UMAP plots showing 27 cell types and 123 cell states identified by fine-grained clustering in **(A)** CD4+ T cell (CD4 T), **(B)** CD8+ and unconventional T cell (CD8/other T), **(C)** natural killer cell (NK), **(D)** B cell, **(E)** monocyte (Mono), and **(F)** dendritic cell (DC) lineages (dataset 1; *N* = 269). The hematopoietic stem and progenitor cell (HSPC) lineage is omitted as it comprises a single cell type and cell state. For each panel, large lineage-level UMAPs are colored by cell types, while small cell-type-level UMAPs are colored and numbered by cell states. Cell-state annotations are listed to the right of each panel. See **table S2** for cell type abbreviations. **(G)** Bar plots showing the number of cell states replicated in **(left)** leave-one-batch-out analysis of dataset 1 and **(right)** independent clustering of dataset 2 and 3.

Fig. 3. Mapping cell state compositional changes in SLE. (A) Distributions of neighborhood correlations for each cell state in disease-state (*n* = 137) or disease-activity (*n* = 61) comparisons (dataset 1, cell-type-level-CNA). 29 cell states with > 50% of neighborhoods positively associated with phenotypes (FDR < 0.05) are shown (See **fig. S5A** for details). (B) UMAP plots of representative cell types, colored by neighborhood correlations. Cell states are highlighted if > 50% (**) or > 25% (*) of their neighborhoods are associated with phenotypes. (C) Scatter plot of median neighborhood correlation (disease-state CNA) for each cell state vs. cell-state proportion within cell type. Shaded regions represent 95% confidence intervals. (D) Comparison of median neighborhood correlation between disease-state (x-axis) and disease-activity (y-axis) CNA. Cell states relevant to each panel title are highlighted in color. (E) Heatmap of median neighborhood correlation for low (LDA) / moderate (MDA) / high disease activity (HDA) vs. Inactive SLE (*n* = 58/17/21/40, respectively). 32 disease-activity-associated cell states are shown. (F) Comparison of median neighborhood correlation for each cell state between dataset 1 (x-axis) and dataset 2 (y-axis; *N* = 58). In (C) and (F), Pearson's *r* and *P* values are indicated.

Fig. 4. Statistical framework to dissect quantitative and qualitative changes within cell types. (A) Schematic representation of two distinct scenarios (quantitative and qualitative changes) driving cell-type-level pseudobulk DEGs. **(B)** Schematic of the framework to dissect quantitative and qualitative changes. **(C)** Representative examples of abundance (*OAS1*) and dysregulated (*IFIH1*) signature genes. UMAPs are colored by expression density. Model statistics for both genes are provided below. **(D)** Heatmap of abundance and dysregulated term Z scores across 18 major cell types for 100 IFN-response genes (dataset 1; disease-state comparison; n = 137). Gene order is based on hierarchical clustering of Z scores. **(E)** Bar plots showing the enrichment of representative pathways for each signature across cell types. *P*, *P* values from one-sided Fisher's exact tests. **(F)** Comparison of SLE risk variant enrichment around abundance vs. dysregulated signature genes for each cell type using stratified LD score regression (S-LDSC).

Fig. 5. Characterization of aberrant immune networks between disease-relevant cell states. (A) Heatmap of cell-state proportions across 18 major cell types for all samples (dataset 1, *N* = 269). Sample and cell-state orders are based on hierarchical clustering of cell-state proportions. **(B)** Bar plot showing the frequencies of disease-activity categories for each sample group (SG). *P*, *P* value from two-sided Fisher's exact test. LDA/MDA/HDA; low/moderate/high disease activity. **(C)** Box plots comparing cell-state proportions between "Flare" and "Treated" status in **(left)** Tph and **(right)** DP EMCD8. Samples from the same donor are connected by lines. *P*, *P* values from MASC (*n* = 10 × 2). **(D)** Bar plot showing the number of potential cell-state-specific ligand-receptor (LR) interactions within each CG. **(E)** Schematic of cell-state-specific LR networks across known pathogenic and newly identified population in cell-state group (CG) 2. **(F-G)** Heatmaps of LR scores for **(F)** IFNG–IFNGR1/2 signaling from EMCD8 to NaiveB cell states and **(G)** CD70–CD27 signaling from EMCD8 to EMCD4 cell states.

Fig. 6. *FOXO1* is a candidate driver of the pathogenic CD4+ T cell state gene program. (A) Bar plots showing the enrichment of representative pathways for upregulated signature genes for NaiveCD4 and CMCD4 cell states. *P*, *P* values from one-sided Fisher's exact tests. **(B)** Dot plot showing scaled expression levels of ligand and receptor genes across NaiveCD4 and CMCD4 cell states. Color intensity represents the proportion of cells expressing each gene. Corresponding interaction partners are indicated on the right. **(C)** Bar plot showing the percentage of two TCR V gene usages skewed in NaiveCD4-5. *P*, *P* values from two-sided Fisher's exact tests. **(D)** Dot plot showing the logFC of top 15 upregulated surface markers for NaiveCD4-5 and CMCD4-4. *P, P* values from Seurat FindMarkers. **(E)** Heatmap showing the logFC of GWAS-prioritized genes across NaiveCD4 and CMCD4 cell states. Up to five genes with the highest logFC per cell state are shown. **(F)** Heatmap showing - log10(enrichment *P*) values of each TF for NaiveCD4-5 or CMCD4-4 signature genes based on (i) pathway/TF enrichment analyses, TF activity inference using (ii) decoupleR and (iii) blood-derived ChIP-seq data, and (iv) an arrayed CRISPR-KO experiment. *P*, *P* values from one-sided Fisher's exact tests. **(G)** UMAP plots of whole CD4+ T cells, colored by **(left)** focused cell states and other cell types, **(upper right)** *FOXO1* expression, and **(lower right)** pseudotime from trajectory analysis. Edges represent trajectory graph.

Fig. 7. DP EMCD8 cells as a candidate pathogenic player in SLE. (A-B) UMAP plots of **(top)** EMCD8 and **(bottom)** EMCD4, colored by **(A)** oxidative phosphorylation and **(B)** exhaustion signature scores. **(C)** Bar plots showing **(top)** the absolute number of EMCD8-5 cells per donor and **(bottom)** the proportion of TCR clonotypes categorized into five size groups within EMCD8-5 (dataset 3; $N = 32$; 23 donors with >10 cells are shown). **(D)** Dot plot showing scaled expression levels of ligand and receptor genes across EMCD8 cell states. Color intensity represents the proportion of cells expressing each gene. Corresponding interaction partners are indicated on the right. **(E)** Volcano plot showing differential expression of surface proteins between EMCD8-5 vs. other EMCD8 cells. P, P values from Seurat FindMarkers. **(F)** Experimental design for flowcytometry (FCM) and FACS of HLA-DR$^+$CD27$^+$CD38$^+$CD28$^+$ EMCD8 cells. **(G)** ATAC-seq and UNIChro-seq tracks around *IFNG* locus in whole EMCD8 and HLA-DR$^+$CD27$^+$CD38$^+$CD28$^+$ cells from an SLE donor. P, logistic regression P values. **(H) (left)** Boxplots comparing the mean fluorescence intensity (MFI) of PD-1 and IFN-γ between HLA-DR$^+$CD27$^+$CD38$^+$CD28$^+$ and whole EMCD8 cells. Samples from the same donor are connected by lines. P, P values from linear mixed models. **(right)** Representative histogram comparing IFN-γ fluorescence intensity between HLA-DR$^+$CD27$^+$CD38$^+$CD28$^+$ and whole EMCD8 cells from an SLE donor.

Fig. S1. Quality control of datasets. (A) Density plot showing cell distributions based on the number of detected genes and the percentage of mitochondrial UMIs (%Mito) in dataset 1 (N = 269). **(B)** UMAP plot showing PBMCs in dataset 3 (N = 32), colored by Souporcell-based singlet/doublet/unassigned classification. **(C)** Scatter plot showing donor-level pseudobulk expression of XIST and Y-chromosome-related genes in dataset 1, colored by metadata-derived sex information. **(D)-(F)** Flowchart illustrating the strategy to partition PBMCs into seven lineages in dataset 1. In **(F)**, dot plots display scaled expression levels of lineage or subcluster marker genes, with color intensity indicating the proportion of expressing cells. **(G)** Boxplot comparing Scrublet-derived doublet scores between singlets and doublets identified in **(E)** and **(F)**. **(H)** Bar plots showing the number of cells retained after each QC step in datasets 1, 2, and 3.

Fig. S2. Definition of 123 robust cell states using fine-grained clustering. (A-F) Flowcharts illustrating the hierarchical separation of lineage cells into cell types in **(A)** CD4+ T cell (CD4 T), **(B)** CD8+ and unconventional T cell (CD8/other T), **(C)** natural killer cell (NK), **(D)** B cell, **(E)** monocyte (Mono), and **(F)** dendritic cell (DC) lineages (dataset 1; $N = 269$). The hematopoietic stem and progenitor cell (HSPC) lineage is omitted as it contains only a single cell type. Dot plots display scaled expression levels of cell-type marker genes, with color intensity representing the proportion of cells expressing each gene. **(G)** Box plot showing the concordance rates of cell-type-level annotation in the original and the current study with the predicted cell-type labels in the reference mapping onto the independent PBMC dataset. **(H) (left)** Distributions of neighborhood correlations for each cell type in disease-state ($n = 137$) or disease-activity ($n = 61$) comparisons (dataset 1, lineage-level-CNA). **(right)** UMAP plots in representative lineages. Cell types are highlighted if > 50% of neighborhoods are associated with phenotypes. See **table S2** for cell type abbreviations.

Fig. S3. Marker gene expression across 123 cell states. (A–F) Dot plots showing scaled expression levels of cell-type and cell-state marker genes for each cell state in **(A)** CD4+ T cell (CD4 T), **(B)** CD8+ and unconventional T cell (CD8/other T), **(C)** natural killer cell (NK), **(D)** B cell, **(E)** monocyte (Mono), and **(F)** dendritic cell (DC) lineages (dataset 1; *N* = 269). Color intensity represents the proportion of cells expressing each gene.

Fig. S4. Robustness and replicability of 123 cell states. (A) UMAP plots of representative cell type (EMCD8), colored by cell states for each leave-one-batch-out condition. **(B)** Bar plots showing the number of cell states replicated in independent clustering of dataset 2 and/or 3. **(C)** UMAP plots of representative cell types in the CD8/other T lineage, colored by cell states in independent clustering of dataset 2 and 3. Cell states not replicated in each condition are listed to the left, and cell-state annotations are listed to the right. **(D)** Dot plots showing scaled expression levels of cell-type and cell-state marker genes for each cell state in the CD8/other T lineage from dataset 2 and 3. Color intensity represents the proportion of cells expressing each gene. **(E)** Heatmap showing the number of overlapping genes among the top 100 upregulated marker genes for each cell state in EMCD8.

Fig. S5. Mapping cell state compositional changes in SLE. (A) Distributions of neighborhood correlations for each cell state in disease-state (*n* = 137) or disease-activity (*n* = 61) comparisons (dataset 1, cell-type-level-CNA). **(B)** Scatter plot of median neighborhood correlation (disease-activity CNA) for each cell state vs. cell-state proportion within cell type. Shaded regions represent 95% confidence intervals. **(C)** Distributions of neighborhood correlations in Asian vs. EUR SLE comparisons. 15 cell states with > 50% of neighborhoods associated with phenotype (FDR < 0.05) are shown. **(D)** Histograms showing the number of cells per SLE sample for representative cell states expanded in SLE. **(E)** Scatter plot of median neighborhood correlation in CNA vs. effect size in MASC for each cell state in disease-state and activity comparisons. **(F)** Scatter plot of median neighborhood correlation of cell-type-level CNA in European vs. in Asian. In **(B)**, **(E)** and **(F)**, Pearson's *r* and *P* values are indicated.

Fig. S6. Statistical framework to dissect quantitative and qualitative changes within cell types. (A) Relative proportion of variance explained by abundance and dysregulated terms for the top 1,000 pseudobulk DEGs across all cell types and comparisons. **(B)** Quantile-quantile plots of observed vs. expected model *P* values for abundance and dysregulated terms across all cell types and comparisons. **(C)** Bar plot showing Pearson's *r* of model effect sizes between the original vs. each leave-one-batch-out analysis for representative cell types. **(D)** Comparison of model effect sizes across genes between dataset 1 and 2. *P*, *P* values from one-sided binomial tests. **(E)** Heatmap of abundance and dysregulated term Z scores across 18 major cell types for 100 IFN-response genes (disease-activity comparison; *n* = 61). Gene order is based on hierarchical clustering of Z scores. **(F)** Bar plots showing the enrichment of representative pathways for each signature across cell types (disease-activity comparison). *P*, *P* values from one-sided Fisher's exact tests. **(G)** Comparison of normalized coefficients in stratified LD score regression (S-LDSC) between EAS vs. EUR SLE-GWAS. Each dot represents a signature for a given cell type. Error bars represent 2×standard errors. Pearson's *r* and *P* values are indicated. **(H)** Comparison of SLE risk variant enrichment around abundance signature genes with positive vs. negative model statistics for each cell type.

Fig. S7. Characterization of aberrant immune networks between disease-relevant cell states. (A) Bar plot showing the frequencies of SLE samples with **(left)** mucocutaneous or renal activity and **(right)** hydroxychloroquine (HCQ) or mycofenolate mofetil (MMF) treatment for each sample group (SG). *P*, *P* values from two-sided Fisher's exact tests. **(B)** Box plots comparing cell-state proportions between "Flare" and "Treated" status in **(top)** CG2 and **(bottom)** CG3 cell states. Samples from the same donor are connected by lines. *P*, *P* values from MASC (*n* = 10 × 2). **(C-E)** Heatmaps of LR scores for **(C)** ADA–DPP4 signaling from EMCD8 to NaiveCD4 cell states, **(D)** CALM1–PTPRA signaling from EMCD8 to CMCD4 cell states, and **(E)** COPA–CD74 signaling from CMCD4 to EMCD8 cell states.

Fig. S8. Multimodal molecular profiling across fine-grained cell states. (A) Schematic of the integration of multimodal dataset 3 (query; *N* = 32) into large scRNA-seq dataset 1 (reference; *N* = 269) through reference mapping. **(B)** Bar plot showing the number of cell-state-specific upregulated surface marker proteins. Colors represent cell-state IDs (e.g., NaiveCD4-0). **(C)** PCA plot of TCR V gene usage frequencies for each cell state, colored by cell type. **(D)** Bar plot showing the number of cell-state-specific (overrepresented) TCR V genes. **(E)** Bar plot showing the enrichment of called transcribed enhancers across all cell types for "PBMC enhancer" compared with **(left)** "Brain enhancer" and **(right)** "PBMC heterochromatin" in Roadmap Epigenomics project annotations.

Fig. S9. *FOXO1* is a candidate driver of the pathogenic CD4+ T cell state gene program. (A) Representative examples of cell-state-specific transcribed enhancers near upregulated marker genes in NaiveCD4-5. **(B)** Dot plot showing the logFC of top 15 upregulated surface markers for NaiveCD4-4. *P*, *P* values from Seurat FindMarkers. **(C)** Experiment design of the arrayed CRISPR-KO experiment in primary CD4$^+$ T cells. **(D)** Heatmap showing -log$_{10}$(enrichment *P*) values of *ARID5B* for NaiveCD4-4 signature genes based on (i) pathway/TF enrichment analyses, TF activity inference using (ii) decoupleR and (iii) blood-derived ChIP-seq data, and (iv) a CRISPR-KO experiment. *P*, *P* values from one-sided Fisher's exact tests.

Fig. S10. DP EMCD8 cells as a candidate pathogenic player in SLE. (A) Heatmap showing the logFC of GWAS-prioritized genes across EMCD8 cell states. Up to five genes with the highest logFC per cell state are shown. **(B)** Bar plots showing the percentage of **(left)** a representative clonotype shared across EMCD8 cell states and **(right)** a clonotype specific to EMCD8-5. *P*, *P* values from two-sided Fisher's exact tests. **(C)** Bar plots showing the percentage of **(left)** three TRAV and **(right)** six TRBV gene usages positively skewed in EMCD8-5. **(D)** Bar plot showing the percentage of EMCD8-5 cells within each in-silico gate using dataset 3. **(E)** Gating strategy for defining HLA-DR⁺CD27⁺CD38⁺CD28⁺ EMCD8 cells in flowcytometry (FCM) and FACS. **(F)** Scatter plot comparing the scRNA-seq dataset 1 logFC (EMCD8-5 vs. other EMCD8 cells) and bulk RNA-seq logFC (HLA-DR⁺CD27⁺CD38⁺CD28⁺ vs. whole EMCD8 cells). **(G)** TF motif logos enriched for differentially accessible peaks in HLA-DR⁺CD27⁺CD38⁺CD28⁺ cells compared with whole EMCD8 cells. *P*, enrichment *P* values from HOMER. **(H) (left)** Boxplots comparing the mean fluorescence intensities (MFI) of TIGIT between HLA-DR⁺CD27⁺CD38⁺CD28⁺ and whole EMCD8 cells. Samples from the same donor are connected by lines. *P*, *P* values from linear mixed models.

Methods

Subjects

In this study, we utilized two publicly available 3'-end scRNA-seq datasets (datasets 1 and 2). For

dataset 1, FASTQ files were downloaded from CELLxGENE (*5*) and raw genotype data were obtained

from dbGaP (phs002812.v1.p1) with authorized access. Samples were recruited from the California

Lupus Epidemiology Study (CLUES) cohort and the ImmVar Consortium. For dataset 2, count data were

downloaded from dbGaP (phs002048.v1.p1) (*4*). For our newly generated 5'-end multimodal dataset

(CITE-seq, TCR-seq, and enhancer RNA profiling; dataset 3) and additional FCM/FACS experiments,

patients were recruited at Okayama University Hospital and Hospital of the University of the

Occupational and Environmental Health under approvals by the Ethics Committees of RIKEN Center for

Integrative Medical Sciences (RIKEN−Y-2025-006, 008). Written informed consent was obtained from all

participants.

Across all datasets, SLE disease activity was categorized into four levels based on the SLE

Disease Activity Index (SLEDAI) (*77*): i) inactive (SLEDAI = 0), ii) low (LDA; $1 \leq$ SLEDAI ≤ 4), iii)

moderate (MDA; $5 \leq$ SLEDAI ≤ 9), and iv) high disease activity (HDA; SLEDAI ≥ 10) (**table S1**). In

dataset 1, "flare" status was categorized into HDA.

Sample processing and quality control

For dataset 3, single-cell suspensions of PBMCs from 32 SLE donors were processed using the 10x

Genomics Chromium system. Gene expression (GEX), surface protein (ADT), and TCR (VDJ) libraries

were prepared using the Chromium Next GEM Single Cell 5' Kit v2 and TotalSeq™-C Human Universal

Cocktail v1.0 (Biolegend, 399905). Eight libraries (16 samples pooled per library; four libraries per batch)

were sequenced on an Illumina NovaSeq 6000 (150-bp paired-end).

FASTQ files for datasets 1 and 3 were processed using Cell Ranger v7.0.1. We excluded low-

quality cells based on their distributions: (i) total unique molecular identifiers (UMIs) (< 500), (ii) number

of genes (nGene), (iii) percentage of mitochondrial genes (% MT), or (iv) percentage of ribosomal protein

genes (% Ribo) (dataset 1: nGene < 300, %MT > 7.5%, %Ribo < 8.5%; dataset 2: nGene < 300, %MT >

8.5%, %Ribo < 8.5%; dataset 3: nGene < 500, %MT > 8.5%, %Ribo < 6%; **fig. S1A**). Cells with high

percentage of hemoglobin genes (> 0.9%; *HBB*, *HBA1*, *HBA2*, *HBD*) or platelet (> 0.6%; *PF4*, *PPBP*,

CAVIN2, *GNG11*) signatures were also removed.

For dataset 1, samples were demultiplexed using Demuxalot (*78*) based on imputed genotypes

(SHAPEIT2, Minimac3 r^2 > 0.95) or Souporcell (*79*) for samples lacking reference genotypes. For dataset

3, Souporcell was used with 1000 Genomes Project common variants (minor allele frequency > 1%),

followed by multiplex PCR and MiSeq sequencing of 20 target variants to link genotype clusters to

specific individuals. Putative doublets from different genotypes were excluded (**fig. S1B**). In dataset 1,

we excluded two samples with < 200 cells recovered and three samples showing sex discordance

between metadata and pseudobulk expressions of *XIST* and Y chromosome genes (**fig. S1C**).

PBMCs were partitioned into seven lineages (CD4 T, CD8/other T, NK, B, Mono, DC, and

HSPC) through iterative Louvain clustering (**fig. S1D−F**). Briefly, gene expression was normalized via

SCTransform (Seurat v5.1.0) (*80*) and integrated using 3,000 highly variable genes (HVGs), followed by

batch correction via Harmony (group.by.vars = "library", "sample") (*81*). Mitochondrial, Y chromosome,

hemoglobin, platelet, and cell-cycle genes were excluded from HVGs. We used 30 Harmony-corrected

PCs for UMAP visualization and shared nearest neighbor (SNN) graph construction. After initial

separation into four major populations (T/NK, B, Myeloid, and HSPC), we performed recursive sub-

clustering to maximize lineage purity based on marker gene expression (**fig. S1D−F**). Hyperparameters

for each step are summarized in **table S3**. Sub-clusters representing remaining doublets with multiple

lineage markers (**fig. S1F**) and higher Scrublet doublet scores (*82*) (**fig. S1G**) were excluded. After multi-

step QC, we retained 1,456,041 (dataset 1), 344,235 (dataset 2), and 268,783 (dataset 3) high-quality

PBMCs (**fig. S1H**).

Definition of 123 fine-grained cell states

To identify granular cell states, we employed a three-layered iterative clustering strategy in dataset 1:

lineage (layer 1), cell type (layer 2), and cell state (layer 3). Following initial lineage separation, we

performed SCTransform normalization, HVG selection, PCA, Harmony batch correction, and Louvain

clustering for each lineage using optimized parameters (**fig. S2A−F** and **table S3**). Sub-clustering was

performed iteratively to ensure cell-type purity based on canonical marker expression. In a few instances

(nine clusters), biologically redundant clusters were merged to define a final set of 123 cell states. To

validate annotation accuracy, we performed reference mapping of dataset 1 onto an independent PBMC

reference (Azimuth) (*19*) and compared concordance rates between our fine-grained annotations and the

original labels (**fig. S2G**).

Cell-state-specific marker genes were identified using Seurat's FindMarkers (v4.3.0) (min.pct =

0.05), comparing each state against all others within the same cell type (**fig. S3** and **table S4**).

Upregulated markers (FDR < 0.05 and logFC > 0.1) were used for downstream pathway/TF enrichment

analyses. To provide more intuitive differential expression values, we also reported logFC calculated with

pseudocount.use = 0 in addition to the default Seurat values (**table S4**).

The robustness of the 123 cell states was assessed via leave-one-batch-out analysis, where

each of the 23 experimental batches was systematically excluded, and the entire pipeline—from

normalization to clustering—was re-executed. We titrated the resolution parameter (in increments of

0.05) to confirm that the original clusters were consistently replicated (**fig. S4A** and **table S5**).

Furthermore, independent replication analyses were performed using dataset 2 and 3, where the entire

pipeline was re-run from scratch for each dataset (**fig. S4B−C**). We confirmed that the resulting clusters

exhibited marker gene profiles comparable to those in dataset 1 (**fig. S4D** and **table S5**).

Co-varying neighborhood analysis

To identify single-cell-resolution transcriptional neighborhoods associated with clinical phenotypes, we

applied co-varying neighborhood analysis (CNA v0.1.6) (*15*), to the 18 major cell types (> 5,000 cells

each; **Fig. 3**) and six cell lineages (**fig. S2H**) in dataset 1. Following our established framework (*6*), we

defined two comparison axes: (i) disease-state (97 healthy controls vs. 40 inactive SLE) and (ii) disease-

activity (40 inactive vs. 21 high-activity [HDA] SLE).

Harmony-corrected PCs were used as inputs to infer the neighborhood abundance matrix

(NAM), followed by PCA to obtain NAM-PCs. We tested the associations between phenotype labels and

NAM-PCs using linear models, adjusting for age, sex, ancestry, and the number of cells per cell type.

Given the substantial correlation between disease activity and age ($r = -0.60$) in dataset 1, age was

excluded from covariates in disease-activity comparisons. For cell-type-level CNAs, significance was

defined as a global permutation $P < 0.05/36$ (Bonferroni correction). CNA-derived neighborhood

correlations (positive for expansion, negative for depletion) were aggregated for each cell state (**Fig. 3**

and **table S8**). States were categorized as showing "remarkable expansion/depletion" if > 50% of

constituent neighborhoods were associated with the phenotype at an empirical FDR < 0.05, and

"moderate" if > 25%. We additionally performed CNA for (1) inactive vs. low-activity (LDA), (2) inactive

vs. moderate-activity (MDA), and (3) Asian vs. European SLE (**Fig. 3E** and **fig. S5C**). MASC (*11*) was

also applied to test cell-state abundance associations, adjusting for age, sex, and ancestry (**fig. S5E**).

621   For replication, CNA was applied to dataset 2 (**Fig. 3F**). Cells were projected into the dataset 1

latent space using Symphony (*53*), and Symphony-corrected PCs served as inputs. To ensure stable

CNA performance (requiring ~30 samples) (*15*), we compared 14 SLE (SLEDAI <= 2) vs. 17 healthy

donors for disease-state, and 15 SLE (SLEDAI >= 6) vs. 14 SLE (SLEDAI <= 2) for disease-activity

comparison.

Statistical framework to dissect quantitative and qualitative changes

To decompose cell-type-level gene expression into quantitative and qualitative changes, we introduced a

single-cell-level generalized linear mixed effect model (GLMM, **Fig. 4B** and **table S9**). For each gene,

the model was fitted using a negative binomial distribution:

$$\log(E_i) = \beta_{abund}X_{i,abund} + \beta_{dysreg}X_{d,dysreg} + \beta_{age}X_{d,age} + \beta_{sex}X_{d,sex}$$

$$+\beta_{ancestry}X_{d,ancestry} + \beta_{nUMI}X_{i,nUMI} + \beta_{MT}X_{i,MT} + (\phi_d|d) + (\kappa_b|b) + \theta$$

where *E* is the expression of the gene in cell *i*, *θ* is an intercept, and *β* coefficients represent fixed effects

for cell *i*, donor *d*, or batch *b*. Donor and batch were modeled as random intercepts. The abundance

signature term ($X_{i,abund}$) represents neighborhood correlation values from cell-type-level CNA, while the

dysregulated signature term ($X_{d,dysreg}$) represents phenotype information (e.g., healthy vs. inactive SLE).

The model was applied to genes passing low-expression filters (>=3 counts in >= 10% of samples)

across 18 major cell types (> 5,000 cells) for disease-state and activity comparisons in dataset 1, using

lme4 (*83*). Significance was determined via likelihood ratio tests (LRT) comparing models with and

without the respective terms. ImmVar samples were excluded from disease-state comparisons. To

quantify the contribution of each term to pseudobulk expression, we calculated Nagelkerke R^2 values for

the top 1,000 DEGs (identified via edgeR (*84*)) per cell type (**fig. S6A**).

643   Type I error calibration was assessed via permutation tests (randomly shuffling neighborhood

correlations or phenotype labels; **fig. S6B**). Robustness was further validated through leave-one-batch-

out analysis (**fig. S6C**) and independent replication in dataset 2, with effect-size replicability tested via

one-sided binomial sign tests (**fig. S6D**). For 100 IFN response genes (IRGs) (*4*) (**table S11**), abundance

and dysregulated Z-scores were used for hierarchical clustering (Ward's method, Euclidean distance) to

visualize patterns across cell types (**Fig. 4C−D** and **fig. S6E**).

Pathway enrichment analysis and gene signature scores

To evaluate the enrichment of pathway annotations for abundance and dysregulated signature genes,

we performed over-representation analysis (ORA) using the top 1,000 genes (by LRT P-values) for each

signature term in each cell type. ORA was conducted using a one-sided Fisher's exact test via

clusterProfiler (*85*) (**Fig. 4E**, **fig. S6F**, and **table S12**), utilizing MSigDB Hallmark and KEGG pathway

collections. To capture cell-type-specific biology, the union of the top 1,000 LRT P-value genes across all

cell types was used as the background gene set.

657   For pathway analysis of cell-state-specific upregulated marker genes (FDR < 0.05, logFC > 0.1,

Seurat FindMarkers), ORA was performed similarly (**Fig. 6A**, **Fig. 7A**, and **table S7**). The background for

each cell type consisted of all genes passing the low-expression filter (>=3 counts in >= 10% of

samples). To calculate classical exhaustion signature scores in EMCD8 and EMCD4 populations, an

exhaustion-related gene set was curated from previous studies (*3*, *5*) (**table S20**). Gene expression was

log-normalized and scaled, and the signature score for each cell was defined as the sum of the scaled

expression values across the gene set (**Fig. 7B**).

Stratified LD score regression analysis

To evaluate the enrichment of SLE heritability (all risk variants irrespective of effect size) around

abundance and dysregulated signature genes, we performed stratified linkage disequilibrium score

regression (S-LDSC) (*40*) using the sc-linker pipeline (*86*) (**Fig. 4F** and **fig. S6H**). We first calculated

min-max-scaled gene scores based on the Z-scores of abundance and dysregulated terms for all tested

genes in each cell type. LD scores were then estimated for East Asian (EAS) and European (EUR)

populations using 1000 Genomes Project reference panels, with the scaled gene scores serving as

annotation weights. In S-LDSC, we examined the enrichment of SLE heritability for common variants

within 100-kb windows flanking the transcription start sites (TSS) of genes with positive Z-scores for both

signatures, adjusting for the baseline model annotations v1.2 (*40*). Two large-scale SLE-GWAS

summary statistics from EAS and EUR ancestries (*44*, *45*) were utilized. To account for differences in

GWAS heritability, regression coefficients were normalized by the mean per-SNP heritability as

previously described (*40*). Finally, a fixed-effect meta-analysis of the EAS and EUR results was

performed using the inverse-variance weighting method based on the normalized coefficients and their

standard errors (**fig. S6G**). We reported one-sided *P* values to test whether the regression coefficients

were significantly positive.

Localization of GWAS prioritized genes

To identify candidate causal genes for SLE-associated genetic variants ($P < 5 \times 10^{-8}$), we used the L2G

(locus-to-gene) scores from the Open Targets Platform (*43*). We defined 124 genes with an L2G score >

0.5 as SLE-GWAS prioritized genes (**table S10**). For each cell state, we identified the overlap between

these 124 prioritized genes and the cell-state-specific upregulated markers (FDR < 0.05, logFC > 0.2;

Seurat FindMarkers) (**Fig. 6E** and **fig. S10A**).

Sample group (SG) and cell-state group (CG) analysis

To investigate the clinical relevance of fine-grained cell states, we performed bi-hierarchical clustering

(Ward's method, Euclidean distance) of all samples in dataset 1 and cell states within 18 major cell types

based on their proportions (**Fig. 5A**). Sample groups (SGs) and cell-state groups (CGs) were defined

based on the resulting dendrograms. The frequencies of disease-activity categories, organ-specific

activity, and treatment agents (**table S1**) were compared across SGs (**Fig. 5B** and **fig. S7A**). For

longitudinal analysis in dataset 1, changes in cell-state proportions between paired "Flare" and "Treated"

samples (n = 10 donors) were tested using MASC (*11*) (**Fig. 5C** and **fig. S7B**).

Cell-cell interaction analysis

To infer the inflammatory networks across all cell states, we performed cell-cell interaction analysis using

LIANA (46) with patemeters method = c("logfc", "sca") and min_cells = 10. From the consensus set of

4,701 ligand–receptor (LR) interactions across all 123 × 123 cell-state pairs, we identified 1,451,770

potentially active signals (defined by sca.LRscore > 0.6, logfc.logfc_comb > -0.1, and expression

proportions of >=1% for ligands and >=5% for receptors). To minimize redundancy, for each unique LR

interaction within each of the 27 × 27 cell-type pairs, we prioritized the cell-state pair with the highest LR

score as the representative cell-state-specific interaction (118,629 signals; **Fig. 4D−G**, **Fig. 6B**, **Fig. 7D**,

and **fig. S7C−E**).

Reference mapping and integration in LuPIN

To integrate the three single-cell datasets into the LuPIN framework, we performed reference mapping

using Symphony (*53*), designating the large-scale dataset 1 as the reference and datasets 2 and 3 as

queries (**fig. S8A**). For dataset 1, lineage- and cell-type-level reference objects were constructed based

on the Harmony-integrated embeddings used in the fine-grained clustering. For the query datasets (2

and 3), PBMCs were first partitioned into seven lineages using the same pipeline as for dataset 1.

Following normalization and scaling, cells from each lineage were projected into the corresponding

lineage-level latent transcriptional space of dataset 1 (lineage-level reference mapping). Cell-type labels

were then assigned to query cells using k-nearest neighbor classifiers (k = 5) based on the dataset 1

reference. This procedure was repeated at the cell-type level (cell-type-level reference mapping) to

annotate each cell in datasets 2 and 3 with the best-predicted cell-state labels.

Surface protein analysis

Following reference mapping of the multimodal dataset 3, we investigated cell-state-specific surface

protein profiles. For dataset 3, 130 surface proteins and seven isotype controls were profiled using the

TotalSeq™-C Human Universal Cocktail v1.0 (Biolegend, 399905). Protein expression for each cell was

normalized using a centered-log ratio (CLR) transformation across all 130 proteins. Cell-state-specific

surface markers were identified using Seurat FindMarkers, comparing each cell state against all others

within the same cell type. Proteins with FDR <0.05 and logFC >0.2 were defined as cell-state-specific

upregulated surface markers (**fig. S8B** and **table S14**).

TCR repertoire analysis

In dataset 3, we utilized TCR UMI counts for cells meeting the following criteria: (i) passed GEX-based

QC; (ii) annotated as NaiveCD4, CMCD4, EMCD4, CTLCD4, Treg, NaiveCD8, CMCD8, or EMCD8 via

reference mapping; and (iii) passed Cell Ranger TCR quality filters. To identify cell-state-specific V gene

usage, frequencies were compared for each state against all others within the same cell type using two-

sided Fisher's exact tests (**Fig. 6C**, **fig. S8D**, **S10C**, and **table S15**). Among all 45 TRAV and 49 TRBV

genes, we tested V genes passing a low expression filter (>= 10 total counts per cell type).

TCR clonal expansion in EMCD8-5 (DP EMCD8) was assessed by quantifying clonotypes

across SLE donors (**Fig. 7C**). Clonotypes were defined based on the combination of the CDR3 amino

acid sequence and the V gene of the β-chain. To identify DP EMCD8-specific clonotypes, frequencies

were compared between states within each donor using two-sided Fisher's exact tests (**fig. S10B** and

table S21). Known cognate epitopes for each clonotype were identified using the VDJdb browser (**table**

S21).

Transcribed enhancer analysis

To detect transcribed enhancers in dataset 3, we employed the ReapTEC pipeline (*18*). 10x Genomics

libraries were sequenced on an Illumina NovaSeq 6000 with 150-bp paired-end reads for both Read 1

and Read 2, ensuring the capture of transcription start sites (TSS) within the 5' regions. Reads were

mapped using STARsolo to identify those containing unencoded G. After deduplication, reads were

assigned to specific cell types based on barcodes. TSS peaks were generated by merging TSSs within

10 bp, and bidirectional transcribed candidate enhancers were identified using the developers' scripts,

with promoter regions masked. To validate these enhancers, we tested their enrichment across

Roadmap Epigenomics core 15-state reference annotations (*57*) for PBMC-derived enhancers (6_EnhG

and 7_Enh in E062), compared with (i) PBMC heterochromatin (9_Het in E062) and (ii) brain

hippocampus middle enhancers (6_EnhG and 7_Enh in E071) using two-sided Fisher's exact tests (**fig.**

S8E).

Cell-state-specific transcribed enhancers were identified using an entropy-based specificity

score (*18*). We first filtered for enhancers with >=5 reads per cell type, aggregated reads across donors,

and calculated entropy and specificity scores for each enhancer within each cell type. Enhancers with a

specificity score > 0.7 were defined as cell-state-specific (**table S16**).

Trajectory analysis

To infer the developmental trajectory across the CD4$^+$ T cell lineage, we employed Monocle3 (*64*) with

default parameters (**Fig. 6G**). For pseudotime estimation, the root was defined as a cell within the

SOX4$^+$TOX$^+$ NaiveCD4-3 state. This selection was based on recent evidence that these cells represent

recent thymic emigrants (RTEs), the most immature population within the peripheral naive T cell

compartment (*22*).

Upstream TF activity inference

To identify transcriptional regulators driving disease-relevant CD4$^+$ T cell states, we integrated three

complementary approaches (**Fig. 6F** and **fig. S9D**): (i) over-representation analysis (ORA) of TF targets,

(ii) TF activity inference via decoupleR (*62*) and (iii) target enrichment analysis using ChIP-Atlas (*63*).

Candidates TFs were selected from those upregulated (logFC > 0.3) in NaiveCD4-5 (n = 22) and

NaiveCD4-4 (n = 1, *ARID5B*). ORA was conducted using a one-sided Fisher's exact test via

clusterProfiler (*85*) for MSigDB C3 and KEGG collections (**table S17**), using signature genes (FDR <

0.05, logFC > 0.1; Seurat FindMarkers) from NaiveCD4-4, NaiveCD4-5, and CMCD4-4. For decoupleR,

TF activities were inferred using the ulm method based on DoRothEA regulons and FindMarkers logFC

values. For the ChIP-Atlas-based analysis, we utilized ChIP-seq datasets restricted to blood tissues

(**table S17**). Promoter regions were defined as -5kb to +2kb from the TSS. TF-target gene annotations

were constructed by intersecting ChIP-seq peaks with these promoter regions. Enrichment of NaiveCD4-

4/5, and CMCD4-4 signature genes within these TF-target sets was evaluated using one-sided Fisher's

exact tests.

Arrayed CRISPR-KO experiment

To validate the identified transcriptional regulators, we performed an arrayed CRISPR knock-out (KO)

experiment targeting 22 candidate TFs (**Upstream TF activity inference**; **fig. S9C**). For each TF, two

independent single guide RNAs (sgRNAs) (*87*) were synthesized (IDT; **table S18**) and complexed with

recombinant Cas9 (IDT, 1081058) to form ribonucleoprotein (RNP) complexes. Two control samples

(four sgRNAs) were prepared for each donor. Editing and frameshift efficiency for each sgRNA was

confirmed via Sanger sequencing (on median 79.7% and 68.7%, respectively; **table S18**).

Primary CD4$^+$ T cells from three healthy donors were isolated from fresh PBMCs using CD4$^+$ T

Cell Isolation Kit (Miltenyi, 130-096-533), expanded for 2 days using Dynabeads Human T-Activator

CD3/CD28 (Thermo, DB11132) in complete CTS OpTmizer T-Cell Expansion SFM (Thermo Fisher

Scientific, A1048501) supplemented with IL-2 (Peprotech, 200-02), IL-7 (Peprotech, 200-07), and IL-15

(Peprotech, 200-15). Seven days after activation, cells were electroporated using the 4D-Nucleofector

(Lonza, program EH-140) with 1×10^6 cells per reaction and subsequently cultured in triplicate. Five

days post-transfection, total RNA was isolated using MagMAX-96 Total RNA Isolation Kit (Thermo Fisher

Scientific, AM1830) and libraries were prepared using the RamDA-seq method (*88*). Sequencing was

performed on an Illumina NovaSeq 6000 (150bp paired-end).

For RNA-seq analysis, adaptors were trimmed using Cutadapt and reads with < 50 bp were

removed. Reads were mapped to GRCh38 using STAR (*89*) and quantified via HTSeq in unstranded

mode. After excluding three low-quality samples (unique reads < 4 × 10^6), TMM-normalized counts (*84*)

were fitted to a negative binomial GLMM using lme4 (*83*):

802
$$\log(E_s) = \beta_{KO}X_{s,KO} + (\phi_d|d) + \text{offset}(\log[libsize]) + \theta$$

where *E* is gene expression in sample *s*, *θ* is an intercept, *β*$_{KO}$ is the fixed effect for KO status, and Donor

(*d*) is a random intercept. Significance was determined via LRT. Enrichment of downregulated genes

(FDR < 0.05; **table S19**) within NaiveCD4-4/5 and CMCD-4 signature genes was tested using one-sided

Fisher's exact tests (**Fig. 6F**).

FCM and FACS experiments on EMCD8 cells

To characterize EMCD8-5 (DP EMCD8) cells, we performed functional assays using FACSMelody (BD

Biosciences) based on the four surface markers with the highest logFC values (HLA-DR, CD27, CD38,

and CD28; **Fig. 7E−F** and **table S14**). Two panels were prepared: Panel 1 for surface markers (PD-1,

TIGIT) and Panel 2 for intracellular IFN-γ (**table S22**).

For FACS, CD8$^+$ T cells were isolated from six SLE donors using the CD8$^+$ T Cell Isolation Kit

(Miltenyi, 130-096-495), blocked with anti-FcγR antibodies (Thermo Fisher Scientific), and stained with

Panel 1. Target EMCD8 (CCR7$^-$HLA-DR$^+$CD27$^+$CD38$^+$CD28$^+$) and total CCR7$^-$ EMCD8 cells (control)

were sorted (5,000 cells per sample; **fig. S10E**) for bulk RNA-seq (SMART-seq, Takara Bio 634772),

ATAC-seq, and UNIChro-seq. ATAC-seq tagmentation used TDE1 (Illumina 20034198) and 1% digitonin

(Promega G944A). Purified tagmented DNA was amplified for 15–18 cycles using KAPA HiFi ReadyMix

(Nippon Genetics KK2602) and Nextera XT Index Kit v2 and size-selected (180–1000 bp) using AMPure

XP beads. UNIChro-seq libraries were prepared with custom primers to quantify chromatin accessibility

(*68*) at five target *IFNG* and one reference *UBE2D2* loci (**table S24**) and sequenced on Miseq (illumina)

using a custom R1 primer (5'- ACACTCTTTCCCTACACGACGCTCTTCAGGTCT).

RNA-seq data were analyzed using the same pipeline as the **Arrayed CRISPR-KO experiment**,

with the following GLMM:

$$\log(E_s) = \beta_{sub} X_{s,sub} + \beta_{age} X_{d,age} + \beta_{sex} X_{d,sex} + (\phi_d|d) + \text{offset}(\log[libsize]) + \theta$$

where β_{sub} is the fixed effect for cell subsets. ATAC-seq and UNIChro-seq reads were mapped using

Bowtie2 (*90*), with low-quality (q < 30), duplicated (Picard), and ENCODE-blacklisted reads removed.

MACS2 (*91*) was used for peak calling. Differentially accessible regions (DARs) were identified using the

same GLMM framework as RNA-seq. Motif enrichment for DARs (FDR < 0.1, logFC >0) was performed

with HOMER (**fig. S10G**). For UNIChro-seq, we extracted UMI counts at target loci as previously

described (*68*). Accessibility at five IFNG loci relative to a UBE2D2 reference was evaluated using a

logistic mixed-effect model, regressing cell status (target/control EMCD8) by genomic region

(target/reference) with donor as a random effect. Genomic tracks were visualized using Integrative

Genomics Viewer (**Fig. 7G**).

For FCM, CD8[+] T cells from 11 SLE donors were stained with Panel 1 (surface) or Panel 2

(intracellular). For Panel 2, cells were stimulated with PMA (50 ng/ml, MP Biomedicals, 16561-29-8) and

ionomycin (1 µg/ml) for four hours, with GolgiStop (BD Biosciences, 554724) added after the first hour.

Intracellular staining was performed using the Cytofix/Cytoperm kit (BD Biosciences, 554714). Mean

fluorescence intensities (MFI) were compared between target and total EMCD8 cells using linear mixed

models (fixed effects: age, sex; random effect: donor; **Fig. 7H** and **fig. S10H**). All data were analyzed

using FlowJo (BD Biosciences).

Reference

1. G. C. Tsokos, M. S. Lo, P. C. Reis, K. E. Sullivan, New insights into the immunopathogenesis of systemic lupus erythematosus. *Nat. Rev. Rheumatol.* **12**, 716–730 (2016).

2. R. Banchereau, S. Hong, B. Cantarel, N. Baldwin, J. Baisch, M. Edens, A.-M. Cepika, P. Acs, J. Turner, E. Anguiano, P. Vinod, S. Khan, G. Obermoser, D. Blankenship, E. Wakeland, L. Nassi, A. Gotte, M. Punaro, Y.-J. Liu, J. Banchereau, J. Rossello-Urgell, T. Wright, V. Pascual, Personalized Immunomonitoring Uncovers Molecular Networks that Stratify Lupus Patients. *Cell* **165**, 551–565 (2016).

3. the Accelerating Medicines Partnership in SLE network, A. Arazi, D. A. Rao, C. C. Berthier, A. Davidson, Y. Liu, P. J. Hoover, A. Chicoine, T. M. Eisenhaure, A. H. Jonsson, S. Li, D. J. Lieb, F. Zhang, K. Slowikowski, E. P. Browne, A. Noma, D. Sutherby, S. Steelman, D. E. Smilek, P. Tosta, W. Apruzzese, E. Massarotti, M. Dall'Era, M. Park, D. L. Kamen, R. A. Furie, F. Payan-Schober, W. F. Pendergraft, E. A. McInnis, J. P. Buyon, M. A. Petri, C. Putterman, K. C. Kalunian, E. S. Woodle, J. A. Lederer, D. A. Hildeman, C. Nusbaum, S. Raychaudhuri, M. Kretzler, J. H. Anolik, M. B. Brenner, D. Wofsy, N. Hacohen, B. Diamond, The immune cell landscape in kidneys of patients with lupus nephritis. *Nat. Immunol.* **20**, 902–914 (2019).

4. D. Nehar-Belaid, S. Hong, R. Marches, G. Chen, M. Bolisetty, J. Baisch, L. Walters, M. Punaro, R. J. Rossi, C.-H. Chung, R. P. Huynh, P. Singh, W. F. Flynn, J.-A. Tabanor-Gayle, N. Kuchipudi, A. Mejias, M. A. Collet, A. L. Lucido, K. Palucka, P. Robson, S. Lakshminarayanan, O. Ramilo, T. Wright, V. Pascual, J. F. Banchereau, Mapping systemic lupus erythematosus heterogeneity at the single-cell level. *Nat. Immunol.* **21**, 1094–1106 (2020).

5. R. K. Perez, M. G. Gordon, M. Subramaniam, M. C. Kim, G. C. Hartoularos, S. Targ, Y. Sun, A. Ogorodnikov, R. Bueno, A. Lu, M. Thompson, N. Rappoport, A. Dahl, C. M. Lanata, M. Matloubian, L. Maliskova, S. S. Kwek, T. Li, M. Slyper, J. Waldman, D. Dionne, O. Rozenblatt-Rosen, L. Fong, M. Dall'Era, B. Balliu, A. Regev, J. Yazdany, L. A. Criswell, N. Zaitlen, C. J. Ye, Single-cell RNA-seq reveals cell type–specific molecular and genetic associations to lupus. *Science* **376**, eabf1970 (2022).

6. M. Nakano, M. Ota, Y. Takeshima, Y. Iwasaki, H. Hatano, Y. Nagafuchi, T. Itamiya, J. Maeda, R. Yoshida, S. Yamada, A. Nishiwaki, H. Takahashi, H. Takahashi, Y. Akutsu, T. Kusuda, H. Suetsugu, L. Liu, K. Kim, X. Yin, S.-Y. Bang, Y. Cui, H.-S. Lee, H. Shoda, X. Zhang, S.-C. Bae, C. Terao, K. Yamamoto, T. Okamura, K. Ishigaki, K. Fujio, Distinct transcriptome architectures underlying lupus establishment and exacerbation. *Cell* **185**, 3375-3389.e21 (2022).

7. S. V. Navarra, R. M. Guzmán, A. E. Gallacher, S. Hall, R. A. Levy, R. E. Jimenez, E. K.-M. Li, M. Thomas, H.-Y. Kim, M. G. León, C. Tanasescu, E. Nasonov, J.-L. Lan, L. Pineda, Z. J. Zhong, W. Freimuth, M. A. Petri, Efficacy and safety of belimumab in patients with active systemic lupus erythematosus: a randomised, placebo-controlled, phase 3 trial. *The Lancet* **377**, 721–731 (2011).

8.   E. F. Morand, R. Furie, Y. Tanaka, I. N. Bruce, A. D. Askanase, C. Richez, S.-C. Bae, P. Z. Brohawn, L. Pineda, A.
Berglind, R. Tummala, Trial of Anifrolumab in Active Systemic Lupus Erythematosus. *N. Engl. J. Med.* **382**, 211–221
(2020).

9.   D. A. Rao, M. F. Gurish, J. L. Marshall, K. Slowikowski, C. Y. Fonseka, Y. Liu, L. T. Donlin, L. A. Henderson, K. Wei, F.
Mizoguchi, N. C. Teslovich, M. E. Weinblatt, E. M. Massarotti, J. S. Coblyn, S. M. Helfgott, Y. C. Lee, D. J. Todd, V. P.
Bykerk, S. M. Goodman, A. B. Pernis, L. B. Ivashkiv, E. W. Karlson, P. A. Nigrovic, A. Filer, C. D. Buckley, J. A. Lederer,
S. Raychaudhuri, M. B. Brenner, Pathologically expanded peripheral T helper cell subset drives B cells in rheumatoid
arthritis. *Nature* **542**, 110–114 (2017).

10. S. A. Jenks, K. S. Cashman, E. Zumaquero, U. M. Marigorta, A. V. Patel, X. Wang, D. Tomar, M. C. Woodruff, Z. Simon, R.
Bugrovsky, E. L. Blalock, C. D. Scharer, C. M. Tipton, C. Wei, S. S. Lim, M. Petri, T. B. Niewold, J. H. Anolik, G. Gibson,
F. E.-H. Lee, J. M. Boss, F. E. Lund, I. Sanz, Distinct Effector B Cells Induced by Unregulated Toll-like Receptor 7
Contribute to Pathogenic Responses in Systemic Lupus Erythematosus. *Immunity* **49**, 725-739.e6 (2018).

11. C. Y. Fonseka, D. A. Rao, N. C. Teslovich, I. Korsunsky, S. K. Hannes, K. Slowikowski, M. F. Gurish, L. T. Donlin, J. A.
Lederer, M. E. Weinblatt, E. M. Massarotti, J. S. Coblyn, S. M. Helfgott, D. J. Todd, V. P. Bykerk, E. W. Karlson, J.
Ermann, Y. C. Lee, M. B. Brenner, S. Raychaudhuri, Mixed-effects association of single cells identifies an expanded
effector CD4$^+$ T cell subset in rheumatoid arthritis. *Sci. Transl. Med.* **10**, eaaq0305 (2018).

12. J. Li, M. Zaslavsky, Y. Su, J. Guo, M. J. Sikora, V. Van Unen, A. Christophersen, S.-H. Chiou, L. Chen, J. Li, X. Ji, J.
Wilhelmy, A. M. McSween, B. A. Palanski, V. V. A. Mallajosyula, N. A. Bracey, G. K. R. Dhondalay, K. Bhamidipati, J. Pai,
L. B. Kipp, J. E. Dunn, S. L. Hauser, J. R. Oksenberg, A. T. Satpathy, W. H. Robinson, C. L. Dekker, L. M. Steinmetz, C.
Khosla, P. J. Utz, L. M. Sollid, Y.-H. Chien, J. R. Heath, N. Q. Fernandez-Becker, K. C. Nadeau, N. Saligrama, M. M.
Davis, KIR$^+$ CD8$^+$ T cells suppress pathogenic T cells and are active in autoimmune diseases and COVID-19. *Science*
376, eabi9591 (2022).

13. F. Zhang, A. H. Jonsson, A. Nathan, N. Millard, M. Curtis, Q. Xiao, M. Gutierrez-Arcelus, W. Apruzzese, G. F. M. Watts, D.
Weisenfeld, S. Nayar, J. Rangel-Moreno, N. Meednu, K. E. Marks, I. Mantel, J. B. Kang, L. Rumker, J. Mears, K.
Slowikowski, K. Weinand, D. E. Orange, L. Geraldino-Pardilla, K. D. Deane, D. Tabechian, A. Ceponis, G. S. Firestein, M.
Maybury, I. Sahbudin, A. Ben-Artzi, A. M. Mandelin, A. Nerviani, M. J. Lewis, F. Rivellese, C. Pitzalis, L. B. Hughes, D.
Horowitz, E. DiCarlo, E. M. Gravallese, B. F. Boyce, Accelerating Medicines Partnership: RA/SLE Network, J. Albrecht, J.
L. Barnas, J. M. Bathon, D. L. Boyle, S. L. Bridges, D. Campbell, H. L. Carr, A. Chicoine, A. Cordle, P. Dunn, L. Forbess,
P. K. Gregersen, J. M. Guthridge, L. B. Ivashkiv, K. Ishigaki, J. A. James, G. Keras, I. Korsunsky, A. Lakhanpal, J. A.
Lederer, Z. J. Li, Y. Li, A. McDavid, M. J. McGeachy, K. Raza, Y. Reshef, C. Ritchlin, W. H. Robinson, S. Sakaue, J. A.
Seifert, A. Singaraju, M. H. Smith, D. Scheel-Toellner, P. J. Utz, M. H. Weisman, A. Wyse, Z. Zhu, L. W. Moreland, S. M.
Goodman, H. Perlman, V. M. Holers, K. P. Liao, A. Filer, V. P. Bykerk, K. Wei, D. A. Rao, L. T. Donlin, J. H. Anolik, M. B.
Brenner, S. Raychaudhuri, Deconstruction of rheumatoid arthritis synovium defines inflammatory subtypes. *Nature* **623**,
616–624 (2023).

14. E. Dann, N. C. Henderson, S. A. Teichmann, M. D. Morgan, J. C. Marioni, Differential abundance testing on single-cell data
using k-nearest neighbor graphs. *Nat. Biotechnol.* **40**, 245–253 (2022).

15. Y. A. Reshef, L. Rumker, J. B. Kang, A. Nathan, I. Korsunsky, S. Asgari, M. B. Murray, D. B. Moody, S. Raychaudhuri, Co-
varying neighborhood analysis identifies cell populations associated with phenotypes of interest from single-cell
transcriptomics. *Nat. Biotechnol.* **40**, 355–363 (2022).

16. M. Stoeckius, C. Hafemeister, W. Stephenson, B. Houck-Loomis, P. K. Chattopadhyay, H. Swerdlow, R. Satija, P. Smibert,
Simultaneous epitope and transcriptome measurement in single cells. *Nat. Methods* **14**, 865–868 (2017).

17. J. Moody, T. Kouno, M. Kojima, I. Koya, J. Leon, A. Suzuki, A. Hasegawa, T. Akiyama, N. Akiyama, M. Amagai, J.-C.
Chang, A. Fukushima-Nomura, M. Handa, K. Hino, M. Hino, T. Hirata, Y. Imai, K. Inoue, H. Kawasaki, T. Kimura, T.
Kinoshita, K. Kubo, Y. Kunii, F. López-Redondo, R. Manabe, T. Miyai, S. Morimoto, A. Nagaoka, J. Nakajima, S. Noma,
Y. Okazaki, K. Ozaki, N. Saeki, H. Sakai, K. Seyama, Y. Shibayama, T. Sujino, M. Tagami, H. Takahashi, M. Takao, M.
Takeshita, T. Takiuchi, C. Terao, C. W. Yip, S. Yoshinaga, H. Okano, K. Yahamoto, T. Kasukawa, Y. Ando, P. Carninci, J.
W. Shin, C.-C. Hon, A single-cell atlas of transcribed *cis* -regulatory elements in the human genome. Genomics [Preprint]
(2023). https://doi.org/10.1101/2023.11.13.566791.

18. A. Oguchi, A. Suzuki, S. Komatsu, H. Yoshitomi, S. Bhagat, R. Son, R. J. P. Bonnal, S. Kojima, M. Koido, K. Takeuchi, K.
Myouzen, G. Inoue, T. Hirai, H. Sano, Y. Takegami, A. Kanemaru, I. Yamaguchi, Y. Ishikawa, N. Tanaka, S. Hirabayashi,
R. Konishi, S. Sekito, T. Inoue, J. Kere, S. Takeda, A. Takaori-Kondo, I. Endo, S. Kawaoka, H. Kawaji, K. Ishigaki, H.
Ueno, Y. Hayashizaki, M. Pagani, P. Carninci, M. Yanagita, ITEC Consortium‡, N. Parrish, C. Terao, K. Yamamoto, Y.
Murakawa, M. Guerrini, H. Hatano, K. Ishigaki, Y. Ishikawa, S. Komatsu, M. Kono, Y. Murakawa, M. Nakano, A. Oguchi,
R. Son, A. Suzuki, K. Takeuchi, N. Tanaka, C. Terao, K. Tomizuka, K. Yamamoto, S. Yoshino, An atlas of transcribed
enhancers across helper T cell diversity for decoding human diseases. *Science* **385**, eadd8394 (2024).

19. Y. Hao, S. Hao, E. Andersen-Nissen, W. M. Mauck, S. Zheng, A. Butler, M. J. Lee, A. J. Wilk, C. Darby, M. Zager, P.
Hoffman, M. Stoeckius, E. Papalexi, E. P. Mimitou, J. Jain, A. Srivastava, T. Stuart, L. M. Fleming, B. Yeung, A. J.
Rogers, J. M. McElrath, C. A. Blish, R. Gottardo, P. Smibert, R. Satija, Integrated analysis of multimodal single-cell data.
Cell **184**, 3573-3587.e29 (2021).

20. M. Watanabe, K. D. Moon, M. S. Vacchio, K. S. Hathcock, R. J. Hodes, Downmodulation of Tumor Suppressor p53 by T
Cell Receptor Signaling Is Critical for Antigen-Specific CD4+ T Cell Responses. *Immunity* **40**, 681–691 (2014).

21. M. C. Castellanos, C. Muñoz, M. C. Montoya, E. Lara-Pezzi, M. López-Cabrera, M. O. de Landázuri, Expression of the
leukocyte early activation antigen CD69 is regulated by the transcription factor AP-1. *J. Immunol. Baltim. Md 1950* **159**,
5463–5473 (1997).

22. P. Bohacova, M. Terekhova, P. Tsurinov, R. Mullins, K. Husarcikova, I. Shchukina, A. U. Antonova, B. Echalar, J. Kossl, A.
Saidu, T. Francis, C. Mannie, L. Arthur, S. D. R. Harridge, D. Kreisel, P. A. Mudd, A. M. Taylor, C. A. McNamara, M.

Cella, S. V. Puram, T. Van Den Broek, F. Van Wijk, P. Eghtesady, M. N. Artyomov, Multidimensional profiling of human T

cells reveals high CD38 expression, marking recent thymic emigrants and age-related naive T cell remodeling. *Immunity*

57, 2362-2379.e10 (2024).

23. Y. Yasumizu, D. Takeuchi, R. Morimoto, Y. Takeshima, T. Okuno, M. Kinoshita, T. Morita, Y. Kato, M. Wang, D. Motooka,

D. Okuzaki, Y. Nakamura, N. Mikami, M. Arai, X. Zhang, A. Kumanogoh, H. Mochizuki, N. Ohkura, S. Sakaguchi, Single-

cell transcriptome landscape of circulating CD4+ T cell populations in autoimmune diseases. *Cell Genomics* **4**, 100473

(2024).

24. J. D. Chan, C. M. Scheffler, I. Munoz, K. Sek, J. N. Lee, Y.-K. Huang, K. M. Yap, N. Y. L. Saw, J. Li, A. X. Y. Chen, C. W.

Chan, E. B. Derrick, K. L. Todd, J. Tong, P. A. Dunbar, J. Li, T. X. Hoang, M. N. De Menezes, E. V. Petley, J. S. Kim, D.

Nguyen, P. S. K. Leung, J. So, C. Deguit, J. Zhu, I. G. House, L. M. Kats, A. M. Scott, B. J. Solomon, S. J. Harrison, J.

Oliaro, I. A. Parish, K. M. Quinn, P. J. Neeson, C. Y. Slaney, J. Lai, P. A. Beavis, P. K. Darcy, FOXO1 enhances CAR T

cell stemness, metabolic fitness and efficacy. *Nature* **629**, 201–210 (2024).

25. A. E. Doan, K. P. Mueller, A. Y. Chen, G. T. Rouin, Y. Chen, B. Daniel, J. Lattin, M. Markovska, B. Mozarsky, J. Arias-

Umana, R. Hapke, I.-Y. Jung, A. Wang, P. Xu, D. Klysz, G. Zuern, M. Bashti, P. J. Quinn, Z. Miao, K. Sandor, W. Zhang,

G. M. Chen, F. Ryu, M. Logun, J. Hall, K. Tan, S. A. Grupp, S. E. McClory, C. A. Lareau, J. A. Fraietta, E. Sotillo, A. T.

Satpathy, C. L. Mackall, E. W. Weber, FOXO1 is a master regulator of memory programming in CAR T cells. *Nature* **629**,

211–218 (2024).

26. M. K. Crow, Type I Interferon in the Pathogenesis of Lupus. *J. Immunol.* **192**, 5459–5468 (2014).

27. J. He, L. M. Tsai, Y. A. Leong, X. Hu, C. S. Ma, N. Chevalier, X. Sun, K. Vandenberg, S. Rockman, Y. Ding, L. Zhu, W.

Wei, C. Wang, A. Karnowski, G. T. Belz, J. R. Ghali, M. C. Cook, D. S. Riminton, A. Veillette, P. L. Schwartzberg, F.

Mackay, R. Brink, S. G. Tangye, C. G. Vinuesa, C. R. Mackay, Z. Li, D. Yu, Circulating Precursor CCR7loPD-1hi CXCR5+

CD4+ T Cells Indicate Tfh Cell Activity and Promote Antibody Responses upon Antigen Reexposure. *Immunity* **39**, 770–

781 (2013).

28. T. Koga, C. M. Hedrich, M. Mizui, N. Yoshida, K. Otomo, L. A. Lieberman, T. Rauen, J. C. Crispín, G. C. Tsokos, CaMK4-

dependent activation of AKT/mTOR and CREM-α underlies autoimmunity-associated Th17 imbalance. *J. Clin. Invest.*

124, 2234–2245 (2014).

29. V. S. Patil, A. Madrigal, B. J. Schmiedel, J. Clarke, P. O'Rourke, A. D. De Silva, E. Harris, B. Peters, G. Seumois, D.

Weiskopf, A. Sette, P. Vijayanand, Precursors of human CD4+ cytotoxic T lymphocytes identified by single-cell

transcriptome analysis. *Sci. Immunol.* **3**, eaan8664 (2018).

30. Y. Luo, C. Xu, B. Wang, Q. Niu, X. Su, Y. Bai, S. Zhu, C. Zhao, Y. Sun, J. Wang, M. Liu, X. Sun, G. Song, H. Cui, X. Chen,

H. Huang, H. Wang, M. Han, E. Jiang, L. Shi, X. Feng, Single-cell transcriptomic analysis reveals disparate effector

differentiation pathways in human Treg compartment. *Nat. Commun.* **12**, 3913 (2021).

31. A. H. Jonsson, F. Zhang, G. Dunlap, E. Gomez-Rivas, G. F. M. Watts, H. J. Faust, K. V. Rupani, J. R. Mears, N. Meednu,
R. Wang, G. Keras, J. S. Coblyn, E. M. Massarotti, D. J. Todd, J. H. Anolik, A. McDavid, Accelerating Medicines
Partnership RA/SLE Network, K. Wei, D. A. Rao, S. Raychaudhuri, M. B. Brenner, Granzyme K[+] CD8 T cells form a core
population in inflamed human tissue. *Sci. Transl. Med.* **14**, eabo0686 (2022).

32. T. Velikkakam, K. J. Gollob, W. O. Dutra, Double-negative T cells: Setting the stage for disease control or progression.
Immunology **165**, 371–385 (2022).

33. M. Colonna, Innate Lymphoid Cells: Diversity, Plasticity, and Unique Functions in Immunity. *Immunity* **48**, 1104–1117
(2018).

34. D. Morgan, V. Tergaonkar, Unraveling B cell trajectories at single cell resolution. *Trends Immunol.* **43**, 210–229 (2022).

35. S. L. Nutt, P. D. Hodgkin, D. M. Tarlinton, L. M. Corcoran, The generation of antibody-secreting plasma cells. *Nat. Rev.*
Immunol. **15**, 160–171 (2015).

36. K. Mulder, A. A. Patel, W. T. Kong, C. Piot, E. Halitzki, G. Dunsmore, S. Khalilnezhad, S. E. Irac, A. Dubuisson, M.
Chevrier, X. M. Zhang, J. K. C. Tam, T. K. H. Lim, R. M. M. Wong, R. Pai, A. I. S. Khalil, P. K. H. Chow, S. Z. Wu, G. Al-
Eryani, D. Roden, A. Swarbrick, J. K. Y. Chan, S. Albani, L. Derosa, L. Zitvogel, A. Sharma, J. Chen, A. Silvin, A.
Bertoletti, C. Blériot, C.-A. Dutertre, F. Ginhoux, Cross-tissue single-cell landscape of human monocytes and
macrophages in health and disease. *Immunity* **54**, 1883-1900.e5 (2021).

37. J. Leffler, A. A. Bengtsson, A. M. Blom, The complement system in systemic lupus erythematosus: an update. *Ann. Rheum.*
Dis. **73**, 1601–1606 (2014).

38. A.-C. Villani, R. Satija, G. Reynolds, S. Sarkizova, K. Shekhar, J. Fletcher, M. Griesbeck, A. Butler, S. Zheng, S. Lazo, L.
Jardine, D. Dixon, E. Stephenson, E. Nilsson, I. Grundberg, D. McDonald, A. Filby, W. Li, P. L. De Jager, O. Rozenblatt-
Rosen, A. A. Lane, M. Haniffa, A. Regev, N. Hacohen, Single-cell RNA-seq reveals new types of human blood dendritic
cells, monocytes, and progenitors. *Science* **356**, eaah4573 (2017).

39. Q. Zhang, Y. He, N. Luo, S. J. Patel, Y. Han, R. Gao, M. Modak, S. Carotta, C. Haslinger, D. Kind, G. W. Peet, G. Zhong,
S. Lu, W. Zhu, Y. Mao, M. Xiao, M. Bergmann, X. Hu, S. P. Kerkar, A. B. Vogt, S. Pflanz, K. Liu, J. Peng, X. Ren, Z.
Zhang, Landscape and Dynamics of Single Immune Cells in Hepatocellular Carcinoma. *Cell* **179**, 829-845.e20 (2019).

40. H. K. Finucane, Y. A. Reshef, V. Anttila, K. Slowikowski, A. Gusev, A. Byrnes, S. Gazal, P.-R. Loh, C. Lareau, N. Shoresh,
G. Genovese, A. Saunders, E. Macosko, S. Pollack, The Brainstorm Consortium, J. R. B. Perry, J. D. Buenrostro, B. E.
Bernstein, S. Raychaudhuri, S. McCarroll, B. M. Neale, A. L. Price, Heritability enrichment of specifically expressed genes
identifies disease-relevant tissues and cell types. *Nat. Genet.* **50**, 621–629 (2018).

41. R. P. Taylor, M. A. Lindorfer, Drug Insight: the mechanism of action of rituximab in autoimmune disease—the immune
complex decoy hypothesis. *Nat. Clin. Pract. Rheumatol.* **3**, 86–95 (2007).

42. F. Müller, J. Taubmann, L. Bucci, A. Wilhelm, C. Bergmann, S. Völkl, M. Aigner, T. Rothe, I. Minopoulou, C. Tur, J. Knitza,
S. Kharboutli, S. Kretschmann, I. Vasova, S. Spoerl, H. Reimann, L. Munoz, R. G. Gerlach, S. Schäfer, R. Grieshaber-
Bouyer, A.-S. Korganow, D. Farge-Bancel, D. Mougiakakos, A. Bozec, T. Winkler, G. Krönke, A. Mackensen, G. Schett,
CD19 CAR T-Cell Therapy in Autoimmune Disease — A Case Series with Follow-up. *N. Engl. J. Med.* **390**, 687–700
(2024).

43. E. Mountjoy, E. M. Schmidt, M. Carmona, J. Schwartzentruber, G. Peat, A. Miranda, L. Fumis, J. Hayhurst, A. Buniello, M.
A. Karim, D. Wright, A. Hercules, E. Papa, E. B. Fauman, J. C. Barrett, J. A. Todd, D. Ochoa, I. Dunham, M. Ghoussaini,
An open approach to systematically prioritize causal variants and genes at all published human GWAS trait-associated
loci. *Nat. Genet.* **53**, 1527–1533 (2021).

44. X. Yin, K. Kim, H. Suetsugu, S.-Y. Bang, L. Wen, M. Koido, E. Ha, L. Liu, Y. Sakamoto, S. Jo, R.-X. Leng, N. Otomo, V.
Laurynenka, Y.-C. Kwon, Y. Sheng, N. Sugano, M. Y. Hwang, W. Li, M. Mukai, K. Yoon, M. Cai, K. Ishigaki, W. T. Chung,
H. Huang, D. Takahashi, S.-S. Lee, M. Wang, K. Karino, S.-C. Shim, X. Zheng, T. Miyamura, Y. M. Kang, D. Ye, J.
Nakamura, C.-H. Suh, Y. Tang, G. Motomura, Y.-B. Park, H. Ding, T. Kuroda, J.-Y. Choe, C. Li, H. Niiro, Y. Park, C.
Shen, T. Miyamoto, G.-Y. Ahn, W. Fei, T. Takeuchi, J.-M. Shin, K. Li, Y. Kawaguchi, Y.-K. Lee, Y. Wang, K. Amano, D. J.
Park, W. Yang, Y. Tada, K. Yamaji, M. Shimizu, T. Atsumi, A. Suzuki, T. Sumida, Y. Okada, K. Matsuda, K. Matsuo, Y.
Kochi, L. C. Kottyan, M. T. Weirauch, S. Parameswaran, S. Eswar, H. Salim, X. Chen, K. Yamamoto, J. B. Harley, K.
Ohmura, T.-H. Kim, S. Yang, T. Yamamoto, B.-J. Kim, N. Shen, S. Ikegawa, H.-S. Lee, X. Zhang, C. Terao, Y. Cui, S.-C.
Bae, Meta-analysis of 208370 East Asians identifies 113 susceptibility loci for systemic lupus erythematosus. *Ann.*
Rheum. Dis. **80**, 632–640 (2021).

45. C. D. Langefeld, H. C. Ainsworth, D. S. C. Graham, J. A. Kelly, M. E. Comeau, M. C. Marion, T. D. Howard, P. S. Ramos, J.
A. Croker, D. L. Morris, J. K. Sandling, J. C. Almlöf, E. M. Acevedo-Vásquez, G. S. Alarcón, A. M. Babini, V. Baca, A. A.
Bengtsson, G. A. Berbotto, M. Bijl, E. E. Brown, H. I. Brunner, M. H. Cardiel, L. Catoggio, R. Cervera, J. M. Cucho-
Venegas, S. R. Dahlqvist, S. D'Alfonso, B. M. Da Silva, I. De La Rúa Figueroa, A. Doria, J. C. Edberg, E. Endreffy, J. A.
Esquivel-Valerio, P. R. Fortin, B. I. Freedman, J. Frostegård, M. A. García, I. G. De La Torre, G. S. Gilkeson, D. D.
Gladman, I. Gunnarsson, J. M. Guthridge, J. L. Huggins, J. A. James, C. G. M. Kallenberg, D. L. Kamen, D. R. Karp, K.
M. Kaufman, L. C. Kottyan, L. Kovács, H. Laustrup, B. R. Lauwerys, Q.-Z. Li, M. A. Maradiaga-Ceceña, J. Martín, J. M.
McCune, D. R. McWilliams, J. T. Merrill, P. Miranda, J. F. Moctezuma, S. K. Nath, T. B. Niewold, L. Orozco, N. Ortego-
Centeno, M. Petri, C. A. Pineau, B. A. Pons-Estel, J. Pope, P. Raj, R. Ramsey-Goldman, J. D. Reveille, L. P. Russell, J.
M. Sabio, C. A. Aguilar-Salinas, H. R. Scherbarth, R. Scorza, M. F. Seldin, C. Sjöwall, E. Svenungsson, S. D. Thompson,
S. M. A. Toloza, L. Truedsson, T. Tusié-Luna, C. Vasconcelos, L. M. Vilá, D. J. Wallace, M. H. Weisman, J. E. Wither, T.
Bhangale, J. R. Oksenberg, J. D. Rioux, P. K. Gregersen, A.-C. Syvänen, L. Rönnblom, L. A. Criswell, C. O. Jacob, K. L.
Sivils, B. P. Tsao, L. E. Schanberg, T. W. Behrens, E. D. Silverman, M. E. Alarcón-Riquelme, R. P. Kimberly, J. B. Harley,

E. K. Wakeland, R. R. Graham, P. M. Gaffney, T. J. Vyse, Transancestral mapping and genetic load in systemic lupus erythematosus. *Nat. Commun.* **8**, 16021 (2017).

46. D. Dimitrov, D. Türei, M. Garrido-Rodriguez, P. L. Burmedi, J. S. Nagai, C. Boys, R. O. Ramirez Flores, H. Kim, B. Szalai, I. G. Costa, A. Valdeolivas, A. Dugourd, J. Saez-Rodriguez, Comparison of methods and resources for cell-cell communication inference from single-cell RNA-Seq data. *Nat. Commun.* **13**, 3224 (2022).

47. E. Zumaquero, S. L. Stone, C. D. Scharer, S. A. Jenks, A. Nellore, B. Mousseau, A. Rosal-Vela, D. Botta, J. E. Bradley, W. Wojciechowski, T. Ptacek, M. I. Danila, J. C. Edberg, S. L. Bridges, R. P. Kimberly, W. W. Chatham, T. R. Schoeb, A. F. Rosenberg, J. M. Boss, I. Sanz, F. E. Lund, IFNγ induces epigenetic programming of human T-bethi B cells and promotes TLR7/8 and IL-21 induced differentiation. *eLife* **8**, e41641 (2019).

48. B. K. Han, N. J. Olsen, A. Bottaro, The CD27–CD70 pathway and pathogenesis of autoimmune disease. *Semin. Arthritis Rheum.* **45**, 496–501 (2016).

49. Q. Zhang, D. A. A. Vignali, Co-stimulatory and Co-inhibitory Pathways in Autoimmunity. *Immunity* **44**, 1034–1051 (2016).

50. S. M. Stanford, M. N. D. Svensson, C. Sacchetti, C. A. Pilo, D. J. Wu, W. B. Kiosses, A. Hellvard, B. Bergum, G. R. A. Muench, C. Elly, Y.-C. Liu, J. den Hertog, A. Elson, J. Sap, P. Mydel, D. L. Boyle, M. Corr, G. S. Firestein, N. Bottini, Receptor Protein Tyrosine Phosphatase α–Mediated Enhancement of Rheumatoid Synovial Fibroblast Signaling and Promotion of Arthritis in Mice. *Arthritis Rheumatol.* **68**, 359–369 (2016).

51. J. Huang, X. Liu, Y. Wei, X. Li, S. Gao, L. Dong, X. Rao, J. Zhong, Emerging Role of Dipeptidyl Peptidase-4 in Autoimmune Disease. *Front. Immunol.* **13**, 830863 (2022).

52. E. Doherty, L. Osmani, J. Bilsborrow, J. Par-Young, S. Choi, P. Tilstam, M. Shin, M. Piecychna, H. Cai, L. Leng, W.-U. Kim, I. Kang, R. Bucala, Pathogenic role of MIF receptor (CD74) expressing T cells in inflammatory arthritis. *Proc. Natl. Acad. Sci.* **123**, e2509156123 (2026).

53. J. B. Kang, A. Nathan, K. Weinand, F. Zhang, N. Millard, L. Rumker, D. B. Moody, I. Korsunsky, S. Raychaudhuri, Efficient and precise single-cell reference atlas mapping with Symphony. *Nat. Commun.* **12**, 5890 (2021).

54. K. A. Lagattuta, J. B. Kang, A. Nathan, K. E. Pauken, A. H. Jonsson, D. A. Rao, A. H. Sharpe, K. Ishigaki, S. Raychaudhuri, Repertoire analyses reveal T cell antigen receptor sequence features that influence T cell fate. *Nat. Immunol.* **23**, 446–457 (2022).

55. Y. Nagafuchi, M. Nakano, K. A. Lagattuta, M. Ota, H. Hatano, H. Takahashi, T. Itamiya, H. Inokuchi, S. Raychaudhuri, T. Okamura, K. Fujio, K. Ishigaki, T cell plasticity in systemic lupus erythematosus revealed by large-scale T cell receptor repertoire and transcriptome studies. Rheumatology [Preprint] (2025). https://doi.org/10.1101/2025.01.06.24319648.

56. O. V. Britanova, K. R. Lupyr, D. B. Staroverov, I. A. Shagina, A. A. Aleksandrov, Y. Y. Ustyugov, D. V. Somov, A. Klimenko, N. A. Shostak, I. V. Zvyagin, A. V. Stepanov, E. M. Merzlyak, A. N. Davydov, M. Izraelson, E. S. Egorov, E. A. Bogdanova, A. K. Vladimirova, P. A. Iakovlev, D. A. Fedorenko, R. A. Ivanov, V. I. Skvortsova, S. Lukyanov, D. M. Chudakov, Targeted depletion of TRBV9+ T cells as immunotherapy in a patient with ankylosing spondylitis. *Nat. Med.* **29**, 2731–2736 (2023).

57. Roadmap Epigenomics Consortium, A. Kundaje, W. Meuleman, J. Ernst, M. Bilenky, A. Yen, A. Heravi-Moussavi, P. Kheradpour, Z. Zhang, J. Wang, M. J. Ziller, V. Amin, J. W. Whitaker, M. D. Schultz, L. D. Ward, A. Sarkar, G. Quon, R. S. Sandstrom, M. L. Eaton, Y.-C. Wu, A. R. Pfenning, X. Wang, M. Claussnitzer, Y. Liu, C. Coarfa, R. A. Harris, N. Shoresh, C. B. Epstein, E. Gjoneska, D. Leung, W. Xie, R. D. Hawkins, R. Lister, C. Hong, P. Gascard, A. J. Mungall, R. Moore, E. Chuah, A. Tam, T. K. Canfield, R. S. Hansen, R. Kaul, P. J. Sabo, M. S. Bansal, A. Carles, J. R. Dixon, K.-H. Farh, S. Feizi, R. Karlic, A.-R. Kim, A. Kulkarni, D. Li, R. Lowdon, G. Elliott, T. R. Mercer, S. J. Neph, V. Onuchic, P. Polak, N. Rajagopal, P. Ray, R. C. Sallari, K. T. Siebenthall, N. A. Sinnott-Armstrong, M. Stevens, R. E. Thurman, J. Wu, B. Zhang, X. Zhou, A. E. Beaudet, L. A. Boyer, P. L. De Jager, P. J. Farnham, S. J. Fisher, D. Haussler, S. J. M. Jones, W. Li, M. A. Marra, M. T. McManus, S. Sunyaev, J. A. Thomson, T. D. Tlsty, L.-H. Tsai, W. Wang, R. A. Waterland, M. Q. Zhang, L. H. Chadwick, B. E. Bernstein, J. F. Costello, J. R. Ecker, M. Hirst, A. Meissner, A. Milosavljevic, B. Ren, J. A. Stamatoyannopoulos, T. Wang, M. Kellis, Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317–330 (2015).

58. V. R. Moulton, G. C. Tsokos, Abnormalities of T cell signaling in systemic lupus erythematosus. *Arthritis Res. Ther.* **13**, 207 (2011).

59. A. Meyer, P. J. Parmar, S. Shahrara, Significance of IL-7 and IL-7R in RA and autoimmunity. *Autoimmun. Rev.* **21**, 103120 (2022).

60. L. Sun, X. Gang, Z. Li, X. Zhao, T. Zhou, S. Zhang, G. Wang, Advances in Understanding the Roles of CD244 (SLAMF4) in Immune Regulation and Associated Diseases. *Front. Immunol.* **12**, 648182 (2021).

61. E. Katsuyama, M. Humbel, A. Suarez-Fueyo, A. Satyam, N. Yoshida, V. C. Kyttaris, M. G. Tsokos, G. C. Tsokos, CD38 in SLE CD4 T cells promotes Ca2+ flux and suppresses interleukin-2 production by enhancing the expression of GM2 on the surface membrane. *Nat. Commun.* **15**, 8304 (2024).

62. P. Badia-i-Mompel, J. Vélez Santiago, J. Braunger, C. Geiss, D. Dimitrov, S. Müller-Dott, P. Taus, A. Dugourd, C. H. Holland, R. O. Ramirez Flores, J. Saez-Rodriguez, decoupleR: ensemble of computational methods to infer biological activities from omics data. *Bioinforma. Adv.* **2**, vbac016 (2022).

63. Z. Zou, T. Ohta, S. Oki, ChIP-Atlas 3.0: a data-mining suite to explore chromosome architecture together with large-scale regulome data. *Nucleic Acids Res.* **52**, W45–W53 (2024).

64. J. Cao, M. Spielmann, X. Qiu, X. Huang, D. M. Ibrahim, A. J. Hill, F. Zhang, S. Mundlos, L. Christiansen, F. J. Steemers, C. Trapnell, J. Shendure, The single-cell transcriptional landscape of mammalian organogenesis. *Nature* **566**, 496–502 (2019).

65. C. Saggau, P. Bacher, D. Esser, M. Rasa, S. Meise, N. Mohr, N. Kohlstedt, A. Hutloff, S.-S. Schacht, J. Dargvainiene, G. R. Martini, K. H. Stürner, I. Schröder, R. Markewitz, J. Hartl, M. Hastermann, A. Duchow, P. Schindler, M. Becker, C. Bautista, J. Gottfreund, J. Walter, J. K. Polansky, M. Yang, R. Naghavian, M. Wendorff, E.-M. Schuster, A. Dahl, A. Petzold, S. Reinhardt, A. Franke, M. Wieczorek, L. Henschel, D. Berger, G. Heine, M. Holtsche, V. Häußler, C. Peters, E. Schmidt, S. Fillatreau, D. H. Busch, K.-P. Wandinger, K. Schober, R. Martin, F. Paul, F. Leypoldt, A. Scheffold, Autoantigen-specific CD4+ T cells acquire an exhausted phenotype and persist in human antigen-specific autoimmune diseases. *Immunity* **57**, 2416-2432.e8 (2024).

66. Z. Zhou, H. He, K. Wang, X. Shi, Y. Wang, Y. Su, Y. Wang, D. Li, W. Liu, Y. Zhang, L. Shen, W. Han, L. Shen, J. Ding, F. Shao, Granzyme A from cytotoxic lymphocytes cleaves GSDMB to trigger pyroptosis in target cells. *Science* **368**, eaaz7548 (2020).

67. A. Perl, L. Morel, Expanding scope of TEMRA in autoimmunity. *eBioMedicine* **90**, 104520 (2023).

68. M. Kono, H. Hatano, K. Asahara, M. Nakano, R. Bagherzadeh, T. Kawashima, T. Arakawa, M. Sato, H. Inokuchi, T. Nishino, T. Itamiya, H. Takahashi, B. Natsumoto, A. Suzuki, K. Yamamoto, K. Ishigaki, Accurate, sensitive, and efficient chromatin accessibility quantification at target loci using UNIChro-seq. *medRxiv*, 2025.07.29.25332340 (2025).

69. L. Chiche, N. Jourde-Chiche, E. Whalen, S. Presnell, V. Gersuk, K. Dang, E. Anguiano, C. Quinn, S. Burtey, Y. Berland, G. Kaplanski, J.-R. Harle, V. Pascual, D. Chaussabel, Modular Transcriptional Repertoire Analyses of Adults With Systemic Lupus Erythematosus Reveal Distinct Type I and Type II Interferon Signatures. *Arthritis Rheumatol.* **66**, 1583–1595 (2014).

70. M. E. Munroe, R. Lu, Y. D. Zhao, D. A. Fife, J. M. Robertson, J. M. Guthridge, T. B. Niewold, G. C. Tsokos, M. P. Keith, J. B. Harley, J. A. James, Altered type II interferon precedes autoantibody accrual and elevated type I interferon activity prior to systemic lupus erythematosus classification. *Ann. Rheum. Dis.* **75**, 2014–2021 (2016).

71. A. Fava, J. Buyon, C. Mohan, T. Zhang, H. M. Belmont, P. Izmirly, R. Clancy, J. M. Trujillo, D. Fine, Y. Zhang, L. Magder, D. A. Rao, A. Arazi, C. C. Berthier, A. Davidson, B. Diamond, N. Hacohen, D. Wofsy, W. Apruzzese, S. Raychaudhuri, M. Petri, Integrated urine proteomics and renal single-cell genomics identify an IFN-γ response gradient in lupus nephritis. *JCI Insight* **5**, e138345 (2020).

72. A. Wagner, A. Regev, N. Yosef, Revealing the vectors of cellular identity with single-cell genomics. *Nat. Biotechnol.* **34**, 1145–1160 (2016).

73. S. Gurajala, N. Sugiarto, M. Curtis, T. Eisenhaure, A. Arazi, A. Fava, Q. Xiao, J. Mears, B. Rovin, C. C. Berthier, P. Izmirly,
J. Barnas, P. J. Hoover, M. Peters, R. Raychowdhury, A. Horsberger, S. Sakaue, Y. Zhao, R. A. Furie, M. Belmont, D.
Hildeman, E. S. Woodle, M. Dall'Era, C. Putterman, D. L. Kamen, M. A. McMahon, J. Grossman, K. Kalunian, J. Hodgin,
F. Payan-Schober, W. Apruzzese, H. Perlman, C. M. Cuda, D. Wofsy, J. M. Guthridge, J. H. Anolik, J. A. James,
Accelerating Medicines Partnerships Rheumatoid Arthritis/Systemic Lupus Erythematosus, D. A. Rao, A. Davidson, M. A.
Petri, J. P. Buyon, N. Hacohen, B. Diamond, S. Raychaudhuri, A population-scale atlas of blood and tissue in lupus
nephritis. *bioRxiv*, 2025.08.11.669754 (2025).

74. A. Missarova, E. Dann, L. Rosen, R. Satija, J. Marioni, Leveraging neighborhood representations of single-cell data to
achieve sensitive DE testing with miloDE. *Genome Biol.* **25**, 189 (2024).

75. T. Lappalainen, D. G. MacArthur, From variant to function in human disease genetics. *Science* **373**, 1464–1468 (2021).

76. A. F. Labrijn, M. L. Janmaat, J. M. Reichert, P. W. H. I. Parren, Bispecific antibodies: a mechanistic review of the pipeline.
Nat. Rev. Drug Discov. **18**, 585–608 (2019).

77. C. Bombardier, D. D. Gladman, M. B. Urowitz, D. Caron, C. H. Chang, A. Austin, A. Bell, D. A. Bloch, P. N. Corey, J. L.
Decker, J. Esdaile, J. F. Fries, E. M. Ginzler, C. H. Goldsmith, M. C. Hochberg, J. V. Jones, N. G. H. L. Riche, M. H.
Liang, M. D. Lockshin, L. R. Muenz, D. L. Sackett, P. H. Schur, Derivation of the sledai. A disease activity index for lupus
patients. *Arthritis Rheum.* **35**, 630–640 (1992).

78. A. Rogozhnikov, P. Ramkumar, K. Shah, R. Bedi, S. Kato, G. S. Escola, Demuxalot: scaled up genetic demultiplexing for
single-cell sequencing. Bioinformatics [Preprint] (2021). https://doi.org/10.1101/2021.05.22.443646.

79. H. Heaton, A. M. Talman, A. Knights, M. Imaz, D. J. Gaffney, R. Durbin, M. Hemberg, M. K. N. Lawniczak, Souporcell:
robust clustering of single-cell RNA-seq data by genotype without reference genotypes. *Nat. Methods* **17**, 615–620
(2020).

80. Y. Hao, T. Stuart, M. H. Kowalski, S. Choudhary, P. Hoffman, A. Hartman, A. Srivastava, G. Molla, S. Madad, C.
Fernandez-Granda, R. Satija, Dictionary learning for integrative, multimodal and scalable single-cell analysis. *Nat.*
Biotechnol. **42**, 293–304 (2024).

81. I. Korsunsky, N. Millard, J. Fan, K. Slowikowski, F. Zhang, K. Wei, Y. Baglaenko, M. Brenner, P. Loh, S. Raychaudhuri,
Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat. Methods* **16**, 1289–1296 (2019).

82. S. L. Wolock, R. Lopez, A. M. Klein, Scrublet: Computational Identification of Cell Doublets in Single-Cell Transcriptomic
Data. *Cell Syst.* **8**, 281-291.e9 (2019).

83. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using **lme4**. *J. Stat. Softw.* **67** (2015).

84. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR : a Bioconductor package for differential expression analysis of digital
gene expression data. *Bioinformatics* **26**, 139–140 (2010).

85. S. Xu, E. Hu, Y. Cai, Z. Xie, X. Luo, L. Zhan, W. Tang, Q. Wang, B. Liu, R. Wang, W. Xie, T. Wu, L. Xie, G. Yu, Using
clusterProfiler to characterize multiomics data. *Nat. Protoc.* **19**, 3292–3320 (2024).

86. K. A. Jagadeesh, K. K. Dey, D. T. Montoro, R. Mohan, S. Gazal, J. M. Engreitz, R. J. Xavier, A. L. Price, A. Regev,
Identifying disease-critical cell types and cellular processes by integrating single-cell RNA-sequencing and human
genetics. *Nat. Genet.* **54**, 1479–1492 (2022).

87. J. W. Freimer, O. Shaked, S. Naqvi, N. Sinnott-Armstrong, A. Kathiria, C. M. Garrido, A. F. Chen, J. T. Cortez, W. J.
Greenleaf, J. K. Pritchard, A. Marson, Systematic discovery and perturbation of regulatory genes in human T cells reveals
the architecture of immune networks. *Nat. Genet.* **54**, 1133–1144 (2022).

88. T. Hayashi, H. Ozaki, Y. Sasagawa, M. Umeda, H. Danno, I. Nikaido, Single-cell full-length total RNA sequencing uncovers
dynamics of recursive splicing and enhancer RNAs. *Nat. Commun.* **9**, 619 (2018).

89. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, T. R. Gingeras, STAR:
ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

90. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–359 (2012).

91. Y. Zhang, T. Liu, C. A. Meyer, J. Eeckhoute, D. S. Johnson, B. E. Bernstein, C. Nusbaum, R. M. Myers, M. Brown, W. Li, X.
S. Liu, Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

1171