Title

Cellular and molecular fine mapping pinpoints new immunopathology of lupus

Authors

Masahiro Nakano[1,21]*, Michihiro Kono[1,2,21], Kenichiro Asahara[1,3], Takayuki Katsuyama[4], Satoshi Kubo[5],

Eri Katsuyama[4,6], Yuya Fujita[5], Hajime Inokuchi[1,2,7], Takahiro Nishino[1,2], Takahiro Arakawa[1], Tsugumi

Kawashima[1], Shohei Noma[1], Reza Bagherzadeh[1,2], Yukiro Matsumoto[1,2,8], Jun Inamo[1,2], Haruka

Takahashi[1,2], Bunki Natsumoto[1], Xuejun Zhang[9,10,11], Sang-Cheol Bae[12,13,14], Akari Suzuki[15], Hiroaki

Hatano[1,2], Chikashi Terao[16,17,18], Yoshiya Tanaka[5,19], Yoshinori Matsumoto[4], Kazuhiko Yamamoto[15],

Kazuyoshi Ishigaki[1,2,20]*

Affiliations

[1] Laboratory for Human Immunogenetics, RIKEN Center for Integrative Medical Sciences, Yokohama,

Japan

[2] Department of Microbiology and Immunology, Keio University School of Medicine, Tokyo, Japan

[3] Department of Ophthalmology, Graduate School of Medical Sciences, Kyushu University, Fukuoka,

Japan

[4] Department of Nephrology, Rheumatology, Endocrinology and Metabolism, Okayama University

Faculty of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan

[5] The First Department of Internal Medicine, University of Occupational and Environmental Health,

Kitakyushu, Japan

[6] Faculty of Health Science, Okayama University Medical School, Graduate School of Health Sciences,

Okayama University, Okayama, Japan

[7] Division of Rheumatology, Department of Internal Medicine, Keio University School of Medicine, Tokyo,

Japan

[8] Department of Dermatology, Keio University School of Medicine, Tokyo, Japan

[9] Department of Dermatology, the First Affiliated Hospital of Anhui Medical University, Hefei, Anhui,

China

[10] Institute of Dermatology, Anhui Medical University, Hefei, Anhui, China

[11] Key Laboratory of Dermatology (Anhui Medical University), Ministry of Education, Hefei, Anhui, China

[12] Department of Rheumatology, Hanyang University Hospital for Rheumatic Diseases, Seoul, Korea

[13] Hanyang Institute of Bioscience and Biotechnology, Seoul, Korea

[14] Hanyang University Institute for Rheumatology Research, Seoul, Korea

[15] Laboratory for Autoimmune Diseases, RIKEN Center for Integrative Medical Sciences, Yokohama,

Japan

[16] Laboratory for Statistical and Translational Genetics, RIKEN Center for Integrative Medical Sciences,

Yokohama, Japan

[17] Clinical Research Center, Shizuoka General Hospital, Shizuoka, Japan

[18] The Department of Applied Genetics, The School of Pharmaceutical Sciences, University of Shizuoka,

Shizuoka, Japan

[19] Department of Molecular Targeted Therapeutics, University of Occupational and Environmental Health,

Kitakyushu, Japan

[20] Keio University Human Biology-Microbiome-Quantum Research Center (WPI-Bio2Q)

[21] These authors contributed equally.

45
46
47
48
49   *Correspondence:

50   Masahiro Nakano, M.D., Ph.D.

51   masahiro.nakano.fc@riken.jp

52   Kazuyoshi Ishigaki, M.D., Ph.D.

53   kazuyoshi.ishigaki@keio.jp

54
55

Abstract

Systemic lupus erythematosus (SLE) is a complex autoimmune disease with unknown etiology. To

pinpoint new disease-relevant cell states in SLE and their detailed molecular profiles, we applied the

deepest investigation strategy for single-cell data, cellular and molecular fine mapping, to the largest-

scale multimodal SLE datasets, comprising ~2.1 million peripheral blood mononuclear cells from 232

SLE cases and 114 healthy controls. Among 123 fine-grained cell states in 27 cell types, we discovered

that previously uncharacterized cell states, such as $GZMK^+GZMH^+HLA\text{-}DR^+$ effector memory CD8$^+$ T

cells (double-positive [DP] EMCD8), $FAM13A^+CDK6^+$ naive CD4$^+$ T cells, and $ARHGAP15^+FOXO1^+$ T

cells, expanded especially in severe SLE. Based on our new statistical model, which efficiently dissects

gene signatures reflecting cell state abundance (quantitative change) and single-cell-level gene

dysregulation (qualitative change) within each conventional cell type, we found that key biological

processes and genetic risks of SLE were linked to these expanded cell states. Finally, by combining

multimodal in-silico analyses, CRISPR perturbations, and cellular function assays, we identified cell-

state-specific transcriptional regulators (e.g., $FOXO1$ in $ARHGAP15^+$ $FOXO1^+$ T cells), key surface

proteins (e.g., HLA-DR, CD27, CD38, and CD28 in DP EMCD8), T cell receptor features, and cis-

regulatory elements. Notably, we newly revealed that DP EMCD8 cells exhibit chronic antigen exposure

signatures and serve as a major source of key cytokines, such as IFN-γ, underscoring this cell state as a

73   central driver of SLE immunopathology. These findings provide new insights into the drug target

74   discovery in SLE.

Introduction

Systemic lupus erythematosus (SLE) is a prototypic autoimmune disease that affects multiple organs,

triggered by genetic and environmental factors[1]. Since the pathophysiology of SLE involves various

immune cell types, many bulk and single-cell transcriptome studies on peripheral blood and affected

organs have been conducted[2–6]. While these efforts have begun to reveal key gene signatures and core

cell populations of SLE[7,8], the fine-resolution pathophysiology remains unresolved, and the treatment

options are still limited; to date, only two molecular-targeted drugs have been approved for this

disease[9,10].

To understand the complex biology of SLE and discover new therapeutic targets, it is crucial to

identify new disease-relevant cell states and profile their key molecules at fine cellular resolution.

Recently, we reported the largest-scale bulk study of SLE with 27 immune cell types and identified cell-

type-specific signatures linked to disease development and exacerbation[6]. However, as an inherent

limitation of bulk studies, we failed to identify granular disease-relevant cell states that may exist within

each cell type. While single-cell RNA sequencing (scRNA-seq) can address this problem, most studies

have not fully leveraged the advantage of single-cell data. First, although scRNA-seq can, in theory,

define hundreds of distinct and heterogeneous cell states, previous studies ended up roughly profiling

only ~20 major cell types[5] or ~40 cell clusters[3,4] and did not pursue more granular and detailed biology.

Second, although scRNA-seq can evaluate quantitative and qualitative changes in disease status at

single-cell resolution, most studies rely on the cell-type-level pseudobulk approach, which aggregates

single-cell expression across each cell type and treats them as if bulk expression data. As a result, this

approach obscures the distinction between altered cell state composition (quantitative changes) and

dysregulated gene expression within individual cells (qualitative changes)[11], both of which may require

different treatment strategies. These critical limitations have hindered us from the comprehensive

identification of disease-relevant cell states and their functional characterization.

To tackle these issues, we took the deepest investigation strategy, cellular and molecular fine

mapping, to pinpoint granular disease-relevant cell states and their key molecules at single-cell

resolution and applied it to the largest-scale multimodal single-cell data of SLE with a total of ~2.1 million

cells from 346 donors (Lupus-derived Peripheral immune cell Integrated panels [LuPIN]; **Fig. 1, left**).

First, by employing a fine-grained clustering approach, we identified 123 distinct cell states in 27 immune

cell types. Among them, we found 11 previously uncharacterized cell states expanded especially in

clinically active SLE ("cellular fine mapping", **Fig. 1, middle**). Next, our new statistical model, which

dissected gene signatures reflecting quantitative and qualitative changes, demonstrated that key

immunological pathways and SLE risk variants were closely linked to cell-state compositional changes

within each cell type. Finally, through multimodal in-silico analyses, CRISPR perturbations, and cellular

function assays of disease-relevant cell state candidates, we successfully revealed their key molecular

profiles, such as transcriptional regulators, cytokine profiles, surface marker proteins, T cell receptor

(TCR) repertoires, and epigenetic status ("molecular fine mapping", **Fig. 1, right**). Overall, our intensive

and comprehensive cellular and molecular fine mapping shed light on the new immunopathology of SLE

and provided guidance to design new treatment strategies.

Results

Overview of this study

LuPIN consists of three datasets, each with a different purpose. To robustly identify disease-relevant cell

states, we first used the largest, publicly available 3'-end scRNA-seq dataset from SLE[5] (dataset 1; **Fig.**

1, left and **Supplementary Table 1**). Our stringent multi-step quality control (QC) process yielded the

final dataset (1,456,041 peripheral blood mononuclear cells [PBMCs] from 97 healthy and 172 SLE

samples [159 unique SLE donors: 11 donors have 2~3 timepoints]; **Extended Data Fig. 1** and

Methods). Among 172 SLE samples, 21 had high disease activity (HDA), and 40 were clinically inactive

(**Methods**). Following our previous study scheme[6], we prepared two comparison axes: (i) disease-state

comparison between healthy controls and inactive SLE patients, reflecting the biology of disease

development, and (ii) disease-activity comparison between inactive vs. HDA SLE, reflecting the biology

of disease exacerbation. For replication analysis, we used another scRNA-seq dataset[4] (dataset 2,

344,235 PBMCs from 17 healthy and 41 SLE samples).

To overlay in-depth molecular information onto the fine-grained cell state map, we additionally

prepared a new multimodal 5'-end dataset with cellular indexing of transcriptomes and epitopes by

sequencing (CITE-seq), a technology profiling gene expression and surface marker proteins from single

cells[12], as well as TCR-seq and enhancer RNA profiling[13,14] (dataset 3, 268,783 PBMCs from 32 SLE

samples; **Methods**). Through the efficient integration with dataset 1, we searched for key surface

proteins, TCR repertoires, and cis-regulatory elements specific to disease-relevant cell states.

Identification of 123 distinct cell states using fine-grained clustering

To identify granular cell states of SLE, we employed a three-step intensive clustering approach in

dataset 1: lineage (layer 1), cell type (layer 2), and cell state (layer 3) (**Fig. 2**). First, by applying Louvain

clustering, we separated 1,456,041 PBMCs into seven cell lineages (layer 1: CD4$^+$ T cells [CD4 T], CD8$^+$

and unconventional T cells [CD8 and other T], natural killer cells [NK], B cells, monocytes [Mono],

dendritic cells [DC], and hematopoietic stem and progenitor cells [HSPC]; **Extended Data Fig. 1d-f**). We

then separated these seven lineages into 27 coarse cell types (layer 2: **Fig. 2a-f**, **Extended Data Fig.**

2a-f, and **Supplementary Table 2**). Our elaborated analytical pipelines (e.g., curated clustering

algorithms; **Supplementary Table 3** and **Methods**) achieved higher cell-type annotation accuracy than

the original study[5] (**Extended Data Fig. 2g**).

While the original study defined only 20 coarse cell types, we further performed clustering on

each cell type (cell-type-level clustering) and identified 123 cell states with unique gene expression

profiles (layer 3: **Fig. 2a-f**, **Extended Data Fig. 3**, and **Supplementary Table 4**). Compared with

conventional lineage-level clustering, which primarily captures global cell-type structures (e.g., naive-

ness vs. memory-ness), cell-type-level clustering can efficiently capture subtle gene-program

heterogeneities within each cell type. As these cell-state definitions form the basis for all downstream

analyses, we performed rigorous approaches to ensure their robustness. First, we set the conservative

clustering resolution parameter for each cell type so that the distinct identity of each cluster was

supported by a wide range of literatures[7,8,15–38] (**Supplementary Table 5**). Indeed, upregulated marker

genes for each state showed minimal overlap within the same cell type (**Fig. 2g**), indicating that these

clusters were not redundant. Second, pathway analysis using these marker genes showed enrichment of

multiple cell-state-specific pathways (**Supplementary Table 6**), supporting their biological relevance.

Finally, our multimodal investigations using the independent dataset 3 identified cell-state-specific

surface proteins and TCR features (**Fig. 5-6**), providing orthogonal validation at the immune phenotypic

level.

Our fine-grained annotations successfully captured multiple disease-relevant cell state

candidates (**Fig. 2**), most of which were not reported in the original study due to insufficient resolution.

We first recapitulated well-known disease-relevant cell states such as (1) follicular and peripheral helper

T cells (Tfh and Tph) in central and effector memory CD4 T cells (CMCD4-3: $ICOS^+PASK^+$ and EMCD4-

7: $PDCD1^+TIGIT^+$) and (2) autoimmune-associated B cells (ABC) in memory B cells (MemB-6:

$ITGAX^+FCRL5^+$), which were important in the autoantibody production[7,8,23]. We also confirmed the

subpopulations highly activated by IFN signaling ($ISG15^+MX1^+$ cell states), the hallmark signature of

SLE[1,22], in most cell types. The shared gene program across cell types is not unique to IFN signaling;

JUN^+CD69^+ cell states with upregulated AP-1 family TFs (e.g., *JUN, FOS*) and early activation markers

(e.g., *CD69* and *NFKBIA*)[17] were identified in most lymphoid cell types (**Extended Data Fig. 3**).

Importantly, we identified multiple previously uncharacterized cell states with distinct gene programs:

some of them were later shown to expand especially in severe SLE (e.g., EMCD8-5: $GZMK^+GZMH^+HLA$-

$DRB1^+$, **Fig. 3**). Moreover, our approach captured very rare subpopulations such as $ANXA2^+CTLA4^+$

activated double negative T cells (dnT-1)[31], $RORA^+GATA3^+$ group 2 innate lymphoid cells (ILC-1)[32],

$MZB1^+PCNA^+$ plasmablasts (PlasmaB-1)[34], and $LAMP3^+CCR7^+$ DCs (cDC-5)[38]. Collectively, our three-

step intensive clustering approach yielded the most granular cell state information for SLE, serving as

the foundation for subsequent analyses.

Cellular fine mapping to pinpoint disease-relevant cell states of SLE

To evaluate which cell states expanded in SLE (cellular fine mapping), we next performed co-varying

neighborhood analysis (CNA), an approach to identify single-cell-resolution transcriptional

neighborhoods associated with clinical phenotypes[39]. The extent of expansion (or depletion) of each cell

state was expressed as continuous values at the single-cell level ("neighborhood correlation", **Fig. 3a-b**).

As mentioned above, we performed CNA with two comparison axes: disease-state (inactive SLE vs.

healthy, reflecting disease development) and disease-activity CNA (HDA vs. inactive SLE, reflecting

disease exacerbation)[6].

We first examined the abundance of 26 coarse cell types across six cell lineages (excluding

HSPC), accounting for age, sex, and ancestry (lineage-level CNA; **Fig. 3a** and **Methods**). Our analysis

recapitulated the depletion of NaiveCD4 and mucosal-associated invariant T cells (MAIT) in disease-

state comparison[5] and newly showed that these depletion patterns were remarkable in disease-activity

comparison (**Fig. 3a** and **Extended Data Fig. 4a**). Similarly, regulatory T cells (Treg) and Proliferating

(Prolif) CD4/CD8/NK cells expanded both in disease-state and -activity CNA. In Mono-lineage,

monocytes with high C1q genes[36] (C1Q$^+$Mono) expanded in disease-activity CNA. Although

neighborhoods in these cell types exhibited universal expansion (or depletion) patterns, many other cell

types showed heterogeneous patterns in lineage-level CNA (**Fig. 3a**), underscoring the need for a more

fine-grained investigation.

To assess the cell state compositional changes within each cell type, we next applied CNA to 18

major cell types (>5,000 cells; cell-type-level CNA). Strikingly, we observed skewed neighborhood

abundance in 34 out of 36 tests (18 cell types × 2 comparisons, global $P < 0.05/36$), and 29 cell states

showed evidence of remarkable expansion (**Fig. 3b**, **Extended Data Fig. 4b**, and **Supplementary Table**

7). These expanded cell states tended to be rare within each cell type, supporting the importance of our

fine-grained strategy (**Fig. 3c** and **Extended Data Fig. 4c**).

203   Among the 29 cell states with remarkable expansion,18 recapitulated known signals. All IFN-

activated (*ISG15*⁺*MX1*⁺) cell states within each cell type expanded in disease-state and -activity CNA

(**Fig. 3d**), supporting our previous report that IFN signal contributes to both disease development and

exacerbation[6]. While Tfh (CMCD4-3) expanded primarily in disease-state, Tph (EMCD4-7) and ABC

(MemB-6) expanded especially in disease-activity CNA. The remaining 11 cell states represented

previously unreported expansions in SLE[3–5], nominating new disease-relevant cell state candidates. An

example is *GZMK*⁺*GZMH*⁺*HLA-DRB1*⁺ cells (EMCD8-5), showing evidence of expansion both in disease-

state and -activity CNA (**Fig. 3b**). While other EMCD8 cell states express either *GZMK* or *GZMH*, this

population expresses both, as well as *GZMB* to a lesser extent (**Extended Data Fig. 3b**; hereafter, we

refer to this cell state as "double-positive [DP] EMCD8"). Notably, this population showed high

expression of multiple T cell activation markers, including *HLA-DR* genes, *PDCD1*, and *TIGIT* (**Extended**

Data Fig. 3b). Another example is *FAM13A*⁺*CDK6*⁺ cells (NaiveCD4-4), which further highlights the

significance of our fine-grained analysis: although NaiveCD4 was on average depleted within the CD4 T

lineage (**Fig. 3a**), this depletion pattern is not uniform across NaiveCD4 cell states. Lastly,

ARHGAP15[+]*FOXO1*[+] cell states also expanded across most T-cell cell types in disease-state and -

activity CNA (**Fig. 3b, d**).

Some cell states exhibited different patterns of abundance changes between disease-state and

activity CNA. While *JUN*[+]*CD69*[+] cell states expanded in disease-state CNA, they consistently depleted in

disease-activity CNA in most cell types (**Fig. 3d**). These results suggest distinct roles of these cell states

in SLE depending on the disease phase[6]. Our extensive analysis also identified ancestry-specific signals:

some *ISG15*[+]*MX1*[+] cell states and ABC (MemB-6) expanded in Asian SLE compared to European (EUR)

SLE[8,40] (**Extended Data Fig. 4d**).

Our CNA results showed high reproducibility in multiple approaches. First, a cluster-based

abundance testing (MASC)[26] showed concordant results with CNA (**Extended Data Fig. 4e**). Second,

most CNA signals were globally concordant between Asian and EUR ancestries (**Extended Data Fig.**

4f). Third, by applying CNA independently to low disease activity (LDA) and moderate disease activity

(MDA) cases that were excluded from the discovery set, we observed the concordant signals with a

gradual increase in disease-activity effects, reflecting the continuous nature of SLE activity (**Fig. 3e** and

Methods). Lastly, disease-state and activity CNA using the independent dataset 2 showed good

replicability of CNA signals detected in dataset 1 (**Fig. 3f** and **Methods**). Overall, our intensive cellular

fine mapping has robustly pinpointed novel disease-relevant cell-state candidates.

A new single-cell model to dissect quantitative and qualitative changes within cell types

Following the successful cellular fine mapping, we proceed to the molecular fine mapping. As a

stepping stone toward pinpointing key molecules in SLE, we introduce a new statistical model to

overcome limitations of the pseudobulk analyses, a standard approach in many studies. Since a typical

pseudobulk analysis aggregates single-cell-level expression within a cell type, the resulting differentially

expressed genes (DEGs) could be mixed up by the following two scenarios **(Fig. 4a)**: (i) a gene is

specifically expressed in a particular cell state whose abundance shifts with phenotype (quantitative

change) or (ii) a gene's per-cell expression is directly dysregulated with phenotype regardless of cell

state abundance (qualitative change). The pseudobulk approach cannot determine which scenario has a

more pathogenic effect on disease biology.

To address this issue, we developed ScQuaD (Single-cell-level Quantitative and Qualitative

change Dissection), a generalized linear mixed effect model to evaluate the association of gene

expression levels with both quantitative and qualitative changes (**Fig. 4b**). In contrast to pseudobulk

analysis, ScQuaD treats each single-cell-level expression as an independent observation based on

negative binomial distribution and regresses this jointly by (i) a phenotype-driven cell-state abundance

term (quantitative change) and (ii) a direct phenotype term (e.g., inactive SLE vs. healthy) capturing per-

cell gene dysregulation (qualitative change), adjusted for donor-/batch-random effect and other fixed

effect covariates (**Methods**). Specifically, we incorporated single-cell-level neighborhood correlation

values in cell-type-level CNA (**Fig. 3**) into ScQuaD as metrics for cell-state abundance changes. We

applied this model to each gene across 18 major cell types. We termed (i) the genes associated with

quantitative changes as "abundance signature genes" and (ii) the genes directly associated with

qualitative changes, even after adjusting for quantitative changes, as "dysregulated signature genes"

(**Supplementary Table 8**). Permutation tests showed that this model is well calibrated (**Extended Data**

Fig. 5a). When we quantified the contribution of qualitative and quantitative changes to the top 1,000

pseudobulk DEGs for each cell type, around 50% of their variation was driven by quantitative change

(**Extended Data Fig. 5b**). For simplicity, we showed disease-state comparison results in **Fig. 4c-f** and

disease-activity results in **Extended Data Fig. 5c-d**.

IFIH1 and *OAS1* were well-known interferon response genes (IRGs), and both are linked to the

significant genetic loci in genome-wide association studies of SLE[41] (SLE-GWAS; **Supplementary Table**

9 and **Methods**). While pseudobulk analysis identified both as DEGs in NaiveCD4 (false discovery rate

[FDR] < 0.05), ScQuaD revealed different behaviors of qualitative and quantitative changes for these two

genes (**Fig. 4c**). While *OAS1* expression localized to NaiveCD4-7 (*ISG15*[+]*MX1*[+]), which expanded in

SLE (abundance signature gene), *IFIH1* did not show such localization and was globally overexpressed

in inactive SLE compared with healthy cells (dysregulated signature gene). Hierarchical clustering of 100

IRGs (**Supplementary Table 10**) based on ScQuaD model statistics across 18 cell types identified a

module representing abundance signature genes and another representing dysregulated signature

genes, suggesting the heterogeneous roles of IRGs on SLE pathophysiology (**Fig. 4d** and **Extended**

Data Fig. 5c).

Next, we performed pathway enrichment analyses on these signatures for each cell type (**Fig.**

4e). We observed enrichment of IFN signaling in both dysregulated and abundance signatures across

almost all cell types (FDR < 0.05; one-sided Fisher's exact test), demonstrating that IFN signaling exerts

its pathogenic effects through both quantitative and qualitative changes. However, other classical

immunological pathways were primarily enriched in abundance signature genes of relevant cell types

(e.g., TCR signaling in T cells and complement signaling in myeloid cells; **Fig. 4e**, **Extended Data Fig.**

5d, and **Supplementary Table 11**), suggesting that key biological processes in SLE were more linked to

abundance signatures rather than dysregulated signatures.

To estimate the causal roles of these two signatures in SLE, we integrated ScQuaD statistics

with the largest-scale SLE-GWASs in East Asian (EAS)[42] and EUR[43]. We tested the polygenic

enrichment of SLE risk variants around the signature genes using stratified linkage disequilibrium score

regression (S-LDSC)[44] for EAS- and EUR-GWAS and meta-analyzed them **(Extended Data Fig. 5e)**.

While SLE risk variants were enriched around both dysregulated and abundance signature genes in

multiple cell types (FDR < 0.05), the extent of enrichment was more remarkable in abundance signature

genes ($P = 5.3 \times 10^{-4}$; Paired Wilcoxon test; **Fig. 4f**). Notably, this enrichment was observed in the

abundance signature genes with the positive statistics, but not in those with the negative statistics

(**Extended Data Fig. 5f**). Therefore, our model demonstrated that cell state compositional changes

within a cell type (especially those of expanded cell states) can exert causal effects on SLE

pathogenesis, rather than reflecting the secondary effects during the chronic disease phase.

Fine mapping of multimodal molecular profiles across cell states

Having delineated cell state abundance and gene programs linked to SLE pathophysiology, we

next sought to profile these cell states in greater detail. To this end, we prepared a new multimodal

dataset with CITE-seq (130 surface marker proteins with gene expression), TCR-seq, and enhancer

RNA profiling (dataset 3; 32 SLE donors). Reference mapping[45] successfully integrated 268,783 PBMCs

from dataset 3 into the latent transcriptional space in dataset 1, enabling us to extend our fine-grained

strategy to the multimodal layers (**Fig. 5a** and **Methods**). First, we comprehensively investigated cell-

state-specific surface proteins and identified, on average, nine upregulated proteins per cell state (FDR <

0.05; **Extended Data Fig. 6a** and **Supplementary Table 12**). We recapitulated the well-known surface

proteins in some cell states: (1) CD278 (ICOS), CD279 (PD1), and TIGIT in Tph (EMCD4-7)[7], (2) CD11c

and CD19 in ABC (MemB-6)[8], and (3) CD38 and CD31 in *SOX4*+ naive T cell states (NaiveCD4-3 and

NaiveCD8-3)[18]. Furthermore, we successfully identified several key proteins of new disease-relevant cell

states, which were utilized in the subsequent investigation (**Fig. 5-6**).

We next investigated TCR repertoire profiles. While we and others have shown that V gene

usage differs across cell types[46,47] and could be a new therapeutic target for autoimmunity[48], a

systematic cell-state-level analysis has been lacking. Principal component analysis (PCA) of V gene

frequency for each cell state confirmed that embeddings globally separated cell types (**Extended Data**

Fig. 6b). We next compared V gene usage across cell states. Notably, out of 94 V genes (45 α- and 49

β-chain), ~3.8 V genes per cell state showed skewed usage (FDR < 0.05; two-sided Fisher's exact test;

Extended Data Fig. 6c and **Supplementary Table 13**), highlighting distinct TCR signatures at the cell-

state level. While most of them were observed in EMCD8 cell states, we successfully identified several

cell-state-specific V genes in other cell types (e.g., NaiveCD4-5; **Fig. 5**).

Leveraging 5' transcript sequencing, we next mapped transcription start sites (TSSs) at single-

cell resolution. Recent studies have shown that bidirectional transcription around TSSs marks enhancer

activity at the locus[13,14]. Using this strategy, we called ~1,564 transcribed enhancers per cell type

(**Methods**), which were enriched in PBMC enhancer marks than others[49] (**Extended Data Fig. 6d**).

Among them, we identified ~68 cell-state-specific enhancers per cell type (**Supplementary Table 14**).

Thus, the efficient integration of our multimodal data with large-scale scRNA-seq data

successfully provided comprehensive molecular information on fine-grained cell states.

FOXO1 **is a candidate driver of the pathogenic CD4+ T cell state gene program**

We next advanced molecular fine-mapping to investigate disease-relevant cell states. Motivated

by our findings that key signal pathways and SLE causal variants were more enriched in abundance

signatures than dysregulated signatures (**Fig. 4e-f**), we focused subsequent analyses on several new

cell states that expanded in both disease-state and activity CNA (**Fig. 3b**).

Although some expanded CD4$^+$ T cell states such as Tph (EMCD4-7) and *ISG15$^+$MX1$^+$* cell

states have been well-characterized, the molecular features of our newly identified cell states remain

unclear: (1) *FAM13A$^+$CDK6$^+$* cells (NaiveCD4-4) in NaiveCD4, and (2) *ARHGAP15$^+$FOXO1$^+$* cell states in

multiple T cell types (e.g., NaiveCD4-5 and CMCD4-4, **Fig. 3b**). Pathway analyses of upregulated

marker genes showed the enrichment of JAK-STAT signaling for *FAM13A$^+$CDK6$^+$* cells and TCR

signaling for *ARHGAP15$^+$FOXO1$^+$* cells, demonstrating the distinct immunological roles for these cell

states (**Fig. 5b** and **Supplementary Table 6**). We observed a unique TCR V gene usage pattern in

ARHGAP15$^+$FOXO1$^+$ cells (NaiveCD4-5; **Fig. 5c** and **Supplementary Table 13**). We also identified cell-

state-specific enhancers near the marker genes, suggesting unique epigenetic regulations (**Extended**

Data Fig. 7a and **Supplementary Table 14**).

We next examined the localization of GWAS-prioritized genes for each cell state. In

FAM13A$^+$CDK6$^+$ cells, *ARID5B*, a transcription coactivator, was linked to one SLE-GWAS locus[41] (**Fig.**

5d and **Supplementary Table 9**). In *ARHGAP15$^+$FOXO1$^+$* cells, *ARHGAP15*, encoding a GTPase-

activating protein, was linked to another SLE-GWAS locus. Furthermore, we identified several key

surface proteins associated with these cell states. CD127 (IL-7 receptor α) was upregulated in

FAM13A⁺CDK6⁺ cells (**Fig. 5e**), reflecting the promotion of their differentiation[50]. On the other hand, a

signaling lymphocyte activation molecule [SLAM] family protein CD244 was consistently upregulated in

multiple *ARHGAP15⁺FOXO1⁺* cell states (NaiveCD4-5, CMCD4-4, and NaiveCD8-4; **Supplementary**

Table 12).

Finally, to identify transcriptional regulators driving the gene program of *FAM13A⁺CDK6⁺* and

ARHGAP15⁺FOXO1⁺ cells, we took an experimental approach. We selected 22 TFs upregulated in these

cells and knocked out (KO) each TF, using a CRISPR array screen in primary CD4 T cells from two

healthy donors (**Fig. 5f**, **Supplementary Table 15**). Genes downregulated by *FOXO1*-KO were closest

to *ARHGAP15⁺FOXO1⁺* upregulated signature genes (Bonferroni corrected *P* < 0.05; one-sided Fisher's

exact test; **Fig. 5g** and **Supplementary Table 16**). A validation experiment from the 3rd donor showed

concordant results (**Extended Data Fig. 7b**). Indeed, *ARHGAP15⁺FOXO1⁺* signature genes showed

enrichment for FOXO signaling pathway (**Extended Data Fig. 7c** and **Supplementary Table 6**). When

focused on EMCD4 cells, *FOXO1* showed the highest expression in Tph (EMCD4-7; **Fig. 5h** and

Extended Data Fig. 3a). Consistently, trajectory analysis[51] on whole CD4⁺ T cells showed the

differentiation of *ARHGAP15⁺FOXO1⁺* cell states toward Tph (and ProlifCD4) rather than other effector

cells. Given that *FOXO1* has recently been reported to be responsible for T cell memory-ness promotion

and clinical efficacy in cancer immunotherapy[20,21], the inhibition of *FOXO1* could be a new treatment

option for autoimmunity by modulating the aberrant gene program in CD4 T cells. On the other hand, we

observed that the genes downregulated by *ARID5B*-KO did not account for the *FAM13A*$^+$*CDK6*$^+$ cell

state, indicating that *ARID5B* does not play a significant role in defining the gene program of this

population (**Extended Data Fig. 7d**). Together, our molecular fine mapping pinpointed key regulator

candidates in disease-relevant CD4 T cell states of SLE.

Multimodal molecular fine mapping on *GZMK*$^+$*GZMH*$^+$*HLA-DRB1*$^+$ EMCD8 cells

367   Among CD8$^+$ T cell states exhibiting remarkable expansions both in disease-state and activity

CNA, the most uncharacterized one is *GZMK*$^+$*GZMH*$^+$*HLA-DRB1*$^+$ EMCD8 cells (EMCD8-5, DP EMCD8).

Among upregulated signature genes in DP EMCD8, *RGS1*, known as a chemokine signaling regulator

and a cell exhaustion marker[52], was linked to one SLE-GWAS locus[41] (**Fig. 6a**). Indeed, we observed the

highest classical exhaustion signature scores (e.g., *PDCD1* and *TIGIT*, **Supplementary Table 17**) on

this cell state in EMCD8 (**Fig. 6b**). Notably, this pattern was comparable to Tph cells in EMCD4,

suggesting chronic antigen exposure to these cell states[53]. Consistently, DP EMCD8 showed evidence of

TCR clonal expansion in multiple SLE donors (**Fig. 6c**). Some of the expanded clonotypes overlapped

with those that reacted to epitopes of cytomegalovirus or Epstein-Barr virus (**Supplementary Tables**

18). While many of these clones were shared with other EMCD8 cell states, we observed 9 clonotypes

and 9 V genes specific to DP EMCD8 (FDR < 0.05; two-sided Fisher's exact test; **Extended Data Fig.**

8a-b, Supplementary Tables 13 and **18**). Pathway analyses showed the enrichment of oxidative

phosphorylation signaling in DP EMCD8 and Tph (**Extended Data Fig. 8c** and **Supplementary Table**

6), suggesting the shared metabolomic reprogramming between these two cell states. We also examined

the cytokine profile of DP EMCD8. Intriguingly, we found that this population served as a major source of

proinflammatory cytokines, including *IFNG*, *IL16*, *IL32*, *CCL5*, and *CD70* (**Fig. 6d**). Accumulating

evidence has suggested that *IFNG* (type II IFN) is mainly produced by CD8$^+$ T cells and plays a pivotal

role in SLE as well as type I IFN[54–56]. Thus, our fine mapping strategy successfully pinpointed the cell-

state origin of this key cytokine.

We next sought to identify key surface proteins of DP EMCD8. Integration of the multimodal

dataset 3 into dataset 1 through reference mapping showed that HLA-DR, CD27, CD38, and CD28 as

the most upregulated proteins in DP EMCD8 (**Fig. 6e** and **Supplementary Table 12**). Notably, CD45RA

was downregulated in DP EMCD8, suggesting this population does not overlap with terminally

differentiated effector memory (TEMRA) CD8$^+$ T cells[57]. An inhibitory marker, GPR56, was also

downregulated in DP EMCD8. In-silico gating on HLA-DR$^+$CD27$^+$CD38$^+$CD28$^+$ successfully achieved a

purity of >90% for DP EMCD8 (**Extended Data Fig. 8d**), enabling us to investigate this population using

flow cytometry (FCM) and fluorescence-activated cell sorting (FACS).

We first FACS-sorted target EMCD8 cells gated on CCR7$^-$HLA-DR$^+$CD27$^+$CD38$^+$CD28$^+$ from

purified CD8$^+$ T cells and performed bulk RNAseq and Assay for Transposase-Accessible Chromatin

with sequencing (ATAC-seq) (**Fig. 6f**, **Extended Data Fig. 8e**, and **Supplementary Table 19**). After

confirming that these sorted cells showed a highly comparable gene expression profile to the DP EMCD8

(**Extended Data Fig. 8f**), we identified 113 differentially accessible regions (DARs) in this population

compared with whole EMCD8 (FDR < 0.1; **Extended Data Fig. 8g** and **Supplementary Table 20**). Motif

analysis for these DARs revealed the enrichment of Fli1 and IRF motifs (**Fig. 6g**). Since there were no

DARs around the *IFNG* region in standard ATAC-seq, we enriched open chromatin peaks around this

region using UNIChro-seq[58] (**Methods** and **Supplementary Table 21**). Around 20-fold enrichment of

peak counts detected one DAR, suggesting the epigenetic dysregulation of *IFNG* in this population. (**Fig.**

6h).

405   In intracellular FCM, we confirmed that a larger proportion of HLA-DR$^+$CD27$^+$CD38$^+$CD28$^+$

406   EMCD8 cells co-express GZMK and GZMB at the single-cell level, compared with whole EMCD8

407   (**Extended Data Fig. 8h**). Importantly, these cells showed high PD1 and TIGIT expression even at the

408   protein level (**Fig. 6i** and **Extended Data Fig. 8h**). Upon stimulation with Phorbol myristate acetate

409   (PMA)/Ionomycin, this population exhibited higher IFN-γ expression than whole EMCD8.

410   Identification of cell-state-specific surface proteins could have significant implications for drug

target discovery. The current therapeutic agents in autoimmunity target ubiquitous surface markers,

which can deplete normal cells as well as autoreactive cells (e.g., CD19 and CD20 in B cells[59,60]).

Narrowing down the target to disease-relevant cell states based on their specific markers could provide

more efficient treatment options. Collectively, our extensive approaches revealed proinflammatory

molecular phenotypes of DP EMCD8, underscoring this population as a potential key player in SLE.

Discussion

In this study, we developed an analytical framework, cellular and molecular fine mapping, to

pinpoint disease-relevant cell states and molecules at single-cell resolution, applying it to ~2.1 million

cells from 346 donors. Fine-grained clustering resolved 123 cell states across 27 cell types, including 11

previously uncharacterized ones that expanded especially in severe SLE. Subsequent in-silico

multimodal analyses, CRISPR perturbations, and functional experiments successfully defined their key

transcriptional regulators, cytokines, TCR features, and surface proteins, guiding the efficient design of

therapeutics for clinical settings.

In cellular fine mapping, we noted an important caveat: to pinpoint pathogenic cell states, careful

fine-grained clustering is essential. The most disease-relevant cell states are often low-frequency

populations and are easily obscured by coarse clustering, consistent with other studies that successfully

identified rare pathogenic populations[7,8,24]. Based on the extensive literature review[7,8,15–38] and rigorous

validation, we reconstructed 123 cell states with distinct immunological features from the largest, publicly

available single-cell data[5] (~1.5 million PBMCs). While the recent single-cell studies on inflamed tissues

in autoimmune diseases have identified several critical parenchymal subpopulations[24,61], the resolution of

immune cells is limited (~60 immune cell states from 0.2~0.4 million cells). To benefit the research

community, we released the most fine-grained immune cell reference panels with multimodal features,

called LuPIN (**Methods**). Researchers can map their datasets onto our reference panels for cell-state

annotations, providing an opportunity to uncover new pathophysiology of multiple diseases.

In this study, we proposed a new statistical model to overcome the limitation of the conventional

cell-type-level pseudobulk approach. To pinpoint molecular therapeutic targets from single-cell data,

distinguishing quantitative and qualitative changes linked to disease is a critical step, since each could

play different roles in disease pathophysiology. Some researchers are tackling this issue with different

statistical concepts from ours[11]. While such approaches are expected to be the mainstream of next-

generation single-cell research, it must be noted that our model relies on the latent space defined by

current sparse single-cell gene expression. The distinction between abundance and dysregulated

signatures may be refined in the future as single-cell technology advances.

Despite rapid growth in single-cell datasets, biological insights have not kept pace with the rising

number of reported association signals. This trajectory mirrors early era of GWAS, when efforts were

focused primarily on variant discovery rather than mechanistic understanding[62]. GWAS ultimately

matured by shifting from simple association to fine-mapping of causal variants followed by functional

dissection[63] (the so-called *post-GWAS era*); similarly, single-cell studies should progress from lists of

coarse associations toward fine-mapping of pathogenic cell states and their key driver molecules (a *post-*

cell atlas era). Given the abundance of public datasets now available, we advocate reallocating

resources from generating ever more data to performing targeted functional characterizations.

This study has some limitations. First, the current study focused on the functional investigation

of three cell states in T cells. Further characterization of other disease-relevant population candidates will

deepen understanding of complex pathophysiology in SLE. Second, because this study focused on

PBMCs, we may have missed key tissue-resident populations. Our fine-mapping approach should next

be extended to inflamed tissues in multiple diseases. Third, our study design lacked in-vivo functional

validation of disease-relevant cell states.

Together, our intensive cellular and molecular fine mapping can contribute to a better

understanding of SLE pathophysiology and the discovery of new treatment targets.

Figure 1. Overview of this study. (left) We applied cellular and molecular fine mapping to the largest-scale multimodal single-cell datasets of SLE, comprising ~2.1 million PBMCs from 346 donors. **(middle)** Through fine-grained clustering, we first identified 123 distinct cell states in 27 immune cell types. Among them, we found previously uncharacterized cell states expanded especially in clinically active SLE. **(right)** Next, our new statistical model, which dissected gene signatures reflecting quantitative and qualitative changes linked to phenotypes, demonstrated that key pathways and SLE risk variants were linked to cell-state compositional changes within each cell type. Finally, through multimodal in-silico and experimental approaches, we successfully revealed key molecular profiles of these disease-relevant cell states. CITE-seq, cellular indexing of transcriptomes and epitopes by sequencing; TCR, T cell receptor; EMCD8, effector memory CD8[+] T cells; HDA, high disease activity; TF, transcription factor; GWAS, genome-wide association studies; eRNA, enhancer RNA; UNIChro-seq, unique molecular identifier counting of regional chromatin accessibility with sequencing; FCM, flowcytometry; FACS, fluorescence-activated cell sorting.

Figure 2. Identification of 123 distinct cell states using fine-grained clustering. (a-f) UMAP plots showing 27 cell types and 123 cell states identified by fine-grained clustering in **(a)** CD4$^+$ T cell (CD4 T), **(b)** CD8$^+$ and unconventional T cell (CD8/other T), **(c)** natural killer cell (NK), **(d)** B cell, **(e)** monocyte (Mono), and **(f)** dendritic cell (DC) lineages (dataset 1; N=256). Hematopoietic stem and progenitor cell (HSPC) lineage is not shown since it has only one cell type and one cell state. In **(a-f)**, large UMAPs represent lineage-level UMAPs, colored by cell types. Small UMAPs represent cell-type-level UMAPs, colored and numbered by cell states. For each panel, cell state annotations are listed on the right. **(g)** Heatmap showing the number of overlaps across 100 upregulated marker genes for each cell state in memory B cells (MemB). See **Supplementary Table 3** for the description of cell type abbreviations.

Figure 3. Cellular fine mapping to pinpoint disease-relevant cell states of SLE. (a-b) (left) Distributions of neighborhood correlations in disease-state or disease-activity comparisons **(a)** for each cell type in lineage-level-CNA and **(b)** for each cell state in cell-type-level-CNA (dataset 1; N=256). In **(b)**, 29 cell states with > 50% of neighborhoods positively associated with phenotypes at false discovery rate (FDR) < 0.05 are shown (See **Extended Data Fig. 4b** for details). **(right)** UMAP plots in representative lineages (or cell types). Cell types (or cell states) are annotated if > 50% (******) or > 25% (*****) of neighborhoods associated with phenotypes. **(c)** A scatter plot showing the association between median neighborhood correlation for each cell state in cell-type-level disease-state CNA and cell-state proportion over cell type. Shaded regions represent 95% confidence intervals. **(d)** Scatter plots comparing median neighborhood correlation for each cell state between cell-type-level disease-state (x-axis) and disease-activity CNA (y-axis). Cell states related to each panel title are colored. **(e)** A heatmap showing the median neighborhood correlation for each cell state for low disease activity (LDA) vs. Inactive SLE, moderate disease activity (MDA) vs. Inactive SLE, and high disease activity (HDA) vs. Inactive SLE comparisons. 32 cell states with > 50% of neighborhoods associated with disease-activity comparison are shown. **(f)** Scatter plots comparing median neighborhood correlation for each cell state between dataset 1 (x-axis) and dataset 2 (y-axis; N=58). In **(c)** and **(f)**, Pearson correlation coefficients (*r*) between x- and y-axis and their *P* values are shown.

Figure 4. A new single-cell model to dissect quantitative and qualitative changes within cell types. (a) Shema of two different scenarios (quantitative and qualitative changes) to drive cell-type-level pseudobulk DEGs (differentially expressed genes) in SLE. **(b)** Shema of a new statistical model (ScQuaD) to dissect quantitative and qualitative changes (dataset 1; N=256). **(c)** Representative examples of abundance (*OAS1*) and dysregulated (*IFIH1*) signature genes. In UMAPs, cells are colored in expression density. Model statistics for both genes are shown in the bottom. **(d)** Heatmap showing the ScQuaD model Z scores of abundance and dysregulated terms across 18 major cell types for 100 IFN-response genes. Gene orders are based on hierarchical clustering using Z scores across cell types. **(e)** Bar plots showing the enrichment of representative pathways for each signature for each cell type. *P*, *P* values in one-sided Fisher's exact test. **(f)** Scatter plots comparing the enrichment of SLE risk variants around abundance (x-axis) and dysregulated signature genes (y-axis) for each cell type in S-LDSC (stratified LD score regression).

Figure 5. *FOXO1* is a candidate driver of the pathogenic CD4⁺ T cell state gene program. (a) Shema of the integration of multimodal dataset 3 (query; N=32) into larger scRNA-seq dataset 1 (reference; N=256) through reference mapping. **(b)** Bar plots showing the enrichment of representative pathways for upregulated signature genes for each NaiveCD4 cell state. *P*, *P* values in one-sided Fisher's exact test. **(c)** A bar plot showing the percentage of two TCR V gene usages that are skewed in NaiveCD4-5 compared with other NaiveCD4 cells. *P*, *P* values in two-sided Fisher's exact test. **(d)** Heatmap showing the logFC of GWAS-prioritized genes between one vs. other NaiveCD4 cell states. Up to three genes with the highest logFC per cell state are shown. Top annotation color represents the median neighborhood correlation of each cell state in disease-state and activity CNA (**Fig. 3**). **(e)** A volcano plot showing the differential expression of surface proteins between NaiveCD4-4 vs. other NaiveCD4 cell states. *P*, *P* values in differential expression with FindMarker function (Seurat; **Methods**). **(f)** Experiment design of CRISPR-KO array screening to primary CD4⁺ T cells. 22 TFs upregulated in NaiveCD4-4 or NaiveCD4-5 are selected for array screening (**Methods**). **(g)** A bar plot showing the enrichment of downregulated genes by each KO condition for NaiveCD4-5 upregulated signature genes. **(h)** Whole CD4⁺ T cell UMAP plots, colored by **(left)** the focused cell states and other cell types, **(upper right)** *FOXO1* expression, and **(lower right)** pseudotime in trajectory analysis. Edges represent trajectory graph.

Figure 6. Multimodal molecular fine mapping on *GZMK⁺GZMH⁺HLA-DRB1⁺* EMCD8 cells. (a) Heatmap showing the logFC of GWAS-prioritized genes between one vs. other EMCD8 cell states (dataset 1; N=256). Top annotation color represents the median neighborhood correlation in CNA. **(b) (top)** EMCD8 and **(bottom)** EMCD4 UMAP plots, colored by exhaustion signature scores. **(c)** Bar plots showing **(top)** the number of EMCD8-5 cells for each donor and **(bottom)** the proportion of TCR clonotypes that are categorized into five clone-size groups in EMCD8-5 (dataset3; N=32; 23 donors with >10 cells are shown). **(d)** A dot plot showing the expressions of upregulated cytokines in EMCD8-5. Color represents the proportion of cells expressing each gene. **(e)** A volcano plot showing the differential expression of surface proteins between EMCD8-5 vs. other EMCD8 cells. *P*, *P* values in differential expression (**Methods**). **(f)** Experiment design of flowcytometry (FCM) and FACS on HLA-DR⁺CD27⁺CD38⁺CD28⁺ EMCD8 cells. **(g)** TF motif logos enriched for differentially accessible peaks in HLA-DR⁺CD27⁺CD38⁺CD28⁺ cells compared with whole EMCD8 cells. *P*, enrichment *P* values in HOMER. **(h)** ATAC-seq and UNIChro-seq peaks around *IFNG* region in whole EMCD8 cells and HLA-DR⁺CD27⁺CD38⁺CD28⁺ cells from a SLE donor. *P*, logistic regression *P* values (N=6; **Methods**). **(i) (left)** Boxplots comparing the mean fluorescence intensities (MFI) of PD1 and IFNG between HLA-DR⁺CD27⁺CD38⁺CD28⁺ and whole EMCD8 cells. Samples from the same donor are connected with lines. *P*, *P* values in linear mixed models (N=11). **(right)** A histogram comparing the IFNG FI between HLA-DR⁺CD27⁺CD38⁺CD28⁺ and whole EMCD8 cells from a SLE donor.

Extended Data Figure 1. Quality control of datasets. (a) A density plot showing the cell density based on the number of genes detected and the percentage of UMIs from mitochondrial genes (%Mito) in dataset 1 (N=256). **(b)** A UMAP plot showing PBMCs in dataset 3 (N=32), colored by the diagnosis of Souporcell. Due to the computational cost, dataset 3 cells are visualized instead of dataset 1. **(c)** A scatter plot showing donor-level pseudobulk gene expression of *XIST* and Y chromosome-related genes in dataset 1. Color represents the sex information from the metadata. **(d)-(f)** Flow chart to separate PBMCs into seven lineages in dataset 1 (**Methods**). In **(f)**, dot plots show the scaled expression levels of lineage or subcluster marker genes. Color represents the proportion of cells expressing each gene. **(g)** A boxplot comparing doublet scores in Scrublet between singlets and doublets detected in **(e)-(f)**. **(h)** Barplots showing the number of cells that passed each QC step in datasets 1, 2, and 3.

Extended Data Figure 2. Identification of 123 distinct cell states using fine-grained clustering. (a-f) Flow chart to separate lineage cells into cell types and cell states in **(a)** CD4+ T cell (CD4 T), **(b)** CD8+ and unconventional T cell (CD8/other T), **(c)** natural killer cell (NK), **(d)** B cell, **(e)** monocyte (Mono), and **(f)** dendritic cell (DC) lineages (dataset 1; N=256; **Methods**). Hematopoietic stem and progenitor cell (HSPC) lineage is not shown since it has only one cell type and one cell state. Dot plots show the scaled expression levels of cell type marker genes. Color represents the proportion of cells expressing each gene. **(g)** A box plot showing the concordance rates of cell-type-level annotation in the original and the current study with the predicted cell-type labels in the reference mapping onto the independent PBMC dataset (**Methods**).

Extended Data Figure 3. Identification of 123 distinct cell states using fine-grained clustering. Dot plots showing the scaled expression levels of cell type and cell state marker genes in **(a)** CD4⁺ T cell (CD4 T), **(b)** CD8⁺ and unconventional T cell (CD8/other T), **(c)** natural killer cell (NK), **(d)** B cell, **(e)** monocyte (Mono), and **(f)** dendritic cell (DC) lineages (dataset 1; N=256). Color represents the proportion of cells expressing each gene.

Extended Data Figure 4. Cellular fine mapping to pinpoint disease-relevant cell states of SLE. (a) A scatter plot comparing the median neighborhood correlation of each cell type between disease-state (x-axis) and activity (y-axis) comparisons in lineage-level CNA (dataset 1; N=256). **(b)** Distributions of neighborhood correlations for each cell state with disease-state and activity comparisons in cell-type-level-CNA. **(c)** A scatter plot showing the association between median neighborhood correlation of each cell state in cell-type-level disease-activity CNA and cell-state proportion over cell type. Shaded regions represent 95% confidence intervals. **(d)** Distributions of neighborhood correlations for each cell state with Asian vs. EUR SLE comparison in cell-type-level-CNA. 15 cell states with > 50% of neighborhoods associated with phenotype at false discovery rate (FDR) < 0.05 are shown. **(e)** A scatter plot comparing median neighborhood correlation in cell-type-level CNA and effect size in MASC for each cell state with disease-state and activity comparisons. **(f)** A scatter plot comparing median neighborhood correlation in cell-type-level CNA in European and cell-type-level CNA in Asian with disease-state and activity comparisons. In **(c)**, **(e)** and **(f)**, Pearson correlation coefficients (*r*) between x- and y-axis and their *P* values are shown.

Extended Data Figure 5. A new single-cell model to dissect quantitative and qualitative changes within cell types. (a) Quantile-quantile plots comparing the observed *P* values in abundance (red) and dysregulated (blue) terms in the ScQuaD model across all cell types and comparisons (dataset 1; N=256). **(b)** Relative proportion of the variance explained by abundance (red) and dysregulated (blue) terms for the union of top 1,000 P-value pseudobulk DEGs in each cell type and each comparison. **(c)** Heatmap showing the ScQuaD model Z scores of abundance and dysregulated terms across 18 major cell types (row) for 100 IFN-response genes (column) in disease-activity comparisons. Gene orders are based on hierarchical clustering using Z scores across cell types. **(d)** Bar plots showing the enrichment of representative pathways for each signature for each cell type with disease-state and activity comparisons. *P*, *P* values in one-sided Fisher's exact test. **(e)** Scatter plots comparing the normalized coefficients in S-LDSC (stratified LD score regression) for EAS and EUR SLE-GWAS. Each dot represents each signature across 18 major cell types. Error bars represent 2×standard errors. Pearson correlation coefficient (*r*) between them and *P* values are shown. **(f)** Scatter plots comparing the enrichment of SLE risk variants around abundance signature genes with positive (x-axis) and negative (y-axis) model statistics for each cell type in S-LDSC.

Extended Data Figure 6. Fine mapping of multimodal molecular profiles across cell states. (a) A bar plot showing the number of cell-state-specific upregulated surface marker proteins (dataset 3; N=32). Color represents cell state IDs for each cell state (e.g., NaiveCD4-0). **(b)** A PCA plot of TCR V gene usage frequency for each cell state, colored by cell types. **(c)** A bar plot showing the number of cell-state-specific TCR V genes. For each cell state, more frequently used V genes are counted. **(d)** A bar plot showing the enrichment of called transcribed enhancers across all cell types for Roadmap project PBMC enhancer annotation compared with (left) Brain enhancer and (right) PBMC heterochromatin.

Extended Data Figure 7. *FOXO1* is a candidate driver of the pathogenic CD4+ T cell state gene program.
(a) Representative examples of cell-state-specific transcribed enhancers near the upregulated marker genes in NaiveCD4-5 (dataset 3; N=32). **(b)** Scatter plots comparing the logFC by *FOXO1*-KO in donors 1-2 and donor 3 (replication) for each gene. Genes with concordant signs between donors 1-2 and donor 3 are colored in red. *P*, *P* values in one-sided binomial sign test. **(c)** A Bar plot showing the enrichment of Kyoto Encyclopedia of Genes and Genomes (KEGG) FoxO signaling pathway for upregulated signature genes for each NaiveCD4 cell state. *P*, *P* values in one-sided Fisher's exact test. **(d)** A bar plot showing the enrichment of downregulated genes by *ARID5B*-KO for NaiveCD4-4 upregulated signature genes.

Extended Data Figure 8. Multimodal molecular fine mapping on *GZMK⁺GZMH⁺HLA-DRB1⁺* EMCD8 cells. (a) Bar plots showing the percentage of **(left)** a representative clonotype that is shared across EMCD8 cell states and **(right)** one that is specific to EMCD8-5 (dataset 3; N=32). *P*, *P* values in two-sided Fisher's exact test. **(b)** Bar plots showing the percentage of **(left)** six TRBV and **(right)** three TRAV gene usages that are positively skewed in EMCD8-5 compared with other EMCD8 cells. *P*, *P* values in two-sided Fisher's exact test. **(c) (top)** EMCD8 and **(bottom)** EMCD4 UMAP plots, colored by oxidative phosphorylation signature scores. **(d)** A bar plot showing the percentage of EMCD8-5 cells on each in-silico gate using dataset 3. **(e)** Gating strategy to define HLA-DR⁺CD27⁺CD38⁺CD28⁺ EMCD8 cells in flowcytometry (FCM) and FACS. **(f)** Scatter plots comparing the scRNA-seq dataset 1 logFC (EMCD8-5 vs. other EMCD8 cells; N=256) and bulk RNA-seq logFC (HLA-DR⁺CD27⁺CD38⁺CD28⁺ vs. whole EMCD8 cells; N=6). **(g)** A volcano plot showing the differential peaks between HLA-DR⁺CD27⁺CD38⁺CD28⁺ vs. whole EMCD8 cells in bulk ATAC-seq (N=6). **(h) (left)** Boxplots comparing the mean fluorescence intensities (MFI) of TIGIT and the percentage of GZMK⁺GZMB⁺ (double-positive) cells between HLA-DR⁺CD27⁺CD38⁺CD28⁺ and whole EMCD8 cells. Samples from the same donor are connected with lines. *P*, *P* values in linear mixed models (N=11). **(right)** A FCM plot to show the percentage of double-positive cells in HLA-DR⁺CD27⁺CD38⁺CD28⁺ EMCD8 cells from a SLE donor.

Methods

Data availability

Single-cell count matrices of gene expression, surface protein, and TCR features in the CITE-seq

dataset will be available at the National Bioscience Database Center (NBDC) Human Database

(https://humandbs.dbcls.jp/en/) as of the date of publication. Public data used in this study are available

at https://cellxgene.cziscience.com/collections. Reference panels for each lineage and each cell type are

available on GitHub (https://github.com/MasahiroNakano-hub/LuPIN).

Code availability

Original codes used in this study are available on GitHub (https://github.com/MasahiroNakano-

hub/LuPIN).

Subjects

In this study, we utilized two publicly available 3'-end scRNA-seq datasets (datasets 1 and 2). For

dataset 1, we downloaded FASQ files from Chan Zuckerberg CELL by GENE

(https://cellxgene.cziscience.com/collections) and raw genotype data from dbGaP (phs002812.v1.p1)

with authorized access[5]. SLE samples were recruited from the California Lupus Epidemiology Study

(CLUES) cohort, and healthy samples were obtained from the CLUES and the ImmVar Consortium. For

dataset 2, we downloaded the count data from dbGaP (phs002048.v1.p1)[4]. For our new 5'-end dataset

with CITE-seq, TCR-seq, and enhancer RNA profiling (dataset 3) and additional FCM and FACS

experiments (**Fig. 6**), patients were recruited at Okayama University Hospital and Hospital of the

University of the Occupational and Environmental Health based on the approval by the Ethics

Committees of RIKEN Center for Integrative Medical Sciences (RIKEN−Y-2025-006, 008). Written

informed consent was obtained from all participants.

Across all datasets, we defined four disease-activity categories in SLE[64]: i) inactive as SLE

disease activity index (SLEDAI) = 0, ii) low disease activity (LDA) as $1 \leq SLEDAI \leq 4$, iii) moderate

disease activity (MDA) as $5 \leq SLEDAI \leq 9$, and iv) high disease activity (HDA) as $SLEDAI \geq 10$

(**Supplementary Table 1**).

Sample processing and quality control

To generate dataset 3, we processed single-cell suspensions of PBMCs from 32 SLE donors through the

10x Genomics Chromium. We prepared gene expression (GEX), antibody-derived tag (ADT), and TCR

(VDJ) libraries using the Chromium Next GEM Single Cell 5' Kit v2, Feature Barcode Kit, Human TCR

Amplification Kit (10x Genomics), and TotalSeq™-C Human Universal Cocktail v1.0 (Biolegend,

399905), following the manufacturer's protocol. This dataset comprises two experimental batches and

eight cell libraries. For each batch, we pooled cells from 16 samples into each of four independent

libraries. These libraries were sequenced on the Illumina NovaSeq 6000 to generate 150-base paired-

end reads.

In datasets 1 and 3, FASTQ files were processed using Cell Ranger v7.0.1 (10x Genomics) with

the developer's reference (refdata-gex-GRCh38-2020-A). We first filtered out the droplets with < 500

total unique molecular identifiers (UMIs) or those annotated as antibody aggregates. We next excluded

low-quality cells with a low number of genes (nGene), a high percentage of UMIs from mitochondrial

genes (% MT), or a low percentage of UMIs from ribosomal protein genes (% Ribo) based on their

distributions for each dataset (dataset 1: nGene < 300, %MT > 7.5%, %Ribo < 8.5%; dataset 2: nGene <

300, %MT > 8.5%, %Ribo < 8.5%; dataset 3: nGene < 500, %MT > 8.5%, %Ribo < 6%; **Extended Data**

Fig. 1a). We also excluded cells with a high percentage of UMIs from hemoglobin genes (> 0.9%, *HBB*,

HBA1, *HBA2*, *HBD*) or platelet genes (> 0.6%, *PF4*, *PPBP*, *CAVIN2*, *GNG11*).

In datasets 1 and 3, multiple samples were pooled in each library (dataset 1: 8-19 samples per

library, dataset 3: 16 samples per library). For dataset 1, we used Demuxalot, a reference-genotype-

based demultiplexing software[65], to demultiplex these samples. We first performed genotype imputation

for autosomal variants using SHAPEIT2[66] and Minimac3[67] after quality control (QC). Next, we ran the

Demultiplexer.learn_genotypes function for each library, using ~5 million variants with Minimac3 R2 >

0.95. We additionally demultiplexed three samples with missing genotype using Souporcell, a reference-

genotype-free demultiplexing software[68]. For dataset 3, we ran Souporcell using common variants (minor

allele frequency > 1%) from the 1000 Genomes Project (GRCh38) and identified 16 genotype clusters

per library. In this step, we identified and excluded putative doublets from different genotypes for each

dataset (**Extended Data Fig. 1b**). For dataset 1, we excluded two samples with < 200 cells recovered

and three samples that showed discordance between the sex estimated from pseudobulk expressions of

XIST and Y chromosome-related genes and the metadata sex information (**Extended Data Fig. 1c**). For

dataset 3, to link the Souporcell genotype clusters to distinct individuals, we performed multiplex PCR,

amplifying genomic regions that cover 20 common variants using genomic DNA from each sample, and

sequenced the libraries on the Illumina MiSeq. For each sample, we identified the corresponding

genotype cluster whose genotypes of these 20 variants matched those obtained from multiplex PCR.

To separate PBMCs into seven cell lineages (CD4 T, CD8/other T, NK, B, Mono, DC, and

HSPC), we performed Louvain clustering with multiple steps (**Extended Data Fig. 1d-f**). First, we

normalized and scaled gene expression of PBMCs for each batch using SCTransform[69] in R package

Seurat (v5.1.0)[70] with parameters vst.flavor = "v1", method = "glmGamPoi", and min_cells = 30 (low-

expression gene filter). We then integrated data from each batch using 3,000 shared highly variable

genes (HVGs) in the SelectIntegrationFeatures function, followed by PCA and batch correction using R

package Harmony (v1.2.3)[71] with parameter group.by.vars = c("library", "sample"). Mitochondrial genes,

Y chromosome-related genes, hemoglobin genes, platelet genes, and cell-cycle genes were excluded

from HVGs. We used 30 harmony-corrected PCs for UMAP visualization and shared nearest neighbor

(SNN) graph construction. Next, we performed Louvain clustering based on the SNN graph and

separated PBMCs into four large cell lineages (T/NK, B, Myeloid, and HSPC, **Extended Data Fig. 1d**).

We then repeated the above procedures for each of T/NK, B, and Myeloid cells and identified 12, 7, and

3 clusters, respectively (**Extended Data Fig. 1e**). Since most of these clusters were mixed by multiple

cell lineages, we further performed sub-clustering for each cluster (**Extended Data Fig. 1f**). We

annotated each sub-cluster as relevant cell lineage based on their lineage marker gene expression. For

some clusters, we iterated this step to maximize the purity of each cell lineage. Detailed

hyperparameters for each step were summarized in **Supplementary Table 3**. During this step, we

identified several sub-clusters representing remaining doublets (even after the exclusion of putative

doublets in Demuxalot or Souporcell), which showed high expression of marker genes from multiple

lineages (**Extended Data Fig. 1f**) and higher doublet scores in Python package Scrublet[72] (**Extended**

Data Fig. 1g), and excluded them from the final dataset.

545   Through these multi-step QCs, we finally kept 1,456,041 PBMCs (159 SLE and 97 healthy

donors) for dataset 1, 344,235 PBMCs (41 SLE and 17 healthy) for dataset 2, and 268,783 PBMCs (32

SLE) for dataset 3 (**Extended Data Fig. 1h**).

Three-step clustering to define fine-grained 123 cell states

To identify fine-grained cell states within each cell type, we employed a three-step intensive clustering

approach in dataset 1: lineage (layer 1), cell type (layer 2), and cell state (layer 3). After separating the

PBMCs into seven cell lineages in the QC step, we again performed SCTransform, HVGs selection,

PCA, Harmony batch correction, and Louvain clustering for each lineage with optimal parameters

(**Extended Data Fig. 2a-f, left** and **Supplementary Table 3**). Since some of the initial coarse clusters

were mixed by multiple cell types, we further performed sub-clustering for each cluster. We then

annotated each sub-cluster as a relevant cell type based on their cell type marker gene expression

(**Extended Data Fig. 2a-f, middle**). Finally, we repeated the above procedures for each cell type using

optimal parameters (**Extended Data Fig. 2a-f, right** and **Supplementary Table 3**) and identified a total

of 123 cell states. To evaluate the cell-type annotation accuracy in this study and the original study, we

performed reference mapping of dataset 1 onto the independent PBMC dataset (Azimuth)[15]. We then

calculated the concordance rates of our annotation and the original annotation with the predicted cell-

type labels in the reference mapping for each cell type (**Extended Data Fig. 2g**).

To identify cell-state-specific marker genes, we ran the FindMarkers function in Seurat (v4.3.0)[15]

with parameter min.pct = 0.05, comparing one cell state vs. other cell states within the same cell type

(**Extended Data Fig. 3** and **Supplementary Table 4**). We used cell-state-specific upregulated marker

genes (FDR < 0.05 and logFC > 0.1) for pathway enrichment analyses (**Pathway enrichment analysis**

and gene signature score). Since the default logFC values in FindMarkers (Seurat) are small due to the

parameter pseudocount.use=1, we also provided logFC with pseudocount.use=0, more intuitive values

for differential expression (**Supplementary Table 4**).

Co-varying neighborhood analysis

To examine cell state abundance within each cell type (or cell type abundance within each lineage), we

applied co-varying neighborhood analysis (Python package CNA v0.1.6), an approach to identify single-

cell-resolution transcriptional "neighborhoods" associated with clinical phenotypes[39], to the 18 major cell

types with a total of > 5,000 cells (cell-type-level CNA) and six cell lineages (lineage-level CNA) in

dataset 1. Following our previous study scheme[6], we prepared two comparison axes: (i) disease-state

comparison between healthy controls and inactive SLE, and (ii) disease-activity comparison between

inactive and HDA SLE.

We used harmony-corrected PCs as inputs and inferred the neighborhood abundance matrix

(NAM) using all CLUES samples and cells, followed by PCA (NAM-PC). ImmVar healthy samples were

excluded from all disease-state comparisons. We then tested the association between individual-level

phenotype labels and NAM-PCs with a linear model, adjusting for age, sex, ancestry, and the number of

cells per cell type. Given the substantial correlation between disease activity and age ($r = -0.60$) in

dataset 1, we excluded age from covariates in disease-activity comparisons. For CNAs across 18 cell

types in disease-state and activity comparisons, we defined a significant association as a global

permutation P < 0.05/36 (Bonferroni correction). CNA calculates "neighborhood correlations" from the

above linear model statistics, which represent the correlation between each neighborhood and

phenotypes (positive values as expansion and negative values as depletion). We aggregated these

single-cell-level neighborhood correlation values into each cell state (**Fig. 3** and **Supplementary Table**

6) and defined the cell states with more than 50% of neighborhoods correlated with phenotypes at

empirical FDR < 0.05 as "remarkable expansion (depletion)" and those with more than 25% of

neighborhoods as "moderate expansion (depletion)".

We additionally applied CNA to dataset 1 with (1) inactive vs. LDA SLE, (2) inactive vs. MDA

SLE, (3) Asian vs. EUR SLE with the same pipelines (**Fig. 3e** and **Extended Data Fig. 4d**). For

replication analysis, we applied CNA to dataset 2 with disease-state and disease-activity comparisons

(**Fig. 3f**). We projected the dataset 2 cells into the latent transcriptional space in dataset 1 using

Symphony[45] and utilized symphony-corrected PCs as inputs of CNA. Since CNA requires ~30 samples

for stable performance[39], we compared 14 SLE patients with 0 <= SLEDAI <= 2 vs. 17 healthy for

disease-state comparison, and 15 patients with SLEDAI >= 6 vs. 14 patients with 0 <= SLEDAI <= 2 for

disease-activity comparison in dataset 2. We also applied MASC[26] to dataset 1 to test the association of

cell-state abundance with disease-state and disease-activity, adjusting for age, sex, and ancestry

(**Extended Data Fig. 4e**).

604 **ScQuaD model to dissect quantitative and qualitative changes**

605 To decompose cell-type-level pseudobulk gene expression into quantitative and qualitative changes, we

606 developed ScQuaD (Single-cell-level Quantitative and Qualitative change Dissection), a single-cell-

607 level generalized linear mixed effect model (GLMM, **Fig. 4a-b** and **Supplementary Table 8**). This model

608 fitted each single-cell-level expression to the following GLMM with a negative binomial distribution:

609 $$\log(E_i) = \beta_{abundance}X_{i,abundance} + \beta_{dysregulated}X_{d,dysregulated} + \beta_{age}X_{d,age} + \beta_{sex}X_{d,sex}$$

610 $$+\beta_{ancestry}X_{d,ancestry} + \beta_{nUMI}X_{i,nUMI} + \beta_{MT}X_{i,MT} + (\phi_d|d) + (\kappa_b|b) + \theta$$

611 where E is the expression of the gene in cell i, θ is an intercept, and all other βs represent fixed effects

612 as indicated (nUMI = number of UMI and MT = % MT) in cell i, donor d, or batch b. Donor and batch are

613 modeled as random effect intercepts. The first exploratory variable is the neighborhood correlation

614 values for cell i in cell-type-level CNA with a specific comparison (e.g., disease-state CNA), representing

615 cell-state abundance changes in that comparison (abundance signature term). The second one is binary

616 phenotype information (e.g., healthy = 0 and inactive SLE = 1), representing single-cell-level gene

617 dysregulation (dysregulated signature term). We applied this model to all genes that passed the low-

618 expression filter (>=3 counts in >= 10% of samples) across 18 major cell types (> 5,000 cells) for

619 disease-state and activity comparisons in dataset 1, using R package lme4[73]. To determine significance,

620 we used a likelihood ratio test (LRT) comparing the models with and without the abundance term or the

621 dysregulated term.

To assess the type I error calibration, we performed permutation tests across all cell types for

each comparison (**Extended Data Fig. 5a**). For the abundance term, we randomly shuffled

neighborhood correlation values across all single cells. For the dysregulated term, we shuffled

phenotype labels across all target samples.

To quantify the contribution of quantitative and qualitative changes to cell-type-level pseudobulk

gene expression, we first selected the top 1,000 P-value pseudobulk DEGs from R package edgeR[74] for

each cell type. For each gene, we calculated the explained variances by abundance and dysregulated

terms as Nagelkerke r2 from the above GLMM. We then calculated the mean across all genes for each

term and their proportions (**Extended Data Fig. 5b**).

We also investigated the abundance and dysregulated term statistics of IRGs for disease-state

and activity comparisons (**Fig. 4c-d** and **Extended Data Fig. 5c**). We selected 100 IRGs based on a

previous study[4] (**Supplementary Table 10**) and performed hierarchical clustering of them with Ward's

method, based on the Euclidean distances of the abundance and dysregulated term Z scores for each

cell type.

Pathway enrichment analysis and gene signature score

To test the enrichment of pathway annotations for abundance and dysregulated signature genes, we

performed over-representation analyses (ORA) using the top 1,000 LRT P-value genes for each

signature term in each cell type, with one-sided Fisher's exact test in R package clusterProfiler[75] (**Fig. 4e**

and **Extended Data Fig. 5d**). For pathway annotations, we used the MsigDB hallmark gene set

collection and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways. To capture the cell-type-

specific biology for both signature genes, we used the union of the top 1,000 LRT P-value genes across

all cell types as the background gene set.

For pathway analyses of cell-state-specific upregulated marker genes (FDR < 0.05 in

FindMarkers [Seurat]), we performed ORA with one-sided Fisher's exact test (**Fig. 5b**, **Extended Data**

Fig. 7c, and **Supplementary Table 6**). We set all genes that passed the low-expression filter (>=3

counts in >= 10% of samples) for the corresponding cell type as the background gene sets.

To calculate classical exhaustion signature scores in EMCD8 and EMCD4 cells, we prepared a

classical exhaustion signature gene set based on previous studies[3,5] (**Supplementary Table 17**). We

first log-normalized and scaled each gene expression in the gene set. We then calculated the signature

score for each cell as the sum of these scaled expressions across all genes (**Fig. 6b**).

Stratified LD score regression analysis

To evaluate the enrichment of the genome-wide distribution of all SLE risk variants irrespective of their

effect sizes (heritability) around abundance and dysregulated signature genes, we performed stratified

linkage disequilibrium score regression (S-LDSC)[44] with the sc-linker pipeline[76] (**Fig. 4f** and **Extended**

Data Fig. 5f). We first inferred min-max-scaled gene scores based on abundance and dysregulated term

Z scores across all tested genes for each cell type. We then calculated LD scores for EAS and EUR,

using the above gene scores as gene weights. In S-LDSC, we examined enrichment of SLE heritability

for common variants within 100-kb windows on either side of the transcription start sites (TSS) of all

genes with positive Z scores for both signatures across all cell types, adjusting for baseline model v1[44].

For this analysis, we used two large-scale SLE GWAS summary statistics from EAS and EUR

ancestries[42,43]. Since the regression coefficients of S-LDSC are influenced by GWAS heritability, we

normalized the coefficients by dividing them by the mean per-SNP heritability as reported in a previous

report[44]; we then reported the normalized coefficients. Finally, we performed fixed-effect meta-analysis of

EAS and EUR results using the inverse-variance weighting method with normalized coefficients and their

standard errors (**Extended Data Fig. 5e**). We reported P values to test whether the regression

coefficient is significantly positive.

Localization of GWAS prioritized genes for each cell state

To assign the most likely causal genes to SLE-GWAS significant variants ($P < 5 \times 10^{-8}$), we used the

L2G score from Open Targets Platform[41]. We defined 124 genes that showed L2G score > 0.5 for SLE-

GWAS variants as SLE-GWAS prioritized genes (**Supplementary Table 9**). For each cell state, we

identified overlapping genes between the SLE-GWAS prioritized genes and cell-state-specific

upregulated marker genes with FDR < 0.05 and logFC > 0.2 (FindMarkers [Seurat]; **Fig. 5d** and **Fig. 6a**).

Reference mapping

To integrate three single-cell datasets in LuPIN, we performed reference mapping using R package

Symphony[45], setting the larger dataset 1 as reference and datasets 2 and 3 as query (**Fig. 5a**). For

dataset 1, we built lineage-level and cell-type-level reference objects based on the corresponding

harmony objects used in fine-grained clustering. For datasets 2 and 3, we first separated PBMCs into

seven lineages with the same pipeline as in dataset 1 (**Sample processing and quality control**). After

normalizing and scaling gene expression with the same parameters in dataset 1, we next projected the

cells from each lineage into the latent transcriptional space of the same lineage in dataset 1 (lineage-

level reference mapping). For each cell in datasets 2 and 3, we annotated the best-predicted cell type

labels using k-nearest neighbor classifiers (k = 5) based on the cell type labels of dataset 1. We then

repeated the above procedure at the cell-type level (cell-type-level reference mapping) and annotated

the best-predicted cell state labels for each cell in datasets 2 and 3.

Surface marker protein analysis

After obtaining fine-grained cell state information for multimodal dataset 3 through the reference

mapping, we investigated the cell-state-specific surface marker profiles. For dataset 3, we profiled 130

surface marker proteins and 7 isotype controls using TotalSeq™-C Human Universal Cocktail v1.0

(Biolegend, 399905). For each cell, we normalized each protein expression with centered-log ratio (CLR)

across 130 proteins.

To identify cell-state-specific surface markers, we ran the FindMarkers function (Seurat),

comparing one cell state vs. other cell states within the same cell type. We reported the proteins with

FDR <0.05 and logFC >0.2 as cell-state-specific upregulated surface markers (**Fig. 5e**, **6e**, **Extended**

Data Fig. 6a, and **Supplementary Table 12**).

TCR analysis

In dataset 3, we used Cell Ranger-called TCR UMI counts of the cells that passed the following criteria:

(1) cells that passed in our GEX-based QC steps (**Sample processing and quality control**), (2) cells

annotated as NaiveCD4, CMCD4, EMCD4, CTLCD4, Treg, NaiveCD8, CMCD8, or EMCD8 (**Reference**

mapping), and (3) cells that passed Cell Ranger TCR QC criteria.

To identify cell-state-specific V gene usage, we compared the V gene frequencies between one

cell state vs. other cell states within the same cell type. To detect both positive and negative

enrichments, we used two-sided Fisher's exact tests (**Fig. 5c**, **Extended Data Fig. 6c**, **8b**, and

Supplementary Table 11). Among all 45 TRAV and 49 TRBV genes, we tested V genes that passed the

low expression filter (>= 10 counts in total) for each cell type.

To assess the TCR clonal expansion in EMCD8-5 (DP EMCD8), we counted the number of

clonotypes in DP EMCD8 across SLE donors (**Fig. 6c**). Clonotypes were defined as the combination of

CDR3 amino acid sequence and V gene on the β-chain. To identify DP EMCD8-specific clonotypes, we

compared the clonotype frequencies between one cell state vs. other cell states within the same cell type

for each donor. To detect both positive and negative enrichments, we used two-sided Fisher's exact

tests (**Extended Data Fig. 8a** and **Supplementary Table 18**). To identify the known cognate epitopes

for each clonotype, we used VDJdb browser (**Supplementary Table 18**).

Enhancer RNA analysis

To detect transcribed enhancers in dataset 3, we followed the ReapTEC (read-level prefiltering and

transcribed enhancer call) pipeline[14]. We sequenced 10X libraries on the Illumina NovaSeq 6000 to

generate 150 base-paired-end reads for Read1 as well as Read2, and mapped them using STARsolo[77]

to obtain reads with unencoded G. Reads were deduplicated, and those with cell barcodes for each cell

type were extracted. TSS peaks were generated by merging TSSs located within 10 bp of each other. To

identify bidirectionally transcribed candidate enhancers for each cell type, TSS peak pairs were detected

with the developers' scripts, masking promoter regions. To test their enrichment for Roadmap

epigenomes core 15 reference annotations[49], we merged called enhancers across all cell types and

compared their overlap with (1) PBMC enhancers (6_EnhG and 7_Enh in E062), (2) PBMC

heterochromatin (9_Het in E062), and (3) Brain hippocampus middle enhancers (6_EnhG and 7_Enh in

E062) using two-sided Fisher's exact tests (**Extended Data Fig. 6d**).

To identify cell-state-specific transcribed enhancers, we calculated cell-state-specific scores

using entropy as previously described[14]. We first excluded enhancers with <5 mapped reads per each

cell type and aggregated reads across donors for each cell state. We then calculated the entropy and

cell-state-specific scores for each enhancer with the following equations:

736
$$\text{Entropy}(x) = -\sum_{i=1}^{n} p_i * log_2(p_i); \ p_i = \frac{x_i}{\sum_{i=1}^{n} x_i}$$

737
$$\text{Specificity score}(x) = 1 - \frac{\text{entropy}(x)}{log_2(n)}$$

where x_i is the transcript level (count-per-million, CPM) for enhancer x in cell state i, and n represents the

number of cell states in each cell type. We defined cell-state-specific enhancers as those with a cell-

state-specific score >0.7 (**Extended Data Fig. 7a** and **Supplementary Table 14**).

Trajectory analysis

To infer the trajectory graph in the whole CD4[+] T cell lineage, we used Monocle3[51] with default

parameters (**Fig. 5h**). We set a cell in NaiveCD4-3 (SOX4[+]TOX[+]) as the root cell for pseudotime

definition based on the recent report that these cells (recent thymic emigrants) are the most immature

naive T cell population[18].

Arrayed CRISPR knock-out screening

To identify transcriptional regulators of disease-relevant cell state candidates in CD4 T cells, we

performed arrayed CRISPR knock-out (KO) screening. We first selected upregulated TFs (FDR < 0.05

and logFC > 0.3 in FindMarkers [Seurat]) in NaiveCD4-5 (22 TFs) and NaiveCD4-4 (1 TF [ARID5B];

overlapped with NaiveCD4-5), which expanded both in disease-state and activity CNA (**Fig. 3b**). Each

TF was targeted with two different guide RNAs (gRNAs) selected from a previous report[78]

(**Supplementary Table 15**; for each donor, two control samples were prepared).

755   We first isolated human primary CD4$^+$ T cells from freshly thawed PBMCs of three healthy

donors using CD4$^+$ T Cell Isolation Kit (Miltenyi, 130-096-533). For expansion, we activated CD4$^+$ T cells

for 2 days with Dynabeads Human T-Activator CD3/CD28 (Thermo, DB11132) at a 1:1 bead-to-cell ratio

and cultured them in T cell medium in the presence of 30 IU/mL IL-2 (Peprotech, 200-02), 5 ng/mL IL-7

(Peprotech, 200-07), and 5 ng/mL IL-15 (Peprotech, 200-15) at 37 °C and 5% CO_2. We replaced the

medium every three days to maintain CD4+ T cells at ~1 × 10^6/mL, and we collected and froze them

seven days after activation.

Prior to electroporation, expanded CD4 T cells from three donors were reactivated with

Dynabeads Human T-Activator CD3/CD28 at a 1:1 bead-to-cell ratio in the presence of IL-2, IL-7, and IL-

15, with beads removed after 24 h. We assembled CRISPR ribonucleoprotein (RNP) complexes by

combining recombinant Cas9 protein (IDT, 1081058) with two synthetic sgRNAs (IDT) at a 1:1:2 molar

ratio, and then incubated them at room temperature for 15 min. We performed electroporation using the

P3 Primary Cell 96-well Nucleofector kit and 4D-Nucleofector (Lonza, program EH-140). Each reaction

contained 1×10^6 expanded CD4 T cells, 3 μL Cas9 RNP, and 0.75 μL 100 μM electroporation enhancer

(IDT). After electroporation, cells were divided into three well replicates and cultured in T cell medium

supplemented with 30 IU/mL IL-2. Control cells were electroporated with equal amounts of non-targeting

Cas9 RNP and electroporation enhancer. For two donors, we knocked out all 22 TFs (discovery data,

Fig. 5f-g). For the 3rd donor, we knocked out *FOXO1* as a replication analysis (**Extended Data Fig. 7b**).

For bulk RNA-seq, we isolated RNA 5 days after transfection using MagMAX-96 Total RNA

Isolation Kit (Thermo Fisher Scientific, AM1830). Libraries were prepared using the RamDA-seq method

as previously described[79] with 100 pg of total RNA as input and sequenced on the Illumina NovaSeq

6000 to generate 150 base paired-end reads.

FCM and FACS experiments on EMCD8 cells

For in-depth characterization for EMCD8-5 cells (DP EMCD8), we performed functional assays of this

population with FCM and FACS using FACSMelody (BD Biosciences) based on four key surface

markers (HLA-DR, CD27, CD38, and CD28) with highest logFCs (FindMarkers [Seurat]; **Fig. 6e-f** and

Supplementary Table 12). We prepared three FCM panels (**Supplementary Table 19**): Panel 1 for

FACS and surface marker FCM (PD1 and TIGIT), Panel 2 for intracellular FCM (GZMK and GZMB), and

Panel 3 for intracellular FCM (IFNG).

785   For the FACS experiment, we first isolated CD8$^+$ T cells from six SLE donors using the CD8$^+$ T

Cell Isolation Kit (Miltenyi, 130-096-495). After blocking non-specific binding with anti-human Fc-gamma

receptor antibodies (Thermo Fisher Scientific) and surface marker staining (Panel 1) with the

manufacturer's protocol, we FACS-sorted target EMCD8 cells gated on CCR7$^-$HLA-

DR$^+$CD27$^+$CD38$^+$CD28$^+$ from each donor (**Extended Data Fig. 8e**) for bulk RNA-seq, ATAC-seq, and

UNIChro-seq (unique molecular identifier counting of regional chromatin accessibility with sequencing;

5,000 cells per sample). We also FACS-sorted whole EMCD8 cells gated on CCR7$^-$ from each donor as

a control (5,000 cells per sample). Total RNA from each sorted sample was extracted using MagMAX-96

Total RNA Isolation Kits. RNA-seq libraries were prepared using SMART-seq mRNA (Takara Bio

634772). For ATAC-seq, we washed sorted cells twice with cold PBS and performed tagmentation using

1.25 μL TDE1, 12.5 μL 2× TD buffer (Illumina 20034198), 0.25 μL 1% digitonin (Promega G944A), and 6

μL ddH$_2$O. We purified the tagmented DNA (Zymo D4014), amplified it for 15−18 cycles with KAPA HiFi

ReadyMix (Nippon Genetics KK2602) and Nextera XT Index Kit v2 (Illumina), and size-selected

fragments (180–1000 bp) using AMPure XP beads. UNIChro-seq libraries were prepared with custom

primers as previously described[58] to quantify chromatin accessibility at the five loci near the *IFNG* gene,

using an accessible region near the *UBE2D2* gene as a reference control (**Supplementary Table 21**).

RNA-seq and ATAC-seq libraries were sequenced on the Illumina NovaSeq 6000 to generate 150 base

paired-end reads. UNIChro-seq libraries were sequenced on the illumina Miseq platform using a custom

R1 sequencing primer (5'- ACACTCTTTCCCTACACGACGCTCTTCAGGTCT).

For FCM experiments, we isolated CD8$^+$ T cells from 11 SLE donors using CD8$^+$ T Cell Isolation

Kit (Miltenyi, 130-096-495). For surface marker staining (Panels 1-3), we took the same procedures as

the above FACS experiment. Dead cells were stained and removed using Zombie Violet Fixable Viability

Kit (Biolegend, 423113). For intracellular staining of GZMK and GZMB (Panel 2), we used Pharmingen™

Transcription Factor Buffer Set (BD Biosciences, 562574), following the manufacturer's protocol. For

intracellular staining of IFNG (Panel 3), we first stimulated CD8$^+$ T cells in culture medium with 50 ng/ml

PMA (MP Biomedicals, 16561-29-8) and 1 μg/ml ionomycin (1 μg/ml). One hour after stimulation, we

added GolgiStop™ Protein Transport Inhibitor (BD Biosciences, 554724) and cultured them for an

additional three hours. We then performed intracellular staining using Cytofix/Cytoperm™

Fixation/Permeabilization Kit (BD Biosciences, 554714), following the manufacturer's protocol. In all

panels, we compared the mean fluorescence intensities (MFI) of target proteins between CCR7$^-$HLA-

DR⁺CD27⁺CD38⁺CD28⁺ cells and whole EMCD8 cells with a linear mixed model, adjusting for age and

sex as fixed effects and donor as a random effect (**Fig. 6i** and **Extended Data Fig. 8h**). All analyses

were performed with FlowJo™ Software (BD Biosciences).

Bulk RNA-seq, ATAC-seq, and UNIChro-seq analyses

For both RNA-seq and ATAC-seq samples, adaptor sequences in FASTQ files were trimmed using

Cutadapt and short reads with < 50 base pairs were removed.

For RNA-seq samples, reads were mapped to the GRCh38 genome with STAR[80] using the

same reference files with Cell Ranger (refdata-gex-GRCh38-2020-A) for single-cell data. Expression was

counted with HTSeq[81], setting the parameter "stranded=no" for the unstranded libraries.

Among a total of 153 CRISPR KO RNA-seq samples (discovery: [22 TFs + 2 controls] × 3

replicates × 2 donors, replication: [1 TFs + 2 controls] × 3 replicates × 1 donors; **Arrayed CRISPR**

knock-out screening), we first excluded three samples with unique read counts < 4 × 10⁶ as low-quality

samples. We kept the genes that passed the low-expression filter (>=10 counts in >= 15% of samples),

followed by a trimmed mean of M values (TMM) normalization[74]. For differential expression analysis

between TF-KO conditions and controls, we fitted each count to the following GLMM with a negative

binomial distribution:

$$\log(E_s) = \beta_{KO} X_{s,KO} + (\phi_d | d) + \text{offset}(\log[libsize]) + \theta$$

where E is the expression of the gene in sample s, θ is an intercept, and β represent fixed effect for KO

condition (one KO condition vs. control). Donor d is modeled as a random effect intercept. TMM-

normalized library size is modeled as an offset term. To determine the significance of this model, we

used LRT comparing the models with or without the KO condition term in lme4[73]. We tested the

enrichment of the downregulated genes by each KO condition (LRT FDR < 0.05; **Supplementary Table**

16) for NaiveCD4-4 and NaiveCD4-5 upregulated signature genes (FDR < 0.05 in Seurat FindMarkers)

using one-sided Fisher's exact test (**Fig. 5g** and **Extended Data Fig. 7d**).

840   For EMCD8 bulk RNA-seq samples, all 12 samples (6 donors × 2 cell subsets [target and

controls]; **FCM and FACS on EMCD8 cells**) passed QCs (unique read counts > 2 × 10^7). We kept the

genes that passed the low-expression filter (>=10 counts in >= 3 samples), followed by TMM

normalization. For differential expression analysis between target (HLA-DR$^+$CD27$^+$CD38$^+$CD28$^+$ EMCD8)

and control (whole EMCD8) cell populations, we fitted each count to the following GLMM with a negative

binomial distribution in lme4[73] (**Extended Data Fig. 8f**):

$$\log(E_s) = \beta_{subset}X_{s,subset} + \beta_{age}X_{d,age} + \beta_{sex}X_{d,sex} + (\phi_d|d) + \text{offset}(\log[libsize]) + \theta$$

where E is the expression of the gene in sample s, θ is an intercept, and β represent fixed effect for cell

subsets (target vs. control). Donor d is modeled as a random effect intercept. TMM-normalized library

size is modeled as an offset term.

For EMCD8 bulk ATAC-seq and UNIChro-seq samples, reads were mapped to the GRCh38

genome with Bowtie2[82]. We filtered out low-quality reads by samtools[83] with parameters "-f 2 -q 30",

duplicated reads by Picard, with and blacklisted regions in ENCODE blacklist for hg38. We then called

ATAC narrow peaks using MACS2[84]. We kept the peaks that passed the low-expression filter (>=5

counts in >= 3 samples), followed by TMM normalization. To detect differentially accessible regions

(DARs) between target and control populations in ATAC-seq, we fitted each count to the same GLMM

equation with RNA-seq (**Extended Data Fig. 8g**). For motif enrichment analysis with HOMER, we used

the DARs with LRT FDR < 0.1 and logFC >0 (**Fig. 6g**). For UNIChro-seq samples, we extracted UMI

counts as previously described[58] at the target five loci near the *IFNG* gene and a reference control

region. To detect DARs between target and control cell populations, we used a logistic mixed effect

model, regressing a binary phenotype vector (control populations: 0, target populations: 1) for each UMI

by region information (reference region: 0, target region: 1), adjusting for donor random effect. We used

Integrative Genomics Viewer for the visualization of focused genomic regions (**Fig. 6h**).

Acknowledgements

This study was supported by the Japan Agency for Medical Research and Development (AMED)

(JP22tm0424223, JP22ek0410099, JP23tm0524005, JP25ek0410139, JP22ama121015 and

JP223fa627010, JP22gm1810002, JP22gm1810005), JSPS Grants-in-Aid for Scientific Research

(JP21K20647, JP22H03114, JP23KJ2185, JP23K15361, JP25K19621), the Uehara Memorial

Foundation, the Nakajima Foundation, Japan Intractable Diseases Research Foundation, GSK Japan

Research Grant, Daiichi Sankyo Foundation of Life Science, Mochida Memorial Foundation for Medical

and Pharmaceutical Research, Takeda COCKPI-T Funding, Kanzawa Medical Research Foundation,

SENSHIN Medical Research Foundation, Japan Research Foundation for Clinical Pharmacology, the

Japan Rheumatism Foundation, the Ichiro Kanehara Foundation, Keio University Academic

Development Funds, and Keio University Program for the Advancement of Next Generation Research

Projects. The super-computing resource was provided by Human Genome Center, Institute of Medical

Sciences, The University of Tokyo. We appreciate the generosity of the donors for a RIKEN solicited

donation project, Support for SLE research and development. We appreciate the RIKEN-IMS Genome

Platform for its help in the sequencing experiments. We thank AIDA consortium and RIKEN-IMS

Laboratory for Gene Structure and Regulation for helpful feedback.

Author contributions

M.N. and K.I. designed the study. M.N., K.I., and M.K. wrote the manuscript with critical inputs from T.N.,

Y.T., and K.Y. M.N. conducted analyses with support from K.I., M.K., K.A., H.I., T.N., J.I., H.T., and H.H.

M.K. M.N., and K.A. conducted experiments with support from T. A., T. K., S. N., R. B., Y.M., B.N., A.S.,

and K.Y. T.K., S.K., E.K., Y.F., Y.T., and Y.M. contributed to sample collection. X.Z., S.B. and C.T.

managed EAS SLE-GWAS data. All authors contributed to the final manuscript.

Competing interests

RIKEN has filed a patent application related to UNIChro-seq (WO/2024/242037). K.I, M.K, and T.A are

inventors on this patent. S.K. has received speaking fees from Eli Lilly, GlaxoSmithKline, Bristol-Myers,

Abbvie, Eisai, Pfizer, Astra-Zeneca, and research grants from Daiichi-Sankyo, Abbvie, Boehringer

Ingelheim, and Astellas.

Materials and Correspondence

Correspondence and requests for materials should be addressed to Masahiro Nakano and Kazuyoshi

Ishigaki.

Reference

1. Tsokos, G. C., Lo, M. S., Reis, P. C. & Sullivan, K. E. New insights into the immunopathogenesis of systemic lupus

erythematosus. *Nat. Rev. Rheumatol.* **12**, 716–730 (2016).

2. Banchereau, R. *et al.* Personalized Immunomonitoring Uncovers Molecular Networks that Stratify Lupus Patients. *Cell* **165**,

551–565 (2016).

3. the Accelerating Medicines Partnership in SLE network *et al.* The immune cell landscape in kidneys of patients with lupus

nephritis. *Nat. Immunol.* **20**, 902–914 (2019).

4. Nehar-Belaid, D. *et al.* Mapping systemic lupus erythematosus heterogeneity at the single-cell level. *Nat. Immunol.* **21**,

1094–1106 (2020).

5. Perez, R. K. *et al.* Single-cell RNA-seq reveals cell type–specific molecular and genetic associations to lupus. *Science* **376**,

eabf1970 (2022).

6. Nakano, M. *et al.* Distinct transcriptome architectures underlying lupus establishment and exacerbation. *Cell* **185**, 3375-

3389.e21 (2022).

7. Rao, D. A. *et al.* Pathologically expanded peripheral T helper cell subset drives B cells in rheumatoid arthritis. *Nature* **542**,

110–114 (2017).

8. Jenks, S. A. *et al.* Distinct Effector B Cells Induced by Unregulated Toll-like Receptor 7 Contribute to Pathogenic

Responses in Systemic Lupus Erythematosus. *Immunity* **49**, 725-739.e6 (2018).

9. Navarra, S. V. *et al.* Efficacy and safety of belimumab in patients with active systemic lupus erythematosus: a randomised,

placebo-controlled, phase 3 trial. *The Lancet* **377**, 721–731 (2011).

10. Morand, E. F. *et al.* Trial of Anifrolumab in Active Systemic Lupus Erythematosus. *N. Engl. J. Med.* **382**, 211–221 (2020).

11. Missarova, A., Dann, E., Rosen, L., Satija, R. & Marioni, J. Leveraging neighborhood representations of single-cell data to

achieve sensitive DE testing with miloDE. *Genome Biol.* **25**, 189 (2024).

12. Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in single cells. *Nat. Methods* **14**, 865–868

(2017).

13. Moody, J. *et al.* A single-cell atlas of transcribed *cis* -regulatory elements in the human genome. Preprint at

https://doi.org/10.1101/2023.11.13.566791 (2023).

14. Oguchi, A. *et al.* An atlas of transcribed enhancers across helper T cell diversity for decoding human diseases. *Science*

385, eadd8394 (2024).

15. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29 (2021).

16. Watanabe, M., Moon, K. D., Vacchio, M. S., Hathcock, K. S. & Hodes, R. J. Downmodulation of Tumor Suppressor p53 by

T Cell Receptor Signaling Is Critical for Antigen-Specific CD4+ T Cell Responses. *Immunity* **40**, 681–691 (2014).

17. Castellanos, M. C. *et al.* Expression of the leukocyte early activation antigen CD69 is regulated by the transcription factor

AP-1. *J. Immunol. Baltim. Md 1950* **159**, 5463–5473 (1997).

18. Bohacova, P. *et al.* Multidimensional profiling of human T cells reveals high CD38 expression, marking recent thymic

emigrants and age-related naive T cell remodeling. *Immunity* **57**, 2362-2379.e10 (2024).

19. Yasumizu, Y. *et al.* Single-cell transcriptome landscape of circulating CD4+ T cell populations in autoimmune diseases. *Cell*

Genomics **4**, 100473 (2024).

20. Chan, J. D. *et al.* FOXO1 enhances CAR T cell stemness, metabolic fitness and efficacy. *Nature* **629**, 201–210 (2024).

21. Doan, A. E. *et al.* FOXO1 is a master regulator of memory programming in CAR T cells. *Nature* **629**, 211–218 (2024).

22. Crow, M. K. Type I Interferon in the Pathogenesis of Lupus. *J. Immunol.* **192**, 5459–5468 (2014).

23. He, J. *et al.* Circulating Precursor CCR7loPD-1hi CXCR5+ CD4+ T Cells Indicate Tfh Cell Activity and Promote Antibody

Responses upon Antigen Reexposure. *Immunity* **39**, 770–781 (2013).

24. Zhang, F. *et al.* Deconstruction of rheumatoid arthritis synovium defines inflammatory subtypes. *Nature* **623**, 616–624

(2023).

25. Koga, T. *et al.* CaMK4-dependent activation of AKT/mTOR and CREM-α underlies autoimmunity-associated Th17

imbalance. *J. Clin. Invest.* **124**, 2234–2245 (2014).

26. Fonseka, C. Y. *et al.* Mixed-effects association of single cells identifies an expanded effector CD4[+] T cell subset in

rheumatoid arthritis. *Sci. Transl. Med.* **10**, eaaq0305 (2018).

27. Patil, V. S. *et al.* Precursors of human CD4[+] cytotoxic T lymphocytes identified by single-cell transcriptome analysis. *Sci.*

Immunol. **3**, eaan8664 (2018).

28. Luo, Y. *et al.* Single-cell transcriptomic analysis reveals disparate effector differentiation pathways in human Treg

compartment. *Nat. Commun.* **12**, 3913 (2021).

29. Li, J. *et al.* KIR[+] CD8[+] T cells suppress pathogenic T cells and are active in autoimmune diseases and COVID-19. *Science*

376, eabi9591 (2022).

30. Jonsson, A. H. *et al.* Granzyme K[+] CD8 T cells form a core population in inflamed human tissue. *Sci. Transl. Med.* **14**,

eabo0686 (2022).

31. Velikkakam, T., Gollob, K. J. & Dutra, W. O. Double-negative T cells: Setting the stage for disease control or progression.

Immunology **165**, 371–385 (2022).

32. Colonna, M. Innate Lymphoid Cells: Diversity, Plasticity, and Unique Functions in Immunity. *Immunity* **48**, 1104–1117

(2018).

33. Morgan, D. & Tergaonkar, V. Unraveling B cell trajectories at single cell resolution. *Trends Immunol.* **43**, 210–229 (2022).

34. Nutt, S. L., Hodgkin, P. D., Tarlinton, D. M. & Corcoran, L. M. The generation of antibody-secreting plasma cells. *Nat. Rev.*

Immunol. **15**, 160–171 (2015).

35. Mulder, K. *et al.* Cross-tissue single-cell landscape of human monocytes and macrophages in health and disease. *Immunity*

54, 1883-1900.e5 (2021).

36. Leffler, J., Bengtsson, A. A. & Blom, A. M. The complement system in systemic lupus erythematosus: an update. *Ann.*

Rheum. Dis. **73**, 1601–1606 (2014).

37. Villani, A.-C. *et al.* Single-cell RNA-seq reveals new types of human blood dendritic cells, monocytes, and progenitors.

Science **356**, eaah4573 (2017).

38. Zhang, Q. *et al.* Landscape and Dynamics of Single Immune Cells in Hepatocellular Carcinoma. *Cell* **179**, 829-845.e20

(2019).

39. Reshef, Y. A. *et al.* Co-varying neighborhood analysis identifies cell populations associated with phenotypes of interest from

single-cell transcriptomics. *Nat. Biotechnol.* **40**, 355–363 (2022).

40. Rector, I. *et al.* Differential regulation of the interferon response in systemic lupus erythematosus distinguishes patients of

Asian ancestry. *RMD Open* **9**, e003475 (2023).

41. Mountjoy, E. *et al.* An open approach to systematically prioritize causal variants and genes at all published human GWAS

trait-associated loci. *Nat. Genet.* **53**, 1527–1533 (2021).

42. Yin, X. *et al.* Meta-analysis of 208370 East Asians identifies 113 susceptibility loci for systemic lupus erythematosus. *Ann.*

Rheum. Dis. **80**, 632–640 (2021).

43. Langefeld, C. D. *et al.* Transancestral mapping and genetic load in systemic lupus erythematosus. *Nat. Commun.* **8**, 16021

(2017).

44. Finucane, H. K. *et al.* Heritability enrichment of specifically expressed genes identifies disease-relevant tissues and cell

types. *Nat. Genet.* **50**, 621–629 (2018).

45. Kang, J. B. *et al.* Efficient and precise single-cell reference atlas mapping with Symphony. *Nat. Commun.* **12**, 5890 (2021).

46. Lagattuta, K. A. *et al.* Repertoire analyses reveal T cell antigen receptor sequence features that influence T cell fate. *Nat.*

Immunol. **23**, 446–457 (2022).

47. Nagafuchi, Y. *et al.* T cell plasticity in systemic lupus erythematosus revealed by large-scale T cell receptor repertoire and

transcriptome studies. Preprint at https://doi.org/10.1101/2025.01.06.24319648 (2025).

48. Britanova, O. V. *et al.* Targeted depletion of TRBV9+ T cells as immunotherapy in a patient with ankylosing spondylitis. *Nat.*

Med. **29**, 2731–2736 (2023).

49. Roadmap Epigenomics Consortium *et al.* Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317–330

(2015).

50. Meyer, A., Parmar, P. J. & Shahrara, S. Significance of IL-7 and IL-7R in RA and autoimmunity. *Autoimmun. Rev.* **21**,

103120 (2022).

51. Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis. *Nature* **566**, 496–502 (2019).

52. Bai, Y., Hu, M., Chen, Z., Wei, J. & Du, H. Single-Cell Transcriptome Analysis Reveals RGS1 as a New Marker and

Promoting Factor for T-Cell Exhaustion in Multiple Cancers. *Front. Immunol.* **12**, 767070 (2021).

53. Saggau, C. *et al.* Autoantigen-specific CD4+ T cells acquire an exhausted phenotype and persist in human antigen-specific

autoimmune diseases. *Immunity* **57**, 2416-2432.e8 (2024).

54. Chiche, L. *et al.* Modular Transcriptional Repertoire Analyses of Adults With Systemic Lupus Erythematosus Reveal Distinct

Type I and Type II Interferon Signatures. *Arthritis Rheumatol.* **66**, 1583–1595 (2014).

55. Munroe, M. E. *et al.* Altered type II interferon precedes autoantibody accrual and elevated type I interferon activity prior to

systemic lupus erythematosus classification. *Ann. Rheum. Dis.* **75**, 2014–2021 (2016).

56. Fava, A. *et al.* Integrated urine proteomics and renal single-cell genomics identify an IFN-γ response gradient in lupus

nephritis. *JCI Insight* **5**, e138345 (2020).

57. Perl, A. & Morel, L. Expanding scope of TEMRA in autoimmunity. *eBioMedicine* **90**, 104520 (2023).

58. Kono, M. *et al.* Accurate, sensitive, and efficient chromatin accessibility quantification at target loci using UNIChro-seq.

medRxiv 2025.07.29.25332340 (2025) doi:10.1101/2025.07.29.25332340.

59. Taylor, R. P. & Lindorfer, M. A. Drug Insight: the mechanism of action of rituximab in autoimmune disease—the immune

complex decoy hypothesis. *Nat. Clin. Pract. Rheumatol.* **3**, 86–95 (2007).

60. Müller, F. *et al.* CD19 CAR T-Cell Therapy in Autoimmune Disease — A Case Series with Follow-up. *N. Engl. J. Med.* **390**,

687–700 (2024).

61. Gurajala, S. *et al.* A population-scale atlas of blood and tissue in lupus nephritis. *bioRxiv* 2025.08.11.669754 (2025)

doi:10.1101/2025.08.11.669754.

62. Ishigaki, K. Beyond GWAS: from simple associations to functional insights. *Semin. Immunopathol.* **44**, 3–14 (2022).

63. Lappalainen, T. & MacArthur, D. G. From variant to function in human disease genetics. *Science* **373**, 1464–1468 (2021).

64. Bombardier, C. *et al.* Derivation of the sledai. A disease activity index for lupus patients. *Arthritis Rheum.* **35**, 630–640

(1992).

65. Rogozhnikov, A. *et al.* Demuxalot: scaled up genetic demultiplexing for single-cell sequencing. Preprint at

https://doi.org/10.1101/2021.05.22.443646 (2021).

66. Delaneau, O., Zagury, J.-F. & Marchini, J. Improved whole-chromosome phasing for disease and population genetic

studies. *Nat. Methods* **10**, 5–6 (2013).

67. Das, S. *et al.* Next-generation genotype imputation service and methods. *Nat. Genet.* **48**, 1284–1287 (2016).

68. Heaton, H. *et al.* Souporcell: robust clustering of single-cell RNA-seq data by genotype without reference genotypes. *Nat.*

Methods **17**, 615–620 (2020).

69. Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq data using regularized negative

binomial regression. *Genome Biol.* **20**, 296 (2019).

70. Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-cell analysis. *Nat. Biotechnol.* **42**, 293–304

(2024).

71. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat. Methods* **16**, 1289–1296

(2019).

72. Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of Cell Doublets in Single-Cell Transcriptomic

Data. *Cell Syst.* **8**, 281-291.e9 (2019).

73. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using **lme4**. *J. Stat. Softw.* **67**, (2015).

74. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR : a Bioconductor package for differential expression analysis of

digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

75. Xu, S. *et al.* Using clusterProfiler to characterize multiomics data. *Nat. Protoc.* **19**, 3292–3320 (2024).

76. Jagadeesh, K. A. *et al.* Identifying disease-critical cell types and cellular processes by integrating single-cell RNA-

1037   sequencing and human genetics. *Nat. Genet.* **54**, 1479–1492 (2022).

1038   77. Kaminow, B., Yunusov, D. & Dobin, A. STARsolo: accurate, fast and versatile mapping/quantification of single-cell and

1039   single-nucleus RNA-seq data. Preprint at https://doi.org/10.1101/2021.05.05.442755 (2021).

1040   78. Freimer, J. W. *et al.* Systematic discovery and perturbation of regulatory genes in human T cells reveals the architecture of

1041   immune networks. *Nat. Genet.* **54**, 1133–1144 (2022).

1042   79. Hayashi, T. *et al.* Single-cell full-length total RNA sequencing uncovers dynamics of recursive splicing and enhancer RNAs.

1043   *Nat. Commun.* **9**, 619 (2018).

1044   80. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

1045   81. Anders, S., Pyl, P. T. & Huber, W. HTSeq—a Python framework to work with high-throughput sequencing data.

1046   *Bioinformatics* **31**, 166–169 (2015).

1047   82. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–359 (2012).

1048   83. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

1049   84. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

1050