Coordinated action of CRK2 and QSK1 regulate osmotic stress response in Arabidopsis

Sunita Jindal[1], Adam Zeiner[1,2], Alexey Bondar[1,2], Michaela Neubergerová[3,4], Sara Christina

Stolze[5], Anne Harzen[5], Francisco J. Colina[1,#], Sara Liekens[1,6], Mirva Pääkkönen[7], Johannes

Merilahti[7], Ivan Kulich[1,2], Roman Pleskot[3], Hirofumi Nakagami[5], Michael Wrzaczek[1,2,*]

[1]Institute of Plant Molecular Biology, Biology Centre, Czech Academy of Sciences, 37005 České
Budějovice, Czechia

[2]Faculty of Science, University of South Bohemia, 37005 České Budějovice, Czechia

[3]Institute of Experimental Botany (IEB), Czech Academy of Sciences, 16502 Prague, Czechia

[4]Faculty of Science, Charles University, 12800 Prague, Czechia

[5]Max Planck Institute for Plant Breeding Research, Cologne, Germany

[6]Thomas More University of Applied Sciences, Campus Geel, 2440 Geel, Belgium

[7]Turku Bioscience Centre, University of Turku and Åbo Akademi University, 20520 Turku, Finland

[#]Current address: Estación Experimental del Zaidín (EEZ-CSIC), 18008, Granada, Spain

*Correspondence: michael.wrzaczek@umbr.cas.cz

Abstract

Regulation of callose deposition at plasmodesmata (PD) is necessary for optimal intercellular
communication during development and stress responses. The Cysteine-rich Receptor-like
Kinase CRK2 and the Leucine-rich repeat Receptor-like kinase QSK1 relocalize from the
general plasma membrane (PM) to PD in response to osmotic stress and regulate callose
deposition. However, the current understanding on regulation of callose accumulation under
osmotic stress remains limited. In this study, we identify a novel role for CRK2 in regulating
osmotic stress-induced callose deposition through controlling the availability of QSK1 at PD. We
show that osmotic stress-induced enrichment of QSK1 at PD requires functional CRK2 and
establish that phosphorylation of QSK1 at serines 621 and 626 is inhibitory for its enrichment at
PD. QSK1 positively regulates stress-induced callose deposition in connection with its
enrichment at PD. Live cell confocal microscopy and fluorescent lifetime imaging further reveal
that CRK2 inhibits PD localization of QSK1 under normal growth conditions by phosphorylating
and sequestering it to the general PM. We propose that following osmotic stress,
dephosphorylation of QSK1 releases it from the complex, enabling it to relocalize to PD. This
regulatory mechanism prevents constitutively high callose synthesis during normal growth
conditions and allows a downstream response only in the presence of osmotic stress stimuli to
precisely control callose levels thus providing stress adaptability.

Introduction

In natural habitats, plant water uptake is compromised frequently by salinity, drought and extreme temperatures leading to osmotic stress which strongly limits normal growth and development. On the physiological level, osmotic adjustments are made by closing stomata, growth inhibition, hydraulic modifications and changes in root system architecture. On the molecular level, rapid signaling events are triggered at the plasma membrane (PM) and include activation of phosphorylation networks, activation of NADPH oxidases (respiratory burst oxidase homologs, RBOHs) and ROS burst (Martinière et al. 2019), Ca^{2+} influx (Knight et al. 1997; Tracy et al. 2008; Stephan et al. 2016) and changes in lipid signaling, for example, accumulation of phosphatidic acid (PA) (Munnik et al. 2000; Katagiri et al. 2001; Testerink and Munnik 2011; Yao and Xue 2018). Another key regulatory element of osmotic stress signaling is the deposition of callose (Hunter et al. 2019; Sangi et al. 2023; Su et al. 2023). Callose, a β-1,3-glucan polysaccharide is deposited primarily in the neck region of plasmodesmata (PD) and controls symplastic connectivity through gating the PD aperture (J Maule et al. 2013; Han et al. 2014). Symplastic connection regulates cell division, expansion and organ formation (Otero et al. 2016; Wu et al. 2016) which could provide growth adaptation especially during extreme stress conditions. For example, mutants accumulating higher levels of callose have reduced primary root (PR) length (Vatén et al. 2011; Hunter et al. 2019) and altered lateral root (LR) patterning (Benitez-Alfonso et al. 2013; J Maule et al. 2013). Therefore, adaptations in the root phenotype during osmotic stress can at least partly result from the changes in symplastic connection.

The amount of callose deposition at PD changes dynamically throughout plant growth, development and during stress conditions. The main determinants of callose amounts are callose synthases (CALS) and β-1,3-glucanases (PdBGs), two antagonistic classes of enzymes responsible for callose synthesis and degradation, respectively (Wu et al. 2018). In addition, several PD-localized proteins (PDLPs), PD-localized chaperons such as PD-callose binding proteins (PDCBs) participate in maintaining callose homeostasis by recruiting or stabilizing CALS or PdBGs (Lee et al. 2011; Wu et al. 2018; Tee et al. 2023). In recent years, receptor-like kinases (RLKs) have emerged as regulating callose accumulation during abiotic stress (Grison et al. 2019; Hunter et al. 2019; Vu et al. 2023). We previously showed that CRK2, a cysteine-rich RLK (CRK) is involved in the regulation of callose deposition and positively regulates seed

germination during salinity stress (Hunter et al. 2019). The mutant *crk2* exhibits an overall small

phenotype and deposits higher callose in unstressed conditions. CRK2 relocalizes to PD in

response to salt and mannitol treatments and also assembles in nanodomain-like clusters upon

flg22 and H_2O_2 exposure (Hunter et al. 2019). Concurrently, Grison *et al.*, 2019 showed that

Qian Shou kinase 1 (QSK1), a leucine-rich repeat RLK (LRR-RLK) is recruited to PD upon

osmotic stress and the *qsk1* mutant accumulates lower callose levels and shows altered root

response to mannitol.

PM components undergo dynamic re-organization upon perception of various stimuli and cluster

into nanodomains (Jaillais et al. 2024) to facilitate the assembly of signaling complexes and

providing signal specificity (Jarsch et al. 2014; Bücherl et al. 2017). Membrane nanodomains are

hallmarked by specific lipid and protein compositions. Scaffold proteins such as Remorins and

Stomatin/Prohibitin/Flotillin/HflK/C (SPFH) domain-containing protein bridge lipids and

membrane receptors, whereas RLKs are dynamically positioned in nanodomains (Jaillais and Ott

2020; Hdedeh et al. 2025). Cytoskeletal elements and cell wall components further regulate

nanodomain mobility and spatial distribution (Hdedeh et al. 2025). The re-organization of

membrane proteins is often coupled to a phosphorylation switch. For example, in resting cells,

BRI1 (Brassinosteroid Insensitive 1)-Associated receptor Kinase 1 (BAK1), an LRR-RLK, is

evenly distributed across the PM, with its catalytic pocket shielded by BAK1-Interacting

Receptor-like kinases (BIRs) (Bücherl et al. 2013). Ligand perception such as brassinosteroids or

pathogen-associated molecular patterns (PAMPs) activates the BRI1-BAK1 or FLAGELLIN-

SENSITIVE 2 (FLS2)-BAK1 signaling axes, triggering growth or immune signaling,

respectively. This activation involves a phosphorylation-dependent shift of BAK1 from resting to

the active state and the formation of receptor nanodomains (Chinchilla et al. 2009). Likewise,

the distribution of Remorin (Perraki et al. 2018; Xu et al. 2024) and QSK1 (Grison et al. 2019)

across the PM is dependent on their phosphorylation status.

Similar to BAK1, a shared co-receptor in growth and immune signaling, QSK1 forms several

protein complexes. For instance, the SIRK1-QSK1 module regulates aquaporins where QSK1

acts as a co-receptor of the RLK Sucrose-Induced Receptor Kinase 1 (SIRK1) (Wu et al. 2019;

Wang et al. 2022). QSK1 also associates with pattern recognition receptor (PRR)-RBOHD

complexes involving FLS2 and Elongation Factor Tu Receptor (EFR) RLKs (Goto et al. 2024).

Isner *et al.*, 2018 proposed an involvement of an additional receptor kinase in QSK1-mediated

regulation of the vacuolar Two-pore potassium channel 1 (TPK1) during stomatal closure.

Similarly, QSK1 associates with the transceptor nitrate transporter NRT1.1 in regulating PM

Arabidopsis H^+-ATPase 2 (AHA2) (Zhu et al. 2024). In addition, we previously identified that

QSK1 also associates with CRK2 (Hunter et al. 2019).

While the re-organization of individual PM components is extensively studied, how different

RLKs coordinate their spatio-temporal behavior and how the shared components are exchanged

between the complexes is much less understood. CRK2 and QSK1 are both primarily present at

the general PM under normal growth conditions and both relocalize to PD upon osmotic stress

stimulus. However, the functional link between these two RLKs in osmotic stress response was

not explored. Here, we aim to identify the connection between CRK2 and QSK1 in their

recruitment at PD to fine-tune callose deposition and eventually plant response to osmotic stress.

Our data suggest that CRK2 and QSK1 antagonistically regulate callose deposition during

osmotic stress responses and CRK2 controls the localization of QSK1. Our work provides

evidence that the dynamics of the re-organization of QSK1 and CRK2 in response to osmotic

stress is coordinated and controlled in space and time.

Results

Role of CRK2 and QSK1 in plant growth and osmotic stress response

The growth phenotype and osmotic stress responses of the *crk2* and *qsk1* mutants have been

previously described (Bourdais et al. 2015; Grison et al. 2019; Hunter et al. 2019). To understand

their possible functional interaction, we generated the *crk2 qsk1* double mutant by crossing the

single mutants *crk2* and *qsk1* and characterized the growth phenotypes including rosette area,

primary root (PR) length and lateral root (LR) density under normal growth conditions and

analyzed their salt stress responses. As reported previously (Hunter et al. 2019), *crk2* displayed

overall smaller rosette size, PR length and LR density (Fig. 1, Table S1). The *qsk1* rosette size

was comparable to Col-0 wild type (WT), while in *crk2 qsk1*, the small rosette phenotype of *crk2*

was rescued with a slight increase of rosette size in comparison to the WT (Fig. 1A, B). While

crk2 exhibited the shortest PR, *qsk1* and *crk2 qsk1* also displayed shorter PR compared to the

WT (Fig. 1C, Fig. S1A). Similar to PR, LR density was lower compared to WT in all the tested

mutants (Fig.1 D). Overall, the observable phenotypic traits of *crk2 qsk1* were more similar to

either WT or *qsk1* rather than *crk2*. To test the effect of salinity on PR growth, we carried out

root elongation assays where 5 DAS (days-after stratification) seedlings were transplanted to

medium containing salt. When normalized to the PR growth of the respective control conditions,

all the mutants exhibited a similar degree of retardation in PR elongation compared to WT (Fig.

S1B, C). The partial reversion of *crk2* phenotypes in *crk2 qsk1* mutant was also observed at the

level of stress responses. Hunter *et al.*, 2019 reported that *crk2* exhibits hypersensitivity to salt

during seed germination compared to the WT. We further extended this work by testing *qsk1* and

crk2 qsk1 mutants and observed that, while *crk2* seeds showed hypersensitivity to salt as

reported previously (Hunter *et al.*, 2019), *qsk1* and *crk2 qsk1* seed germination was comparable

to WT (Fig. S1D). Taken together, these results suggest, that CRK2 and QSK1 functionally

interact to regulate plant growth and stress responses, and moderation of *crk2* phenotype by

introduction of additional *qsk1* mutation points towards antagonistic roles for those two RLKs.

CRK2 and QSK1 fine-tune callose deposition in response to osmotic stress

The mutants *crk2* and *qsk1* have been previously reported to accumulate altered levels of callose

(Hunter et al., 2019, Grison et al., 2019). To further understand their role and interaction in

osmotic stress-induced callose deposition, we quantified callose accumulation in WT, *crk2*, *qsk1*

and *crk2 qsk1* seedlings by staining callose with aniline blue followed by microscopy-based

detection of the aniline blue fluorescence signal (Fig. 2, Table S2) (Xu et al. 2017; Hunter et al.

2019; Huang et al. 2022). Since osmotic stress is a common component of various abiotic

stresses, including salinity, drought, waterlogging, and extreme temperatures, we used mannitol

to specifically impose osmotic stress without introducing additional ionic effects. Two-week-old

hydroponically-grown Arabidopsis plants were treated with mannitol for 7 hours before aniline

blue staining and imaging. We quantified mean fluorescence intensity of the spots (Fig. 2B), the

number of deposits (Fig. 2C) and total callose deposition by taking into consideration both the

parameters (intensity and number) (Fig. 2D). In the *crk2* mutant, both intensity and number of

deposits were found to be higher than the WT in control conditions which further increased upon

mannitol treatment. This suggested a negative influence of CRK2 on callose accumulation during

steady-state as well as osmotic stress conditions. In the *qsk1* mutant, the intensity of deposits was

slightly higher than in WT but the number of deposits was similar, and an overall comparable

level of total callose during control conditions was observed. Surprisingly, mannitol-treated *qsk1*

showed a higher intensity of deposits but reduced number and lower total callose signal compared to normal growth conditions suggesting that QSK1 might not play a role in basal callose deposition but positively regulates osmotic stress-induced callose deposition (Fig. 2). Interestingly, the *crk2 qsk1* double mutant seemed to negate the reciprocal impact of each other in the *crk2 qsk1* plants resulting in similar callose deposition compared to the WT under normal growth conditions as well as upon mannitol treatment (Fig. 2). Together, these findings suggest opposing roles of CRK2 and QSK1 in regulating osmotic stress-induced callose deposition.

Functional CRK2 is essential for osmotic stress-induced enrichment of QSK1 at PD

CRK2 and QSK1 were previously shown to relocalize to PD in response to osmotic stress (Grison et al. 2019; Hunter et al. 2019) and QSK1 was identified as an *in planta* interactor of CRK2 (Hunter et al. 2019). We hypothesized that their interaction might be necessary for the enrichment of both proteins at PD. Therefore, we tested whether the absence of functional CRK2 or QSK1 would impact the recruitment of the respective other RLK to PD. To this end, we generated transgenic plants expressing QSK1 C-terminally fused to mCherry driven by the native *QSK1* promoter in *crk2* as well as in WT background and imaged leaf epidermal cells. Interestingly, while QSK1 was enriched at PD following mannitol or NaCl treatment in WT background, this enrichment was abolished in *crk2* (Fig. 3A, B, Fig. S2A, Table S3). In QSK1-mCherry/WT, the enrichment index, which is the fluorescence intensity ratio between the enriched spot (PD) and an adjacent non-enriched spot on the bulk PM, increased from 0.940 to 1.693 upon mannitol treatment, whereas it did not increase for QSK1-mCherry/*crk2* and remained approximately 1.0 for both control and mannitol treatment (Fig. 3B). Since callose is particularly present at PD, it is used to mark PD by aniline blue staining (Xu et al. 2017; Hunter et al. 2019; Huang et al. 2022). To test if the enriched QSK1-mCherry spots following osmotic stress treatment corresponded with PD, we applied osmotic stress and callose staining by simultaneously treating leaves with salt and aniline blue. The co-localization of aniline blue-stained callose with enriched mCherry signal supported that the punctate pattern of QSK1-mCherry correspond to PD (Fig. S2B). Consistently, the genetic cross between QSK1-mCherry/*crk2* and CRK2-mVenus/*crk2* rescued QSK1 enrichment at PD in response to mannitol treatment (Fig. 3C). At this timepoint after mannitol treatment, we did not observe PD enrichment for CRK2 possibly due to a lag in the PD relocalization of CRK2 (Fig. 3C). The

mutation D450N in CRK2 (CRK2^{D450N}) in the catalytic core of the CRK2 kinase domain is

known to abolish the phosphorylation activity of CRK2, and inactive CRK2 (CRK2^{D450N}) is not

able to relocalize to PD (Hunter et al. 2019). To determine whether the kinase activity of CRK2

was also essential for the recruitment of QSK1 at PD, we crossed QSK1-mCherry/*crk2* with

CRK2^{D450N}-mVenus/*crk2* plants. Inactive CRK2 (CRK2^{D450N}-mVenus) was not able to direct

QSK1 to PD in response to mannitol treatment (Fig. 3D). Next, to investigate whether CRK2

requires QSK1 for its own PD localization, we generated Arabidopsis transgenic plants

expressing CRK2-mVenus under native *CRK2* promoter in WT and *qsk1* backgrounds.

Interestingly, we found that CRK2 relocalized to PD upon mannitol treatment even in the

absence of functional QSK1 (Fig. 3E, F). These results established that enzymatically active

CRK2 is essential for the relocalization of QSK1 to PD in response to osmotic stress. However,

the relocalization of CRK2 is independent of the presence of QSK1 pointing to an upstream role

of CRK2 in PD-targeting of itself and QSK1.

200 QSK1 and CRK2 enrich at PD in a sequential manner following osmotic stress

RLKs are dynamically re-organized at the PM in response to various internal and external cues

(Jarsch et al. 2014; Bücherl et al. 2017). Despite the dependence of QSK1 on CRK2 in its

recruitment at PD, we observed a temporal disconnection in their relocalization to PD. QSK1

accumulation at PD peaked within 30-45 min of salt or mannitol treatment and subsequently the

punctate pattern faded with almost no observable enrichment 60 min post-treatment (Fig. S2A,

Fig. 4, Fig. S3). In contrast to QSK1, the punctate pattern corresponding to CRK2-mVenus

emerged at PD only 60 min following the treatment. By the time of peak CRK2 accumulation at

PD, QSK1 no longer displayed PD localization (Fig. 4). Using QSK1-mCherry and CRK2-

mVenus co-expressing lines, we carried out a time-course analysis and showed that, enrichment

of QSK1 and CRK2 at PD was not simultaneous (Fig. 4, Fig. S3). Upon treatment with mannitol

QSK1-mCherry relocalized to PD while CRK2-mVenus remained at the general PM. Following

peak enrichment of QSK1-mCherry at PD, CRK2-mVenus started to relocalize to PD. At peak

enrichment of CRK2-mVenus at PD we did not observe localization of QSK1-mCherry at PD

anymore. Overall, we did not observe simultaneous peak enrichment for both QSK1 and CRK2

at PD. The temporal separation of QSK1 and CRK2 enrichment at PD suggested that the two

RLKs might co-ordinate to precisely fine-tune callose deposition to control the PD aperture.

CRK2 and QSK1 exist in a pre-formed complex at the bulk PM that dissociates upon osmotic stress

We previously detected QSK1 as in *in vivo* interactor of CRK2 (Hunter et al. 2019) implying that the two RLKs co-localize at the general PM in close proximity by interacting either directly or in a multi-protein complex. However, the sequential recruitment of QSK1 and CRK2 at PD suggested a dissociation of the two upon stress perception allowing sequential relocalization. To test this, we carried out Förster resonance energy transfer by fluorescence lifetime imaging (FRET-FLIM) of Arabidopsis plants stably co-expressing QSK1-mCherry and CRK2-mVenus proteins (QSK1-mCherry/*crk2* x CRK2-mVenus/*crk2*) with and without mannitol treatment. As negative control, we used equivalent plants with kinase inactive variant of CRK2 (QSK1-mCherry/*crk2* x CRK2^{D450N}-mVenus/*crk2*), since CRK2^{D450N} did not relocalize following stress application (Hunter et al. 2019). In plants expressing the donor-acceptor pair (CRK2-mVenus x QSK1-mCherry), donor life-time was found to be lower in comparison to the donor-only and negative control plants under control conditions (Fig. 5A, Table S4). This suggested the existence of a preformed complex of CRK2 and QSK1 in control conditions which would reduce the fluorescence lifetime of the donor mVenus. Interestingly, upon mannitol treatment, the donor lifetime in QSK1-mCherry x CRK2-mVenus plants increased suggesting dissociation of the complex upon mannitol treatment. This finding was in-line with the sequential enrichment of QSK1 and CRK2 at PD (Fig. 4, Fig. S3). In contrast, in the QSK1-mCherry x CRK2^{D450N}-mVenus plants, the donor lifetime was higher under both control and mannitol-treated conditions suggesting that the kinase inactive CRK2 might have weaker affinity for the protein complex formation (Fig. 5A). Further, we performed structural modeling to predict the interaction between CRK2 and QSK1 using Colabfold (version 1.5.2) implementation of AlphaFold2 (Jumper et al. 2021; Mirdita et al. 2022). We modeled the full-length proteins, as well as the combinations of their individual structural segments including extracellular domain (ExD), transmembrane domain (TMD) and the cytosolic kinase domain. The modeled combinations were i) full-length proteins, ii) ExD, iii) ExD+TMD, iv) TMD, v) TMD+Kinase domain, and vi) Kinase domain for both CRK2 and QSK1. Based on the predicted aligned error (PAE) values (Fig. 5B, C, Fig. S5) and interface predicted template modelling (iPTM) scores (Table S5), ExD+TMD regions showed high confidence for interaction (Fig. 5B). The modeling of the full-length proteins suggested that the two RLKs engage specific segments for interaction (Fig. 5C)

248   which mainly resulted from the ExD and TMD regions (Fig. S4). Collectively, these results

249   suggested that CRK2 and QSK1 are present in close proximity at the general PM and form a

250   complex *in planta,* potentially involving additional proteins, and this complex dissociates upon

251   osmotic stress perception.

CRK2 phosphorylates QSK1 on its C-terminus, and salt stress reduces this
phosphorylation

The role of QSK1 in PD regulation was reported to be regulated by its phosphorylation (Grison

et al. 2019) specifically of two serine residues in the C-terminus of the cytosolic region of QSK1

(Ser-621 and Ser-626). However, the responsible kinase was not investigated. Since we

previously identified QSK1 as an *in vivo* interactor of CRK2 (Hunter et al. 2019), we asked if

CRK2 could phosphorylate QSK1, thereby controlling its plasmodesmal relocalization. We

recombinantly expressed the cytosolic parts of CRK2 (6His-GST-CRK2cyto and 6His-GST-

CRK2^{D450N}cyto) (Hunter et al. 2019) and QSK1 (6His-MBP-QSK1cyto) in *E. coli*. To identify

CRK2-targeted phosphorylation sites in QSK1, we carried out LC-ESI-MS/MS analysis of in-gel

trypsin-digested peptides after the kinase reaction of CRK2cyto and QSK1cyto. The identified *in*

vitro phosphorylation sites are shown in Table S6 and Fig. 6A. Interestingly, Ser-626 was among

the identified phosphosites (Fig. 6A), suggesting that CRK2 was able to directly interact with

and phosphorylate this peptide. Due to the importance of Ser-621 and Ser-626 in osmotic stress

response (Grison *et al.*, 2019; Wu *et al.*, 2019; Wang *et al.*, 2022), we generated a variant of

QSK1 with phospho-negative substitutions at positions 621 and 626 (QSK1$^{S621A-S626A}$) as a

phosphorylation target of CRK2. Since 6His-GST-CRK2cyto and 6His-MBP-QSK1cyto are

close in their molecular weights, we digested 6His-MBP-QSK1cyto and 6His-MBP- QSK1^{S621A-}

^{S626A}cyto with Factor Xa protease to cleave the MBP tag from the MBP-QSK1cyto fusion protein

thereby resulting in a clear resolution of CRK2 and QSK1 cytosolic fragments before the kinase

reaction. The QSK1cyto was phosphorylated by 6His-GST-CRK2cyto but not by the kinase

inactive 6His-GST-CRK2^{D450N}cyto (Fig. 6B) in the *in vitro* kinase assay. Further, we detected an

overall comparable phosphorylation signal of the WT QSK1 and QSK1$^{S621A-S626A}$, probably due

to the multiple other phosphorylation sites present in this part of the protein. (Fig. 6B). Given the

role of QSK1 phosphorylation in its relocalization to PD, its *in vitro* phosphorylation by CRK2

suggested a phosphorylation-dependent regulatory function of CRK2 in controlling the
subcellular localization of QSK1.

To determine this *in planta*, we applied a quantitative phosphoproteomics approach and
compared the levels of QSK1 phosphopeptides isolated from WT and *crk2* plants under control
and salt stress conditions. We detected the dually phosphorylated peptide
LIEEVSHSSGSPNPVSD in both WT and *crk2* under normal and stress conditions. This
phosphopeptide contains the Ser-621 and Ser-626 that have been previously shown to be
phosphorylated in phosphoproteomics studies (Niittylä et al. 2007; Hsu et al. 2009; Chang et al.
2012; Xue et al. 2013). Interestingly, the abundance of the LIEEVSHSSGSPNPVSD
phosphopeptide was reduced following salt treatment in WT but not in *crk2*. Moreover, the
abundance of the phosphopeptide was lower in *crk2* compared to WT under normal growth
conditions (Fig. 6C, Table S7). Lower abundance in *crk2* may suggest that CRK2 phosphorylates
QSK1 in the peptide LIEEVSHSSGSPNPVSD. This finding aligns with the results of the FRET-
FLIM assay where we observed that mannitol stress disrupted the binding of CRK2 and QSK1
and placed a role of phosphorylation-dependent coupling of the components in osmotic stress
signaling.

Phosphorylation of QSK1 at serine 621 and serine 626 inhibits its stress-induced
enrichment at PD

QSK1 relocalization in response to osmotic stress is associated with its phosphorylation status at
Ser-621 and Ser-626 (Grison et al. 2019). SIRK1 phosphorylates QSK1 at Ser-621 and Ser-626
during sucrose-induced osmotic conditions (Wu et al. 2019). Therefore, we examined if QSK1
relocalization to PD was dependent on SIRK1 by testing QSK1-mCherry/*sirk1* plants. We found
that QSK1 exhibited osmotic stress-induced PD enrichment in QSK1-mCherry/*sirk1* plants
without impairment. An enrichment index of 1.967 was observed in *sirk1* following osmotic
stress compared to 1.092 in control conditions (Fig. S5A, B, Table S3). This finding suggested,
that while phosphorylation of Ser-621 and Ser-626 plays an important role in QSK1 recruitment
to PD, SIRK1 does not appear to be the only kinase regulating this process.

Since CRK2 also targeted QSK1 phosphorylation in the phosphopeptide
LIEEVSHSSGSPNPVSD (Fig. 6) and kinase active CRK2 was required for QSK1 recruitment at
PD (Fig. 3C, D), we further explored the role of Ser-621 and Ser-626 phosphorylation in context

of CRK2. We generated transgenic Arabidopsis expressing phosphomimetic QSK1 at Ser-621

and Ser-626 by replacing both serines by glutamic acid and aspartic acid in *crk2* (QSK1$^{(S621D-}$

$^{S626D)}$-mCherry/*crk2*; QSK1$^{(S621E-S626E)}$-mCherry/*crk2*) and in WT background as control

(QSK1$^{(S621D-S626D)}$-mCherry/WT; QSK1$^{(S621E-S626E)}$-mCherry/WT). We found that enrichment of

phosphomimetic QSK1$^{(S621D-S626D)}$-mCherry was negligible and statistically insignificant, in the

WT background and only showed an increase of enrichment index from 1.045 to 1.119 upon

mannitol treatment (Fig. 7A, B). Similarly, negligible relocalization of QSK1$^{(S621D-S626D)}$-

mCherry was observed in the *crk2* mutant background (Fig. 7A, B). Similarly, in the

phosphomimetic variant QSK1$^{(S621E-S626E)}$-mCherry, we observed a mild enrichment of 1.336

following mannitol treatment as compared to 1.017 under control conditions in the WT

background and poor enrichment in the *crk2* mutant background (1.011 to 1.117) (Fig. S5C, D).

This enrichment of QSK1$^{(S621E-S626E)}$-mCherry/WT at PD was still weaker and delayed compared

to that in the QSK1-mCherry/WT plants. These findings were in line with the normal

relocalization of QSK1 in the *sirk1* mutant plants (Fig. S4A, B). This also led us to test the PD

recruitment of non-phosphorylatable QSK1, in which the Ser-621 and Ser-626 were replaced by

alanines, in WT and *crk2* backgrounds (QSK1$^{(S621A-S626A)}$-mCherry/WT; QSK1$^{(S621A-S626A)}$-

mCherry/*crk2*). We observed a higher enrichment of non-phosphorylatable QSK1 at PD in

response to mannitol treatment (Fig. 7C, D). To a lesser extent, non-phosphorylatable QSK1 was

also present at PD even without mannitol treatment (Fig. 7C, D). The enrichment index for

QSK1$^{(S621A-S626A)}$-mCherry/WT in control condition was 1.338 and increased to 2.018 which was

even higher than QSK1-mCherry/WT under mannitol treatment (Fig. 3B). We also validated this

result in QSK1$^{(S621A-S626A)}$-mCherry/*qsk1* lines and the findings were consistent with a very clear

and intense enrichment of QSK1$^{(S621A-S626A)}$-mCherry in the *qsk1* mutant background (Fig. S5E,

F). These findings, together with the unaltered QSK1 relocalization in *sirk1* confirmed that

phosphorylation of QSK1 at Ser-621 and Ser-626 was inhibitory for its recruitment to PD.

Further, attenuated or complete inability of QSK1 relocalization to PD in the *crk2* and

CRK2^{D450N} backgrounds suggested an essential role for CRK2-mediated phosphorylation of

further and yet unknown factor(s) involved in this process.

Phosphorylation of QSK1 at serine 621 and serine 626 suppresses osmotic stress-induced

callose deposition and regulates plant stress response

A positive correlation between QSK1 availability at PD and callose accumulation in the root

epidermal layer is already established (Grison et al. 2019). Since in our conditions, non-

phosphorylatable QSK1 (QSK1$^{(S621A-S626A)}$) was highly enriched at PD in leaf epidermal cells,

we tested the effect of QSK1 recruitment at PD on callose accumulation in the leaf epidermis by

aniline blue staining. Even though the re-organization of RLKs at PM is a relatively fast

response, changes in callose accumulation have a time lag. Therefore, we quantified callose

levels after 7 hours of mannitol treatment to QSK1-mCherry/*qsk1*, QSK1$^{(S621A-S626A)}$-

mCherry/*qsk1* and QSK1$^{(S621D-S626D)}$-mCherry/*qsk1* plants (Fig. 8A-D, Table S8). Plants

expressing WT or non-phosphorylatable QSK1 (QSK1$^{(S621A-S626A)}$) accumulated higher callose

upon mannitol treatment compared to their respective control conditions. However, plants

expressing phosphomimetic QSK1 (QSK1$^{(S621D-S626D)}$) responded weakly to mannitol treatment

(Fig. 8A-D). In control conditions, slightly higher callose levels were detected in the QSK1$^{(S621A-}$

$^{S626A)}$ plants in terms of both intensity and number of deposits compared to the plants with WT

QSK1 (Fig. 8A-D)) which is consistent with a higher basal presence of QSK1$^{(S621A-S626A)}$ at PD

even without stress treatment (Fig. 7C, D), however, the differences in the overall basal callose

accumulation were statistically insignificant among the three genotypes which is consistent with

our finding that QSK1 might not play a regulatory role in basal callose deposition (Fig. 2).

Collectively, these results provide evidence that QSK1 plays a positive role in stress-induced

callose deposition and its relocalization to PD is necessary in this process which is controlled by

changes in its phosphorylation status.

To test the physiological relevance of QSK1 relocalization to PD and callose deposition during

osmotic stress, we carried out root growth assays of plants with WT QSK1 (QSK1-

mCherry/*qsk1*), phospho-negative QSK1 (QSK1$^{(S621A-S626A)}$-mCherry/*qsk1*) and phospho-

mimetic QSK1 (QSK1$^{(S621D-S626D)}$-mCherry/*qsk1*) plants with and without salinity stress. While

the PR length of all three genotypes was similar under control conditions (Fig. 8E, Table S9), the

QSK1$^{(S621A-S626A)}$ plants exhibited maximum retardation in the PR elongation upon salinity stress

compared to their respective control PR growth. In contrast, the phospho-mimetic QSK1 plants

(QSK1$^{(S621D-S626D)}$) showed even higher PR elongation than the plants with WT QSK1 when

subjected to salinity stress for one week (Fig. 8F). The LR density, did not statistically vary

among the genotypes under both control and salt stress conditions (Fig. 8G, H), nevertheless, a

slightly higher LR density was observed in the QSK1$^{(S621A-S626A)}$-mCherry/*qsk1* plants under salt

stress and an opposite trend was observed for the QSK1$^{(S621D-S626D)}$-mCherry/*qsk1* plants (Fig.

8H). We also tested salt sensitivity at the seed germination stage and found that the

phosphorylation state of QSK1 did not impact seed germination in saline environment (Fig. 8I)

implying a functional specificity of QSK1 phosphorylation during post-germination plant

growth. These results established the physiological relevance of the QSK1 relocalization to PD

during osmotic stress and highlighted protein phosphorylation as a regulatory mechanism in

growth adjustments.

Discussion

Symplastic communication through plasmodesmata (PD) is crucial for coordinating plant growth

and stress responses (Stahl and Simon 2013; Otero et al. 2016; Wu et al. 2016; Cheval and

Faulkner 2018; Li et al. 2022). Dynamic regulation of symplastic connectivity is achieved

through PD-associated proteins (Pérez-Sancho et al. 2025), lipid composition (Grison et al.

2015) and callose deposition (Wu et al. 2018). Current knowledge on the changes of callose

accumulation and symplastic communication is largely focused on immune response (Lee et al.

2011; Wang et al. 2013; German et al. 2023; Tee et al. 2023; Li et al. 2025). Studies show altered

symplastic communication during abiotic stress conditions such as metal toxicity (Sivaguru et al.

2000; O'Lexy et al. 2018), low temperature (Bilska and Sowinski 2010) and mechanical

wounding (Cui and Lee 2016). Despite the widespread occurrence of osmotic stress in plants,

understanding of how callose accumulation and PD connectivity respond to it remains limited

(Grison et al. 2019; Hunter et al. 2019; Sangi et al. 2023; Su et al. 2023).

RLKs are involved in sensing and integrating extracellular signals and they typically function in

complexes rather than acting alone. For example, FLS2-BAK1, EFR-BAK1 and CERK1-LYK5

(Chitin elicitor receptor kinase 1- Lysin motif-containing receptor-like kinase 5) (Cao et al. 2014)

in immune signaling (Chinchilla *et al.*, 2007; Mühlenbeck *et al.*, 2024), BRI1-BAK1 in BR

signaling (Nam and Li 2002), SIRK1-QSK1 in regulation of aquaporins (Wu et al. 2019; Wang et

al. 2022). Importantly, the combination of different RLKs can determine the functional outcome.

The RLKs CRK2 and QSK1 are known to regulate plant growth and osmotic stress response

through plasmodesmal regulation (Grison et al. 2019; Hunter et al. 2019), however, how they

coordinate to precisely control callose levels during osmotic stress conditions is so far not

understood. In this study, we identify a regulatory mechanism that precisely fine-tunes callose

deposition during osmotic stress response, a key part of plant growth adaptation to changing
environmental conditions. We report that under normal growth conditions, CRK2 inhibits callose
deposition by engaging QSK1 at the general PM where the CRK2-QSK1 complex potentially
has another role specific to the general PM.

The growth phenotypes of *crk2*, *qsk1* and the *crk2 qsk1* double mutant suggested that CRK2 and
QSK1 are important for normal plant growth and stress response. The *crk2* mutant exhibits
strongest phenotypic alterations compared to WT Arabidopsis which may be due to pleotropic
functions of CRK2 (Bourdais et al. 2015; Hunter et al. 2019; Kimura et al. 2020; Colina et al.
2025). The smaller rosette size and salt hypersensitivity during germination in the *crk2* mutant
was rescued in the *crk2 qsk1* double mutant suggesting the antagonistic roles of CRK2 and
QSK1 in the biological processes that result in observed phenotypes. Further, as shown
previously, mutants defective in callose regulation have defects in LR patterning and PR length
due to altered symplastic connection (Benitez-Alfonso et al. 2013; J Maule et al. 2013; Hunter et
al. 2019). In our analyses, the root phenotype of the *crk2 qsk1* double mutant was milder
compared to the *crk2* and did not rescue to the WT level implying that the absence of both
functional CRK2 and QSK1 results in defects in processes regulating PR growth and LR
emergence.

Further, our callose estimation results revealed antagonistic roles of CRK2 and QSK1 in
regulating callose deposition especially under osmotic stress conditions. In *crk2*, callose
deposition was higher compared to WT suggesting an inhibitory role of CRK2 on callose
deposition. This could result from multiple yet unknown actions of CRK2. For example, CRK2-
targeted phosphorylation of CALLOSE SYNTHASES (CALS) (Hunter et al. 2019) could be one
of the regulatory mechanisms determining callose levels. Further, CRK2 also regulates ROS
production through RBOHD (Kimura et al. 2020) which could add another layer of CRK2-
mediated regulation of callose production. In the *qsk1* mutant, on the other hand, a reduced
callose deposition upon mannitol treatment suggested that the inhibitory action of CRK2 is
ensued following osmotic stress despite the absence of QSK1 which gets further support from
the result that QSK1 was not found to be necessary to recruit CRK2 at the PD. In the *crk2 qsk1*
double mutant, the WT-like response indicated alleviation of the inhibitory effect of CRK2 and

QSK1 on each other and possibly the presence of compensatory or feedback mechanisms that
maintain callose homeostasis.

Protein phosphorylation is a common regulatory mechanism across living organisms for
dynamically reshaping the binding repertoire of a given protein and thereby providing functional
specificity (Wang et al. 2011; Skinner et al. 2017). The changes in the protein-protein interaction
networks by phosphorylation can also lead to changes in the localization and protein abundance
(Bah et al. 2015; Perraki et al. 2018). Phosphorylated QSK1 at Ser-621 and Ser-626 has been
identified in several phosphoproteomics analyses (Niittylä et al. 2007; Hsu et al. 2009; Chang et
al. 2012) implying a regulatory role of this phosphorylation in QSK1 function. For example,
SIRK1, an LRR RLK is known to phosphorylate QSK1 at Ser-621 and Ser-626 which leads to
stabilization of SIRK1-QSK1 protein complex involved in the regulation of aquaporins in roots
(Wu et al. 2019). QSK1 phosphorylation at these two residues also enhances its interaction with
the nitrate transceptor NRT1.1 which eventually transduces low nitrate signal to PM H^+-ATPase
AHA2 (Zhu et al. 2024).

Through *in vitro* kinase assays, we identified that CRK2 phosphorylates multiple residues in the
cytosolic region of QSK1 including Ser-626. Further, *in planta* phosphorylation status of QSK1
within LIEEVSHSSGSPNPVSD was altered in the *crk2* mutant. QSK1 has previously been
shown to be an *in planta* interactor of CRK2 (Hunter et al. 2019), and FRET-FLIM analyses
together with structural modeling supported that CRK2 and QSK1 exist in a preformed complex
at the bulk PM (Fig. 9 panel A). Interestingly, the CRK2 and QSK1 affinity was abolished when
CRK2 was replaced by CRK2^{D450N}. Together, these results and existing literature suggested that
phosphorylation of QSK1 at Ser-621 and Ser-626 could be a common regulatory mechanism to
stabilize QSK1 protein complexes, including the CRK2-QSK1 complex. *In vivo*
phosphoproteomics analyses suggested reduced phosphorylation of the
LIEEVSHSSGSPNPVSD peptide at two serine residues following salt treatment in WT.
Consistently, mannitol treatment reduced the binding affinity of the CRK2-QSK1 complex in the
FRET-FLIM assays. These findings suggest, that an unknown protein phosphatase might act on
QSK1 to allow dissociation of the CRK2-QSK1 complex (Fig. 9 panel B) leading to the PD
localization of QSK1 prior to the relocalization of CRK2. Consistently, our microscopy-based
relocalization studies showed that phosphorylation of QSK1 at Ser-621 and Ser-626 is inhibitory

for the recruitment of QSK1 to PD and consequently callose accumulation following mannitol

treatment. Non-phosphorylatable QSK1$^{S621A-S626A}$, efficiently relocalized to PD while QSK1

phosphomimetic substitutions at Ser-621 and Ser-626 abated its recruitment to PD. Collectively,

these results demonstrated a phosphorylation-dependent subcellular dichotomy for QSK1

conditionally associating with and dissociating from CRK2 and probably other proteins. This

mechanism could direct QSK1 towards distinct molecular pathways to promote growth

adaptability under osmotic stress.

QSK1 was recruited to PD following osmotic stress in a CRK2-dependent manner. The non-

phosphorylatable QSK1$^{S621A-S626A}$, the preferred form of QSK1 to relocalize to PD, also did not

relocalize efficiently in the *crk2* mutant background. Furthermore, enzymatically active CRK2

was essential for the recruitment of QSK1 to PD. This suggested the possibility that CRK2 might

phosphorylate another protein(s) in addition to QSK1, whose phosphorylation by CRK2 would

be necessary to facilitate or allow the relocalization of QSK1 to PD (Fig. 9 panel B). The PM

could be biochemically divided into detergent resistant membrane (DRM) and detergent soluble

membrane (DSM) domains, and PD-associated membranes are considered to be DRMs. Proteins

identified in PD proteomes and DRM fractions show a large overlap (Grison et al. 2019) and

QSK1 has been found in several DRM and PD proteomes (compiled by Grison *et al.*, 2019).

Szymanski *et al.*, 2015 showed that the actin cytoskeleton is necessary for the distribution of

proteins between DRM and DSM and disruption of the actin cytoskeleton depletes QSK1 from

DRMs (Szymanski et al. 2015). We previously identified actin isoforms, as well as other DRM-

localized proteins as interactors of CRK2 in pull-down assays followed by MS analyses (Hunter

et al. 2019). Therefore, we speculate on regulation of the actin cytoskeleton or other DRM-

localized proteins by CRK2 to enable QSK1 relocalization after its dephosphorylation, however,

more studies are required to support this.

Adjusting root system architecture is one of the main strategies plants employ during extended

periods of salinity stress for efficient water uptake (Julkowska et al. 2014). QSK1

phosphorylation at Ser-621 and Ser-626 affects callose deposition in the epidermal layer of root

(Grison et al. 2019) as well as leaves resulting in altered symplastic connectivity and therefore

root phenotype. Adjusting callose levels through QSK1 can therefore be a contributory

mechanism plants employ for osmotic stress resilience. In our physiological analyses, we found

that the plants accumulating higher callose levels upon osmotic stress, exhibit a root phenotype
adjusted for osmotic stress conditions.

Based on these findings, we propose a working model in which symplastic connectivity is
dynamically regulated through temporally coordinated signaling between CRK2 and QSK1 (Fig.
9). We propose that CRK2 works as a moderator of QSK1-mediated osmotic stress responses.
CRK2 phosphorylates QSK1 to form a stable protein complex at the general PM. This complex
might have a function specific to their loacalization at the general PM. At the same time,
existence of such a complex inhibits unnecessary QSK1 function at PD in the absence of osmotic
stress. Dephosphorylation of QSK1 at the C-terminus upon stress stimuli dissociates it from
CRK2 allowing it to relocalize to PD to positively regulate callose deposition. At a later stage,
CRK2 moves to PD itself where it finds different phosphorylation targets than at the PM and
eventually negatively regulates callose deposition to switch it off and prevent unnecessary
clogging of PD. In conclusion, our work shows that CRK2 controls the dynamic and stimulus-
dependent relocalization of QSK1, and possibly other proteins, to PD. PD and other membrane
nanodomains contain constitutive protein components (Jaillais et al. 2024), however, the
dynamics of QSK1 and CRK2 demonstrate the necessity and potential for spatiotemporal
regulation of the cellular response machinery. In the future, it will be highly interesting to
investigate the spatiotemporal dynamics of other proteins showing conditional clustering in
nanodomains vs. the general plasma membrane.

Material and methods

Plant materials and growth conditions

The *Arabidopsis thaliana* ecotype Col-0 was used as wild-type control. The seeds of mutants
qsk1 (SALK_019840; AT3G02880), *crk2* (SALK_012659C; At1g70520), and *sirk1*
(SALK_125543C; At5g10020) were obtained from NASC (University of Nottingham). The *crk2*
qsk1 double mutant was generated by crossing *qsk1* and *crk2*. For growing plants under sterile
conditions, seeds were surface-sterilized and stratified in dark for 3 days at 4°C and plated on
0.5X Murashige and Skoog media (Sigma-Aldrich) supplemented with 0.8% (w/v) agar, 1%
(w/v) Suc, and 0.1% (w/v) MES, pH 5.8. This media composition was used as standard growth
media for all the *in vitro* experiments unless specified. Plants were grown in climate-controlled
growth chambers with a 16-h light, 8-h dark photoperiod. For experiments with adult stages,

seed propagation, and growth of selected transgenics, the plants were grown in substrate mix (2:1

peat:vermiculite) in a Weiss Gallenkamp growth chamber or greenhouse with long-day

conditions at 21°C.

Physiological assays

For comparative phenotypic analysis of WT, *crk2*, *qsk1*, and *crk2 qsk1* plants, 10 plants for each

genotype were planted in individual pots with substrate mix (2:1 peat:vermiculite). Plants were

allowed to grow in a Weiss Gallenkamp growth chamber with long-day conditions at 21°C with

equal irrigation and randomization of pot positioning. Images were captured at various growth

stages. Related data and statistical analysis are given in Table S1.

For root elongation assays, 5-day-old seedlings grown in vertically positioned square petri plates

in Photon Systems Instruments (PSI) growth chamber were transferred to standard 0.5X MS

medium supplemented with 100 mM NaCl and the primary root tip was marked. For the control

group, the seedlings were transferred to medium without NaCl. The images were captured after

another 7 days and the root elongation was measured with the program Fiji

(https://imagej.net/software/fiji/). The data obtained were the average of at least 13-15 seedlings

and all the experiments were repeated three times unless specified. Related data and statistical

analysis are given in Table S1.

For the seed germination assay, surface-sterilized seeds were plated evenly in petri dishes with

standard 0.5X MS medium supplemented with 150 mM NaCl. For the control group, the seeds

were sown on the medium without NaCl. The seedlings were allowed to grow in a PSI growth

chamber and the images of the plates were captured after 12 days. For each genotype and

treatment, 3 plates were taken as biological replicates and in each plate 30-35 seedlings were

taken. The percentage of seed germination was first estimated in control plates to normalize the

effect of non-viable seeds. The total number of seeds and the viable germinated seeds with

clearly visible root and green cotyledon/leaves were counted. Related data and statistical analysis

are given in Table S1.

Osmotic stress treatment and callose estimation

For estimation of osmotic stress-induced callose deposition, seeds were sown hydroponically for

two weeks on a mesh in liquid 0.5X MS with 1% sucrose in standard growth conditions. For

inducing callose deposition under osmotic stress, the roots were subjected to 50 mM mannitol in

liquid medium for 7 hours. After treatment, the plants were pulled out of the mesh and infiltrated

with 0.2%-0.5% aniline blue (constant for a single experiment) and the first pair of true leaves

were imaged in a laser confocal microscope. For each genotype, 5 plants were imaged and 5-6

areas were scanned from first pair of true leaves of each plant. The control plants underwent the

same process without mannitol treatment. The mean intensity and number of deposits were

measured using Fiji. To quantify overall callose levels that include both intensity and number of

deposits, a composite metric was derived by multiplying mean intensity of deposits and number

of deposits and dividing it with a constant scaling factor to yield a number suitable for

comparative analysis. This is termed as total callose deposition. Related data and statistical

analysis are given in Table S2 and S8.

Plasmid vector construction and transgenic lines

Plasmid constructs for P_{QSK1}:*QSK1*-mCherry, P_{QSK1}:*QSK1*$^{(S621A-S626A)}$-mCherry, P_{QSK1}:*QSK1*$^{(S621D-}$

$^{S626D)}$-mCherry and P_{QSK1}:*QSK1*$^{(S621E-S626E)}$-mCherry were generated by the MultiSite Gateway

technology (Invitrogen; Thermo Fisher Scientific). The donor clones for promoter regions of

QSK1 and coding sequence of *QSK1* were cloned in the vectors pDONR™P4-P1R and

pDONR™221, respectively. The donor clone for mCherry in the vector pDONR™P2R-P3 was

already available in the laboratory. The donor clones were combined in an LR reaction to obtain

the final constructs in the binary vector pH7m34GW (VIB-UGENT). The phospho-mutant

versions of QSK1 (S621A-S626A, S621D-S626D and S621E-S626E) were generated by

introducing point mutation by DpnI-mediated site-directed mutagenesis (Fisher and Pei 1997).

The primers are listed in Table S10. P_{CRK2}:*CRK2*- mVenus and P_{CRK2}:*CRK2*^{D450N}-mVenus

constructs were available in the binary vector pBm43GW (Invitrogen; Thermo Fisher Scientific)

(Hunter et al 2019). The final clones in the binary vectors were introduced in *Agrobacterium*

tumefaciens GV3101_pSoup and Arabidopsis plants of desired genotypes were transformed by

floral dip method (Clough and Bent 1998). The CRK2/CRK2^{D450N}-mVenus and QSK1-mCherry

co-expressing lines were generated by crossing P_{CRK2}:*CRK2*-mVenus/*crk2* and P_{QSK1}:*QSK1*-

mCherry/*crk2*, and, P_{CRK2}:*CRK2*^{D450N}-mVenus/*crk2* and P_{QSK1}:*QSK1*-mCherry/*crk2* plants.

Transformants were selected on 0.5X MS medium with appropriate antibiotic or herbicide

[(hygromycin 20 ug/ml (CARL ROTH) or Basta 10 ug/ml (DL-phosphinothricin; Duchefa

Biochemie)].

Osmotic stress treatment and live confocal imaging

For visualizing localization of the fluorescently-tagged proteins in response to osmotic stress, the

mature leaves of 4-week-old Arabidopsis were detached from the plant and submerged in 0.4 M

mannitol or 150 mM NaCl in 0.5X MS solution. For detecting co-localization with aniline blue,

the samples were treated with 0.2% aniline blue simultaneously with mock or 0.4 M mannitol

and 150 mM NaCl treatments. After the incubation period, the same area of all the leaf samples

was targeted for imaging and mounted in the same solution and the abaxial side was imaged. A

mock-treated leaf of the same plant was used as a control. All the live-imaging experiments were

conducted on an Olympus FLUOVIEW FV3000 confocal laser scanning microscope with 60.0 X

or 40.0 X oil-immersion objectives and was operated by Olympus FV31S-SW software. The

excitation wavelengths for mCherry, mVenus and DAPI (for aniline blue) were 561, 488 and

405, respectively. The detection wavelengths were 570-620 nm for mCherry, 500-540 nm for

mVenus and 450-550 nm for DAPI. The laser transmittivity was kept constant for a single

experiment and for the comparisons of fluorescence intensities unless specified. The co-

localizations were detected in sequential scanning mode to prevent cross bleeding of emission

spectra.

Quantification of protein enrichment

To quantify the extent of protein enrichment, we used the method described in (Grison et al.

2020) with modification. We assumed that the intense dots of the fluorescently-tagged protein

correspond to PD based on the fact that these dots co-localized with the aniline blue-stained

callose dots in our analysis (Fig. S2B). Under salt and mannitol treatments, the punctate pattern

of CRK2 and QSK1 correspond to their PD localization and previous studies have confirmed this

with co-localization with PD-located protein markers (PDLPs) in addition to aniline blue (Grison

et al. 2019; Hunter et al. 2019). Using Fiji, we selected a circular region of interest (ROI) on such

intensity spots resembling PD. The ROI adjacent to the enrichment spots was considered as bulk

PM and the ratio of the two was referred to as 'enrichment index'. In each image, 5-6 pairs of

such ROIs were measured for the intensities and enrichment indexes were determined. 3 images

for each treatment/genotype were analyzed. The relocalization experiments were conducted on 3

independent transgenic lines for each genotype. The data related to enrichment index and
statistical analysis are given in Table S3.

FRET-FLIM

The experiments on Förster resonance energy transfer detected by fluorescence lifetime (FRET-
FLIM) were performed on the confocal microscope FV3000 (Olympus) equipped with the 40.0
X UApo N water immersion objective lens NA 1.15 (Olympus), MultiHarp 150 Time-Correlated
Single Photon Counting device, pulsed 488 nm laser, and two Hybrid Photomultiplier detectors
(all Picoquant). All data were acquired at 40 MHz pulse repetition rate. Emitted fluorescence was
filtered using the FF02-525/40-25 emission filter (Semrock). Data analysis was done using
Picoquant SymphoTime64 software package. Membrane regions were manually selected in the
images, and lifetime (τ) values were calculated from decay curves using exponential tailfit with
the fixed time window for analysis. The acceptable goodness of fit χ2 values were between 0.9
and 1.2. The stage of the plants and treatment used in FRET-FLIM analysis are as described in
the method section 'osmotic stress treatment and live confocal imaging'. Related data and
statistical analysis are given in Table S4.

Structural modeling

621 The interactions of full-length and truncated CRK2 and QSK1 were predicted using Colabfold
622 (version 1.5.2) implementation of AlphaFold2 (Jumper et al. 2021) multimer version 1 with a
623 maximum of 48 recycles or 0.5 tolerance. The amino acid boundaries selected for both the
624 proteins are given in Table S5.

Identification of phosphopeptides from *in vitro* kinase assay

The cytosolic region of QSK1 coding sequence QSK1cyto was amplified and cloned into pMal-
c2X (Addgene, #75286) using *BamH*I and *Sal*I restriction sites. The primers are given in Table
S10. The phospho-negative QSK1cyto(S621A-S626A) was generated by *Dpn*I-mediated site-
directed mutagenesis and cloned in the same way as for the WT construct. The construct for the
cytosolic region of CRK2 was available (CRK2cyto-pOPINM; Kimura et al. 2020). GST-tagged
CRK2cyto and MBP-tagged QSK1cyto were expressed in the *E. coli* strain SHuffle T7 (New
England Biolabs, C3026J) and BL21-CodonPlus-RIL (Agilent, 230240), respectively, and
eventually purified with glutathione sepharose 4B (GE Healthcare, 17-0756-01) and amylose

resin (New England Biolabs E8021S) beads, respectively. 10mM maltose was used for the elution of MBP-tagged proteins, while 20mM reduced glutathione for the elution of GST-tagged proteins. In vitro kinase assay was performed as described in Kimura et al. 2020, with minor modifications. Briefly, expressed and purified recombinant proteins were incubated in the ratio 1: 3-5 (kinase: substrate) for 30 min at room temperature in reaction buffer containing γ-32P labelled ATP (0.2µl/20µl of labelled ATP with declared activity 9.25MBq), 50mM HEPES (pH 7.4), 1mM DTT, 10mM MgCl2, 60µM unlabeled ATP. Any specific procedure applied to a specific Figure panel is given in its Figure legend. Enzymatic reactions were terminated by boiling the samples in protein loading dye (98°C, 5 minutes) and proteins were separated by SDS-PAGE. Gels were stained with Coomassie Brilliant Blue and dried. Dried gels were exposed to photosensitive layer and documented with imaging system Typhoon 9410 (Molecular Dynamics). For the purpose of the identification of phosphorylation sites, labelled ATP was replaced by the respective volume of deionized water. Separation and staining of the gels were followed by in-gel trypsin digestion (Thermo Fisher Scientific, 90057). Digested peptide samples were dissolved in 10 µl of 0.1% formic acid and 5 µl was injected for LC-ESI-MS/MS analysis. The LC-ESI-MS/MS analysis was performed on a nanoflow HPLC system (Easy-nLC1000, Thermo Fisher Scientific) coupled to the Q Exactive HF mass spectrometer (Thermo Scientific, Bremen, Germany) equipped with a nano-electrospray ionization source. Peptides were first loaded on a trapping column and subsequently separated inline on a 15 cm C18 column (75 µm x 15 cm, ReproSil-Pur 3 µm 120 Å C18-AQ, Dr. Maisch HPLC GmbH, Ammerbuch-Entringen, Germany). The mobile phase consisted of water with 0.1% formic acid (solvent A) and acetonitrile/water (80:20 (v/v)) with 0.1% formic acid (solvent B). Peptides were eluted using a 30 min gradient: 5% to 21% solvent B in 11 min, 21% to 36% solvent B in 9 min, and 36% to 100% solvent B in 5 min, followed by a 5 min wash with 100% solvent B. MS data was acquired automatically by using Thermo Xcalibur 4.1 software (Thermo Scientific). The data-dependent acquisition method consisted of an Orbitrap MS survey scan of mass range 350–1750 m/z with a resolution of 120000 and AGC target of 3000000. This was followed by HCD fragmentation for 10 most intense peptide ions with resolution of 15000 and AGC target of 5000. Raw files were searched for protein identifications using Proteome Discoverer 2.5 software (Thermo Fisher Scientific) connected to an in-house Mascot 2.8.2 server (Matrix Science). Data was searched

against a SwissProt database (version 2022_03) with taxonomy restricted to *Arabidopsis*

thaliana.

Phosphoproteomics

Plant treatment- Arabidopsis seeds were sterilized by incubating with 1.5% NaClO 0.02%

Triton X-100 solution for 5 min and vernalized at 4 °C for 2 days. Sterilized seeds were

germinated and grown in liquid culture on 12 well plates (30 seeds/well) in 1/2 MS medium with

1% sucrose and 0.5g MES/l at 23 °C under continuous light (100 µmol m^{-2} s^{-1}) in a Percival

growth chamber. Plates with 7-day old seedlings were transferred from the growth chamber to a

workbench and kept o/n for acclimatization before treatments. Seedlings were treated with either

150 mM NaCl or sterile water for 30 minutes after which seedlings were immediately collected

and flash-frozen in liquid nitrogen and stored at -80 °C. ***Sample preparation and***

phosphopeptide enrichment- Frozen seedlings were disrupted using a Retsch mill with 2 steel

beads at 50 Hz for 5 min. To the finely ground material was added 500 µL urea extraction buffer

(8M in 100 mM Tris-HCl pH 8.5, 20 µL/mL phosphatase inhibitor cocktail 2 (Sigma, P5726), 20

µL/mL phosphatase inhibitor cocktail 3 (Sigma, P0044), 5 mM DDT), samples were vortexed

thoroughly to ensure wetting of all powder while thawing. Samples were incubated at RT for 30

min with shaking. Next samples were centrifuged for 5 min at 10k x g, supernatants were

transferred to fresh tubes and centrifuged at full speed for 10 mins. This procedure was repeated

until no visible pellet was retained. Protein concentration of the extract was determined using

Pierce 660 mn protein assay. Samples were alkylated using chloroacetamide (CAA) (550 mM

stock, 14 mM final concentration) for 30 min at RT in the dark, after which samples were

subjected to an in-solution digestion. In brief, samples (400 µL) were diluted with 400 µL

100mM Tris, pH 8.5, 1mM CaCl$_2$, then 5 µg Lys-C (Merck) was added. Samples were incubated

for 4h at room temperature, after which 2.4 mL 100mM Tris, pH 8.5, 1 mM CaCl$_2$ and 5 µg

trypsin were added. Samples were incubated o/n at 37 °C. Samples were acidified with 20 µL

TFA and desalted using C18 SepPaks (1cc cartridge, 100 mg (WAT023590)). In brief, SepPaks

were conditioned using methanol (1 mL), buffer B (80% acetonitrile, 0.1% TFA) (1 mL) and

buffer A (0.1% TFA) (2 mL). Samples were loaded by gravity flow, washed with buffer A (1 x 1

mL, 1x 2 mL) and eluted with buffer B (2 x 400 µL). 40 µL of eluates were used for peptide

measurement and concentrated as input control. For phosphopeptide enrichment by metal-oxide

chromatography (MOC) (adapted from: Nakagami 2014) the samples were evaporated to a

sample volume of 50 µL and diluted with sample buffer (2 mL AcN, 820 µL lactic acid (LA), 2.5

µL TFA / 80% ACN, 0.1% TFA, 300mg/ml LA, final concentrations) (282 µL). MOC tips were

prepared by loading 100 µL of a well-mixed slurry of 15mg metal oxide beads (Titansphere TiO_2

beads 10 µm (GL Science Inc, Japan, Cat. No. 5020-75010) in 500 µL MeOH onto 4 C2 micro

columns/condition (3 mg/column) and centrifugation for 5 min at 1500g. Tips were washed with

centrifugation at 1500g for 5 min using 80 µL of solution B (80% acetonitrile, 0.1% TFA) and 80

µL of solution C (300 mg/mL LA in solution B). To simplify the processing, samples tips were

fitted onto a 96/500 µL deep well plate (Protein LoBind, (Eppendorf Cat. No. 0030504100).

After washing MOC tips were transferred to a fresh plate, samples were loaded onto the

equilibrated tips and centrifuged for 10 min at 1000g. The flow through was reloaded onto the

tips and centrifugation was repeated. Tips were washed with centrifugation at 1500g for 5 min

using 80 µL of solution C and 3x 80 µL of solution B. For the elution of the enriched

phosphopeptides the tips were transferred to a fresh 96/500 µL deep well plate containing 100

µL/well of acidification buffer (20% phosphoric acid). Peptides were eluted first with 50 µL

elution buffer 1 (5% NH_4OH) and centrifugation for 5 min at 800g, then with 50 µL of elution

buffer 2 (10% piperidine) and centrifugation for 5 min at 800g. Next, the samples were desalted

using StageTips with C18 Empore disk membranes (3 M) (Rappsilber et al. 2003), dried in a

vacuum evaporator, and dissolved in 10 µL 2% ACN, 0.1% TFA (A* buffer) for MS analysis.

Data acquisition- Samples were analysed using an Ultimate 3000 RSLC nano (Thermo Fisher)

coupled to an Orbitrap Exploris 480 mass spectrometer equipped with a FAIMS Pro interface for

Field asymmetric ion mobility separation (Thermo Fisher). Peptides were pre-concentrated on an

Acclaim PepMap 100 pre-column (75 µM x 2 cm, C18, 3 µM, 100 Å, Thermo Fisher) using the

loading pump and buffer A** (water, 0.1% TFA) with a flow of 7 µl/min for 5 min. Peptides

were separated on 16 cm frit-less silica emitters (New Objective, 75 µm inner diameter), packed

in-house with reversed-phase ReproSil-Pur C18 AQ 1.9 µm resin (Dr. Maisch). Peptides were

loaded on the column and eluted for 130 min using a segmented linear gradient of 5% to 95%

solvent B (0 min : 5%B; 0-5 min -> 5%B; 5-65 min -> 20%B; 65-90 min ->35%B; 90-100 min -

> 55%; 100-105 min ->95%, 105-115 min ->95%, 115-115.1 min -> 5%, 115.1-130 min ->5%)

(solvent A 0% ACN, 0.1% FA; solvent B 80% ACN, 0.1%FA) at a flow rate of 300 nL/min. Mass

spectra were acquired in data-dependent acquisition mode with a TOP_S method using a cycle

725 time of 2 seconds. For field asymmetric ion mobility separation (FAIMS) two compensation

726 voltages (-45 and -65) were applied, the cycle time for the CV-45 experiment was set to 1.2

727 seconds and for the CV-65 experiment to 0.8 sec. MS spectra were acquired in the Orbitrap

728 analyzer with a mass range of 320–1200 m/z at a resolution of 60,000 FWHM and a normalized

729 AGC target of 300%. Precursors were filtered using the MIPS option (MIPS mode = peptide),

730 the intensity threshold was set to 5000, Precursors were selected with an isolation window of 1.6

731 m/z. HCD fragmentation was performed at a normalized collision energy of 30%. MS/MS

732 spectra were acquired with a target value of 75% ions at a resolution of 15,000 FWHM, at an

733 injection time of 120 ms and a fixed first mass of m/z 120. Peptides with a charge of +1, greater

734 than 6, or with unassigned charge state were excluded from fragmentation for MS^2. ***Data***

735 ***analysis phosphoproteome analysis-***Raw data were processed using MaxQuant software

736 (version 1.6.3.4, http://www.maxquant.org/) (Cox et al., Nat. Biotechnol. 2008, 26, 1367.) with

737 label-free quantification (LFQ) and iBAQ enabled (Cox et al., Nat. Protoc. 2016, 11, 2301.).

738 MS/MS spectra were searched by the Andromeda search engine against a combined database

739 containing the sequences from *A. thaliana* (TAIR10_pep_20101214 database), and sequences of

740 248 common contaminant proteins and decoy sequences. Trypsin specificity was required and a

741 maximum of two missed cleavages allowed. Minimal peptide length was set to seven amino

742 acids. Carbamidomethylation of cysteine residues was set as fixed, phosphorylation of serine,

743 threonine and tyrosine, oxidation of methionine and protein N-terminal acetylation as variable

744 modifications. The match between runs option was enabled. Peptide-spectrum-matches and

745 proteins were retained if they were below a false discovery rate of 1% in both cases. Statistical

746 analysis of the intensity values obtained for the phospho-modified peptides

747 ("modificationSpecificPeptides.txt" output file) was carried out using Perseus (version 1.5.8.5,

748 http://www.maxquant.org/). Intensities were filtered for reverse and contaminant hits and the

749 data was filtered to retain only phospho-modified peptides. Next. intensity values were log2

750 transformed. After grouping samples by condition only those sites were retained for the

751 subsequent analysis that had three valid values in one of the conditions. Two-sample t-tests were

752 performed using a permutation-based FDR of 0.05. Alternatively, the data was filtered for 4 valid

753 values in one of the conditions, the valid value-filtered data was median-normalized and missing

754 values were imputed from a normal distribution, using the default settings in Perseus (1.8

755 downshift, separately for each column). Volcano plots were generated in Perseus using an FDR
756 of 5% an S0=1. The Perseus output was exported and further processed using Excel.

Quantification and statistical analysis

Fiji (ImageJ; (https://imagej.net/software/fiji/) was used for image analyses. The information on
statistical analyses is given in the corresponding figure legends and supplementary tables. Instant
Clue (Nolte et al. 2018) and Graph Pad Prism v10 were used for graph preparation and statistical
analyses. In general, data normality was tested by Shapiro-Wilk test. Kruskal-Wallis test
followed by Dunn's multiple comparison test was used for those datasets that did not pass the
normality test. For normal data, one-way ANOVA with Tukey's post-hoc test was used. For the
datasets with two variables (genotype and treatment), two-way ANOVA was applied with
Tukey's post-hoc multiple comparison test. Any other specific tests are mentioned in figure
legends and corresponding supplementary tables.

Acknowledgments

This work was supported by the Czech Science Foundation grants nr. 23-04866S (MW) and 22-
35680M (RP). FC was funded by EMBO (award number: ALTF 1115-2021; 2022-2024) and the
Czech Ministry of Youth and Education MSCA Fellowships Interfellows (Project number:
CZ.02.01.01/00/22_010/0003414; 2024-2025). We acknowledge Prof. Jiří Friml and Dmitrii
Vladimirtsev (Institute of Science and Technology, Austria) for assistance in FRET-FLIM. We
thank Laboratory of Microscopy and Histology, Biology Centre CAS, supported by project
LM2023050 funded by the Ministry of Education, Youth and Sports of the Czech Republic, with
the instrumental equipment co-financed by the European Union. We also acknowledge Growth
Facility (Jan Kadlec), BC Core Facilities, Institute for Plant Molecular Biology, Biology Centre
CAS. Computational resources used for ColabFold predictions were provided by the e-INFRA
CZ project (ID:90254), supported by the Ministry of Education, Youth and Sports of the Czech
Republic. Identification of phosphopeptides from *in vitro* kinase assay was performed at the
Turku Proteomics Facility supported by Biocenter Finland. We thank Dr. Julia Krasensky-
Wrzaczek for her research input and constructive feedback on the manuscript.

Data availability

All data is available in the supplementary files provided alongside this manuscript, including
used methodologies, statistical procedures, oligonucleotide sequences and *in vitro*
phosphopeptide data. Materials are available from the authors upon request. *In vivo* mass
spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the
PRIDE (Perez-Riverol et al. 2025) partner repository with the dataset identifier PXD070133.

Author contributions

SJ and MW conceived the study and designed the research. AZ and SJ performed *in vitro* kinase
assays and callose staining, imaging and quantification. AB, SJ and IK conducted FRET-FLIM
experiment. SCS, AH and HN conducted and analyzed phosphoproteomics experiment. FC
performed crossing. SJ and SL carried out physiological assays. MN and RP performed structural
analysis. MP and JM carried out LC-MS/MS analyses. SJ conducted all other experiments. SJ
wrote the first draft and MW edited and reviewed the paper. MW acquired the research grant. All
authors participated in this research and contributed to the final draft of the paper.

References

Bah A, Vernon RM, Siddiqui Z, Krzeminski M, Muhandiram R, Zhao C, Sonenberg N, Kay LE, and
Forman-Kay JD. Folding of an intrinsically disordered protein by phosphorylation as a regulatory
switch. Nature. 2015:**519**(7541):106–109. https://doi.org/10.1038/nature13999

Benitez-Alfonso Y, Faulkner C, Pendle A, Miyashima S, Helariutta Y, and Maule A. Symplastic
Intercellular Connectivity Regulates Lateral Root Patterning. Developmental Cell.
2013:**26**(2):136–147. https://doi.org/10.1016/j.devcel.2013.06.010

Bilska A and Sowinski P. Closure of plasmodesmata in maize (Zea mays) at low temperature: a new
mechanism for inhibition of photosynthesis. Ann Bot. 2010:**106**(5):675–686.
https://doi.org/10.1093/aob/mcq169

Bourdais G, Burdiak P, Gauthier A, Nitsch L, Salojärvi J, Rayapuram C, Idänheimo N, Hunter K, Kimura S,
Merilo E, et al. Large-Scale Phenomics Identifies Primary and Fine-Tuning Roles for CRKs in
Responses Related to Oxidative Stress. PLOS Genetics. 2015:**11**(7):e1005373.
https://doi.org/10.1371/journal.pgen.1005373

Bücherl CA, van Esse GW, Kruis A, Luchtenberg J, Westphal AH, Aker J, van Hoek A, Albrecht C, Borst
JW, and de Vries SC. Visualization of BRI1 and BAK1(SERK3) Membrane Receptor
Heterooligomers during Brassinosteroid Signaling1[W][OPEN]. Plant Physiol. 2013:**162**(4):1911–
1925. https://doi.org/10.1104/pp.113.220152

Bücherl CA, Jarsch IK, Schudoma C, Segonzac C, Mbengue M, Robatzek S, MacLean D, Ott T, and Zipfel
C. Plant immune and growth receptors share common signalling components but localise to

distinct plasma membrane nanodomains. eLife. 2017:**6**:e25114.
https://doi.org/10.7554/eLife.25114

Cao Y, Liang Y, Tanaka K, Nguyen CT, Jedrzejczak RP, Joachimiak A, and Stacey G. The kinase LYK5 is a
major chitin receptor in Arabidopsis and forms a chitin-induced complex with related kinase
CERK1. eLife. 2014:**3**:e03766. https://doi.org/10.7554/eLife.03766

Chang I-F, Hsu J-L, Hsu P-H, Sheng W-A, Lai S-J, Lee C, Chen C-W, Hsu J-C, Wang S-Y, Wang L-Y, et al.
Comparative phosphoproteomic analysis of microsomal fractions of *Arabidopsis thaliana* and
Oryza sativa subjected to high salinity. Plant Science. 2012:**185–186**:131–142.
https://doi.org/10.1016/j.plantsci.2011.09.009

Cheval C and Faulkner C. Plasmodesmal regulation during plant–pathogen interactions. New Phytologist.
2018:**217**(1):62–67. https://doi.org/10.1111/nph.14857

Chinchilla D, Shan L, He P, de Vries S, and Kemmerling B. One for all: the receptor-associated kinase
BAK1. Trends Plant Sci. 2009:**14**(10):535–541. https://doi.org/10.1016/j.tplants.2009.08.002

Chinchilla D, Zipfel C, Robatzek S, Kemmerling B, Nürnberger T, Jones JDG, Felix G, and Boller T. A
flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. Nature.
2007:**448**(7152):497–500. https://doi.org/10.1038/nature05999

Clough SJ and Bent AF. Floral dip: a simplified method for *Agrobacterium* -mediated transformation of
Arabidopsis thaliana. The Plant Journal. 1998:**16**(6):735–743. https://doi.org/10.1046/j.1365-
313x.1998.00343.x

Colina FJ, Krasensky-Wrzaczek J, Pérez-Guillén A, and Wrzaczek M. CRK2 modulates flowering in
Arabidopsis together with GLYCINE-RICH RNA BINDING PROTEIN 7 (GRP7).
2025:2025.08.05.668606. https://doi.org/10.1101/2025.08.05.668606

Cui W and Lee J-Y. Arabidopsis callose synthases CalS1/8 regulate plasmodesmal permeability during
stress. Nat Plants. 2016:**2**(5):16034. https://doi.org/10.1038/nplants.2016.34

Fisher CL and Pei GK. Modification of a PCR-based site-directed mutagenesis method. Biotechniques.
1997:**23**(4):570–571, 574. https://doi.org/10.2144/97234bm01

German L, Yeshvekar R, and Benitez-Alfonso Y. Callose metabolism and the regulation of cell walls and
plasmodesmata during plant mutualistic and pathogenic interactions. Plant Cell Environ.
2023:**46**(2):391–404. https://doi.org/10.1111/pce.14510

Goto Y, Kadota Y, Mbengue M, Lewis JD, Matsui H, Maki N, Ngou BPM, Sklenar J, Derbyshire P, Shibata
A, et al. The leucine-rich repeat receptor kinase QSK1 regulates PRR-RBOHD complexes targeted
by the bacterial effector HopF2Pto. Plant Cell. 2024:**36**(12):4932–4951.
https://doi.org/10.1093/plcell/koae267

Grison MS, Brocard L, Fouillen L, Nicolas W, Wewer V, Dörmann P, Nacir H, Benitez-Alfonso Y, Claverol
S, Germain V, et al. Specific membrane lipid composition is important for plasmodesmata
function in Arabidopsis. The Plant Cell. 2015:**27**(4):1228–1250.
https://doi.org/10.1105/tpc.114.135731

Grison MS, Kirk P, Brault ML, Wu XN, Schulze WX, Benitez-Alfonso Y, Immel F, and Bayer EM. Plasma
Membrane-Associated Receptor-like Kinases Relocalize to Plasmodesmata in Response to
Osmotic Stress1[OPEN]. Plant Physiol. 2019:**181**(1):142–160.
https://doi.org/10.1104/pp.19.00473

Grison MS, Petit JD, Glavier M, and Bayer EM. Quantification of Protein Enrichment at Plasmodesmata.
Bio Protoc. 2020:**10**(5):e3545. https://doi.org/10.21769/BioProtoc.3545

Han X, Hyun TK, Zhang M, Kumar R, Koh E, Kang B-H, Lucas WJ, and Kim J-Y. Auxin-Callose-Mediated
Plasmodesmal Gating Is Essential for Tropic Auxin Gradient Formation and Signaling.
Developmental Cell. 2014:**28**(2):132–146. https://doi.org/10.1016/j.devcel.2013.12.008

Hdedeh O, Mercier C, Poitout A, Martinière A, and Zelazny E. Membrane nanodomains to shape plant
cellular functions and signaling. New Phytologist. 2025:**245**(4):1369–1385.
https://doi.org/10.1111/nph.20367

Hsu J-L, Wang L-Y, Wang S-Y, Lin C-H, Ho K-C, Shi F-K, and Chang I-F. Functional phosphoproteomic
profiling of phosphorylation sites in membrane fractions of salt-stressed Arabidopsis thaliana.
Proteome Sci. 2009:**7**:42. https://doi.org/10.1186/1477-5956-7-42

Huang C, Mutterer J, and Heinlein M. In Vivo Aniline Blue Staining and Semiautomated Quantification of
Callose Deposition at Plasmodesmata. Methods Mol Biol. 2022:**2457**:151–165.
https://doi.org/10.1007/978-1-0716-2132-5_9

Hunter K, Kimura S, Rokka A, Tran HC, Toyota M, Kukkonen JP, and Wrzaczek M. CRK2 Enhances Salt
Tolerance by Regulating Callose Deposition in Connection with PLDα1. Plant Physiol.
2019:**180**(4):2004–2021. https://doi.org/10.1104/pp.19.00560

Isner JC, Begum A, Nuehse T, Hetherington AM, and Maathuis FJM. KIN7 Kinase Regulates the Vacuolar
TPK1 K+ Channel during Stomatal Closure. Current Biology. 2018:**28**(3):466-472.e4.
https://doi.org/10.1016/j.cub.2017.12.046

J Maule A, Gaudioso-Pedraza R, and Benitez-Alfonso Y. Callose deposition and symplastic connectivity
are regulated prior to lateral root emergence. Commun Integr Biol. 2013:**6**(6):e26531.
https://doi.org/10.4161/cib.26531

Jaillais Y, Bayer E, Bergmann DC, Botella MA, Boutté Y, Bozkurt TO, Caillaud M-C, Germain V,
Grossmann G, Heilmann I, et al. Guidelines for naming and studying plasma membrane domains
in plants. Nat Plants. 2024:**10**(8):1172–1183. https://doi.org/10.1038/s41477-024-01742-8

Jaillais Y and Ott T. The Nanoscale Organization of the Plasma Membrane and Its Importance in
Signaling: A Proteolipid Perspective1[OPEN]. Plant Physiol. 2020:**182**(4):1682–1696.
https://doi.org/10.1104/pp.19.01349

Jarsch IK, Konrad SSA, Stratil TF, Urbanus SL, Szymanski W, Braun P, Braun K-H, and Ott T. Plasma
Membranes Are Subcompartmentalized into a Plethora of Coexisting and Diverse Microdomains
in Arabidopsis and Nicotiana benthamiana. Plant Cell. 2014:**26**(4):1698–1711.
https://doi.org/10.1105/tpc.114.124446

Julkowska MM, Hoefsloot HCJ, Mol S, Feron R, de Boer G-J, Haring MA, and Testerink C. Capturing
Arabidopsis root architecture dynamics with ROOT-FIT reveals diversity in responses to salinity.
Plant Physiol. 2014:**166**(3):1387–1402. https://doi.org/10.1104/pp.114.248963

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek
A, Potapenko A, et al. Highly accurate protein structure prediction with AlphaFold. Nature.
2021:**596**(7873):583–589. https://doi.org/10.1038/s41586-021-03819-2

Katagiri T, Takahashi S, and Shinozaki K. Involvement of a novel Arabidopsis phospholipase D,
AtPLDdelta, in dehydration-inducible accumulation of phosphatidic acid in stress signalling. Plant
J. 2001:**26**(6):595–605. https://doi.org/10.1046/j.1365-313x.2001.01060.x

Kimura S, Hunter K, Vaahtera L, Tran HC, Citterico M, Vaattovaara A, Rokka A, Stolze SC, Harzen A,
Meißner L, et al. CRK2 and C-terminal Phosphorylation of NADPH Oxidase RBOHD Regulate
Reactive Oxygen Species Production in Arabidopsis[OPEN]. Plant Cell. 2020:**32**(4):1063–1080.
https://doi.org/10.1105/tpc.19.00525

Knight H, Trewavas AJ, and Knight MR. Calcium signalling in Arabidopsis thaliana responding to drought
and salinity. Plant J. 1997:**12**(5):1067–1078. https://doi.org/10.1046/j.1365-
313x.1997.12051067.x

Lee J-Y, Wang X, Cui W, Sager R, Modla S, Czymmek K, Zybaliov B, van Wijk K, Zhang C, Lu H, et al. A
plasmodesmata-localized protein mediates crosstalk between cell-to-cell communication and
innate immunity in Arabidopsis. Plant Cell. 2011:**23**(9):3353–3373.
https://doi.org/10.1105/tpc.111.087742

Li J, Lu P, Pan Q, Wang B, Wang Y, and Li J. ER-PM tether Syt1 limits cell-to-cell connectivity via
plasmodesmata during innate immune responses in Arabidopsis. Cell Reports. 2025:**44**(5).
https://doi.org/10.1016/j.celrep.2025.115672

Li M, Wang M, Lin Q, Wang M, Niu X, Cheng J, Xu M, Qin Y, Liao X, and Wu JX and S. Symplastic
communication in the root cap directs auxin distribution to modulate root development. Journal
of Integrative Plant Biology. 2022:**64**(4):859. https://doi.org/10.1111/jipb.13237

Martinière A, Fiche JB, Smokvarska M, Mari S, Alcon C, Dumont X, Hematy K, Jaillais Y, Nollmann M,
and Maurel C. Osmotic Stress Activates Two Reactive Oxygen Species Pathways with Distinct
Effects on Protein Nanodomains and Diffusion. Plant Physiol. 2019:**179**(4):1581–1593.
https://doi.org/10.1104/pp.18.01065

Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, and Steinegger M. ColabFold: making protein
folding accessible to all. Nat Methods. 2022:**19**(6):679–682. https://doi.org/10.1038/s41592-
022-01488-1

Mühlenbeck H, Tsutsui Y, Lemmon MA, Bender KW, and Zipfel C. Allosteric activation of the co-receptor
BAK1 by the EFR receptor kinase initiates immune signaling. eLife. **12**:RP92110.
https://doi.org/10.7554/eLife.92110

Munnik T, Meijer HJ, Ter Riet B, Hirt H, Frank W, Bartels D, and Musgrave A. Hyperosmotic stress
stimulates phospholipase D activity and elevates the levels of phosphatidic acid and

diacylglycerol pyrophosphate. Plant J. 2000:**22**(2):147–154. https://doi.org/10.1046/j.1365-
313x.2000.00725.x

Nakagami H. StageTip-based HAMMOC, an efficient and inexpensive phosphopeptide enrichment
method for plant shotgun phosphoproteomics. Methods Mol Biol. 2014:**1072**:595–607.
https://doi.org/10.1007/978-1-62703-631-3_40

Nam KH and Li J. BRI1/BAK1, a receptor kinase pair mediating brassinosteroid signaling. Cell.
2002:**110**(2):203–212. https://doi.org/10.1016/s0092-8674(02)00814-0

Niittylä T, Fuglsang AT, Palmgren MG, Frommer WB, and Schulze WX. Temporal Analysis of Sucrose-
induced Phosphorylation Changes in Plasma Membrane Proteins of Arabidopsis. Molecular &
Cellular Proteomics. 2007:**6**(10):1711–1726. https://doi.org/10.1074/mcp.M700164-MCP200

Nolte H, MacVicar TD, Tellkamp F, and Krüger M. Instant Clue: A Software Suite for Interactive Data
Visualization and Analysis. Sci Rep. 2018:**8**(1):12648. https://doi.org/10.1038/s41598-018-
31154-6

O'Lexy R, Kasai K, Clark N, Fujiwara T, Sozzani R, and Gallagher KL. Exposure to heavy metal stress
triggers changes in plasmodesmatal permeability via deposition and breakdown of callose. J Exp
Bot. 2018:**69**(15):3715–3728. https://doi.org/10.1093/jxb/ery171

Otero S, Helariutta Y, and Benitez-Alfonso Y. Symplastic communication in organ formation and tissue
patterning. Current Opinion in Plant Biology. 2016:**29**:21–28.
https://doi.org/10.1016/j.pbi.2015.10.007

Perez-Riverol Y, Bandla C, Kundu DJ, Kamatchinathan S, Bai J, Hewapathirana S, John NS, Prakash A,
Walzer M, Wang S, et al. The PRIDE database at 20 years: 2025 update. Nucleic Acids Res.
2025:**53**(D1):D543–D553. https://doi.org/10.1093/nar/gkae1011

Pérez-Sancho J, Smokvarska M, Dubois G, Glavier M, Sritharan S, Moraes TS, Moreau H, Dietrich V,
Platre MP, Paterlini A, et al. Plasmodesmata act as unconventional membrane contact sites
regulating intercellular molecular exchange in plants. Cell. 2025:**188**(4):958-977.e23.
https://doi.org/10.1016/j.cell.2024.11.034

Perraki A, Gronnier J, Gouguet P, Boudsocq M, Deroubaix A-F, Simon V, German-Retana S, Legrand A,
Habenstein B, Zipfel C, et al. REM1.3's phospho-status defines its plasma membrane
nanodomain organization and activity in restricting PVX cell-to-cell movement. PLOS Pathogens.
2018:**14**(11):e1007378. https://doi.org/10.1371/journal.ppat.1007378

Rappsilber J, Ishihama Y, and Mann M. Stop and go extraction tips for matrix-assisted laser
desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in proteomics. Anal
Chem. 2003:**75**(3):663–670. https://doi.org/10.1021/ac026117i

Sangi S, Olimpio GV, Coelho FS, Alexandrino CR, Da Cunha M, and Grativol C. Flagellin and mannitol
modulate callose biosynthesis and deposition in soybean seedlings. Physiologia Plantarum.
2023:**175**(2):e13877. https://doi.org/10.1111/ppl.13877

Sivaguru M, Fujiwara T, Samaj J, Baluska F, Yang Z, Osawa H, Maeda T, Mori T, Volkmann D, and
Matsumoto H. Aluminum-induced 1-->3-beta-D-glucan inhibits cell-to-cell trafficking of
molecules through plasmodesmata. A new mechanism of aluminum toxicity in plants. Plant
Physiol. 2000:**124**(3):991–1006. https://doi.org/10.1104/pp.124.3.991

Skinner JJ, Wang S, Lee J, Ong C, Sommese R, Sivaramakrishnan S, Koelmel W, Hirschbeck M,
Schindelin H, Kisker C, et al. Conserved salt-bridge competition triggered by phosphorylation
regulates the protein interactome. Proceedings of the National Academy of Sciences.
2017:**114**(51):13453–13458. https://doi.org/10.1073/pnas.1711543114

Stahl Y and Simon R. Gated communities: apoplastic and symplastic signals converge at plasmodesmata
to control cell fates. J Exp Bot. 2013:**64**(17):5237–5241. https://doi.org/10.1093/jxb/ert245

Stephan AB, Kunz H-H, Yang E, and Schroeder JI. Rapid hyperosmotic-induced Ca2+ responses in
Arabidopsis thaliana exhibit sensory potentiation and involvement of plastidial KEA transporters.
Proceedings of the National Academy of Sciences. 2016:**113**(35):E5242–E5249.
https://doi.org/10.1073/pnas.1519555113

Su L, Zhang Y, Yu S, Geng L, Lin S, Ouyang L, and Jiang X. RcbHLH59-RcPRs module enhances salinity
stress tolerance by balancing Na+/K+ through callose deposition in rose (Rosa chinensis). Hortic
Res. 2023:**10**(3):uhac291. https://doi.org/10.1093/hr/uhac291

Szymanski WG, Zauber H, Erban A, Gorka M, Wu XN, and Schulze WX. Cytoskeletal Components Define
Protein Location to Membrane Microdomains. Mol Cell Proteomics. 2015:**14**(9):2493–2509.
https://doi.org/10.1074/mcp.M114.046904

Tee EE, Johnston MG, Papp D, and Faulkner C. A PDLP-NHL3 complex integrates plasmodesmal immune
signaling cascades. Proceedings of the National Academy of Sciences.
2023:**120**(17):e2216397120. https://doi.org/10.1073/pnas.2216397120

Testerink C and Munnik T. Molecular, cellular, and physiological responses to phosphatidic acid
formation in plants. J Exp Bot. 2011:**62**(7):2349–2361. https://doi.org/10.1093/jxb/err079

Tracy FE, Gilliham M, Dodd AN, Webb AAR, and Tester M. NaCl-induced changes in cytosolic free Ca2+
in Arabidopsis thaliana are heterogeneous and modified by external ionic composition. Plant Cell
Environ. 2008:**31**(8):1063–1073. https://doi.org/10.1111/j.1365-3040.2008.01817.x

Vatén A, Dettmer J, Wu S, Stierhof Y-D, Miyashima S, Yadav SR, Roberts CJ, Campilho A, Bulone V,
Lichtenberger R, et al. Callose Biosynthesis Regulates Symplastic Trafficking during Root
Development. Developmental Cell. 2011:**21**(6):1144–1155.
https://doi.org/10.1016/j.devcel.2011.10.006

Vu MH, Hyun TK, Bahk S, Jo Y, Kumar R, Thiruppathi D, Iswanto ABB, Chung WS, Shelake RM, and Kim
J-Y. ROS-mediated plasmodesmal regulation requires a network of an Arabidopsis receptor-like
kinase, calmodulin-like proteins, and callose synthases. Front Plant Sci. 2023:**13**.
https://doi.org/10.3389/fpls.2022.1107224

Wang J, Xi L, Wu XN, König S, Rohr L, Neumann T, Weber J, Harter K, and Schulze WX. PEP7 acts as a
peptide ligand for the receptor kinase SIRK1 to regulate aquaporin-mediated water influx and

lateral root growth. Molecular Plant. 2022:**15**(10):1615–1631.
https://doi.org/10.1016/j.molp.2022.09.016

Wang X, Sager R, Cui W, Zhang C, Lu H, and Lee J-Y. Salicylic Acid Regulates Plasmodesmata Closure
during Innate Immune Responses in Arabidopsis[C][W]. Plant Cell. 2013:**25**(6):2315–2329.
https://doi.org/10.1105/tpc.113.110676

Wang Z, Ding G, Geistlinger L, Li H, Liu L, Zeng R, Tateno Y, and Li Y. Evolution of protein
phosphorylation for distinct functional modules in vertebrate genomes. Mol Biol Evol.
2011:**28**(3):1131–1140. https://doi.org/10.1093/molbev/msq268

Wu S, O'Lexy R, Xu M, Sang Y, Chen X, Yu Q, and Gallagher KL. Symplastic signaling instructs cell
division, cell expansion, and cell polarity in the ground tissue of Arabidopsis thaliana roots.
Proceedings of the National Academy of Sciences. 2016:**113**(41):11621–11626.
https://doi.org/10.1073/pnas.1610358113

Wu S-W, Kumar R, Iswanto ABB, and Kim J-Y. Callose balancing at plasmodesmata. J Exp Bot.
2018:**69**(22):5325–5339. https://doi.org/10.1093/jxb/ery317

**Wu XN, Chu L, Xi L, Pertl-Obermeyer H, Li Z, Sklodowski K, Sanchez-Rodriguez C, Obermeyer G, and
Schulze WX**. Sucrose-induced Receptor Kinase 1 is Modulated by an Interacting Kinase with
Short Extracellular Domain*,. Molecular & Cellular Proteomics. 2019:**18**(8):1556–1571.
https://doi.org/10.1074/mcp.RA119.001336

Xu B, Cheval C, Laohavisit A, Hocking B, Chiasson D, Olsson TSG, Shirasu K, Faulkner C, and Gilliham M.
A calmodulin-like protein regulates plasmodesmal closure during bacterial immune responses.
New Phytologist. 2017:**215**(1):77–84. https://doi.org/10.1111/nph.14599

**Xu Z, Schahl A, Jolivet M-D, Legrand A, Grélard A, Berbon M, Morvan E, Lagardere L, Piquemal J-P,
Loquet A, et al.** Dynamic pre-structuration of lipid nanodomain-segregating remorin proteins.
Commun Biol. 2024:**7**(1):1620. https://doi.org/10.1038/s42003-024-07330-y

Xue L, Wang P, Wang L, Renzi E, Radivojac P, Tang H, Arnold R, Zhu J-K, and Tao WA. Quantitative
measurement of phosphoproteome response to osmotic stress in arabidopsis based on Library-
Assisted eXtracted Ion Chromatogram (LAXIC). Mol Cell Proteomics. 2013:**12**(8):2354–2369.
https://doi.org/10.1074/mcp.O113.027284

Yao H-Y and Xue H-W. Phosphatidic acid plays key roles regulating plant development and stress
responses. Journal of Integrative Plant Biology. 2018:**60**(9):851–863.
https://doi.org/10.1111/jipb.12655

Zhu Z, Krall L, Li Z, Xi L, Luo H, Li S, He M, Yang X, Zan H, Gilbert M, et al. Transceptor NRT1.1 and
receptor-kinase QSK1 complex controls PM H+-ATPase activity under low nitrate. Current
Biology. 2024:**34**(7):1479-1491.e6. https://doi.org/10.1016/j.cub.2024.02.066

1036

1037 **Figure legends**

Figure 1. CRK2 and QSK1 regulate plant growth and osmotic stress response.

(A) Rosette of 2-week-old Arabidopsis plants grown in long-day conditions. Mutation of QSK1 reverts small growth phenotype of *crk2* in the double mutant *crk2 qsk1*. (B) rosette area of 12-days-old plants. (C) Primary root (PR) length of 12-days-old plants. (D) Lateral root (LR) number per cm of PR (LR density) was counted under a stereozoom from the same plants in panel C. The boxes represent the interquartile range (25th–75th percentile), the horizontal line indicates the median, the whiskers extend to 1.5× the IQR, and dots represent individual data points. Panel B, n=59, one-way ANOVA, Tukey's test; Panel C, n=34, Kuskal-Wallis test followed by Dunn's multiple comparison; Panel D, n=34, one-way ANOVA, Tukey's test. Alphabets represent statistically significant differences. Scale bar 1 cm. See methods and Table S1.

Figure 2. QSK1 and CRK2 play antagonistic roles in osmotic stress-induced callose deposition.

(A) Visualization of aniline blue-stained callose deposits in leaf epidermal cells of Arabidopsis with or without osmotic stress. Roots of 2-week-old plants were hydroponically subjected to 50 mM mannitol for 7 hours before imaging in the treated groups. Images acquired with a laser-scanning confocal microscope under the DAPI filter with emission wavelength 461 nm and laser wavelength 405 nm. Detection wavelength 430-530 nm; laser transmissivity 5.0 %; objective lens 60.0 X oil immersion, scale bar 10 µm. (B) Mean intensity of callose deposits. (C) Number of callose deposits. (D) Relative callose deposition. Boxplots as described in Figure 1, n ≥ 22 for all the plots. Two-way ANOVA, Tukey's multiple comparison test. See methods and Table S2.

Figure 3. Functional CRK2 is required for osmotic stress-induced enrichment of QSK1 at plasmodesmata.

(A) Relocalization QSK1-mCherry in Arabidopsis leaf epidermal cells. Detached leaves from plants expressing QSK1-mCherry in WT or *crk2* mutant backgrounds were treated with 0.4 M mannitol. The abaxial surface was visualized and imaged with a laser-scanning confocal microscope after 30 min of treatment. QSK1 failed to enrich in the *crk2* mutant background. (B) Boxplot showing the extent of PD enrichment of QSK1 which is signal intensity ratio between PD and PM ROI. (C) The disruption of QSK1 enrichment in the *crk2* mutant background was rescued by complementing QSK1-mCherry/*crk2* plants with CRK2-mVenus (QSK1-mCherry/*crk2* crossed with CRK2-mVenus/*crk2*). (D) The kinase inactive CRK2^{D450N} failed to complement this function in the cross of QSK1-mCherry/*crk2* and CRK2 ^{D450N}-mVenus/*crk2* plants. (E) The mutation in *QSK1* did not impact the relocalization of CRK2 in the CRK2-mVenus/*qsk1* plants. Images are acquired after 60 min of 0.4 M mannitol treatment. (F) Boxplot showing enrichment index related to panel E. The images were captured with 60.0 X oil immersion lens. Laser and emission wavelengths were 561 nm and 610 nm, respectively for mCherry; 514 nm and 527 nm for mVenus. Detection wavelengths were 570-620 nm and 530-630 nm for mCherry and mVenus, respectively. Laser transmissivity in the shown images is 7.0%-15% which was kept constant between the comparisons. Scanning of signals were done in a sequential manner to avoid signal cross-contamination between the channels. Scale bar 10 µm. At least 3 independent transgenic lines were tested for each genotype. Boxplots as described in Figure 1. n ≥ 15, two-way ANOVA, Tukey's test. See methods and Table S3.

Figure 4. QSK1 and CRK2 relocalize and enrich at plasmodesmata in a sequential manner during osmotic stress.

Arabidopsis leaf epidermal cells showing relocalization of QSK1 and CRK2. The images shown are captured within a window of 0-60 min. QSK1 enrichment takes place first and reaches maximum enrichment at approximately 30 min of treatment and subsequently starts to decline to basal levels. Objective lens 40.0 X oil immersion. Treatment and laser settings as described for Figure 3. Scale bar 10 µm.

Figure 5. QSK1 and CRK2 are present at the bulk plasma membrane in a preformed complex.

(A) FRET-FLIM analysis of CRK2-mVenus (donor-only), CRK2-mVenus+QSK1-mCherry (donor-acceptor) and CRK2^{D450N}-mVenus+QSK1-mCherry (donor-acceptor negative control) plants with or without mannitol. A lower mVenus lifetime in the donor-acceptor pair CRK2-mVenus+QSK1-mCherry indicated the presence of a preformed complex of CRK2 and QSK1. An increase in the mVenus lifetime in this pair after mannitol treatment suggested dissociation of the complex. Boxplot as described in Figure 1. $n \geq 9$. Two-way ANOVA, Tukey's test. See methods and Table S4. (B, C) AlphaFold2 prediction of interaction between CRK2 and QSK1 extracellular and transmembrane domains (ExD+TMD) (B) and full-length proteins (C). Contacts under 6 Å between the two proteins are shown. Lines are colored according to the PAE values, with most contacts showing high confidence.

Figure 6. CRK2 phosphorylates QSK1

In vitro kinase reaction between the recombinantly expressed cytosolic regions of CRK2 and QSK1. (A) CRK2-targeted phosphosites in the QSK1 cytosolic region as identified by LC-ESI-MS/MS analysis. The highlighted residues in pink are consolidated from 3 experiments-high confidence only. The shown amino acid sequence is the cytosolic region of QSK1 and the grey highlighted part corresponds to the pseudokinase domain. The arrows represent the two serine residues at position 621 and 626 which were selected for further studies. See Supplementary Table S6. (B) *In vitro* phosphorylation of QSK1cyto by CRK2cyto. The QSK1cyto-MBP and QSK1$^{S621A-S626A}$ cyto-MBP were digested with Factor Xa protease for 24 h to separate the MBP tag for size resolution before carrying out the kinase reaction. The QSK1 was phosphorylated by CRK2 but not with the kinase inactive CRK2^{D450N}. No difference was observed in the phosphorylation of QSK1$^{S621A-S626A}$. The right panel shows the Coomassie Blue-stained gel which was destained and developed for the detection of labeled ATP (left panel). (C) *In vivo* phosphorylation of QSK1. The shown phosphopeptide was detected in the WT and *crk2* seedlings. NaCl treatment and mutation in *CRK2* reduced the intensity of phosphopeptide. The detected phosphopeptide was predicted to be phosphorylated at two serine residues. Boxplot description as Figure 1. n=3-4. Two-way ANOVA, Tukey's test. See methods and Table S7.

Figure 7. Phosphorylation of QSK1 at Ser-621 and Ser-626 inhibits its recruitment to PD.

Relocalization of phospho-mutant versions of mCherry-tagged QSK1 in the leaf epidermal cells of Arabidopsis WT or *crk2* mutant backgrounds. (A, B) The phospho-mimetic QSK1 (QSK1$^{(S621D-S626D)}$) showed reduced enrichment in both WT and *crk2* mutant backgrounds. (C, D) The phospho-negative QSK1 (QSK1$^{(S621A-S626A)}$) showed much higher enrichment in the WT background upon mannitol treatment. A weak enrichment of phospho-negative QSK1 was also

observed in control conditions. The phospho-negative QSK1 also enriched weakly in the *crk2* mutant background after mannitol treatment. Objective lens 60.0 X oil immersion. Laser transmissivity 7.0%-15%; constant between the comparisons. Experimental conditions and other laser settings for mCherry as Figure 3. Scale bar 10 µm. Boxplots as described in Figure 1. Two-way ANOVA, Tukey's test. n ≥ 12. See methods and Table S3.

Figure 8. Phosphorylation of QSK1 at Ser-621 and Ser-626 suppresses stress-induced callose deposition and regulates plant stress response.

(A) Visualization of aniline blue-stained callose deposits in leaf epidermal cells of Arabidopsis with or without osmotic stress. Treatment and imaging as described for Figure 2. Scale bar 10 µm. (B) Mean intensity of callose deposits. (C) Number of callose deposits. (D) Relative callose deposition. Boxplots as described in Figure 1, n = 30 for plots B, C and D. Two-way ANOVA, Tukey's test. (E) Primary root (PR) length, n =23-24. Kruskal-Wallis test, Dunn's multiple comparison test. (F) PR elongation. 5-days-old seedlings were transferred to media containing NaCl and were imaged after another 7 days. The PR elongation is calculated respective to the control PR length. n=22-23, two-way ANOVA, Tukey's test. (G) LR density (no. of lateral root per cm of PR) was counted under a stereozoom from the same plants in panel F. n=22, Kruskal-Wallis test, Dunn's multiple comparison test. (H) Percentage of viable seed germination on media containing NaCl. The germination efficiency was normalized according to the respective control plates. n=8, two-way ANOVA, Tukey's test. Boxplots description as Figure 1. QSK1, QSK1^{S-A} and QSK1^{S-D} are the plants expressing WT, QSK1$^{(S621A-S626A)}$ and QSK1$^{(S621D-S626D)}$ variant of QSK1, respectively, in the *qsk1* mutant background. Data and statistical details for panels A-D in Table S8 and for panels E-I in S9.

Figure 9. Proposed model- CRK2 modulates plasmodesmal function of QSK1.

Under normal growth conditions, CRK2 phosphorylates QSK1 and the two RLKs form a complex at the bulk plasma membrane (PM) inhibiting unnecessary recruitment of QSK1 to PD (A). Exposure to salt or mannitol promotes an unknown protein phosphatase activity at QSK1 releasing it from the CRK2-QSK1 complex and consequently QSK1 relocalization to PD. CRK2 also phosphorylates another unknown protein(s), a necessary event for QSK1 enrichment at PD. QSK1 positively regulates callose deposition at PD (B). Subsequently, QSK1 is removed from PD and returns to basal level. CRK2 enrichment at PD follows (C).

Supplementary Figure legends

Figure S1 (related to Figure 1). CRK2 and QSK1 regulate plant growth and osmotic stress response.

(A) Comparison of PR length under normal growth conditions. (B) Root elongation assay on NaCl. 5 DAS (days-after-stratification) seedlings were transplanted on media containing NaCl and imaged after 7 days. (C) Percentage of PR elongation of the plants shown in panel B. Root elongation is calculated respective of the control PR length. An equal retardation of PR elongation was observed on media containing salt for all the genotypes. (D) Percentage of viable seed germination on media containing NaCl. The hypersensitive response of *crk2* mutant on salt-containing media was reverted in the *crk2 qsk1* double mutants. The germination efficiency was normalized according to the respective control plates. The boxes represent the interquartile range (25th–75th percentile), the horizontal line indicates the median, the whiskers extend to 1.5× the

1164 IQR, and dots represent individual data points. Panel C, n=34, Kuskal-Wallis test followed by
1165 Dunn's multiple comparison; Panel D, n=3, one-way ANOVA, Tukey's test. See methods and
1166 Table S1.

Figure S2 (related to Figure 3). Enrichment of QSK1 at plasmodesmata requires CRK2.

(A) Relocalization QSK1-mCherry in Arabidopsis leaf epidermal cells. Detached leaves from
plants expressing QSK1-mCherry in WT and *crk2* backgrounds (QSK1-mCherry/WT, QSK1-
mCherry/*crk2*) were treated with 150 mM NaCl and imaged at 30- and 60-minute post-treatment.
Control samples were without salt treatment. In the QSK1-mCherry/WT plants, the punctate
pattern of mCherry signal was prominent at 30 min but faded after 60 min of treatment. In the
QSK1-mCherry/*crk2* plants, QSK1-mCherry did not enrich upon salt treatment. (B) Co-
localization of aniline blue-stained callose deposits with enriched QSK1-mCherry signal
showing that QSK1 enriched at PD upon osmotic stress treatment. Leaves were co-treated with
150 mM NaCl and 0.2 % aniline blue and abaxial surface was imaged with laser-scanning
confocal microscope after 30 min of treatment. Objective lens 40.0 X oil immersion, emission
wavelength for DAPI filter 461 nm and laser wavelength 405 nm. Detection wavelength 430-530
nm, laser transmittivity 2.0%, Scale bar 5 µm. Other laser settings for mCherry as Figure 3.

Figure S3 (related to Figure 4). QSK1 and CRK2 relocalize and enrich at plasmodesmata
in a sequential manner during osmotic stress.

Arabidopsis leaf epidermal cells showing relocalization of QSK1 and CRK2 in the QSK1-
mCherry/CRK2-mVenus plants. The images shown are captured within a window of 0-60 min of
0.4 M mannitol treatment. QSK1 enrichment is followed by CRK2 enrichment. Objective lens
40.0 X oil immersion. Laser wavelengths were 561 nm and 514 nm for mCherry and mVenus,
respectively. Emission wavelengths were 610 nm and 527 nm for mCherry and mVenus,
respectively. Detection wavelengths were 570-620 nm and 530-630 nm for mCherry and
mVenus, respectively. Scanning of signals were done in a sequential manner to avoid signal
cross-contamination between the channels. Scale bar 10 µm.

Figure S4 (related to Figure 5). Structural modeling prediction of the specific domains of
CRK2 and QSK1.

1192 AlphaFold2 prediction of interaction between CRK2 and QSK1 segments- extracellular domains
1193 (ExD), transmembrane domains (TMD), TMD + Kinase domains, and Kinase domains.
1194 Predicted Aligned Error (PAE) plot shows model's confidence. Lower PAE values indicate
1195 higher confidence of interaction. Lines are colored according to the PAE values. The modeled
1196 boundaries and predicated scores in Table S5.

Figure S5 (related to Figure 7). Phosphorylation of QSK1 at Ser-621 and Ser-626 inhibits
its recruitment to PD.

Relocalization of QSK1-mCherry and QSK1 phospho-mutants in different genetic backgrounds
of Arabidopsis leaf epidermal cells. (A, B) Enrichment of QSK1 at PD is independent of
Sucrose-Induced Receptor Kinase 1 (SIRK1). Arabidopsis plants expressing QSK1-mCherry in
sirk1 mutant background show normal recruitment of QSK1 at PD upon mannitol treatment. A
basal presence of QSK1 at PD was visible in control samples as well. (C, D) QSK1$^{S621E-S626E}$
showed a weak enrichment at PD in WT background upon mannitol treatment. (E, F) Similar to
WT background in Fig. 7C, D, the phospho-negative QSK1 (QSK1$^{(S621A-S626A)}$) showed much

higher enrichment in the *qsk1* background upon mannitol treatment. For plots B and F, statistical significance was assessed using unpaired test with Welch's correction; for plots D and E, statistical significance was assessed using two-way ANOVA followed by Tukey's multiple comparisons test. In panels A, C; laser transmissivity 7.0%, objective lens 60.0 X oil immersion; D; laser transmissivity 2.0%, objective lens 40.0 X oil immersion. Scale bar 10 um. 3 independent transgenic lines were tested. See methods and Table S3.

Supplementary Tables

Table S1. Phenotypic analysis and stress response-raw data (related to Fig. 1B, 1C, 1D and Fig. S1C, S1D)

Table S2. Callose quantification-raw data (related to Fig. 2B, 2C, 2D)

Table S3. Enrichment index-raw data (related to Fig. 3B, 3F, 7B, 7D; Fig. S5B, S5D, S5F)

Table S4. FRET-FLIM-raw data (related to Fig. 5A)

Table S5: Amino acid boundaries of full-length, ExD, ExD+TMD, TMD, TMD+Kinase domain, and Kinase domain CRK2 and QSK1 used for ColabFold predictions

Table S6. Phosphopeptides identified in LC-MS/MS after *in vitro* kinase reaction between CRK2 and QSK1 cytosolic regions

Table S7. Intensity of QSK1 phosphopeptide in phosphoproteomics (related to Fig. 6C)

Table S8. Callose quantification-raw data (related to Fig. 8B, 8C, 8D)

Table S9. Phenotypic analysis and stress response-raw data (related to Fig. 8E, 8F, 8G, 8H, 8I)

Table S10-Primers used in the study

A

donor lifetime (ns)

CRK2-mVenus

CRK2-mVenus +
QSK1-mCherry

CRK2^{D450N}-mVenus +
QSK1-mCherry

Control
Mannitol

B CRK2 QSK1

ExD+TMD

90°

CRK2

QSK1

C

Full-length

ExD

TMD

90°

Kinase

Predicted Aligned Error (Å)

0   15   30

CRK2

QSK1

A

```
CRKRKKEENVPSRNVEAPVAAATSSAAIPKETVVVVPPAK
ATGSESGAVNKDLTFFVKSFGEFDLDGLLKASAEVLGKGT
VGSSYKASFEHGLVVAVKRLRDVVPEKEFRERLHVLGSM
SHANLVTLIAYYFSRDEKLLVFEYMSKGSLSAILHGNKGN
GRTPLNWETRAGIALGAARAISYLHSRDGTTSHGNIKSSN
ILLSDSYEAKVSDYGLAPIISSTSAPNRIDGYRAPEITDA
RKISQKADVYSFGVLILELLTGKSPTHQQLNEEGVDLPRW
VQSVTEQQTPSDVLDPELTRYQPEGNENIIRLLKIGMSCT
AQFPDSRPSMAEVTRLIEEVSHSSGSPNPVSD
```

B

MBP	-	-	-	-	+	+
QSK1^{S621A-S626A}cyto	-	-	+	+	+	+
QSK1cyto	+	-	-	-	-	-
CRK2^{D450N}cyto	-	+	-	+	-	+
CRK2cyto	+	-	+	-	+	-

autoradiogram — CRK2cyto, QSK1cyto

CBB — KDa 180 130 100 70 55 40 35 25

C

In vivo phosphorylated QSK1 peptide

LIEEVSHSSGSPNPVSD

Intensity of phosphopeptide (a.u.)

Control / NaCl

WT *crk2*

Normal growth conditions

A

PD

PM

Cytoplasm

CRK2

QSK1

P

NaCl/Mannitol

B

PD

CRK2

QSK1

P

QSK1 relocalization to PD

QSK1

PP activity

P

C

PD

CRK2

CRK2 relocalization to PD

CRK2

QSK1

QSK1 removal
from PD

P

Time after stress treatment

 Unknown phosphorylation target(s) of CRK2　　

 Callose　　🔄 P Phosphorylation

PP　Protein Phosphatase
PD　Plasmodesmata