CRK2 controls the spatiotemporal distribution of QSK1 at plasma membrane during osmotic stress

Running title: CRK2 controls plasmodesmal RLK recruitment

Sunita Jindal[1], Adam Zeiner[1,2], Alexey Bondar[1,2], Michaela Neubergerová[3,4], Sara Christina Stolze[5], Anne Harzen[5], Francisco J. Colina[1,8], Sara Liekens[1,6], Mirva Pääkkönen[7], Johannes Merilahti[7], Ivan Kulich[1,2], Roman Pleskot[3], Hirofumi Nakagami[5], Michael Wrzaczek[1,2,9]

[1]Institute of Plant Molecular Biology, Biology Centre, Czech Academy of Sciences, 37005 České Budějovice, Czechia

[2]Faculty of Science, University of South Bohemia, 37005 České Budějovice, Czechia

[3]Institute of Experimental Botany (IEB), Czech Academy of Sciences, 16502 Prague, Czechia

[4]Faculty of Science, Charles University, 12800 Prague, Czechia

[5]Max Planck Institute for Plant Breeding Research, Cologne, Germany

[6]Thomas More University of Applied Sciences, Campus Geel, 2440 Geel, Belgium

[7]Turku Bioscience Centre, University of Turku and Åbo Akademi University, 20520 Turku, Finland

[8]Present address: Estación Experimental del Zaidín (EEZ-CSIC), 18008, Granada, Spain

[9]Lead contact: michael.wrzaczek@umbr.cas.cz

Abstract

Precise control of intercellular communication is essential for normal growth and stress responses in all multicellular organisms. In Arabidopsis, two membrane-localized receptor like kinases (RLKs), the Cysteine-rich RLK CRK2 and the Leucine-rich repeat (LRR) RLK QSK1 relocalize from the general plasma membrane (PM) to plasmodesmata (PD) in response to osmotic stress. Both these RLKs regulate callose deposition thereby modulating PD permeability. However, unchecked callose deposition can block the PD and disrupt proper intercellular communication. Here, we show that under normal growth conditions, CRK2 phosphorylates and sequesters QSK1 at the general PM, preventing unnecessary callose deposition at PD. We show that osmotic stress-induced enrichment of QSK1 at PD requires functional CRK2 and establish that phosphorylation of QSK1 in its C-terminal region is inhibitory in this process. We propose that osmotic stress triggers dephosphorylation and release of QSK1 from the CRK2-QSK1 complex, enabling its relocalization from general PM to PD, where it promotes stress-induced callose deposition. Subsequently, CRK2 relocalizes to PD where it negatively influences callose deposition. Our work reveals a tightly coordinated distribution of QSK1 and CRK2 at PM, establishing a dynamic gating mechanism that balances growth and stress responsiveness.

Introduction

In natural habitats, plant water uptake is compromised by salinity, drought and extreme

temperatures leading to osmotic stress which limits normal growth and development. On the

physiological level, osmotic adjustments are made by closing stomata, growth inhibition,

hydraulic modifications and changes in root system architecture. On the molecular level, rapid

signaling events are triggered at the plasma membrane (PM) and include activation of

phosphorylation networks, activation of NADPH oxidases (respiratory burst oxidase homologs,

RBOHs) and ROS burst[1], Ca^{2+} influx[2–4] and changes in lipid signaling, for example,

accumulation of phosphatidic acid (PA)[5–8]. Another key regulatory element of osmotic stress

signaling is the deposition of callose[9–11]. Callose, a β-1,3-glucan polysaccharide is deposited

primarily in the neck region of plasmodesmata (PD) and controls symplastic connectivity

through gating the PD aperture[12,13]. Symplastic connection regulates cell division, expansion and

organ formation[14,15] which could provide growth adaptation especially during extreme stress

conditions. For example, mutants accumulating higher levels of callose have reduced primary

root (PR) length[9,16] and altered lateral root (LR) patterning[12,17]. Therefore, adaptations in the root

phenotype during osmotic stress can at least partly result from the changes in symplastic

connection.

The amount of callose at PD changes dynamically throughout plant growth, development and

during stress conditions. The main determinants of callose amounts are callose synthases

(CALSs) and β-1,3-glucanases (PdBGs), responsible for callose synthesis and degradation,

respectively[18]. Several PD-localized proteins (PDLPs), PD-localized chaperons such as PD-

callose binding proteins (PDCBs) participate in maintaining callose homeostasis by recruiting or

stabilizing CALSs or PdBGs[18–20]. In recent years, receptor-like kinases (RLKs) have emerged as

regulating callose accumulation during abiotic stress[9,21,22]. We previously showed that CRK2, a

cysteine-rich RLK (CRK) is involved in the regulation of callose deposition and positively

regulates seed germination during salinity stress[9]. The mutant *crk2* exhibits an overall small

phenotype and deposits higher callose. CRK2 relocalizes to PD in response to salt and mannitol

treatments and also assembles in nanodomain-like clusters upon flg22 and H_2O_2 exposure[9].

Concurrently,[21] showed that Qian Shou kinase 1 (QSK1), a leucine-rich repeat RLK (LRR-RLK)

is recruited to PD upon osmotic stress and the *qsk1* mutant accumulates lower callose levels and

shows altered root response to mannitol.

PM components undergo dynamic re-organization upon perception of various stimuli and cluster

into nanodomains[23] to facilitate the assembly of signaling complexes and providing signal

specificity[24,25]. Membrane nanodomains are hallmarked by specific lipid and protein

compositions. Scaffold proteins such as Remorins and Stomatin/Prohibitin/Flotillin/HflK/C

(SPFH) domain-containing protein bridge lipids and membrane receptors, whereas RLKs are

dynamically positioned in nanodomains[26,27]. Cytoskeletal elements and cell wall components

further regulate nanodomain mobility and spatial distribution[27]. The re-organization of

membrane proteins is often coupled to a phosphorylation switch. For example, in resting cells,

BRI1 (Brassinosteroid Insensitive 1)-Associated receptor Kinase 1 (BAK1), an LRR-RLK, is

evenly distributed across the PM, with its catalytic pocket shielded by BAK1-Interacting

Receptor-like kinases (BIRs)[28]. Ligand perception activates the BRI1-BAK1 or FLAGELLIN-

SENSITIVE 2 (FLS2)-BAK1 signaling axes, triggering growth or immune signaling,

respectively. This involves a phosphorylation-dependent shift of BAK1 from resting to active

state and the formation of receptor nanodomains[29]. Likewise, the distribution of Remorin[30,31]

and QSK1[21] across the PM is dependent on their phosphorylation status.

Similar to BAK1 as a shared co-receptor in growth and immune signaling, QSK1 forms several

protein complexes. For instance, the SIRK1-QSK1 module regulates aquaporins where QSK1

acts as a co-receptor of the RLK Sucrose-Induced Receptor Kinase 1 (SIRK1)[32,33]. QSK1 also

associates with pattern recognition receptor (PRR)-RBOHD complexes involving FLS2 and

Elongation Factor Tu Receptor (EFR) RLKs[34]. QSK1 also regulates the vacuolar Two-pore

potassium channel 1 (TPK1) during stomatal closure by phosphorylating TPK1 with

involvement of an unidentified receptor kinase[35]. Similarly, QSK1 associates with the transceptor

nitrate transporter NRT1.1 in regulating PM Arabidopsis H^+-ATPase 2 (AHA2)[36].

While the re-organization of individual PM components is extensively studied, how different

RLKs coordinate their spatiotemporal behavior and how the shared components are exchanged

between the complexes is much less understood. CRK2 and QSK1 are both primarily present at

the general PM under normal growth conditions but both relocalize to PD upon osmotic stress.

We previously identified that QSK1 associates with CRK2[9]. However, the functional link

between these two RLKs in osmotic stress response was not explored. Here, we elucidate their

connection in fine-tuning callose deposition and plant responses to osmotic stress. We show that

CRK2 controls the plasmodesmal relocalization of QSK1 and both RLKs antagonistically

regulate callose deposition during osmotic stress responses. Our work highlights that the

dynamics of the re-organization of QSK1 and CRK2 in response to osmotic stress is

spatiotemporally coordinated and involves a phosphorylation switch.

Results

Role of CRK2 and QSK1 in plant growth and osmotic stress response

The growth phenotype and osmotic stress responses of the *crk2* and *qsk1* have been previously

described[9,21,37]. We generated the *crk2 qsk1* double mutant by crossing *crk2* and *qsk1* single

mutants and characterized the growth phenotype under normal growth conditions salinity stress.

As reported previously[9], *crk2* displayed overall smaller rosette, shorter primary root (PR) and

lower lateral root (LR) density (Fig. 1A-C, Table S1). The *qsk1* rosette area was comparable to

Col-0 wild type (WT), while in *crk2 qsk1*, the small rosette phenotype of *crk2* was rescued with

a slight increase in comparison to the WT (Fig. 1A, Fig. S1A). While *crk2* exhibited the shortest

PR, *qsk1* and *crk2 qsk1* also displayed shorter PR compared to the WT (Fig. 1B, Fig. S1B).

Similarly, LR density was lower in all the tested mutants compared to WT (Fig. 1C). Overall, the

observable phenotypic traits of *crk2 qsk1* were more similar to either WT or *qsk1* rather than

crk2. The partial reversion of *crk2* phenotypes in *crk2 qsk1* mutant was also observed at the level

of stress responses. We tested WT, *crk2*, *qsk1* and *crk2 qsk1* seeds for salt-sensitivity during seed

germination and observed that, *crk2* seeds showed hypersensitivity to salt as reported

previously[9], whereas *qsk1* and *crk2 qsk1* seed germination was comparable to WT (Fig. 1D).

Taken together, these results suggest, that CRK2 and QSK1 functionally interact to regulate plant

growth and stress responses, and moderation of *crk2* phenotype by introduction of additional

qsk1 mutation points towards antagonistic roles for those two RLKs.

CRK2 and QSK1 fine-tune callose deposition in response to osmotic stress

The mutants *crk2* and *qsk1* have been previously reported to accumulate altered levels of

callose[9,21]. We quantified callose accumulation in WT, *crk2*, *qsk1* and *crk2 qsk1* seedlings by

aniline blue staining followed by microscopy-based detection of the aniline blue fluorescence

signal (Fig. 1E-F, Fig. SC-D, Table S2)[9,38,39]. Since osmotic stress is a common component of

various abiotic stresses, including salinity, drought, waterlogging, and extreme temperatures, we

used mannitol to specifically impose osmotic stress without introducing additional ionic effects.

Arabidopsis plants were treated with mannitol for 7 hours before aniline blue staining and

imaging. We quantified total callose deposition (Fig. 1F) by taking into consideration mean

fluorescence intensity of the spots and the number of deposits (Fig. S1C-D, Table S2). In the

crk2 mutant, total callose deposition was higher than the WT in control conditions which further

increased upon mannitol treatment (Fig. 1F). This suggested a negative influence of CRK2 on

callose accumulation during steady-state as well as osmotic stress conditions. In the *qsk1* mutant

under control conditions, total callose was comparable to the WT (Fig. 1F). Surprisingly,

mannitol-treated *qsk1* showed a lower total callose signal compared to normal growth conditions

suggesting that QSK1 might not play a role in basal callose deposition but positively regulates

osmotic stress-induced callose deposition (Fig. 1F). Interestingly, the reciprocal impact of CRK2

and QSK1 seemed to negate each other in the *crk2 qsk1* mutant as this mutant accumulated

similar callose levels as WT under control and mannitol treatment (Fig. 1F, Fig. SC-D). Together,

these findings suggest opposing roles of CRK2 and QSK1 in regulating osmotic stress-induced

callose deposition.

Functional CRK2 is essential for osmotic stress-induced enrichment of QSK1 at PD

CRK2 and QSK1 were shown to relocalize to PD in response to osmotic stress[9,21] and QSK1 was

identified as an *in planta* interactor of CRK2[9]. We hypothesized that their interaction might be

necessary for the enrichment of both proteins at PD. Therefore, we tested whether the absence of

functional CRK2 or QSK1 would impact the recruitment of the respective other RLK to PD. We

generated transgenic plants expressing QSK1 fused to mCherry under native *QSK1* promoter in

crk2 and WT backgrounds and imaged leaf epidermal cells. Interestingly, while QSK1 enriched

at PD following mannitol or NaCl treatment in WT background, this enrichment was abolished

in *crk2* background (Fig. 2A-B, Fig. S2A, Table S3). In QSK1-mCherry/WT, the enrichment

index, which is the fluorescence intensity ratio between the enriched spot (PD) and an adjacent

non-enriched spot on the bulk PM, increased from 0.940 to 1.693 upon mannitol treatment. It did

not increase for QSK1-mCherry/*crk2* and remained approximately 1.0 for both control and

mannitol treatment (Fig. 2B). Since callose is particularly present at PD, it is used to mark PD by

aniline blue staining[9,38,39]. To test if the enriched QSK1-mCherry spots following osmotic stress

treatment corresponded with PD, we simultaneously applied osmotic stress and aniline blue

staining. The co-localization of aniline blue-stained callose with enriched mCherry signal

supported that the punctate pattern of QSK1-mCherry correspond to PD (Fig. S2B). Consistently,

the genetic cross between QSK1-mCherry/*crk2* and CRK2-mVenus/*crk2* rescued QSK1

enrichment at PD in response to mannitol treatment (Fig. 2C). At this timepoint after mannitol

treatment, we did not observe PD enrichment for CRK2 possibly due to a lag in the PD

relocalization of CRK2 (Fig. 3B). The mutation D450N in the catalytic core of CRK2 kinase

domain (CRK2^{D450N}) abolishes the phosphorylation activity of CRK2, and CRK2^{D450N} is not able

to relocalize to PD[9]. To determine whether the kinase activity of CRK2 was also essential for the

recruitment of QSK1 at PD, we crossed QSK1-mCherry/*crk2* with CRK2^{D450N}-mVenus/*crk2*

plants. In the resulting plants, QSK1 failed to relocalize to PD in response to mannitol treatment

(Fig. 2D). To investigate whether CRK2 requires QSK1 for its own PD localization, we stably

expressed CRK2-mVenus under native *CRK2* promoter in WT and *qsk1* backgrounds.

Interestingly, we found that CRK2 relocalized to PD upon mannitol treatment without functional

QSK1 (Fig. 2E, F). These results established that enzymatically active CRK2 is essential for the

relocalization of QSK1 to PD in response to osmotic stress. However, the relocalization of CRK2

is independent of QSK1 pointing to an upstream role of CRK2 in PD-targeting of itself and

QSK1.

QSK1 and CRK2 enrich at PD in a sequential manner following osmotic stress

RLKs are dynamically re-organized at the PM in response to various internal and external

cues[24,25]. Despite the dependence of QSK1 on CRK2 in its relocalization to PD, we observed a

temporal disconnection in their relocalization. QSK1 accumulation at PD peaked within 30-45

min of salt or mannitol treatment and subsequently the punctate pattern faded with almost no

observable enrichment 60 min post-treatment (Fig. 2A, Fig. S2A, Fig. 3A). In contrast to QSK1,

the punctate pattern corresponding to CRK2-mVenus peaked at PD only after 60 min of

treatment (Fig. 2E). Using QSK1-mCherry and CRK2-mVenus co-expressing lines, we carried

out a time-course analysis and showed that, enrichment of QSK1 and CRK2 at PD was

sequential (Fig. 3A, Fig. S3A). Following peak enrichment of QSK1-mCherry at PD, CRK2-

mVenus began to relocalize to PD. As CRK2-mVenus reached its saturation at PD, we did not

observe localization of QSK1-mCherry at PD anymore (Fig. 3A). Overall, we did not observe

simultaneous peak enrichment for both QSK1 and CRK2 at PD. This finding suggested that the

two RLKs might co-ordinate to precisely fine-tune callose deposition to control the PD aperture.

CRK2 and QSK1 exist in a pre-formed complex at the bulk PM that dissociates upon

osmotic stress

We previously detected QSK1 as an *in vivo* interactor of CRK2[9] implying that the two RLKs co-

localize at the general PM in close proximity by interacting either directly or in a multi-protein

complex. However, the sequential recruitment of QSK1 and CRK2 at PD suggested a

dissociation of the two upon stress perception. To test this, we performed Förster resonance

energy transfer by fluorescence lifetime imaging (FRET-FLIM) of Arabidopsis plants stably co-

expressing QSK1-mCherry and CRK2-mVenus proteins (QSK1-mCherry/*crk2* x CRK2-

mVenus/*crk2*) with and without mannitol treatment. As a negative control, we used equivalent

plants with kinase inactive CRK2 (QSK1-mCherry/*crk2* x CRK2^{D450N}-mVenus/*crk2*), since

CRK2^{D450N} did not relocalize following stress application[9]. In plants expressing the donor-

acceptor pair (CRK2-mVenus + QSK1-mCherry), donor life-time was lower in comparison to the

donor-only and negative control plants under control conditions (Fig. 3B, Table S4). This

suggested the existence of a preformed complex of CRK2 and QSK1 in normal conditions which

would reduce the fluorescence lifetime of the donor mVenus. Interestingly, upon mannitol

treatment, the donor lifetime in QSK1-mCherry + CRK2-mVenus plants increased suggesting

dissociation of the complex upon mannitol treatment. This finding was in-line with the sequential

enrichment of QSK1 and CRK2 at PD (Fig. 3A, Fig. S3A). In contrast, in the QSK1-mCherry +

CRK2 ^{D450N}-mVenus plants, the donor lifetime was higher under both control and mannitol-

treated conditions suggesting that the kinase inactive CRK2 might have weaker affinity for the

protein complex formation (Fig. 3B). Further, we performed structural modeling to predict the

interaction between CRK2 and QSK1 using Colabfold (version 1.5.2) implementation of

AlphaFold2[40,41]. We modeled the full-length proteins, and the combinations of their individual

structural segments. The modeled combinations were i) full-length proteins, ii) ExD

(Extracellular domains), iii) ExD+TMD (transmembrane domains), iv) TMD, v) TMD+Kinase

domain, and vi) Kinase domain for both CRK2 and QSK1. Based on the predicted aligned error

(PAE) values (Fig. 3C-D, Fig. S3B) and interface predicted template modelling (iPTM) scores

(Table S5), ExD+TMD regions showed high confidence for interaction (Fig. 3C). The modeling

of the full-length proteins suggested that the two RLKs engage specific segments for interaction

(Fig. 3D) which mainly resulted from the ExD and TMD regions (Fig. S3B). Collectively, these

results confirmed that CRK2 and QSK1 are present in close proximity at the general PM and

form a complex *in planta,* and this complex dissociates upon osmotic stress perception.

CRK2 phosphorylates QSK1 on its C-terminus, and salt stress reduces this
phosphorylation

The role of QSK1 in PD regulation was reported to be regulated by its phosphorylation[21]

specifically of two serine residues in the C-terminus of the cytosolic region of QSK1 (Ser-621

and Ser-626). However, the responsible kinase was not investigated. Since we previously

identified QSK1 as an *in vivo* interactor of CRK2[9], we asked if CRK2 could phosphorylate

QSK1, thereby controlling its plasmodesmal relocalization. We recombinantly expressed the

cytosolic parts of CRK2 (6His-GST-CRK2cyto and 6His-GST-CRK2^{D450N}cyto)[9] and QSK1

(6His-MBP-QSK1cyto) in *E. coli*. To identify CRK2-targeted phosphorylation sites in QSK1, we

carried out LC-ESI-MS/MS analysis of in-gel trypsin-digested peptides after the kinase reaction

of CRK2cyto and QSK1cyto. The identified *in vitro* phosphorylation sites are shown in Table S6

and Fig. 4A. Interestingly, Ser-626 was among the identified phosphosites (Fig. 4A), suggesting

that CRK2 was able to directly interact with and phosphorylate the QSK1 peptide containing

Ser-626. Due to the importance of Ser-621 and Ser-626 in osmotic stress response[21,32,33,42], we

generated a variant of QSK1 with phospho-negative substitutions at positions 621 and 626

(QSK1$^{S621A\text{-}S626A}$) as a phosphorylation target of CRK2. Since 6His-GST-CRK2cyto and 6His-

MBP-QSK1cyto have similar molecular weights, we digested 6His-MBP-QSK1cyto and 6His-

MBP-QSK1$^{S621A\text{-}S626A}$cyto with Factor Xa protease before the kinase reaction to cleave the MBP

tag off from the MBP-QSK1cyto fusion protein resulting in a clear resolution of CRK2 and

QSK1 cytosolic fragments. The QSK1cyto was phosphorylated by 6His-GST-CRK2cyto but not

by the kinase inactive 6His-GST-CRK2^{D450N}cyto (Fig. 4B) in *in vitro* kinase assay. Further, we

detected an overall comparable phosphorylation signal of the WT QSK1 and QSK1$^{S621A\text{-}S626A}$,

probably due to multiple other phosphorylation sites present in this part of the protein. (Fig. 4B).

Given the role of QSK1 phosphorylation in its relocalization to PD, its *in vitro* phosphorylation

by CRK2 suggested a phosphorylation-dependent regulatory function of CRK2 in controlling the

subcellular localization of QSK1.

To determine this *in vivo*, we applied a quantitative phosphoproteomics approach and compared

the levels of QSK1 phosphopeptides isolated from WT and *crk2* plants under control and salt

stress conditions. We detected the dually phosphorylated peptide LIEEVSHSSGSPNPVSD in

both WT and *crk2* under normal and stress conditions. This phosphopeptide contains the Ser-621

and Ser-626 that are previously shown to be phosphorylated in phosphoproteomics studies[42–45].

Interestingly, the abundance of the LIEEVSHSSGSPNPVSD phosphopeptide was reduced

following salt treatment in WT but not in *crk2*. Moreover, the abundance of the phosphopeptide

was lower in *crk2* compared to WT under normal growth conditions (Fig. 4C, Table S7). Lower

abundance in *crk2* may suggest that CRK2 phosphorylates QSK1 in the peptide

LIEEVSHSSGSPNPVSD. This finding aligns with the FRET-FLIM assay where we observed

that mannitol stress disrupted the binding of CRK2 and QSK1 and placed a role of

phosphorylation-dependent coupling of the RLKs in osmotic stress signaling.

Phosphorylation of QSK1 at serines 621 and 626 inhibits its stress-induced enrichment at

PD

QSK1 relocalization in response to osmotic stress is associated with its phosphorylation status at

Ser-621 and Ser-626[21]. SIRK1 phosphorylates QSK1 at Ser-621 and Ser-626 during sucrose-

induced osmotic conditions[32]. Therefore, we examined if QSK1 relocalization to PD was

dependent on SIRK1 by testing QSK1-mCherry/*sirk1* plants. We found that QSK1 exhibited

osmotic stress-induced PD enrichment in QSK1-mCherry/*sirk1* plants without impairment. An

enrichment index of 1.967 was observed in *sirk1* following osmotic stress compared to 1.092 in

control conditions (Fig. S4A-B, Table S3). This finding suggested, that while phosphorylation of

Ser-621 and Ser-626 plays an important role in QSK1 recruitment to PD, SIRK1-mediated

phosphorylation does not appear to regulate PD recruitment of QSK1.

Since we found that CRK2 also targeted QSK1 phosphorylation in the phosphopeptide

LIEEVSHSSGSPNPVSD (Fig. 4) and kinase active CRK2 was required for QSK1 recruitment at

PD (Fig. 2C-D), we explored the role of Ser-621 and Ser-626 phosphorylation in context of

CRK2. We stably expressed phosphomimetic QSK1 at Ser-621 and Ser-626 by replacing both

serines by glutamic acid and aspartic acid in *crk2* (QSK1$^{(S621D-S626D)}$-mCherry/*crk2*; QSK1$^{(S621E-}$

$^{S626E)}$-mCherry/*crk2*) and in WT background as control (QSK1$^{(S621D-S626D)}$-mCherry/WT;

QSK1$^{(S621E-S626E)}$-mCherry/WT). We found that enrichment of phosphomimetic QSK1$^{(S621D-}$

$^{S626D)}$-mCherry in the WT background was negligible and statistically insignificant and only

showed an increase of enrichment index from 1.045 to 1.119 upon mannitol treatment (Fig. 5A-

B). Similarly, negligible relocalization of QSK1$^{(S621D-S626D)}$-mCherry was observed in the *crk2*

mutant background (Fig. 5A-B). Consistently, the phosphomimetic variant QSK1$^{(S621E-S626E)}$-

mCherry, showed a mild enrichment of 1.336 from 1.017 following mannitol treatment in the

WT background and poor enrichment in the *crk2* mutant background (1.011 to 1.117) (Fig. S4C-

D). This enrichment of QSK1$^{(S621E-S626E)}$-mCherry in WT background was still weaker and

delayed compared to that in the QSK1-mCherry/WT plants. This led us to test the PD

recruitment of non-phosphorylatable QSK1, in which the Ser-621 and Ser-626 were replaced by

alanines, in WT and *crk2* backgrounds (QSK1$^{(S621A-S626A)}$-mCherry/WT; QSK1$^{(S621A-S626A)}$-

mCherry/*crk2*). We observed a higher enrichment of QSK1$^{(S621A-S626A)}$-mCherry at PD in

response to mannitol treatment (Fig. 5C-D) in QSK1$^{(S621A-S626A)}$-mCherry/WT plants. To a lesser

extent, QSK1$^{(S621A-S626A)}$-mCherry was also present at PD even without mannitol treatment (Fig.

5C-D). The enrichment index for QSK1$^{(S621A-S626A)}$-mCherry/WT in control condition was 1.338

and increased to 2.018 which was even higher than QSK1-mCherry/WT under mannitol

treatment (Fig. 2B). We also validated this result in QSK1$^{(S621A-S626A)}$-mCherry/*qsk1* lines and the

findings were consistent with a clear and intense enrichment of QSK1$^{(S621A-S626A)}$-mCherry in the

qsk1 mutant background (Fig. S5E-F). These findings confirmed that phosphorylation of QSK1

at Ser-621 and Ser-626 was inhibitory for its recruitment to PD. Further, attenuated or complete

inability of QSK1 relocalization to PD in the *crk2* and CRK2^{D450N} backgrounds suggested an

essential role for CRK2-mediated phosphorylation of yet unknown factor(s) involved in this

process.

Phosphorylation of QSK1 at serines 621 and 626 regulates plant stress response and
suppresses osmotic stress-induced callose deposition

302 To test the physiological relevance of QSK1 relocalization to PD and callose deposition during

303 osmotic stress, we carried out root growth assays of plants with WT QSK1 (QSK1-

304 mCherry/*qsk1*), non-phosphorylatable QSK1 (QSK1$^{(S621A-S626A)}$-mCherry/*qsk1*) and phospho-

305 mimetic QSK1 (QSK1$^{(S621D-S626D)}$-mCherry/*qsk1*) with and without salinity stress. While the PR

length of all three genotypes was similar under control conditions (Fig. S4I, Table S8), the QSK1$^{(S621A-S626A)}$ plants exhibited maximum retardation in the PR elongation upon salinity stress (Fig. 5E). In contrast, the phospho-mimetic QSK1 plants (QSK1$^{(S621D-S626D)}$) showed even higher PR elongation than the plants with WT QSK1 under salinity stress (Fig. 5E). The LR density, did not statistically vary among the genotypes under both control and salt stress conditions (Fig. S4J, Fig. 5F), nevertheless, a slightly higher LR density was observed in the QSK1$^{(S621A-S626A)}$-mCherry/*qsk1* plants under salt stress and an opposite trend was observed for the QSK1$^{(S621D-S626D)}$-mCherry/*qsk1* plants (Fig. 5F). A positive correlation between QSK1 availability at PD and callose accumulation in the root epidermal layer is already established[21]. Since in our conditions, non-phosphorylatable QSK1 (QSK1$^{(S621A-S626A)}$) was highly enriched at PD in leaf epidermal cells, we tested the effect of QSK1 recruitment at PD on callose accumulation in the leaf epidermis by aniline blue staining. We quantified callose levels after 7 hours of mannitol treatment to QSK1-mCherry/*qsk1*, QSK1$^{(S621A-S626A)}$-mCherry/*qsk1* and QSK1$^{(S621D-S626D)}$-mCherry/*qsk1* plants (Fig. 5G-H, Fig. S4G-H, Table S9). Plants expressing WT or non-phosphorylatable QSK1 (QSK1$^{(S621A-S626A)}$) accumulated higher callose upon mannitol treatment compared to their respective control conditions. However, plants expressing phosphomimetic QSK1 (QSK1$^{(S621D-S626D)}$) responded weakly to mannitol treatment (Fig. 5G-H, Fig. S4G-H). In control conditions, slightly higher callose levels were detected in the QSK1$^{(S621A-S626A)}$ plants in terms of both intensity and number of deposits compared to the plants with WT QSK1 (Fig. 5G-H, Fig. S4G-H) which is consistent with a higher basal presence of QSK1$^{(S621A-S626A)}$ at PD even without stress treatment (Fig. 5C-D), however, the differences in the overall basal callose accumulation were statistically insignificant among the three genotypes which is consistent with our finding that QSK1 might not play a regulatory role in basal callose deposition (Fig. 1E-F). Collectively, these results established the physiological relevance of the QSK1 relocalization to PD during osmotic stress and highlighted protein phosphorylation as a regulatory mechanism in growth adjustments.

Discussion

Symplastic communication through PD is crucial for coordinating plant growth and stress responses[14,15,46–48]. Dynamic regulation of symplastic connectivity is achieved through PD-associated proteins[49], lipid composition[50] and callose deposition[18]. Current knowledge on the

changes of callose accumulation and symplastic communication is largely focused on immune

response[19,20,51–53]. Studies show altered symplastic communication during abiotic stress

conditions such as metal toxicity[54,55], low temperature[56] and mechanical wounding[57]. Despite the

widespread occurrence of osmotic stress in plants, understanding of how callose accumulation

and PD connectivity respond to it remains limited[9–11,21].

RLKs are involved in sensing and integrating extracellular signals and they typically function in

complexes rather than acting alone. For example, FLS2-BAK1, EFR-BAK1 and CERK1-LYK5

(Chitin elicitor receptor kinase 1-Lysin motif-containing receptor-like kinase 5)[58] in immune

signaling[59,60], BRI1-BAK1 in BR signaling[61], SIRK1-QSK1 in regulation of aquaporins[32,33].

Importantly, the combination of different RLKs can determine the functional outcome. The

RLKs CRK2 and QSK1 mediate plant growth and osmotic stress response through

plasmodesmal regulation[9,21], however, how they coordinate to control callose levels during

osmotic stress is so far not understood. In this study, we identify a regulatory mechanism that

precisely fine-tunes callose deposition during osmotic stress response, a key part of plant growth

adaptation to changing environmental conditions. We report that under normal growth

conditions, CRK2 inhibits callose deposition by engaging QSK1 at the general PM where the

CRK2-QSK1 complex potentially has another role specific to the general PM.

The growth phenotypes of *crk2*, *qsk1* and the *crk2 qsk1* mutants suggested that CRK2 and QSK1

are important for normal plant growth and stress response. The *crk2* mutant exhibits strongest

phenotypic alterations compared to WT Arabidopsis which may be due to pleotropic functions of

CRK2[9,37,62,63]. The smaller rosette and salt hypersensitivity during germination in the *crk2*

mutant was rescued in the *crk2 qsk1* mutant suggesting antagonistic roles of CRK2 and QSK1 in

the biological processes that result in observed phenotypes. Further, mutants defective in callose

regulation have defects in LR patterning and PR length due to altered symplastic

connection[9,12,17]. In our analyses, the root phenotype of the *crk2 qsk1* mutant was milder

compared to the *crk2* and did not rescue to the WT level implying that the absence of both

functional CRK2 and QSK1 results in defects in processes regulating PR growth and LR

emergence.

Further, CRK2 and QSK1 also showed antagonistic effects on callose deposition especially

under osmotic stress conditions. A higher callose level in *crk2* suggested an inhibitory role in

callose deposition which could result from multiple yet unknown actions of CRK2. For example,

CRK2-targeted phosphorylation of CALLOSE SYNTHASES (CALS)[9] could be one of the

regulatory mechanisms determining callose levels. Further, CRK2 also regulates ROS production

through RBOHD[62] which could add another layer of CRK2-mediated regulation of callose

production. In the *qsk1* mutant, a reduced callose deposition upon mannitol treatment suggested

that the inhibitory action of CRK2 is ensued following osmotic stress despite the absence of

QSK1 which gets further support from the result that QSK1 was not found to be necessary to

recruit CRK2 at the PD. In the *crk2 qsk1* mutant, the WT-like response indicated alleviation of

the inhibitory effect of CRK2 and QSK1 on each other and possibly the presence of

compensatory or feedback mechanisms that maintain callose homeostasis.

Protein phosphorylation is a common regulatory mechanism across living organisms for

dynamically reshaping the binding repertoire of a given protein and thereby providing functional

specificity[64,65]. The changes in the protein-protein interaction networks by phosphorylation can

also lead to changes in the localization and protein abundance[30,66]. A regulatory role of QSK1

phosphorylation at Ser-621 and S-626 has been previously describd[42–44]. For example, QSK1

phosphorylation at Ser-621 and Ser-626 by SIRK1, an LRR RLK, stabilizes the SIRK1-QSK1

protein complex involved in the regulation of aquaporins in roots[32]. QSK1 phosphorylation at

these two residues also enhances its interaction with the nitrate transceptor NRT1.1 which

eventually transduces low nitrate signals to PM H^+-ATPase AHA2[36].

Through *in vitro* kinase assays, we identified that CRK2 phosphorylates multiple residues in the

cytosolic region of QSK1 including Ser-626. Further, *in vivo* phosphorylation status of QSK1

within LIEEVSHSSGSPNPVSD peptide was altered in the *crk2* mutant. QSK1 was shown to

interact with CRK2 *in planta*[9], and FRET-FLIM analyses together with structural modeling

supported that CRK2 and QSK1 exist in a preformed complex at the general PM (Fig. 6 panel

A). Interestingly, the CRK2 and QSK1 affinity was abolished when CRK2 was replaced by

CRK2[D450N]. Together, these results and existing literature suggested that phosphorylation of

QSK1 at Ser-621 and Ser-626 could be a common regulatory mechanism to stabilize QSK1

protein complexes, including the CRK2-QSK1 complex. *In vivo* phosphoproteomics analyses

suggested reduced phosphorylation of the LIEEVSHSSGSPNPVSD peptide at two serine

residues following salt treatment in WT. Consistently, mannitol treatment reduced the binding

affinity of the CRK2-QSK1 complex in the FRET-FLIM assays. These findings suggest, that an
unknown protein phosphatase might act on QSK1 to allow dissociation of the CRK2-QSK1
complex (Fig. 6 panel B) leading to the PD localization of QSK1 prior to the relocalization of
CRK2. Consistently, our microscopy-based relocalization studies showed that phosphorylation of
QSK1 at Ser-621 and Ser-626 is inhibitory for the recruitment of QSK1 to PD and consequently
callose accumulation following mannitol treatment. Non-phosphorylatable QSK1$^{\text{S621A-S626A}}$,
efficiently relocalized to PD while QSK1 phosphomimetic substitutions at Ser-621 and Ser-626
abated its recruitment to PD. Collectively, these results demonstrated a phosphorylation-
dependent subcellular dichotomy for QSK1 conditionally associating with and dissociating from
CRK2 and probably other proteins. This mechanism could direct QSK1 towards distinct
molecular pathways to promote growth adaptability under osmotic stress.

Although phosphorylation of QSK1 was inhibitory for its relocalization to PD (Fig. 5A-D),
enzymatically active CRK2 was still essential for QSK1 recruitment at PD (Fig. 2A-D). This
suggested the possibility that CRK2 might phosphorylate another protein(s) in addition to QSK1,
whose phosphorylation by CRK2 would be necessary to facilitate the relocalization of QSK1 to
PD (Fig. 6 panel B). The PM could be biochemically divided into detergent resistant membrane
(DRM) and detergent soluble membrane (DSM) domains, and PD-associated membranes are
considered to be DRMs. Proteins identified in PD proteomes and DRM fractions show a large
overlap[21] and QSK1 has been found in several DRM and PD proteomes (compiled in[21]). The
actin cytoskeleton is necessary for the distribution of proteins between DRM and DSM and
disruption of the actin cytoskeleton depletes QSK1 from DRMs[67]. We previously identified actin
isoforms, as well as other DRM-localized proteins as interactors of CRK2 in pull-down assays
followed by MS analyses[9]. Therefore, we speculate on regulation of the actin cytoskeleton or
other DRM-localized proteins by CRK2 to enable QSK1 relocalization after its
dephosphorylation, however, more studies are required to support this.

Adjusting root system architecture is one of the main strategies plants employ during extended
periods of salinity stress for efficient water uptake[68]. QSK1 phosphorylation at Ser-621 and Ser-
626 affects callose deposition in the epidermal layer of root [21] as well as leaves, resulting in
altered symplastic connectivity and root phenotype. Adjusting callose levels through QSK1 can
therefore be a contributory mechanism plants employ for osmotic stress resilience. In our

physiological analyses, we found that the plants accumulating higher callose levels upon osmotic

stress, exhibit a root phenotype adjusted for osmotic stress.

Based on these findings, we propose a working model in which symplastic connectivity is

dynamically regulated through temporally coordinated signaling between CRK2 and QSK1 (Fig.

6). We propose that CRK2 works as a moderator of QSK1-mediated osmotic stress responses.

CRK2 phosphorylates QSK1 and sequesters it at the general PM. This complex might have a

function specific to their loacalization at the general PM. At the same time, existence of such a

complex inhibits unnecessary QSK1 function at PD in the absence of osmotic stress.

Dephosphorylation of QSK1 at the C-terminus upon stress stimuli dissociates it from CRK2

allowing it to relocalize to PD to positively regulate callose deposition. At a later stage, CRK2

moves to PD itself where it likely finds different phosphorylation targets than at the PM and

eventually negatively regulates callose deposition to switch it off and prevent unnecessary

clogging of PD. In conclusion, our work shows that CRK2 controls the dynamic and stimulus-

dependent relocalization of QSK1, and likely also other proteins, to PD. PD and other membrane

nanodomains contain constitutive protein components[23], however, the dynamics of QSK1 and

CRK2 demonstrate the necessity and potential for the precise spatiotemporal regulation of the

cellular response machinery. In the future, it will be highly interesting to investigate the

spatiotemporal dynamics of other proteins showing conditional clustering in nanodomains vs. the

general plasma membrane.

Material and methods

Plant materials and growth conditions

The *Arabidopsis thaliana* ecotype Col-0 was used as wild-type control. The seeds of mutants

qsk1 (SALK_019840; AT3G02880), *crk2* (SALK_012659C; At1g70520), and *sirk1*

(SALK_125543C; At5g10020) were obtained from NASC (University of Nottingham). The *crk2*

qsk1 double mutant was generated by crossing *qsk1* and *crk2*. For growing plants under sterile

conditions, seeds were surface-sterilized and stratified in dark for 3 days at 4°C and plated on

0.5X Murashige and Skoog media (Sigma-Aldrich) supplemented with 0.8% (w/v) agar, 1%

(w/v) Suc, and 0.1% (w/v) MES, pH 5.8. This media composition was used as standard growth

media for all the *in vitro* experiments unless specified. Plants were grown in climate-controlled

growth chambers with a 16-h light, 8-h dark photoperiod. For experiments with adult stages,

seed propagation, and growth of selected transgenics, the plants were grown in substrate mix (2:1
peat:vermiculite) in a Weiss Gallenkamp growth chamber or greenhouse with long-day
conditions at 21°C.

Physiological assays

For comparative phenotypic analysis of WT, *crk2*, *qsk1*, and *crk2 qsk1* plants, 10 plants for each
genotype were planted in individual pots with substrate mix (2:1 peat:vermiculite). Plants were
allowed to grow in a Weiss Gallenkamp growth chamber with long-day conditions at 21°C with
equal irrigation and randomization of pot positioning. Images were captured at various growth
stages. Related data and statistical analysis are given in Table S1.

For root elongation assays, 5-day-old seedlings grown in vertically positioned square petri plates
in Photon Systems Instruments (PSI) growth chamber were transferred to standard 0.5X MS
medium supplemented with 100 mM NaCl and the primary root tip was marked. For the control
group, the seedlings were transferred to medium without NaCl. The images were captured after
another 7 days and the root elongation was measured with the program Fiji
(https://imagej.net/software/fiji/). The data obtained were the average of at least 13-15 seedlings
and all the experiments were repeated three times unless specified. Related data and statistical
analysis are given in Table S1.

For the seed germination assay, surface-sterilized seeds were plated evenly in petri dishes with
standard 0.5X MS medium supplemented with 150 mM NaCl. For the control group, the seeds
were sown on the medium without NaCl. The seedlings were allowed to grow in a PSI growth
chamber and the images of the plates were captured after 12 days. For each genotype and
treatment, 3 plates were taken as biological replicates and in each plate 30-35 seedlings were
taken. The percentage of seed germination was first estimated in control plates to normalize the
effect of non-viable seeds. The total number of seeds and the viable germinated seeds with
clearly visible root and green cotyledon/leaves were counted. Related data and statistical analysis
are given in Table S1.

Osmotic stress treatment and callose estimation

For estimation of osmotic stress-induced callose deposition, seeds were sown hydroponically for
two weeks on a mesh in liquid 0.5X MS with 1% sucrose in standard growth conditions. For

inducing callose deposition under osmotic stress, the roots were subjected to 50 mM mannitol in
liquid medium for 7 hours. After treatment, the plants were pulled out of the mesh and infiltrated
with 0.2%-0.5% aniline blue (constant for a single experiment) and the first pair of true leaves
were imaged in a laser confocal microscope. For each genotype, 5 plants were imaged and 5-6
areas were scanned from first pair of true leaves of each plant. The control plants underwent the
same process without mannitol treatment. The mean intensity and number of deposits were
measured using Fiji. To quantify overall callose levels that include both intensity and number of
deposits, a composite metric was derived by multiplying mean intensity of deposits and number
of deposits and dividing it with a constant scaling factor to yield a number suitable for
comparative analysis. This is termed as total callose deposition. Related data and statistical
analysis are given in Table S2 and S9.

Plasmid vector construction and transgenic lines

Plasmid constructs for P_{QSK1}:$QSK1$-mCherry, P_{QSK1}:$QSK1^{(S621A-S626A)}$-mCherry, P_{QSK1}:$QSK1^{(S621D-}$
$^{S626D)}$-mCherry and P_{QSK1}:$QSK1^{(S621E-S626E)}$-mCherry were generated by the MultiSite Gateway
technology (Invitrogen; Thermo Fisher Scientific). The donor clones for promoter regions of
QSK1 and coding sequence of *QSK1* were cloned in the vectors pDONR™P4-P1R and
pDONR™221, respectively. The donor clone for mCherry in the vector pDONR™P2R-P3 was
already available in the laboratory. The donor clones were combined in an LR reaction to obtain
the final constructs in the binary vector pH7m34GW (VIB-UGENT). The phospho-mutant
versions of QSK1 (S621A-S626A, S621D-S626D and S621E-S626E) were generated by
introducing point mutation by DpnI-mediated site-directed mutagenesis[70]. The primers are listed
in Table S10. P_{CRK2}:$CRK2$- mVenus and P_{CRK2}:$CRK2^{D450N}$-mVenus constructs were available in
the binary vector pBm43GW (Invitrogen; Thermo Fisher Scientific) (Hunter et al 2019). The
final clones in the binary vectors were introduced in *Agrobacterium tumefaciens* GV3101_pSoup
and Arabidopsis plants of desired genotypes were transformed by floral dip method[71]. The
CRK2/CRK2^{D450N}-mVenus and QSK1-mCherry co-expressing lines were generated by crossing
P_{CRK2}:$CRK2$-mVenus/*crk2* and P_{QSK1}:$QSK1$-mCherry/*crk2*, and, P_{CRK2}:$CRK2^{D450N}$-mVenus/*crk2*
and P_{QSK1}:$QSK1$-mCherry/*crk2* plants. Transformants were selected on 0.5X MS medium with
appropriate antibiotic or herbicide [(hygromycin 20 ug/ml (CARL ROTH) or Basta 10 ug/ml
(DL-phosphinothricin; Duchefa Biochemie)].

Osmotic stress treatment and confocal imaging

For visualizing localization of the fluorescently-tagged proteins in response to osmotic stress, the mature leaves of 4-week-old Arabidopsis were detached from the plant and submerged in 0.4 M mannitol or 150 mM NaCl in 0.5X MS solution. For detecting co-localization with aniline blue, the samples were treated with 0.2% aniline blue simultaneously with mock or 0.4 M mannitol and 150 mM NaCl treatments. After the incubation period, the same area of all the leaf samples was targeted for imaging and mounted in the same solution and the abaxial side was imaged. A mock-treated leaf of the same plant was used as a control. All the live-imaging experiments were conducted on an Olympus FLUOVIEW FV3000 confocal laser scanning microscope with 60.0 X or 40.0 X oil-immersion objectives and was operated by Olympus FV31S-SW software. The excitation wavelengths for mCherry, mVenus and DAPI (for aniline blue) were 561, 488 and 405, respectively. The detection wavelengths were 570-620 nm for mCherry, 500-540 nm for mVenus and 450-550 nm for DAPI. The laser transmittivity was kept constant for a single experiment and for the comparisons of fluorescence intensities unless otherwise specified. Co-localizations were detected in sequential scanning mode to prevent cross bleeding of emission spectra.

Quantification of protein enrichment

To quantify the extent of protein enrichment, we used the method described in [72] with modification. We assumed that the intense dots of the fluorescently-tagged protein correspond to PD based on the fact that these dots co-localized with the aniline blue-stained callose dots in our analysis (Fig. S2B). Under salt and mannitol treatments, the punctate pattern of CRK2 and QSK1 correspond to their PD localization and previous studies have confirmed this with co-localization with PD-located protein markers (PDLPs) in addition to aniline blue[9,21]. Using Fiji, we selected a circular region of interest (ROI) on such intensity spots resembling PD. The ROI adjacent to the enrichment spots was considered as bulk PM and the ratio of the two was referred to as 'enrichment index'. In each image, 5-6 pairs of such ROIs were measured for the intensities and enrichment indexes were determined. 3 images for each treatment/genotype were analyzed. The relocalization experiments were conducted on 3 independent transgenic lines for each genotype. The data related to enrichment index and statistical analysis are given in Table S3.

FRET-FLIM

The experiments on Förster resonance energy transfer detected by fluorescence lifetime (FRET-

FLIM) were performed on the confocal microscope FV3000 (Olympus) equipped with the 40.0

X UApo N water immersion objective lens NA 1.15 (Olympus), MultiHarp 150 Time-Correlated

Single Photon Counting device, pulsed 488 nm laser, and two Hybrid Photomultiplier detectors

(all Picoquant). All data were acquired at 40 MHz pulse repetition rate. Emitted fluorescence was

filtered using the FF02-525/40-25 emission filter (Semrock). Data analysis was done using

Picoquant SymphoTime64 software package. Membrane regions were manually selected in the

images, and lifetime (τ) values were calculated from decay curves using exponential tailfit with

the fixed time window for analysis. The acceptable goodness of fit $\chi 2$ values were between 0.9

and 1.2. The stage of the plants and treatment used in FRET-FLIM analysis are as described in

the method section 'osmotic stress treatment and live confocal imaging'. Related data and

statistical analysis are given in Table S4.

Structural modeling

558 The interactions of full-length and truncated CRK2 and QSK1 were predicted using Colabfold

559 (version 1.5.2) implementation of AlphaFold2[40] multimer version 1 with a maximum of 48

560 recycles or 0.5 tolerance. The amino acid boundaries selected for both the proteins are given in

561 Table S5.

Identification of phosphopeptides from *in vitro* kinase assay

The cytosolic region of QSK1 coding sequence QSK1cyto was amplified and cloned into pMal-

c2X (Addgene, #75286) using *BamH*I and *Sal*I restriction sites. The primers are given in Table

S10. The phospho-negative QSK1cyto(S621A-S626A) was generated by *Dpn*I-mediated site-

directed mutagenesis and cloned in the same way as for the WT construct. The construct for the

cytosolic region of CRK2 was available (CRK2cyto-pOPINM;[62]). GST-tagged CRK2cyto and

MBP-tagged QSK1cyto were expressed in the *E. coli* strain SHuffle T7 (New England Biolabs,

C3026J) and BL21-CodonPlus-RIL (Agilent, 230240), respectively, and eventually purified with

glutathione sepharose 4B (GE Healthcare, 17-0756-01) and amylose resin (New England Biolabs

E8021S) beads, respectively. 10mM maltose was used for the elution of MBP-tagged proteins,

while 20mM reduced glutathione for the elution of GST-tagged proteins. In vitro kinase assay

was performed as described in Kimura et al. 2020, with minor modifications. Briefly, expressed

and purified recombinant proteins were incubated in the ratio 1: 3-5 (kinase: substrate) for 30

min at room temperature in reaction buffer containing γ-32P labelled ATP (0.2µl/20µl of labelled ATP with declared activity 9.25MBq), 50mM HEPES (pH 7.4), 1mM DTT, 10mM MgCl2, 60µM unlabeled ATP. Any specific procedure applied to a specific Figure panel is given in its Figure legend. Enzymatic reactions were terminated by boiling the samples in protein loading dye (98°C, 5 minutes) and proteins were separated by SDS-PAGE. Gels were stained with Coomassie Brilliant Blue and dried. Dried gels were exposed to photosensitive layer and documented with imaging system Typhoon 9410 (Molecular Dynamics). For the purpose of the identification of phosphorylation sites, labelled ATP was replaced by the respective volume of deionized water. Separation and staining of the gels were followed by in-gel trypsin digestion (Thermo Fisher Scientific, 90057). Digested peptide samples were dissolved in 10 µl of 0.1% formic acid and 5 µl was injected for LC-ESI-MS/MS analysis. The LC-ESI-MS/MS analysis was performed on a nanoflow HPLC system (Easy-nLC1000, Thermo Fisher Scientific) coupled to the Q Exactive HF mass spectrometer (Thermo Scientific, Bremen, Germany) equipped with a nano-electrospray ionization source. Peptides were first loaded on a trapping column and subsequently separated inline on a 15 cm C18 column (75 µm x 15 cm, ReproSil-Pur 3 µm 120 Å C18-AQ, Dr. Maisch HPLC GmbH, Ammerbuch-Entringen, Germany). The mobile phase consisted of water with 0.1% formic acid (solvent A) and acetonitrile/water (80:20 (v/v)) with 0.1% formic acid (solvent B). Peptides were eluted using a 30 min gradient: 5% to 21% solvent B in 11 min, 21% to 36% solvent B in 9 min, and 36% to 100% solvent B in 5 min, followed by a 5 min wash with 100% solvent B. MS data was acquired automatically by using Thermo Xcalibur 4.1 software (Thermo Scientific). The data-dependent acquisition method consisted of an Orbitrap MS survey scan of mass range 350–1750 m/z with a resolution of 120000 and AGC target of 3000000. This was followed by HCD fragmentation for 10 most intense peptide ions with resolution of 15000 and AGC target of 5000. Raw files were searched for protein identifications using Proteome Discoverer 2.5 software (Thermo Fisher Scientific) connected to an in-house Mascot 2.8.2 server (Matrix Science). Data was searched against a SwissProt database (version 2022_03) with taxonomy restricted to *Arabidopsis thaliana*.

Phosphoproteomics

Plant treatment- Arabidopsis seeds were sterilized by incubating with 1.5% NaClO 0.02% Triton X-100 solution for 5 min and vernalized at 4 °C for 2 days. Sterilized seeds were

germinated and grown in liquid culture on 12 well plates (30 seeds/well) in 1/2 MS medium with

1% sucrose and 0.5g MES/l at 23 °C under continuous light (100 µmol m^{-2} s^{-1}) in a Percival

growth chamber. Plates with 7-day old seedlings were transferred from the growth chamber to a

workbench and kept o/n for acclimatization before treatments. Seedlings were treated with either

150 mM NaCl or sterile water for 30 minutes after which seedlings were immediately collected

and flash-frozen in liquid nitrogen and stored at -80 °C. ***Sample preparation and***

phosphopeptide enrichment- Frozen seedlings were disrupted using a Retsch mill with 2 steel

beads at 50 Hz for 5 min. To the finely ground material was added 500 µL urea extraction buffer

(8M in 100 mM Tris-HCl pH 8.5, 20 µL/mL phosphatase inhibitor cocktail 2 (Sigma, P5726), 20

µL/mL phosphatase inhibitor cocktail 3 (Sigma, P0044), 5 mM DDT), samples were vortexed

thoroughly to ensure wetting of all powder while thawing. Samples were incubated at RT for 30

min with shaking. Next samples were centrifuged for 5 min at 10k x g, supernatants were

transferred to fresh tubes and centrifuged at full speed for 10 mins. This procedure was repeated

until no visible pellet was retained. Protein concentration of the extract was determined using

Pierce 660 mn protein assay. Samples were alkylated using chloroacetamide (CAA) (550 mM

stock, 14 mM final concentration) for 30 min at RT in the dark, after which samples were

subjected to an in-solution digestion. In brief, samples (400 µL) were diluted with 400 µL

100mM Tris, pH 8.5, 1mM CaCl$_2$, then 5 µg Lys-C (Merck) was added. Samples were incubated

for 4h at room temperature, after which 2.4 mL 100mM Tris, pH 8.5, 1 mM CaCl$_2$ and 5 µg

trypsin were added. Samples were incubated o/n at 37 °C. Samples were acidified with 20 µL

TFA and desalted using C18 SepPaks (1cc cartridge, 100 mg (WAT023590)). In brief, SepPaks

were conditioned using methanol (1 mL), buffer B (80% acetonitrile, 0.1% TFA) (1 mL) and

buffer A (0.1% TFA) (2 mL). Samples were loaded by gravity flow, washed with buffer A (1 x 1

mL, 1x 2 mL) and eluted with buffer B (2 x 400 µL). 40 µL of eluates were used for peptide

measurement and concentrated as input control. For phosphopeptide enrichment by metal-oxide

chromatography (MOC) (adapted from: Nakagami 2014) the samples were evaporated to a

sample volume of 50 µL and diluted with sample buffer (2 mL AcN, 820 µL lactic acid (LA), 2.5

µL TFA / 80% ACN, 0.1% TFA, 300mg/ml LA, final concentrations) (282 µL). MOC tips were

prepared by loading 100 µL of a well-mixed slurry of 15mg metal oxide beads (Titansphere TiO$_2$

beads 10 µm (GL Science Inc, Japan, Cat. No. 5020-75010) in 500 µL MeOH onto 4 C2 micro

columns/condition (3 mg/column) and centrifugation for 5 min at 1500g. Tips were washed with

centrifugation at 1500g for 5 min using 80 µL of solution B (80% acetonitrile, 0.1% TFA) and 80

µL of solution C (300 mg/mL LA in solution B). To simplify the processing, samples tips were

fitted onto a 96/500 µL deep well plate (Protein LoBind, (Eppendorf Cat. No. 0030504100).

After washing MOC tips were transferred to a fresh plate, samples were loaded onto the

equilibrated tips and centrifuged for 10 min at 1000g. The flow through was reloaded onto the

tips and centrifugation was repeated. Tips were washed with centrifugation at 1500g for 5 min

using 80 µL of solution C and 3x 80 µL of solution B. For the elution of the enriched

phosphopeptides the tips were transferred to a fresh 96/500 µL deep well plate containing 100

µL/well of acidification buffer (20% phosphoric acid). Peptides were eluted first with 50 µL

elution buffer 1 (5% NH_4OH) and centrifugation for 5 min at 800g, then with 50 µL of elution

buffer 2 (10% piperidine) and centrifugation for 5 min at 800g. Next, the samples were desalted

using StageTips with C18 Empore disk membranes (3 M) [74], dried in a vacuum evaporator, and

dissolved in 10 µL 2% ACN, 0.1% TFA (A* buffer) for MS analysis. ***Data acquisition-*** Samples

were analysed using an Ultimate 3000 RSLC nano (Thermo Fisher) coupled to an Orbitrap

Exploris 480 mass spectrometer equipped with a FAIMS Pro interface for Field asymmetric ion

mobility separation (Thermo Fisher). Peptides were pre-concentrated on an Acclaim PepMap 100

pre-column (75 µM x 2 cm, C18, 3 µM, 100 Å, Thermo Fisher) using the loading pump and

buffer A** (water, 0.1% TFA) with a flow of 7 µl/min for 5 min. Peptides were separated on 16

cm frit-less silica emitters (New Objective, 75 µm inner diameter), packed in-house with

reversed-phase ReproSil-Pur C18 AQ 1.9 µm resin (Dr. Maisch). Peptides were loaded on the

column and eluted for 130 min using a segmented linear gradient of 5% to 95% solvent B (0 min

: 5%B; 0-5 min -> 5%B; 5-65 min -> 20%B; 65-90 min ->35%B; 90-100 min -> 55%; 100-105

min ->95%, 105-115 min ->95%, 115-115.1 min -> 5%, 115.1-130 min ->5%) (solvent A 0%

ACN, 0.1% FA; solvent B 80% ACN, 0.1%FA) at a flow rate of 300 nL/min. Mass spectra were

acquired in data-dependent acquisition mode with a TOP_S method using a cycle time of 2

seconds. For field asymmetric ion mobility separation (FAIMS) two compensation voltages (-45

and -65) were applied, the cycle time for the CV-45 experiment was set to 1.2 seconds and for

the CV-65 experiment to 0.8 sec. MS spectra were acquired in the Orbitrap analyzer with a mass

range of 320–1200 m/z at a resolution of 60,000 FWHM and a normalized AGC target of 300%.

Precursors were filtered using the MIPS option (MIPS mode = peptide), the intensity threshold

was set to 5000, Precursors were selected with an isolation window of 1.6 m/z. HCD

fragmentation was performed at a normalized collision energy of 30%. MS/MS spectra were

acquired with a target value of 75% ions at a resolution of 15,000 FWHM, at an injection time of

120 ms and a fixed first mass of m/z 120. Peptides with a charge of +1, greater than 6, or with

unassigned charge state were excluded from fragmentation for MS^2. ***Data analysis***

phosphoproteome analysis-Raw data were processed using MaxQuant software (version 1.6.3.4,

http://www.maxquant.org/)[75] with label-free quantification (LFQ) and iBAQ enabled[76]. MS/MS

spectra were searched by the Andromeda search engine against a combined database containing

the sequences from *A. thaliana* (TAIR10_pep_20101214 database), and sequences of 248

common contaminant proteins and decoy sequences. Trypsin specificity was required and a

maximum of two missed cleavages allowed. Minimal peptide length was set to seven amino

acids. Carbamidomethylation of cysteine residues was set as fixed, phosphorylation of serine,

threonine and tyrosine, oxidation of methionine and protein N-terminal acetylation as variable

modifications. The match between runs option was enabled. Peptide-spectrum-matches and

proteins were retained if they were below a false discovery rate of 1% in both cases. Statistical

analysis of the intensity values obtained for the phospho-modified peptides

("modificationSpecificPeptides.txt" output file) was carried out using Perseus (version 1.5.8.5,

http://www.maxquant.org/). Intensities were filtered for reverse and contaminant hits and the

data was filtered to retain only phospho-modified peptides. Next. intensity values were log2

transformed. After grouping samples by condition only those sites were retained for the

subsequent analysis that had three valid values in one of the conditions. Two-sample t-tests were

performed using a permutation-based FDR of 0.05. Alternatively, the data was filtered for 4 valid

values in one of the conditions, the valid value-filtered data was median-normalized and missing

values were imputed from a normal distribution, using the default settings in Perseus (1.8

downshift, separately for each column). Volcano plots were generated in Perseus using an FDR

of 5% an S0=1. The Perseus output was exported and further processed using Excel.

Quantification and statistical analysis

Fiji (ImageJ; (https://imagej.net/software/fiji/) was used for image analyses. The information on

statistical analyses is given in the corresponding figure legends and supplementary tables. Instant

Clue[77] and GraphPad Prism v10 were used for graph preparation and statistical analyses. In

general, data normality was tested by Shapiro-Wilk test. Kruskal-Wallis test followed by Dunn's

multiple comparison test was used for those datasets that did not pass the normality test. For
normal data, one-way ANOVA with Tukey's post-hoc test was used. For the datasets with two
variables (genotype and treatment), two-way ANOVA was applied with Tukey's post-hoc
multiple comparison test. Any other specific tests are mentioned in figure legends and
corresponding supplementary tables.

Resource availability

Contact

704 Requests for further information and resources in this study should be directed to and will be
705 fulfilled by the lead contact, Michael Wrzaczek (michael.wrzaczek@umbr.cas.cz).

Materials availability

707 Plasmids and transgenic plant seeds generated in this study are available from the lead contact
708 upon request.

Data and code availability

710 • This paper does not report original code.
711 • The raw MS data from this study (phosphoproteomics profiling) was deposited at the
712   ProteomeXchange Consortium via the PRIDE[69] partner repository with the dataset
713   identifier PXD070133.

Acknowledgments

This work was supported by the Czech Science Foundation grants nr. 23-04866S (MW) and 22-
35680M (RP). FC was funded by EMBO (award number: ALTF 1115-2021; 2022-2024) and the
Czech Ministry of Youth and Education MSCA Fellowships Interfellows (Project number:
CZ.02.01.01/00/22_010/0003414; 2024-2025). We acknowledge Prof. Jiří Friml and Dmitrii
Vladimirtsev (Institute of Science and Technology, Austria) for assistance with FRET-FLIM. We
thank Laboratory of Microscopy and Histology, Biology Centre CAS, supported by project
LM2023050 funded by the Ministry of Education, Youth and Sports of the Czech Republic, with
the instrumental equipment co-financed by the European Union. We also acknowledge Growth
Facility (Jan Kadlec), BC Core Facilities, Institute for Plant Molecular Biology, Biology Centre
CAS. Computational resources used for ColabFold predictions were provided by the e-INFRA

CZ project (ID:90254), supported by the Ministry of Education, Youth and Sports of the Czech Republic. Identification of phosphopeptides from *in vitro* kinase assay was performed at the Turku Proteomics Facility supported by Biocenter Finland. We thank Dr. Julia Krasensky-Wrzaczek for her research input and constructive feedback on the manuscript.

Author contributions

SJ and MW conceived the study and designed the research. AZ and SJ performed *in vitro* kinase assays and callose staining, imaging and quantification. AB, SJ and IK conducted FRET-FLIM experiment. SCS, AH and HN conducted and analyzed phosphoproteomics experiment. FC performed crossing. SJ and SL carried out physiological assays. MN and RP performed structural analysis. MP and JM carried out LC-MS/MS analyses. SJ conducted all other experiments. SJ wrote the first draft and MW edited and reviewed the paper. MW acquired the research grant. All authors participated in this research and contributed to the final draft of the paper.

References

1. Martinière, A., Fiche, J.B., Smokvarska, M., Mari, S., Alcon, C., Dumont, X., Hematy, K., Jaillais, Y., Nollmann, M., and Maurel, C. (2019). Osmotic Stress Activates Two Reactive Oxygen Species Pathways with Distinct Effects on Protein Nanodomains and Diffusion. Plant Physiol *179*, 1581–1593. https://doi.org/10.1104/pp.18.01065.

2. Knight, H., Trewavas, A.J., and Knight, M.R. (1997). Calcium signalling in Arabidopsis thaliana responding to drought and salinity. Plant J *12*, 1067–1078. https://doi.org/10.1046/j.1365-313x.1997.12051067.x.

3. Tracy, F.E., Gilliham, M., Dodd, A.N., Webb, A.A.R., and Tester, M. (2008). NaCl-induced changes in cytosolic free Ca2+ in Arabidopsis thaliana are heterogeneous and modified by external ionic composition. Plant Cell Environ *31*, 1063–1073. https://doi.org/10.1111/j.1365-3040.2008.01817.x.

4. Stephan, A.B., Kunz, H.-H., Yang, E., and Schroeder, J.I. (2016). Rapid hyperosmotic-induced Ca2+ responses in Arabidopsis thaliana exhibit sensory potentiation and involvement of plastidial KEA transporters. Proceedings of the National Academy of Sciences *113*, E5242–E5249. https://doi.org/10.1073/pnas.1519555113.

5. Munnik, T., Meijer, H.J., Ter Riet, B., Hirt, H., Frank, W., Bartels, D., and Musgrave, A. (2000). Hyperosmotic stress stimulates phospholipase D activity and elevates the levels of phosphatidic acid and diacylglycerol pyrophosphate. Plant J *22*, 147–154. https://doi.org/10.1046/j.1365-313x.2000.00725.x.

6. Katagiri, T., Takahashi, S., and Shinozaki, K. (2001). Involvement of a novel Arabidopsis phospholipase D, AtPLDdelta, in dehydration-inducible accumulation of phosphatidic acid in stress signalling. Plant J *26*, 595–605. https://doi.org/10.1046/j.1365-313x.2001.01060.x.

7. Testerink, C., and Munnik, T. (2011). Molecular, cellular, and physiological responses to phosphatidic
acid formation in plants. J Exp Bot *62*, 2349–2361. https://doi.org/10.1093/jxb/err079.

8. Yao, H.-Y., and Xue, H.-W. (2018). Phosphatidic acid plays key roles regulating plant development
and stress responses. Journal of Integrative Plant Biology *60*, 851–863.
https://doi.org/10.1111/jipb.12655.

9. Hunter, K., Kimura, S., Rokka, A., Tran, H.C., Toyota, M., Kukkonen, J.P., and Wrzaczek, M. (2019).
CRK2 Enhances Salt Tolerance by Regulating Callose Deposition in Connection with PLDα1. Plant
Physiol *180*, 2004–2021. https://doi.org/10.1104/pp.19.00560.

10. Sangi, S., Olimpio, G.V., Coelho, F.S., Alexandrino, C.R., Da Cunha, M., and Grativol, C. (2023).
Flagellin and mannitol modulate callose biosynthesis and deposition in soybean seedlings.
Physiologia Plantarum *175*, e13877. https://doi.org/10.1111/ppl.13877.

11. Su, L., Zhang, Y., Yu, S., Geng, L., Lin, S., Ouyang, L., and Jiang, X. (2023). RcbHLH59-RcPRs
module enhances salinity stress tolerance by balancing Na+/K+ through callose deposition in rose
(Rosa chinensis). Hortic Res *10*, uhac291. https://doi.org/10.1093/hr/uhac291.

12. J Maule, A., Gaudioso-Pedraza, R., and Benitez-Alfonso, Y. (2013). Callose deposition and
symplastic connectivity are regulated prior to lateral root emergence. Commun Integr Biol *6*, e26531.
https://doi.org/10.4161/cib.26531.

13. Han, X., Hyun, T.K., Zhang, M., Kumar, R., Koh, E., Kang, B.-H., Lucas, W.J., and Kim, J.-Y.
(2014). Auxin-Callose-Mediated Plasmodesmal Gating Is Essential for Tropic Auxin Gradient
Formation and Signaling. Developmental Cell *28*, 132–146.
https://doi.org/10.1016/j.devcel.2013.12.008.

14. Otero, S., Helariutta, Y., and Benitez-Alfonso, Y. (2016). Symplastic communication in organ
formation and tissue patterning. Current Opinion in Plant Biology *29*, 21–28.
https://doi.org/10.1016/j.pbi.2015.10.007.

15. Wu, S., O'Lexy, R., Xu, M., Sang, Y., Chen, X., Yu, Q., and Gallagher, K.L. (2016). Symplastic
signaling instructs cell division, cell expansion, and cell polarity in the ground tissue of Arabidopsis
thaliana roots. Proceedings of the National Academy of Sciences *113*, 11621–11626.
https://doi.org/10.1073/pnas.1610358113.

16. Vatén, A., Dettmer, J., Wu, S., Stierhof, Y.-D., Miyashima, S., Yadav, S.R., Roberts, C.J., Campilho,
A., Bulone, V., Lichtenberger, R., et al. (2011). Callose Biosynthesis Regulates Symplastic
Trafficking during Root Development. Developmental Cell *21*, 1144–1155.
https://doi.org/10.1016/j.devcel.2011.10.006.

17. Benitez-Alfonso, Y., Faulkner, C., Pendle, A., Miyashima, S., Helariutta, Y., and Maule, A. (2013).
Symplastic Intercellular Connectivity Regulates Lateral Root Patterning. Developmental Cell *26*,
136–147. https://doi.org/10.1016/j.devcel.2013.06.010.

18. Wu, S.-W., Kumar, R., Iswanto, A.B.B., and Kim, J.-Y. (2018). Callose balancing at plasmodesmata.
J Exp Bot *69*, 5325–5339. https://doi.org/10.1093/jxb/ery317.

19. Lee, J.-Y., Wang, X., Cui, W., Sager, R., Modla, S., Czymmek, K., Zybaliov, B., van Wijk, K., Zhang,
C., Lu, H., et al. (2011). A plasmodesmata-localized protein mediates crosstalk between cell-to-cell

communication and innate immunity in Arabidopsis. Plant Cell *23*, 3353–3373.
https://doi.org/10.1105/tpc.111.087742.

20. Tee, E.E., Johnston, M.G., Papp, D., and Faulkner, C. (2023). A PDLP-NHL3 complex integrates
plasmodesmal immune signaling cascades. Proceedings of the National Academy of Sciences *120*,
e2216397120. https://doi.org/10.1073/pnas.2216397120.

21. Grison, M.S., Kirk, P., Brault, M.L., Wu, X.N., Schulze, W.X., Benitez-Alfonso, Y., Immel, F., and
Bayer, E.M. (2019). Plasma Membrane-Associated Receptor-like Kinases Relocalize to
Plasmodesmata in Response to Osmotic Stress1[OPEN]. Plant Physiol *181*, 142–160.
https://doi.org/10.1104/pp.19.00473.

22. Vu, M.H., Hyun, T.K., Bahk, S., Jo, Y., Kumar, R., Thiruppathi, D., Iswanto, A.B.B., Chung, W.S.,
Shelake, R.M., and Kim, J.-Y. (2023). ROS-mediated plasmodesmal regulation requires a network of
an Arabidopsis receptor-like kinase, calmodulin-like proteins, and callose synthases. Front. Plant Sci.
13. https://doi.org/10.3389/fpls.2022.1107224.

23. Jaillais, Y., Bayer, E., Bergmann, D.C., Botella, M.A., Boutté, Y., Bozkurt, T.O., Caillaud, M.-C.,
Germain, V., Grossmann, G., Heilmann, I., et al. (2024). Guidelines for naming and studying plasma
membrane domains in plants. Nat. Plants *10*, 1172–1183. https://doi.org/10.1038/s41477-024-01742-
8.

24. Jarsch, I.K., Konrad, S.S.A., Stratil, T.F., Urbanus, S.L., Szymanski, W., Braun, P., Braun, K.-H., and
Ott, T. (2014). Plasma Membranes Are Subcompartmentalized into a Plethora of Coexisting and
Diverse Microdomains in Arabidopsis and Nicotiana benthamiana. Plant Cell *26*, 1698–1711.
https://doi.org/10.1105/tpc.114.124446.

25. Bücherl, C.A., Jarsch, I.K., Schudoma, C., Segonzac, C., Mbengue, M., Robatzek, S., MacLean, D.,
Ott, T., and Zipfel, C. (2017). Plant immune and growth receptors share common signalling
components but localise to distinct plasma membrane nanodomains. eLife *6*, e25114.
https://doi.org/10.7554/eLife.25114.

26. Jaillais, Y., and Ott, T. (2020). The Nanoscale Organization of the Plasma Membrane and Its
Importance in Signaling: A Proteolipid Perspective1[OPEN]. Plant Physiol *182*, 1682–1696.
https://doi.org/10.1104/pp.19.01349.

27. Hdedeh, O., Mercier, C., Poitout, A., Martinière, A., and Zelazny, E. (2025). Membrane nanodomains
to shape plant cellular functions and signaling. New Phytologist *245*, 1369–1385.
https://doi.org/10.1111/nph.20367.

28. Bücherl, C.A., van Esse, G.W., Kruis, A., Luchtenberg, J., Westphal, A.H., Aker, J., van Hoek, A.,
Albrecht, C., Borst, J.W., and de Vries, S.C. (2013). Visualization of BRI1 and BAK1(SERK3)
Membrane Receptor Heterooligomers during Brassinosteroid Signaling1[W][OPEN]. Plant Physiol
162, 1911–1925. https://doi.org/10.1104/pp.113.220152.

29. Chinchilla, D., Shan, L., He, P., de Vries, S., and Kemmerling, B. (2009). One for all: the receptor-
associated kinase BAK1. Trends Plant Sci *14*, 535–541. https://doi.org/10.1016/j.tplants.2009.08.002.

30. Perraki, A., Gronnier, J., Gouguet, P., Boudsocq, M., Deroubaix, A.-F., Simon, V., German-Retana,
S., Legrand, A., Habenstein, B., Zipfel, C., et al. (2018). REM1.3's phospho-status defines its plasma

membrane nanodomain organization and activity in restricting PVX cell-to-cell movement. PLOS
Pathogens *14*, e1007378. https://doi.org/10.1371/journal.ppat.1007378.

31. Xu, Z., Schahl, A., Jolivet, M.-D., Legrand, A., Grélard, A., Berbon, M., Morvan, E., Lagardere, L.,
Piquemal, J.-P., Loquet, A., et al. (2024). Dynamic pre-structuration of lipid nanodomain-segregating
remorin proteins. Commun Biol *7*, 1620. https://doi.org/10.1038/s42003-024-07330-y.

32. Wu, X.N., Chu, L., Xi, L., Pertl-Obermeyer, H., Li, Z., Sklodowski, K., Sanchez-Rodriguez, C.,
Obermeyer, G., and Schulze, W.X. (2019). Sucrose-induced Receptor Kinase 1 is Modulated by an
Interacting Kinase with Short Extracellular Domain*,. Molecular & Cellular Proteomics *18*, 1556–
1571. https://doi.org/10.1074/mcp.RA119.001336.

33. Wang, J., Xi, L., Wu, X.N., König, S., Rohr, L., Neumann, T., Weber, J., Harter, K., and Schulze,
W.X. (2022). PEP7 acts as a peptide ligand for the receptor kinase SIRK1 to regulate aquaporin-
mediated water influx and lateral root growth. Molecular Plant *15*, 1615–1631.
https://doi.org/10.1016/j.molp.2022.09.016.

34. Goto, Y., Kadota, Y., Mbengue, M., Lewis, J.D., Matsui, H., Maki, N., Ngou, B.P.M., Sklenar, J.,
Derbyshire, P., Shibata, A., et al. (2024). The leucine-rich repeat receptor kinase QSK1 regulates
PRR-RBOHD complexes targeted by the bacterial effector HopF2Pto. Plant Cell *36*, 4932–4951.
https://doi.org/10.1093/plcell/koae267.

35. Isner, J.C., Begum, A., Nuehse, T., Hetherington, A.M., and Maathuis, F.J.M. (2018). KIN7 Kinase
Regulates the Vacuolar TPK1 K+ Channel during Stomatal Closure. Current Biology *28*, 466-472.e4.
https://doi.org/10.1016/j.cub.2017.12.046.

36. Zhu, Z., Krall, L., Li, Z., Xi, L., Luo, H., Li, S., He, M., Yang, X., Zan, H., Gilbert, M., et al. (2024).
Transceptor NRT1.1 and receptor-kinase QSK1 complex controls PM H+-ATPase activity under low
nitrate. Current Biology *34*, 1479-1491.e6. https://doi.org/10.1016/j.cub.2024.02.066.

37. Bourdais, G., Burdiak, P., Gauthier, A., Nitsch, L., Salojärvi, J., Rayapuram, C., Idänheimo, N.,
Hunter, K., Kimura, S., Merilo, E., et al. (2015). Large-Scale Phenomics Identifies Primary and Fine-
Tuning Roles for CRKs in Responses Related to Oxidative Stress. PLOS Genetics *11*, e1005373.
https://doi.org/10.1371/journal.pgen.1005373.

38. Xu, B., Cheval, C., Laohavisit, A., Hocking, B., Chiasson, D., Olsson, T.S.G., Shirasu, K., Faulkner,
C., and Gilliham, M. (2017). A calmodulin-like protein regulates plasmodesmal closure during
bacterial immune responses. New Phytologist *215*, 77–84. https://doi.org/10.1111/nph.14599.

39. Huang, C., Mutterer, J., and Heinlein, M. (2022). In Vivo Aniline Blue Staining and Semiautomated
Quantification of Callose Deposition at Plasmodesmata. Methods Mol Biol *2457*, 151–165.
https://doi.org/10.1007/978-1-0716-2132-5_9.

40. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction with
AlphaFold. Nature *596*, 583–589. https://doi.org/10.1038/s41586-021-03819-2.

41. Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M. (2022).
ColabFold: making protein folding accessible to all. Nat Methods *19*, 679–682.
https://doi.org/10.1038/s41592-022-01488-1.

42. Hsu, J.-L., Wang, L.-Y., Wang, S.-Y., Lin, C.-H., Ho, K.-C., Shi, F.-K., and Chang, I.-F. (2009).
Functional phosphoproteomic profiling of phosphorylation sites in membrane fractions of salt-
stressed Arabidopsis thaliana. Proteome Sci *7*, 42. https://doi.org/10.1186/1477-5956-7-42.

43. Niittylä, T., Fuglsang, A.T., Palmgren, M.G., Frommer, W.B., and Schulze, W.X. (2007). Temporal
Analysis of Sucrose-induced Phosphorylation Changes in Plasma Membrane Proteins of Arabidopsis.
Molecular & Cellular Proteomics *6*, 1711–1726. https://doi.org/10.1074/mcp.M700164-MCP200.

44. Chang, I.-F., Hsu, J.-L., Hsu, P.-H., Sheng, W.-A., Lai, S.-J., Lee, C., Chen, C.-W., Hsu, J.-C., Wang,
S.-Y., Wang, L.-Y., et al. (2012). Comparative phosphoproteomic analysis of microsomal fractions of
Arabidopsis thaliana and *Oryza sativa* subjected to high salinity. Plant Science *185–186*, 131–142.
https://doi.org/10.1016/j.plantsci.2011.09.009.

45. Xue, L., Wang, P., Wang, L., Renzi, E., Radivojac, P., Tang, H., Arnold, R., Zhu, J.-K., and Tao, W.A.
(2013). Quantitative measurement of phosphoproteome response to osmotic stress in arabidopsis
based on Library-Assisted eXtracted Ion Chromatogram (LAXIC). Mol Cell Proteomics *12*, 2354–
2369. https://doi.org/10.1074/mcp.O113.027284.

46. Stahl, Y., and Simon, R. (2013). Gated communities: apoplastic and symplastic signals converge at
plasmodesmata to control cell fates. J Exp Bot *64*, 5237–5241. https://doi.org/10.1093/jxb/ert245.

47. Cheval, C., and Faulkner, C. (2018). Plasmodesmal regulation during plant–pathogen interactions.
New Phytologist *217*, 62–67. https://doi.org/10.1111/nph.14857.

48. Li, M., Wang, M., Lin, Q., Wang, M., Niu, X., Cheng, J., Xu, M., Qin, Y., Liao, X., and Wu, J.X. and
S. (2022). Symplastic communication in the root cap directs auxin distribution to modulate root
development. Journal of Integrative Plant Biology *64*, 859. https://doi.org/10.1111/jipb.13237.

49. Pérez-Sancho, J., Smokvarska, M., Dubois, G., Glavier, M., Sritharan, S., Moraes, T.S., Moreau, H.,
Dietrich, V., Platre, M.P., Paterlini, A., et al. (2025). Plasmodesmata act as unconventional membrane
contact sites regulating intercellular molecular exchange in plants. Cell *188*, 958-977.e23.
https://doi.org/10.1016/j.cell.2024.11.034.

50. Grison, M.S., Brocard, L., Fouillen, L., Nicolas, W., Wewer, V., Dörmann, P., Nacir, H., Benitez-
Alfonso, Y., Claverol, S., Germain, V., et al. (2015). Specific membrane lipid composition is
important for plasmodesmata function in Arabidopsis. The Plant Cell *27*, 1228–1250.
https://doi.org/10.1105/tpc.114.135731.

51. Wang, X., Sager, R., Cui, W., Zhang, C., Lu, H., and Lee, J.-Y. (2013). Salicylic Acid Regulates
Plasmodesmata Closure during Innate Immune Responses in Arabidopsis[C][W]. Plant Cell *25*,
2315–2329. https://doi.org/10.1105/tpc.113.110676.

52. German, L., Yeshvekar, R., and Benitez-Alfonso, Y. (2023). Callose metabolism and the regulation of
cell walls and plasmodesmata during plant mutualistic and pathogenic interactions. Plant Cell
Environ *46*, 391–404. https://doi.org/10.1111/pce.14510.

53. Li, J., Lu, P., Pan, Q., Wang, B., Wang, Y., and Li, J. (2025). ER-PM tether Syt1 limits cell-to-cell
connectivity via plasmodesmata during innate immune responses in Arabidopsis. Cell Reports *44*.
https://doi.org/10.1016/j.celrep.2025.115672.

54. Sivaguru, M., Fujiwara, T., Samaj, J., Baluska, F., Yang, Z., Osawa, H., Maeda, T., Mori, T.,
Volkmann, D., and Matsumoto, H. (2000). Aluminum-induced 1-->3-beta-D-glucan inhibits cell-to-
cell trafficking of molecules through plasmodesmata. A new mechanism of aluminum toxicity in
plants. Plant Physiol *124*, 991–1006. https://doi.org/10.1104/pp.124.3.991.

55. O'Lexy, R., Kasai, K., Clark, N., Fujiwara, T., Sozzani, R., and Gallagher, K.L. (2018). Exposure to
heavy metal stress triggers changes in plasmodesmatal permeability via deposition and breakdown of
callose. J Exp Bot *69*, 3715–3728. https://doi.org/10.1093/jxb/ery171.

56. Bilska, A., and Sowinski, P. (2010). Closure of plasmodesmata in maize (Zea mays) at low
temperature: a new mechanism for inhibition of photosynthesis. Ann Bot *106*, 675–686.
https://doi.org/10.1093/aob/mcq169.

57. Cui, W., and Lee, J.-Y. (2016). Arabidopsis callose synthases CalS1/8 regulate plasmodesmal
permeability during stress. Nat Plants *2*, 16034. https://doi.org/10.1038/nplants.2016.34.

58. Cao, Y., Liang, Y., Tanaka, K., Nguyen, C.T., Jedrzejczak, R.P., Joachimiak, A., and Stacey, G.
(2014). The kinase LYK5 is a major chitin receptor in Arabidopsis and forms a chitin-induced
complex with related kinase CERK1. eLife *3*, e03766. https://doi.org/10.7554/eLife.03766.

59. Chinchilla, D., Zipfel, C., Robatzek, S., Kemmerling, B., Nürnberger, T., Jones, J.D.G., Felix, G., and
Boller, T. (2007). A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant
defence. Nature *448*, 497–500. https://doi.org/10.1038/nature05999.

60. Mühlenbeck, H., Tsutsui, Y., Lemmon, M.A., Bender, K.W., and Zipfel, C. Allosteric activation of the
co-receptor BAK1 by the EFR receptor kinase initiates immune signaling. eLife *12*, RP92110.
https://doi.org/10.7554/eLife.92110.

61. Nam, K.H., and Li, J. (2002). BRI1/BAK1, a receptor kinase pair mediating brassinosteroid
signaling. Cell *110*, 203–212. https://doi.org/10.1016/s0092-8674(02)00814-0.

62. Kimura, S., Hunter, K., Vaahtera, L., Tran, H.C., Citterico, M., Vaattovaara, A., Rokka, A., Stolze,
S.C., Harzen, A., Meißner, L., et al. (2020). CRK2 and C-terminal Phosphorylation of NADPH
Oxidase RBOHD Regulate Reactive Oxygen Species Production in Arabidopsis[OPEN]. Plant Cell
32, 1063–1080. https://doi.org/10.1105/tpc.19.00525.

63. Colina, F.J., Krasensky-Wrzaczek, J., Pérez-Guillén, A., and Wrzaczek, M. (2025). CRK2 modulates
flowering in Arabidopsis together with GLYCINE-RICH RNA BINDING PROTEIN 7 (GRP7).
Preprint at bioRxiv, https://doi.org/10.1101/2025.08.05.668606
https://doi.org/10.1101/2025.08.05.668606.

64. Wang, Z., Ding, G., Geistlinger, L., Li, H., Liu, L., Zeng, R., Tateno, Y., and Li, Y. (2011). Evolution
of protein phosphorylation for distinct functional modules in vertebrate genomes. Mol Biol Evol *28*,
1131–1140. https://doi.org/10.1093/molbev/msq268.

65. Skinner, J.J., Wang, S., Lee, J., Ong, C., Sommese, R., Sivaramakrishnan, S., Koelmel, W.,
Hirschbeck, M., Schindelin, H., Kisker, C., et al. (2017). Conserved salt-bridge competition triggered
by phosphorylation regulates the protein interactome. Proceedings of the National Academy of
Sciences *114*, 13453–13458. https://doi.org/10.1073/pnas.1711543114.

66. Bah, A., Vernon, R.M., Siddiqui, Z., Krzeminski, M., Muhandiram, R., Zhao, C., Sonenberg, N., Kay,
L.E., and Forman-Kay, J.D. (2015). Folding of an intrinsically disordered protein by phosphorylation
as a regulatory switch. Nature *519*, 106–109. https://doi.org/10.1038/nature13999.

67. Szymanski, W.G., Zauber, H., Erban, A., Gorka, M., Wu, X.N., and Schulze, W.X. (2015).
Cytoskeletal Components Define Protein Location to Membrane Microdomains. Mol Cell Proteomics
14, 2493–2509. https://doi.org/10.1074/mcp.M114.046904.

68. Julkowska, M.M., Hoefsloot, H.C.J., Mol, S., Feron, R., de Boer, G.-J., Haring, M.A., and Testerink,
C. (2014). Capturing Arabidopsis root architecture dynamics with ROOT-FIT reveals diversity in
responses to salinity. Plant Physiol *166*, 1387–1402. https://doi.org/10.1104/pp.114.248963.

69. Perez-Riverol, Y., Bandla, C., Kundu, D.J., Kamatchinathan, S., Bai, J., Hewapathirana, S., John,
N.S., Prakash, A., Walzer, M., Wang, S., et al. (2025). The PRIDE database at 20 years: 2025 update.
Nucleic Acids Res *53*, D543–D553. https://doi.org/10.1093/nar/gkae1011.

70. Fisher, C.L., and Pei, G.K. (1997). Modification of a PCR-based site-directed mutagenesis method.
Biotechniques *23*, 570–571, 574. https://doi.org/10.2144/97234bm01.

71. Clough, S.J., and Bent, A.F. (1998). Floral dip: a simplified method for *Agrobacterium* -mediated
transformation of *Arabidopsis thaliana*. The Plant Journal *16*, 735–743.
https://doi.org/10.1046/j.1365-313x.1998.00343.x.

72. Grison, M.S., Petit, J.D., Glavier, M., and Bayer, E.M. (2020). Quantification of Protein Enrichment
at Plasmodesmata. Bio Protoc *10*, e3545. https://doi.org/10.21769/BioProtoc.3545.

73. Nakagami, H. (2014). StageTip-based HAMMOC, an efficient and inexpensive phosphopeptide
enrichment method for plant shotgun phosphoproteomics. Methods Mol Biol *1072*, 595–607.
https://doi.org/10.1007/978-1-62703-631-3_40.

74. Rappsilber, J., Ishihama, Y., and Mann, M. (2003). Stop and go extraction tips for matrix-assisted
laser desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in proteomics. Anal
Chem *75*, 663–670. https://doi.org/10.1021/ac026117i.

75. Cox, J., and Mann, M. (2008). MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol *26*, 1367–
1372. https://doi.org/10.1038/nbt.1511.

76. Tyanova, S., Temu, T., and Cox, J. (2016). The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. Nat Protoc *11*, 2301–2319.
https://doi.org/10.1038/nprot.2016.136.

77. Nolte, H., MacVicar, T.D., Tellkamp, F., and Krüger, M. (2018). Instant Clue: A Software Suite for
Interactive Data Visualization and Analysis. Sci Rep *8*, 12648. https://doi.org/10.1038/s41598-018-
31154-6.

Figure legends

Figure 1. CRK2 and QSK1 regulate plant growth and play antagonistic roles in osmotic

stress-induced callose deposition.

(A) Rosette of 2-week-old Arabidopsis plants grown in long-day conditions. Mutation of *QSK1* reverts small growth phenotype of *crk2* in the double mutant *crk2 qsk1*. Scale bar 1 cm. (B) Primary root (PR) length of 12-days-old plants. (C) Lateral root (LR) number per cm of PR (LR density). (D) Percentage of viable seed germination on media containing NaCl. The hypersensitive response of *crk2* mutant on salt-containing media was reverted in the *crk2 qsk1* double mutant. The germination efficiency was normalized according to the respective control plates. (E-F) Visualization and quantification of callose deposition in aniline blue-stained leaf epidermal cells of Arabidopsis. Roots of 2-week-old plants were subjected to 50 mM mannitol for 7 hours before imaging in the treated groups. Images acquired with a laser-scanning confocal microscope under the DAPI filter with emission wavelength 461 nm and laser wavelength 405 nm. Detection wavelength 430-530 nm; laser transmissivity 5.0 %; objective lens 60.0 X oil immersion, scale bar 10 µm. The boxes represent the interquartile range (25th–75th percentile), the horizontal line indicates the median, the whiskers extend to 1.5× the IQR, and dots represent individual data points. Panel B, n=34, Kuskal-Wallis test followed by Dunn's multiple comparison; Panel C, n=34, one-way ANOVA, Tukey's test; Panel D, n=3, one-way ANOVA, Tukey's test; Panel E, n≥22, two-way ANOVA, Tukey's multiple comparison test. Alphabets represent statistically significant differences. See also Fig. S1, Table S1 and Table S2.

Figure 2. Functional CRK2 is required for osmotic stress-induced enrichment of QSK1 at plasmodesmata.

(A) Relocalization QSK1-mCherry in Arabidopsis leaf epidermal cells. Detached leaves from plants expressing QSK1-mCherry in WT or *crk2* mutant backgrounds were treated with 0.4 M mannitol. The abaxial surface was visualized and imaged with a laser-scanning confocal microscope after 30 min of treatment. QSK1 failed to enrich in the *crk2* mutant background. (B) Boxplot showing the extent of PD enrichment of QSK1 which is signal intensity ratio between PD and PM ROI. (C) The disruption of QSK1 enrichment in the *crk2* mutant background was rescued by complementing QSK1-mCherry/*crk2* plants with CRK2-mVenus (QSK1-mCherry/*crk2* crossed with CRK2-mVenus/*crk2*). (D) The kinase inactive CRK2^{D450N} failed to complement this function in the cross of QSK1-mCherry/*crk2* and CRK2^{D450N}-mVenus/*crk2* plants. (E) The mutation in *QSK1* did not impact the relocalization of CRK2 in the CRK2-mVenus/*qsk1* plants. Images are acquired after 60 min of 0.4 M mannitol treatment. (F) Boxplot

showing enrichment index related to panel E. The images were captured with 60.0 X oil

immersion lens. Laser and emission wavelengths were 561 nm and 610 nm, respectively for

mCherry; 514 nm and 527 nm for mVenus. Detection wavelengths were 570-620 nm and 530-

630 nm for mCherry and mVenus, respectively. Laser transmissivity in the shown images is

7.0%-15% which was kept constant between the comparisons. Scanning of signals were done in

a sequential manner to avoid signal cross-contamination between the channels. Scale bar 10 µm.

At least 3 independent transgenic lines were tested for each genotype. Boxplots as described in

Figure 1. Panels B and F, n ≥ 15, two-way ANOVA, Tukey's test. See also Fig. S2 and Table S3.

Figure 3. QSK1 and CRK2 are present at the general plasma membrane in a pre-formed

complex and relocalize and enrich at plasmodesmata in a sequential manner during

osmotic stress.

(A) Arabidopsis leaf epidermal cells showing relocalization of QSK1 and CRK2. The images

shown are captured within a window of 0-60 min. QSK1 enrichment takes place first and reaches

maximum enrichment at approximately 30 min of treatment and subsequently starts to decline.

Objective lens 40.0 X oil immersion. Treatment and laser settings as described for Figure 2.

Scale bar 10 µm. (B) FRET-FLIM analysis of CRK2-mVenus (donor-only), CRK2-

mVenus+QSK1-mCherry (donor-acceptor) and CRK2^{D450N}-mVenus+QSK1-mCherry (donor-

acceptor negative control) plants with or without mannitol treatment. A lower mVenus lifetime in

the donor-acceptor pair CRK2-mVenus+QSK1-mCherry indicated the presence of a preformed

complex of CRK2 and QSK1. An increase in the mVenus lifetime in this pair after mannitol

treatment suggested dissociation of the complex. Boxplot as described in Figure 1. n ≥ 9. Two-

way ANOVA, Tukey's test. (C- D) AlphaFold2 prediction of interaction between CRK2 and

QSK1 extracellular and transmembrane domains (ExD+TMD) (C) and full-length proteins (D).

Contacts under 6 Å between the two proteins are shown. Lines are colored according to the PAE

values, with most contacts showing high confidence. See also Fig. S3, Table S4 and Table S5.

Figure 4. CRK2 phosphorylates QSK1.

In vitro kinase reaction between the recombinantly expressed cytosolic regions of CRK2 and

QSK1. (A) CRK2-targeted phosphosites in the QSK1 cytosolic region as identified by LC-ESI-

MS/MS analysis. The highlighted residues in pink are consolidated from 3 experiments-high

confidence only. The shown amino acid sequence is the cytosolic region of QSK1 and the grey

highlighted part corresponds to the pseudokinase domain. The arrows represent the two serine

residues at positions 621 and 626 which were selected for further studies. (B) *In vitro*

phosphorylation of QSK1cyto by CRK2cyto. The QSK1cyto-MBP and QSK1$^{S621A-S626A}$ cyto-

MBP were digested with Factor Xa protease for 24 h to separate the MBP tag for size resolution

before carrying out the kinase reaction. The QSK1 was phosphorylated by CRK2 but not with

the kinase inactive CRK2^{D450N}. No difference was observed in the phosphorylation of

QSK1$^{S621A-S626A}$. The lower blot shows the Coomassie Brilliant Blue (CBB)-stained gel which

was destained and developed for the detection of labeled ATP (upper blot). (C) *In vivo*

phosphorylation of QSK1. The shown phosphopeptide was detected in the WT and *crk2*

seedlings. NaCl treatment and mutation in *CRK2* reduced the intensity of phosphopeptide. The

detected phosphopeptide was predicted to be phosphorylated at two serine residues. Boxplot

description as Figure 1. n=3-4. Two-way ANOVA, Tukey's test. See also Table S6 for panel A

and Table S7 for panel C.

Figure 5. Phosphorylation of QSK1 at Ser-621 and Ser-626 inhibits its recruitment to PD

and modulates plant response to osmotic stress.

Relocalization of phospho-mutant versions of mCherry-tagged QSK1 in the leaf epidermal cells

of Arabidopsis WT or *crk2* mutant backgrounds. (A-B) The phospho-mimetic QSK1^{S-D}

(QSK1$^{(S621D-S626D)}$) showed reduced enrichment in both WT and *crk2* mutant backgrounds. (C-D)

The phospho-negative QSK1^{S-A} (QSK1$^{(S621A-S626A)}$) showed much higher enrichment in the WT

background upon mannitol treatment. A weak enrichment of phospho-negative QSK1 was also

observed in control conditions. The phospho-negative QSK1^{S-A} also enriched weakly in the *crk2*

mutant background after mannitol treatment. Objective lens 60.0 X oil immersion. Laser

transmissivity 7.0%-15%; constant between the comparisons. Experimental conditions and other

laser settings for mCherry as Fig. 2. Scale bar 10 μm. Boxplots as described in Figure 1. Two-

way ANOVA, Tukey's test. n ≥ 12. (E-F) Comparison of root response to salt stress. PR

elongation is the primary root growth post-treatment. 5-days-old seedlings were transferred to

media containing NaCl and were imaged after another 7 days. The PR elongation is calculated

respective to the control PR length. n=22-23, two-way ANOVA, Tukey's test (E). (F) LR density

upon NaCl treatment same as for panel E. n=22, Kruskal-Wallis test, Dunn's multiple

comparison test. QSK1, QSK1^{S-A} and QSK1^{S-D} are the plants expressing WT, QSK1$^{(S621A-S626A)}$

and QSK1$^{(S621D-S626D)}$ variant of QSK1, respectively, in the *qsk1* mutant background. (G-H)

Visualization and quantification of callose deposition in aniline blue-stained leaf epidermal cells

of Arabidopsis. Treatment and imaging as described for Figure 1. Boxplot as described in Figure

1. n = 30, two-way ANOVA, Tukey's test. See also Fig. S4; Table S3 for panels B, D, Table S8

for panels E-F and Table S9 for panels G-H.

Figure 6. Proposed model- CRK2 modulates plasmodesmal function of QSK1.

Under normal growth conditions, CRK2 phosphorylates QSK1 and the two RLKs form a

complex at the bulk plasma membrane (PM) inhibiting unnecessary recruitment of QSK1 to PD

(A). Exposure to salt or mannitol promotes an unknown protein phosphatase activity at QSK1

releasing it from the CRK2-QSK1 complex and consequently QSK1 relocalization to PD. CRK2

also phosphorylates another unknown protein(s), a necessary event for QSK1 enrichment at PD.

QSK1 positively regulates callose deposition at PD (B). Subsequently, QSK1 is removed from

PD and returns to basal level. CRK2 enrichment at PD follows (C).

Supplementary Figure legends

Figure S1 (related to Figure 1). CRK2 and QSK1 regulate plant growth and osmotic stress

response.

(A) Rosette area of 12-days-old plants. (B) Root phenotype of the mutants. (C-D) Quantification

of callose deposition in epidermal cells of aniline blue-stained leaves of plants treated with or

without mannitol. The boxes represent the interquartile range (25th–75th percentile), the

horizontal line indicates the median, the whiskers extend to 1.5× the IQR, and dots represent

individual data points. Panel A, n=59, one-way ANOVA, Tukey's test; Panel C-D, n ≥ 22, two-

way ANOVA, Tukey's multiple comparison test. See also Table S1 for panel A and Table S2 for

panels C-D.

Figure S2 (related to Figure 2). Enrichment of QSK1 at plasmodesmata requires CRK2.

(A) Relocalization QSK1-mCherry in Arabidopsis leaf epidermal cells. Detached leaves from

plants expressing QSK1-mCherry in WT and *crk2* backgrounds (QSK1-mCherry/WT, QSK1-

mCherry/*crk2*) were treated with 150 mM NaCl and imaged at 30- and 60-minute post-treatment.

Control samples were without salt treatment. In the QSK1-mCherry/WT plants, the punctate

pattern of mCherry signal was prominent at 30 min but faded after 60 min of treatment. In the
QSK1-mCherry/*crk2* plants, QSK1-mCherry did not enrich upon salt treatment. (B) Co-
localization of aniline blue-stained callose deposits with enriched QSK1-mCherry signal
showing that QSK1 enriched at PD upon osmotic stress treatment. Leaves were co-treated with
150 mM NaCl and 0.2 % aniline blue and abaxial surface was imaged with laser-scanning
confocal microscope after 30 min of treatment. Objective lens 40.0 X oil immersion, emission
wavelength for DAPI filter 461 nm and laser wavelength 405 nm. Detection wavelength 430-530
nm, laser transmittivity 2.0%, Scale bar 5 μm. Other laser settings for mCherry as Fig. 2.

Figure S3 (related to Figure 3). Spatio-temporal dynamics of QSK1 and CRK2 association
at plasma membrane and structural modeling prediction of their interaction.

(A) Arabidopsis leaf epidermal cells showing relocalization of QSK1 and CRK2 in the QSK1-
mCherry/CRK2-mVenus plants. The images shown are captured within a window of 0-60 min of
0.4 M mannitol treatment. QSK1 enrichment is followed by CRK2 enrichment. Objective lens
40.0 X oil immersion. Laser wavelengths were 561 nm and 514 nm for mCherry and mVenus,
respectively. Emission wavelengths were 610 nm and 527 nm for mCherry and mVenus,
respectively. Detection wavelengths were 570-620 nm and 530-630 nm for mCherry and
mVenus, respectively. Scanning of signals were done in a sequential manner to avoid signal
cross-contamination between the channels. Scale bar 10 μm. (B) AlphaFold2 prediction of
interaction between CRK2 and QSK1 segments- extracellular domains (ExD), transmembrane
domains (TMD), TMD + Kinase domains, and Kinase domains. Predicted Aligned Error (PAE)
plots show model's confidence. Lower PAE values indicate higher confidence of interaction.
Lines are colored according to the PAE values. The modeled boundaries and predicated scores in
Table S5.

Figure S4 (related to Figure 5). Phosphorylation of QSK1 at Ser-621 and Ser-626 inhibits
its recruitment to PD and alters plant response to osmotic stress.

Relocalization of QSK1-mCherry and QSK1 phospho-mutants in different genetic backgrounds
of Arabidopsis leaf epidermal cells. (A-B) Enrichment of QSK1 at PD is independent of Sucrose-
Induced Receptor Kinase 1 (SIRK1). Arabidopsis plants expressing QSK1-mCherry in *sirk1*
mutant background show normal recruitment of QSK1 at PD upon mannitol treatment. A basal
presence of QSK1 at PD was visible in control samples as well. (C-D) The phospho-mimetic

QSK1^{S-E} (QSK1$^{S621E-S626E}$) showed a weak enrichment at PD in WT background upon mannitol treatment. (E-F) Similar to WT background in Fig. 5C-D, the phospho-negative QSK1^{S-A} (QSK1$^{(S621A-S626A)}$) showed high enrichment in the *qsk1* background upon mannitol treatment. In panels A, C- laser transmissivity 7.0%, objective lens 60.0 X oil immersion; panel D- laser transmissivity 2.0%, objective lens 40.0 X oil immersion. Scale bar 10 um. 3 independent transgenic lines were tested. Panel B, n=18, unpaired test with Welch's correction; Panel D, n=18, two-way ANOVA, Tukey's test; Panel F, n=17-18, unpaired test with Welch's correction. (G-H) Quantification of callose deposition in epidermal cells of aniline blue-stained leaves of plants treated with or without mannitol. $n \geq 22$, two-way ANOVA, Tukey's multiple comparison test. (I-J) PR length and LR density of QSK1 phospho-mutants at serines 621 and 626. The phospho-mutation of QSK1 at serines 621 and 626 does not impact the root phenotype under normal growth conditions. Panels I-J, n=22, Kruskal-Wallis test, Dunn's multiple comparison test. QSK1, QSK1^{S-A} and QSK1^{S-D} are the plants expressing WT, QSK1$^{(S621A-S626A)}$ and QSK1$^{(S621D-S626D)}$ variant of QSK1, respectively, in the *qsk1* mutant background. See also Table S3 for panels A-F, Table S8 for G-H, and Table S9 for I-J. Boxplots as described in Fig. 1 and Fig. S1.

Supplementary Tables

Table S1. Phenotypic analysis and stress response-raw data (related to Fig. 1B-D; Fig. S1A)

Table S2. Callose quantification-raw data (related to Fig. 1F; Fig. S2C, S2D)

Table S3. Enrichment index-raw data (related to Fig. 2B, 2F, 5B, 5D; Fig. S4B, S4D, S4F)

Table S4. FRET-FLIM-raw data (related to Fig. 3B)

Table S5: Amino acid boundaries of full-length, ExD, ExD+TMD, TMD, TMD+Kinase domain, and Kinase domain CRK2 and QSK1 used for ColabFold predictions (related to Fig. 3C-D, Fig. S3B)

Table S6. Phosphopeptides identified in LC-MS/MS after *in vitro* kinase reaction between CRK2 and QSK1 cytosolic regions (related to Fig. 4A)

Table S7. Intensity of QSK1 phosphopeptide in phosphoproteomics (related to Fig. 4C)

Table S8. Phenotypic analysis and stress response-raw data (related to Fig. 5E-F; Fig. S4I-J)

Table S9. Callose quantification-raw data (related to Fig. 5H; Fig. S4G-H)

Table S10-Primers used in the study

A

0 min → Time after mannitol treatment → 60 min

mCherry | QSK1

CRK2 | mVenus

QSK1-mCherry/CRK2-mVenus

— QSK1 — CRK2

Enrichment

B

Donor lifetime (ns)

CRK2-mVenus
CRK2-mVenus + QSK1-mCherry
CRK2^{D450N}-mVenus + QSK1-mCherry

Control
Mannitol

C

CRK2 QSK1

ExD+TMD

90°

CRK2

QSK1

0   200   400

D

Full-length

ExD

TMD

90°

Kinase

Predicted Aligned Error (Å)

0   15   30

CRK2

QSK1

0   500   1000

A

CRKRKKEENVP S RNVEAPVAAA T SSAAIPKE T VVVVPPAK
ATGSE S GAVNKDLTFFVKSFGEFDLDGLLKA S AEVLGKGT
VGS S YKASFEHGLVVAVKRLRDVVVPEKEFRERLHVLGSM
SHANLVTLIAYYFSRDEKLLVFEYMSKGSLSAILHGNKGN
GRTPLNWETRAGIALGAARAISYLHSRDGTT S HGNIKSSN
ILLSDSYEAKVSDYGLAPIISSTSAPNRIDGYRAPEITDA
RKISQKADVYSFGVLILELLTGKSPTHQQLNEEGVDLPRW
VQSVTEQQTPSDVLDPEL T RYQPEGNENIIRLLKIGM S CT
AQFPDSRPSMAEVTRLIEEVSHSSGSPNPV S D

B

	MBP	-	-	-	-	+	+
QSK1^{S621A-S626A}cyto	-	-	+	+	-	-
QSK1cyto	+	+	-	-	-	-
CRK2^{D450N}cyto	-	+	-	+	-	+
CRK2cyto	+	-	+	-	+	-

CRK2cyto ▶ (yellow)
QSK1cyto ▶ (green)
autoradiogram

70 KDa
55 KDa
40 KDa
CBB

C

In vivo phosphorylated
QSK1 peptide

LIEEVSHSSGSPNPVSD

Control   NaCl

Intensity of phosphopeptide (a.u.)

a
b
ab
ab

WT   *crk2*

A

QSK1^{S-D}/WT | Control | Mannitol
QSK1^{S-D}/*crk2* | Control | Mannitol

B

Enrichment index

a   a   a   a

QSK1^{S-D}/WT   QSK1^{S-D}/*crk2*

Control   Mannitol

C

QSK1^{S-A}/WT | Control | Mannitol
QSK1^{S-A}/*crk2* | Control | Mannitol

D

Enrichment index

b   a   c   b

QSK1^{S-A}/WT   QSK1^{S-A}/*crk2*

Control   Mannitol

E

Primary root elongation (%)

100 mM NaCl

ab   b   a

QSK1/*qsk1*   QSK1^{S-A}/*qsk1*   QSK1^{S-D}/*qsk1*

F

No. of LR/cm of PR

100 mM NaCl

a   a   a

QSK1/*qsk1*   QSK1^{S-A}/*qsk1*   QSK1^{S-D}/*qsk1*

G

QSK1/*qsk1*   QSK1^{S-A}/*qsk1*   QSK1^{S-D}/*qsk1*

Control

Mannitol

H

Total callose deposition (a.u.)

b   a   ab   a   b   ab

QSK1/*qsk1*   QSK1^{S-A}/*qsk1*   QSK1^{S-D}/*qsk1*

Control   Mannitol

Normal growth conditions

NaCl/Mannitol

Time after stress treatment

| Unknown phosphorylation target(s) of CRK2 | Callose | Phosphorylation |

PP Protein Phosphatase
PD Plasmodesmata