# Time-dependent bistability leads to critical slowing down during floral transition in Arabidopsis

**G. Rodríguez-Maroto[1,2], K. Wang[1,2], P. Casanova-Ferrer[1], M. Cerise[1], G. Coupland[1,*], and P. Formosa-Jordan[1, *]**

[1] Department of Plant Developmental Biology, Max Planck Institute for Plant Breeding Research, Cologne, Germany
[2] These authors contributed equally
* Corresponding authors
Correspondence: pformosa@mpipz.mpg.de, coupland@mpipz.mpg.de

## Abstract

Developmental transitions occur in the life cycles of all multicellular organisms. Despite their fundamental relevance, the underlying dynamics remain poorly understood. In plants, floral transition is a key developmental process whereby the shoot apical meristem changes from producing leaves to flowers. Using quantitative imaging, developmental genetics and dynamical systems theory, we show that a time-dependent bistable switch between *APETALA2*, a key floral inhibitor, and the floral activators *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1* and *FRUITFULL*, can explain the dynamics of floral transition in Arabidopsis. Notably, we detect a slowing down of the inhibitor dynamics, consistent with having a system crossing a critical point of a bistable switch and transiently experiencing a ghost attractor. We demonstrate that only this time-dependent bistability can generate the range of dynamical behaviours measured across genotypes, including oscillations in the inhibitor. We propose that time-dependent bistability is a robust dynamical mechanism for regulating developmental transitions.

*Keywords*: developmental transitions, plant development, flowering, Arabidopsis, meristems, dynamical systems theory, criticality, quantitative image analysis.

# Introduction

Biological system progress through their developmental programmes through a succession of distinct states. These transitions between distinct states are fundamental across life, and include, for example, transitions at the cellular level, such as cell-fate decisions[1], transitions at the multicellular level, such as phase transitions in morphogenetic processes[2], and transitions at the organism level, such as the metamorphosis in amphibians[3]. Long-standing questions are to explain the dynamics and timing of these transitions, and how they can generate appropriate responses against changing stimuli. Conrad Waddington was a pioneer in investigating the theoretical basis of this question[4]. In his iconic illustration known as the Waddington landscape, the state of a developing system is represented by a ball rolling down an undulated surface. Each valley represents a distinct developmental route, and the system's trajectory through a valley represents the development of the system[4]. A key feature of Waddington's landscape is that it can be dynamic — genetic or environmental signals can alter its topography[4,5] (Figure 1A). This time-dependent nature is particularly relevant in the context of developmental transitions, where understanding the system's transient behaviour is essential. Furthermore, Waddington's landscape can be connected to bifurcation theory in dynamical systems, where it is defined as a potential or quasi-potential function derived from the underlying dynamical system equations[5–9].

In the context of the Waddington landscape, a developmental transition can be characterised by two main different dynamical scenarios (Figure 1A). One corresponds to a monostable transition, where a single valley contains a stable steady state (attractor) to which the system converges throughout development (Figure 1A, top row). In this scenario, the valley can shift its position and change shape over time, with the developmental system's change in identity being represented either by a continuous shift or a sharp change upon crossing a positional threshold (Figure 1A, top row). The second scenario involves the coexistence, at some point of development, of two valleys, each containing a stable steady state. This is known as bistability[6,8–11] and we refer to the switch from one stable state to the other as a bistable transition (Figure 1A, bottom row). Assuming the developmental trajectory is deterministic, i.e. the system trajectory does not undergo random fluctuations, this sharp and discontinuous transition occurs when the initial stable attractor disappears, forcing the system's trajectory out of the first valley, and eventually converging on the second stable attractor[8] (Figure 1A, bottom row). The change in the number or type of attractors (i.e. stable and unstable steady states) when the system is perturbed (i.e. changes in a model parameter) is known as a bifurcation[12,13] (saddle-node bifurcation, in the case above[12]) and the state at the bifurcation point itself is referred to as a critical state. Thus, a bistable transition would involve undergoing a saddle-node bifurcation over time, passing through criticality, which has important dynamic consequences. After the stable attractor disappears, its influence on the system's dynamics persists due to the flattened landscape surface, called a ghost attractor[14–17] (Figure 1A, third panel in lower row). This phenomenon, which results in a slowing down of the system's dynamics, is known as critical slowing down[14–17], and has been proposed to occur in several biological processes, from neural spiking[18,19], to embryonic development[20,21] and population dynamics[22].

Bistability requires a nonlinear positive feedback loop[8,10] and can confer robustness and hysteresis to the system[5,10,11,23,24]. These emerging properties make bistability a key dynamical

mechanism[25,11], particularly in contexts such as cell signalling[1,26,27], and differentiation[28], where it can explain how an undifferentiated cell progresses to one of the available differentiated cell fates during development[8,29]. Yet, despite its theoretical relevance in biology, proving the existence of bistability in a biological system remains a significant challenge, especially in contexts where it enables developmental transitions. Furthermore, there is still limited experimental evidence supporting critical slowing down and its relevance in biology. Although the gene regulatory networks (GRN) driving key developmental transitions are increasingly known, the underlying dynamical mechanisms that enable robust yet flexible transitions against genetic and environmental variations remain poorly understood.

The plant life cycle includes several developmental transitions, but only a few studies have investigated their underlying dynamic behaviour[30–33]. Floral transition is one of these crucial steps in plant development, marking the progression from vegetative growth to the reproductive stage, and is strongly influenced by endogenous and environmental cues that regulate its timing[34,35]. Only a precise spatiotemporal coordination between gene expression dynamics and morphological and physiological changes enables reproductive success and completion of seed development under favourable conditions[34]. In *Arabidopsis thaliana*, these simultaneous processes converge at the shoot apical meristem (SAM) (Figure 1B), which contains a population of stem cells whose progeny give rise to aerial organs[36] — leaves in the vegetative state and lateral branches and flowers in the inflorescence phase, also known as flowering phase. The SAM is also the tissue where relevant changes in GRN dynamics controlling floral transition occur[37–40]. During the vegetative phase, flowering is strongly repressed by APETALA2 (AP2) and APETALA2-LIKE (AP2-LIKE) transcription factors (TFs). These repressors inhibit the transcription of the floral activators *FRUITFULL* (*FUL*) and *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1* (*SOC1*), both encoding MADS box TFs, and *MIR172* genes, which encode microRNA miR172 that targets the mRNAs of *AP2* and *AP2-LIKE* genes[41,42] (Figure 1B, left panel). Exposure to inductive long-day (LD) conditions (16 h light and 8 h darkness) activates the photoperiodic pathway, which triggers *FLOWERING LOCUS T* (*FT*) transcription in leaf vasculature. FT protein is then transported to the SAM, where it binds to the b-ZIP transcription factor FD to form the florigen activation complex (FAC) and activates the transcription of genes such as *FUL* and *SOC1*[47]. The ageing pathway also contributes to the activation of these floral activator genes[44,45]. As plants age, expression of the developmental timer miR156 decreases[44,45]. This attenuates the inhibition of a set of *SQUAMOSA PROMOTER BINDING PROTEIN-LIKE* (*SPL*) genes, whose encoded TFs induce in turn the expression of *FUL*, *SOC1*, and *MIR172*. Under short-day (SD) conditions (8 h light and 16 h darkness), *SPL15* is particularly important, and floral transition occurs at a slower pace via the ageing pathway[44,46–48]. The combined effect of age and daylength signals thus modifies the network's dynamics as they induce the transcription of *FUL*, *SOC1* and *MIR172* (Figure 1B, middle panel), which in turn repress *AP2* and *AP2-LIKE* genes transcriptionally[49,50] and post-transcriptionally[51]. This mutual inhibition between floral activators and repressors defines an intermediate, transition phase. As floral transition progresses, the repression of floral inhibitors becomes more pronounced, thereby weakening the meristem's vegetative identity and leading to the initiation of floral primordia and the flowering phase.

In this study, we combine experimental data and theory to elucidate the dynamics underlying the SAM transition from a vegetative to an inflorescence state. Specifically, we focus on determining whether bistability is involved in this process, and its dynamical consequences.

We provide *in vivo* and *in silico* evidence that a time-dependent bistable switch, modelled as a toggle switch modulated by an age signal over time, underlies floral transition by explaining the temporal expression dynamics in wild type and mutants. We show that late-flowering phenotypes emerge naturally due to the influence of a ghost attractor, which explains long transients associated with disruptions to floral primordium development. Furthermore, we demonstrate that AP2 self-repression can cause fluctuations in its abundance at early stages. Collectively, our results suggest that floral transition at the SAM is driven by a time-dependent bistable switch mechanism, whereby different meristematic regions undergo distinct dynamic regimes.

# Results

## A time-dependent toggle-switch model for the floral transition

To investigate the floral transition mechanism at the SAM under the regulation of the photoperiodic and ageing pathways, we analysed the dynamics of the AP2 inhibitor and the SOC1 activator *in vivo*. We expanded previous reports[38,40,52,53] by reanalysing existing data[40] and by acquiring new data for AP2 expression through imaging the SAM of *AP2::AP2:VENUS* LD-grown plants from 7 to 24LDs (Figures 2A, S1, and S2A). To improve accuracy, our new computational pipeline accounts for meristem size by defining characteristic expression domains (see Methods and Figure S1). Consistently, both datasets showed a decrease in AP2 abundance over time until 17LDs, when SOC1 also reached its maximum (Figures 2, S2A, and S2D–S2F). From 17LDs onwards, the concentrations of both the activator and inhibitor stabilised (Figure 2C), and mature floral primordia began to appear on the meristem flanks (Figures 2A, 2B, S2B, and S2C).

The developmental trajectories of the SAM from the vegetative to the inflorescence state can be inferred from the evolution of AP2 and SOC1 concentration (Figure 2C). Here, we proposed a time-dependent toggle-switch model[25,5] to study the dynamics of the GRN underlying the floral transition and the effects of age and light signals (Figure 2D; see Methods). This single-compartment model consists of three ordinary differential equations that capture the mutual repression between floral activators (*SOC1*, *FUL; F* variable) and inhibitor (*AP2*; *A* variable) genes (see Methods). This double-negative feedback loop is influenced by a time-varying age signal, *S*, which is independent of *A* and *F*. *S* represents the age of the plant and serves as a proxy for the increasing SPL levels as miR156 declines with age[47]. In the model, *S* dynamics increases over time until saturation and mimics how SPLs activate *F* (*FUL* and *SOC1*) and *MIR172,* all of which lead to direct repression of *A* (*AP2*)[37,52]. The influence of light in the system was introduced for simplicity through the activator variable, distinguishing its contribution from that of the ageing pathway.

To analyse the behaviour of the model over time, we treated the age variable *S* as a parameter, *S\**, and studied the corresponding phase portraits and bifurcation diagrams, whose stable states correspond to the valleys of the Waddington landscape (see Methods). In parallel, we performed numerical simulations of the model dynamics over time and studied their correspondence with the bifurcation diagrams (Figures 2E–2J, S3, and S4; Methods).

Bifurcation analyses as a function of *S\** showed that a monostable transition with a single decreasing inhibitor stable branch and an increasing activator stable branch can take place

(Figure 2E). This behaviour was consistent with the corresponding simulations, showing a smoothly decreasing inhibitor and increasing activator (Figures 2F and S4A). For other parameter values, bistable transitions were observed: bistability exists within a range of age values, and eventually the system becomes monostable, undergoing a saddle-node bifurcation and transitioning discontinuously from a vegetative (high repressor, low activator) to an inflorescence (high activator, low repressor) state (Figure 2H). The strength of both light and age signals determines the appearance of bistability (Figures 2H and S4C–S4H). Numerical simulations in the bistable scenario also showed consistent antagonistic dynamics between the activator and repressor variables (Figures 2I, 2J, S3D–S3I, and S4B). The discontinuity of the transition in the bistable case observed in the bifurcation diagrams (Figure 2H, S3D, and S3G) was not always evident in the numerical simulations, and only appeared for certain model parameters (Figures 2I, 2J, S3F, and S3I). When present, it was accompanied by long transients where the temporal dynamics are slowed down (Figures 2J, S3I, S4B-S4H, S5A, and S5B).

In the bistable case, the system's dynamics continue to be transiently influenced by the initial state even after its disappearance via a ghost attractor[14–17], leading to a critical slowing down[15–17]. This phenomenon can cause regulatory variables to remain at a transient concentration plateau, whose duration depends on the distance ($\Delta S$) between the final steady state of the age variable ($S_{ss}$) and the saddle-node bifurcation ($S_{SN}$) (Figures S4C–S4H, S5A, and S5B). When critical slowing down is present, the duration of a transition follows a power law whose characteristic exponent is m = -½ (i.e., the duration of the transition follows $\tau \propto \Delta S^m$, with $\Delta S = S_{ss}\text{-}S_{SN}$; Figure S5C). This behaviour is consistent with systems at criticality close to a saddle-node bifurcation, whereby an additional temporal scale in their dynamics appears, which follows a similar power-law behaviour[14–17].

Model numerical simulations confirmed that higher AP2 production rates ($\alpha_1$) led to delayed floral transition (Figures S4G and S4H), consistent with late-flowering phenotypes observed in the inhibitor overexpression lines[40,52].

## AP2 protein dynamics at the SAM are consistent with an underlying bistable mechanism during floral transition

To determine whether the experimental data were consistent with a monostable or bistable transition, we closely analysed the reduction in AP2 abundance at the SAM over time (Figure 2C). In the vegetative meristem, AP2 declined steeply between 7 and 10 LDs, but the reduction rate became significantly slower between 10 and 14LDs (Figures 2C and S2D–S2F). At this point (14LDs), leaf primordium formation ceases at the SAM tip, SAM elongation continues, and cauline leaves are detectable (Figures 2A, 2B, S2B, and S2C), and as stated above, at 17LDs, floral primordia start to appear. We hypothesised that this fast-slow-fast decline could be a signature of time-dependent bistability in the system (Figure 2I), where the slow dynamics result from the critical slowing down, consistent with the flat Waddington landscape after the vegetative state becomes destabilised (Figure 1B). This introduces an intermediate state with prolonged AP2 expression levels before floral primordia appear.

Simulations across the parameter space revealed that these dynamics could only be obtained in the bistable transition scenario (Figures S5D–S5F, see Methods). Although a monostable

transition can generate progressively slower decay of the inhibitor, a subsequent re-acceleration was consistently absent (Figures S5D–S5F). Thus, the second rapid decay of the inhibitor is then most likely caused by the saddle-node bifurcation itself, specifically, when the system exits the ghost state and is attracted again by the inflorescence state. The discontinuous nature of this bifurcation would explain the measured second rapid reduction in the inhibitor (Figure 2C). Hence, a wild-type SAM is expected to be in a region of the parameter space where time-dependent bistability exists. As the age signal, $S$, increases, if age is treated as a parameter, the high inhibitor and low activator steady state concentrations smoothly decrease and increase, respectively (Figure 2K). Numerical simulation of system dynamics confirmed that the deceleration in AP2 reduction occurs exactly at the value predicted by the bifurcation analysis (Figures S3D, S3G, S5A, and S5B), and that AP2 is fully depleted only once the saddle-node is crossed, in conjunction with a completed floral transition.

For wt, numerical simulations qualitatively recapitulated the experimental data (Figures 2L, 2M, and S5G), showing an initial rapid reduction of AP2, followed by a transient, slower decreasing phase (10-14LDs Figure 2M), and subsequently by another strong reduction in AP2 expression (14-17LDs Figure 2M). Taken together, these data suggest that the slowdown and subsequent acceleration of AP2 reduction at the SAM constitute a distinct signature of bistability and a feature demonstrably absent in monostable transitions. This evidence provides compelling support for a bistable switch mechanism underlying floral transition.

## Single activator mutants exhibit delayed floral transition due to prolonged AP2 expression and the influence of the ghost attractor

To test the bistable nature of the floral transition, we studied how mutations in floral activator genes (*FUL* and *SOC1*), known to cause delayed flowering times[40,54,55], affect the critical slowing down dynamics associated with the bistable transition. From confocal images, we quantified AP2 abundance at the SAM of *ful* mutant plants grown under LDs (Figure 3A). In addition, we re-analysed previously published data on AP2 expression in *soc1* mutants[40] using our new computational pipeline (Figure S1). Confocal images of AP2 translational reporter lines in both *ful* and *soc1* demonstrated that AP2 expression at the SAM persists longer than in wt under LDs, consistent with their delayed flowering phenotypes (Figures 3A and 3B). In *ful,* the decrease in AP2 expression was very similar to wt and with only complete AP2 depletion slightly delayed compared to wt (19LDs in *ful* versus 17LDs in wt, Figure 3C), matching its slight late-flowering phenotype under LDs[54]. These results contrast with those of *soc1*, in which AP2 expression persists much longer than in wt (depletion of AP2 in *soc1* occurs at 24LDs versus 17LDs in wt; Figure 3C), in agreement also with the late-flowering phenotype of *soc1* compared with *ful* and wt under LDs[40]. Despite an initial similar decline in expression, the reduction in AP2 at the SAM in *soc1* ceases from 17LDs until 21LDs, and only at 24LDs is AP2 completely depleted from the SAM (Figure 3C). This prolonged expression of AP2 in *soc1* directly correlated with severely delayed floral primordia development, a robust proxy for floral transition completion, with visible floral primordia appearing only from 24LDs onwards (Figures 3B, S6A, and S6B). Such an extremely late formation of floral primordia was thus preceded by the quantified plateau phase in decline of AP2 expression (Figure 3C). These data collectively suggest that AP2 depletion at the SAM is necessary for proper floral primordium formation.

Numerical simulations of the *ful* and *soc1*, modelled by reduced expression rates in the activator variable, qualitatively recapitulated the AP2 decline shown in the experimental data (Figures 3C and 3D, and S6C–S6E). A stronger age signal (higher SPL levels) was required in the mutants to reach the saddle-node bifurcation and to perform the vegetative-to-inflorescence transition at the SAM (Figure 3E). This shift in the position of the saddle-node bifurcation amplifies the influence of the ghost attractor, naturally explaining the delayed floral transition observed in both mutants (more pronounced in *soc1*) and the longer plateau in the AP2 decline observed in *soc1* (17-21 LDs, Figure 3C), consistent with the critical slowing down predicted by the model. Prolonged transient dynamics, linked to impaired development of floral primordia in late-flowering mutants such as *soc1*, are naturally derived from the influence of the ghost attractor. Taken together, these results support the hypothesis of floral transition proceeding via a bistable-switch mechanism.

## *soc1 ful* double activator mutants and plants carrying a version of *AP2* resistant to miR172 display fluctuations in AP2 levels in the vegetative meristem

To further address our hypothesis, the dynamics of AP2 expression were assessed under LDs in two extreme genotypes: the *soc1 ful* double mutant and an *AP2::rAP2:VENUS ap2-12* transgenic line that expresses a version of AP2 resistant to miR172 (*rAP2*) (Figures 4F–4I). In *soc1 ful*, AP2 depletion at the SAM was even more delayed compared with that in single mutants (AP2 complete depletion only occurred at 26LDs in *soc1 ful*; Figures 3F, 3H, S6H, and S6I), consistent with its extremely late-flowering phenotype[55,56]. Similarly to *soc1*, *soc1 ful* showed a long plateau phase in AP2 reduction between 17 and 24LDs (Figures 3F, 3H, S6H, and S6I), which was also associated with a severe impairment of floral primordium initiation (Figures 3F, S6F, and S6G). AP2 was depleted from the SAM at 26LDs (Figures 3F, 3H, and S6H–S6J), and floral primordia were only observed at 29LDs (Figures S6F, S6G, and S6J). This extended plateau in AP2 expression before complete depletion of the inhibitor was reflected in the numerical simulations of *soc1 ful* (Figure 3J). In this genotype, the bistable region expands towards increased age-signal values, which reduces the gap between the saddle-node point and the final age value, thereby amplifying the influence of the ghost attractor (Figures S7A–S7F). The double mutant and *soc1* single mutant show a clear increase in the time the system is trapped by the ghost attractor ($\tau_g$) compared with the time needed to reach the saddle-node bifurcation ($\tau_0$) in wt or *ful* (Figures S7C–S7F), highlighting the emergence of a time scale in the system through the critical slowing down phenomenon. These results strengthen the hypothesis of time-dependent bistability in the system and highlight the key role of the ghost attractor in explaining the long transient dynamics in *soc1 ful*.

In *rAP2*, AP2 expression at the SAM effectively declined at 21-24LDs, but persisted even when mature floral primordia were clearly visible (Figures 3G, 3H, S6F, and S6G). Notably, rAP2 maintained comparatively high AP2 abundance even at later stages of the floral transition (17-19LDs in Figures 3H, S6H, and S6I). To further understand the rAP2 dynamics, we compared rAP2 expression levels across different SAM domains, such as the CLV3-expressing central zone (CZ) and the peripheral zone (PZ), where new primordia are initiated. We found that rAP2 signal intensity at the adjacent region of newly formed primordia was consistently lower than that at the central region (Figures 3G, S8A–S8C). These preliminary observations

suggest that the PZ region undergoes the strongest reduction in AP2 levels during the floral transition, potentially facilitating floral primordium initiation despite high overall AP2 expression in the CZ. Such a spatial gradient suggests a distinct mode of action of AP2 in different regions of the SAM.

Both *soc1 ful* and *rAP2* genotypes, however, exhibited a feature in their AP2 dynamics during early vegetative stages (7–12LDs) that was very different from that in wt and single mutants (Figures 3C and 3H). In *rAP2* and *soc1 ful*, AP2 protein levels increased significantly between 10 and 12 LDs, and only decreased again at 12LDs (Figure 3H). To confirm that this fluctuating behaviour between 10 and 14 LDs was not due to an artifact of the quantification pipeline or an uncontrolled event, we quantified single-cell nuclear AP2 concentration in SAMs from both genotypes during floral transition (Figures 3I, S8D–S8F, S9, and S10). Single-cell quantification confirmed, first, the overall increase in AP2 expression between 10 and 12 LDs in *soc1 ful* and *rAP2* genotypes (Figures 3I, S9D, S9E, S10D, and S10E) and, second, the absence of fluctuations in AP2 levels at early vegetative stages in wt (Figures S9A and S10A) and single activator mutants (Figures S9B, S9C, S10B, and S10C). However, these initial fluctuations in AP2 dynamics could not be replicated in *rAP2* or *soc1 ful* numerical simulations (Figures 3J and S7B). This discrepancy raised an obvious question: how can this fluctuating behaviour be theoretically generated, and which underlying biological interactions could explain it?

## A negative feedback loop in AP2 resolves the fluctuating behaviour and still recapitulates wt dynamics

A negative feedback loop with delay is sufficient to generate oscillations in a system[6,12,57,58]. Based on this principle, we hypothesised that an autoinhibition in the repressor variable could explain the fluctuating behaviour in AP2 expression dynamics observed in *soc1 ful* and *rAP2* (Figure 4A). Indeed, AP2 protein can bind to its own DNA-coding sequence[59,60], and some studies have proposed its self-repression[41,50,59,60]. However, the physiological relevance of this regulation remains unknown. We propose that the influence of this autoinhibition on the system's dynamics is attenuated in wt under LDs due to the strong repression of AP2 by both miR172 and FUL/SOC1, and therefore becomes relevant only in deregulated contexts in which AP2 dynamics are severely impaired, such as in *soc1 ful* or *rAP2* backgrounds.

The delay in AP2 self-repression was implemented into the model using three variables mediating the feedback: *AP2* mRNA (*Am*), an intermediate AP2 protein state (*Ac*), and the final AP2 transcription factor (*Ap*), which represses its own transcription and SOC1-FUL variable (*Fp*) (Figure 4A; see Methods). As a result of this autoinhibition in the repressor, oscillations emerged in the system for certain values of the age signal, both in a monostable and a bistable scenario (Figures S11A–S11G) due to the appearance of a stable limit cycle[12,13], which is an attractor that presents sustained oscillations. In the bistable scenario, the stable limit cycle can coexist with the stable inflorescence attractor (Figures 4B, S11A–S11C, S11F, and S11G). As age increases, the limit cycle eventually becomes a non-oscillatory vegetative attractor through a Hopf bifurcation, followed by a saddle-node bifurcation (Figure 4B). The amplitude of these oscillations, determined by the amplitude of the limit cycle, is significantly higher in the repressor variables (*Am*, *Ac* and *Ap*) than in the

activator variable (Figure 4B), as determined by the initial low concentration of this variable in a vegetative SAM.

Can oscillations function as an alternative mechanism to recapitulate the experimentally observed AP2-wt dynamics? To answer this question, we first examined with numerical simulations whether monostable transitions having a limit cycle –with or without a Hopf bifurcation– could recover the slowing down and re-acceleration of the AP2 decay. Our simulations in those cases showed either sustained or damped oscillations (Figures S11D, S11E, S11H, and S11I). In a monostable scenario with a Hopf bifurcation, damped oscillations turned into a non-oscillatory behaviour after a short period of time for very high age signal steady state values (which led to a very rapid age increase; $S_{ss}$ = 2000 in Figures S11H and S11I). Conversely, oscillatory behaviour could easily be prevented in the presence of bistability, as the age signal steady state increased (Figures S11J and S11K). Noticeably, although a monostable transition with a Hopf bifurcation could emulate the wt AP2 dynamics for high age steady state values (Figure S11H), this scenario failed to recapitulate the dynamics of the activator mutants that were experimentally measured, such as their late flowering phenotypes, the increase in length of AP2 stagnation or the fluctuations in the inhibitor in *soc1 ful* (Figures S12A and S12B). This lack of flexibility in the monostable scenario contrasts with the variety of dynamic behaviours yielded by a bistable mechanism (Figures S11J and S11K).

Consequently, the wt-genotype in this extended model remains within a bistable transition parametric region (Figure 4B). Only this transition simultaneously recovers its fast–slow–fast dynamics and enables the possibility to generate oscillations detected in other genotypes for other parameter values (Figures S11J and S11K). Numerical simulations showed opposite non-oscillatory temporal dynamics in both the activator (*Fp*) and the active form of the inhibitor (*Ap*) and recapitulated the wt behaviour (Figure 4C and 4D). Consistent with the experiments, the effect of AP2 self-repression in wt was attenuated. The slowing down observed in the decay of the inhibitor *Ap* continues to reflect the influence of the ghost attractor ($Ap \gtrsim 0.5$ in Figure 4C). Overall, the extended model shows a good qualitative agreement between theory and experiments for wt (Figure 4D; differences from 19LDs onwards are attributed to the background AP2 signal in the experiments). These results also demonstrate the context-dependent relevance of AP2 autoinhibition, key in *soc1 ful* and *rAP2* and minimised in wt and single mutants, highlighting the role of bistability in providing robustness necessary to suppress oscillations during the transition in wt.

## AP2 dynamics during floral transition in different genotypes can be explained by the extended toggle-switch model

The dynamics of AP2 expression in different mutant genotypes were assessed analytically and numerically with the extended model, using the previously modelled wt-genotype as reference (see Methods and Figure 5). Figure 5A shows the position of each genotype as a function of the inhibitor (*Am*) and activator (*Fp*) basal production rates ($\alpha_1$ and $\alpha_0$, respectively). For each pair of parameters ($\alpha_0$, $\alpha_1$), we computed the dynamical regimes crossed as the age signal variable increases to the steady state reached in the numerical simulations. The resulting diagram is coloured to distinguish regions by crossed regimes (B: bistable, M: monostable), their order, and the presence or absence of a limit cycle (oscillations). The reduction of the activator expression rate to emulate floral activator mutants shifts the system

in the parameter space towards the M-B-M region (Figure 5A), moving the Hopf and saddle-node bifurcation towards higher age-signal values (Figure 5B). This shift has two consequences in the temporal dynamics: it expands the range of the stable limit cycle and extends the upper branch of the saddle-node after the Hopf bifurcation, increasing the likelihood of observing oscillations and the influence of the ghost attractor (Figures 5B and S12C–S12G).

Numerical simulations consistently reproduced the delayed AP2 reduction and late-flowering phenotypes of the single mutants (Figures 5C–5E; the inflorescence SAMs appear at 17LDs, 17-19LDs, and 24LDs for wt, *ful* and *soc1,* respectively). The simulated *ful* mutant exhibited a more pronounced plateau during AP2 reduction (Figures 5B and 5D), which was further accentuated in the *soc1* mutant (Figures 5B and 5E).

In the *soc1 ful* double mutant, these features were amplified, resulting in a very high age-signal requirement to reach the saddle-node bifurcation and a remarkably long phase with oscillations (Figures 5B, 5F, and S12E). Numerical simulations of the double mutant effectively captured both the fluctuating behaviour of AP2 decline in a vegetative SAM and its extreme late-flowering phenotype (Figures 5B, 5F, and S12E). These dynamics were also recapitulated in the simulated *rAP2* genotype, in which an initial oscillation in AP2 levels is followed by a plateau at very high inhibitor values (Figure 5G). Such a plateau at high AP2 expression in *rAP2* is possible due to the absence of the repression of this inhibitor by the age signal (see Methods and Figure 5G).

These results demonstrate that the time-dependent bistable switch model with autoinhibition can better recapitulate the different behaviours across all studied genotypes (Figures 5 and S12F-S12K). The absence of different dynamic behaviours across the numerically simulated genotypes in the monostable scenario highlights the fundamental role of bistability in floral transition. Bistability ensures flexibility of the system upon perturbations via the expanding range of possible dynamical behaviours. It is this bistability that allows the new model to simultaneously account for the stagnation and late flowering phenotypes in wt and single floral activator mutants, and also the oscillatory regime in *soc1 ful* and *rAP2* genotypes, underscoring the relevance of AP2 autoinhibition in contexts in which this inhibitor is severely deregulated.

## Spatial modelling reveals distinct dynamic regimes along the SAM

AP2 and SOC1 expression patterns also exhibit spatial heterogeneity within the SAM during floral transition (Figure 2). To accurately capture and analyse this phenomenon, we quantified their spatial patterns of expression throughout the transition (Figures 6A, S13, and S14A; see Methods). At early stages (7-12 LDs), AP2 protein was strongly expressed near the apex (Figures 2A, 6A, and S13A). By contrast, SOC1 expression initiated at a lower SAM position (10LDs), and progressively expanded towards the apex (17 and 19LDs; Figures 2B, 6A, and S13B). Notably, a double peak of expression along the longitudinal axis of the SAM was observed in the SOC1 spatial expression profile at 14LDs (Figures 6A and S13E). AP2 spatial profiles were also quantified in *ful*, *soc1*, *soc1 ful* and *rAP2* genotypes (Figures S13C and S13D). A longer persistence of the inhibitor was observed in all of them within the meristem (Figures S13C and S13D), as well as initial variations in AP2 levels in the meristem in *soc1 ful* and *rAP2* (Figure S13D), as previously described when analysing the global temporal

dynamics of the SAM (Figure 3). We therefore investigated whether this observed spatiotemporal heterogeneity in AP2 and SOC1 wt-expression patterns indicates distinct meristematic regions undergoing different dynamic regimes. To test this, we extended the previous single-compartment model into a 1D spatially extended and continuous model representing the longitudinal axis of the SAM (Figure 6).

The model equations incorporated a diffusion term for the age signal S (see Methods), which is biologically supported by the movement of small microRNAs, such as miR172, through plasmodesmata[37,61]. Accordingly, deeper SAM meristematic regions were initialised with higher age signal levels (Figure S14B), and age signal production was assumed to occur only upon reaching a specific age signal concentration threshold. Although the SAM continuously grows during floral transition, explicit growth or cell division was not included for simplicity. To replicate the attenuation of reporter signals at distant regions from the SAM apex, the expression rates of the activator and inhibitor ($Ap$ and $Fp$) were progressively reduced as a function of the distance to the apex (see Methods).

Numerical simulations of the spatial model demonstrated strong qualitative agreement with experimental data. The activator profile emerged close to the apex, subsequently displaying two distinct peaks, recapitulating the double peak found in SOC1 expression (10–12LDs and t=20 in Figures 6A and 6B). Only at later time points was the activator expression profile flattened near the apex, as measured in experiments (Figure 6B). The double peak in the activator is consistent with a coherent feedforward loop motif[62], in which the age-signal activates directly and indirectly (by repressing AP2) the activator, and the differing strengths of these two activation mechanisms in space generate the two peaks along the SAM longitudinal axis. Theoretical analysis of the spatial model revealed that a non-homogeneous spatiotemporal distribution of the age signal confers distinct dynamic properties to different meristematic regions (Figures 6C–6E and S14B). Regions near the apex initially exhibit transient oscillatory behaviour (section 1 in Figures 6C–6E and S14). As the bifurcation age parameter, $S^*$, increases over time, this transient oscillatory phase terminates via a Hopf bifurcation (Figures S14C and S14D). Sustained oscillations would only occur if the final variable value of $S$ in the simulations remained within the oscillatory region of the bifurcation diagram. Figure 6E summarises these distinct scenarios comprehensively. For regions farther from the apex, higher age signal levels reduce the likelihood of observing oscillations. This causes the system to pass earlier through the non-oscillatory branch with high AP2 levels after the Hopf bifurcation, and to undergo the saddle-node bifurcation earlier (section 2 in Figures 6C–6E, S14C, and S14D). At even greater distances from the apex, the initial age signal concentration positions the system within a monostable region (sections 3 and 4 in Figures 6C–6E, and S14).

This spatial version of the toggle-switch model with AP2 self-inhibition exhibits qualitative behaviours that are highly consistent with experimental observations, highlighting the robustness and flexibility of the model. Importantly, these results suggest that different meristematic regions can undergo distinct dynamic regimes during floral transition, a consequence of time-dependent bistability influenced by spatial gradients of the age signal.

# Discussion

The question of how biological systems transition between distinct developmental states has long intrigued scientists since Waddington's seminal work. Although this question is often studied in the context of cell signalling and differentiation, it has remained, with a few exceptions[30–33], largely unexplored in plant development. Here, we provide evidence for a time-dependent bistable mechanism underlying the floral transition, whereby the SAM changes between vegetative and inflorescence states. To achieve this, we comprehensively quantified and analysed the spatiotemporal dynamics of key regulatory elements within the GRN that regulates floral transition at the SAM under both ageing and photoperiodic pathways.

Our analysis of AP2 and SOC1 temporal dynamics consistently supports a time-dependent bistable mechanism underlying floral transition. A toggle-switch model modulated over time by an age signal revealed that the characteristic slowing-down of AP2 reduction measured in wt is the theoretical signature of a bistable transition (Figure 2). This deceleration is caused by the influence of a ghost attractor, which introduces a temporal scale in the transition[14,15,17,21,63]. This phenomenon directly accounts for the prolonged AP2 expression quantified in floral activator mutants (*ful*, *soc1,* and *soc1 ful*). The remarkable correlation between these long transient plateaus in AP2 reduction and the severe impairments in floral primordium development makes the morphological implications of this second timescale visible (Figures 3A–3E). Furthermore, the incorporation of AP2 self-repression in the extended model, supported by previous studies[41,50,59,60], could recapitulate the fluctuations in AP2 levels at early stages in *soc1 fu*l and *rAP2* genotypes (Figures 3H–3J, 5F, 5G, S9, and S10). The model demonstrates that a bistable transition is necessary to ensure the broad range of distinct system's dynamical behaviours and flexibility upon genetic perturbation, a property not easily achieved with a monostable transition. The broad range of behaviors is also due to the coexistence between a non-oscillatory and an oscillatory attractor[64], a kind of bistability recently proposed in animal development modelling[58]. But importantly, the critical slowing down regime itself confers dynamic flexibility to prolong the transition, which could be used in certain environmental conditions.

For environmentally sensitive systems such as plants, the mechanism of a developmental transition can determine susceptibility or robustness to environmental perturbations. A bistable switch-like floral transition could thus explain how the SAM maintains its commitment to the inflorescence state, and does not return to a vegetative state, even under suboptimal light conditions. This robustness against noisy signals is a hallmark of bistability[11,65–68] that has been investigated in other developmental processes such as seed germination in plants[68,69], and in the patterning of neuronal subtype specification at the vertebrate neural tube[65]. Hysteresis is another property linked to bistable systems, whereby a system's response depends on its previous behaviour, conferring a "memory effect"[11,65]. It occurs, for instance, when a system remains committed to a certain state upon surpassing an activation threshold (i.e., the SAM retains commitment to flower) and only returns to the initial state (i.e., vegetative state) if the stimulus decreases below a different, lower deactivation threshold. Examples of reversion of floral transition[55,70,71] and the inflorescence state commitment in a double shift LD-SD-LD experiment[72] suggest the presence of hysteresis, further supporting the existence of a bistable mechanism. Noticeably, our modelled *soc1 ful* double mutant has an additional saddle-node bifurcation at relatively high age levels that makes the bistability region

constrained to a small signal age range, facilitating the reversion to a vegetative state upon a decrease in the age signal (Figure 4G). This feature again connects this time-dependent bistability to the robustness of the transition. A quantitative analysis of the floral transition GRN under perturbations is yet required to demonstrate the hysteretic behaviour of the system.

Previous models of animal developing tissues have proposed that gene regulatory networks and the underlying dynamics can be modulated over the spatial dimension[63,65,66,73]. Our work provides *in silico* and *in vivo* evidence for the relevance of the spatial component in the system during the transition, showing that different meristematic regions can be subjected to distinct dynamic regimes. Detailed image analysis in the *rAP2* genotype, which maintains high AP2 levels while producing floral primordia, revealed persistently lower AP2 expression at the boundary regions of newly formed primordia (Figures S8A–S8C). This suggests a distinct, localised mode of AP2 action across SAM regions, explaining how floral primordia development can proceed despite high inhibitor levels at SAM central regions. Extending the model into the spatial dimension unveiled a rich variety of dynamical behaviours simultaneously occurring at different meristematic positions (Figures 6 and S14). Therefore, the spatial dimension offers an additional route to the dynamical system for being poised at criticality[20], given that different regions of the meristem undergo different age signal trajectories. Future advances in live-imaging protocols will enable a more comprehensive assessment of the spatial dimension of the vegetative-to-inflorescence switch at the SAM.

While this study offers significant mechanistic insights into the transition, the analysis necessarily involves certain assumptions and simplifications. These include grouping SOC1 and FUL into a single activator entity despite differences in their activation timing[38,40,51,74], and omitting the possible influence of *AP2-LIKE* genes[39,50,60]. The addition of other factors in the model, such as the incorporation of an individual variable for miR172 or using a wave-like light signal to capture photoperiodic periodicity, could also increase the model's accuracy. However, the primary goal of this study was not to perfectly fit experimental data, but rather to understand the fundamental nature of the floral transition dynamics and explain the range and likelihood of its possible behaviours. In the future, it would be relevant to explore how the time-dependent bistable switch is modulated in other environmental conditions and genetic perturbations.

Our work highlights the importance of analysing time-dependent and transient behaviours in developmental contexts[5,58], which can reveal relevant behaviours in plants and in other biological systems that are entirely missed in traditional steady-state analyses[75]. A time-dependent bistable switch introduces a new timescale due to the emergence of a critical slowing down close to the bifurcation point. This critical behaviour confers dynamical flexibility and can be used to prolong developmental transitions upon genetic and environmental perturbations. Furthermore, this slowing down drives the emergence of an intermediate, transient state, which might be associated with a new developmental state. Simultaneously, time-dependent bistability serves as a dynamical motif that makes the committed new developmental state robust against endogenous and external fluctuations. In the future, it will be important to develop new combined theoretical and experimental approaches to address how the underlying dynamics of developing tissues can be modulated simultaneously in both the temporal and spatial dimensions.

# Methods

## Model equations

The first two formulated models are non-spatial, single-compartment phenomenological models that describe the rate of change of protein (and RNA in model 3) concentrations of AP2 (*A*) and SOC1/FUL (*F*), using ordinary differential equations. The variable *S* emulates the age-related developmental signal itself as a proxy of the developmental age of the plant. In biological terms, it could be interpreted to represent the levels of expression of SPL (SPL15), and, to some extent, also to MIR172. Their activations occur as a consequence of the weakening of their repression by MIR156. This microRNA acts as a timer in the system because its expression level decreases as the plant ages. The age signal is thus modelled through a simple basal production and a degradation proportional to its concentration. In all models, the rate of protein synthesis is assumed to be proportional to the corresponding activity of the promoter. Only pairwise interactions are considered and all proteins are assumed to be linearly degraded for simplicity. The following subsections describe the specific characteristics of each model. For the three models presented here, we performed a minimal qualitative parameter exploration searching for dynamical behaviours whose temporal dynamics were similar to those quantified in experiments. In the case of the model with AP2-autoinhibition, we required the system to display oscillatory behaviour (sustained oscillations) in the absence of age and activator variables. To achieve that the Hill exponent of AP2-autoinhibition must be greater than eight (see Model 2 for more details). All parameter values can be found in See Suppl. Tables 1 and 2.

### Model 1: Toggle-switch model

In this case, AP2 and SOC1/FUL proteins mutually repress each other through Hill functions. They are additionally inhibited and activated by the age signal, *S*, respectively. The quasi-steady assumption for mRNA has been implemented (i.e., mRNA dynamics are faster than protein dynamics). The dynamic equations over time (*t*) for AP2 (*A*), SOC1/FUL (*F*) and the age-signal (*S*) concentrations can be written as:

$$\frac{dA}{dt} = \frac{\alpha_1}{1 + \left(\frac{F}{k_{AF}}\right)^n + \left(\frac{S}{k_{AS}}\right)^n} - d_1 A \tag{1.1}$$

$$\frac{dF}{dt} = \alpha_0 \left( \frac{1}{1 + \left(\frac{A}{k_{FA}}\right)^n} + \frac{\alpha_4 \left(\frac{S}{k_{FS}}\right)^n}{1 + \left(\frac{S}{k_{FS}}\right)^n + \left(\frac{A}{k_{FA}}\right)^n} \right) - d_4 F \tag{1.2}$$

$$\frac{dS}{dt} = \beta_s - d_s S \, , \tag{1.3}$$

where $\alpha_1, \alpha_0$ and $\beta_s$ are the basal production rates of AP2, SOC1/FUL and age; $d_x$ represents the degradation rate constants for each variable *X*; $k_{ZY}$ are the concentrations of *Y* inhibiting or activating *Z* at which the production is half the saturated value; and exponent *n* sets the cooperativity ($n = 2$). The *F* production term – i.e., the first term within the parentheses – is formed by two parts to decouple the contributions of the photoperiodic and ageing pathways, whereby $\alpha_4$ is the relative strength between the ageing pathway with respect to the

photoperiodic pathway. Unless otherwise stated, we consider $\alpha_4<1$, implying that the photoperiodic pathway has a stronger contribution. In the results shown in Figures S4C–S4D, we chose another parametrisation of Equation 1.2, namely,

$$\frac{dF}{dt} = \alpha_0 \cdot \left( \frac{\alpha_L}{1 + \left(\frac{A}{k_{FA}}\right)^n} + \frac{\alpha_S \left(\frac{S}{k_{FS}}\right)^n}{1 + \left(\frac{S}{k_{FS}}\right)^n + \left(\frac{A}{k_{FA}}\right)^n} \right) - d_4 F \tag{1.5}$$

where $\alpha_L$ and $\alpha_S$ represent the strength of the photoperiodic pathway and ageing pathways, respectively. Nullclines in Figures 3E (right), S4A and S4B were obtained by setting equations 1.1 and 1.2 to zero (d$A$/dt = d$F$/dt= 0), and simultaneously treating S as a parameter (S*) as follows:

$$A^N(F) = \frac{\alpha_1}{d_1} \cdot \frac{1}{1 + \left(\frac{F}{k_{AF}}\right)^n + \left(\frac{S^*}{k_{AS}}\right)^n} \tag{1.6}$$

$$F^N(A) = \frac{\alpha_0}{d_4} \left( \frac{1}{1 + \left(\frac{A}{k_{FA}}\right)^n} + \frac{\alpha_4 \left(\frac{S^*}{k_{FS}}\right)^n}{1 + \left(\frac{S^*}{k_{FS}}\right)^n + \left(\frac{A}{k_{FA}}\right)^n} \right) \tag{1.7}$$

where $A^N(F)$ and $F^N(A)$ correspond to AP2 and SOC1/FUL nullclines, respectively. For Figure 3E, $A^N$ is plotted as a function of $A$, by first computing 1.7 and then introducing it into 1.6, leading to $A^N(F) = A^N(F^N(A))$. A multiplicative factor is used for visualization purposes ($c = 25$).

Floral activator mutants were modelled through a reduced activation expression rate ($\alpha_0$), with late flowering phenotypes paired with lower $\alpha_0$ values (i.e., $\alpha_0^{wt} > \alpha_0^{ful} > \alpha_0^{soc1} > \alpha_0^{soc1\,ful}$), whereas for $rAP2$, the inhibition of AP2 by the age signal was completely removed, mimicking the resistance to miR172 binding, and therefore using the following equation:

$$\frac{dA}{dt} = \frac{\alpha_1}{1 + \left(\frac{F}{k_{AF}}\right)^n} - d_1 A \tag{1.5}$$

See Suppl. Table 1 for parameter values used in the simulations. For making the comparison between theory and experiments in the figures, note that the $A$ variable is represented as [$AP2$], and the $F$ variable is represented as [$SOC1$].

## Model 2: Toggle-switch model with a negative feedback loop

This model is based on Model 1, except that the quasi-steady assumption for AP2 dynamics is absent because a time delay is required to generate oscillations in the system, and this is achieved by considering RNA explicitly. The dynamic equations over time ($t$) for $AP2$ mRNA ($Am$), AP2 protein intermediate-I as cytosolic translated protein ($Ac$), AP2 protein intermediate-II as functional nuclear transcription factor ($Ap$), SOC1/FUL protein ($Fp$) and the age-signal ($S$) concentrations can be written as:

$$\frac{dA_m}{dt} = \frac{\alpha_1}{1 + \left(\frac{A_p}{K_{AA}}\right)^n + \left(\frac{F_p}{k_{AF}}\right)^{n2}} - d_1 A_m \tag{2.1}$$

$$\frac{dA_c}{dt} = \frac{\alpha_2 A_m}{1 + \left(\frac{S}{k_{AS}}\right)^{n2}} - d_2 A_c \tag{2.2}$$

$$\frac{dA_p}{dt} = \alpha_3 A_c - d_3 A_p \tag{2.3}$$

$$\frac{dF_p}{dt} = \alpha_0 \left( \frac{1}{1 + \left(\frac{A_p}{k_{FA}}\right)^{n2}} + \frac{\alpha_4 \left(\frac{S}{k_{FS}}\right)^{n2}}{1 + \left(\frac{S}{k_{FS}}\right)^{n2} + \left(\frac{A_p}{k_{FA}}\right)^{n2}} \right) - d_4 F_p \tag{2.4}$$

$$\frac{dS}{dt} = \beta_s - d_s S \tag{2.5}$$

where $\alpha_i$ (i = 0,..,3) and $\beta_s$ are the basal production rates of *Am, Ac, Ap, Fp* and age; $d_x$ represents the degradation constants; $k_{ZY}$ are the concentrations of *Y* inhibiting or activating *Z* at which the production half the saturated value (notice that $K_{AA}$ is the equilibrium constant associated to AP2 auto-inhibition); and exponents $n$ and $n_2$ set the cooperativity ($n$ = 10 and $n_2$ = 2). AP2 is repressed by miR172, which is in turn activated by SPLs. This indirect inhibition is modelled here through the repression of *Ac* variable by the age signal, *S*. Although miR172 can also target AP2-RNA for degradation (cleavage mechanism[59,60]), most of its inhibition on AP2 occurs at the translational level[60]. This justifies why *S* was chosen to repress *Ac* and not *Am* in this model. Delay in AP2 self-repression was implemented, including two additional variables (*Am* and *Ac*). Without an explicit time delay, a negative feedback loop requires a minimum of three components to produce oscillations, provided there is sufficient nonlinearity[6,76]. To obtain limit cycle oscillations in such a simple three-component network, the Hill coefficient must be greater than eight[76]. See Suppl. Table 1 for parameter values. For *rAP2*, the inhibition of AP2 by the age signal was completely removed, mimicking the resistance to miR172 binding, and therefore using the following equation:

$$\frac{dA_c}{dt} = \alpha_2 A_m - d_2 A_c \tag{2.6}$$

**Spatial model**

This model consists of the 1D-spatial extension of Model 2. The dependence on the space variable has been introduced into the system through the age signal, which is now assumed to be able to diffuse in space.

$$\frac{\partial S(x,t)}{\partial t} = \beta_s(x, S) D \frac{\partial^2 S(x,t)}{\partial x^2} - d_s S \tag{3.1}$$

where the age signal production was considered just for a range of values of the spatial domain or when the age signal was above a certain threshold, specifically, $\beta_s > 0$ if $x \in [30,70]$ or if $S(x,t) > 0$. The diffusion of the age variable can be interpreted as a proxy of miR172 spatial movement through plasmodesmata during floral transition[37]. Zero flux boundary conditions were assumed, so the age variable did not diffuse out of the modelled space. The rest of the

equations from Model 3 are identical to Eqns. 2.1-2.4 with the adjustment that if x > 70 μm, $\alpha_i = 1/(\alpha_i \cdot (1+0.1 \cdot |x-70|))$ for i =1 and i =3. See Suppl. Table 1 for parameter values.

## Bifurcation and stability diagrams

The analytical stability diagrams showing the stability regimes of the system (Figures 5A, S7G and S7H) have been computed through continuation algorithms of the saddle-node and Hopf bifurcations in Python using [PyDSTool](77). A bifurcation diagram for a given $(\alpha_0, \alpha_1)$ as a function of $S^*$ is computed and then, from the saddle-node and/or Hopf bifurcation points, a continuation algorithm is run varying both parameters $\alpha_0$ and $\alpha_1$. Such continuation of the bifurcation point yields a 2D-stability diagram as shown in Figure S7G, which shows the different stability regions (monostable or bistable in this case) for the system for a particular $S^*$. In such a diagram, the boundaries between regions correspond to bifurcations.

Because $S$ is a system variable that depends on time, $S^*$-bifurcation diagrams can also be plotted as a function of time. This temporary dependency enables the study of the system's stability at each time. Similarly, as S changes along the development, different 2D-stability diagrams are computed for different $S^*$-values (Figure S7G). All these diagrams can be collapsed into a single one that distinguishes among regions in the space with different stability regimes as $S^*$ changes. These three-parameter stability diagrams are shown in Figures 5A and S7H. The initial conditions of the numerical simulations are always the steady-state concentrations when $S = 0$.

## Numerical integration analysis

Numerical simulations were performed in Python using also PyDSTool[77] toolkit with the VODE integrator, which can also be used for stiff systems. Although the output resembles that of a fixed time-step integrator, an adaptive time-step is used internally by this integrator. The creation of a Trajectory object is needed to later extract values at a desired resolution with the sample() method. Numerical simulations were initialised by setting the inhibitor variables ($A$, for Model 1; and $Am$, $Ac$ and $Ap$, for Model 2) at its corresponding steady state for $S = 0$ (i.e., $\beta_s$=0). If a limit cycle existed at $S = 0$, the center coordinate of the stable limit cycle of the corresponding inhibitor variable was used as the initial condition. Initial conditions for SOC1/FUL ($F$) and age ($S$) variables were set to zero for both models. Dynamics of $rAP2$ in Model 2 were initialised with [$Am$, $Ac$, $Ap$, $Fp$, $S$] = [0, 2, 15, 0, 0]. Model 3 initial conditions were set to zero for $Am$, $Ac$ and $Fp$ and $Ap$(x,0) $=15e^{(-2 \cdot 10^{-4} x^2)}$, fitting the spatial profile of AP2-VENUS fluorescence protein along the SAM at 7LDs. For the age variable, $S(0, x \geq 50)$ = $-0.25x^2 + 35x - 1125$ and $S(0, x \geq 70)$ = 100 to form an initial $S$ profile as shown in Figure 6B.

See Suppl. Tables 1 and 2 for parameters. To better understand how the bifurcation diagrams related to the actual numerical results, we visualised the bifurcation diagrams as a function of time, by assuming that $S^*$ is a variable $S(t)$ governed by equation 1.3 (or 2.5) with initial conditions at 0, whose solution is $S(t) = (\beta_s/d_s)(1 - e^{-d_s t})$. The simultaneous plot of bifurcation diagrams and numerical simulations highlights a strong agreement between simulations and theory (Figures S3, S11D–S11G and S12C–S12E). This bifurcation over time provides a notion of instantaneous stability of the trajectory of the system, whose accuracy will depend on the relative timescales of the age signal progression with respect to the timescales of the activator and inhibitor variables. Henceforth, the computed steady states should be interpreted as instantaneous stability of the system at a certain time point.

The first and second derivatives in Figures S5E and S5F were also computed numerically on the normalised numerical simulations of AP2 dynamics that had been previously smoothed to avoid numerical errors in the derivative with a Gaussian 1D-filter (*gaussian_filter1d* in Python). The derivative was performed by second order central accurate differences in the interior points of the array and first order accurate one-side differences at the boundaries (*gradient* function). The second derivative was computed using a second time the previous function. For visualization purposes, the first derivative was normalised by itself. See code for more details.

## Plant material and growth conditions

All plants in this study were in the *Arabidopsis thaliana* Columbia-0 (Col-0) background. Mutant alleles were previously described: *ap2-12*[41]*, soc1-2*[78] and *ful-2*[43]. The following transgenic lines were used: AP2::AP2-VENUS *ap2-12*[40], AP2::AP2-VENUS *ap2-12 soc1-2*[40] and AP2::rAP2-VENUS *ap2-12*[52]. AP2::AP2-VENUS *ap2-12 ful-2* and AP2::AP2-VENUS *ap2-12 soc1-2 ful-2* genotypes were generated in this study by crossing. An image dataset of AP2::AP2-VENUS *ap2-12*, and both AP2::AP2-VENUS *ap2-12 soc1-2* and *SOC1::SOC1-GFP soc1-2* images were obtained from a previous publication[40]. Plants were grown on soil under controlled conditions of LDs (16 h light/8 h dark).

## Confocal imaging

Shoot apices of plants at different developmental stages were dissected under a stereo microscope and fixed with 4% (w/v) paraformaldehyde (PFA; Electron Microscopy Sciences) by vacuum infiltration. Samples were incubated overnight in the dark and washed twice for 1 min in phosphate-buffered saline (PBS), to be transferred to ClearSee for 5 – 6 days[79]. Cell walls were stained with Renaissance 2200 [0.1% (v/v) in ClearSee][80] at least 1 day before imaging. Confocal microscopy was performed with a Stellaris 5 confocal microscope (Leica). Renaissance was excited at 405 nm, and image collection was performed at 445–475 nm (Figures 2A, 3A, 3B, 3E, 3F, S6J and S8A). VENUS was excited at 514 nm, and the signal was detected at 520–600 nm. Images of SOC1-GFP were obtained from published data[40].

## Fluorescence quantification

In this work, to analyse AP2 and SOC1 protein dynamics at the SAM during floral transition under LDs, we quantified previously published data[40] (AP2::AP2-VENUS ap2-12 and *SOC1::SOC1-GFP soc1-2)* and added further experimental repeats and genotypes (AP2::AP2-VENUS *ap2-12,* AP2::AP2-VENUS *ap2-12 ful-2,* AP2::AP2-VENUS *ap2-12 soc1-2 ful-2, and* AP2::rAP2-VENUS *ap2-12).* Previously, at the tissue level, we quantified normalised levels of translational reporters, referred to as effective concentrations, exclusively in fixed-volume regions in the SAM[40]. Here, to increase the accuracy of the analysis at the tissue level and determine how SAM growth influences such measurements, we reanalysed the data using a new computational pipeline that accounts for meristem size by defining characteristic expression domains from each reporter. Furthermore, we performed quantification at the single-cell level for some of the data to corroborate that tissue and single-cell level quantifications were showing equivalent results.

### Tissue quantification

All confocal fluorescence z-stacks were processed and analysed using our custom-made MATLAB code[38,40] (https://gitlab.com/slcu/teamHJ/pau/RegionsAnalysis), which we extended

it from previous versions as explained below. The main objective of this analysis was to obtain reproducible measures of fluorescence intensity within the SAM. A normalised fluorescence intensity measure, together with spatial SAM-longitudinal-axis expression profiles, was computed as a proxy for the concentration of protein at the SAM. A semi-automated pipeline was developed and refined for this purpose (Figure S1). To achieve homogenous volumetric resolution, z-stacks were resized via bicubic interpolation, increasing the number of slices in the z-direction to account for differences in the xz-plane and z-direction.

To exclude fluorescence signals outside the region of interest and to quantify only the fluorescence intensity within the meristematic region, a pre-processing step was performed: a 3D paraboloid mask was automatically generated using the curvature of the meristem (Figure S1). First, a stack-slice interval containing the meristem tip was selected, and the cell wall signal within this interval was projected onto the orthogonal xy and yz planes (Figure S1B). Then, two curved lines following the parabolic outline of the SAM were drawn from the sum-of-slice projection of each plane. Later, a parabolic fitting was performed on these two drawn lines (Figure S1B), accounting for potential SAM tilting relative to the vertical axis.

Specifically, the code recursively fits parabolas in different orientations of the drawn outline and selects the one that minimises the $R^2$ value. The coordinates of the apex, necessary to derive the 3D paraboloid equation, were computed from the two orthogonal parabolas fitted for each z-stack. The $z_O$ coordinate of the paraboloid was determined by averaging the apices of the orthogonal parabolas (Figure S1B). The parameter $a$ in the parabola equation (i.e the curvature) was used to replace the denominator terms in the paraboloid equation ($c_1^2$ and $c_2^2$; Figure S1 legend) such that the paraboloid equation matched the linear and quadratic terms of their respective equations at $y = y_0$ and $x = x_0$. Since z-stacks did not always encompass the entire meristem in the yz plane (lateral view), only the xy curvature was used ($c_2^2 = c_1^2$). Finally, to extract the fluorescence signal within the SAM using the 3D paraboloid, a 2D parabolic mask was created for each z-stack slice, and, for each slice, all intensity values of pixels outside the paraboloid were set to 0 (Figure S1C).

To exclude fluorescence signal at the boundaries of the SAM and primordia, the paraboloid's curvature was increased relative to the original (Figure S1C), such that $a' = a/\alpha$, where $a'$ is the curvature of the new paraboloid and $\alpha$ is the image resolution ($\alpha < 1$ µm) (Figure S1 legend). The 3D paraboloid was then divided into consecutive 3D subsections of 10 µm-height along the longitudinal axis of the SAM starting from the apex (Figure S1D). A fluorescence intensity concentration profile along the SAM longitudinal axis was extracted by computing the concentration of fluorescence signal within each section. Concentration was defined as the ratio of total intensity (sum of voxel intensities) to the total volume (sum of voxel volumes).

To determine the effective fluorescence concentrations in each meristem, a characteristic domain was defined based on the median concentration profile for each time point in each experiment (Figure S1E). The boundaries of these characteristic domains were defined by the points at which the normalised median concentration profile crossed a predefined threshold. Two or three thresholds were used to demonstrate the robustness of this quantification: 0.25, 0.50 and 0.75, always including the last two. The effective fluorescence concentrations per meristem were then calculated by dividing the total intensity within all sections included in the defined characteristic domain by the sum of the volume of all the voxels in these sections

(Figure S1E). The discrete nature of the SAM-paraboloid partitioned sections was handled by shifting the boundaries of the characteristic domains to those of the nearest paraboloid subsection.

To quantify SOC1::SOC1:GFP *soc1-2 ap2-12* fluorescence, a Gaussian filter ($\sigma$ = 2.5) was applied in the region within the paraboloid to homogenise a weak reporter signal. Two independent experiments for AP2::AP2:VENUS *ap2-12* were combined (Figures S2A–S2C) for all non-spatial AP2 data shown. The 7LDs data points from the *wt-3* experiment (Figures S2A–S2C) were not included in the analysis due to poor signal quality in the samples. An additional tissue quantification analysis was also performed for AP2::rAP2:VENUS *ap2-12* (Figures S8A–S8C). In Fiji, the fluorescence signal from 19, 21 and 24LDs samples was projected and summed within 10 z-stack slices (40 µm) containing the central region of the SAM, and then homogenised with a 3D-Gaussian filter (radius 2.5 µm). A curved line was drawn between the two uppermost primordia at opposite sides of the main shoot (Figure S8C). Normalised intensity profiles by time point were then extracted to highlight the lower expression of rAP2 at regions adjacent to the SAM primordia, compared with that in the central part of the SAM (Figure S8C).

### Single-cell quantification
All confocal fluorescence z-stacks with AP2-VENUS or rAP2-VENUS reporters were also analysed at the single-cell level. Confocal fluorescence z-stack .lif files were converted to .tif files, and prior to segmentation, a 3D-Gaussian filter (r = 2 µm) was applied. Nuclei expressing the reporters were segmented using Stardist[81] with [StarDist Plant Nuclei 3D ResNet](#) model[82] (Figures S8D and S8E). In MATLAB, the segmented images were then re-labelled and single-nuclear features (nuclear concentration, volume and position) were computed (Figures S8F, S9 and S10) using the original and labelled images, together with the 3D-paraboloid generated for each SAM by the tissue quantification pipeline (Figure S1).

## SAM morphology quantification
To quantify the morphology of the meristem, its height and width were measured. The parabolic fits, obtained from the parabolic outline of the SAM drawn in the *xy* and *yz* planes between the uppermost primordia on opposite sides of the SAM (Figure S1), were used to obtain the coordinate of the apex and calculate the horizontal (width) and vertical (height) distance from the apex to the nearest primordium (Figures S2B, S2C, S6A, S6B, S6F and S6G).

## Statistics and reproducibility
Plants from a single experiment were grown together in the same chamber, and their positions in the growth chamber were randomised to avoid bias due to non-homogeneous growth conditions. To minimise potential effects of the clearing treatment on fluorescence signal quantification, samples of each genotype were included in each imaging session when time point acquisition extended beyond one day. Meristems exhibiting severe defects (i.e. tissue breakage, poor cell wall dye penetration, or blocking of the fluorescence signal by leaf primordia) and/or strong developmental differences compared to the other samples from the same time point were not considered for fluorescence quantification. A detailed list of the meristems analysed in each experiment is provided in the code files. Investigators were not blinded in allocation during experiments and outcome assessment, because the material was rigorously labelled, thus making blinding impossible. See Suppl. Tables 3-9 for statistics.

## Data availability

The original confocal microscopy images for one experiment of AP2-VENUS expression in *ap2* and *ap2 soc1* and SOC1-GFP in *soc1* and in *soc1 ap2* have been obtained from this study[40]. The remaining confocal microscopy images can be provided upon request. Source data used for the analysis is available in the OSF repository https://osf.io/jyq58/.

## Code availability

Source code for the the image analysis at both tissue and single cell levels can be found in the OSF repository https://osf.io/jyq58/ and also in the following GitLab repositories https://gitlab.gwdg.de/devplantpatterning/Publications/2025-rodriguez-maroto_wang_et_al and https://gitlab.com/slcu/teamHJ/pau/RegionsAnalysis/-/tags/Rodriguez_Maroto_et_al_2025_Region_Analysis_code.

## Acknowledgments

We thank Josep Mercadal, John Chandler, and Ruben Perez-Carrasco for critical comments on the manuscript, and Marta Ibañes Miguez and James Locke for fruitful discussions. G.R.-M. acknowledges funding from the International Max Planck Research School on "Understanding Complex Plant Traits Using Computational and Evolutionary Approaches", and K.W. received a studentship from the China Scholarship Council. P.C. and M.C. received a post-doctoral fellowship from the Alexander von Humboldt foundation (ESP - 1238682 - HFST-P and ITA 1216206 HFST-P, respectively), P.C. also received a post-doctoral Marie Curie fellowship (101150752 from HORIZON-MSCA-2023-PF-01). G.C. received funding from the Deutsche Forschungsgemeinschaft (DFG) through grant CO 318/14-1. G.C. and P.F.-J. also received funding from the DFG through the Cluster of Excellence CEPLAS (EXC 2048/1 Project ID: 390686111) and from a Core Grant from the Max Planck Society. P.F.-J. also acknowledges funding from the National Science Foundation (US NSF, https://www.nsf.gov), Division of Biological Infrastructure DBI-232051.

## Author contributions

G.R.-M., K.W., G.C., and P.F.-J. conceived and designed the study. K.W. generated the AP2::AP2-VENUS *ap2-12 ful-2* and AP2::AP2-VENUS *ap2-12 soc1-2 ful-2* transgenic lines. G.R.-M., K.W. and M.C. performed confocal imaging. G.R.-M., P.F.-J. and P.C.-F. designed the imaging quantification pipelines. G.R.-M. built and performed the quantification analyses. P.C.-F. assessed primordia identity. G.R.-M. and P.F.-J. formulated and analysed the mathematical model. G.R.-M., G.C. and P.F.-J. wrote the manuscript. P.F.-J., G.C., M.C. and P.C.-F. provided scientific guidance throughout the project. All the authors discussed the results and commented on the manuscript.

## Declaration of interests

The authors declare no competing interests.

## References

1. Sáez, M., Briscoe, J. & Rand, D. A. Dynamical landscapes of cell fate decisions. *Interface Focus* **12**, (2022).
2. Lenne, P.-F. & Trivedi, V. Sculpting tissues by phase transitions. *Nat. Commun.* **13**, 664 (2022).

3. Force, E., Lamy, D., Debernard, S., Savouré, A. & Dacher, M. Developmental transitions involve common biological processes across living beings. *Heliyon* **11**, e42995 (2025).

4. Waddington, C. H. *The Strategy of the Genes. A Discussion of Some Aspects of Theoretical Biology*. (Routledge, London and New York, 1957).

5. Verd, B., Crombach, A. & Jaeger, J. Classification of transient behaviours in a time-dependent toggle switch model. *BMC Syst. Biol.* **8**, (2014).

6. Goodwin, B. C. Oscillatory behavior in enzymatic control processes. *Adv. Enzyme Regul.* **3**, 425–437 (1965).

7. Jaeger, J. & Crombach, A. Life's Attractors. in *Evolutionary Systems Biology. Advances in Experimental Medicine and Biology* vol. 751 (Springer, New York, NY., 2012).

8. Ferrell, J. E. Bistability, bifurcations, and Waddington's epigenetic landscape. *Curr. Biol.* **22**, (2012).

9. Huang, S. The molecular and mathematical basis of Waddington's epigenetic landscape: A framework for post-Darwinian biology? *BioEssays* **34**, 149–157 (2012).

10. Ferrell, J. E. Self-perpetuating states in signal transduction: positive feedback, double-negative feedback and bistability. *Curr. Opin. Cell Biol.* **14**, 140–148 (2002).

11. Rombouts, J. & Gelens, L. Dynamic bistable switches enhance robustness and accuracy of cell cycle transitions. *PLoS Comput. Biol.* **17**, (2021).

12. Strogatz, S. H. *NONLINEAR DYNAMICS AND CHAOS: With Applications to Physics, Biology, Chemistry, and Engineering*. *Nonlinear Dynamics and Chaos: With Applications to Physics, Biology, Chemistry, and Engineering* (2018). doi:10.1201/9780429492563.

13. Kadiyala, U. *et al.* From genes to patterns: five key dynamical systems concepts to decode developmental regulatory mechanisms. *Development* **152**, dev204617 (2025).

14. Strogatz, S. H. & Westervelt, R. M. Predicted power laws for delayed switching of charge-density waves. *Phys. Rev. B* **40**, 10501–10508 (1989).

15. Sardanyés, J., Raich, C. & Alarcón, T. Noise-induced stabilization of saddle-node ghosts. *New J. Phys.* **22**, 093064 (2020).

16. Karin, O., Miska, E. A. & Simons, B. D. Epigenetic inheritance of gene silencing is maintained by a self-tuning mechanism based on resource competition. *Cell Syst.* **14**, 24-40.e11 (2023).

17. Koch, D. *et al.* Ghost Channels and Ghost Cycles Guiding Long Transients in Dynamical Systems. *Phys. Rev. Lett.* **133**, 047202 (2024).

18. Meisel, C., Klaus, A., Kuehn, C. & Plenz, D. Critical Slowing Down Governs the Transition to Neuron Spiking. *PLOS Comput. Biol.* **11**, e1004097 (2015).

19. Maturana, M. I. *et al.* Critical slowing down as a biomarker for seizure susceptibility. *Nat. Commun.* **11**, 2172 (2020).

20. Krotov, D., Dubuis, J. O., Gregor, T. & Bialek, W. Morphogenesis at criticality. *Proc. Natl. Acad. Sci.* **111**, 3683–3688 (2014).

21. Tufcea, D. E. & François, P. Critical Timing without a Timer for Embryonic Development. *Biophys. J.* **109**, 1724–1734 (2015).

22. Dakos, V. & Bascompte, J. Critical slowing down as early warning for the onset of collapse in mutualistic communities. *Proc. Natl. Acad. Sci.* **111**, 17546–17551 (2014).

23. Ferrell, J. E. & Xiong, W. Bistability in cell signaling: How to make continuous processes discontinuous, and reversible processes irreversible. *Chaos* **11**, (2001).

24. Tsai, T. Y. C. *et al.* Robust, tunable biological oscillations from interlinked positive and negative feedback loops. *Science* **321**, (2008).

25. Gardner, T. S., Cantor, C. R. & Collins, J. J. Construction of a genetic toggle switch in Escherichia coli. *Nature* **403**, 339–342 (2000).

26. Sáez, M. *et al.* Statistically derived geometrical landscapes capture principles of decision-making dynamics during cell fate transitions. *Cell Syst.* **13**, (2022).

27. Huang, S., Guo, Y. P., May, G. & Enver, T. Bifurcation dynamics in lineage-commitment in bipotent progenitor cells. *Dev. Biol.* **305**, (2007).

28. Simons, B. D. & Karin, O. Cell cycle criticality as a mechanism for robust cell population control. *Mol. Syst. Biol.* 1–18 (2025) doi:10.1038/s44320-025-00164-8.

29. Gilbert, S. F. *Epigenetic Landscaping: Waddington's Use of Cell Fate Bifurcation Diagrams. Biology and Philosophy* vol. 6 135–154 (1991).

30. Davila-Velderrain, J., Martinez-Garcia, J. C. & Alvarez-Buylla, E. R. Modeling the epigenetic attractors landscape: Toward a post-genomic mechanistic understanding of development. *Front. Genet.* **6**, (2015).

31. Davila-Velderrain, J., Martinez-Garcia, J. C. & Alvarez-Buylla, E. R. Dynamic network modelling to understand flowering transition and floral patterning. *J. Exp. Bot.* **67**, 2565–2572 (2016).

32. Azpeitia, E. *et al.* Cauliflower fractal forms arise from perturbations of floral gene networks. *Science* **373**, 192–197 (2021).

33. Coen, E. & Prusinkiewicz, P. Developmental timing in plants. *Nat. Commun.* **15**, 2674 (2024).

34. Bäurle, I. & Dean, C. The Timing of Developmental Transitions in Plants. *Cell* **125**, 655–664 (2006).

35. Zhu, Y. *et al.* TERMINAL FLOWER 1-FD complex target genes and competition with FLOWERING LOCUS T. *Nat. Commun.* **11**, (2020).

36. Fuchs, M. & Lohmann, J. U. Aiming for the top: non-cell autonomous control of shoot stem cells in Arabidopsis. *J. Plant Res.* **133**, (2020).

37. Ó'Maoiléidigh, D. S. *et al.* Systematic analyses of the MIR172 family members of Arabidopsis define their distinct roles in regulation of APETALA2 during floral transition. *PLoS Biol.* **19**, (2021).

38. Cerise, M. *et al.* Two modes of gene regulation by TFL1 mediate its dual function in flowering time and shoot determinacy of Arabidopsis. *Development* **150**, dev202089 (2023).

39. Kinoshita, A. *et al.* Regulation of shoot meristem shape by photoperiodic signaling and phytohormones during floral induction of arabidopsis. *eLife* **9**, (2020).

40. Bertran Garcia de Olalla, E. *et al.* Coordination of shoot apical meristem shape and identity by APETALA2 during floral transition in Arabidopsis. *Nat. Commun.* **15**, 6930 (2024).

41. Yant, L. *et al.* Orchestration of the floral transition and floral development in arabidopsis by the bifunctional transcription factor APETALA2. *Plant Cell* **22**, 2156–2170 (2010).

42. Immink, R. G. H. *et al.* Characterization of SOC1's central role in flowering by the identification of its upstream and downstream regulators. *Plant Physiol.* **160**, 433–449 (2012).

43. Romera-Branchat, M. *et al.* Functional Divergence of the Arabidopsis Florigen-Interacting bZIP Transcription Factors FD and FDP. *Cell Rep.* **31**, (2020).

44. Wang, J. W., Czech, B. & Weigel, D. miR156-Regulated SPL Transcription Factors Define an Endogenous Flowering Pathway in Arabidopsis thaliana. *Cell* **138**, 738–749 (2009).

45. Hyun, Y., Richter, R. & Coupland, G. Competence to flower: Age-controlled sensitivity to environmental cues. *Plant Physiol.* **173**, 36–46 (2017).

46. Gandikota, M. *et al.* The miRNA156/157 recognition element in the 3′ UTR of the

Arabidopsis SBP box gene SPL3 prevents early flowering by translational inhibition in seedlings. *Plant J.* **49**, (2007).

47. Hyun, Y. *et al.* Multi-layered Regulation of SPL15 and Cooperation with SOC1 Integrate Endogenous Flowering Pathways at the Arabidopsis Shoot Meristem. *Dev. Cell* **37**, (2016).

48. Yamaguchi, A. *et al.* The MicroRNA-Regulated SBP-Box Transcription Factor SPL3 Is a Direct Upstream Activator of LEAFY, FRUITFULL, and APETALA1. *Dev. Cell* **17**, (2009).

49. Chen, X. A MicroRNA as a Translational Repressor of APETALA2 in Arabidopsis Flower Development. *Science* **303**, (2004).

50. Mathieu, J., Yant, L. J., Mürdter, F., Küttner, F. & Schmid, M. Repression of flowering by the miR172 target SMZ. *PLoS Biol.* **7**, (2009).

51. Balanzà, V. *et al.* Genetic control of meristem arrest and life span in Arabidopsis by a FRUITFULL-APETALA2 pathway. *Nat. Commun.* **9**, (2018).

52. Sang, Q. *et al.* MicroRNA172 controls inflorescence meristem size through regulation of APETALA2 in Arabidopsis. *New Phytol.* **235**, 356–371 (2022).

53. Jaeger, K. E., Pullen, N., Lamzin, S., Morris, R. J. & Wigge, P. A. Interlocking Feedback Loops Govern the Dynamic Behavior of the Floral Transition in Arabidopsis. *Plant Cell* **25**, 820–833 (2013).

54. Ferrándiz, C., Gu, Q., Martienssen, R. & Yanofsky, M. F. Redundant regulation of meristem identity and plant architecture by FRUITFULL, APETALA1 and CAULIFLOWER. *Development* **127**, (2000).

55. Melzer, S. *et al.* Flowering-time genes modulate meristem determinacy and growth form in Arabidopsis thaliana. *Nat. Genet.* **40**, (2008).

56. Andrés, F. *et al.* SHORT VEGETATIVE PHASE reduces gibberellin biosynthesis at the Arabidopsis shoot apex to regulate the floral transition. *Proc. Natl. Acad. Sci.* **111**, (2014).

57. Elowitz, M. B. & Leibier, S. A synthetic oscillatory network of transcriptional regulators. *Nature* **403**, (2000).

58. Perez-Carrasco, R. *et al.* Combining a Toggle Switch and a Repressilator within the AC-DC Circuit Generates Distinct Dynamical Behaviors. *Cell Syst.* **6**, 521-530.e3 (2018).

59. Schwab, R. *et al.* Specific effects of microRNAs on the plant transcriptome. *Dev. Cell* **8**, 517–527 (2005).

60. Aukerman, M. J. & Sakai, H. Regulation of Flowering Time and Floral Organ Identity by a MicroRNA and Its APETALA2-Like Target Genes. *Plant Cell* **15**, (2003).

61. Fouracre, J. P. & Scott Poethig, R. Role for the shoot apical meristem in the specification of juvenile leaf identity in Arabidopsis. *Proc. Natl. Acad. Sci.* **116**, (2019).

62. Alon, U. Network motifs: theory and experimental approaches. *Nat. Rev. Genet.* **8**, 450–461 (2007).

63. Perez-Carrasco, R., Guerrero, P., Briscoe, J. & Page, K. M. Intrinsic Noise Profoundly Alters the Dynamics and Steady State of Morphogen-Controlled Bistable Genetic Switches. *PLOS Comput. Biol.* **12**, e1005154 (2016).

64. Maretvadakethope, S. *et al.* Let there be multifunctionality: Uncovering the criticality zoo of the AC-DC genetic circuit. *bioRxiv* (2025).

65. Balaskas, N. *et al.* Gene Regulatory Logic for Reading the Sonic Hedgehog Signaling Gradient in the Vertebrate Neural Tube. *Cell* **148**, 273–284 (2012).

66. Verd, B. *et al.* A damped oscillator imposes temporal order on posterior gap gene expression in Drosophila. *PLOS Biol.* **16**, e2003174 (2018).

67. Nagata, S. & Kikuchi, M. Emergence of cooperative bistability and robustness of gene

regulatory networks. *PLOS Comput. Biol.* **16**, e1007969 (2020).

68. Abley, K. *et al.* An ABA-GA bistable switch can account for natural variation in the variability of Arabidopsis seed germination time. *eLife* **10**, e59485 (2021).

69. Abley, K., Goswami, R. & Locke, J. C. W. Bet-hedging and variability in plant development: seed germination and beyond. *Philos. Trans. R. Soc. B Biol. Sci.* **379**, 20230048 (2024).

70. Tooke, F., Ordidge, M., Chiurugwi, T. & Battey, N. Mechanisms and function of flower and inflorescence reversion. *J. Exp. Bot.* **56**, 2587–2599 (2005).

71. McCullough, E. *et al.* Photoperiod-dependent floral reversion in the natural allopolyploid Arabidopsis suecica. *New Phytol.* **186**, 239–250 (2010).

72. Torti, S. *et al.* Analysis of the Arabidopsis Shoot Meristem Transcriptome during Floral Transition Identifies Distinct Regulatory Patterns and a Leucine-Rich Repeat Protein That Promotes Flowering. *Plant Cell* **24**, 444–462 (2012).

73. Verd, B., Crombach, A. & Jaeger, J. Dynamic Maternal Gradients Control Timing and Shift-Rates for Drosophila Gap Gene Expression. *PLOS Comput. Biol.* **13**, e1005285 (2017).

74. Balanzà, V., Martínez-Fernández, I. & Ferrándiz, C. Sequential action of FRUITFULL as a modulator of the activity of the floral regulators SVP and SOC1. *J. Exp. Bot.* **65**, 1193–1203 (2014).

75. Koch, D., Nandan, A., Ramesan, G. & Koseska, A. Biological computations: Limitations of attractor-based formalisms and the need for transients. *Biochem. Biophys. Res. Commun.* **720**, 150069 (2024).

76. Griffith, J. S. Mathematics of cellular control processes I. Negative feedback to one gene. *J. Theor. Biol.* **20**, 202–208 (1968).

77. Clewley, R., Sherwood, W. E., LaMar, M. & Guckenheimer, J. PyDSTool, a software environment for dynamical systems modeling. (2007).

78. Lee, H. *et al.* The AGAMOUS-lIKE 20 MADS domain protein integrates floral inductive pathways in Arabidopsis. *Genes Dev.* **14**, 2366–2376 (2000).

79. Kurihara, D., Mizuta, Y., Sato, Y. & Higashiyama, T. ClearSee: a rapid optical clearing reagent for whole-plant fluorescence imaging. *Development* **142**, 4168–4179 (2015).

80. Musielak, T. J., Schenkel, L., Kolb, M., Henschen, A. & Bayer, M. A simple and versatile cell wall staining protocol to study plant reproduction. *Plant Reprod.* **28**, 161–169 (2015).

81. Weigert, M., Schmidt, U., Haase, R., Sugawara, K. & Myers, G. Star-convex Polyhedra for 3D Object Detection and Segmentation in Microscopy. in *2020 IEEE Winter Conference on Applications of Computer Vision (WACV)* 3655–3662 (2020). doi:10.1109/WACV45572.2020.9093435.

82. Vijayan, A. *et al.* A deep learning-based toolkit for 3D nuclei segmentation and quantitative analysis in cellular and tissue context. *Development* **151**, dev202800 (2024).

# Main Figures

**Figure 1. Monostable versus bistable transitions in the context of floral transition at the SAM.** A) Waddington/quasi-potential landscape cartoons illustrating a monostable (top row) and a bistable (bottom row) transition. The system's state is depicted as a marble that is attracted to stable steady states (stable attractors), located at the bottom of the valleys. In the monostable transition (top row), the only existing valley changes over time. A threshold (grey dashed line) arbitrarily separates developmental state A (yellow) from developmental state B (blue). In the bistable transition (bottom row), two different valleys coexist for some time, each containing a stable steady state representing a different developmental state. The system only moves to the new stable attractor when the initial stable attractor disappears. The disappearance of the attractor flattens the landscape, and the system undergoes a slowing down of the dynamics, remaining in a transient state reminiscent of the previous unstabilised state, known as a ghost state. B) Scheme of simplified regulatory network underlying floral transition at the SAM. Light and age factors modulate mutual inhibition between floral activators and inhibitors over time. Normal arrows indicate activation; blunt arrows represent inhibition.

**Figure 2. A time-dependent toggle-switch can recapitulate AP2 protein dynamics at the SAM in wt, indicating that time-dependent bistability underlies floral transition.** A and B) Expression of AP2::AP2::VENUS (A) and SOC1::SOC1:GFP (B) at the SAM in *ap2-12* and *soc1*-2 backgrounds, respectively, grown under LDs. Scale bars = 20 μm. Arrowheads indicate visible floral primordia. C) Quantification of the temporal dynamics of AP2 (yellow) and SOC1

(cyan) effective concentrations (see Methods). Circles represent normalised median protein concentrations; shaded regions indicate the inter-quartile range (IQR). The arrowhead indicates floral transition completion (visible floral primordial, see Figure S6A). D) Toggle-switch model network diagram for studying floral transition and the influence of environmental/endogenous signals. $A$: AP2; $F$: FUL and SOC1; $S$: age endogenous signal. E–M) Model analysis. E, H, and K) Bifurcation diagrams for $A$ (yellow) and $F$ (cyan) as a function of the age-signal ($S^*$) for (E) a monostable transition, (H) a bistable transition, and (K) a bistable transition representing the wt-genotype. Solid lines represent stable steady states, dashed lines represent unstable states. Triangles on the x-axes indicate the steady state value of $S$ ($S_{ss}$). Bistability occurs in the regions in grey. Coloured circles indicate saddle-node bifurcation points. F, I, J and L) Normalised numerical simulations showing the temporal dynamics of the activator (cyan) and inhibitor (yellow) (F) a monostable transition, associated to (E); (I, J) bistable transitions for two different parameter values of the age signal production, associated to (H), in which $S_{ss}$ is indicated with a white and black triangle on the x-axis for (I) and (J), respectively; and (L) a bistable transition representing wt, associated to (K). G) Age-signal ($S$) dynamics for (F, I) in red, and for J in dark red. In (J), the saturation of the age signal is closer to the saddle-node bifurcation, resulting in an extremely long transient due to the increased influence of the ghost attractor. M) Comparison between experimental (continuous yellow) and model (dashed black) results for AP2 dynamics. See Suppl. Table 1 for parameter values.

**Figure 3. Mutations of floral activator genes prolong the transient-expressing phase of the inhibitor and reveal its fluctuating behaviour.** A and B) Expression pattern of AP2:AP2::VENUS in (A) *ap2-12 ful-2* and (B) *ap2-12 soc1-2* at the SAM of plants grown under LDs. Scale bars = 20 μm. C) Quantification of the temporal dynamics of AP2 concentration at the SAM in wt (yellow), *ful* (orange), and *soc1* (red). Circles and shaded regions are defined as in Figure 2C. Arrows indicate floral transition completion (visible floral primordia, see Figure S6). D) Numerical simulations for wt, *ful*, and *soc1* showing the *AP2* model variable dynamics. E) AP2-age bifurcation diagram (left) and nullclines (right) for wt, *ful,* and *soc1*. The bifurcation

diagram is represented as in Figure 2. F and G) Expression pattern of (F) AP2:AP2::VENUS in *ap2-12 soc1-2 ful-2* and (G) AP2::rAP2:VENUS *ap2-12* at the SAM under LDs. Scale bars = 20 μm. H) Quantification of the temporal dynamics of AP2 concentrations at the SAM in wt (yellow), *soc1 ful* (pink), and *rAP2* (purple). I) Single-cell nuclear quantification of AP2 protein expression in *soc1 ful* (pink) and *rAP2* (purple) at the SAM during floral transition, for the same SAMs analysed in (H). The inset shows the median trends. J) AP2 model temporal dynamics for wt (yellow) and *soc1 ful* (pink). See Suppl. Table 1 for parameter values.

**Figure 4. An extended toggle-switch model incorporating AP2 autoinhibition driving oscillations can still recapitulate the wt-dynamics during floral transition.** A) Extended toggle-switch network-model diagram incorporating AP2-autoinhibition (left) and a detailed version of the diagram (right), introducing new variables to describe the AP2 autoinhibition, to ensure sufficient time delay to drive oscillations. B) Bifurcation diagrams for model variables *Am* (orange-brown), *Ac* (light brown), *Ap* (dark yellow) and *Fp* (cyan) as a function of the age-associated signal ($S^*$) for wt. Bifurcation diagrams are represented as in Figure 2; here, circles represent saddle-node and Hopf bifurcations. C) Numerical simulations for wt showing the temporal dynamics of the activator (Fp, cyan) and inhibitor (Ap, yellow) proteins. *S* dynamics are in red in the top-right corner. D) Comparison between experimental (yellow) and the extended model's (dashed black) results for AP2 expression at the SAM. See Suppl. Table 1 for parameter values.

**Figure 5. The extended model explains genotype-specific dynamics and qualitative features of floral transition**. A) Stability diagram illustrating the stability regimes (M: monostable, B: bistable) crossed for a given $(\alpha_0, \alpha_1)$ parameter pair (representing *Fp* and *Ap* expression rates, respectively) across the *S\** range [0, 750]. Genotypes are represented by coloured dots. Stability regimes include: M, where the system is always monostable; B-M,

where the system starts bistable and transitions to monostable; M-B-M, involving two saddle-node bifurcations; and M-B, where $S^*$ is insufficient to reach the second saddle-node. The red line marks the boundary above which oscillations occur. B) *Ap* bifurcation diagrams (left) for wt, *ful*, *soc1,* and *soc1 ful* as a function of $S^*$. Normalised numerical simulations (right) show AP2 and $S$ dynamics across genotypes. C–G) Comparison between the modelled genotypes and the experimental data for (C) wt, (D) *ful,* (E) *soc1*, (F) *soc1 ful,* and (G) *rAP2.* Network diagrams (first column), bifurcation diagrams (second and third columns), numerical simulations reached for $S_{ss}$ = 750 (fourth column), and comparison between experimental (solid line) and simulated (dashed black line) AP2 dynamics (fifth column). Bifurcation diagrams are represented as in Figure 4B. See Suppl. Table 1 for parameter values.

**Figure 6. A spatially extended model reveals different dynamical scenarios within the SAM during floral transition.** A) Temporal and spatial dynamics of normalised AP2 (yellow) and SOC1 (cyan) expression during floral transition at the SAM in wt. $d_{apex}$ represents the

distance to the SAM apex. B) Numerical simulations for *Ap*, *Fp,* and *S* wt variables along one spatial dimension at different time points. Only diffusion of the age signal, *S*, has been added to the model analysed in Figures 4 and 5. C, D) Extracted spatial (C) and (D) temporal profiles from (B), showing *Ap*, *Fp,* and *S*. In (C), spatial profiles as a function of the distance to the SAM apex are normalised by their global maximum. In (D), temporal dynamics are shown at different positions along SAM's longitudinal axis, and coloured dots correspond to those in (C). E) Extracted temporal profiles at different positions along SAM's longitudinal axis from the spatial model for different $S_{SS}$. *Ap* and *Fp* are normalised by their maximum values across the four scenarios displayed; *S* has been normalised within each scenario. The modelled spatial and concentration variables are in arbitrary units. See information about parameters and equations in Methods and in Suppl. Table 1.