Compaction and swelling of single stretched DNAs driven by molecular crowding

Paritosh Gupta[1,2,3], John F. Marko[‡,4], and Vittore F. Scolari[‡,1,2]

[1]*Institut Curie, PSL Research University, Sorbonne Université, CNRS UMR168, Laboratoire Physico Chimie Curie, 75005 Paris, France.*
[2]*Institut Curie, PSL Research University, Sorbonne Université, CNRS UMR3664, Laboratoire Dynamique du Noyau, 75005 Paris, France.*
[3]*Department of Physics, Indian Institute of Science Education and Research, Dr. Homi Bhabha Road, Pune, 411 008, India*
[4]*Department of Molecular Biosciences and Department of Physics and Astronomy, Northwestern University, Evanston, Illinois 60208, USA*

December 9, 2025

Abstract

We present a theory for the effects of osmotic pressure exerted by macromolecular crowders, on a double-stranded DNA (or any other semi-flexible polymer) extended by tension. Our results predict DNA force extension curves. The lowest-order effect is a crowder-dependent compression that counteracts the external stretching force. This compression effect can collapse the polymer at a crowder-dependent critical force f^*. This compression is dependent on crowder radius r and density ϕ, with higher densities and smaller radii having greater effects on the force-extension curve. At first order in perturbation theory we also find a fluctuation-dependent correction to the depletion volume, which can overwhelm the simple compression effect for large crowders, leading to expansion of the polymer.

The genetic material of living cells is confined in compartments where it shares its space with a large number of other macromolecules fundamental for cellular functions. Many of these macromolecules, such as transcription factors, DNA and RNA polymerases, nucleosomes, specific enzymes and molecular motors associate with DNA, forming nucleoprotein complexes along *chromatin* in eukaryotes, or the *nucleoid* in bacteria. In both these cases, one can consider the chromosome fiber as a polymer: in eukaryotes, chromatin is at base a string of nucleosomes, while in bacteria the nucleoid fiber consists of DNA decorated by DNA-binding proteins. However, many other macromolecules near the genome do not bind to the nucleoid or chromatin fibers, but do share the

---

[‡] e-mail: vittore.scolari@cnrs.fr, john-marko@northwestern.edu

same constrained space inside the cell, and interact with the genome via steric exclusion.

These "free molecules", that we call *crowders*, are characterized by a high diffusive mobility, and produce osmotic pressure. This pressure is exerted towards the exterior of the cell, on the plasma or cell membrane, and is a key factor in the regulation of overall cellular volume [RJS23]. The same pressure is applied towards the interior, and ultimately on the genetic material. Analogous to how depletion forces by macromolecules can induce attractive interactions between larger colloidal particles [AO54], these osmotic depletion forces can compact chromatin and induce its phase separation [Odi98, BBD⁺12].

Experiments *in vivo* have observed osmotic shock-driven condensation of the nucleoid in *Escherichia coli*, in a reduction of nucleiod volume following an increase of salt concentration in the extracellular medium [YMRM20]. The osmotic shock drives water out of the cell and causes a direct reduction in cell volume. In turn, this induces an increase in concentrations of the crowders in the cytoplasm. It was further shown that the volume reduction is considerably smaller for the cytoplasm than for the nucleoid, hinting at a cooperative effect which can be thought about as a type of phase separation (i.e., via considering the energetic cost of expanding the nucleoid volume).

Complementary *in vitro* experiments using purified bacterial nucleoids and a set of crowders solutions covering a large range of densities and sizes have shown that crowders at physiological levels can directly compact the nucleoid [PHH⁺12, RD15, JNCP16], indicating that crowding can be a key factor in chromosome condensation. These observations indicate that crowders at physiological densities can regulate the level of condensation of the bacterial chromosome fiber into the nucleoid, allowing its membraneless compartmentalization.

Similar phenomena have been observed for eukaryotic chromatin in human cell lines: an increase in density of molecular crowders has been correlated to chromosomal mitotic condensation [IIT⁺24] following the dissolution of the nuclear envelope. Osmotic shock has been shown to condense mitotic chromosomes similar to how it affects the nucleoid in *E. coli* [IIT⁺24]. Further, it is possible that the effect of crowding-induced liquid-liquid phase separation of key chromatin-associated proteins [AAL⁺02] is relevant for the chromosome's three dimensional compartmentalization in silent and transcribed regions.

Experimental results and theory have been complemented by molecular dynamics simulations reproducing crowders mediated compaction of polymer models *in silico* [SBdH⁺15, KPHT15, JJH16, YMRM20, CCG⁺24]. Typical simulations of this type consist of L spherical beads of radius r_0 connected by springs of a similar extension, immersed in in a bath of small spherical crowders of radius r with density c. These simulations identified that the crowder volume fraction $\phi = 4/3\,\pi r^3 c$ as well as the relative crowder size r/r_0 are fundamental in setting the strength of the crowder-induced compacting force. While simulations has demonstrated that crowders have the ability to compact and collapse polymers, there are limits of this approach towards a quantitative theory. In general, molecular dynamics simulation results, as all empirical derivations, are valid only on the range of parameters that were investigated, and this methodology

Figure 1: Effective force in the rigid cylinder approximation. **(A)** Force extension relationship, without crowders ($\phi = 0$) *vs* with crowders ($r/r_0 = 2$ and $\phi = 0.2$). The crowders reduce the effective stretching force by a factor f^*, indicated by the dashed red line. **(B)** Rigid cylinder approximation for the stretched WLC. Neglecting fluctuations, the volume depleted by the chain to crowders is a cylinder of extension z and radius equal to the sum of the cross-radius of the polymer r_0 and the crowder's radius r. **(C)** Collapse force f^* as function of crowder density ϕ, for varied crowder size r. Smaller crowders have a stronger collapsing effect. Values of $f^*/f_c = 1.1$ are indicated by dashed black line; below this the collapsing effect of crowders on the force-extension relationship is small and difficult to detect.

does not discern the theoretical determinants of the range of validity. Furthermore, it is a challenge to improve this type of simulation, since a reduction of r/r_0 by a factor x at fixed ϕ entails an increase of the number of simulated particles of $\mathcal{O}\left(x^3\right)$. Similarly, an increase of the polymer length by a factor x entails, due to scaling of polymer dynamics, an increase of the simulated volume and number of simulated particles of $\mathcal{O}\left(x^{5/2}\right)$. On a more fundamental level, the bead-spring model is not a good polymer to simulate DNA (or DNA-based chromosome fibers), as DNA is a significantly slender polymer with bending stiffness (persistence length) much longer than its cross section, and this makes quantitative matching of parameters in published molecular dynamics simulations to experiments problematic.

Alternately, one carry out molecular pulling experiments with high precision *in vitro*. These types of experiments can be used as a tool for quantitative study of crowding agents, by quantifying how the crowders change the force-extension behavior of DNA, or in principle isolated chromosome fibers. As an example, a recent experiment has shown how crowding affects DNA supercoiling, suppressing its tendency to strand-separate under negative torsional stress [DM25].

Here we present a theory of effects of molecular crowding on the force-extension relation for DNA, or any other semiflexible polymer, in the rigid rod regime. The DNA molecule is represented by a worm like chain (WLC) under pulling stress, described by its tangent vector $\hat{\mathbf{t}}(s)$ over the arc-length coordi-

nate s spanning the full length L of the chain. In a coordinates with cylindrical symmetry around the $\hat{\mathbf{z}}$ pulling direction, we write the energy as the sum of the bending energy of the WLC with bare persistence length A, the usual coupling of the externally imposed pulling force to the molecule extension $z = \int_0^L \mathrm{d}s\, \hat{\mathbf{z}} \cdot \hat{\mathbf{t}}$, and an osmotic term due to the crowders in the solvent. This osmotic contribution is simply the product of the crowder concentration c and the volume made sterically inaccessible to the crowders by the presence of the polymer chain δV:

$$\beta E\left[\hat{\mathbf{t}}(s)\right] = \int_0^L \mathrm{d}s \left[\frac{A}{2}\left(\partial_s\hat{\mathbf{t}}\right)^2 - \beta f \hat{\mathbf{z}} \cdot \hat{\mathbf{t}}\right] + c\,\delta V, \tag{1}$$

for more details on the derivation and underlying assumptions see Appendix A.

In the zero-crowder-concentration limit ($c \to 0$ or equivalently, volume fraction $\phi = vc \to 0$) this becomes the standard WLC model under pulling force [MS95]. The model has a characteristic force $f_c = k_B T/A$ that corresponds to the force at which the Pincus blob radius [Gen79] is equal to the persistence length of the chain. The force f_c corresponds to WLC extension of roughly 50% of its maximum. For pulling forces $f > f_c$, an expansion in inverse powers of f gives z as the full chain extension L, minus a force-dependent term that is proportional to the intensity of the chain fluctuations perpendicular to the pulling force $\langle z \rangle / L = 1 - (4f/f_c)^{-1/2} + \cdots$ as depicted in Fig. 1A, black line.

To calculate δV in Eq. 1, we consider first the approximation where we neglect the effects of the transverse fluctuations of the chain, taking it to be a rigid rod of radius r_0 and elongation z. In this configuration, the volume δV_{cyl} inaccessible to the crowders is a cylinder of radius $R = r_0 + r$ and of the same length as DNA extension, as depicted in Fig. 1B:

$$c\,\delta V\left[\hat{\mathbf{t}}(s)\right] \approx c\,\delta V_{cyl}\left[\hat{\mathbf{t}}(s)\right] = \int_0^L \mathrm{d}s\,\pi c R^2\,\hat{\mathbf{z}} \cdot \hat{\mathbf{t}} \tag{2}$$

Here the crowders are considered to be spheres of radius r, but the theory can be generalized to other shapes of crowding molecules. This is the zeroth-order starting point for dealing with the transverse fluctuations, which modify the volume excluded to the crowders.

The osmotic force acting on the DNA chain as a whole tends to reduce its extension z by acting against the effect of the external force f with a term that has exactly the same dependence on the conformation $\hat{\mathbf{t}}$ as the pulling term (see Eq. 1). This corresponds to an additive reduction of the effective pulling force by a crowder-dependent f^*, as depicted in Fig. 1A, red line, with

$$\beta f^* = \pi c R^2 = \frac{3}{4}\left(1 + \frac{r_0}{r}\right)^2 \frac{\phi}{r}; \tag{3}$$

the dashed red line in Fig. 1A indicates the value of the ordinate corresponding to f^*. All crowding-dependent terms in this rigid cylinder approximation are

in this effective force term $\tilde{f}$:

$$\tilde{f} = f - f^*, \qquad \text{and} \qquad \frac{\langle z \rangle}{L} = 1 - \sqrt{\frac{f_c}{4\tilde{f}}} + \cdots . \qquad (4)$$

This result predicts a chain collapse instability for external forces $f < f^*$, where the net effective pulling force goes to zero. The larger the critical force, the larger is the effect of crowders on the force-extension relationship. The effects are proportional to the crowder volume fraction ϕ, and become more significant for smaller crowder sizes r/r_0, as depicted in Fig. 1C. While we expect the chain to collapse at effective forces $\tilde{f} < 0$, once collapse occurs, we expect higher-order fluctuation terms, effects of DNA looping, surface tension and nematic effects to become important in defining the condensed "phase". These effects are not accounted for by our theory, since its domain of validity is $\tilde{f} > f_*$ (and f_c); in this regime the DNA is extended so these nonlinear effects do not play a role.

We can estimate effects of crowders on DNA in physiological salt conditions, considering that the characteristic force $f_c \simeq 0.08$ pN, and that the radius of DNA is $r_0 \simeq 2$ nm, for crowders of radius $r = 16$ nm it takes a volume density of at around $\phi > 0.4$ to observe relative effects $(f^* - f_c)/f_c$ above 10% (crossing of red and dashed black lines in Fig. 1C), and it takes larger volume densities to see effects at higher forces. For substantially larger crowders, the jamming of the solvent, significant at $\phi > 0.5$, would make it very difficult to see any polymer collapse at any values of concentration and pulling forces. On the other hand, smaller crowders have an observable effect at lower densities, *i.e.* for $r = 3$ nm, we observe a 10% effect already at $\phi > 0.06$. This size of crowders is comparable in size to the radius of proteins and protein complexes in the nucleoplasm: therefore we expect crowding-induced collapse to be relevant *in vivo* where protein concentrations are in the range of $\phi > 0.1$.

We now add the effect of WLC bending fluctuations that were neglected in the previous computation. The polymer, pulled by an effective force $\tilde{f}$ in the $\hat{\mathbf{z}}$ direction, has fluctuations in the $\hat{\mathbf{x}}$ and $\hat{\mathbf{y}}$ directions orthogonal to the force, as depicted in Fig. 2A. The orthogonal displacement of the chain $\mathbf{u}_\perp(s)$, defined by the orthogonal component of the tangent vector $\mathbf{t}_\perp(s)$ through $\partial_s \mathbf{u}_\perp(s) = \mathbf{t}_\perp$ has fluctuations with a force-dependent correlation length $\xi = A/(\tilde{f}/f_c)^{1/2}$ [MS95], or explicitly $\langle \mathbf{t}_\perp(s_0) \cdot \mathbf{t}_\perp(s) \rangle = (\xi/A) \exp[-(s - s_0)/\xi]$ relative to any point s_0 along the polymer. Consider a crowder touching the chain at some particular position s_0: the trajectory of the chain $\mathbf{u}_\perp(s)$ will have a rigid-rod conformation in the direction $\mathbf{t}_\perp(s_0)$ for $|s - s_0| < \xi$, and will then begin to random walk in the transverse directions for larger values of $|s - s_0|$. This configuration will be excluded if this trajectory "overlaps" with the same crowder at a second point away from s_0.

To account for this we consider this effect as an increase of the radius of the cylinder of depletion by an additional fluctuation dependent distance ρ, representing the additional volume inaccessible to the crowders as depicted in

Figure 2: Effects of chain fluctuations. **(A)** The polymer fluctuates perpendicular to the pulling force, generating additional excluded volume δV not accounted for by the rigid cylinder approximation. We account for this by an additional fluctuation dependent radius ρ. **(B)** 1D illustration of calculation of $\langle \rho^2 \rangle$ at fixed r/r_0 and ϕ. Two spherical crowders of different radius touch the polymer at $s = 0$. The non-zero mean component of the WLC trajectory along the $\mathbf{t}_\perp(0)$ direction does not contribute to depletion while standard deviation does. **(C)** Force extension relationship, as in Fig. 1A ($\phi = 0.2$ and $r/r_0 = 2$). Dashed line shows zeroth-order contribution of crowders; solid lines accounts for fluctuation-dependent corrections. **(D)**. Polymer fluctuation-dependent contribution to the force-extension relationship as function of r/A at fixed effective force $(f_c/\tilde{f})^{1/2} = 0.5$, from Eq. 10 (solid lines, red line corresponding to DNA r_0 value). Dashed lines show $\lambda \ll 1$ approximation (Eq. 11).

Fig. 2A. The depletion volume is

$$
\begin{aligned}
c \, \delta V \, \hat{\mathbf{t}}(s) &\approx c \int_0^L \mathrm{d}s \, \pi \left(r_0 + r + \rho \right)^2 \, \hat{\mathbf{z}} \cdot \hat{\mathbf{t}} \\
&\approx c \, \delta V_{cyl} + \pi c L \left[\langle \rho^2 \rangle + 2R \sqrt{\langle \rho^2 \rangle} \right],
\end{aligned}
\tag{5}
$$

where we have used a perturbation expansion of the partition function to first order in crowder volume fraction ϕ.

To calculate $\langle \rho^2 \rangle$ we consider the orthogonal displacement $\mathbf{u}_\perp(s)$ of the WLC conformation over distances comparable to the crowder size r, which should be compared to the correlation length for bending and displacement fluctuations ξ. For small crowders $r < \xi$ care must be taken: the tangent vector has strongly correlated tilt over distances of the crowder size, expressed by a nonzero value of $\langle \mathbf{t}_\perp(0) \cdot \mathbf{u}_\perp(s) \rangle / \langle |\mathbf{t}_\perp(0)|^2 \rangle^{1/2}$. This quantity is the component of orthogonal displacement in the direction of the tangent vector tilt, $\mathbf{t}_\perp(0)/|\mathbf{t}_\perp(0)|$, and can be computed to be

$$
\frac{\langle \mathbf{t}_\perp(0) \cdot \mathbf{u}_\perp(s) \rangle}{\langle |\mathbf{t}_\perp(0)|^2 \rangle^{1/2}} = \int_0^r \mathrm{d}s \frac{\langle \mathbf{t}_\perp(0) \cdot \mathbf{t}_\perp(s) \rangle}{\langle |\mathbf{t}_\perp(0)|^2 \rangle^{1/2}} = \left(\tilde{f}/f_c \right)^{1/4} \xi \left(1 - e^{-s/\xi} \right).
\tag{6}
$$

Subtracting this from $\mathbf{u}_\perp(s)$ yields displacement fluctuations in the tilted reference frame, correctly expressing the area transverse to the polymer that is excluded to the crowders (Fig. 2B).

The fluctuation-corrected effective radius valid for all r follows as

$$
\langle \rho^2 \rangle = \left\langle \left| \mathbf{u}_\perp(r) - \frac{\langle \mathbf{t}_\perp(0) \cdot \mathbf{u}_\perp(r) \rangle \mathbf{t}_\perp(0)}{\langle |\mathbf{t}_\perp(0)|^2 \rangle} \right|^2 \right\rangle,
\tag{7}
$$

which can be written in terms of a dimensionless scaling function $g(\lambda)$ of scaled crowder size $\lambda = r/\xi$:

$$
\begin{aligned}
\langle \rho^2 \rangle &= \int_0^r \mathrm{d}s \int_0^r \mathrm{d}s' \langle \mathbf{t}_\perp(s') \cdot \mathbf{t}_\perp(s) \rangle - \left(\int_0^r \mathrm{d}s \frac{\langle \mathbf{t}_\perp(0) \cdot \mathbf{t}_\perp(s) \rangle}{\langle |\mathbf{t}_\perp(0)|^2 \rangle^{1/2}} \right)^2 \\
&= \frac{\xi^3}{A} g(\lambda), \quad \text{where} \quad g(\lambda) = 2\lambda - 3 + 4e^{-\lambda} - e^{-2\lambda}.
\end{aligned}
\tag{8}
$$

The exclusion radius depends on fluctuations and is force-dependent; for sufficiently large force and large crowders ($r > \xi$ or $\lambda > 1$), $g(\lambda) = 2\lambda + \cdots$, with asymptotic behavior $\langle \rho^2 \rangle = 2k_B T r \xi^2 / A + \cdots$, compatible with the usual WLC transverse size result [MS95]. In the opposite small-crowder limit ($r < \xi$ or $\lambda < 1$), $g(\lambda) = 2\lambda^3/3 + \cdots$ one has $\langle \rho^2 \rangle = 2r^3/(3A) + \cdots$. In this latter $\lambda \to 0$ limit, the force-dependence disappears, compatible with the analysis of transverse fluctuations of a WLC confined to a narrow tube [Odi83] (note that without removal of tangent tilt one has the incorrect small-λ behavior $g(\lambda) \approx \lambda^2$ and $\langle \rho^2 \rangle \approx r^2 \xi / A$, resulting in unphysical force dependence persisting as $r \to 0$, i.e., as the crowders vanish).

The partition function for the WLC including crowders can now be written down, shifting $f \to \tilde{f}$ to account for δV_{cyl}, and adding the excluded volume from (5) including (8):

$$\frac{1}{L} \log Z\left(\lambda, \xi, A\right) = \beta \tilde{f} - \sqrt{\beta \tilde{f}/A} - \pi c \left(\langle \rho^2 \rangle + 2R\sqrt{\langle \rho^2 \rangle}\right) + \mathcal{O}(\tilde{f}^{-3/2}), \quad (9)$$

The force-extension relationship follows as $\langle z \rangle / L = \partial \log Z / \partial(\beta \tilde{f})$, leading to correction of the zeroth-order force extension relationship of Eq. 4 ($z_0/L = 1 - (f_c/[4\tilde{f}])^{1/2}$):

$$\frac{\langle z \rangle}{L} = \frac{z_0}{L} + \frac{3\phi}{8} \left[\frac{\xi^2}{A^2} + \frac{(r_0 + \lambda \xi)\, \xi^{1/2}}{\sqrt{g(\lambda)} A^{3/2}} \right] \frac{3g(\lambda) - \lambda g'(\lambda)}{\lambda^3}, \quad (10)$$

where $\langle z \rangle / L$ is a function of the force-dependent bending fluctuation correlation length ξ, the volume fraction of the crowders ϕ, their size via $\lambda = r/\xi$, the polymer stiffness (persistence length) A, and the bare polymer radius r_0. This result is plotted in Fig. 2C.

The fluctuation correction to $\langle z \rangle / L$ is always positive, since $3g(\lambda) - \lambda g'(\lambda)$ is positive for all λ. We define a characteristic force $f_\lambda = f_c A^2/r^2$ at which $\lambda = 1$, which is the boundary between the two scaling regimes for the fluctuation corrections. For $\tilde{f} \gg f_\lambda$, both $g(\lambda)$ as well as $3g(\lambda) - \lambda g'(\lambda)$ vary $\sim \lambda$. In this regime, the fluctuation correction goes to zero as $\mathcal{O}[(f_c/\tilde{f})^{3/2}]$ asymptotically, as such the polymer's extension always attains its contour length for infinite pulling force. It also monotonically decreases toward zero for large crowders as $\mathcal{O}[(A/r)^{3/2}]$ as is physically reasonable.

In the opposite limit $\tilde{f} \ll f_\lambda$, $g(\lambda)$ varies $\sim 2\lambda^3/3$ while $3g(\lambda) - \lambda g'(\lambda)$ varies $\sim \lambda^4/2$, leading to:

$$\frac{\langle z \rangle - z_0}{L\phi} \simeq \frac{3}{16} \left[\frac{r}{A} + \sqrt{\frac{3}{2}} \left(\frac{r}{A} + \frac{r_0}{A}\right) \sqrt{\frac{A}{r}} \right] \sqrt{\frac{f_c}{\tilde{f}}}. \quad (11)$$

In this regime, the fluctuation dependent correction varies with force as $\sim (f_c/\tilde{f})^{1/2}$, which is the same behavior as z_0/L in $\tilde{f}/f_c$; the fluctuation correction can be interpreted as changing the polymer persistence length A. The bulkier is the polymer (depending on the fraction r_0/A) the greater is the contribution of fluctuations. Notably, as a function of r/A, the function has a minimum for crowders of radius $r \approx r_0$, and increases for larger r/A.

Requiring a smooth match of the two asymptotic behaviors of the function in r/A implies that, at fixed $\tilde{f}/f_c$, the full form of $(\langle z \rangle - z_0)/(L\phi)$ from Eq. 10 has a maximum for the crowders of size close to the force-dependent correlation length $r \approx \xi$. In fact, this is true only for slender enough polymers (when $r_0/A \ll (f_c/\tilde{f})^{1/2}$), as can be seen in Fig. 2D. This also implies that, at high enough forces, the swelling effect of fluctuations generated by small crowders is always larger to the one by larger crowders, although it is so small in absolute terms that it cannot be seen in Fig. 2C.

When we consider the combined effect of the zeroth-order collapse and the fluctuation dependent swelling, strikingly, we predict that overall, crowders can *extend* the chain above the values for $\phi = 0$ if they are large enough such that $f^* < f_c$ and small enough such that the effect of fluctuations is significant (Fig. 2C, brown line), or for very high forces. This unexpected result belies the simple notion that neutral (purely steric) crowders can only compact a polymer. Experiments using Förster resonance energy transfer (FRET) experiments on protein disordered domains indicate that the same crowders can induce either increased folding or unfolding depending on the tested domain [SKB+14]. It is possible that these experimental results are due to physical effects discussed here, with compaction or decompaction dependent on the the size of crowders relative to the polymer bending stiffness, as well as the force applied to the chain. At present quantitative comparison to that experimental data is not possible, since the theory of this paper is valid only in the large pulling force regime (for $f > f_c$) while solution FRET experiments are performed on unstretched polymers.

In this paper we have developed a theory of effects of steric crowders of radius r on the force-extension response of a strongly stretched DNA or other semiflexible polymer, including fluctuation effects through first order in crowder fraction ϕ. Our results predict *in vitro* experiments which could be used as a sensitive probe of crowder effects relevant to compaction of chromosomes or other polymer structures *in vivo*.

As a function of crowder size and volume fraction ϕ we find that there is an additive shift of net force on the polymer by $f^* \approx k_B T \phi / r$ which can drive a collapse against external force, plus changes in the high-force response including a regime where the apparent persistence length is changed. These effects will be important to experimentalists since the force exerted by the apparatus will no longer be the net tension stretching the polymer once crowders are present. For example, at fixed external force f, addition of crowders can lead to a shift down of the apparent force and enhanced transverse fluctuations in experiments that calibrate force via fluctuations (e.g., magnetic tweezers [DM25]), or alternately a shift up in force in cantilever-type experiments that essentially maintain constant extension (e.g., optical tweezers, atomic force microscopes). For optical tweezers, attention should also be paid to constant-force system dynamics, as well as to laser heating and trapping effects on the crowders themselves. Exploration of the different regimes of crowder size and force relative to the force characterizing the boundary between different fluctuation effect scaling, $f_\lambda \approx k_B T A / r^2$, can modulate these basic effects and in some cases reverse them.

Our analysis uses approximations that need to be considered carefully when comparing with experiments. We use the limit of small crowder volume fraction when computing osmotic pressure, and the perturbation expansion for the WLC partition function, using the strong stretching regime $\langle z \rangle > L/2$ where bending fluctuations are relatively small. Considering the broad range of forces over which we predict quantitative and qualitative effects, our calculations should be useful in understanding *in vivo* compaction of bacterial nucleoids

and other intracellular structures. For instance, while it is accepted that RNA and DNA polymerases, structural maintenance of chromosomes complexes, and other molecular motors exerting forces on DNA can change chromosome condensation due to motor effects [CW24], our theory demonstrates that effective forces are sensed by a polymer, in response to varied crowder size and volume fraction.

We also observe condensation effects without any globular phase for physiological tensions, which raises the question of whether the bacterial nucleoid is condensed on its own, or rather is decondensed with size dependent on activities and local environment. These two options are not necessarily mutually exclusive, since it is possible to have a polymer coexist in an activity induced decondensed phase as well as more compacted regions in a different genomic-scale compartment, consistent with results of Hi-C and ATACseq genomic experiments on human cell lines [BGZ+13, LvBW+09].

Consideration of *in vivo* conditions is complicated by the fact that the cellular environment is not simply a solution of spherical monodisperse crowders. More specifically the intracellular environment possesses *(i)* a high degree of polydispersity, *(ii)* heterogeneity of genomic domains in terms of binding proteins and their post-translational modifications, and *(iii)* presence of molecular activities in the cytoplasm. These factors are an objective for further development of the theory of the present paper to understand experimental results [CCG+24].

Acknowledgments

The authors acknowledge helpful discussions with Quentin Chaboche, Marco Cosentino-Lagomarsino and Sumitabha Brahmachari. VFS acknowledges Laura Finzi, David Dunlap, Jin Qian, and Narendar Kolimi for sharing preliminary optical trap force-extension data. This work was supported by the Centre National de la Recherche Scientifique (CNRS), France (VFS), and by the NIH (US) through grant R01-GM105847 (JFM). JFM and VFS also acknowledge travel support from the CNRS "Physics Meets Genomes" Conference and Summer School in Cargese (France) where this work was initiated.

Appendix A

The effective energy in Eq. 1 can be derived from a Hamiltonian that considers the configuration of the crowders in space. We add an interaction term between each crowder of position $\mathbf{x}_i$ and the polymer chain to a classical WLC energy for the polymer chain. Individual crowders are indexed by i spanning from 1 to N, such that the concentration of crowders is $c = N/V$, where V is the full experimental volume. This results in:

$$\beta E \left[\hat{\mathbf{t}}(s), \{\mathbf{x}_i\} \right] = \beta E_{ch} + \beta E_{cro} \tag{12}$$

with

$$\beta E_{ch}\left[\hat{\mathbf{t}}(s)\right] = \int_0^L ds \left[\frac{A}{2}\left(\partial_s \hat{\mathbf{t}}\right)^2 - \beta f \hat{\mathbf{z}} \cdot \hat{\mathbf{t}}\right], \tag{13}$$

and

$$\beta E_{cro}\left[\hat{\mathbf{t}}(s), \{\mathbf{x}_i\}\right] = \int_0^L ds \left[\sum_i^N U\left(\mathbf{x}_i - \mathbf{u}\right)\right], \quad \text{with} \quad \frac{\partial \mathbf{u}(s)}{\partial s} = \hat{\mathbf{t}}; \tag{14}$$

this is valid for low crowder densities (low enough ϕ), since it ignores two-body interactions between crowders, as well as higher orders; further, this is valid for overdamped conditions, when the inertial mass of the crowders and the chain are both negligible. Given monodisperse crowders of radius d, we set the steric interaction potential as:

$$U(\mathbf{x}) = \begin{cases} \infty & \text{if } |\mathbf{x}| < d \\ 0 & \text{otherwise.} \end{cases} \tag{15}$$

we can then derive the corresponding partition function from the Gibbs distribution, that is a path integral over all worm like chain configurations, integrated over all the positions of the crowders over the volume V:

$$Z(f, \beta, N, d, L) = \int \mathcal{D}\left[\hat{\mathbf{t}}(s)\right] \prod_i \int_{V \times N} d\mathbf{x}_i \exp\left[-\beta E\right]. \tag{16}$$

We then integrate the partition function over the crowder degrees of freedom $\{\mathbf{x}_i\}$. This allows calculation of the reduction in configuration space available due to the interaction with the polymer chain, i.e., the strength of depletion interactions. Considering the shape of the potential $U(\mathbf{x})$, the problem reduces to calculating the reduction of volume accessible to the crowders $\delta V\left(\hat{\mathbf{t}}(s)\right)$. In the specifics,

$$\frac{Z'\left[\hat{\mathbf{t}}(s)\right]}{Z_{cro}(N)} = \exp\left[N \log\left(1 - \frac{\delta V}{V}\right)\right] \approx \exp\left[-\frac{N \, \delta V}{V}\right] = e^{-c \, \delta V} \tag{17}$$

that is the van't Hoff entropy approximation [Mar04], valid for $\delta V \ll V$. More realistic forms of $U(\mathbf{x})$ would lead to a similar result. Of important note, this approximation not only simplifies the calculation, but also assures that c is not affected by variations in volume δV. Substituting, we obtain:

$$Z(f, \beta, c, r, L) = \int \mathcal{D}\left[\hat{\mathbf{t}}(s)\right] e^{-c \, \delta V} \exp\left[-\beta E_{ch}\right] = \int \mathcal{D}\left[\hat{\mathbf{t}}(s)\right] \exp\left[-\beta E_{eff}\right]. \tag{18}$$

as defined in Eq. 1.

References

[AAL$^+$02]   Onofrio Annunziata, Neer Asherie, Aleksey Lomakin, Jayanti Pande, Olutayo Ogun, and George B. Benedek. Effect of polyethy-

lene glycol on the liquid–liquid phase transition in aqueous protein solutions. *Proc Natl Acad Sci U S A*, 99(22):14165, October 2002.

[AO54] Sho Asakura and Fumio Oosawa. On Interaction between Two Bodies Immersed in a Solution of Macromolecules. *The Journal of Chemical Physics*, 22(7):1255–1256, July 1954.

[BBD+12] Vincenzo G. Benza, Bruno Bassetti, Kevin D. Dorfman, Vittore F. Scolari, Krystyna Bromek, Pietro Cicuta, and Marco Cosentino Lagomarsino. Physical descriptions of the bacterial nucleoid at large scales, and their biological implications. *Reports on Progress in Physics. Physical Society (Great Britain)*, 75(7):076602, July 2012.

[BGZ+13] Jason D. Buenrostro, Paul G. Giresi, Lisa C. Zaba, Howard Y. Chang, and William J. Greenleaf. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nature Methods*, 10(12):1213–1218, December 2013.

[CCG+24] Quentin Chaboche, Gerardo Campos-Villalobos, Giuliana Giunta, Marjolein Dijkstra, Marco Cosentino Lagomarsino, and Vittore F. Scolari. A mean-field theory for predicting single polymer collapse induced by neutral crowders. *Soft Matter*, February 2024.

[CW24] Zhiyu Cao and Peter G. Wolynes. Motorized chain models of the ideal chromosome. *Proceedings of the National Academy of Sciences of the United States of America*, 121(28):e2407077121, July 2024.

[DM25] Parth Rakesh Desai and John F. Marko. Molecular crowding suppresses mechanical stress-driven DNA strand separation. *Biophysical Journal*, 124(12):1886–1890, June 2025.

[Gen79] P. G. de Gennes. *Scaling Concepts in Polymer Physics*. Cornell University Press, 1979.

[IIT+24] Shiori Iida, Satoru Ide, Sachiko Tamura, Masaki Sasai, Tomomi Tani, Tatsuhiko Goto, Michael Shribak, and Kazuhiro Maeshima. Orientation-independent-DIC imaging reveals that a transient rise in depletion attraction contributes to mitotic chromosome condensation. *Proc Natl Acad Sci U S A*, 121(36):e2403153121, September 2024.

[JJH16] Chanil Jeon, Youngkyun Jung, and Bae-Yeun Ha. Effects of molecular crowding and confinement on the spatial organization of a biopolymer. *Soft Matter*, 12(47):9436–9450, 2016.

[JNCP16]   Andreia F. Jorge, Sandra C.C. Nunes, Tânia F.G.G. Cova, and Alberto A.C.C. Pais. Cooperative action in DNA condensation. *Current Opinion in Colloid & Interface Science*, 26:66–74, December 2016.

[KPHT15]   Hongsuk Kang, Philip A. Pincus, Changbong Hyeon, and D. Thirumalai. Effects of macromolecular crowding on the collapse of biopolymers. *Physical Review Letters*, 114(6):068303, February 2015. arXiv:1409.5663 [cond-mat, physics:physics, q-bio].

[LvBW+09]   Erez Lieberman-Aiden, Nynke L. van Berkum, Louise Williams, Maxim Imakaev, Tobias Ragoczy, Agnes Telling, Ido Amit, Bryan R. Lajoie, Peter J. Sabo, Michael O. Dorschner, Richard Sandstrom, Bradley Bernstein, M. A. Bender, Mark Groudine, Andreas Gnirke, John Stamatoyannopoulos, Leonid A. Mirny, Eric S. Lander, and Job Dekker. Comprehensive Mapping of Long-Range Interactions Reveals Folding Principles of the Human Genome. *Science*, 326(5950):289–293, October 2009.

[Mar04]   Geoffrey Martin. On van't Hoff's Law of Osmotic Pressure. *Nature*, 70(1822):531–532, September 1904.

[MS95]   John F. Marko and Eric D. Siggia. Stretching DNA. *Macromolecules*, 28(26):8759–8770, December 1995.

[Odi83]   Theo Odijk. The statistics and dynamics of confined or entangled stiff polymers. *Macromolecules*, 16(8):1340–1344, August 1983.

[Odi98]   Theo Odijk. Osmotic compaction of supercoiled DNA into a bacterial nucleoid. *Biophysical Chemistry*, 73(1):23–29, July 1998.

[PHH+12]   James Pelletier, Ken Halvorsen, Bae-Yeun Ha, Raffaella Paparcone, Steven J. Sandler, Conrad L. Woldringh, Wesley P. Wong, and Suckjoon Jun. Physical manipulation of the Escherichia coli chromosome reveals its soft nature. *Proceedings of the National Academy of Sciences*, 109(40):E2649–E2656, October 2012.

[RD15]   Sravani Keerthi Ramisetty and Rita S. Dias. Synergistic role of DNA-binding protein and macromolecular crowding on DNA condensation. An experimental and theoretical approach. *Journal of Molecular Liquids*, 210:64–73, October 2015.

[RJS23]   Romain Rollin, Jean-François Joanny, and Pierre Sens. Physical basis of the cell size scaling laws. *eLife*, 12:e82490, May 2023.

[SBdH+15]   Tyler N. Shendruk, Martin Bertrand, Hendrick W. de Haan, James L. Harden, and Gary W. Slater. Simulating the Entropic Collapse of Coarse-Grained Chromosomes. *Biophysical Journal*, 108(4):810–820, February 2015. arXiv:1407.2850 [cond-mat, physics:physics, q-bio].

[SKB⁺14]   Andrea Soranno, Iwo Koenig, Madeleine B. Borgia, Hagen Hofmann, Franziska Zosel, Daniel Nettels, and Benjamin Schuler. Single-molecule spectroscopy reveals polymer effects of disordered proteins in crowded environments. *Proceedings of the National Academy of Sciences of the United States of America*, 111(13):4874–4879, April 2014.

[YMRM20]   Da Yang, Jaana Männik, Scott T. Retterer, and Jaan Männik. The effects of polydisperse crowders on the compaction of the *Escherichia coli* nucleoid. *Molecular Microbiology*, 113(5):1022–1037, May 2020.