How to Become Invisible to Mosquitoes:
A Computational Study of Host Signal Collapse

Zi-Ru JIANG*

Independent Researcher, Nagoya, Japan
Email: ziru.jiang@gmail.com
ORCID: https://orcid.org/0000-0003-1378-1646

* Corresponding Author

ABSTRACT

Mosquitoes locate vertebrate hosts by following coherent gradients of carbon dioxide, heat, and humidity. Nevertheless, a persistent empirical observation is that ordinary electric fans drastically reduce mosquito bites, even in the absence of chemical repellents. Here, we introduce a physics-guided, agent-based simulation demonstrating that mosquito host-seeking is not chemically inevitable but physically fragile. The model integrates directional airflow, vortex-induced chaotic mixing, and thermal decoy fields, and tracks 250 autonomous mosquito agents over 600 time-steps under Monte-Carlo sampling without fixed random seeds. Three aerodynamic mechanisms were quantified. Mild upward airflow already reduces successful host localization by more than 80%, despite not mechanically preventing flight. Increasing vortex circulation produces a continuous phase transition, collapsing localization probability below 1% at moderate strength. Thermal decoys cause only a linear dilution of success, indicating misinformation alone cannot trigger collapse. A two-dimensional phase map reveals a robust "invisibility region" where airflow and vortex perturbations interact multiplicatively, eliminating host detection even when neither factor individually reaches threshold. These results show that mosquito host-seeking operates as a gradient-dependent failure system: once scalar fields lose spatial coherence, navigation collapses independent of sensory capability. This provides a quantitative theoretical basis for fan-mediated mosquito protection and suggests that non-chemical, low-energy airflow strategies can induce physical invisibility to hematophagous insects.

Keywords: Mosquito host-seeking; Aerodynamic invisibility; Vortex-induced chaos; CO2 plume disruption; Agent-based simulation

1. INTRODUCTION

Mosquitoes remain the world's deadliest animals, responsible for more than 700,000 human deaths each year through malaria, dengue, chikungunya and other vector-borne diseases (WHO, 2023). Although long-lasting insecticidal nets (LLINs), indoor residual spraying (IRS), and emerging gene-drive or Wolbachia-based approaches have significantly reduced transmission, global malaria burden has plateaued since 2015 and is now resurging across high-transmission regions (Bhatt et al., 2015; Lawal et al., 2024; Weiss et al., 2025; Umugwaneza et al., 2025). Widespread pyrethroid resistance, declining insecticide durability, and escalating operational costs increasingly indicate that chemical or genetic interventions alone are insufficient for achieving long-term elimination (Ranson & Lissenden, 2016).

In parallel, there is growing interest in low-cost, non-chemical environmental manipulation, particularly airflow. Field studies have shown that ordinary household fans can substantially reduce mosquito landings and biting indoors, even without repellents, apparently by disrupting host-seeking behavior (Sutcliffe & Yin, 2021; Carrasco-Tenezaca et al., 2023). However, airflow remains absent from global vector-control frameworks, and major reviews still treat it as

a laboratory nuisance rather than a behaviorally meaningful suppressor (Kline, 2006). This persists despite convergent experimental evidence demonstrating that airflow significantly modifies mosquito navigation, sensory integration, and plume detection.

Mechanistically, airflow impacts host seeking on several sensory and physical levels. Wind-tunnel experiments show that airflow distorts CO_2 plume topology, filament intermittency, and gradient stability, thereby impairing chemotactic tracking and heat localization (Cooperband et al., 2006; McMeniman et al., 2014; Sutcliffe & Yin, 2021; Carrasco-Tenezaca et al., 2023; Hinze et al., 2021). Complementing these findings, Cribellier et al. (2018) demonstrated that even in the absence of bulk wind, local recirculation vortices around odour-baited traps fragment odor filaments, generate intermittent "lost–found–lost" trajectories, and induce repeated reorientation events—direct behavioral evidence that small-scale turbulence alone can degrade plume coherence and destabilize host-approach paths. Beyond olfaction, recent work demonstrates that thermal and olfactory cues act jointly, with human scent strongly shaping mosquito thermotaxis via conserved sensory pathways (Giraldo et al., 2023). These findings integrate with broader multisensory reviews indicating that mosquitoes use a rich suite of olfactory, thermal, hygrosensory, visual, and inertial cues to navigate complex environments (Montell, 2025). Even background CO_2 concentration can significantly alter search performance, pointing to a critical role for plume geometry, not just odor intensity (Majeed et al., 2014).

Despite these advances, airflow has typically been treated experimentally as binary "low vs. high" perturbations, without quantifying why host detection sometimes declines gradually whereas in other cases it collapses abruptly. Three fundamental questions therefore remain unresolved: What determines the collapse threshold at which host detection probability falls below 1%? Do directional airflow and vortex turbulence suppress host seeking through distinct mechanisms? Do airflow and thermal misinformation combine additively or multiplicatively?

Recent developments in plume physics and search theory conceptualize mosquito host seeking as an information-limited gradient-climbing process, rather than a deterministic olfactory response. Similar to bacterial chemotaxis and theoretical infotaxis (Vergassola et al., 2007), mosquitoes track scalar fields whose structure is highly vulnerable to turbulence. Stretch–fold–shear cascades can destroy scalar coherence faster than insects can integrate temporal cues (Shraiman & Siggia, 2000). Under this framework, airflow may render a host functionally invisible despite mosquitoes retaining full flight ability and intact sensory systems—a hypothesis directly supported by indoor cross-flow experiments showing sharp behavioral collapse at surprisingly low wind speeds (Sutcliffe & Yin, 2021).

Although prior studies have independently examined directionality (Hoffmann & Miller, 2002), ventilation-driven suppression (Carrasco-Tenezaca et al., 2023), or turbulence-induced fragmentation (Lei et al., 2023), no unified mechanistic simulation has yet integrated directional airflow, vortex mixing, and thermal misinformation under a common physics-based ruleset. As a result, the field lacks an integrative explanation for the nonlinear—sometimes catastrophic—failure of mosquito host seeking in disturbed airflow environments.

To address this gap, we developed a physics-guided, agent-based simulation integrating: (i) 2D CO_2 and heat scalar fields governed by a $1/(1+r)$ decay model; (ii) 250 autonomous mosquito agents performing noisy gradient-biased walks; (iii) three aerodynamic perturbation modules—uniform drift, vortex-induced mixing, and thermal-decoy fields; (iv) 30 Monte-Carlo repetitions per parameter set (>2.4 million total flights). This unified framework enables quantification of collapse thresholds, comparison of aerodynamic suppression modes, and identification of how turbulence, directionality, and thermal misinformation interact to determine whether a host remains detectable—or becomes aerodynamically invisible.

2. METHODS

2.1. Model Overview

We developed an agent-based navigation model to quantify how airflow and vortex-like perturbations disrupt odor–heat guided host-seeking flight. The model simulates N mosquitoes moving within a 2D arena toward a stationary host that emits a chemothermal gradient. At each discrete time step, individuals update their heading based on the combined influence of (i) deterministic CO2–heat gradients, (ii) deterministic aerodynamic forcing (wind or vortex), and (iii) stochastic sensory noise. To focus solely on navigational information, the model does not include mechanical blow-off or loss of flight stability; all mosquitoes remain fully flight-capable under all airflow intensities, consistent with classical observations that mosquitoes can sustain upwind flight at >1–1.5 m□s□¹ when odor cues remain intact (Kennedy, 1940; Gillies & Wilkes, 1970). The main output variable is the proportion of mosquitoes that reach the host region within a fixed time horizon. All simulations are performed using the Python implementation in *mos_phase_models_v19.py* (mos_phase_models_v19). This architecture isolates deterministic environmental structure from stochastic sensory uncertainty, enabling clear identification of airflow-induced collapses.

2.2. Spatial Domain and Host Field Construction

A square lattice of size GRID × GRID (default 256×256) defines the navigable space. The host is positioned at the center (HOST = [GRID/2, GRID/2]). Two scalar fields are precomputed following the script's build_fields () function:

Thermal field:

$$T(r) = \frac{1}{1+r}$$

CO2-like odor field:

$$C(r) = \frac{1}{1+r}$$

Combined attractant:

$$S(x, y) = T(x, y) + C(x, y)$$

Centered finite differences:

$$g_x = S(x+1, y) - S(x-1, y), g_y = S(x, y+1) - S(x, y-1)$$

implemented as:

gx = total[x+1, y] - total[x-1, y]
gy = total[x, y+1] - total[x, y-1]
grad_move = np.stack([gx, gy], axis=1)

The gradient term provides the deterministic chemothermal drive.

2.3. Deterministic Aerodynamic Forcing

Two airflow regimes were examined independently.

(a) Uniform Wind Field

A spatially uniform advective field from calc_wind_velocity():

$$v_x = 0, v_y = -w$$

(b) Vortex Field

Rotational airflow from calc_vortex_velocity ():

$$\mathbf{u} = \frac{\mathbf{p} - \mathbf{H}}{\| \mathbf{p} - \mathbf{H} \|}, \mathbf{t} = (-u_y, u_x), \mathbf{F}_v = v\mathbf{t}$$

The vortex field represents a simplified, deterministic analogue of the recirculation zones and odor-filament fragmentation documented around odour-baited traps (Cribellier et al., 2018), where small-scale rotational flow causes intermittent plume contact and repeated orientation failures. Both airflow components are strictly deterministic; no random gusts or turbulence realizations are included, allowing threshold behavior to be attributed solely to aerodynamic structure.

2.4. Sensory Noise

Sensory noise is modeled as additive Gaussian perturbation applied to the movement
vector:
noise_move = SENSORY_NOISE * rng.normal(size=grad_move.shape)
This is the only source of stochasticity; no randomness exists in the airflow or the scalar
fields.
2.5. Movement Update Rule
At each time step:

$$\mathbf{v} = \nabla S + \mathbf{F}_{air} + \eta$$

Movement is direction-dominant:
move = grad_move + flow_move + noise_move
step = np.sign(move)
pos = pos + step
Positions are clipped within the arena. Airflow alters heading direction only; it does not
increase step magnitude, ensuring that collapse reflects loss of navigational information rather
than forced kinematics.
2.6. Initialization and Success Criterion
At the start of each simulation, $N_{MOSQ} = 200$ mosquito agents are initialized at random
positions within the 2D arena:

$$pos = rng.integers\ (0, GRID, size = (n\_mosq, 2))$$

160 Each simulation proceeds for a maximum of 1,500 discrete time-steps, unless all agents
161 have either reached the host or exited the effective search region. An individual mosquito is
162 classified as successful if, at any time during the simulation, its position enters a circular target
163 zone of radius 8 pixels centered on the host location.
164   The success rate for a single simulation is defined as:

$$\text{Success} = \frac{N_{\text{hit}}}{N_{\text{MOSQ}}}$$

where N_{hit} denotes the number of agents that reached the host zone.
2.7. Monte-Carlo Protocol and Phase Scans
To characterize the effect of airflow parameters, we conducted Monte-Carlo simulations
across a predefined parameter space. For each wind or vortex intensity level in the interval [0, 3]
with increments of 0.1, we performed 50 independent realizations, each using a distinct but
reproducible random seed to ensure statistical robustness.
For a given airflow parameter w (wind) or v (vortex), the mean success probability is
computed as:

$$\bar{S}(w) = \frac{1}{50} \sum_{i=1}^{50} S_i(w)$$

where $S_i(w)$ is the success rate observed in the i-th simulation. The ensemble-averaged
values $\bar{S}(w)$ and $\bar{S}(v)$ were used to construct the phase diagrams presented in the main text.
2.8. Rationale for Model Structure
The model isolates airflow–gradient interactions while minimizing biological and
computational complexity. Key simplifying assumptions (kept minimal): 2D navigation (indoor
CO2 plumes are quasi-planar); Static scalar fields; no temporal fluctuations; No inter-agent
interactions; No additional sensory modalities (visual, humidity, skin volatiles); No mechanical
blow-off, tumbling, or flight loss.
These assumptions ensure that observed failures originate from physical destruction or
displacement of the chemothermal gradient, not from extraneous behavioral mechanisms. This
abstraction is justified by empirical evidence showing that turbulence-induced
intermittency—not mechanical impairment—is the primary driver of navigation failure near
vortex-dominated flows (Cribellier et al., 2018; Sutcliffe & Yin, 2021).
186

3. RESULTS

Figure 1. Effect of wind strength on mosquito navigation success.
Mean host-detection probability declines sharply from 0.34 at zero airflow to 0.06 at wind
strength w=0.3, representing an 82% reduction in successful host localization. Beyond w>1.0,
performance plateaus at ≈0.05, indicating that directional drift displaces the plume without fully
destroying gradient topology. The red dashed line denotes the aerodynamic invisibility threshold
(P_{detect}<0.01). Error bars show standard deviation from 30 Monte-Carlo runs.
195

3.1. Weak wind perturbation triggers an abrupt navigational collapse
Across all tested wind strengths, host-seeking success exhibited a highly nonlinear,
threshold-like drop (Fig. 1). A minimal airflow of w=0.3 caused an 82% reduction (0.34→0.06),
and success reached a low plateau near ~0.045 above w≈1.0. The fact that collapse happens at
very weak airflow agrees with wind-tunnel experiments showing that *Aedes aegypti* approach
rates can fall by 60–70% at only 0.2–0.3 m/s (Sutcliffe & Yin, 2021).
Importantly, mosquitoes in these experiments showed no mechanical impairment of flight,
consistent with classic observations that mosquitoes can still maintain stable upwind flight at >1
m/s when odor cues remain intact (Kennedy, 1940; Gillies & Wilkes, 1970). The abruptness
therefore reflects loss of navigational information, not loss of aerodynamic capability.
Convergence tests (Figs. S1–S2) verify that the collapse is not caused by simulation noise
or insufficient runtime. Success-time distributions (Fig. S6) further reveal that the rare successes
under wind occur almost exclusively during early steps (10–20), resembling the isolated, chance
encounters occasionally reported in behavioral assays when plume structure becomes
intermittent (Sutcliffe & Yin, 2021; Carrasco-Tenezaca et al., 2023).

Together, the numerical results and empirical data indicate that weak directional airflow is
sufficient to push the system past a reliability threshold where gradient-following behavior
collapses abruptly.
214

Figure 2. Vortex-induced collapse of mosquito host detection.
Success probability decreases monotonically with vortex strength and crosses the collapse
threshold at v≈2.0, where mean detection falls to 0.009. Unlike wind, no plateau is observed,
indicating a true continuous phase transition. The mechanism is consistent with scalar
stretch–fold destruction predicted by turbulent plume theory (Shraiman & Siggia, 2000). The
red arrow marks the critical transition point.
222

3.2. Vortex flow induces a continuous, second-order–like transition

223
Vortex forcing produced a fundamentally different pattern: success declined smoothly with
v, crossing the collapse threshold at v≈2.0 (Fig. 2). No plateau appeared, and success continued
trending toward zero with increasing vortex strength. This gradual decay contrasts with the
step-like drop under wind and resembles scalar mixing and filament stretching observed in
fluid-mechanics studies of odor plumes (Sutcliffe & Yin, 2021; Carrasco-Tenezaca et al., 2023).
Cribellier et al. (2018) showed similar behavior around odour-baited traps, where
small-scale recirculation vortices fragmented odor filaments into intermittent packets and
induced repeated "lost–found–lost" trajectories. This pattern represents the same progressive
degradation of plume coherence reproduced by our vortex simulations.
Human CO2 Schlieren reconstructions show similar vortex-driven intermittency in real
indoor environments (Sutcliffe & Yin, 2021; Carrasco-Tenezaca et al., 2023), and our
simulations reproduce this continuous degradation rather than a step-transition. Temporal and

population-size robustness tests (Figs. S1, S3) confirm that the collapse point is stable across
different sampling regimes.
Success-time distributions (Fig. S6) also show that vortex-affected individuals maintain
longer trajectories than wind-affected individuals at comparable suppression levels, consistent
with shear-driven dispersion rather than forced directional drift. This agrees with laboratory
plume observations showing that vortex-like motions fragment odor filaments into intermittent
packets rather than biasing motion (Geier et al., 1999).
Thus, vortex perturbations induce a progressive erosion of gradient coherence, producing a
continuous, second-order–like navigational decline.
245

247   Figure 3. Effect of thermal decoys on mosquito host seeking.
248   Introducing 0–8 heat-only decoy sources produces a linear decrease in success probability
249 from 0.34 to 0.27, without any collapse boundary. This demonstrates entropy dilution, not
250 gradient destruction: the host remains the global maximum in the scalar field. Error bars
251 indicate SD across 30 Monte-Carlo simulations.
252
3.3. Thermal decoys dilute detection probability without causing collapse
Adding up to eight thermal decoys produced only a linear reduction in host-detection
success (0.34→0.27), with no abrupt failure (Fig. 3). Standard deviations remained below 0.03,
indicating that the decline primarily reflects increased choice entropy, not plume destruction.
This pattern aligns with empirical reports that heat-only traps attract mosquitoes but cannot
prevent human detection unless airflow or odor disruption is also present (Takken & Knols
1999; Cribellier et al., 2018). In Cribellier et al. (2018), heat enhanced approach only when odor
cues remained coherent; when odor coherence was degraded by local vortex motions, heat alone
did not produce stable upwind orientation—matching our finding that thermal clutter weakens
contrast but does not induce collapse.In experimental hut studies, thermal cues enhance

attraction but do not dominate CO2 without accompanying airflow modifications (Mmbando et
al., 2022).
Robustness tests (Figs. S3–S4) show that adjusting the weighting of CO2 versus heat or
increasing the agent count does not change this linear trend. In all conditions, the host remains
the global scalar maximum, meaning the navigational framework remains intact.
Thus, temperature clutter alone cannot induce navigational collapse; it merely weakens the
effective contrast of the scalar landscape.
270

272　　　Figure 4. Combined aerodynamic invisibility phase map under wind–vortex superposition.
273 Color scale shows mean detection probability across 121 parameter combinations. The
274 collapse boundary follows v+0.8w≥2.0, confirmed by the red dashed line (< 1% region). The
275 result reveals multiplicative disruption: neither wind nor vortex must reach its individual critical
276 value if both are present. Contour lines indicate equal-detection regions.
277

3.4. Combined perturbations reveal a robust aerodynamic invisibility region
When wind and vortex forcing were combined, a two-dimensional collapse surface
emerged (Fig. 4). The collapse region is well described by the linear boundary:

Moderate wind and moderate vortex were sufficient for full collapse even when each
perturbation alone was subcritical (e.g., v=1.6, w=0.5). This interaction resembles multi-factor
plume failure regimes reported in robotics and chemotaxis search theory, where directional drift
and rotational mixing jointly degrade search reliability more strongly than either alone.
This combined-failure pattern is consistent with architectural and ventilation studies
showing that airflow geometry—not just speed—can strongly influence mosquito entry

(Carrasco-Tenezaca et al., 2023; Francisco & Watanabe, 2024). Practical claims that multiple
airflow sources (e.g., opposing fans) may outperform a single high-speed stream are common in
field settings, though systematic experimental validation is still lacking. To our knowledge, no
controlled study has directly compared opposing versus unidirectional fan configurations in
mosquito behavior.
Robustness analyses (Figs. S1–S6) confirm that the collapse boundary persists across
population sizes, sensory noise levels, and runtime durations, demonstrating that this
"aerodynamic invisibility zone" is a stable consequence of airflow-driven plume degradation.
These distinct collapse modes—step-like suppression under wind, continuous decay under
vortex flow, linear dilution from thermal clutter, and the true invisibility zone arising from
combined perturbations—are synthesized in **Table 1**.
298
299 **Table 1 Summary of Computational Findings**

Perturbation Type	Behavior	Collapse Threshold	Empirical Consistency
Wind only	Step-like suppression	w≈0.3	Matches reductions at 0.1–0.4 m/s (Sutcliffe & Yin, 2021)
Vortex only	Continuous decline	v≈2.0	Consistent with CO_2 intermittency (Sutcliffe & Yin, 2021; Cribellier et al. 2018; Carrasco-Tenezaca et al., 2023)
Thermal decoys	Linear dilution	None	Matches heat-only trap behavior (Takken & Knols 1999; Cribellier et al. 2018)
Wind + vortex	True invisibility zone	v+0.8w≥2	Matches fan + ventilation studies (Carrasco-Tenezaca 2023)

300   Note: Collapse thresholds correspond to $P_{success} \leq 0.01$ and summarize qualitative
301 behaviors across all parameter sweeps.
302
Across all tests, collapse thresholds remained stable, indicating that the aerodynamic
invisibility phenomenon arises from airflow-driven destruction or dilution of chemothermal
gradients rather than numerical artifacts.

4. DISCUSSION

4.1. Physical meaning of the navigational collapse
Our simulations show that mosquito host detection undergoes a reproducible, abrupt
collapse under structured airflow. This pattern resembles a phase-transition–like breakdown of
navigational reliability rather than a smooth degradation. Directional wind primarily induces
bias displacement—it shifts the apparent position of the scalar maximum without destroying
gradient continuity. Behavioral assays similarly report that *Aedes aegypti* can orient upwind
under modest flow but exhibit reduced trajectory precision and intermittent success (Gibson &
Torr, 1999; Spitzen et al., 2013).
In contrast, vortex forcing generates topological destruction of the plume. Rotational shear
stretches and folds odor filaments at rates that exceed diffusive smoothing and the temporal
integration limits of insect olfaction. This mechanism is predicted by scalar mixing theory
(Shraiman & Siggia, 2000) and directly confirmed by Schlieren observations of human CO_2
plumes, which exhibit intermittent fragmentation above mosquito sensory integration thresholds
(Sutcliffe & Yin, 2021; Carrasco-Tenezaca et al., 2023). Cribellier et al. (2018) further provide
behavioral evidence for this mechanism: odor-baited traps surrounded by small-scale
recirculation vortices generate highly intermittent "lost–found–lost" trajectories, demonstrating
that rotational mixing disrupts plume coherence even without strong bulk wind. Their findings

offer a direct biological analogue to the continuous, second-order–like decay produced in our vortex simulations.

Thus, wind displaces information, whereas vortices destroy information. The combined perturbation regime—co-occurring drift and rotational shear—lies beyond compensation by any known sensory or motor strategy. This agrees with the emerging view that olfactory failure in flying insects is fundamentally *physics-limited* rather than *sensory-limited* (van Breugel et al., 2015; Szyszka et al., 2023).

4.2. Why household fans provide real protection

The common belief that "mosquitoes cannot fly against wind" is incomplete. *Aedes* and *Anopheles* routinely sustain upwind flight at 1–1.5 m s^{-1} when odor cues remain coherent (Kennedy, 1940; Gillies & Wilkes, 1970; Cooperband et al., 2006). Therefore, mechanical blow-off cannot explain why ordinary table fans consistently reduce biting.

Instead, airflow destroys the CO_2 + heat plume as coherent information. Wind-tunnel studies show ≥80% reduction in landings once airflow disrupts odor filament structure (van Breugel et al., 2015; Sutcliffe & Yin, 2021). Our model reproduces this effect: a wind of w≈0.3 (~0.3 m s^{-1}) produces an 82% collapse in host detection, matching empirical measurements.

The continuous-degradation component may arise from vortex-like motions naturally generated by household fans. Cribellier et al. (2018) showed that even modest recirculating vortices fragment odor filaments into intermittent packets, preventing stable upwind tracking. This provides a mechanistic explanation for anecdotal field reports that multiple smaller fans—or opposing fans—sometimes outperform a single strong fan: the added vortex shear pushes the system across the stretch–fold threshold required for plume annihilation.

This behavior is consistent with indoor CO_2 plume reconstructions (Sutcliffe & Yin, 2021; Carrasco-Tenezaca et al., 2023), architectural airflow measurements (Francisco & Watanabe, 2024), and household fan efficacy studies showing ventilation geometry, not just speed, strongly determines mosquito entry (Carrasco-Tenezaca et al., 2023; Sutcliffe & Yin, 2021).

In summary, fans do not repel mosquitoes—they erase the information mosquitoes need to find humans.

4.3. Why thermal or CO2 decoys do not make humans invisible

CO_2 and heat traps attract mosquitoes but do not render humans invisible. Our simulations reproduce this: decoys increase the entropy of the attractor landscape but do not remove the human host as the global maximum unless plume topology collapses.

Classic trap studies demonstrated that lactic acid, CO_2, and skin volatiles modulate attraction strength but cannot eliminate the host signal if the gradient is preserved (Takken & Knols 1999; Cribellier et al. 2018). Modern infrared thermotaxis work further shows that temperature cues modulate behavior only when CO_2 plume coherence remains intact (Giraldo et al., 2023). Cribellier et al. (2018) also showed that heat alone cannot compensate for odor intermittency when the plume is fragmented by local vortex motions, reinforcing our finding that thermal clutter reduces contrast but cannot induce collapse.

Thus, invisibility depends on *gradient destruction*, not *competition strength*. This resolves the familiar complaint: "My trap catches many mosquitoes, but I still get bitten." This is not a failure of attraction, but a failure to induce plume collapse.

4.4. Predictions for real-world deployment

To assess real-world relevance, we compared model predictions with empirical studies on airflow, ventilation, and host-seeking disruption.

Full-scale house experiments show that household or ceiling fans reduce mosquito entry by 60–80%, despite wind speeds far below blow-off thresholds (Carrasco-Tenezaca et al., 2023; Mmbando et al., 2022). This supports our prediction that weak, coherent airflow triggers informational collapse. Low-speed cross-draughts around bed nets alter approach trajectories in predictable ways (Sutcliffe & Yin, 2021), matching the model's gradient-displacement mechanism. Directed airflow combined with vapor-phase repellents reduces orientation and

landing (Hoffmann & Miller, 2002), consistent with airflow–chemical synergy predicted by the model. Architectural studies show that well-ventilated house structures reduce mosquito entry by altering CO2 distribution and flow topology (Francisco & Watanabe, 2024), not merely by increasing velocity.

Finally, recent work demonstrates that background wind *gates olfaction-driven search states* in *Aedes aegypti* (Stupski & van Breugel, 2024), confirming that airflow affects not only trajectories but also sensory reliability.

Together, these data indicate that the key mechanism identified by our model—airflow-induced destruction or redirection of the navigational gradient—is strongly supported across laboratory, semi-field, and architectural studies.

Our predictions also align closely with controlled laboratory observations from Sutcliffe & Yin (2021), who showed that even low-speed cross-draughts (~0.1–0.4 m/s) around bed nets induce abrupt shifts in mosquito activity zones, reducing roof contacts while increasing side approaches. These behavioral redistributions are consistent with our model's gradient-displacement and vortex-shear hypotheses, and reinforce the conclusion that airflow disrupts host-seeking by corrupting plume topology rather than impairing flight itself.

Table 2. Empirical studies supporting airflow-based disruption of mosquito host seeking

Mechanism / Setup	Model Prediction	Empirical Evidence
Household/ceiling fans	Weak airflow induces informational collapse → 60–80% reduction	Carrasco-Tenezaca et al., 2023; Mmbando et al., 2022
Low-speed indoor cross-draught	Small, coherent flows disrupt approach near bed nets	Sutcliffe & Yin, 2021
Directed wind + repellent	Airflow destabilizes plume → synergy with chemical screens	Hoffmann & Miller, 2002
Well-ventilated house geometry	Changes CO2 topology → reduces entry	Francisco & Watanabe, 2024
Wind gating sensory states	Background wind alters olfactory processing	Stupski & van Breugel, 2024

4.5. Experimental validation of model predictions

Prediction 1 — Gradient destruction is measurable.

Schlieren or CO2 visualization can quantify plume breakup under controlled airflow. Recent studies already demonstrate measurable intermittency (Sutcliffe & Yin, 2021; Carrasco-Tenezaca et al., 2023), consistent with scalar-mixing theory (Shraiman & Siggia, 2000).

Prediction 2 — Vortex-induced failure is continuous, not binary. 3-D flight tracking using IR markers should reveal progressive degradation with increasing vortex intensity (van Breugel et al., 2015; Pang et al., 2018).

Prediction 3 — Opposing-fan collapse at subcritical wind speeds. Semi-field trials comparing (i) one strong unidirectional fan versus (ii) two moderate opposing fans should yield superior suppression for (ii), since vortex shear—not velocity—is the key determinant. No classical mosquito model predicts this; thus it provides a strong falsification test.

4.6. Limitations and Future Applications

Our model is intentionally minimal, isolating the physical coupling between airflow and chemothermal gradients. Although it omits multimodal sensory integration, species differences, and detailed turbulence, the collapse mechanism it reveals—*topological destruction of the navigational gradient*—is broadly applicable across mosquito species and even non-biological chemotactic systems.

Future directions include: (1) Airflow-based vector control: Quantitative collapse boundaries can guide the design of fan arrays, ventilation curtains, and passive airflow geometries that destroy plume coherence, offering sustainable alternatives to chemical repellents. (2) Integration with indoor airflow engineering: Because collapse depends on flow *geometry* rather than velocity, coupling the model with CFD, Schlieren imaging, and architectural ventilation design can generate engineered "mosquito invisibility zones.". (3) Hybrid biological–physical models: Incorporating 3-D kinematics, turbulence statistics, and multimodal cue gating will refine predictions without altering the core mechanism. (4) Broader applicability beyond mosquitoes: The same physics governs plume search in moths, flies, crustaceans, and chemical-sensing robots, making this framework relevant to robotics, environmental monitoring, and search-and-rescue algorithms.

Overall, despite its simplicity, the model provides a unified and physically grounded explanation for how airflow disrupts host-seeking—a foundation for practical vector-control innovations and for advancing the physics of odor-guided navigation.

5. CONCLUSION

Our study demonstrates that mosquito host seeking is governed not simply by sensory acuity or behavioral strategy but by the physical integrity of the CO_2–heat plume itself. Using a minimal agent-based model, we identify a robust aerodynamic collapse in which weak directional wind displaces the gradient and vortex-like flows destroy its topology. The resulting "invisibility zone'' emerges reliably across Monte-Carlo ensembles, parameter variations, and sensitivity tests, indicating that the failure stems from fundamental scalar-mixing physics rather than algorithmic or biological limitations.

These findings unify several long-standing observations: household fans offer substantial protection independent of chemical repellents; opposing fans outperform single high-speed fans due to added shear; and heat or CO_2 decoys alone cannot eliminate bites without plume disruption. By quantifying the wind–vortex parameter space that triggers collapse, we provide a predictive framework for designing effective airflow-based interventions.

More broadly, the work reveals a general mechanism for chemotactic failure applicable to flying insects, autonomous robots, and any agent navigating scalar fields under turbulence. Manipulating airflow topology—not merely its magnitude—offers a powerful, non-chemical strategy for vector control, with immediate implications for indoor protection, humanitarian shelters, and future engineered repellents.

In summary, mosquito invisibility is a solvable physical problem: when airflow destroys the navigational gradient, the host becomes unreachable, even though the insect remains fully capable of flight and sensing. This principle offers a new foundation for developing sustainable, physics-driven approaches to disease-vector suppression.

Code Availability

All simulation code used in this study is permanently archived on Zenodo (DOI: 10.5281/zenodo.17645050) and publicly available on GitHub at: https://github.com/sugkp112/mos_phase_models_v19.

The Zenodo record preserves the exact version (v19) used to generate all results presented in the manuscript.

Data Availability

All data used in this study were generated directly by the simulation code described above. No external datasets were used.

Acknowledgments

This research was conducted independently without external funding. The author is grateful for constructive discussions within the open-source scientific computing community, which supported the development and refinement of the simulation framework. The author welcomes scientific discussions or potential collaborations related to this work.

Correspondence: ziru.jiang@gmail.com.

Author Contributions

Conceptualization, methodology, software, formal analysis, visualization, and writing: Zi-Ru Jiang.

Declaration of Interests

The author declares no competing interests.

476
477 Figure S1. Simulation time-step convergence test at vortex strength v=2.0. Detection
478 probability remains stable ($\Delta \approx 0.0008$) between 300 and 900 time steps, confirming that 500 is
479 sufficient for asymptotic settling. Error bars show SD across Monte-Carlo replicates.

480
481 Figure S2. Convergence of mean success rate as a function of simulation repetitions.
482 Both weak and strong airflow conditions converge by N=50 runs. Variance shrinkage
483 confirms that 30-run sampling (used in main text) slightly overestimates noise, making results
484 conservative.
485

486
487   Figure S3. Joint sensitivity of detection probability to sensory noise and agent population.
488   Collapse behavior at v=2.0 is robust across all biologically plausible settings. Only extreme
489 high-noise, low-density configurations (>0.6 noise, N=150) show partial recovery, consistent
490 with stochastic lucky walkers.
491

492
493
494   Figure S4. Wind-disruption curves remain consistent across all CO2–heat weighting ratios
495 tested. Despite altering the relative contribution of odor and thermal cues (2:1, 1:1, 1.5:1, 1:1.5,
496 1:2), all curves exhibit an identical suppression cliff between w = 0.2–0.4. This demonstrates
497 that the aerodynamic collapse threshold is structurally determined by airflow and does not
498 depend on the dominance of any single sensory channel.

499
500

501
502 Figure S5. Distribution of navigation time-to-success under different airflow perturbations.
503 Baseline insects exhibit broad arrival times (mean $\approx$ 254 steps). Wind and vortex conditions
504 compress distribution and reduce arrivals, revealing a dual suppression mechanism: fewer
505 successes and faster failures.
506

508
Figure S6. Distribution of navigation times across airflow conditions.
Time-to-success histograms (30 runs × 250 agents) illustrating how airflow reshapes the
temporal dynamics of host localization. In the baseline, arrivals span the full 0–600-step window
(mean ≈ 254.5 steps), reflecting a mixture of long exploratory trajectories and failures. Under
moderate wind (w = 1.0), arrivals become fast and sharply concentrated (mean ≈ 18.6 steps),
consistent with directional advection that channels a small subset of agents efficiently toward the
host. Under moderate vortex forcing (v = 1.0), the distribution remains broad (mean ≈ 211.2 steps)
with fewer late arrivals, indicating partial filament fragmentation and disrupted gradient coherence.
Under strong wind (w = 2.0), successes remain rare but occur very rapidly (mean ≈ 16.8 steps),
revealing that only drift-aligned trajectories reach the host. Vertical dashed lines denote mean arrival
times. Collectively, these distributions reveal two distinct suppression modes:
(1) Gradient redirection (wind): fast but sparse successes.
(2) Gradient scrambling (vortex): slow, decaying, and increasingly rare successes.
522
523

REFERENCES

1. Bhatt, S. et al. (2015). The effect of malaria control on Plasmodium falciparum in Africa 2000–2015. Nature, 526, 207–211. https://doi.org/10.1038/nature15535

2. Carrasco-Tenezaca, M., Mshamu, S., Moore, J. D., et al. (2023). Effect of passive and active ventilation on malaria mosquito house entry and human comfort. Journal of the Royal Society Interface, 20, 20220794. https://doi.org/10.1098/rsif.2022.0794

3. Cooperband, M. F., McElfresh, J. S. & Mullens, B. A. (2006). Comparison of plume structures of CO2 emitted from mosquito traps in a wind tunnel. Journal of the American Mosquito Control Association, 22, 515–522.

4. Cribellier, A., van Erp, J. A., Hiscox, A., Lankheet, M. J., van Leeuwen, J. L., Spitzen, J., & Muijres, F. T. (2018). Flight behaviour of malaria mosquitoes around odour-baited traps: capture and escape dynamics. *Royal Society open science*, *5*(8), 180246.

5. Majeed S, Hill SR, Ignell R. (2014). Impact of elevated CO2 background levels on the host-seeking behaviour of Aedes aegypti. Journal of Experimental Biology, 217, 598–604.

6. Francisco, M. E. & Watanabe, K. (2024). Innovative house structures for mosquito entry prevention and indoor comfort. Frontiers in Public Health, 12, 1404493. https://doi.org/10.3389/fpubh.2024.1404493

7. Geier, M., Bosch, O. J., & Boeckh, J. (1999). Influence of odour plume structure on upwind flight of mosquitoes towards hosts. Journal of Experimental Biology, 202(12), 1639–1648.

8. Gibson, G. & Torr, S. J. (1999). Visual and olfactory responses of African malaria vectors to host-mimicking traps. Medical and Veterinary Entomology, 13, 123–128. https://doi.org/10.1046/j.1365-2915.1999.00163.x

9. Gillies, M. T. & Wilkes, T. J. (1970). The range of attraction of single men to Anopheles gambiae and A. funestus. Bulletin of Entomological Research, 60, 225–235. https://doi.org/10.1017/S000748530004075X

10. Giraldo, D. et al. (2023). Human scent guides mosquito thermotaxis and host selection under naturalistic conditions. Current Biology, 33, 2367–2382. https://doi.org/10.1016/j.cub.2023.04.050

11. Greppi, C. et al. (2020). Mosquito heat-seeking is driven by an ancestral cooling receptor. Science, 367(6478), 681–684.

12. GREPPI, C.; GARRITY, P. A. (2022). Mosquito heat seeking: the molecular and cellular basis of a key driver of blood-feeding behaviour. In: Sensory ecology of disease vectors. Wageningen Academic, p. 605–619.

13. Hemingway, J. & Ranson, H. (2000). Insecticide resistance in insect vectors of human disease. Annual Review of Entomology, 45, 371–391. https://doi.org/10.1146/annurev.ento.45.1.371

14. Hinze, A., Kucher, T. & van Breugel, F. (2021). Mosquito host seeking in 3D using a climate-controlled wind tunnel. Frontiers in Behavioral Neuroscience, 15, 643693.

15. Hoffmann, E. J. & Miller, J. R. (2002). Reduction of mosquito attacks by combining wind and vapor-phase DEET. Journal of Medical Entomology, 39, 935–938. https://doi.org/10.1603/0022-2585-39.6.935

16. Kennedy, J. S. (1940). The visual responses of flying mosquitoes. Proceedings of the Zoological Society of London A, 109, 221–242. https://doi.org/10.1111/j.1096-3642.1940.tb00831.x

17. Kline, D. L. (2006). Traps and trapping techniques for adult mosquito control. Journal of the American Mosquito Control Association, 22(3), 490–496. https://doi.org/10.2987/8756-971X(2006)22[490:TATTFA]2.0.CO

18. Lahondère, C. & Lazzari, C. R. (2020). Mosquito heat-seeking and thermal ecology. Trends in Parasitology, 36, 907–921. https://doi.org/10.1016/j.pt.2020.07.010

19. Lawal, L., Jimoh, A., Musa, J., et al. (2024). Lingering challenges in malaria elimination

efforts in sub-Saharan Africa. Infectious Diseases of Poverty, 13, 81. https://doi.org/10.1186/s40249-024-01081-9

20. Lei, M. et al. (2023). Numerical investigation of odor-guided navigation under turbulence. Biomimetics, 8, 593. https://doi.org/10.3390/biomimetics8080593

21. McMeniman, C. J. et al. (2014). Multimodal integration of carbon dioxide and other sensory cues drives mosquito attraction to humans. Cell, 156(5), 1060–1071. https://doi.org/10.1016/j.cell.2013.12.044

22. Mmbando, A. S. et al. (2022). The effect of light and ventilation on house entry by Anopheles arabiensis. Malaria Journal, 21, 36. https://doi.org/10.1186/s12936-022-04063-3

23. Montell, C. (2025). The sensory arsenal mosquitoes use to find us. Trends Parasitol, 41, 591–602. https://doi.org/10.1016/j.pt.2025.05.004.

24. Pang, R., van Breugel, F., Dickinson, M. H. & Riffell, J. A. (2018). History dependence in insect flight decisions during odor plume tracking. PLoS Computational Biology, 14, e1005969.

25. Ranson, H. & Lissenden, N. (2016). Insecticide resistance in Anopheles: A worsening situation. Trends in Parasitology, 32, 187–196. https://doi.org/10.1016/j.pt.2015.11.010

26. Shraiman, B. I. & Siggia, E. D. (2000). Scalar turbulence. Nature, 405, 639–646. https://doi.org/10.1038/35015000

27. Stupski, S. D. & van Breugel, F. (2024). Wind gates olfaction-driven search states in free flight. Current Biology. https://doi.org/10.1016/j.cub.2024.07.009

28. Sutcliffe, J. F. & Yin, S. (2021). Effects of indoor air movement and ambient temperature on mosquito (Anopheles gambiae) behaviour around bed nets: implications for malaria prevention initiatives. Malaria Journal, 20, 427. https://doi.org/10.1186/s12936-021-03957-y

29. Szyszka, P. & Galizia, C. G. (2023). Extracting spatial information from temporal odor patterns. Current Opinion in Insect Science, 60, 100–107. https://doi.org/10.1016/j.cois.2023.100107

30. Takken, W., & Knols, B. G. (1999). Odor-mediated behavior of Afrotropical malaria mosquitoes. Annual review of entomology, 44(1), 131-157.

31. Umugwaneza, A., Niyigena, J., Karema, C., et al. (2025). Malaria trends and impact of interventions from 2018 to 2023 in Rwanda. Malaria Journal, 24, [article pending]. https://doi.org/10.1186/s12936-025-05278-w

32. van Breugel, F., Riffell, J., Fairhall, A. & Dickinson, M. H. (2015). Mosquitoes use vision to associate odor plumes with thermal targets. Current Biology, 25, 2123–2129. https://doi.org/10.1016/j.cub.2015.06.046

33. Vergassola, M., Villermaux, E. & Shraiman, B. I. (2007). Infotaxis as a strategy for searching without gradients. Nature, 445, 406–409. https://doi.org/10.1038/nature05464

34. Weiss, D. J., Bhatt, S., Gething, P. W., et al. (2025). Mapping the global prevalence, incidence, and mortality of malaria: a systematic analysis. The Lancet. https://doi.org/10.1016/S0140-6736(25)00038-8

35. WHO. (2023). World Malaria Report 2023. https://www.who.int/publications/i/item/9789240083831