Low quality evidence dominates discussion of carbon benefits of alternative grazing strategies

Jonathan Sanderman[1,*], Colleen Partida[1], Yushu Xia[2], Jocelyn M. Lavallee[3], Mark A. Bradford[4,5]

[1]Woodwell Climate Research Center, 149 Woods Hole Road, Falmouth MA 02540 USA
[2]Lamont Doherty Earth Observatory, Columbia University, New York NY
[3]Environmental Defense Fund, Boulder CO
[4]The Forest School, Yale School of the Environment, New Haven CT
[5]Yale Center for Natural Carbon Capture, Yale University, New Haven CT

*corresponding author: jsanderman@woodwellclimate.org

Abstract

A critical review of alternative grazing strategies and their impact on soil organic carbon (SOC) found most studies (n = 47) failed minimum quality criteria. Ten rigorous studies showed no significant SOC change, yet thirteen less robust studies suggested SOC increases. This divergence highlights that low-quality evidence is prevalent, underscoring a critical need for stronger evidence before asserting climate change mitigation benefits from alternative grazing practices.

For submission to: Nature Climate Change

Article type: Brief Communication

Main text

Grazing of domesticated ruminant animals is the largest land use globally. There are ~32 million km^2 of land that cannot support crop production but produce about one third of global protein[1]. Inappropriate management of rangelands has led to large-scale degradation and desertification[2], resulting in losses of 10's of billions of tons of soil organic carbon (SOC)[3]. Reversing rangeland degradation through regenerative grazing practices is being pursued by governments and civil society as a key nature-based climate solution (NbCS), with an estimated potential to sequester 100s - 1000s millions of tons of CO_2 annually[4,5] and exploding interest in paying producers as part of various carbon programs. Yet, conflicting evidence of the impacts of grazing management on SOC has caused deep polarization of the scientific community, and there is a critical need for clarity to drive consensus around this potential NbCS pathway.

Regenerative grazing is a vague term and often refers to a suite of practices including adaptive grazing, holistic planned grazing, cell grazing, multi-paddock grazing, time-controlled grazing or short-duration rotational grazing[6]. At the core of these practices, which we call 'alternative grazing strategies', is development of grazing management plans that achieve various production and/or conservation outcomes. These plans typically have three key objectives[7]: 1) Improve grass productivity and species composition through adequate rest periods during the growing season; 2) Encourage livestock to consume a broader range of vegetation by increasing stocking density; and 3) More uniformly distribute grazers across the ranch through fencing and improved water distribution. When these objectives are met, there is evidence of improved floristic and productivity benefits[8,9] which could, in theory, translate into gains in SOC or at least reductions in soil degradation and associated SOC loss seen in many rangeland ecosystems.

There is a wealth of evidence that optimal grazing intensity (forage consumption per unit time), which for much of the world means lower livestock numbers per unit area can increase SOC levels[10]. In contrast, alternative grazing strategies do not decrease grazing intensity per se but instead spread grazing more evenly, meaning results from grazing intensity studies cannot be used to understand the SOC benefits of alternative grazing management. Further, quantitative syntheses of "improved grazing management" largely comprise individual studies that are not equivalent to alternative grazing strategies[11], questioning the generalizability of the reported management effects. The first meta-analysis[12] to specifically include a comparison of alternative

versus traditional continuous (or season-long) grazing strategies and found an ~28%

improvement in SOC under alternative strategies, whereas a more recent meta-analysis

focused on Australia[13] found no discernible SOC effect. Several recent syntheses, including Bai

and Cotrufo[4], cite the values reported in Byrnes et al.[12] as the potential of alternative grazing

strategies to increase SOC levels.

A notable shortcoming of these past meta-analyses is that, as is common in meta-analysis[14,15],

the quality of evidence in primary studies is not rigorously considered. Building confidence in the

accuracy of the estimated treatment effect – in this case, the effect of alternative grazing

management – requires a balance of evidence between highly controlled, well-replicated

interventional studies (high internal validity but often conducted at a small scale) and well-

designed observational studies conducted at the scale of real-world grazing operations (high

external validity). After conducting a systematic review and meta-analysis of the peer-reviewed

literature and interpreting the findings using a causal inference lens[15,16,17], we conclude that low

quality evidence dominates discussion of SOC benefits of alternative grazing strategies.

Our systematic review identified 70 primary research studies comparing SOC outcomes of

conventional (continuous or season-long) versus alternative grazing strategies. Few studies had

high internal or external validity, with 10 studies meeting all inclusion criteria (detailed in Table

S1), whereas many failed multiple criteria. The most common reason for exclusion was lack of

representative control (i.e. baseline) comparisons (Figure 1), a category including observational,

paired-site studies where SOC was only measured once post-intervention and non-target

controls were not demonstrated to be comparable. Where a paired-site observational study was

excluded solely for a lack of – or inability to evaluate – representative controls (n = 13), we

created a secondary category of studies. For 66% of secondary studies, site history information

was not reported. Our quality criteria resulted in 27 comparisons of SOC stocks from 10 primary

quality studies and 25 from 13 secondary quality studies (see Supplemental Materials).

Figure 1. Counts of reasons for individual studies omitted from primary analysis of grazing management impacts on SOC change. Lack of representative control was most common exclusions criteria. Note: an individual study could fail multiple criteria.

A typical strength of meta-analysis is the ability discern small but real effects when the
underlying studies often do not have the statistical power to differentiate between treatments.
However, in our analysis when all primary and secondary quality studies were pooled together
the mean ln(RR) of SOC was 0.05 (95% CI: -0.01 – 0.10) and the mean SOC sequestration rate
was 0.27 (95% CI: -0.01 – 0.54) tC ha^{-1} yr^{-1} (Figure 2). When the results were narrowed to only
primary studies, the mean SOC response to adoption of alternative grazing strategies was
indistinguishable from zero (Figure 2). The net positive effect for 'All' studies was then driven by
the secondary category, comprising paired-site studies with only post-intervention
measurements and questions about the representativeness of the controls used for
comparisons (Table S4).
97

Figure 2. Change in soil organic carbon (SOC) after adoption of alternative grazing

management measured as (A) the natural-log of the response ratio and (B) annual

sequestration rate. Means shown as bold lines within box-and-whisker plots. Results are

grouped by study quality while the results for All include both the primary and secondary

studies. P-value results from two-tailed t-tests denoted above each box plot as * <0.1, **<0.05,

***<0.01

Despite hypotheses suggesting wetter, more productive rangelands should be able to sequester

more SOC[18], our findings suggest no difference in efficacy between dryland and non-dryland

aridity classes (Table S5) and no difference between native and improved/sown grasslands

(Table S6). Whether the alternative grazing practice was adaptive (i.e. animal movements and

density based upon observations of forage availability and quality) or prescribed also did not

help explain the variability in the primary study quality category (Table S7).

Our findings suggest that the discussion of SOC benefits of grazing management has been

predominantly informed by lower quality evidence. When these lower quality studies are filtered

out, the ten remaining studies do not suggest that SOC responds to adoption of alternative

grazing management (Figure 2). Reanalyzing previous meta-analyses (Supplement S8) using

our systemic criteria suggest that conclusions in those studies of alternative grazing strategies

are not justified.

An ideal grazing experiment would adopt a before-after-control-intervention design implemented

on a commercial scale and follow SOC changes for 5-10 or more years to provide strong causal

evidence. Because this is logistically difficult to achieve, especially given typical short funding

cycles, most investigations have taken a paired site approach with a one-time sampling
campaign (i.e. a space-for-time substitution). Paired site studies can provide high quality
evidence when there is careful consideration of differences between treatment and control sites.
In addition to the need to pair sites by climate, vegetation, soil type and topographic position,
the underlying assumption of a paired site study is that the treatment and control sites were
managed similarly for at least several decades prior to the implementation of the alternative
grazing plan, and the control site continues the same management. In our reading of the
literature, 34 studies did not meet this criterion; albeit many of those studies simply did not
report this critical information. When they did[19], the information exemplified the challenge of
finding representative controls. For example, in one pairing the alternative grazing site was
converted from cropland 29 years prior to soil sampling whereas the continuous grazing control
was converted 17 years prior. The difference in SOC stocks found in this pairing may simply be
due to the extra 12 years under perennial pasture relative to cropland.
136
Controlled experimental studies had representative controls but primarily because these studies
were conducted at scales far smaller than the typical scale of management for commercial
grazing operations[20], with some plots as small as a few hundred square meters[21,22,23,24].
Additionally, our finding of smaller effects size in controlled experiments than in observational
studies runs counter to typical agronomic results where highly controlled plot-scale experiments
have larger effect sizes than seen in farm-scale commercial settings[25]. It is difficult to know the
scale-dependence of the SOC responses to grazing management, but it makes for a difficult
case to say the study was representative of real-world conditions[26] and should stand as a call to
the research community that large-scale interventional studies are needed.
146
A common thread in both the primary literature and influential meta-analyses on the topic of
SOC response to grazing management[12] is lack of data transparency. Most of the studies only
reported mean data, with 10 out of 23 neglecting to report variance metrics. Only one study
provides the raw data behind the reported mean responses, and more than half did not include
a data availability statement (see Supplemental Materials). Much has been written on this
topic[27] but it bears repeating that science needs to be replicable and reproducible to build
confidence that recommendations will have desired outcomes, requiring open sharing of
underlying data.
155

The lack of strong evidence of SOC benefits upon adoption of alternative grazing management
leaves a void where proponents of these grazing strategies can continue to either cite lower
quality studies or rely on anecdotal accounts of positive experiences. Notably, the secondary
quality studies were cited on average three times as often per year as our higher quality studies
(Table S9). Given the strong and growing interest in alternative grazing management practices
as a NbCS, there is a critical need to establish a much stronger evidence base[28]. Such evidence
should include controlled longitudinal experiments, using fit-for-purpose causal designs with
protocols and tools that can quantify SOC change due to management at the scale of whole
ranch management[29]. Such evidence will permit confident evaluation of which grazing practices
should be adopted where to give the greatest potential climate benefits.
166

References

1. Ritchie, H. & Roser, M. Half of the world's habitable land is used for agriculture. *Our World in Data* https://ourworldindata.org/global-land-for-agriculture (2019).

2. Bestelmeyer, B. T. *et al.* Desertification, land use, and the transformation of global drylands. *Frontiers in Ecol & Environ* **13**, 28–36 (2015).

3. Sanderman, J., Hengl, T. & Fiske, G. J. Soil carbon debt of 12,000 years of human land use. *Proc. Natl. Acad. Sci. U.S.A.* **114**, 9575–9580 (2017).

4. Bai, Y. & Cotrufo, M. Grassland soil carbon sequestration: Current understanding, challenges, and solutions. *SCIENCE* **377**, 603–608 (2022).

5. Roe, S. *et al.* Land-based measures to mitigate climate change: Potential and feasibility by country. *Global Change Biology* **27**, 6025–6058 (2021).

6. Stanley, P. L., Wilson, C., Patterson, E., Machmuller, M. B. & Cotrufo, M. F. Ruminating on soil carbon: Applying current understanding to inform grazing management. *Global Change Biology* **30**, e17223 (2024).

7. Briske, D. D. *et al.* Rotational Grazing on Rangelands: Reconciliation of Perception and Experimental Evidence. *Rangeland Ecology & Management* **61**, 3–17 (2008).

8. Teague, W. R. *et al.* Grazing management impacts on vegetation, soil biota and soil chemical, physical and hydrological properties in tall grass prairie. *Agriculture, Ecosystems & Environment* **141**, 310–322 (2011).

9. Kahn, L. P., Earl, J. M. & Nicholls, M. Herbage mass thresholds rather than plant phenology are a more useful cue for grazing management decisions in the mid-north region of South Australia. *The Rangeland Journal* **32**, 379–388 (2010).

10. Ren, S. *et al.* Historical impacts of grazing on carbon stocks and climate mitigation opportunities. *Nat. Clim. Chang.* **14**, 380–386 (2024).

11. Conant, R. T., Cerri, C. E. P., Osborne, B. B. & Paustian, K. Grassland management impacts on soil carbon stocks: a new synthesis. *Ecological Applications* **27**, 662–668 (2017).

12. Byrnes, R. C., Eastburn, D. J., Tate, K. W. & Roche, L. M. A Global Meta-Analysis of Grazing Impacts on Soil Health Indicators. *J of Env Quality* **47**, 758–765 (2018).

13. McDonald, S. E. *et al.* Grazing management for soil carbon in Australia: A review. *Journal of Environmental Management* **347**, 119146 (2023).

14. Simonsohn, U., Simmons, J. & Nelson, L. D. Above averaging in literature reviews. *Nat Rev Psychol* **1**, 551–552 (2022).

15. Spake, R. *et al.* Improving quantitative synthesis to achieve generality in ecology. *Nat*
Ecol Evol **6**, 1818–1828 (2022).

16. Rubin, D. B. Estimating causal effects of treatments in randomized and nonrandomized
studies. *Journal of Educational Psychology* **66**, 688–701 (1974).

17. Holland, P. W. Statistics and Causal Inference. *Journal of the American Statistical*
Association **81**, 945–960 (1986).

18. Abdalla, M. *et al.* Critical review of the impacts of grazing intensity on soil organic carbon
storage and other soil quality indicators in extensively managed grasslands. *Agriculture,*
Ecosystems & Environment **253**, 62-81 (2018).

19. Mosier, S. *et al.* Adaptive multi-paddock grazing enhances soil carbon and nitrogen
stocks and stabilization through mineral association in southeastern U.S. grazing lands.
Journal of Environmental Management **288**, 112409 (2021).

20. Roche, L. M., Cutts, B. B., Derner, J. D., Lubell, M. N. & Tate, K. W. On-Ranch Grazing
Strategies: Context for the Rotational Grazing Dilemma. *Rangeland Ecology &*
Management **68**, 248–256 (2015).

21. Hoogsteen, M. J. J. *et al.* Do Grazing Systems and Species Composition Affect Root
Biomass and Soil Organic Matter Dynamics in Temperate Grassland Swards?
Sustainability **12**, 1260 (2020).

22. Oliveira, P. P. A. *et al.* Soil carbon dynamics in Brazilian Atlantic forest converted into
pasture-based dairy production systems. *Agronomy Journal* **113**, 1136–1149 (2021).

23. Xu, D., Harder, J. K., Xu, W. & Guo, X. Evaluating the Impact of Grazing Cessation and
Reintroduction in Mixed Prairie Using Raster Time Series Analysis of Landsat Data.
Remote Sensing **13**, 3397 (2021).

24. Dong, W. et al. Effect of rest-grazing management on soil water and carbon storage in
an arid grassland (China). Journal of Hydrology 527, 754–760 (2015).

25. Kravchenko, A. N., Snapp, S. S. & Robertson, G. P. Field-scale experiments reveal
persistent yield gaps in low-input and organic cropping systems. *Proc. Natl. Acad. Sci.*
U.S.A. **114**, 926–931 (2017).

26. Teague, R., Provenza, F., Kreuter, U., Steffens, T. & Barnes, M. Multi-paddock grazing
on rangelands: Why the perceptual dichotomy between research results and rancher
experience? *Journal of Environmental Management* **128**, 699–717 (2013).

27. Committee on Reproducibility and Replicability in Science et al. Reproducibility and
Replicability in Science. (National Academies Press, Washington, D.C., 2019).
doi:10.17226/25303.

28. Bradford, M. A., Kuebbing, S. E., Polussa, A., Sanderman, J. & Oldfield, E. E. Upstream
data need to prove soil carbon as a climate solution. *Nat. Clim. Chang.* **15**, 1013–1016
(2025).
29. Xia, Y. *et al.* Coupling remote sensing with a process model for the simulation of
rangeland carbon dynamics. *Journal of Advances in Modeling Earth Systems*, **17**,
e2024MS004342 (2025).

Supplemental Information

Contents
243

258
259

S1. Methods

Literature search

Peer-reviewed investigations of the impact of switching from conventional continuous grazing to alternative grazing strategies on SOC were compiled using the four different sources. Slightly different search strings were used for each source due to limitations on the number of Boolean terms,

Web of Science search string:
search for:((ALL=(soil)) AND ALL=("grazing management" OR "rotational grazing" OR "continuous grazing" or "adaptive management" OR "time controlled grazing" OR "deferred grazing" OR "cell grazing" OR "holistic management" OR "grazing systems" OR "grazing strategies")) AND ALL=("soil organic carbon" OR "soil carbon" OR "carbon sequestration" OR "soil organic matter" OR "C sequestration" OR "SOC" OR "SOM" OR "soil C" OR "OC" OR "TOC"OR "chemical characteristics")

Elsevier Science Direct search string:
searched: ALL=(soil) AND ALL=("grazing management" OR "rotational grazing" OR "continuous grazing") AND ALL=("soil organic carbon" OR "soil carbon" OR "carbon sequestration")

Google Scholar (via Publish or Perish[1]) search string:
search for: Soil soil organic carbon "rotational grazing" OR "continuous grazing" OR "adaptive management" OR "time controlled grazing" OR "deferred grazing" OR "cell grazing" OR "holistic management" OR "grazing systems" OR "grazing strategy" AND "carbon sequestration" OR "SOC" OR "C stocks"

Scopus (via Publish or Perish[1]) search string:
search for: Soil soil organic carbon "rotational grazing" OR "continuous grazing" OR "adaptive management" OR "time controlled grazing" OR "deferred grazing" OR "cell grazing" OR "holistic management" OR "grazing systems" OR "grazing strategy" AND "carbon sequestration" OR "SOC" OR "C stocks"

These searches were conducted in June 2023. The completeness of our search was assessed by the successful retrieval of a short list of key studies including all of the papers used in the Byrnes et al.[2] meta-analysis[3,4,5,6,7,8,9,10,11,12,13,14]. Five additional primary publications were found in a meta-analysis of grazing impacts on SOC in Australia[15] that were not included in our initial search, but none were found to pass our inclusion criteria.

Filtering and Inclusion criteria

Abstracts from all returned studies were analyzed by asking the following three questions:

1. Is there a comparison of alternative (defined using any of our keywords above) and conventional (i.e. continuous or season-long) grazing strategy? Studies that only tracked SOC under alternative practice without a comparable continuous control baseline were

excluded because of the difficulty in disentangling management from climate induced
changes in SOC[16]
2. Are SOC outcomes reported?
3. Is this primary research? Often multiple studies will use the same primary dataset in
different ways such as for modeling or life cycle analyses. We wanted to be sure that the
same field measurements weren't being utilized more than once in our meta-analysis.

Papers were only retained if the answer to all three of these questions was yes.

Table S1. Inclusion or exclusion criteria.

CRITERIA	OMIT	INCLUDE AS PRIMARY
Is the carbon data available in a table or figure, or as a supplemental dataset?	- No	- Yes
What methods were used to determine carbon content?	- Missing methodological detail	- Uses loss on ignition (LOI) - Uses dry combustion and reported carbonate removal if carbonates may be present - Uses Walkley-Black
How long have the treatments been in place?	- Less than 3 years - Cannot be determined	- Greater than or equal to 3 years
To what depth was soil collected?	- Depth of 10 cm or less	- Greater than 10-cm depth
Are there significant external factors that may affect SOC sequestration between treatments that are NOT accounted for in data analysis?	- Yes, i.e. differing climate, topography, parent material, soil type NOT accounted for in data analysis (raw means reported)	- No, i.e. no significant factors OR factors are accounted for in data analysis (conditional means reported)
Is the control representative of the pre-intervention management in the treatment plot?	- No - Cannot be determined	- Yes

After this filtering, the retained papers were read in detail to determine whether the study met
our predefined inclusion criteria (Table S1). After reading the papers in detail, it was determined
that several studies actually failed at least one of the initial questions – e.g. an alternative
grazing management might have been mentioned in the abstract but upon reading the methods,
there wasn't an actual comparison with conventional grazing.

Knowledge of the method of carbon determination, especially in semi-arid systems where
carbonates are often present, is important for determining the quality of the SOC data. We

chose three years as the minimum study duration as SOC changes slowly and it is extremely
unlikely that real differences can arise in treatments in less than three years. The topsoil (≤ 10
cm) in grasslands only contains a small fraction of the total SOC stocks and changes in this
layer may not be representative of changes in the entire profile. The question on significant
external factors was included because the raw means in regional survey studies cannot be
compared until potential confounding factors (climate, soil, topographic, etc.) that also influence
soil carbon stocks and their rate of change are accounted for (see Allen et al.[7] for a good
example in the grazing-SOC literature). Accounting for these factors builds confidence that
estimates of soil carbon change are due to the grazing management and not non-target causal
variables like mean annual precipitation. Similarly, the representativeness of the control site in
observational paired-site studies must satisfy expectations that treated and control sites are
similar enough with regards other potential causal variables, such as soil texture and longer-
term management history, to permit differences in soil carbon stocks to be attributed to grazing
management. When such paired-site studies are conducted at a single point in time, as
opposed to including pre-intervention data on soil carbon stocks that can be used to calculate
differences due to treatment versus control of changes in time, appropriate matching of pairs by
other potential causal variables becomes critical because SOC levels are unknown at the time
of practice adoption, invoking the assumption that they were approximately equal at that time.
That is, we have to be confident that the baselines (i.e. controls) for the sample of sites that are
measured are representative enough in terms of environmental, soil, land use and historical
management variables for them to allow confidence that estimated effects of grazing
management are not due to non-random bias in observed and/or non-observed non-target
causal variables.
340
341   Studies that met all these criteria are termed primary studies in all subsequent analyses. If a
342   study met all the criteria except the final one on representativeness of the control site, the study
343   was retained but flagged as secondary quality. These secondary studies were kept in the
344   analysis because the majority of studies of alternative grazing management studies have this
345   paired-site design.
346

Data extraction

For all retained studies, the following information was extracted or at least attempted to be
extracted: study year, experimental design, treatments, whether the alternative grazing strategy
was adaptive or prescriptive (i.e. decisions on animal density and movements made based on
observations of forage availability/quality vs. following a set plan regardless of seasonal
conditions), controls used, land use history, duration of treatment on land prior to sampling,
grassland type (native or improved) and species, methods for improved pasture, size of
treatment area/study scale, grazing animal type, stocking rate, tools used for soil sampling,
maximum depth of sampling, soil sampling strategy, method for determining SOC content,
carbonate removal information, SOC content and stock data, bulk density (BD) data, country,
coordinates, mean annual precipitation (MAP), mean annual temperature (MAT), and world soil
order. If climate data or soil order was not available it was extracted using location data from

ClimateCharts.net[17] and SoilGrids[18]. Aridity index (AI) was also extracted using location data
from the Global-Aridity_ET0[19].

A mean SOC concentration (wt%) was calculated for the 0-30 cm and 0 to maximum depth
increments by taking the BD-weighted means for each horizon within the specified depth range.
SOC stocks (tC ha^{-1}) were either directly extracted from the study or calculated based on SOC
concentration and BD and depth. No attempt was made to correct for equivalent soil mass.

All data from this study, including the full list of studies retrieved from the literature search, the
short list of 70 studies along with their assessment against our inclusion criteria, and the
extracted data from the final list of studies are included in the supplementary materials. The
locations of the final primary and secondary studies are shown in Figure S1.

372
Figure S1. Locations of primary and secondary studies.

Data analysis

Differences between continuous and alternative grazing practices were summarized in two
ways. First, an effect size measured as the natural log of the alternative grazing strategy SOC
response divided by the continuous grazing SOC response (ln(RR)) was calculated for different
categories. Second, when all the necessary data was available, relative SOC sequestration
rates (tC ha^{-1} yr^{-1}) were also calculated for the 0-30 cm and 0-maximum depth ranges.

Initial analyses indicated that the effect size calculated from SOC concentration and SOC stocks
were nearly identical (see Figure S2) and that there were no differences between studies that
only reported SOC to < 30 cm, 30-50 and >50 cm (see Table S3). These two findings allowed
us to simplify our analyses by only reporting the one effect size as the average of ln(RR) of SOC

concentration and ln(RR) of SOC stocks and by only reporting the results for 0-maximum depth
range.
387
Two-tailed t-tests were then conducted to explore whether individual effect sizes were
statistically different from zero for the following categories: study quality, experimental design,
alternative grazing strategy, aridity index and rangeland type (natural versus
introduced/improved grasses). For experimental design, the groups were limited to controlled
experiment and paired site design because those two groups represented 94% of the data.
Similarly for pasture type, the groups were limited to improved or native pasture. All analyses
and plotting was done in Python using the Pandas[20], Seaborn [21], Matplotlib[22] and SciPy[23]
libraries.
396

397   ***S2. Reporting SOC concentration or stocks did not matter***

398   The ln(RR) results from SOC concentration (wt%) and SOC stocks were nearly identical (Figure
399   S2. As such, in subsequent analyses the ln(RR) was calculated as the average of the ln(RR) for
400   SOC% and SOC stocks. This allowed the maximum number of comparisons to be included in
401   statistical tests because not all studies reported SOC% and stocks.
402

403
404   **Figure S2**. Comparison of natural-log of the response ratio (ln(RR)) calculated for SOC
405   concentration (wt%) and for SOC stocks (tC ha^{-1}) where data was available to calculate ln(RR)
406   both ways (n = 22) indicates that there is no systematic difference in ln(RR) allowing pooling of
407   data to maximize sample size in the meta-analysis.
408
409

S3. No clear effect of soil measurement depth

Perhaps due to our inclusion criteria that sampling depth needed to be greater than 10 cm, the
mean sampling depth was 50 cm and, importantly, there was no difference in results between
studies measuring to less than 30 cm, those that measured to between 30-50 and those that
measured greater than 50 cm (Table S3). Single factor ANOVA p-values were 0.99, 0.88 and
0.81 for primary studies, secondary studies and all studies, respectively. Results for SOC
sequestration rates were similar (data not shown). This result gave us confidence in grouping all
depths together for the main statistical analyses shown in Table 1.

Table S3. Natural-log of the response ratio for soil organic carbon grouped by depth and study
quality.

Depth range (cm)	Primary			Secondary			All		
	n	mean	95% CI	n	mean	95% CI	n	mean	95% CI
<30	6	-0.02	(-0.27, 0.22)	5	0.02	(-0.25, 0.29)	11	-0.01	(-0.18, 0.17)
30-50	10	-0.01	(-0.05, 0.04)	12	0.14	(0.02, 0.26)	22	0.07	(0.00, 0.15)
>50	11	0.00	(-0.13, 0.13)	8	0.11	(0.03, 0.18)	19	0.04	(-0.04, 0.13)
all	27	-0.01	(-0.08, 0.07)	25	0.11	(0.02, 0.19)	52	0.05	(-0.01, 0.10)

421
422
423
424

S4. Paired site studies dominate positive responses

There were four study designs found in our final paper list: controlled experiments where an area of was divided into multiple control and intervention plots, unreplicated experiments where an area was divided into just one replicate of each control and intervention, paired site designs where two "paired" fields under contrasting management were compared, and survey designs where a number of ranches with varying management practices across a region were measured. Table S4 indicates that nearly all the positive SOC responses were associated with paired site studies rather than controlled experiments (p = 0.02 and 0.04 for ln(RR) and SOC sequestration rate, respectively) but these were the same studies that dominated the secondary class based on inclusion/exclusion criteria.

Table S4. Soil organic carbon (SOC) response measured as the natural-log of the response ratio and annual sequestration rate in response to adoption of alternative grazing practices. Results combined for primary and secondary studies and grouped by experimental design.

Experimental design	ln(RR)			SOC sequestration (tC/ha/yr)		
	n	mean	95% CI	n	mean	95% CI
Controlled experiment	13	-0.01	(-0.07, 0.05)	10	-0.07	(-0.34, 0.26)
Paired sites	36	0.07	(-0.01, 0.15)	34	0.40	(0.04, 0.76)
Survey	1	0.15				
Unreplicated experiment	2	-0.05	(-0.08, -0.02)	2	-0.30	(-0.51, -0.09)

S5. Aridity index

It has been hypothesized that there might be more of a positive SOC response in more mesic environments where productivity isn't so constrained by available soil moisture[24]. When grouping studies by aridity class - where AI > 0.65 was considered 'non-dryland', and AI < 0.65 considered 'dryland' - there was little difference in ln(RR) between categories (Table S5). Results for SOC sequestration rates were nearly identical with no separation between aridity class (data not shown).

Table S5. Soil organic carbon (SOC) response measured as the natural-log of the response ratio response to adoption of alternative grazing practices. Results grouped by aridity class.

Aridity class	**Primary**			**Secondary**			**All**		
	n	**mean**	**95% CI**	**n**	**mean**	**95% CI**	**n**	**mean**	**95% CI**
non-dryland	8	-0.03	(-0.11, 0.17)	13	0.11	(0.06, 0.17)	21	0.06	(0.01, 0.11)
dryland	19	0.00	(-0.10, 0.10)	12	0.10	(-0.07, 0.26)	31	0.04	(-0.05, 0.13)

S6. Native versus improved rangeland

It has also been hypothesized that alternative grazing strategies may be more responsive on improved grasslands (some combination of sowing improved varieties, fertilizing or irrigating) that typically have greater productivity than native grasslands in the same climatic zone[25]. Overall, there was no difference in SOC effect size between improved or native grasses (Table S6). However, when just focusing on the secondary study class the improved grasses but not the native grasses had ln(RR) values significantly greater than zero (Table S6). This trend also held for SOC sequestration rates (data not shown).

Table S6. Soil organic carbon (SOC) response measured as the natural-log of the response ratio response to adoption of alternative grazing practices. Results grouped by grassland type.

Grass type	Primary			Secondary			All		
	n	mean	95% CI	n	mean	95% CI	n	mean	95% CI
Improved	20	-0.01	(-0.09, 0.07)	14	0.15	(0.06, 0.25)	34	0.06	(-0.01, 0.12)
Native	7	0.00	(-0.19, 0.20)	8	0.07	(-0.12, 0.26)	15	0.04	(-0.09, 0.17)
Unclear				3	-0.03	(-0.07, 0.02)	3	-0.03	(-0.07, 0.19)

S7. Specific management practices

One of our objectives was to try to understand if specific grazing management practices had
more or less of an SOC benefit after implementation. To achieve this objective, the improved
grazing management styles were defined as either adaptive or prescribed based on information
provided in each publication. One of the secondary studies[26] could not be categorized and as
such was excluded from this analysis. Five of the nine primary studies were classified as using
adaptive grazing management, but these studies represented only about one third of the
primary pairs of data. All but one of the secondary studies were defined as adaptive making it
difficult to make any conclusions about the impact of grazing management style within this study
quality category.
477
478   Despite the mean SOC sequestration rate for the primary studies appearing to separate out by
479   grazing management style (+0.22 vs. -0.12 tC ha^{-1} yr^{-1} for adaptive and prescribed,
480   respectively), the ln(RR) for these two categories were both essentially zero and a single-factor
481   analysis of variance suggests the mean sequestration rates are not different from each other (p
482   = 0.50) due to the large variance within each category (Table S7).
483
484   **Table S7**. Soil organic carbon (SOC) response measured as the natural-log of the response
485   ratio and annual sequestration rate in response to adoption of alternative grazing practices.
486   Results presented by grazing management style nested within study quality category.

Study quality	Grazing management style	ln(RR)			SOC sequestration (tC/ha/yr)		
		n	mean	95% CI	n	mean	95% CI
Primary	Adaptive	9	-0.01	(-0.17, 0.16)	6	0.22	(-0.32, 0.73)
	Prescribed	18	-0.01	(-0.09, 0.07)	18	-0.07	(-0.59, 0.45)
Secondary	Adaptive	22	0.13	(0.06, 0.20)	22	0.53	(0.19, 0.87)
	Prescribed	2	-0.24	(-0.75, 0.27)	0	n/a	n/a

487
488

S8. Reconsidering existing literature

Given that Byrnes et al.[2] has been one of the main studies cited when supporting the notion that
the adoption of alternative grazing strategies leads to SOC sequestration, we tried to determine
why our conclusions varied so much from the conclusions in that study. Byrnes et al.[2] supply a
list of studies that were used in estimating effect sizes but not the detailed data themselves
making it difficult to reproduce their results. We re-extracted the data and attempted to recreate
their analysis but couldn't replicate their findings (Table S8). Byrnes et al.[2] report a mean effect
size of 0.25 (95% CI: 0.10 - 0.41) for 44 comparisons of alternative and conventional grazing
strategies but we could only reconstruct a mean effect size of 0.15 for the same number of
comparisons for the Byrnes et al.[2] dataset.

More importantly, most of the data sources used by Byrnes et al.[2] did not meet our inclusion
criteria (Table S8), highlighting the need to consider the quality of the studies being assembled
for quantitative syntheses if one is to have confidence in the average effect across studies[27,28].
In fact, 7 of the 12 studies listed were omitted from our meta-analysis for failing at least one of
our inclusion criteria or repeating data already included in another study. Another 4 studies
made our secondary list and only one study was on our primary study list. To approximate the
same sample size for each study as reported in Byrnes et al.[2], it appears that each soil depth
increment must have been included as an independent observation (i.e. if a study measured 0-
5, 5-15 and 5-30 cm there were three data points used instead of just one data point for 0-30
cm).

A notable inclusion by Byrnes et al.[2], which we excluded because it failed exclusion criteria #5
(Table S2), was Allen et al.[7]. In this regional survey in Queensland, Australia, the authors
measured much higher SOC stocks under alternative grazing strategies than under
conventional grazing. However, after the authors of the original paper conducted a statistical
analysis that removed the confounding effect of climatic differences between sites (all the
alternative grazing sites were in wetter regions of the state), they concluded that pastures under
alternative grazing strategies had less SOC than conventionally grazed pastures. Byrnes et al.[2]
apparently extracted and included the raw data without consideration of the need to account for
confounding in the study design.

There are a few high-profile papers that examined a shift from cropland to alternative grazing
management of grasses[29,30] which are often incorrectly used to support the notion that
alternative grazing strategies can have large SOC benefits. Both studies focused on
chronosequences of improved pastures seeded into former cropland in the SE of the United
States. Large gains in SOC typically occur when croplands are converted to pastures regardless
of grazing management strategies[31]. A third highly cited paper concluding large sequestration
rates under an adaptive grazing plan[32] was also excluded from our study because there was no
comparison to continuous grazing making attribution of any change over time in SOC to the
grazing management strategy difficult.

Table S8. Comparison of our attempt to reconstruct results from Byrnes et al.[2] and our
extraction (this study) of the same data sources. Studies were classified using our inclusion
criteria (see methods).
536

Study quality	Byrnes reconstruction			this study		
	n	Mean	95% CI	n	mean	95% CI
Primary	9	-0.01	(-0.31, 0.29)	3	0.01	(-0.57, 0.59)
Secondary	17	0.22	(0.15, 0.29)	6	0.24	(0.13, 0.35)
Omitted	18	0.17	(0.05, 0.28)			
All	**44**	**0.15**	**(0.07, 0.23)**	**9**	**0.16**	**(-0.04, 0.36)**

537
538
539

S9. Citation frequency

We utilized Google Scholar to capture the number of times each study was cited as of June 2025 because Google Scholar picks up both peer-reviewed and grey literature articles and reports making it a more inclusive measure of the impact of a particular study. The 10 primary studies were cited 225 times, while the 13 secondary studies were cited a total of 1508 times with, on average, 3 times as many citations per year (Table S9).

Table S9. Google Scholar citation rate for studies group by study quality.

Study quality	Sum of citations	Average citations per study	Average citations per study per year
Primary	287	29	4.3
Secondary	1508	116	12
All	**1795**	**78**	**8.7**

References

1. Harzing -, A.-W. Publish or Perish. *Harzing.com* https://harzing.com/resources/publish-or-perish (2016).

2. Byrnes, R. C., Eastburn, D. J., Tate, K. W. & Roche, L. M. A Global Meta-Analysis of Grazing Impacts on Soil Health Indicators. *J of Env Quality* **47**, 758–765 (2018).

3. Sandhage-Hofmann, A. *et al.* Rangeland management effects on soil properties in the savanna biome, South Africa: A case study along grazing gradients in communal and commercial farms. *Journal of Arid Environments* **120**, 14–25 (2015).

4. Wang, Y. *et al.* Effects of land use and precipitation on above- and below-ground litter decomposition in a semi-arid temperate steppe in Inner Mongolia, China. *Applied Soil Ecology* **96**, 183–191 (2015).

5. Hou, X., Wang, Z., Michael, S. P., Ji, L. & Yun, X. The response of grassland productivity, soil carbon content and soil respiration rates to different grazing regimes in a desert steppe in northern China. *The Rangeland Journal* **36**, 573–582 (2014).

6. Cowie, A. L. *et al.* Impact of carbon farming practices on soil carbon in northern New South Wales. *Soil Research* **51**, 707–718 (2013).

7. Allen, D. E. *et al.* What determines soil organic carbon stocks in the grazing lands of north-eastern Australia? *Soil Research* **51**, 695–706 (2013).

8. Kotzé, E., Sandhage-Hofmann, A., Meinel, J.-A., Du Preez, C. C. & Amelung, W. Rangeland management impacts on the properties of clayey soils along grazing gradients in the semi-arid grassland biome of South Africa. *Journal of Arid Environments* **97**, 220–229 (2013).

9. Garcia, M. R. L., Sampaio, A. A. M. & Nahas, E. Impact of different grazing systems for bovine cattle on the soil microbiological and chemical characteristics. *R. Bras. Zootec.* **40**, 1568–1575 (2011).

10. Teague, W. R. *et al.* Soil and herbaceous plant responses to summer patch burns under continuous and rotational grazing. *Agriculture, Ecosystems & Environment* **137**, 113–123 (2010).

11. Teague, W. R. *et al.* Grazing management impacts on vegetation, soil biota and soil chemical, physical and hydrological properties in tall grass prairie. *Agriculture, Ecosystems & Environment* **141**, 310–322 (2011).

12. Chan, K. Y. *et al.* Soil carbon stocks under different pastures and pasture management in the higher rainfall areas of south-eastern Australia. *Soil Research* **48**, 7–15 (2010).

13. Banerjee, M. R., Burton, D. L., McCaughey, W. P. & Grant, C. A. Influence of pasture management on soil biological quality. *Journal of Range Management* **53**, 127–133 (2000).

14. J. E. Gilley, B. D. Patton, P. E. Nyren, & J. R. Simanton. Grazing and Haying Effects On Runoff and Erosion from a Former Conservation Reserve Program Site. *Applied Engineering in Agriculture* **12**, 681–684 (1996).

15. McDonald, S. E. *et al.* Grazing management for soil carbon in Australia: A review. *Journal of Environmental Management* **347**, 119146 (2023).

16. Mitchell, E. *et al.* Making soil carbon credits work for climate change mitigation. *Carbon Management* **15**, 2430780 (2024).

17. Zepner, L., Karrasch, P., Wiemann, F. & Bernard, L. ClimateCharts.net – an interactive
climate analysis web platform. *International Journal of Digital Earth* **14**, 338–356 (2021).
18. Poggio, L. et al. SoilGrids 2.0: producing soil information for the globe with quantified
spatial uncertainty. SOIL 7, 217–240 (2021).
19. Zomer, R. & Trabucco, A. Global Aridity Index and Potential Evapotranspiration (ET0)
Database: Version 3. 6086884631 Bytes figshare
https://doi.org/10.6084/M9.FIGSHARE.7504448.V6 (2022).
20. McKinney, W. Data Structures for Statistical Computing in Python. in 56–61 (Austin,
Texas, 2010). doi:10.25080/Majora-92bf1922-00a.
21. Waskom, M. seaborn: statistical data visualization. *JOSS* **6**, 3021 (2021).
22. Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Comput. Sci. Eng.* **9**, 90–95
(2007).
23. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python.
Nat Methods **17**, 261–272 (2020).
24. Booker, K., Huntsinger, L., Bartolome, J. W., Sayre, N. F. & Stewart, W. What can
ecological science tell us about opportunities for carbon sequestration on arid
rangelands in the United States? Global Environmental Change 23, 240–251 (2013).
25. Bidwell, T.G. & Woods, B. Management Strategies for Rangeland and Introduced
Pastures. Oklahoma Cooperative Extension Service NREM-2869 (2017).
26. Yang, H. *et al.* Grazing disturbance significantly decreased soil organic carbon contents
of alpine grasslands on the Tibetan Plateau. *Front. Environ. Sci.* **11**, 1113538 (2023).
27. Simonsohn, U., Simmons, J. & Nelson, L. D. Above averaging in literature reviews. *Nat*
Rev Psychol **1**, 551–552 (2022).
28. Spake, R. *et al.* Improving quantitative synthesis to achieve generality in ecology. *Nat*
Ecol Evol **6**, 1818–1828 (2022).
29. Machmuller, M. B. *et al.* Emerging land use practices rapidly increase soil organic
matter. *Nat Commun* **6**, 6995 (2015).
30. Rowntree, J., Stanley, P., Beede, D., DeLonge, M. & Hamm, M. 143 Impacts of soil
carbon sequestration on life cycle greenhouse gas emissions in midwestern USA beef
finishing systems. *Journal of Animal Science* **97**, 147–148 (2019).
31. Post, W. M. & Kwon, K. C. Soil carbon sequestration and land-use change: processes
and potential. *Global Change Biology* **6**, 317–327 (2000).
32. Stanley, P. L., Rowntree, J. E., Beede, D. K., DeLonge, M. S. & Hamm, M. W. Impacts
of soil carbon sequestration on life cycle greenhouse gas emissions in Midwestern USA
beef finishing systems. *Agricultural Systems* **162**, 249–258 (2018).
628
629