Machine learning-based prediction of human structural variation and characterization of associated sequence determinants

Daven Lim[1,2], Runyang Nicolas Lou[3], Nilah Ioannidis[4,5], Peter H Sudmant[3,5]

1 Department of Biosystems Science and Engineering, ETH Zürich, Zürich, Switzerland
2 Department of Electrical Engineering and Computer Sciences, UC Berkeley, Berkeley, California, USA
3 Department of Integrative Biology, University of California Berkeley, Berkeley, CA, USA
4 Department of Applied Math, University of California Santa Cruz, Santa Cruz, CA, USA
5 Center for Computational Biology, University of California Berkeley, Berkeley, CA, USA

Correspondence should be addressed to Peter H Sudmant: psudmant@berkeley.edu

ABSTRACT

Structural variants (SVs) represent a major source of genetic diversity and play key roles in human disease and evolution. Yet, the extent to which local sequence context shapes the likelihood of structural variant formation remains poorly quantified. Here, we develop machine learning models to predict the occurrence of SVs across the human genome and characterize genomic determinants associated with their formation. We developed both a sequence only-based convolutional neural network (CNN) model as well as a random forest approach integrating diverse genomic annotations. Both models achieve high predictive performance individually (>90% AUROC) which can be further improved in an ensemble. The predictive ability of these models demonstrates that SV-prone regions can be accurately inferred from sequence context. Model interpretability techniques reveal key genomic contributors to SVs, including effects of sequence motifs such as microhomology and non-canonical DNA structures, as well as the presence of SV hotspots. We find that different classes of SVs exhibit distinct sequence determinants, with inversions displaying particularly unique signatures. Moreover, predicted SV probability correlates with allele frequency and gene functional constraint, highlighting the utility of the model for variant effect prediction. These findings demonstrate that machine learning models trained on local sequence features can identify unstable genomic regions and provide a framework for quantifying SV susceptibility and SV variant effects in personalized genomics.

INTRODUCTION

Structural variants (SVs) - including insertions, deletions, duplications, and inversions - are large-scale genomic rearrangements that represent a major source of genetic diversity in humans[1,2]. SVs can affect molecular and cellular processes which can cause genetic diseases, affect gene function, and contribute to evolution and trait complexity[3]. For example, de novo duplications at the 7q11.23 locus are strongly associated with autism spectrum disorder, and the reciprocal deletion causes Williams-Beuren syndrome[4]. Rare SVs can alter genes in neurodevelopmental pathways and contribute to schizophrenia[5]. SVs are also an important source of adaptive traits. For instance, increased copy number of salivary amylase genes (*AMY1*) is correlated with increased salivary amylase protein levels, and individuals from populations with high starch diets have been shown to have on average more *AMY1* copies [6] [7].

SVs arise through various sources, each leaving characteristic signatures in the local sequence context. A major source of SVs is non-allelic homologous recombination (NAHR), which occurs when highly similar low-copy repeats or segmental duplications misalign during meiosis, producing deletions, duplications, and inversions[8]. Transposable element activity, such as SINE/Alu and LINE/L1-mediated retrotransposition, also contributes substantially to human structural variation through insertion events. Furthermore, these sequences can serve as substrates for homology mediated rearrangements[9]. In addition, replication-associated processes such as fork stalling and template switching can generate complex duplications and rearrangements without requiring extended homology[10]. Many human SVs also originate from non-homologous end joining (NHEJ), a repair mechanism that rejoins double-strand breaks with minimal or no homology and frequently introduces short insertions or deletions at the junctions[11]. Breakpoint analyses have consistently shown enrichment of short microhomology tracts at SV junctions relative to random expectation, highlighting the role of microhomology-mediated repair and the importance of local sequence composition in predisposing genomic regions to structural instability[12]. Collectively, these mechanisms point toward sequence-dependent determinants of SV formation, motivating the use of computational models to quantify how local sequence features shape genome fragility.

Recent work in a eukaryotic pathogen and subsequently in *Arabidopsis thaliana* has shown that machine learning models can predict structural rearrangements from local genomic features (e.g. proximity to TEs and genes), demonstrating that eukaryotic genomes contain identifiable signatures of chromosomal instability[13]. However, these studies were limited to smaller and less complex genomes and focussed on annotated genomic features instead of raw sequence. The degree to which such predictive signatures exist in the human genome and the extent to which local sequence context influences SV formation is also unclear. Furthermore, there are currently no quantitative frameworks to estimate the likelihood that SVs arise at specific genomic positions and the extent to which raw sequence content can serve as an accurate predictor.

Here, we develop machine learning models to assess SV susceptibility across a large, population-scale SV dataset generated from long-read haplotype-resolved genomes. We integrate an expanded genomic feature space that includes both raw sequence and diverse genomic annotations such as conservation metrics, base composition, neighboring genomic

variation, repeat annotations, and gene-related features. By examining feature contributions and
learned sequence determinants, we identify genomic and sequence factors associated with
SVs. Predictive models of species-wide SV occurrence have the potential to support disease
variant applications, advance our understanding of genome evolution, and guide future studies
on the mechanisms of structural rearrangement.

RESULTS

Development of machine learning models to predict structural variants
Human structural variants are a diverse form of genetic variation spanning five orders of
magnitude in length (**Fig 1A**). Nevertheless, SVs are known to be associated with distinct
sequence features (e.g. stretches of homology). We set out to develop machine learning models
capable of predicting the occurrence of SVs from DNA sequence and associated DNA features.
We employed a population-scale SV dataset from Phase 3 of the Human Genome Structural
Variation Consortium[14]. This is the first population-scale set of human genome assemblies
resolved to near T2T completeness, encompassing 173,824 distinct variants discovered in 65
individuals (130 haplotypes) from diverse ancestries. We first set out to develop a sequence-
only deep learning model for SV prediction utilizing a residual convolutional neural network
architecture similar to models that have been developed to predict splice site usage[15], gene
expression[16,17], mRNA expression[18], and accessible genomic sites[19]. This approach estimates a
probability that any given basepair location is the breakpoint of a structural variant given its
adjacent flanking sequence (**Fig 1B, C**) (**Methods**). Sequence context lengths ranging from
100bp to 2000bp were explored. We also constructed a random forest-based SV prediction
model in which the probability of an SV is determined from the relative distribution of sequence
features from the nucleotide of interest within the context window. These features include the
density of SNPs and neighboring SVs, GC content, constraint (PhyloP score), gene density, and
mobile element density, among others (**Fig 1C, D**). Finally, we constructed an ensemble model
in which the CNN predicted probability was incorporated as a feature into the random forest
thereby combining both sequence content and genomic features into a single model (**Fig 1C**,
see **Methods**). These methods were compared against a simple logistic regression approach
which incorporated all of the features of the random forest. All models were trained and tested
on 347,648 total samples (173,824 SVs and 173,824 background regions), with metrics
reported on the test set which is not encountered during training (**Methods**).
107
We tested the performance of our models on the GRCh38 human genome equally split between
true SVs and randomly selected control regions (**Methods**). Model performance was quantified
using both the true positive and false positive rates assessed as the area under the receiver
operating characteristic curve (AUROC) and the precision and recall rates, assessed as the
area under the precision recall curve (AUPRC) (**Fig 1E, F**). All of our models outperformed the
logistic regression approach, though this simple approach did perform surprisingly well,
highlighting that local genomic features are strong determinants of the probability of SV
formation (AUROC 0.890, AUPRC 0.916). The random forest approach modestly outperforms
this basic method (AUROC 0.904, AUPRC 0.930), capturing non-linear relationships and
higher-order interactions among genomic features that the linear logistic regression cannot
represent. Remarkably however, the CNN approach, which only uses local sequence, achieved

similar performance (AUROC 0.903, AUPRC 0.926). The ensemble approach combining the
CNN with the random forest further improved performance (AUROC 0.916, AUPRC 0.936). The
CNN and ensemble models were also both improved by increasing the sequence context size
up to 400bp, though diminishing beyond this point (**Fig 1F**). Together these results demonstrate
that both genomic features as well as raw local genomic sequence are strong predictors of SV
formation.
125
Feature contributions to model predictions
Identifying and quantifying the impact of features to our model predictions provides insight into
the mechanisms underlying SV formation. We conducted several analyses to quantify the
contribution of individual features to model predictions. We first assessed the correlation
between input features, the predicted SV probabilities, and SV presence/absence across
models using Spearman correlation coefficients (**Fig 2A**). Several input features were strongly
correlated with each other, as expected. For example, phyloP scores were negatively correlated
with SNV and indel presence and DNase accessibility was correlated with transcription factor
binding peaks. Feature correlations across models were highly consistent, with model
probability strongly positively correlated with SNVs, indels, and the presence of other SVs, and
negatively correlated with phyloP score and the distance to the nearest simple repeat. These
features are similarly strongly correlated with the presence of SVs ("True label" **Fig 2A**). While it
is expected that the random forest would pick up such features, surprisingly, the sequence-only
CNN was also able to distinguish them. PhyloP score, for instance, exhibited a strong negative
correlation with CNN output probability, indicating that the model is able to distinguish
conserved regions of the genome from sequence alone.
142
We next evaluated the relative importance of each feature in the ensemble model, which
includes the CNN-derived probability as an input (**Fig 2B**). Gini importance measures how much
a feature reduces node impurity across all trees in the random forest, where impurity reflects
how mixed the classes are within a node and a split that makes nodes more homogeneous
decreases impurity. Distributions of each feature are shown alongside their Gini importances.
The number of neighboring SVs within the 400 bp window contributed most strongly to
predictions, followed by the number of SNPs, conservation scores, and GC content. The CNN-
derived prediction also ranked highly, indicating that sequence-level patterns captured by the
CNN provide complementary predictive signals beyond genomic annotations.
152
To determine which labeled features the CNN is able to capture using raw sequence, we
computed permutation importances in the ensemble model with and without the CNN as a
feature on the test set. Permutation importances quantify the decrease in predictive
performance when each feature is permuted. The decrease in predictive performance of a
feature after permutation indicates the importance of the feature. Without the CNN as an input,
neighboring variants - both SVs and SNPs - showed the highest contributions, followed by
distance to simple repeats, conservation scores, GC content, and distance to Alu elements,
largely consistent with relative feature importance measured with Gini impurity. However, after
introducing the CNN predictions as an additional feature, we observed a redistribution of
importances, reflecting the extent to which the CNN captures signal previously explained by

annotated features (**Fig 2C**). Conservation scores and GC content showed the largest
reductions in importance, suggesting that the CNN effectively captures these sequence
properties. The CNN also appeared to capture the number of SNPs and small indels, and the
distance to simple repeats, Alu, and L1 elements. In contrast, the importance of neighboring
large SVs increased, albeit slightly, indicating that local structural variant density provides
information that is not fully captured by sequence context alone and remains a distinct predictor
of SV likelihood. Although the CNN performs strongly on its own, its permutation importance is
modest when added to the random forest, indicating that the ensemble is able to recover much
of the same predictive signal from existing genomic annotations. The CNN therefore appears to
provide more localized sequence-level cues in addition to the broader genomic features, such
as short motifs and structural sequence patterns, which serves as a significant complementary
signal.
175
Effect of microhomologies, non-canonical DNA structures, and motifs of varying
composition
We next sought to identify the more fine-grained sequence features which the CNN identified as
being associated with SVs. We performed a series of ablation analyses to quantify the effects of
microhomologies, non-canonical DNA structures, and sequence motifs of varying composition
on the predicted likelihood of SV formation. We first computed the enrichment of k-mers of
specific lengths in SV regions relative to background sequences and identified k-mers that were
significantly overrepresented near SV breakpoints. To test whether these enriched motifs
influence model predictions, we inserted each k-mer at random positions within background 400
bp windows (**Methods**). For each modified sequence, we computed the change in predicted SV
probability relative to the original unmodified background sequence and used the average
change as a measure of the k-mer's effect. We observed a positive correlation between k-mer
enrichment and the magnitude of prediction increase induced by insertion (**Fig 3A**), indicating
that the CNN has learned sequence patterns associated with elevated SV susceptibility. To
examine relationships among motifs with similar functional effects, we embedded k-mers in a 2-
dimensional representation space and found clusters of sequence-similar k-mers that produced
comparable effects on predicted SV probabilities, reflecting the contributions of specific
sequence patterns. We further identified a large cluster of k-mers that lead to a consistent
positive net increase in predicted SV probability and examined the shared features of these k-
mers. We found that they are significantly enriched in GC content and CpG islands, suggesting
a shared mechanism leading to structural instability (**Fig 3B**).
197
Given the well-established role of microhomology in double-strand break repair and SV
formation[12], we next quantified the model's sensitivity to short regions of breakpoint-spanning
homology. Empirically, SV regions showed a higher frequency of k-mers with higher homology
across breakpoints compared to background windows (**Fig 3C**). To determine if the CNN was
able to detect such microhomologies, we then inserted pairs of k-mers symmetrically around the
breakpoint at random distances from the center. These insertions changed the model probability
regardless of the similarity between kmers, indicating that the model is sensitive to artificial
looking DNA sequences. Remarkably, however, increased homology between kmers was
significantly associated with an increase in predicted SV probability (**Fig 3D**). This effect

increased in strength as a function of motif length. These results are consistent with the model
capturing sequence features relevant to microhomology-mediated repair.
209
Finally, because non-canonical DNA structures have been observed to be enriched within the
vicinity of SV breakpoints[20], we assessed the model's response to the insertion of sequence
elements capable of forming G-quadruplex (G4) and Z-DNA structures. We inserted these
motifs into background sequences, similar in approach to k-mer insertions, and compared the
resulting change in predicted SV likelihood against random control sequences of matched
length. Both G4 and Z-DNA insertions led to higher predicted SV probabilities (**Fig 3E**),
suggesting that the CNN recognizes sequence features associated with structures that can
increase fragility at SV breakpoints.
218
219   Distinct sequence signatures underlie different SV mechanisms, and together, these results
220   highlight that the CNN model accurately captures these features associated with SVs.
221   Furthermore, these features exhibit different scales; the CNN identified sequence interactions
222   indicative of microhomology as well as individual motifs associated with SVs.
223
Model performance across SV classes
SVs span a wide range of different sizes and are associated with many different mechanisms
(**Fig 4A**). We thus evaluated the performance of the models across different SV classes:
transposable element (TE) associated SVs, simple repeat associated, non-repeat associated,
and inversions. For each subclass, we trained the CNN, random forest, logistic regression, and
the ensemble model independently using the same procedures described above and compared
their predictive performance.
231
Prediction accuracy varied markedly across SV types (**Fig 4B**). SVs associated with simple
repeats were the most accurately predicted, with the CNN achieving an AUROC of 0.994,
slightly outperforming the random forest (0.992), logistic regression (0.986), and the ensemble
model (0.993). Similarly, TE-associated SVs showed strong sequence-driven predictability. For
SINE/Alu SVs, the CNN reached an AUROC of 0.925, substantially higher than the random
forest (0.614), logistic regression (0.626), and ensemble (0.807); LINE/L1 SVs showed a
comparable pattern (CNN: 0.823; random forest: 0.676; logistic regression: 0.677; ensemble:
0.745). These results indicate that TE-mediated SVs are strongly characterized by local
sequence signals rather than broader genomic context, and that sequence models can
accurately predict TE-mediated SVs. In contrast, the CNN underperforms against the random
forest model with non-repeat SV and in inversions. In particular, while none of the models
performed particularly well on inversions, the CNN was extremely poor (0.480). These results
highlight the varying importance of local sequence features to distinct SV classes.
245
To identify the most influential predictors for each SV class, we computed permutation
importances for all features within each subclass and compared them to importance rankings for
the full SV set (**Fig 4C**). CNN-derived sequence scores were among the strongest predictors for
LINE/L1 and especially SINE/Alu associated SVs, reinforcing the role of TE-specific sequence
patterns in driving these events. GC content was the dominant predictor for LINE/L1 SVs and

was also highly informative for retroposon/SVA SVs, consistent with known compositional
biases of these TE insertion targets. Distance to simple repeats was the strongest predictor for
simple repeat-associated SVs, reflecting the tendency of these events to arise within repetitive
regions[8].
255
Inversions showed a notably different signature. Many features contributed with similar
importance, and no single predictor dominated. The number of neighboring inversions emerged
as a strong feature, suggesting the presence of inversion hotspots. Distance to LINE/L1
elements was also informative, consistent with mechanisms in which L1 retrotransposition
facilitates rearrangements in adjacent sequences. The number of SNPs was additionally
predictive. Interestingly, several features exhibited negative permutation importance values,
suggesting that permuting these features slightly improved performance. This may indicate
redundancy among highly correlated predictors or that certain features add noise rather than
informative structure.

Distinct sequence and genomic feature distributions across SV classes
To evaluate how sequence features differ between SV regions and background genomic
windows, we compared the distributions of multiple genomic annotations across SV classes and
computed the Jensen-Shannon divergence (JSD) between SV and background distributions
(**Fig 4D**). Higher JSD values indicate greater distributional separation and therefore stronger
enrichment or depletion of that feature near SV breakpoints. Each feature includes the
distribution of all SVs for comparison.

Across all SVs, breakpoint windows consistently harboured higher local variant density than
background. SV regions contained more SNPs, small indels, and large insertions/deletions, with
moderate-to-high JSD values for each (around 0.30-0.50), indicating that SVs tend to arise in
already variant-rich regions, or that the SV formation can induce local mutations, which has
been previously observed[21]. This enrichment was particularly pronounced for inversions, which
showed the largest divergence in SNP counts (JSD = 0.667). In contrast, the number of
inversions in the surrounding window showed only modest shifts for most classes, but was
clearly elevated around inversion breakpoints themselves (JSD = 0.196), supporting the
presence of localized inversion hotspots.
281
Conservation and base composition also differed substantially between SV and background
windows. Average phyloP scores were markedly lower at SV breakpoints than in background,
especially for non-repetitive and simple-repeat SVs (JSD ≈ 0.7), consistent with reduced
evolutionary constraint at rearrangement-prone loci. TE-mediated SVs GC content likewise
showed strong divergence for several classes, including retroposon/SVA events and inversions
(JSD up to ~0.6), indicating that GC-rich or GC-poor environments contribute differentially to
specific SV mechanisms.
289
Overall, SVs occurred closer to simple repeats than background regions, with the largest effect
for simple repeat-associated SVs themselves (JSD = 0.570), confirming that these events are
highly concentrated within repeat-rich regions. Distances to Alu/SINE and L1/LINE elements
showed modest shifts when all SVs were considered together, but clearer class-specific biases

can be observed - LINE/L1 and SINE/Alu SVs, as well as inversions, exhibited substantial
divergence from background in their proximity to the two transposable elements. By contrast,
distances to the nearest exon showed relatively small JSD values across all classes. These
comparisons reveal that different SV classes occupy distinct regions of genomic feature space.
Overall, breakpoints are preferentially located in regions of elevated local variation, reduced
conservation, characteristic GC content, and specific arrangements of repetitive elements, with
particularly strong and class-specific signatures for repeat- and TE-associated SVs and for
inversions.
302
Gene functional constraint and tolerance
Structural variants play key roles in human disease and phenotypic variation[3] with strong
selection observed against large SVs[1]. However, establishing the relationship between
phenotype and genotype remains a key challenge in human genetics. Machine learning is
providing opportunities for understanding these high-dimensional relationships[22], but most
existing methods focus on SNV[23] while SVs are still underexplored. To assess the potential of
machine learning models for predicting the functional impact of SVs, we explore how our model
predictions interact with features such as allele frequency, genomic constraint, and variant effect
predictions.
312
Deleterious SVs are subject to negative selection and are thus enriched for rare variants[1]. We
thus examined the association between our model predictions and allele frequency. We note
that the training and testing tests were randomly split, alleviating any allele frequency biases in
the generation or testing of our model. Predicted SV probability increased with allele frequency
with common SVs receiving substantially higher predicted probability than rare SVs (**Fig 5A**).
This was true for both the CNN (p=1.6e-7) and ensemble models (p=1.2e-22), indicating that
the model may be able to identify features related to SV persistence or genomic mutability in the
population, thus favoring common SVs versus rarer ones. Indeed, phyloP score is strongly
correlated to both CNN and random-forest predicted probabilities (**Fig 2A**).
322
To further explore the relationship between constraint and our models, we examined model
predictions in genes with constraint quantified using LOEUF (gene loss-of-function
observed/expected upper bound fraction)[24] and pLI (probability of loss intolerance)[25]. LOEUF
increases as a function of relaxed constraint while pLI increases as a function of increased
constraint. We sampled 30 random locations within each gene and calculated the averaged
predicted SV probability. Model predictions over genes were strongly correlated with both
LOEUF (**Fig 5B,** ρ = 0.205, p-value = 4.42e-168) and pLI (**Fig 5C,** ρ = -0.166, p-value = 8.63e-
39) indicating the model is identifying features associated with SV persistence and likelihood.
Indeed, model predictions over genes were correlated with the number of true observed SVs
over those genes (**Fig 5D,** ρ = 0.330, p = 0). This indicates the model's ability to capture
genuine patterns of SV occurrence from genomic context alone, without variant or allele
frequency data. When genes were colored by their pLI values, we observed that even among
constrained genes, some loci sit in SV-prone genomic neighborhoods. However, highly
constrained genes tended to exhibit lower predicted SV probabilities. This suggests that beyond

raw SV density, the model is able to distinguish intrinsic sequence fragility from the selective
pressures that shape the persistence of SVs near essential genes.
339
Finally, we compared the relationship between CADD-SV variant prediction scores[26] and our
model predictions for all SVs in the test set. CADD-SV scores reflect putative variant
pathogenicity with higher scores indicating more deleterious impact. Both the CNN and
ensemble models showed significant negative correlations with CADD-SV PHRED scores (**Fig
5E**, CNN: ρ = -0.294, p = 0; Ensemble: ρ = -0.335, p = 0) indicating that predicted deleterious
SVs have lower probability in our model predictions. These observations show the model has
putative utility for variant effect prediction despite not being specifically designed for this
purpose.
348
Our analyses did reveal several outlier genes and SVs showing markedly higher predicted SV
probabilities than expected for their level of constraint or predicted variant impact (**Fig 5C, D**).
These outliers indicate genomic regions where local sequence architecture is inherently more
prone to structural instability despite strong functional constraint, suggesting possible
decoupling between sequence fragility and selective pressure. An example is the *TBC1D3B*
gene which exhibits extreme structural heterozygosity, with haplotypes differing by hundreds of
kilobases and in gene copy number[27]. This gene plays important roles in human brain evolution
and resides within highly variable segmental duplications. Similarly, Nanog genes are flanked by
Alu elements that modulate their regulation and expression[28]. Nevertheless, such sequence
features can predispose this region to SVs and highlight how certain loci may occupy genomic
environments predisposed to recurrent SV formation despite being under substantial functional
constraint.
361

DISCUSSION

In this study, we demonstrate that SV susceptibility in the human genome can be predicted with
high accuracy using local sequence context and genomic annotations. The strong performance
of both the CNN and the random forest model, and the further improvement achieved by
combining them in an ensemble, indicates that sequence-derived features contain substantial
information about the likelihood of SV formation in human genomes. These results provide
evidence that intrinsic genomic properties shape the distribution of SVs across the genome and
demonstrate the potential to quantify this susceptibility at base-pair resolution.

Our feature interpretation analyses provide insight into the specific genomic properties
associated with elevated SV likelihood. Correlation patterns were broadly consistent with
established mechanisms of genome instability, where regions with lower evolutionary
conservation and increased density of nearby genetic variants showed higher predicted SV
probabilities, reflecting relaxed functional constraint and a greater tolerance for rearrangement.
The strong influence of nearby SVs and SNPs, observed in both Gini and permutation
importances, suggests that SVs tend to accumulate in genomic neighborhoods predisposed to
breakage or rearrangement, forming local hotspot environments. This observation is supported
by empirical evidence highlighting meiotic breaks as key drivers of SNV and SV formation[21], as

well as segmental duplications[1,29,30]. Importantly, our analyses of the CNN model demonstrate
that local sequence patterns carry biological signals sufficient to identify such hotspots.

We performed extensive analyses to identify the features learned by the CNN model. These
analyses indicated that CNN can effectively identify short sequence signatures that contribute to
local fragility. These include microhomology sequences flanking SV breakpoints, as well as
sequences capable of forming non-canonical DNA structures, such as G-quadruplex and Z-
DNA, which can disrupt replication or repair processes. These findings suggest that the model
captures the landscape of genomic fragility shaped by local sequence composition.

The likelihood of structural variation is also shaped by distinct and interpretable genomic
signatures that differ across SV classes. The high predictive performance for simple repeat and
transposable element-associated SVs indicates that these events are strongly driven by local
sequence composition and recognizable motif patterns, rather than broader genomic context.
This reinforces mechanistic models in which polymerase slippage and retrotransposition
generate rearrangements at sequence-encoded instability sites. In contrast, inversions
displayed much lower CNN performance, suggesting that their formation depends less on short
motifs and more on larger-scale genomic architecture.

These findings suggest that human SV landscapes emerge in part from two interacting forces -
locally encoded fragility (driven by motifs, base composition, and repeats), and regional
tolerance for rearrangement (reflected in variant-rich and weakly conserved contexts). This
framework explains why sequence-based models are accurate at predicting repeat- and TE-
driven SVs but perform less effectively for inversions, which depend on long-range homology
not captured within a 400 bp window. By demonstrating class-specific predictability and
genomic determinism, we provide a foundation for SV risk annotation and suggest that
integrating broader structural context may further improve predictions for inversion and other
non-local SV mechanisms.

The relationship between model predictions, SV allele frequency, and gene constraint provides
insight into how the human genome is organized into regions of structural stability and fragility.
The higher predicted probabilities assigned to common SVs suggest that these variants arise in
genomic environments that are both intrinsically prone to rearrangement and evolutionarily
permissive to their persistence and that common SVs mark fragile sequence contexts. Regions
near LoF-tolerant genes showed higher predicted SV probabilities than those near constrained
genes. This association implies that the local sequence environments surrounding functionally
less essential genes tend to harbor the structural and compositional features linked to SV
formation. These findings suggest that the genomic landscape reflects a degree of coupling
between functional importance and structural permissiveness. However, this association,
though highly significant, is weak.

The overall predictive framework introduced here has several implications. First, it provides a
quantitative tool to estimate SV susceptibility at scale and identify functionally constrained
genes that are in SV-prone regions. Thus, these models can support studies seeking to
understand putatively pathogenic rearrangements and in the construction of SV mutability maps

of the human genome. Next, the identification of specific sequence determinants can inform
mechanistic studies into SV formation pathways. In addition, the relationship between predicted
SV probability, observed SV allele frequency, and functional constraints suggests that the model
highlights regions where rearrangements not only arise but also persist within human
populations, pointing to a potential role in long-term genome evolution.

Future work may extend this approach to additional structural variant types and to long-read
sequencing data that capture a broader spectrum of rearrangements. Integrating replication
timing, chromatin organization, or 3D genome architecture may further improve predictions,
particularly for complex events. Additionally, a major limitation of our models is that they
incorporate only highly localized sequence features, contributing to their failure to predict
specific classes of SVs accurately (e.g. inversions). Accounting for both long and short-range
sequence features and homology has the potential to vastly extend the utility of our approaches.
Finally, beyond the human genome, applying our framework across species may reveal
species-specific and conserved determinants of chromosomal instability providing insights into
the evolutionary forces shaping genome structure.

Overall, this work provides a genome-wide framework for quantifying SV susceptibility at
nucleotide resolution. The models and analyses presented advance our understanding of the
sequence architectures that predispose the genome to rearrangement, offering potential
applications in variant interpretation, disease gene discovery, and the identification of rare but
high-risk loci relevant to personalized genomics.
439

METHODS

Data sources
Structural variant calls were obtained from Phase 3 of the Human Genome Structural Variation
Consortium (HGSVC3), using variants reported against the GRCh38 reference genome. Variant
callsets included large insertions and deletions, large inversions, small indels, and SNPs. In the
various callsets, there are 111,803 large insertions, 64,428 large deletions, 300 inversions,
3,128,354 small insertions, 3,049,659 small deletions, and 23,462,179 SNPs.
447

448   Genomic annotations were retrieved from the UCSC Genome Browser, including phyloP
449   conservation scores, recombination rate estimates, RepeatMasker repeat annotations,
450   candidate cis-regulatory elements (cCREs), DNase accessibility, and transcription factor ChIP–
451   seq peak clusters. Reference genome sequences and gene annotations were obtained for
452   GRCh38.
453

Dataset construction
The final dataset consisted of 173,824 SVs, paired with an equal number of randomly sampled
background genomic locations. Only fully contiguous 400bp sequences are included in the
dataset. For each SV, we extracted a 400 bp context window centered on the breakpoint,
comprising 200 bp upstream and 200 bp downstream (**Fig 1B**). For deletions and inversions,

the internal deleted sequence was omitted from the window, where only the flanking sequences
were used.
461
To ensure that our approach does not introduce biases for deletions and inversions, we ran an
additional analysis to sample breakpoints reflecting their true occurrences. Background windows
for insertions consisted of the same 400 bp flanks surrounding the control positions. To
construct inversion-specific negative examples, we generated simulated inversions matched in
genomic length to the real events. First, we derived an empirical inversion length distribution
from the set of true inversions and randomly sampled an inversion length for each negative
example. A chromosome was selected at random, after which we sampled a first breakpoint
uniformly along the chromosome, ensuring that neither the first breakpoint nor the
corresponding second breakpoint (computed as first breakpoint + sampled inversion length)
overlapped any known SV breakpoint. For each simulated inversion, we extracted a single 400
bp sequence window consisting of 200 bp immediately upstream of the first breakpoint and 200
bp immediately downstream of the second breakpoint, mirroring the windowing strategy used for
true inversions. The internal sequence between the two breakpoints was excluded to match the
representation of real inversion events. Genomic annotation features were computed over the
two flanking windows and combined, producing negative examples that preserve the
chromosomal context, breakpoint structure, and inversion length distribution of true inversions
while ensuring no overlap with annotated SVs. Background windows for deletions were
generated with the same method. We find that there is little to no difference in the results we
obtained, compared to our initially described approach, for subsequent analyses.
481
Overall, this yielded 347,648 total samples (173,824 SVs and 173,824 background regions).
The dataset was partitioned into 60% training (n=208,588), 20% validation (n=69,530), and 20%
test sets (n=69,530), with no genomic positions shared across splits. All model selection and
hyperparameter optimization were performed on the training and validation sets. Only the test
set was used for final evaluation and was never seen during training. Genomic annotation
features - including conservation, GC content, repeat proximity, local variant density, gene
features, and regulatory annotations - were calculated or extracted for each window. Sequence
windows were one-hot encoded over four channels (ATCG). Spearman correlations across all
features, model predictions, and true labels were calculated with SciPy.
491
Convolutional neural network architecture
Sequence classification was performed using a custom 1D residual convolutional neural
network (CNN) implemented in PyTorch 2.0.1. The model accepts a 4-channel one-hot–
encoded 400 bp sequence. The architecture consists of an initial convolutional block of 1×1
convolution projecting 4 channels to 128, followed by batch normalization and ReLU activation.
This is followed by a stack of 2 residual blocks, each consisting of Conv1d (kernel size 11,
padding 5), BatchNorm1d, ReLU, second Conv1d (kernel size 11, padding 5), residual skip
connection, and final ReLU. A 1×1 convolution then reduces dimensionality to 32 channels. The
flattened output (12,800 units) is passed through a linear layer (128 units, ReLU), followed by a
final linear layer mapping to a single output neuron. A sigmoid activation produces a probability
estimate of SV occurrence.

The network was trained with binary cross-entropy loss, AdamW optimizer (with learning rate
5×10^{-6}, weight decay 1×10^{-3}), and a ReduceLROnPlateau scheduler (with factor 0.5, patience
3). Training used early stopping with patience of 5 epochs. The model was trained with batch-
based stochastic optimization. The weights corresponding to the lowest validation loss were
selected for all analyses.

Random forest, logistic regression, and ensemble models
In addition to the CNN, we trained a random forest classifier using scikit-learn 1.6.1, a logistic
regression baseline, and an ensemble model, implemented as a random forest receiving both
genomic annotation features and the CNN-predicted probabilities as input to improve predictive
performance.

For conventional machine-learning models, we combined the training and validation sets for
model fitting after CNN hyperparameter selection. All models were evaluated on the held-out
test set. Random forests used 1000 trees, balanced class weights, maximum number of
features when splitting as the square root of the total number of features, and default
bootstrapping. Logistic Regression was trained with L2 regularization.

Feature importance analysis
Two complementary measures of feature importance were used - Gini-based importance,
obtained from `sklearn.ensemble.RandomForestClassifier.feature_importances_`, and
permutation importance, computed using `sklearn.inspection.permutation_importance`, which
measures the decrease in model performance when individual features are permuted.
Permutation importance was calculated on the test set to provide an out-of-sample estimate,
with 5 repetitions. For analyses involving the ensemble model, the CNN probability was treated
as an additional feature and its importance was evaluated alongside annotation features.

K-mer, motif, microhomology, and non-canonical DNA structure analyses
To quantify the contribution of specific sequence patterns to SV likelihood, we performed
targeted insertion experiments using the trained CNN model. All analyses were conducted on
background (non-SV) 400 bp windows to ensure that observed prediction changes reflected the
inserted sequence element rather than surrounding SV-associated context.

K-mer insertion analysis: We evaluated the effect of short sequence motifs by inserting all 1,024
permutations of 5-bp k-mers into 50 random positions within each background window. After
each insertion, the sequence was truncated at the ends to maintain the fixed 400 bp window
centered on the reference position. For each k-mer, we computed the average change in
predicted SV probability across all insertions as a measure of its influence on model predictions.

Microhomology analysis: To assess the effect of short homologous tracts, we inserted 100 pairs
of k-mers - either perfectly matching (symmetric microhomology) or mismatched (asymmetric
controls) - symmetrically around a background location at 50 random distances from the center.
As with the k-mer analysis, sequence windows were truncated to 400 bp after insertion. The

change in predicted SV probability was averaged across all inserted microhomology pairs to
quantify the model's sensitivity to breakpoint-spanning homology.
549
Non-canonical DNA structure insertion: We next tested whether sequence elements capable of
forming non-B DNA structures influence predicted SV likelihood. We inserted 100 distinct
sequences predicted to form G-quadruplexes (G4) and 100 predicted Z-DNA -forming
sequences into background windows at 50 random positions each. To ensure proper controls,
we also generated 100 random sequences matched in length each to the G4 and Z-DNA motifs
and inserted these in parallel. All modified sequences were truncated to retain a fixed 400 bp
window centered at the reference site.
557
558 For all analyses, the mean change in CNN-predicted probability relative to the unmodified
559 background sequence was used as the quantitative measure of motif effect size.
560
561 **Model construction for SV subclasses**
562 Repeat annotations are performed using RepeatMasker version 4.1.5 on the SV callset. We
563 label the SV with the repeat annotation if the repeat covers at least 80% of the SV sequence. In
564 the case of multiple repeat annotations per SV, the repeat that spans at least 80% of the
565 repetitive content in the SV sequence is utilized. Otherwise, the SV is labelled non-repetitive.
566
To evaluate whether different structural variant classes exhibit distinct sequence determinants,
we trained separate models for individual SV subclasses. SVs were grouped into six categories:
inversions, SINE/Alu-associated SVs (n = 4,111), LINE/L1-associated SVs (n = 652), non-
repetitive SVs (n = 37,601), Retroposon/SVA SVs (n = 2,081), and simple-repeat SVs (n =
14,650). For each subclass, we constructed a balanced dataset in which the positive set
consisted exclusively of SVs from the focal subclass, and the negative set consisted of an equal
number of randomly sampled background locations drawn from regions accessible in the
reference genome. Context windows and annotations were extracted following the same
procedures described for the full dataset.
576
577 For each subclass, we trained the same set of models used in the main analysis - the CNN, the
random forest, logistic regression, and the ensemble model integrating the CNN prediction as
an additional feature. The training, validation, and test splits followed the same strategy as for
the full dataset, with no shared samples across splits. Model performance was assessed using
AUROC and additional standard metrics. For interpretability, we computed permutation
importances on the ensemble model with the test set for each SV subclass to identify the most
influential genomic features and to characterize class-specific sequence determinants.
583
584 **Constraint and tolerance**
585 Gene constraint analyses used the gnomAD v4.0 gene constraint dataset, specifically the
updated loss-of-function observed/expected upper bound fraction (LOEUF) metric and the
probability of being loss-of-function (LoF) intolerant (pLI) score. Genes were matched using
gene names from the gene annotation file and the gnomAD dataset. LOEUF and pLI values
were matched to quantify how local constraint relates to predicted SV probability. Spearman

correlations were calculated with SciPy. We quantified the number of SVs within each gene's
surrounding region (±200 bp).

Software and reproducibility
All analyses were conducted in Python 3.10.14, random seed of 42, PyTorch 2.0.1, scikit-learn
1.6.1, transformers 4.41.2, SciPy 1.14.1, RepeatMasker version 4.1.5, and standard scientific
libraries (NumPy, pandas, matplotlib). All code and processing pipelines were executed on
GPU-enabled hardware when available.
596

597   **DECLARATION OF INTERESTS**
598   The authors declare no competing interests.
599

ACKNOWLEDGEMENTS
This work was supported by NIH National Institute of General Medicine award
R35GM142916 to PHS and NIH National Human Genome Research Institute award
R01HG013017 to PHS. This manuscript is the result of funding in whole or in part by the
National Institutes of Health (NIH). It is subject to the NIH Public Access Policy. Through
acceptance of this federal funding, NIH has been given a right to make this manuscript publicly
available in PubMed Central upon the Official Date of Publication, as defined by NIH. The
content is solely the responsibility of the authors and does not necessarily represent the official
views of the National Institutes of Health.
609
610

611   **AUTHOR CONTRIBUTIONS**
612   PHS, NI, and DL conceived of the study. DL performed all analyses. DL, RNL, and PHS wrote
613   the manuscript.
614

615   **DATA AND CODE AVAILABILITY**
616   https://github.com/sudmantlab/sv_prediction.git
617

618   **FIGURES TITLES AND LEGENDS**
619

620
Figure 1 Details of the model and the prediction task. **(A)** Length distributions of insertions (left)
and deletions (right) called against GRCh38 from HGSVC Phase 3, colored by repeat type. **(B)**
Schematic of the model prediction task. At a target location in the human genome, models are
trained to predict the probability of a structural variant event given sequence contexts of 100bp,
400bp, and 2000bp. **(C)** Schematic of the model architectures and predictions. **(D)** Genomic
tracks along a 500kb region in chromosome 1 comparing SV density with a subset of genomic
features. Bins of 5kb are used. **(E)** Receiver operating characteristic and precision-recall curves
of the CNN, random forest, and the ensemble model, plotted against those of a baseline logistic
regression model. **(F)** Comparisons of AUROC across sequence context lengths and model
types.
631

632
Figure 2 Analysis of model features. **(A)** Spearman correlation coefficients between genomic
features, predicted SV probabilities, and true SV presence/absence of each model. **(B)** (Left)
Distribution of genomic features used in the ensemble model in SV and background regions.
Lines represent the interquartile range, and dots represent medians. Values are normalized to
plot on the same scale. (Right) The Gini importances indicating the extent to which the model
structurally relies on each feature for classification. Features are sorted with descending Gini
importance from top to bottom. **(C)** (Left) Permutation importances of genomic features used in
the ensemble model, which measures how much each feature affects model performance when

randomly permuted. (Right) Change in permutation importance between before and after adding
CNN predictions to the random forest model.
643

644
Figure 3 Effect of sequence motifs on CNN predictions. **(A)** Relationship between k-mer
(length=12) enrichment in SV vs background regions, and the change in predicted SV
probability post-insertion of each k-mer. A linear regression is shown as a dotted line, along with
Spearman correlation and p-value. **(B)** (Left) DNABERT token-space clustering of k-mers of
length 12, colored by change in SV probability post-insertion of each k-mer. The marked box
shows a cluster of k-mers that all induce a positive change in SV probability. (Right) Density
distributions of GC content and CpG count in the k-mers highlighted in the box and all plotted k-
mers. **(C)** Empirical frequency of k-mer (length=17) pairs that share various degrees of
similarity, across SV breakpoints (red) vs background locations (blue). Similarity scores
increase from left to right. **(D)** Relationship between similarity between k-mer pairs inserted
across a background location and mean change in predicted SV probability, for k-mers of length
10, 14, and 22. Similarities are binned, and fully symmetric insertions are included as the last
bin. Spearman correlation and p-value are labelled as well. **(E)** Mean change in predicted SV

probability after inserting G4-associated sequences vs. random sequences of G4-length, and Z-
DNA-associated sequences vs. random sequences of Z-DNA-length.
660

661
Figure 4 Model performance and feature distributions of SV subclasses. **(A)** Length profiles of
insertions (blue) and deletions (red) of each SV subclass. Length distribution of inversions are
shown as a separate plot. **(B)** Model AUROC performance across various subclasses and
model types, with the dotted line representing baseline model performance for all SVs. **(C)**
Permutation importance of features in the ensemble model for different SV subclasses. **(D)**
Distributions of various features across SV and background regions, categorized by SV
subclass. Feature distribution of all SVs are plotted in red for comparison.
669

670
671 **Figure 5** Interactions between model prediction and the selective constraint and functional
672 impact of SVs. **(A)** Relationship between allele frequency in HGSVC3 and predicted SV
673 probability of the CNN and ensemble model. Allele frequency is binned into singletons, low
674 frequency variants, and intermediate variants. **(B)** Relationship between gene LOEUF and
675 predicted SV probability of the CNN and ensemble. **(C)** Ensemble predicted probability plotted

against pLI, with outlier genes that have high constraint and high predicted SV probability highlighted in red. **(D)** Relationship between ensemble predicted probability and observed SV count in each gene region, colored by pLI. Points are jittered slightly along the x-axis for visualization. **(E)** Correlation between CADD-SV PHRED-scores and model predictions from both the CNN and ensemble. Density plots are plotted for each axis. A linear regression is shown as a dotted line.

REFERENCES

1. Sudmant, P. H. *et al.* An integrated map of structural variation in 2,504 human genomes. *Nature* **526**, 75–81 (2015).
2. Logsdon, G. A. *et al.* Author Correction: Complex genetic variation in nearly complete human genomes. *Nature* **645**, E6 (2025).
3. Weischenfeldt, J., Symmons, O., Spitz, F. & Korbel, J. O. Phenotypic impact of genomic structural variation: insights from and for human disease. *Nat Rev Genet* **14**, 125–138 (2013).
4. Sanders, S. J. *et al.* Multiple recurrent de novo CNVs, including duplications of the 7q11.23 Williams syndrome region, are strongly associated with autism. *Neuron* **70**, 863–885 (2011).
5. Walsh, T. *et al.* Rare structural variants disrupt multiple genes in neurodevelopmental pathways in schizophrenia. *Science* **320**, 539–543 (2008).
6. Bolognini, D. *et al.* Recurrent evolution and selection shape structural diversity at the amylase locus. *Nature* **634**, 617–625 (2024).
7. Perry, G. H. *et al.* Diet and the evolution of human amylase gene copy number variation. *Nat Genet* **39**, 1256–1260 (2007).
8. Carvalho, C. M. B. & Lupski, J. R. Mechanisms underlying structural variant formation in genomic disorders. *Nat Rev Genet* **17**, 224–238 (2016).
9. Payer, L. M. & Burns, K. H. Transposable elements in human genetic disease. *Nature Reviews Genetics* **20**, 760–772 (2019).
10. Lee, J. A., Carvalho, C. M. B. & Lupski, J. R. A DNA replication mechanism for generating nonrecurrent rearrangements associated with genomic disorders. *Cell* **131**, 1235–1247 (2007).
11. Lieber, M. R. The mechanism of double-strand DNA break repair by the nonhomologous DNA end-joining pathway. *Annu Rev Biochem* **79**, 181–211 (2010).
12. Ottaviani, D., LeCain, M. & Sheer, D. The role of microhomology in genomic structural variation. *Trends Genet* **30**, 85–94 (2014).
13. Badet, T., Fouché, S., Hartmann, F. E., Zala, M. & Croll, D. Machine-learning predicts genomic determinants of meiosis-driven structural variation in a eukaryotic pathogen. *Nature Communications* **12**, 3551 (2021).
14. Fairley, S., Lowy-Gallego, E., Perry, E. & Flicek, P. The International Genome Sample Resource (IGSR) collection of open human genomic variation resources. *Nucleic Acids Res* **48**, D941–D947 (2020).
15. Jaganathan, K. *et al.* Predicting Splicing from Primary Sequence with Deep Learning. *Cell* **176**, 535–548.e24 (2019).
16. Kelley, D. R. *et al.* Sequential regulatory activity prediction across chromosomes with convolutional neural networks. *Genome Res* **28**, 739–750 (2018).
17. Zhou, J. & Troyanskaya, O. G. Predicting effects of noncoding variants with deep learning–based sequence model. *Nature Methods* **12**, 931–934 (2015).

18. Agarwal, V. & Shendure, J. Predicting mRNA Abundance Directly from Genomic Sequence
Using Deep Convolutional Neural Networks. *Cell Rep* **31**, 107663 (2020).
19. Kelley, D. R., Snoek, J. & Rinn, J. L. Basset: learning the regulatory code of the accessible
genome with deep convolutional neural networks. *Genome Res* **26**, 990–999 (2016).
20. Makova, K. D. & Weissensteiner, M. H. Noncanonical DNA structures are drivers of
genome evolution. *Trends Genet* **39**, 109–124 (2023).
21. Hinch, R., Donnelly, P. & Hinch, A. G. Meiotic DNA breaks drive multifaceted mutagenesis
in the human germ line. *Science* **382**, eadh2531 (2023).
22. Benegas, G., Ye, C., Albors, C., Li, J. C. & Song, Y. S. Genomic language models:
opportunities and challenges. *Trends Genet* **41**, 286–302 (2025).
23. Benegas, G., Albors, C., Aw, A. J., Ye, C. & Song, Y. S. A DNA language model based on
multispecies alignment predicts the effects of genome-wide variants. *Nat Biotechnol* (2025)
doi:10.1038/s41587-024-02511-w.
24. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in
141,456 humans. *Nature* **581**, 434–443 (2020).
25. Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**,
285–291 (2016).
26. Kleinert, P. & Kircher, M. A framework to score the effects of structural variants in health
and disease. *Genome Res* **32**, 766–777 (2022).
27. Guitart, X. *et al.* Independent expansion, selection, and hypervariability of the gene family in
humans. *Genome Res* **34**, 1798–1810 (2024).
28. González-Rico, F. J. *et al.* Alu retrotransposons modulate Nanog expression through
dynamic changes in regional chromatin conformation via aryl hydrocarbon receptor.
Epigenetics Chromatin **13**, 15 (2020).
29. Campbell, C. D. *et al.* Population-genetic properties of differentiated human copy-number
polymorphisms. *Am J Hum Genet* **88**, 317–332 (2011).
30. Vollger, M. R. *et al.* Increased mutation and gene conversion within human segmental
duplications. *Nature* **617**, 325–334 (2023).

753