Complete NMR assignment for 275 of the most common dipeptides in intrinsically disordered proteins

Tobias Rindfleisch[1,2,3,*]**, Emilie Fjeldberg Taule**[2]**, Markus S. Miettinen**[1,2,*,†]**, and Jarl Underhaug**[2,*,†]

[1]Computational Biology Unit, Department of Informatics, University of Bergen, 5008 Bergen, Norway
[2]Department of Chemistry, University of Bergen, 5007 Bergen, Norway
[3]Max Planck Institute of Molecular Plant Physiology, 14476 Potsdam, Germany
[*]corresponding authors: Tobias Rindfleisch (tobias.rindfleisch@uib.no), Markus S. Miettinen
(markus.miettinen@iki.fi), Jarl Underhaug (jarl.underhaug@uib.no)
[†]these authors contributed equally to this work

ABSTRACT

Accurate NMR chemical shift assignments are essential for atomic-resolution characterization of proteins. Especially for intrinsically disordered proteins (IDPs) and regions (IDRs), however, the assignment remains a labor-intensive task due to spectral overlap and conformational heterogeneity. Consequently, complete side-chain assignments are rare. Here, we present a comprehensive reference dataset, comprising the complete NMR chemical shift assignments for 275 of the most prevalent dipeptides in the IDPome, covering 93% of it. The dataset contains all NMR-accessible backbone and side-chain nuclei, in total 9 408 validated data points, as well as the 1D (^{1}H, ^{13}C) and 2D (^{1}H-^{15}N HSQC, ^{1}H-^{13}C HSQC, TOCSY, NOESY, ^{1}H-^{13}C HMBC) spectra used for assignment, making it a rich resource for the training, testing, and benchmarking of tools for data-driven protein assignment, peak picking, and synthetic spectrum generation. To facilitate such machine learning applications, all data are delivered in standardized, machine-readable formats.
(The DOI link to access the dataset will be activated once the article is accepted.)

Background & Summary

Accurate chemical shift assignments lay the foundation for nuclear magnetic resonance (NMR) analysis, enabling the structural and dynamic characterization of proteins at atomic resolution. However, the assignment process remains one of the most labor-intensive steps in NMR studies, often complicated by signal overlap, peak broadening, and conformational heterogeneity[1,2]. These issues are particularly pronounced in intrinsically disordered proteins (IDPs), where the lack of a single stable tertiary structure leads to spectral congestion and poor chemical shift dispersion[3,4]. As a result, complete side-chain assignments are rarely achieved for IDPs, leaving a substantial gap in available reference data. To address this gap, a consistent and comprehensive reference set would be required, particularly one that includes all side-chain nuclei. Here, rather than relying on heterogeneous IDP datasets with incomplete assignments, we adopted a reductionist approach and focused on the smallest nontrivial building blocks of proteins—the dipeptides.

The 20 canonical amino acids, each with its unique physicochemical properties, are the elementary building blocks of proteins. Within these large macromolecules, responsible for numerous essential functions in all aspects of life, already the simple connection of just two L-amino acids via a peptide bond leads to 400 possible combinations for L-α-dipeptides. So far, these smallest nontrivial units have been poorly studied in comparison to single amino acids, longer peptides, and complete proteins[5]. However, the key characteristics of polypeptides, such as solubility and reactivity in enzymatic context, emerge from the comprising dipeptides, whose torsion angle preferences naturally also influence the protein folding tendencies. Concerning the specific case of IDPs, dipeptides are known to exhibit structural and dynamic properties similar to those of disordered chains[6], providing an ideal minimal model for exploring disorder-related phenomena.

IDPs lack a unique and fixed tertiary (3D) structure, mainly because of the underrepresentation of hydrophobic residues, which prevents the formation of a hydrophobic core, coupled with the enrichment of charged and polar amino acids, which attracts protein–solvent interactions[7,8]. However, their broad structural ensembles do possess characteristic sequence-dependent distributions over the conformational space[4]. Not all IDPs are fully unstructured; the term also includes proteins with intrinsically disordered regions (IDRs) embedded in otherwise well-folded domains[9]. Bioinformatic analyses predict more than 40% of

eukaryotic proteins to contain IDRs[10] with approximately two thirds of these being fully disordered[11]. Achieving atomistic-resolution characterization of IDP ensembles remains a major challenge, as their continuous rapid interconversion between innumerable conformations prevents the use of traditional structure-determination techniques such as X-ray crystallography or cryo-electron microscopy[8,12–14].

Among experimental techniques, NMR spectroscopy uniquely enables atomistic interrogation of IDP ensembles[15], especially when coupled with all-atom molecular dynamics (MD) simulations[16–20]. Achieving this, however, requires precise and reliable NMR assignments, in which observed chemical shifts are unambiguously matched to their corresponding atoms. Previous studies on mono-[21], di-[6], tri-[22], and small oligopeptides[23] have shown most of these molecules to reflect the behavior of IDPs, suggesting dipeptides as the simplest unit that can capture the fundamental sequence-dependent correlations governing the intrinsic disorder.

Here, we provide complete chemical shift assignments for 275 of the most frequently occurring dipeptides in the IDP proteome (IDPome), covering 93% of it. The dataset, comprising 9 408 unique quality-controlled data points, includes all NMR-accessible nuclei, in both the backbone and side-chains, thereby bridging the IDP-side-chain data gap. Crucially, we also provide the complete set of acquired and processed spectra—1D (^{1}H and ^{13}C) as well as 2D (^{1}H-^{15}N HSQC, ^{1}H-^{13}C HSQC, TOCSY, NOESY, and ^{1}H-^{13}C HMBC)—used for assignment. These enable the development and benchmarking of algorithms for automated all-atom assignment of IDPs and the generation of synthetic NMR spectra for complete protein sequences. The dataset also provides a benchmark for improving automated peak-picking methods, including the deconvolution of overlapping resonances. By delivering standardized, high-quality, machine-readable data, this resource advances reproducibility in both computational and experimental NMR studies, facilitating progress across the protein disorder spectrum.

Methods

Sample selection

To ensure a comprehensive representation of IDPs and efficient use of resources, we aimed to include the most frequently occurring dipeptides in the IDP proteome. This strategy inherently excludes the most hydrophobic dipeptides, minimizing potential solubility issues and allowing all samples to be measured under standardized experimental conditions using a single, uniform solvent.

To evaluate the occurrence of all 400 canonical dipeptides in the IDPome, we analyzed the DisProt[24] database (disprot.org, version 2025-06) that exclusively contains experimentally confirmed IDP sequences. For each dipeptide, we counted the absolute occurrence across all non-redundant sequences and determined the corresponding relative frequencies. As DisProt includes redundant entries derived from multiple experimental reads under the same protein-sequence identifier (ID), only the first sequence associated with each ID was considered (scripts are provided; see section Code Availability).

The measurement order followed the frequency ranking; all proline-containing dipeptides were included due to the distinctive conformational characteristics of proline in IDPs[25,26].

Sample preparation

Unlabeled dipeptides with N-terminal acetylations and C-terminal amidations were purchased from Peptide Protein Research Ltd, Fareham, United Kingdom. The capping groups were added to eliminate the terminal charges and to introduce a third peptide bond as well as a third C$^\alpha$, thereby mimicking the chemical context within a polypeptide chain. All lyophilized samples were dissolved unbuffered in 90% H$_2$O, 10% D$_2$O, and 15 μM DSS (sodium trimethylsilylpropanesulfonate) at concentrations of approximately 5 mg/mL. The measurements were performed in 5-mm Bruker (Billerica, MA, United States) NMR tubes with a volume of 600 μL per sample.

NMR experiments

High-throughput NMR measurements for dipeptide assignments were performed on two Bruker BioSpin spectrometers (Table S1): an Ascend 600 MHz incorporating a QCI-P CryoProbe (219 samples) and an Ascend 850 MHz with a TCI CryoProbe (56 samples), both equipped with an AVANCE NEO console. Samples were handled automatically with a SampleJet sample changer and equilibrated to 298 K for at least 2 minutes prior to measurement and otherwise stored at 4°C. Table 1 lists the performed NMR experiments, along with key acquisition parameters.

Spectra processing and assignments

Acquired NMR spectra were processed using the MestReNova software (version 14.02.0-26256, Mestrelab Research) and referenced to DSS. Bruker datasets (Bruker BioSpin), containing the free induction decays (FIDs), were imported into MestReNova and processed with standard routines; the processing steps included zero-filling (doubling the number of data points in the indirect dimension), Fourier transformation, application of window functions, phase correction, and baseline correction. For 1D spectra, an exponential multiplication (EM) window function was applied with a line broadening of 0.5 Hz

NMR Experiment	Pulse Sequence	Number of Scans	1TD (F1) Points	2TD (F2) Points	Spectral Width [ppm]	Spectral Center [ppm]	NUS [%]
1D ^{1}H (zgesgppe)	zgesgppe	8	65 536	-	11.9	4.698	-
1D ^{1}H (1D NOESY)	noesygppr1d	8	142 856	-	29.8	4.698	-
1D ^{1}H (q-NMR)	noesygppr1d	8	142 856	-	29.8	4.698	-
1D ^{13}C (C13CPD)	zgpg30	256	65 536	-	237	100	-
2D ^{1}H–^{15}N HSQC	hsqcetfpf3gpsi	2	258	2048	35 (F1), 16.0 (F2)	101 (F1), 4.698 (F2)	25
2D ^{1}H–^{13}C HSQC	hsqcedetgpsp.3	2	512	2048	180 (F1), 15.1 (F2)	90 (F1), 4.698 (F2)	25
2D ^{1}H–^{1}H TOCSY	mlevphpr.2	8	256	2048	10.4 (F1), 10.4 (F2)	4.698 (F1), 4.698 (F2)	25
2D ^{1}H–^{1}H NOESY	noesyphpr	8	64	2048	10.4 (F1), 10.4 (F2)	4.698 (F1), 4.698 (F2)	50
2D ^{1}H–^{13}C HMBC	hmbcgplpndprqf	32	1024	4096	180 (F1), 23.2 (F2)	100 (F1), 4.698 (F2)	6.25

Table 1. Overview of selected acquisition parameters for dipeptide NMR assignment experiments: number of scans (NS), number of acquisition points in the F1 and F2 dimensions (1TD and 2TD), spectral width (SW), spectral center (O1P), and the applied percentage of non-uniform sampling (NUS). All pulse sequences were included in TopSpin (version 4.1.4, Bruker BioSpin). For the complete list of acquisition and processing parameters per sample see the Edmond repository https://doi.org/10.17617/3.CBCNQG.

or lower for ^{1}H and 1 Hz for ^{13}C. For 2D spectra, a squared cosine function was used. All spectra were baseline corrected using a third-order polynomial fit, except for the 1D ^{13}C (C13CPD) experiments, where a Whittaker smoother was applied. In 2D experiments, the residual water resonance was suppressed using MestReNova's signal suppression tool, selecting the solvent (water) frequency and applying a narrow suppression filter to remove the water-signal interference.

Chemical shifts of dipeptides were manually assigned for all detectable ^{1}H, ^{13}C, and ^{15}N nuclei using established sequential assignment strategies. Exact values for ^{1}H and ^{13}C chemical shifts were derived from the corresponding 1D spectra and ^{15}N shifts were obtained from the F1 (^{15}N) dimension of ^{1}H–^{15}N HSQC experiments. Clearly identifiable multiplets were analyzed with MestReNova's multiplet manager, and final assignments were validated by cross-checking correlations among several spectra.

NMR spectra with respective acquisition and processing parameters were stored in the machine-readable NMReData[27] format, and the corresponding peak lists were exported as Structure Data Files (SDF) and converted to the NMR-STAR[28], YAML, and chemical shift table formats, all compliant with the Biological Magnetic Resonance Bank (BMRB, bmrb.io) nuclei naming standards[29].

Data Records

The chemical shift datasets for 275 of the most common dipeptides in IDPs, along with the corresponding NMR spectra (Table 1), are permanently openly available in the Edmond Open Research Data Repository (edmond.mpg.de) entry: https://doi.org/10.17617/3.CBCNQG.

The top level of the deposition contains a directory with all the scripts used in this study, along with a second directory including 20 subdirectories named after the canonical amino acids, each representing the first residue of the dipeptides. Within each of the 20 directories, the corresponding dipeptide datasets are stored. Each dipeptide dataset includes the acquired, processed, and analysis-related files in machine-readable formats (see Table 2), facilitating reuse of the NMR data.

The scripts directory contains a fully automated Snakemake[30] workflow that executes the complete data-processing, analysis, and visualization pipeline. In addition, it contains three subdirectories that include the individual scripts used for (i) dipeptide ranking, (ii) creation of the dipeptide dataset files, and (iii) generation of all figures presented in this publication.

Data Overview

The frequencies of all 400 canonical dipeptides in IDPs are visualized in Figure 1 and listed in Table S2. These frequencies were derived from an analysis of 3 201 non-redundant and experimentally confirmed IDP sequences included in the DisProt

File Type	Extension	Description
MestReNova project	.mnova	Original MestReNova project file (version 14.02.0-26256, Mestrelab Research) documenting manual chemical shift assignments. Enables inspection and reproducibility.
NMReData project	_nmredata.zip	Contains all acquired and processed spectra, including complete acquisition and processing parameters (Table 1) in the machine-readable NMReData[27] format.
NMReData molecular structure	_nmredata.sdf	Machine-readable Structure Data File (SDF)[31] reporting the molecular structure of each dipeptide, assignable chemical shifts, and experimental metadata. Included in _nmredata.zip.
PDB structure file	.pdb	Provides dipeptide structures in machine-readable Protein Data Bank (PDB) format[32] using the IUPAC atom naming. Heavy atom order matches the SDF file for direct mapping.
^{15}N assignment table	_15N_shifts.txt	As MestReNova cannot technically store ^{15}N chemical shifts, they are provided separately as a plain-text table with atom numbering consistent with the SDF file.
Complete assignments	_Chemical_Shifts.txt	Comprehensive chemical shift tables for each dipeptide (machine-readable), following the BMRB atom-naming convention[29].
Complete assignments	_Chemical_Shifts.yml	Data from the complete assignment table in YAML format, a human- and machine-readable serialization format. Atom naming follows the BMRB convention[29].
Complete assignments	_nmrstar.str	Experimental metadata and chemical shifts in machine-readable NMR-STAR format[28] used by the BMRB.

Table 2. Overview of the deposited dipeptide dataset files and their contents.

database[24]. These contained 221 637 individual dipeptides, providing a quantitative overview of dipeptide occurrences across the IDP proteome. The 275 dipeptides we study here collectively account for 93% of the IDPome, faithfully reflecting the characteristic amino acid composition and physicochemical properties of disordered proteins and regions.

Dipeptide frequencies are non-uniformly distributed, with only a small subset of a few dozen examples approaching the 0.25% average occurrence. Approximately 30% of dipeptides appear more frequently than this, while about 55% occur more seldomly (Figure S1). The 86 most common dipeptides—roughly one fifth of all the 400—account for half of the entire IDPome (Figure S2), underscoring the compositional bias in disordered proteins. As expected, the most frequently observed dipeptides contain small and/or non-hydrophobic residues (Figure S1 and Table S2), whereas those incorporating one large hydrophobic residue are less common and those containing two are rare. Cysteine- and tryptophan-containing dipeptides are among the least represented, consistent with their structure-stabilizing roles that are incompatible with the disordered nature of IDPs[33]. Consequently, although roughly 100 dipeptides—one quarter of all possible combinations—contain large hydrophobic residues or cysteine, they collectively account for less than 7% of the IDP proteome, resulting in high coverage of the presented dataset.

Note, however, that a few high-ranking dipeptides were left out of the dataset, because the samples either (i) contained substantial impurities (AT) preventing reliable chemical shift assignments, (ii) were insoluble (FL, IV, IN) under the chosen solvent conditions, or (iii) exhibited extensive peak overlap (II).

Technical Validation

All chemical shifts were assigned manually and systematically verified across multiple spectra. Any ambiguous cases were discussed and resolved by at least two independent researchers to minimize subjective bias. To ensure completeness and consistency, a verification script was used to examine the assignment tables for missing or malformed entries, confirming that all the expected NMR-active nuclei were present (scripts are provided; see section Code Availability). The measurement uncertainty of chemical shifts under constant experimental conditions is typically minimal to negligible[34–36], and referencing to DSS ensures that systematic errors are effectively eliminated. Cross-validation was achieved using a comprehensive set of 2D experiments (Table 1), with TOCSY connectivity ensuring that assignments between identical residues in homodipeptides could not be inadvertently swapped.

We further validated the assigned chemical shifts by direct comparison to the BMRB's amino-acid-wise statistics[29] (bmrb.io, accessed October 2025), which include a BMRB-wide average for every detectable NMR-active nuclei: For each assignable nucleus across the 275 dipeptides, we calculated the difference between the observed chemical shift (δ_{obs}) and the corresponding

Figure 1. Relative frequency distribution of the 400 canonical dipeptides in the IDPome, as derived from the DisProt[24] database, compared to the uniform distribution (red line). Blue bars show dipeptides included in the presented dataset, gray the non-included ones; key shows also the total percentages of both populations. Figure S1 shows the same plot with the explicit dipeptide names instead of the rank on the x-axis. Figure S2 shows the cumulative relative frequency as function of the dipeptide rank. Table S2 connects the dipeptide rank with the dipeptide names and additional details, including relative frequency and absolute occurrence.

BMRB average shift (reference; δ_{ref}) as $\Delta\delta = \delta_{\text{obs}} - \delta_{\text{ref}}$. Figure 2 shows the individual nuclei-distributions of $\Delta\delta$ (points) for each of the 20 amino acid residues, with shaded intervals marking the standard deviation (SD) of the corresponding BMRB reference distribution as ±1SD (light gray) and ±2SD (dark gray). All our observed data points are within ±2SD, and 95.8% within ±1 SD of the BMRB reference (Figure 2). This indicates a minimal likelihood of misassignments, which is further supported by the absence of major outliers (Figure 2); only a few, reliably explainable exceptions are present.

Backbone ^{15}N nuclei exhibit the largest systematic deviations, reflecting their dependence on the preceding residue[37]. Note that in the ^{1}H–^{15}N HSQC experiments, the ^{15}N spectral range did not cover the chemical shifts of arginine N$^\varepsilon$ nuclei (signal at roughly 85 ppm), leading this resonance to appear near 120 ppm due to aliasing (folding)[38]; we corrected this by subtracting the spectral width (35 ppm) from the observed resonances. As this correction is frequently omitted in the BMRB, biasing the averages reported there toward higher values, we used 85 ppm as the reference value for arginine N$^\varepsilon$ nuclei in our technical validation (Figure 2). Another common issue in BMRB reference data concerns the side-chain amide groups of asparagine (H$^{\delta21}$ and H$^{\delta22}$) and glutamine (H$^{\varepsilon21}$ and H$^{\varepsilon22}$), where the two protons are often swapped in assignments. To avoid this ambiguity, we used the centers of the BMRB-reported H$^{\delta21}$/H$^{\delta22}$ and H$^{\varepsilon21}$/H$^{\varepsilon22}$ populations as the reference values.

Closer examination reveals consistent sequence-dependent trends. Across all residue-specific plots, H$^\alpha$, C, and C$^\alpha$ have a single data point deviating from the main cluster(s). All these arise from dipeptides with proline in the second position (X–PRO-motif, with X being any amino acid; Figure S3). This systematic behavior reflects the altered chemical environment introduced by proline, which lacks an amide proton and modifies the N–C connectivity within the peptide bond.

The main clusters of carbonyl C, backbone amide N, and C$^\alpha$ chemical shifts separate into two distinct populations (Figure S4), corresponding to the residue's position within the dipeptide (N-terminal versus C-terminal). The absence of such splitting beyond C$^\beta$ in side chains is consistent with the expectation that residual-positional effects should primarily affect the backbone nuclei.

Distinct ^{1}H resonance populations are consistently observed near aromatic residues (PHE, TRP, and TYR), reflecting the influence of a local environment with an increased electron density. This effect affects all nuclei of residues connected to an aromatic residue and is most pronounced for H$^\beta$ and H$^\gamma$ nuclei, showing decreased ^{1}H chemical shifts. Smaller but still pronounced effects are observed on backbone H, H$^\alpha$, H$^\delta$, and H$^\varepsilon$ nuclei (Figure S4).

The effect of non-adjusted pH during sample preparation—enabling a straightforward use of the data for benchmark systems in molecular modeling—did not introduce marked uncertainty in the determination of the exact chemical shift values. For dipeptides with non-charged side chains, this was expected, as the pH remains close to 7.0 after dissolving the capped, and thus non-charged, samples. But also for dipeptides containing charged residues the observed protonation states correspond to those expected at pH 7, indicating that the chemical environments remained largely unchanged in terms of electrostatic interactions within the dipeptides. Furthermore, no major divergence in BMRB-difference distribution widths are observed between charged and non-charged residues across the ^{1}H, ^{13}C, and ^{15}N shifts (Figure 2). Overall, the BMRB-difference distributions are narrow, particularly when residue-position effects are taken into account (Figure S4).

In summary, these validations corroborate the technical quality and reliability of the reported chemical-shift dataset.

Figure 2. Comparison to BMRB: The 20 residue-specific panels show $\Delta\delta = \delta_{\mathrm{obs}} - \delta_{\mathrm{ref}}$ for each assignable nucleus, defined as the difference between the observed chemical shifts (δ_{obs}) and the corresponding BMRB reference values[29] (δ_{ref}). Note that each panel has two y-axis scales, since ^{1}H chemical shifts (blue data points and labels) span a much narrower range than ^{13}C and ^{15}N (red). Shaded regions represent the standard deviation (SD) of the respective BMRB reference distributions[29]: ± 1 SD (light gray) and ± 2 SD (dark gray). The total number of observations (i.e. measured dipeptides) per amino acid residue is given as N. ^{1}H resonances are classified as H, Q, and M, where H represents protons in CH groups, Q corresponds to both CH$_2$ protons, and M to methyl protons.

Usage Notes

The presented dataset delivers complete and standardized chemical shift assignments for 275 of the most common dipeptides found in IDPs, establishing a unique reference resource comprising 9 408 rigorously quality-controlled data points. As the corresponding 1D and 2D spectra are included, it further supports robust training and validation of artificial intelligence (AI) and machine learning approaches in protein NMR spectroscopy.

In contrast to most protein NMR datasets, which typically report only backbone resonances together with C^β/H^β nuclei, our dataset provides complete assignments for all NMR-accessible nuclei in each dipeptide. To our knowledge, this represents the first protein-related dataset that achieves truly all-atom assignments, thereby bridging the side-chain data gap. Notably, the chemical shift distributions of longer side-chain atoms (Figure S5) are remarkably narrow, reflecting sequence independence of these chemical shifts, which supports the feasibility of extending all-atom assignment and prediction strategies to disordered proteins.

In addition to the chemical shift assignments, the dataset includes the complete set of acquired and processed spectra for all the 275 dipeptides. For each sample, eight spectra are provided, covering 1D ^{1}H, 1D ^{13}C, and a diverse range of 2D experiments, including ^{1}H–^{15}N HSQC, ^{1}H–^{13}C HSQC, TOCSY, NOESY, and ^{1}H–^{13}C HMBC. This comprehensive spectral coverage greatly expands the scope of potential applications: The data can be used to train models not only to perform atom-specific assignments from experimental spectra, but also to generate synthetic NMR spectra for entire protein sequences, encompassing all detectable nuclei. Beyond these applications, the dataset provides a benchmark resource for advancing peak-picking algorithms, with particular potential to improve the deconvolution of overlapping peaks—a long-standing challenge in protein NMR spectroscopy[1,2].

We expect the dataset to be especially valuable for advancing AI methods tailored to IDPs, a class of proteins whose conformational heterogeneity poses significant challenges for traditional NMR approaches. The dataset also provides a rigorous benchmark for comparative studies, enabling systematic evaluation of existing NMR methods in terms of accuracy, completeness, and robustness. Its standardized nature ensures reproducibility and facilitates integration with other structural biology resources—such as BMRB[29], due to the use of the NMR-STAR-format[28].

Files reporting chemical shifts (_Chemical_Shifts.txt, _Chemical_Shifts.yml, and _nmrstar.str; see Table 2) are provided in widely used formats and can be inspected with any standard text editor. For programmatic access, _Chemical_Shifts.txt files can be imported in Python using the `pandas` package, _Chemical_Shifts.yml files with `PyYAML`, and _nmrstar.str files with `pynmrstar`. All acquired and processed spectra, together with full acquisition and processing parameters (Table 1), are stored in NMReData format (_nmrdata.zip). This standardized format facilitates interoperability and can be opened in several software environments, including MestReNova, TopSpin (Bruker BioSpin), and the NMReData project tools[27] (NMReData Initiative). In addition, programmatic access and downstream processing can be carried out in Python using the `nmrglue` package[39]. Examples for programmatic access are given in the provided scripts (see section Code availability).

Data availability

The dataset containing the chemical shifts for 275 of the most common dipeptides in IDPs, along with the corresponding NMR spectra used for their assignment, is permanently and openly available in the Edmond Open Research Data Repository (edmond.mpg.de) under the entry: https://doi.org/10.17617/3.CBCNQG.

Code availability

The code of the complete computational pipeline—including (i) the analysis of dipeptide occurrence frequencies in DisProt, (ii) the generation of NMR chemical-shift report files (_Chemical_Shifts.txt, _Chemical_Shifts.yml, and _nmrstar.str; see Table 2), and (iii) the creation of all figures—is permanently and publicly available at the Edmond repository (edmond.mpg.de), entry: https://doi.org/10.17617/3.CBCNQG.

The implementation within a reproducible Snakemake[30] workflow ensures transparent and automated execution of all analysis and visualization steps. The code used to generate Figure 2 is additionally provided as an example template for accessing the dataset.

Versions 4.3, 4.4, and 4.5 of TopSpin (Bruker BioSpin) and version 14.02.0-26256 of MestReNova (Mestrelab Research) were used for data acquisition and processing, while Python 3.9.13 was employed to generate all figures and files reporting chemical shifts. The following Python packages were used in this study: Biopython (1.78), Matplotlib (3.5.2), NumPy (1.21.5), os (3.9.13), Pandas (1.4.4), pynmrstar (3.3.5), re (3.9.13), tqdm (4.64.1), PyYAML (6.0), and Zipfile (3.9.13). These details are also included in the headers of the provided scripts.

References

1. Bax, A. & Grzesiek, S. Methodological advances in protein NMR. *Accounts Chem. Res.* **26**, 131–138 (1993).

2. Dyson, H. J. & Wright, P. E. Unfolded proteins and protein folding studied by nmr. *Chem. Rev.* **104**, 3607–3622 (2004).

3. Brutscher, B. *et al.* NMR methods for the study of instrinsically disordered proteins structure, dynamics, and interactions: General overview and practical guidelines. *Intrinsically Disord. Proteins Studied by NMR Spectrosc.* 49–122 (2015).

4. Jensen, M. R., Zweckstetter, M., Huang, J.-R. & Blackledge, M. Exploring free-energy landscapes of intrinsically disordered proteins at atomic resolution using NMR spectroscopy. *Chem. Rev.* **114**, 6632–6660 (2014).

5. Yagasaki, M. & Hashimoto, S. Synthesis and application of dipeptides; current status and perspectives. *Appl. Microbiol. Biotechnol.* **81**, 13–22 (2008).

6. Oh, K.-I. *et al.* A comprehensive library of blocked dipeptides reveals intrinsic backbone conformational propensities of unfolded proteins. *Proteins: Struct. Funct. Bioinforma.* **80**, 977–990 (2012).

7. Uversky, V. N. & Dunker, A. K. Understanding protein non-folding. *Biochimica et Biophys. Acta (BBA)-Proteins Proteomics* **1804**, 1231–1264 (2010).

8. Van Der Lee, R. *et al.* Classification of intrinsically disordered regions and proteins. *Chem. Rev.* **114**, 6589–6631 (2014).

9. Dunker, A. K. *et al.* Intrinsically disordered protein. *J. Mol. Graph. Model.* **19**, 26–59 (2001).

10. Babu, M. M. The contribution of intrinsically disordered regions to protein function, cellular complexity, and human disease. *Biochem. Soc. Transactions* **44**, 1185–1200 (2016).

11. Oldfield, C. J. *et al.* Comparing and combining predictors of mostly disordered proteins. *Biochemistry* **44**, 1989–2000 (2005).

12. Necci, M., Piovesan, D. & Tosatto, S. C. Critical assessment of protein intrinsic disorder prediction. *Nat. Methods* **18**, 472–481 (2021).

13. Trivedi, R. & Nagarajaram, H. A. Intrinsically disordered proteins: An overview. *Int. J. Mol. Sci.* **23**, 14050 (2022).

14. Evans, R., Ramisetty, S., Kulkarni, P. & Weninger, K. Illuminating intrinsically disordered proteins with integrative structural biology. *Biomolecules* **13**, 124 (2023).

15. Maiti, S., Maji, T., Saibo, N. V., De, S. *et al.* Experimental methods to study the structure and dynamics of intrinsically disordered regions in proteins. *Curr. Res. Struct. Biol.* 100138 (2024).

16. Borthakur, K., Sisk, T. R., Panei, F. P., Bonomi, M. & Robustelli, P. Determining accurate conformational ensembles of intrinsically disordered proteins at atomic resolution. *Nat. Commun.* **16**, 9036 (2025).

17. Kummerer, F., Orioli, S. & Lindorff-Larsen, K. Fitting force field parameters to NMR relaxation data. *J. Chem. Theory Comput.* **19**, 3741–3751 (2023).

18. Rindfleisch, T. *et al.* Molecular dynamics of the intrinsically disordered protein COR15A—A force field validation on structure and dynamics. *J. Chem. Theory Comput.* **21**, 9147–9163 (2025).

19. Ollila, O. S., Heikkinen, H. A. & Iwaï, H. Rotational dynamics of proteins from spin relaxation times and molecular dynamics simulations. *The J. Phys. Chem. B* **122**, 6559–6569 (2018).

20. Salvi, N., Abyzov, A. & Blackledge, M. Multi-timescale dynamics in intrinsically disordered proteins from NMR relaxation and molecular simulation. *The J. Phys. Chem. Lett.* **7**, 2483–2489 (2016).

21. Avbelj, F., Grdadolnik, S. G., Grdadolnik, J. & Baldwin, R. L. Intrinsic backbone preferences are fully present in blocked amino acids. *Proc. Natl. Acad. Sci.* **103**, 1272–1277 (2006).

22. Eker, F., Cao, X., Nafie, L. & Schweitzer-Stenner, R. Tripeptides adopt stable structures in water. A combined polarized visible raman, FTIR, and VCD spectroscopy study. *J. Am. Chem. Soc.* **124**, 14330–14341 (2002).

23. Shi, Z., Woody, R. W. & Kallenbach, N. R. Is polyproline II a major backbone conformation in unfolded proteins? *Adv. Protein Chem.* **62**, 163–240 (2002).

24. Aspromonte, M. C. *et al.* DisProt in 2024: Improving function annotation of intrinsically disordered proteins. *Nucleic Acids Res.* **52**, D434–D441 (2024).

25. Theillet, F.-X. *et al.* The alphabet of intrinsic disorder: I. Act like a pro: On the abundance and roles of proline residues in intrinsically disordered proteins. *Intrinsically Disord. Proteins* **1**, e24360 (2013).

26. Mateos, B. *et al.* The ambivalent role of proline residues in an intrinsically disordered protein: From disorder promoters to compaction facilitators. *J. Mol. Biol.* **432**, 3093–3111 (2020).

27. Pupier, M. *et al.* NMReDATA, a standard to report the NMR assignment and parameters of organic compounds. *Magn. Reson. Chem.* **56**, 703–715 (2018).

28. Ulrich, E. L. *et al.* NMR-STAR: Comprehensive ontology for representing, archiving and exchanging data from nuclear magnetic resonance spectroscopic experiments. *J. Biomol. NMR* **73**, 5–9 (2019).

29. Hoch, J. C. *et al.* Biological magnetic resonance data bank. *Nucleic Acids Res.* **51**, D368–D376 (2023).

30. Mölder, F. *et al.* Sustainable data analysis with Snakemake. *F1000Research* **10**, 33 (2025).

31. Dalby, A. *et al.* Description of several chemical structure file formats used by computer programs developed at molecular design limited. *J. Chem. Inf. Comput. Sci.* **32**, 244–255 (1992).

32. Bernstein, F. C. *et al.* The protein data bank: A computer-based archival file for macromolecular structures. *J. Mol. Biol.* **112**, 535–542 (1977).

33. Uversky, V. N. *et al.* Unfoldomics of human diseases: Linking protein intrinsic disorder with diseases. *BMC Genomics* **10**, S7 (2009).

34. Ziarek, J. J., Baptista, D. & Wagner, G. Recent developments in solution nuclear magnetic resonance (NMR)-based molecular biology. *J. Mol. Medicine* **96**, 1–8 (2018).

35. Wishart, D. S. Interpreting protein chemical shift data. *Prog. Nucl. Magn. Reson. Spectrosc.* **58**, 62–87 (2011).

36. Granger, P., Bourdonneau, M., Assémat, O. & Piotto, M. NMR chemical shift measurements revisited: High precision measurements. *Concepts Magn. Reson. Part A: An Educ. J.* **30**, 184–193 (2007).

37. Wang, Y. & Jardetzky, O. Investigation of the neighboring residue effects on protein chemical shifts. *J. Am. Chem. Soc.* **124**, 14075–14084 (2002).

38. Eggenberger, U., Pfäbdker, P. & Bodenhausen, G. Folding and pattern recognition in two-dimensional NMR spectra. *J. Magn. Reson.* **77**, 192–196 (1988).

39. Helmus, J. J. & Jaroniec, C. P. Nmrglue: An open source python package for the analysis of multidimensional NMR data. *J. Biomol. NMR* **55**, 355–367 (2013).

Acknowledgements

We thank Prof. Dr. Reinhard Lipowsky and Dr. Hanne Antila (Theory and Biosystems, Max Planck Institute of Colloids and Interfaces) for invaluable support in purchasing the dipeptide samples.

Funding

M.S.M. acknowledges support by the Trond Mohn Foundation (BFS2017TMT01). The NMR measurements were supported by the Research Council of Norway through the Norwegian NMR Platform, NNP (226244 and 322373), by the Bergen Research Foundation (BFS-NMR-1), and the Sparebankstiftinga Sogn og Fjordane (509-42/16).

Author contributions statement

J.U., T.R., and M.S.M. conceived the NMR experiments; T.R., E.F.T., and J.U. conducted the NMR experiments; T.R., E.F.T., and J.U. analyzed the NMR results; T.R. and J.U. performed technical validation and deposited the NMR data; M.S.M provided the samples; T.R. performed computational analysis for sample selection. J.U. and M.S.M. supervised the research; T.R. wrote and M.S.M edited the manuscript with inputs from all co-authors. All authors reviewed the final manuscript.

Competing interests

The authors declare no competing interests.