**A Conserved Landscape of Chemokine Receptor Co-expression Defines the Functional States of CD8+ T Cells in Melanoma**

**Authors:** Rodney Macedo[1,2]*, David. W. Harle[1,2], Kevin Hoffer-Hawlik[3], Ximi K. Wang[1,2], Thomas McMahon-Skates[1], Alexandra Matschiner[1], Kirubel Belay[1], Yvonne M. Saenger[4], Ben Izar[1,2,5], Elham Azizi[2,3,4], Ran Reshef[1,2,5]*

**Affiliations:**

[1]Columbia Center for Translational Immunology, Department of Medicine, Columbia University Irving Medical Center, New York, NY 10032, USA.

[2]Herbert Irving Comprehensive Cancer Center, Columbia University Irving Medical Center, New York, NY 10032, USA

[3]Department of Biomedical Engineering, Columbia University, New York, NY, USA. Irving Institute for Cancer Dynamics, Columbia University, New York, NY 10027, USA.

[4]Department of Oncology, Albert Einstein College of Medicine, Bronx, NY 10461, USA.

[5]Department of Medicine, Division of Hematology & Oncology, Columbia University Irving Medical Center, New York, NY, USA.

*To whom correspondence should be addressed: ran.reshef@columbia.edu, rmacedo@igtp.cat

**One Sentence Summary:** Co-expression of CCR5, CXCR6, and CXCR3 provides a functional 'code' that separates T-cell-mediated anti-tumor efficacy from off-target toxicity, enabling the selection of superior cells for safer and more effective cancer immunotherapies.

**Graphical Abstract**

**Highlights**

• CXCR6, CXCR3 and CCR5 co-expression signature stratifies patient survival in human
melanoma
• CD8[+] T cells co-expressing CXCR6, CXCR3 and CCR5 are critical effectors with high
proliferative, cytotoxic, and activation profile in human and mice melanoma
• CD8[+] T cells co-expressing CXCR6, CXCR3 and CCR5 drive anti-tumoral responses during
checkpoint blockade in both human and mice

**Abstract**:

Cancer immunotherapies, from checkpoint blockade to adoptive cell therapies like tumor-
infiltrating lymphocytes (TILs), have revolutionized cancer treatment but are limited by variable
efficacy and significant toxicities. A central challenge is identifying ideal T-cell populations that
effectively eliminate tumors without causing off-target damage, a distinction not captured by
existing biomarkers. We show that co-expression patterns of chemokine receptors (CRs) CXCR3,
CCR5, and CXCR6 on CD8[+] T cells provide a functional "code" defining subsets with divergent roles
in on-target immunity versus off-target inflammation. In mouse and human melanoma, a triple-
positive (CXCR3[+]CCR5[+]CXCR6[+]) T-cell subset is essential for tumor control, and its genetic signature
correlates with positive clinical response, while a distinct CCR5[+]CXCR6[+] subset drives liver immune-
related adverse events (IRAEs). Crucially, this CR code reveals that immunotherapy actively
reshapes T-cell trafficking patterns, uncovering profound heterogeneity within conventional
populations and distinguishing potent anti-tumor progenitors from cells predisposed to exhaustion
or off-target migration. This work establishes CR co-expression as a practical tool, providing a
surface marker-based strategy to identify and enrich optimized T cells for adoptive therapies,
thereby offering a framework to uncouple efficacy from toxicity.

**Article**:

Combined PD-1 and CTLA-4 blockade (P1C4) leads to impressive and durable clinical responses in melanoma [1], and many other cancers [2,3]. Though transformative, only a subset of patients responds, and immune-related adverse events (IRAEs) account for the high incidence of grade 3-4 adverse events, treatment-related mortality and high discontinuation rates, limiting broader application of combination therapies [4-11]. Both tumor rejection and IRAEs induced by checkpoint inhibitors are fundamentally governed by lymphocyte migration into tumors and healthy tissues [12-14]. The recent approval of autologous tumor-infiltrating lymphocytes (TILs) in metastatic melanoma and the fact that the majority of patients do not achieve long-term benefit from this approach [15-17] further underscore the need for deeper understanding of T-cell populations with an ideal profile that combines effective trafficking to the tumor, potent effector function, high proliferative capacity (stemness), ability to resist suppressive signals, and low exhaustion state, resulting in an effective anti-tumor response.

Chemokines are proteins that control the trafficking of T cells and other leukocytes into different tissues by interacting with chemokine receptors (CR) expressed on the cell surface [18]. Previous studies have investigated the role of CRs individually. Elevated expression of certain chemokine receptors such as CXCR3 and CCR5 or their ligands is associated with increased intratumoral T-cell infiltration and favorable outcomes in melanoma [19-21] and other cancers [22-27]. These studies led to the experimental use of CR inhibitors in cancer patients with mixed results [28-31], possibly due to failure to account for the biologic reality that leukocytes co-express multiple CRs which collectively orchestrate their migration and potentially other functions [32]. This highlights a significant gap in our understanding and an unmet need in identifying T-cell populations with an ideal profile - combining potent tumor trafficking and effector function while avoiding healthy tissues. This may facilitate the development of novel pharmacologic or adoptive cell therapy approaches in cancer.

Here, we address this challenge by investigating the co-expression patterns of key inflammatory CRs. We provide evidence in both mice and humans that the coordinated expression of CXCR3, CCR5, and CXCR6 defines distinct CD8$^+$ T-cell subsets with divergent roles in mediating tumor clearance and immune-related toxicity following dual checkpoint blockade. Our findings may help design future therapies that modulate leukocyte migration to improve the efficacy and reduce immune-related toxicity of cancer immunotherapies.

**A T-cell Chemokine Receptor Signature Stratifies Patient Survival in Melanoma**

To identify CRs central to the anti-tumor T-cell response and conserved across multiple cancers, we conducted an *in-silico* analysis of 11,323 tumor transcriptomes from 33 human cancers in the Cancer Genome Atlas (TCGA). This screen revealed that the inflammatory CR genes *CXCR3*, *CCR5*, and *CXCR6* show both high expression and variance (Extended Data 1a). This receptor triad formed a distinct signature that correlated strongly with T-cell specific genes in comparison with other leukocyte populations (Fig. 1a top and Supplementary Table 1). The same was true when analyzing 475 human samples from TCGA melanoma (Fig. 1a bottom and Extended Data 1b). Consistent with this, the primary ligands for CXCR3, CCR5 and CXCR6 correlated with T-cell- and macrophage-specific genes, but not with tumor cells, B cells or fibroblasts (Extended Data 1c).

While the expression level of each receptor individually was higher in "brisk" tumors [33]
(characterized by dense lymphocytic infiltrate) versus "non-brisk" PANCAN and melanoma
tumors, the combinatorial CR score proved superior. Specifically, CR scores (based on *CXCR6*,
*CXCR3* and *CCR5* expression) were not only associated with significant differences between these
tumor types (Fig. 1b, Extended Data 1d) but were also linked to improved overall survival
uniquely within brisk tumors (Fig. 1c). Similarly, CR scores were associated with higher abundance
of CD8+ T cells [34] and IFN-γ [35] in tumors, identifying a survival benefit even within CD8+ and IFN-γ
dominant tumors (Fig. 1b, Extended Data 1e-g). This was reinforced by an analysis of
transcriptional data from metastatic [36] and primary [37] human melanoma where CR scores were
higher in non-recurrent (Extended Data 1h) or responder (Extended Data 1i) tumors.
Interestingly, differentially expressed genes (DEGs) from samples with higher co-expression of
*CXCR6*, *CXCR3*, and *CCR5* are related to T cell activation pathways (Extended Data 1j and 1k).
Taken together, these data establish that *CXCR3*, *CCR5*, and *CXCR6* form a core inflammatory T-
cell gene signature, and their distinct co-expression correlates with better survival across tumor
types.

**Combinatorial Chemokine Receptor Expression Reveals a Hierarchy of T-cell Effector Programs**

To determine if different CR co-expression patterns define distinct functional states, we
performed *in-silico* single-cell analysis of treated and untreated human metastatic melanoma
tumors [38]. As shown in Extended Data 2a, after identifying eight major cell populations using cell
type-specific genes, we focused on CD8+ T cells (CD8+), classifying them into nine unique subsets
based on the combinatorial expression of *CXCR6*, *CXCR3* and *CCR5* (using *CCR7* to demarcate the
naive subset within the triple negative cells) (Fig. 1d left). This approach revealed extensive CR
heterogeneity in the TME (Fig. 1d right) and established that multi-receptor patterns, rather than
single markers, were required to delineate functionally distinct cell states.

Transcriptional profiling of CR-defined subsets uncovered a functional continuum across CD8+ T
cells, establishing a hierarchy from naive to terminally differentiated states (Fig. 1e and 1f,
Extended Data 2b and 2c, and Supplementary Table 2). At one end of the spectrum, the triple-
negative (3N+) and CXCR3-only (R3) subsets were the least differentiated, expressing markers of
stemness and a naive state. These cells also expressed chemokines that attract dendritic cells and
showed signs of early activation and regulation, suggesting a progenitor-like role with early
cytotoxic potential. In contrast, the co-expression of CCR5 and/or CXCR6 marked cells with potent
effector programs. Dual- and triple-expressing subsets were enriched for cytotoxic molecules,
cytokines, and chemokines. The **triple-positive (R6R3R5) subset** emerged as a critical effector
phenotype, showing the highest expression of genes related to proliferation, cytotoxicity, and
activation. Other dual-expressing subsets represented distinct intermediate states; for example,
R3R5 TILs were highly proliferative, while R6R3 TILs displayed a resident-effector profile with high
expression of cytotoxic molecules but lower proliferation. Finally, CXCR3-negative TILs appeared
more terminally differentiated, characterized by high expression of exhaustion markers. Within
this group, subsets displayed unique profiles: R5 cells resembled an immature memory state,
while R6R5 and R6 subsets were more akin to short-lived effector cells, with R6R5 being more
cytotoxic and R6 showing more tissue residency markers. These findings corroborate that specific

combinations of CXCR3, CCR5, and CXCR6 expressions define a detailed functional map of CD8$^+$
T cell diversity within the tumor microenvironment.

**Chemokine Receptor Expression Marks a Differentiation and Activation Trajectory in TILs**

To map the evolution of CR expression during T cell maturation, we analyzed a second human
metastatic melanoma single-cell RNA-seq dataset [39]. We used pseudotime inference, a
computational approach which orders cells along a continuous phenotypic trajectory, in this case
from naive (*TCF7*-expressing) to terminally differentiated (*TOX*-expressing) CD8$^+$ T cells [40]. To
better study the increase/decrease of CR expression and its variability across pseudotime, we
performed a rank sort (Extended Data 2d), and then we used a sliding window (300 cells) to
smooth the CR expression for the four genes across the inferred pseudotime (Extended Data 2e).
The analysis revealed a distinct and sequential pattern of CR expression along this differentiation
trajectory (Fig. 1g). The trajectory begins with naive cells expressing high levels of *CCR7*, which is
progressively lost as they mature (Fig. 1h). *CXCR3* expression emerges early and is sustained
throughout most of the cell's journey, whereas *CCR5* appears and increases to peak expression
in intermediate states. Finally, *CXCR6* expression is acquired late in the trajectory, marking the
most differentiated cells and correlating with *TOX* expression. Taken together, these results
suggest that as T-cells mature within the tumor, they undergo programmed, sequential co-
expression of *CXCR3*, *CCR5* and *CXCR6* and subsequently a gradual loss of these receptors at the
terminally differentiated phase. This evolution underscores that combinatorial dynamic
expression patterns, rather than static expression of a single receptor, are critical for accurately
characterizing the functional state of TIL subsets and their distinct roles in mediating anti-tumor
efficacy.

**A Conserved Chemokine Receptor Code Defines Functional T-cell Programs in Mice and**
**Humans**

To confirm if the CR-defined subsets identified in tumors represent a fundamental aspect of T-
cell biology, we analyzed their protein expression in healthy human and mouse lymphocytes
using multiparametric flow cytometry (Fig. 1a, Extended Data 3 and Supplementary Table 3). This
analysis revealed that the CR co-expression "code" is a conserved feature that defines a
functional hierarchy (Fig. 2b-c, Extended Data 4a-c). In both species, the least differentiated,
naive-like subsets (3N+ and R3) were the most abundant and largely corresponded to traditional
Naive (T$_N$) and Central Memory (T$_{CM}$) populations. In contrast, the Effector Memory (T$_{EM}$)
population was highly heterogeneous, comprising the full spectrum of CR-defined subsets (Fig.
2d-e, Extended Data 4d-f).

Crucially, our CR framework uncovered profound functional divergence even within the
classically defined TEM population (Fig. 2f and Extended Data 4g). For example, R3 T$_{EM}$ cells
retained stem-like properties (high TCF1/7), while subsets co-expressing multiple CRs, like
R6R3R5, were primed for cytotoxicity and proliferation. Other subsets showed exhausted
phenotypes (high PD-1, LAG3) or possessed distinct tissue-homing potentials, expressing
receptors that support either tumor infiltration (CCR2) or mucosal sites (CCR6, CCR9). Altogether,
these results confirm at the protein level that CR co-expression defines conserved, functionally
distinct T-cell subsets. This framework uncovers critical heterogeneity within the traditional T$_{EM}$

population, distinguishing cells with potent anti-tumor potential (R6R3R5) from those
predisposed to exhaustion or migration to other tissues.

**Distinct T-cell CR Subsets Drive On-Target Efficacy Versus Off-Target Inflammation During Checkpoint Blockade**

To dissect the *in vivo* roles of CR-defined subsets in the context of immunotherapy, we studied
their response to dual PD-1 and CTLA-4 blockade (P1C4) in a B16-F10 melanoma model (Fig. 3a).
This combination was chosen as it was most effective at increasing CD8$^+$ T$_{EM}$ cells compared to
single-agent therapy (Extended Data 5a) and is clinically relevant due to its high efficacy in human
melanoma. Following treatment, we observed increased CD8$^+$ T and T$_{EM}$ cell percentages in both
tumor and liver (Fig. 3b). While liver enzymes remained unchanged in this model,
immunohistochemistry revealed a significant increase in CD3$^+$ T cells in the liver, indicating
subclinical inflammation (Extended Data 5c-e). These results established a robust model to
simultaneously investigate the mechanisms of tumor control and the tissue-specific drivers of
immune-related toxicity.

P1C4 therapy profoundly and selectively altered the distribution of CR-defined subsets in
peripheral tissues (Figure 3c). In the tumor, treatment drove a dramatic expansion of the triple-
positive R6R3R5 subset, while causing a contraction of single-expressing R3 and R5 cells.
Conversely, in the liver, P1C4 selectively increased the R6R5 subset. These changes were not
observed in the spleen or tumor-draining lymph nodes (TdLNs), highlighting a tissue-specific
effect (Extended Data 5f). This divergence suggests that distinct CR programs mediate on-target
anti-tumor activity versus off-target immune infiltration, a hypothesis supported by a
corresponding increase in the ligands for CXCR3, CCR5 (e.g., CXCL9, CXCL10, CCL5) in both tumor
and liver supernatants after treatment, whereas CXCR6 ligand (CXCL16) is higher only in the
tumor (Extended Data 5g).

Genetic knockout models proved that this co-expression is essential for the therapeutic effect. In
the context of P1C4, mice lacking either *CCR5* or *CXCR6* lost the ability to increase their
intratumoral CD8$^+$ T$_{EM}$ cells, losing any significant change in CR-defined subsets, an effect not
seen in untreated knockout mice (Fig. 3d-e), demonstrating the role is specific to the
immunotherapy context. Furthermore, the P1C4-mediated increase in the tumor's favorable
CD8/Treg ratio was modestly reduced in *CCR5* knockout mice and completely eliminated in *CXCR6*
knockout mice (Fig. 3f). These data demonstrate that the full anti-tumor response to P1C4
requires the functional contribution of T cells co-expressing all three receptors.

A competitive short-term homing assay further clarified these roles (Fig. 3g). As expected, CXCR3
was confirmed to be critical for T-cell trafficking into the tumor, TdLNs, and liver specifically under
P1C4 treatment (Fig. 3h and Extended Data 5h-i). The roles of the other receptors were more
complex; for instance, CXCR6-deficient cells accumulated in lymphoid tissues after P1C4
treatment, suggesting impaired migration to the tumor or inability to establish tissue residency
status. Altogether, these experiments establish that while CXCR3 acts as a key director of T-cell
trafficking, the functional outcome of P1C4 therapy—driving either potent tumor rejection or

immune-mediated toxicity—is dictated by the specific co-expression patterns of CXCR3, CCR5,
and CXCR6 on responding T cells.

**Anti-Tumor T-cell Subsets Defined by CR Co-expression Expand with Checkpoint Blockade and**
**Correlate with Clinical Response**

To validate our findings in a clinical context, we first analyzed PBMC samples from melanoma
patients, comparing dual Immune Checkpoint Blockade (ICB)-treated and ICB-naive patients
(Supplementary Table 4). ICB-treated patients had significantly higher percentages of CD8+ T cells
in their blood compared to ICB-naive patients (Extended Data 6a-b). Strikingly, dual ICB treatment
led to a significant increase in the circulating triple positive R6R3R5 subset (Fig 4a), which our
preclinical data identified as having potent effector function. This expansion of triple positive
cells in human blood mirrors our findings in splenic CD8+ from mice treated with P1C4 (Extended
Data 6c). The overall profile of CR-defined subsets is modified by ICB (Fig. 4b and Supplementary
Fig. 4a), which also altered their immunophenotype (Fig 4c). Similar to what we found in normal
donors, 3N+ and R3 have higher naive/stemness markers such as TCF1/7, R6R3R5, R3R5, R6R5
and R5 have a greater cytotoxicity signature and increased expression of KLRG1, and those
exhausted are mainly R6R3, R6R5, R6 and R6R3R5 expressing high amounts of PD-1, LAG-3, and
TIM-3.

Interestingly, certain CR-defined subsets in ICB-treated patients have an altered profile compared
to ICB-naïve patients: 3N+ and R3 subsets have a significant decrease in AhR while R6 subsets
show a significant increase in KLRG1. Moreover, R5, R3R5, R6R3R5 and R6R5 subsets show a
significant decline of PD-1 expression, suggesting receptor occupancy by therapeutic antibody [41].
These results confirm in humans that P1C4 treatment increases the proportion of CR-defined
subsets responsible for tumor control but also modifies the functional properties of these
subsets.

To determine if CR-defined subsets within the human TME are associated with clinical outcomes,
we analyzed the single-cell transcriptional dataset from metastatic melanoma patients treated
with ICB [39]. This analysis revealed that *CXCR6* expression is significantly higher (Mann-Whitney
test; *P*=0.0095) in responder patients than non-responders, whereas *CCR7*, *CXCR3* and *CCR5* show
no association with response (Extended Data 6d). More importantly, the gene signatures for the
CXCR6-co-expressing subsets - R6R3R5, R6R3 and R6R5- are all significantly enriched in tumors
of responder patients (Mann-Whitney test; *P*<0.05) (Fig. 4d). These results confirm in humans
that ICB reshapes the landscape of circulating T cells to favor the expansion of the highly
functional R6R3R5 subset. Critically, they also establish that the enrichment of CXCR6-co-
expressing T-cell subsets within the tumor itself is a key feature of a successful anti-tumor
response to immunotherapy.

**Discussion**

A central challenge in cancer immunotherapy—from checkpoint blockade to adoptive cell
therapies like TILs—is uncoupling potent anti-tumor efficacy from treatment-limiting, off-target
toxicity. This requires a deeper understanding of the T-cell subsets that mediate these opposing
outcomes. While the roles of individual chemokine receptors in T-cell trafficking are well-studied,

the combinatorial "code" they form to dictate specific functional states has remained unclear.
Our study addresses this gap, demonstrating that the co-expression of CXCR3, CCR5, and CXCR6
provides a conserved functional map of CD8$^+$ T cells in both mice and humans, allowing for the
prospective identification of T-cell subsets responsible for on-target efficacy versus off-target
inflammation.

Our analysis reveals that this CR code defines a fundamental and conserved T-cell developmental
trajectory. In healthy individuals, we found that the hierarchy progresses from naive-like,
progenitor cells expressing primarily CXCR3 (R3 subset) to highly differentiated effector cells co-
expressing all three receptors (R6R3R5 subset). Crucially, this framework uncovers profound
functional heterogeneity within traditionally defined populations like effector memory T cells
(T$_{EM}$). Within the T$_{EM}$ pool, our code distinguishes stem-like progenitors (R3) from highly cytotoxic
and proliferative effectors (R6R3R5) and exhausted subsets, a level of functional resolution not
achievable with conventional markers. Importantly, our analysis demonstrates that specific
combinations of these three receptors, rather than single markers, are required to accurately
define the functional state and potential of a T cell.

In the context of cancer immunotherapy, this framework provides a powerful lens to dissect the
mechanisms of both tumor response and IRAEs. Our preclinical data show that dual ICB
selectively expands the triple-positive R6R3R5 subset in the tumor, a population essential for
therapeutic efficacy. Conversely, the therapy promoted the accumulation of a distinct
CCR5$^+$CXCR6$^+$ (R6R5) subset in the liver, a common site of IRAEs. This finding mechanistically links
a specific, identifiable T-cell subset to off-target toxicity, providing a cellular target for future
strategies aimed at mitigating IRAEs. The fact that genetic deletion of either CCR5 or CXCR6
abrogated the therapeutic effect of ICB underscores that the coordinated function of all three
receptors on the R6R3R5 subset is non-redundant and essential for a successful anti-tumor
response.

Ultimately, our findings set up this CR code not just as a prognostic biomarker but as a practical
tool for developing better immunotherapies. The prognostic value of single chemokine receptors
has been inconsistent across studies, likely because, as we show, the functional output of T cells
is associated with receptor combinations. The true translational potential of our work lies in its
direct application to cell-based therapies. The surface markers of this code are readily detectable
by high-dimensional flow cytometry, offering a straightforward strategy to identify and
prospectively isolate the most desirable T-cell populations. For adoptive cell therapies, such as
TILs, this framework provides a clear rationale for enriching the final product with the highly
proliferative and cytotoxic R6R3R5 subset while potentially depleting populations like R6R5 that
may contribute to toxicity. This approach offers a tangible path forward to engineer more potent
and safer immunotherapies, improving the therapeutic window by maximizing anti-tumor
activity while minimizing collateral damage.

While the expression profile of CRs and other trafficking molecules in immune cells has been
previously described in healthy humans [42,43] and in cancer [44,45], the functional impact of their co-
expression on the anti-tumor response requires multiparametric, single-cell analysis. Using

genetic datasets and high-dimensional flow cytometry, our study simultaneously analyzed multiple CRs with functional markers and transcription factors to clarify the critical role of CR co-expression in shaping anti-tumor T-cell responses.

**Data and code availability**

All data generated in this study are included in this published article and its supplementary information. Code relating to data processing and figure generation is available in our GitHub repository. Patient information is available on the supplementary table 4. Any additional information required to reanalyze the data reported in this paper is available from the corresponding authors upon request.

**Acknowledgements**

We thank the CCTI Biobank and Flow Cytometry Cores. We gratefully acknowledge the Columbia Institutional Review Board and all patients who generously consented to participate in this study, as well as the Institutional Animal Care and Use Committee. We are also grateful to Megan Sykes, Siu-Hong Ho, Audrey Baeyens, and Claudia Gutierrez-Chavez for helpful discussions.

**Funding**

E.A. is supported by NIH NCI R00CA230195, NHGRI R01HG012875, and grant number 2022-253560 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation. E.A. and R.R. are supported by a Columbia University HICCC programmatic pilot award. R.R. is supported by NIH R01-HL143424, P30-CA013696, and CA171008(DOD) grants. J.L.M. is supported by NIH NHGRI R35HG011941 and NSF CBET 2146007. The CCTI/HICCC Flow Cytometry Core is supported by NIH grants S10OD020056 and S10OD030282.

**Author contributions**

R.M., D.H., and R.R., conceived the study. R.M., D.H., K.H.H., X.W., A.M., and K.B. performed sample processing and data collection. R.M., D.H., K.H.H., X.W., T.M.S., K.B., Y.S., B.I., E.A., and R.R. provided clinical samples and data and maintained regulatory oversight. R.M., D.H., K.H.H., X.W., and T.M.S., analyzed the data. R.M., D.H., K.H.H., X.W., T.M.S., A.M., K.B., Y.S., B.I., E.A., and R.R. interpreted the data. R.M., R.R., E.A. and K.H.H. wrote the manuscript.

**Declaration of interests**

R.R. reports consulting or advisory role with Allogene, Bayer, Gilead Sciences, Incyte, TScan, Orca Bio, Pierre Fabre Pharmaceuticals, CareDx, Sana Biotechnology, Sail Biomedicines and Autolus, and research funding from Atara Biotherapeutics, Incyte, Sanofi, Immatics, Abbvie, Takeda, Gilead Sciences, CareDx, TScan, Cabaletta, Synthekine, BMS, J&J, Allogene, Genentech, Vittoria Therapeutics, AstraZeneca, Kinomica and Imugene.

**Figure 1.**

**Figure 1. The landscape of chemokine receptor (CR) gene expression in human cancer reveals clustered T-cell expression of CXCR3, CCR5 and CXCR6 which define functional subsets and have prognostic importance. a and b.** *In-silico* analysis of tumor transcriptome from Pan-Cancer Atlas samples (TCGA-PANCAN; n=11323) and melanoma patients (TCGA-SKCM; n=475). **a.** Hierarchical-Cluster heatmap reveals correlation of CR gene expression with different cell types in all tumors (top) or melanoma (bottom) tumor microenvironment. Cells are defined by sets of cell-type-specific genes (Supplementary Table 1). **b.** Heatmaps show the expression of *CXCR6*, *CXCR3* and *CCR5* in TCGA-PANCAN and melanoma samples based on their "brisk" classification [33]. Dot plot reveals higher CR score (based on CXCR6, CXCR3 and CCR5 expression) on brisk versus non-brisk samples from PANCAN or melanoma patients. **c.** Kaplan-Meier survival curves categorized by CR score in TCGA (PANCAN or SKCM) samples confirm their prognostic potential in only brisk, CD8[+] high [34] or IFN-γ tumors [35]. **d to f**. *In-silico* analysis of single-cell transcriptome from metastatic melanoma (GSE72056). **d.** Gating strategy used to define cell subsets based on the co-expression of *CXCR6*, *CXCR3*, *CCR5*, and *CCR7* genes. Bar plot showing the higher heterogeneity of CR subsets in T cells in comparison to other populations in the metastatic melanoma microenvironment. **e**. UMAP multidimensional analysis showing CR-defined subsets with different genetic profiles in CD8+ T-cells. **f**. Dot plot gene expression of markers of naïve, cytotoxicity, exhaustion, transcription factors and cytokines reveal transcriptional differences of CR-positive subsets in CD8+ T-cells. **g and h**. *In-silico* analysis of single-cell transcriptome from metastatic melanoma (EGAD00001009291). Diffusion mapping and pseudotime inference analysis reveal differential enrichment (**g**) and pseudotemporal expression patterns (**h**) of *CCR7*, *CXCR3*, *CCR5* and *CXCR6* during *TCF7* to *TOX* CD8[+] T cell trajectory. <u>CR-defined subsets</u>: 3N+ (*CXCR6*[-], *CXCR3*[-], *CCR5*[-], *CCR7*[+]), R3 (*CXCR6*[-], *CXCR3*[+], *CCR5*[-]), R3R5 (*CXCR6-*, *CXCR3*[+], *CCR5*[+]), R5 (*CXCR6*[-], *CXCR3*[-], *CCR5*[+]), R6R3R5 (*CXCR6*[+], *CXCR3*[+], *CCR5*[+]), R6R5 (*CXCR6*[+], *CXCR3*[-], *CCR5*[+]), R6R3 (*CXCR6*[+], *CXCR3*[+], *CCR5*[-]), R6 (*CXCR6*[+], *CXCR3*[-], *CCR5*[-]) and 3N- (*CXCR6*[-], *CXCR3*[-], *CCR5*[-], *CCR7*[-]).

**Extended Data Fig. 1.**

**Extended Data Fig. 1. a to c. Gene expression of chemokine receptors and their ligands in**
**human cancer by sample type.** *In-silico* analysis of tumor transcriptome from Pan-Cancer Atlas
samples (TCGA-PANCAN; n=11323) and melanoma patients (TCGA-SKCM; n=475). **a.** Mean-
Variance Plot graph showing high expression of *CXCR3*, *CCR5* and *CXCR6* genes in PANCAN, or
melanoma. Black dots represent genes for other chemokine receptors and chemokines. **b.**
Hierarchical-Cluster heatmap reveals correlation of chemokine receptor gene expression with
different tumor microenvironment cell-type-specific genes in metastatic and primary melanoma.
**c**. Hierarchical-Cluster heatmap reveals correlation of CXCR3, CCR5 and CXCR6 ligands gene
expression with different cell types in the tumor microenvironment in PANCAN and melanoma
samples. Cell types were defined by cell-type-specific genes. **d.** Dot plots show the expression of
*CXCR6, CXCR3, CCR5,* and CR scores (based on CXCR6, CXCR3 and CCR5 expression) in TCGA-
PANCAN and melanoma samples based on their "brisk" classification [33]. **e and f**. Dot plots show
the expression of CR scores (based on CXCR6, CXCR3 and CCR5 expression) in TCGA-PANCAN and
melanoma samples based on their "CD8$^+$ score" [34] (**e**) or immune subtype [35] (**f**) classification. **g**.
Dot plot represents the Hazard ratio (HR) and "p" value of CR scores and CR ratios (*CXCR3*/*CCR5*,
*CXCR6*/*CCR5*, *CXCR6*/*CXCR3*) for survival prediction in only brisk, CD8 high or IFN-γ tumor from
PANCAN or melanoma samples. **h and i**. Analysis of transcriptional data from human melanoma
reveals higher CR scores in non-recurrent vs recurrent metastatic tumors [36] (**h**) and responders
vs progressors primary tumors [37] (**i**). **j**. Identification of upregulated differentially expressed genes
(DEGs) from TCGA and Liu *et al* primary melanoma samples with higher expression of *CXCR6*,
*CXCR3*, and *CCR5* using SeqGeq. **j**. Dot plot graph showing the gene ontology analysis using shared
upregulated DEGs from TCGA and Liu *et al* samples.

**Extended Data Fig. 2.**

**Extended Data Fig. 2. a to c. Single-cell Gene expression of chemokine receptors at a single cell**
**level in human metastatic melanoma** (GSE72056). **a.** UMAP multidimensional analysis identifies
different immune populations based on immune gene expression in the tumor. **b.** UMAP
multidimensional analysis of gene expression of naïve, cytotoxicity, exhaustion markers,
transcription factors and chemokine receptors observed in CD8[+] T cells. **c**. Hierarchical-Clustering
heatmap of gene expression of markers of naïve, cytotoxicity, exhaustion, transcription factors
and cytokines reveal transcriptional differences of CR-positive subsets in CD8+ T-cells. **d and e.**
**Trajectory of chemokine receptors in Melanoma.** *In-silico* single-cell gene expression analysis of
human metastatic melanoma (EGAD00001009291). **d**. Rank sort and sliding window 300 cells of
*CCR7*, *CXCR3*, *CCR5* and *CXCR6* gene expression across CD8[+] T cell trajectory inferred by diffusion
pseudotime [46]. Pseudotime intervals are shown as colored bars. **e.** Distributions of normalized
chemokine receptor (CR) expressions among CD8[+] T cell samples.
388

**Figure 2.**

**Figure 2. Co-expression of CXCR3, CCR5 and CXCR6 proteins in human PBMCs and murine splenocytes reveals subsets with distinct differentiation profiles.** PBMCs from healthy human donors (n=11) and splenocytes from healthy C57BL/6J mice (n=5) were analyzed using flow cytometry. **a.** Hierarchical-Cluster heatmap analysis of T-cell subsets from human and mouse reveals co-expression of CXCR3, CCR5, CXCR6 and other chemokine receptors (CRs) in healthy conditions. **b.** Gating strategy used to identify CR subsets based on the surface co-expression of CXCR6, CXCR3, CCR5, and CCR7 (human) or CD62L (mouse). The percentages of T-cell subsets defined and color-coded by CR expression are represented in scatter dot plots in human and mouse. **c.** UMAP multidimensional analysis showing CR-defined subsets with different protein profiles in human CD8$^+$ T cells. **d.** Hierarchical-Cluster heatmap of normalized (N) protein

expression of CRs, transcription factors, activation and exhaustion markers reveal phenotypic
differences of CR-defined subsets in human and mouse CD8$^+$ T cells. **e**. Bar graphs showing the
distribution of CR-defined subsets within T-cell memory subsets reveal that T-effector memory
(T$_{EM}$) cells can be subdivided based on their co-expression of CRs in both human and mice. **f**.
Hierarchical-Cluster heatmap of normalized (N) protein expression of CRs, transcription factors,
activation and exhaustion markers also reveal phenotypic differences of CR-defined subsets in
human and murine CD8$^+$ T$_{EM}$ populations. The significance was calculated using unpaired Mann-
Whitney tests, *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. <u>CR-defined subsets</u>: 3N+
(CXCR6$^-$, CXCR3$^-$, CCR5$^-$, CCR7$^+$), R3 (CXCR6$^-$, CXCR3$^+$, CCR5$^-$), R3R5 (CXCR6$^-$, CXCR3+, CCR5$^+$), R5
(CXCR6$^-$, CXCR3$^-$, CCR5$^+$), R6R3R5 (CXCR6$^+$, CXCR3$^+$, CCR5$^+$), R6R5 (CXCR6$^+$, CXCR3$^-$, CCR5$^+$), R6R3
(CXCR6$^+$, CXCR3$^+$, CCR5$^-$), R6 (CXCR6$^+$, CXCR3$^-$, CCR5$^-$) and 3N- (CXCR6$^-$, CXCR3$^-$, CCR5$^-$, CCR7$^-$).

**Extended Data Fig. 3.**

**Extended Data Fig. 3. Gating strategy used for multiparametric full spectral flow cytometry.**
PBMCs from healthy human donors (n=11) and splenocytes from healthy C57BL/6J female mice
(n=5) were analyzed using spectral flow cytometry. For human cells, we used multiparametric
spectral flow cytometry with a 43-color extracellular panel and 42-color intracellular panel and
for murine cells we used a 36-color extracellular panel and a 35-color intracellular panel to study

419   multiple leukocyte populations, chemokine receptor (CRs), and other markers. **a.** Classic gating
420   strategy and UMAP multidimensional visualization used to identify human immune subsets. **b.**
421   Representative plots showing CR expression in human PBMCs. **c, d.** Classic and multidimensional
422   gating strategy used to identify murine immune subsets. **e.** Representative plots showing CR
423   expression in murine splenocytes.
424

**Extended Data Fig. 4.**

**Extended Data Fig. 4. Co-expression of CXCR3, CCR5 and CXCR6 proteins in human PBMCs and murine splenocytes reveals subsets with distinct function and differentiation profiles.** PBMCs from healthy human donors (n=11) and splenocytes from healthy C57BL/6J female mice (n=5) were analyzed using spectral flow cytometry. **a**. Heatmap analysis of immune subsets from humans and mice reveals the heterogeneity of expression and co-expression of chemokine receptors (CRs). **b**. UMAP multidimensional analysis showing protein expression of naïve, memory, activation and exhaustion markers in human CD8+ T cells. **c**. UMAP multidimensional analysis showing CR-defined subsets with different naïve, memory, activation and exhaustion markers profiles in CD8+ T-cells. **d to f**. CR-defined subsets were sorted from human PBMCs and murine splenocytes and stimulated for 2h to study activation and 2d for proliferation. **d**. Gating strategy to define CR subsets, green color refers to R6R3R5, R6R5, R6R3, and R6 subsets. Plots show that CR-defined subsets in CD8+ T cells have different activation (**e**) and proliferation (**f**) profiles. **g**. Scatter dot plot of CD8+ T_EM cells showing the expression of proteins that are differentially expressed between a specific CR subset and the rest of subsets (white dots). Blacked out cells in heatmap represent blank values. The significance was calculated using unpaired

ANOVA or Mann-Whitney multiple comparisons tests, *P < 0.05, **P < 0.01, ***P < 0.001, ****P

< 0.0001. <u>CR-defined subsets</u>: 3N+ (CXCR6$^-$, CXCR3$^-$, CCR5$^-$, CCR7$^+$), R3 (CXCR6$^-$, CXCR3$^+$, CCR5$^-$),

R3R5 (CXCR6$^-$, CXCR3+, CCR5$^+$), R5 (CXCR6$^-$, CXCR3$^-$, CCR5$^+$), R6R3R5 (CXCR6$^+$, CXCR3$^+$, CCR5$^+$),

R6R5 (CXCR6$^+$, CXCR3$^-$, CCR5$^+$), R6R3 (CXCR6$^+$, CXCR3$^+$, CCR5$^-$), R6 (CXCR6$^+$, CXCR3$^-$, CCR5$^-$) and

3N- (CXCR6$^-$, CXCR3$^-$, CCR5$^-$, CCR7$^-$).

**Figure 3.**

448
**Figure 3. Triple expression of CXCR3, CCR5 and CXCR6 is required for PD-1/CTLA4-induced**
**effector T-cell increase in melanoma tumors. a**. B16 tumor-bearing C57BL/6J female mice (n=15)
were treated with PD-1/CTLA-4 mAb on days 5, 8 and 11 (P1C4). Cells from tumors and liver were
isolated on day 14 and analyzed using spectral flow cytometry. **b.** P1C4 treatment increases the
numbers of tumor-infiltrating CD45$^+$ cells with no effect on liver, whereas levels of CD8$^+$ T or T$_{EM}$
cells in tumor and liver are increased after treatment. **c.** Nightingale Rose's charts show changes

of chemokine receptor (CR)-defined subsets after P1C4 treatment in tumor and liver in comparison to the control group (* to *** representing significant differences between P1C4 and PBS). **d to f**. B16 tumor-bearing C57BL/6J female wild-type (WT), CCR5$^{-/-}$ (R5$^{-/-}$) or CXCR6$^{-/-}$ (R6$^{-/-}$) mice (n=10) were treated with P1C4 as described above. **d top**. P1C4 increase of CD8$^+$ T$_{EM}$ infiltration in the tumor was lost in either CCR5 or CXCR6 lacking mice. **d bottom**. P1C4-induced increase of CD8$^+$ T$_{EM}$ infiltration in the liver was eliminated in R6$^{-/-}$ but not in R5$^{-/-}$ mice. **e**. R6R3R5 and R6R5 subsets are increased in CD8$^+$ T$_{EM}$ cells in tumor and liver respectively, after P1C4 treatment. Mice lacking CCR5 or CXCR6 showed no significant change in their CR-defined subsets after P1C4 treatment. **f**. P1C4 treatment increased CD8/Treg ratio in WT tumors; this effect is reduced in R5$^{-/-}$ mice and lost in R6$^{-/-}$ mice. No changes are observed in the liver. **g.** Female mice (n=3) with C57BL/6J background (WT, R5$^{-/-}$, R6$^{-/-}$ and CXCR3$^{-/-}$ (R3$^{-/-}$)) were treated with P1C4 on days 5, 8 and 11. T cells from spleen were isolated on day 12, labelled with CellTrace (excluding GFP$^+$ R6$^{-/-}$), and injected at the same proportion into the tail of B16 tumor-bearing (Day 12) PepBoy female mice (n=6) treated with P1C4 with the same regimen and the same groups (PBS in PBS, P1C4 in P1C4). Mice were sacrificed after 48 hours and cells from tumors, liver, spleen and tumor-draining lymph nodes (TdLNs) were isolated and analyzed using spectral flow cytometry. **h**. CD45.2$^+$ CD3$^+$ (tissue: CD90.2$^-$) cells lacking CXCR3 showed reduced tissue/injection ratio in P1C4-treated TdLNs, tumor and liver, while cells lacking CXCR6 are increased in TdLNs from P1C4 mice. Significance was calculated using unpaired ANOVA or Mann-Whitney multiple comparisons tests, *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. <u>CR-defined subsets</u>: 3N+ (CXCR6$^-$, CXCR3$^-$, CCR5$^-$, CCR7$^+$), R3 (CXCR6$^-$, CXCR3$^+$, CCR5$^-$), R3R5 (CXCR6$^-$, CXCR3+, CCR5$^+$), R5 (CXCR6$^-$, CXCR3$^-$, CCR5$^+$), R6R3R5 (CXCR6$^+$, CXCR3$^+$, CCR5$^+$), R6R5 (CXCR6$^+$, CXCR3$^-$, CCR5$^+$), R6R3 (CXCR6$^+$, CXCR3$^+$, CCR5$^-$), R6 (CXCR6$^+$, CXCR3$^-$, CCR5$^-$) and 3N- (CXCR6$^-$, CXCR3$^-$, CCR5$^-$, CCR7$^-$).

 **Extended Data Fig. 5.**

**Extended Data Fig. 5. Dual PD-1/CTLA4 blockade increases T-effector memory (T$_{EM}$) cells in B16**
**tumors and in target tissues of Immune-Related Adverse Events. a and b.** B16 tumor-bearing
C57BL/6J female mice (n=5-15) were treated with PD-1 (P1), CTLA-4 (C4) or both (P1C4) mAb on
days 5, 8 and 11. Cells from the spleen, tumor-draining lymph nodes (TdLNs), tumor and liver,
were isolated on day 14 and analyzed using spectral flow cytometry. **a**. Dual P1C4 blockade
showed the same tumor growth control at day 14 (**a left**, Vt = tumor volume at time t and Vo =
volume at baseline) but higher tumor infiltration of CD8$^+$ T$_{EM}$ in comparison with monotherapy
(**a center and right**). **b**. P1C4 treatment has no effect on levels of CD8$^+$ T cells in the spleen or
TdLNs but increases levels of CD8$^+$ T$_{EM}$ in these tissues. **c**. Livers from non-tumor bearing mice
show increased CD8$^+$ T$_{EM}$ levels after P1C4 treatment. **d**. Serum from mice treated with P1C4
reveals lower levels of lactate dehydrogenase (LDH) in comparison with control group, but other
serum markers such as alanine transaminase (ALT), aspartate transaminase (AST), alkaline
phosphatase (AL), albumin (ALB) and total bilirubin (BILT) were similar. **e**. Immunochemistry
staining of CD3 in the liver reveals increased levels of T cells in the tissue after P1C4 treatment in
comparison with PBS. **f**. Nightingale Rose's charts show no changes of chemokine receptor (CR)-
defined subsets after P1C4 treatment in spleen and TdLNs in comparison to the control group. **g**.
Scatter dot plots reveal an increase in the concentrations of CCR5, CXCR3 and CXCR6 ligands in
tumor and liver supernatant after P1C4 in comparison with controls. **h and i**. Female mice (n=3)
with C57BL/6J background (WT, CCR5$^{-/-}$, CXCR6$^{-/-}$ and CXCR3$^{-/-}$) were treated with P1C4 on days
5, 8 and 11 as described in Fig. 3i. **h**. A similar proportion of CD8$^+$ T cells from WT and genetic
knockout (KO) mice was confirmed before injection, with a vascular increase (CD90.2$^+$) of KO cells
after 48 hours. **i**. KO CD8$^+$ T cells have different percentages and tissue/injection ratio in the
spleen from recipient mice when compared to WT cells. Significance was calculated using
unpaired ANOVA or Mann-Whitney multiple comparisons tests. CR-defined subsets: 3N+ (CXCR6$^-$
, CXCR3$^-$, CCR5$^-$, CCR7$^+$), R3 (CXCR6$^-$, CXCR3$^+$, CCR5$^-$), R3R5 (CXCR6$^-$, CXCR3+, CCR5$^+$), R5 (CXCR6$^-$,
CXCR3$^-$, CCR5$^+$), R6R3R5 (CXCR6$^+$, CXCR3$^+$, CCR5$^+$), R6R5 (CXCR6$^+$, CXCR3$^-$, CCR5$^+$), R6R3 (CXCR6$^+$,
CXCR3$^+$, CCR5$^-$), R6 (CXCR6$^+$, CXCR3$^-$, CCR5$^-$) and 3N- (CXCR6$^-$, CXCR3$^-$, CCR5$^-$, CCR7$^-$).
507

**Figure 4.**

**Figure 4. PD-1/CTLA4 blockade changes CXCR6, CCR5 and CXCR3 co-expression in peripheral T**
**cells from melanoma patients.** PBMCs samples from melanoma patients (n=15) following
Immune Checkpoint Blockade (ICB, inverted triangles: early draw related to ICB, triangles: draw
in the middle of ICB, star: at the end of ICB) or not treatment (No Tx, circles) were analyzed using
full-spectral flow cytometry. **a**. Scatter dot plots showing that CD8$^+$ R6R3R5, and R6R5 subsets
are increased in the peripheral blood after P1C4. **b**. UMAP multidimensional analysis and
connecting dot plots reveal how ICB treatment changes CR subset profile of CD8$^+$ T cells. **c**.
Connecting dot plots showing molecular expression of chemokine receptors (CRs), transcription
factors, activation and exhaustion markers reveal phenotypic differences of CR-defined subsets
after treatment with ICB in CD8$^+$ T cells.  **d**. Box plots showing gene expression (MWU Test) from
EGAD000019291 dataset reveal higher expression of R6R3R5, R6R3, and R6R5 signatures in ICB
responder patients in comparison to non-responders. The significance was calculated using
unpaired ANOVA or Mann-Whitney multiple comparisons tests, *P < 0.05, **P < 0.01, ***P <
0.001, ****P < 0.0001. <u>CR-defined subsets</u>: 3N+ (CXCR6$^-$, CXCR3$^-$, CCR5$^-$, CCR7$^+$), R3 (CXCR6$^-$,

CXCR3$^+$, CCR5$^-$), R3R5 (CXCR6$^-$, CXCR3$^+$, CCR5$^+$), R5 (CXCR6$^-$, CXCR3$^-$, CCR5$^+$), R6R3R5 (CXCR6$^+$,
CXCR3$^+$, CCR5$^+$), R6R5 (CXCR6$^+$, CXCR3$^-$, CCR5$^+$), R6R3 (CXCR6$^+$, CXCR3$^+$, CCR5$^-$), R6 (CXCR6$^+$,
CXCR3$^-$, CCR5$^-$) and 3N- (CXCR6$^-$, CXCR3$^-$, CCR5$^-$, CCR7$^-$).
527

     **Extended Data Fig. 6.**

**Extended Data Fig. 6. PD-1/CTLA4 blockade changes CXCR6, CCR5 and CXCR3 co-expression in peripheral T cells from melanoma patients. a and b.** PBMCs samples from melanoma patients (n=15) following Immune Checkpoint Blockade (ICB, inverted triangles: early draw related to ICB, triangles: draw in the middle of ICB, star: at the end of ICB) or not treatment (No Tx, circles) were analyzed using full-spectral flow cytometry. ICB treatment significantly increases % of CD8 T$^+$ cells as observed in the UMAP multidimensional analysis (**a**) and plot graphs (**b**). **c**. Scatter dot plots show that CD8$^+$ R6R3R5 subsets are increased in the spleen of mice treated with P1C4 as described in Fig. 3a. **d**. Box plots showing gene expression (MWU Test) from EGAD000019291 dataset reveals higher expression of *CXCR6* in ICB responder patients in comparison to non-responders. The significance was calculated using unpaired ANOVA or Mann-Whitney multiple comparisons tests. <u>CR-defined subsets</u>: 3N+ (CXCR6$^-$, CXCR3$^-$, CCR5$^-$, CCR7$^+$), R3 (CXCR6$^-$, CXCR3$^+$, CCR5$^-$), R3R5 (CXCR6$^-$, CXCR3$^+$, CCR5$^+$), R5 (CXCR6$^-$, CXCR3$^-$, CCR5$^+$), R6R3R5 (CXCR6$^+$, CXCR3$^+$, CCR5$^+$), R6R5 (CXCR6$^+$, CXCR3$^-$, CCR5$^+$), R6R3 (CXCR6$^+$, CXCR3$^+$, CCR5$^-$), R6 (CXCR6$^+$, CXCR3$^-$, CCR5$^-$) and 3N- (CXCR6$^-$, CXCR3$^-$, CCR5$^-$, CCR7$^-$).

## References

Wolchok, J. D. *et al.* Final, 10-Year Outcomes with Nivolumab plus Ipilimumab in Advanced Melanoma. *N Engl J Med* (2024). https://doi.org/10.1056/NEJMoa2407417

Morad, G., Helmink, B. A., Sharma, P. & Wargo, J. A. Hallmarks of response, resistance, and toxicity to immune checkpoint blockade. *Cell* **184**, 5309-5337 (2021). https://doi.org/10.1016/j.cell.2021.09.020

Doki, Y. *et al.* Nivolumab Combination Therapy in Advanced Esophageal Squamous-Cell Carcinoma. *N Engl J Med* **386**, 449-462 (2022). https://doi.org/10.1056/NEJMoa2111380

Wolchok, J. D. *et al.* Overall Survival with Combined Nivolumab and Ipilimumab in Advanced Melanoma. *N Engl J Med* **377**, 1345-1356 (2017). https://doi.org/10.1056/NEJMoa1709684

Sznol, M. *et al.* Pooled Analysis Safety Profile of Nivolumab and Ipilimumab Combination Therapy in Patients With Advanced Melanoma. *J Clin Oncol* **35**, 3815-3822 (2017). https://doi.org/10.1200/JCO.2016.72.1167

Motzer, R. J. *et al.* Nivolumab plus Ipilimumab versus Sunitinib in Advanced Renal-Cell Carcinoma. *N Engl J Med* (2018). https://doi.org/10.1056/NEJMoa1712126

Sharma, P. *et al.* Nivolumab Plus Ipilimumab for Metastatic Castration-Resistant Prostate Cancer: Preliminary Analysis of Patients in the CheckMate 650 Trial. *Cancer Cell* **38**, 489-499 e483 (2020). https://doi.org/10.1016/j.ccell.2020.08.007

Hellmann, M. D. *et al.* Nivolumab plus Ipilimumab in Lung Cancer with a High Tumor Mutational Burden. *N Engl J Med* **378**, 2093-2104 (2018). https://doi.org/10.1056/NEJMoa1801946

Overman, M. J. *et al.* Durable Clinical Benefit With Nivolumab Plus Ipilimumab in DNA Mismatch Repair-Deficient/Microsatellite Instability-High Metastatic Colorectal Cancer. *J Clin Oncol*, JCO2017769901 (2018). https://doi.org/10.1200/JCO.2017.76.9901

Gao, J. *et al.* Neoadjuvant PD-L1 plus CTLA-4 blockade in patients with cisplatin-ineligible operable high-risk urothelial carcinoma. *Nat Med* **26**, 1845-1851 (2020). https://doi.org/10.1038/s41591-020-1086-y

Scherpereel, A. *et al.* Nivolumab or nivolumab plus ipilimumab in patients with relapsed malignant pleural mesothelioma (IFCT-1501 MAPS2): a multicentre, open-label, randomised, non-comparative, phase 2 trial. *Lancet Oncol* **20**, 239-253 (2019). https://doi.org/10.1016/S1470-2045(18)30765-4

Johncilla, M. *et al.* Ipilimumab-associated Hepatitis: Clinicopathologic Characterization in a Series of 11 Cases. *Am J Surg Pathol* **39**, 1075-1084 (2015). https://doi.org/10.1097/PAS.0000000000000453

Chen, P. L. *et al.* Analysis of Immune Signatures in Longitudinal Tumor Samples Yields Insight into Biomarkers of Response and Mechanisms of Resistance to Immune Checkpoint Blockade. *Cancer Discov* **6**, 827-837 (2016). https://doi.org/10.1158/2159-8290.CD-15-1545

Daud, A. I. *et al.* Tumor immune profiling predicts response to anti-PD-1 therapy in human melanoma. *J Clin Invest* **126**, 3447-3452 (2016). https://doi.org/10.1172/JCI87324

Rosenberg, S. A. *et al.* Durable complete responses in heavily pretreated patients with metastatic melanoma using T-cell transfer immunotherapy. *Clin Cancer Res* **17**, 4550-4557 (2011). https://doi.org/10.1158/1078-0432.CCR-11-0116

Rohaan, M. W. *et al.* Tumor-Infiltrating Lymphocyte Therapy or Ipilimumab in Advanced Melanoma. *N Engl J Med* **387**, 2113-2125 (2022). https://doi.org/10.1056/NEJMoa2210233

Sarnaik, A. A. *et al.* Lifileucel, a Tumor-Infiltrating Lymphocyte Therapy, in Metastatic Melanoma. *J Clin Oncol* **39**, 2656-2666 (2021). https://doi.org/10.1200/JCO.21.00612

Schulz, O., Hammerschmidt, S. I., Moschovakis, G. L. & Forster, R. Chemokines and Chemokine Receptors in Lymphoid Tissue Dynamics. *Annu Rev Immunol* **34**, 203-242 (2016). https://doi.org/10.1146/annurev-immunol-041015-055649

Bedognetti, D. *et al.* CXCR3/CCR5 pathways in metastatic melanoma patients treated with adoptive therapy and interleukin-2. *Br J Cancer* **109**, 2412-2423 (2013). https://doi.org/10.1038/bjc.2013.557

Chheda, Z. S., Sharma, R. K., Jala, V. R., Luster, A. D. & Haribabu, B. Chemoattractant Receptors BLT1 and CXCR3 Regulate Antitumor Immunity by Facilitating CD8+ T Cell Migration into Tumors. *J Immunol* **197**, 2016-2026 (2016). https://doi.org/10.4049/jimmunol.1502376

Hoch, T. *et al.* Multiplexed imaging mass cytometry of the chemokine milieus in melanoma characterizes features of the response to immunotherapy. *Sci Immunol* **7**, eabk1692 (2022). https://doi.org/10.1126/sciimmunol.abk1692

| 596 | 22 | Bell, B., Flores-Lovon, K., Cueva-Chicana, L. A. & Macedo, R. Role of chemokine receptors in gastrointestinal |
| 597 | | mucosa. *Int Rev Cell Mol Biol* **388**, 20-52 (2024). https://doi.org/10.1016/bs.ircmb.2024.02.003 |
| 598 | 23 | Valdivia-Silva, J. & Chinney-Herrera, A. Chemokine receptors and their ligands in breast cancer: The key |
| 599 | | roles in progression and metastasis. *Int Rev Cell Mol Biol* **388**, 124-161 (2024). |
| 600 | | https://doi.org:10.1016/bs.ircmb.2024.07.002 |
| 601 | 24 | Tang, Y., Hu, Y., Niu, Y., Sun, L. & Guo, L. CCL5 as a Prognostic Marker for Survival and an Indicator for |
| 602 | | Immune Checkpoint Therapies in Small Cell Lung Cancer. *Front Med (Lausanne)* **9**, 834725 (2022). |
| 603 | | https://doi.org:10.3389/fmed.2022.834725 |
| 604 | 25 | Seitz, S. *et al.* CXCL9 inhibits tumour growth and drives anti-PD-L1 therapy in ovarian cancer. *Br J Cancer* |
| 605 | | **126**, 1470-1480 (2022). https://doi.org:10.1038/s41416-022-01763-0 |
| 606 | 26 | Han, Y. *et al.* CXCL10 and CCL5 as feasible biomarkers for immunotherapy of homologous recombination |
| 607 | | deficient ovarian cancer. *Am J Cancer Res* **13**, 1904-1922 (2023). |
| 608 | 27 | Reschke, R. *et al.* Immune cell and tumor cell-derived CXCL10 is indicative of immunotherapy response in |
| 609 | | metastatic melanoma. *J Immunother Cancer* **9** (2021). https://doi.org:10.1136/jitc-2021-003521 |
| 610 | 28 | Fei, L., Ren, X., Yu, H. & Zhan, Y. Targeting the CCL2/CCR2 Axis in Cancer Immunotherapy: One Stone, Three |
| 611 | | Birds? *Front Immunol* **12**, 771210 (2021). https://doi.org:10.3389/fimmu.2021.771210 |
| 612 | 29 | Velasco-Velazquez, M. *et al.* CCR5 antagonist blocks metastasis of basal breast cancer cells. *Cancer research* |
| 613 | | **72**, 3839-3850 (2012). https://doi.org:10.1158/0008-5472.CAN-11-3917 |
| 614 | 30 | Sicoli, D. *et al.* CCR5 receptor antagonists block metastasis to bone of v-Src oncogene-transformed |
| 615 | | metastatic prostate cancer cell lines. *Cancer research* **74**, 7103-7114 (2014). https://doi.org:10.1158/0008- |
| 616 | | 5472.CAN-14-0612 |
| 617 | 31 | Halama, N. *et al.* Tumoral Immune Cell Exploitation in Colorectal Cancer Metastases Can Be Targeted |
| 618 | | Effectively by Anti-CCR5 Therapy in Cancer Patients. *Cancer Cell* **29**, 587-601 (2016). |
| 619 | | https://doi.org:10.1016/j.ccell.2016.03.005 |
| 620 | 32 | Bachelerie, F. *et al.* International Union of Basic and Clinical Pharmacology. [corrected]. LXXXIX. Update on |
| 621 | | the extended family of chemokine receptors and introducing a new nomenclature for atypical chemokine |
| 622 | | receptors. *Pharmacol Rev* **66**, 1-79 (2014). https://doi.org:10.1124/pr.113.007724 |
| 623 | 33 | Saltz, J., Gupta, R., Hou, L., Kurc, T., Singh, P., Nguyen, V., Samaras, D., Shroyer, K. R., Zhao, T., Batiste, R., |
| 624 | | Van Arnam, J., The Cancer Genome Atlas Research Network, Shmulevich, I., Rao, A. U. K., Lazar, A. J., |
| 625 | | Sharma, A., & Thorsson, V. Tumor-Infiltrating Lymphocytes Maps from TCGA H&E Whole Slide Pathology |
| 626 | | Images [Data set]. (2018). https://doi.org:10.7937/K9/TCIA.2018.Y75F9W1 |
| 627 | 34 | Wolf, D. Immune Signature Scores [Data set, https://pancanatlas.xenahubs.net]. (2016). |
| 628 | 35 | Thorsson, V. *et al.* The Immune Landscape of Cancer. *Immunity* **51**, 411-412 (2019). |
| 629 | | https://doi.org:10.1016/j.immuni.2019.08.004 |
| 630 | 36 | Gartrell, R. D. *et al.* Validation of Melanoma Immune Profile (MIP), a Prognostic Immune Gene Prediction |
| 631 | | Score for Stage II-III Melanoma. *Clin Cancer Res* **25**, 2494-2502 (2019). https://doi.org:10.1158/1078- |
| 632 | | 0432.CCR-18-2847 |
| 633 | 37 | Liu, D. *et al.* Integrative molecular and clinical modeling of clinical outcomes to PD1 blockade in patients |
| 634 | | with metastatic melanoma. *Nat Med* **25**, 1916-1927 (2019). https://doi.org:10.1038/s41591-019-0654-5 |
| 635 | 38 | Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. |
| 636 | | *Science* **352**, 189-196 (2016). https://doi.org:10.1126/science.aad0501 |
| 637 | 39 | Pozniak, J. *et al.* A TCF4-dependent gene regulatory network confers resistance to immunotherapy in |
| 638 | | melanoma. *Cell* **187**, 166-183 e125 (2024). https://doi.org:10.1016/j.cell.2023.11.037 |
| 639 | 40 | Saelens, W., Cannoodt, R., Todorov, H. & Saeys, Y. A comparison of single-cell trajectory inference methods. |
| 640 | | *Nat Biotechnol* **37**, 547-554 (2019). https://doi.org:10.1038/s41587-019-0071-9 |
| 641 | 41 | Yanagihara, T., Tanaka, K. & Matsumoto, K. A measuring method for occupancy of immune checkpoint |
| 642 | | inhibitors in the cell surface. *Biochem Biophys Res Commun* **527**, 213-217 (2020). |
| 643 | | https://doi.org:10.1016/j.bbrc.2020.04.122 |
| 644 | 42 | Wong, M. T. *et al.* A High-Dimensional Atlas of Human T Cell Diversity Reveals Tissue-Specific Trafficking |
| 645 | | and Cytokine Signatures. *Immunity* **45**, 442-456 (2016). https://doi.org:10.1016/j.immuni.2016.07.007 |
| 646 | 43 | Berahovich, R. D., Lai, N. L., Wei, Z., Lanier, L. L. & Schall, T. J. Evidence for NK cell subsets based on |
| 647 | | chemokine receptor expression. *J Immunol* **177**, 7833-7840 (2006). |
| 648 | | https://doi.org:10.4049/jimmunol.177.11.7833 |

44 Franciszkiewicz, K., Boissonnas, A., Boutet, M., Combadiere, C. & Mami-Chouaib, F. Role of chemokines and
chemokine receptors in shaping the effector phase of the antitumor immune response. *Cancer Res* **72**,
6325-6332 (2012). https://doi.org:10.1158/0008-5472.CAN-12-2027
45 Ozga, A. J., Chow, M. T. & Luster, A. D. Chemokines and the immune response to cancer. *Immunity* **54**, 859-
874 (2021). https://doi.org:10.1016/j.immuni.2021.01.012
46 Haghverdi, L., Buettner, F. & Theis, F. J. Diffusion maps for high-dimensional single-cell analysis of
differentiation data. *Bioinformatics* **31**, 2989-2998 (2015). https://doi.org:10.1093/bioinformatics/btv325
47 Villablanca, E. J. & Mora, J. R. Competitive homing assays to study gut-tropic t cell migration. *J Vis Exp* (2011).
https://doi.org:10.3791/2619
48 Anderson, K. G. *et al.* Intravascular staining for discrimination of vascular and tissue leukocytes. *Nat Protoc*
**9**, 209-222 (2014). https://doi.org:10.1038/nprot.2014.005
49 Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic
macrophage. *Nat Immunol* **20**, 163-172 (2019). https://doi.org:10.1038/s41590-018-0276-y
50 Lopez, R., Regier, J., Cole, M. B., Jordan, M. I. & Yosef, N. Deep generative modeling for single-cell
transcriptomics. *Nat Methods* **15**, 1053-1058 (2018). https://doi.org:10.1038/s41592-018-0229-2
664

**Methods**

**Human samples**

All study participants signed a written informed consent before inclusion in the study, and the protocols were approved by the Columbia University institutional review board (IRB). Peripheral blood mononuclear cells (PBMCs) were obtained from melanoma patients by Ficoll gradients and cryopreserved. PBMCs from healthy donors were obtained by Ficoll gradients and freshly analyzed using spectral flow cytometry. Characteristics of patients and healthy donors are summarized in Supplementary Table 4.

**Cell lines**

B16-F10 cell line was a gift from Dr. Vonderheide. B16-F10 cells were cultured at 37°C in Dulbecco's Modified Eagle Media, supplemented with 10% FBS, 100 U ml$^{-1}$ penicillin and 100 µg ml$^{-1}$ streptomycin, 2 mM L-glutamine. All cell lines were determined to be free of Mycoplasma (Lonza) and common mouse pathogens (IDEXX).

***In vivo* mouse studies**

All mice were used according to approved protocols by Columbia University Institutional Animal Care and Use Committee (IACUC). Female mice, including C57BL/6J (B6) (Jax #000664), B6.129P2-*Ccr5$^{tm1Kuz}$*/J (Jax #005427), B6.129P2-*Cxcr6$^{tm1Litt}$*/J (Jax #005693), B6.129P2-*Cxcr3$^{tm1Dgen}$*/J (Jax #005796), and B6.SJL-*Ptprc$^a$Pepc$^b$*/BoyJ (Jax #002014) were obtained from The Jackson Laboratory at 7-12 weeks of age and were used for studies one week or more after their arrival in the animal barrier facility of the Columbia Center for Translational Immunology. For B16-F10 melanoma, $1 \times 10^5$ B16-F10 cells diluted in PBS were mixed with an equal volume of Matrigel (Corning) and subcutaneously injected in the right flank of B6 mice on day 0. Blocking antibodies were given on days 5, 8 and 11 after tumor implantation. Antibodies used for *in vivo* immune checkpoint blockade experiments were given intraperitoneally at a dose of 200 mg per mouse and included: CTLA4 (9H10) and PD-1 (RMP1-14) (BioXCell). Tumor diameters were measured using calipers. Volume was calculated using the formula: (length × width$^2$)/2. Mice were euthanized when tumor burden reached a volume of 1500 mm³. Normalized tumor response to treatment is the measured volume at time t (Vt) relative to volume at baseline (Vo).

**Preparation of murine single-cell suspensions for flow cytometry**

Murine cell suspensions were obtained from tumors or livers by a combination of GentleMACS (Miltenyi Biotec) dissociation and enzymatic dissociation using 1 mg/mL of collagenase IV (Sigma-Aldrich) and 0.2 mg/mL of DNase (Roche), followed by incubation with ACK Lysis buffer (Life Technologies) for 1 minute to remove red blood cells. For tumor-draining lymph nodes, mechanical and enzymatic dissociation was performed. After enzymatic incubation, samples were filtered through a 70-µm nylon cell strainer (Falcon). Splenocytes were obtained by dissociation using a 70-µm nylon cell strainer (Falcon) followed by ACK lysis buffer incubation for 1 minute. Cell suspensions were counted using Trypan Blue for cell viability.

**Flow cytometry and cell sorting**

For flow cytometric analysis, live/dead cell discrimination was performed using Zombie NIR Fixable Viability dye (Biolegend) or eFluor780 Fixable Viability Dye (eBiosciences). Fc receptors were blocked using 2.4G2 anti-mouse CD16/CD32 (BD Pharmingen) or Human TruStain FcX (Biolegend). Cell surface staining was done for 15 min at 36°C, followed by 15 min at 4°C. Intracellular staining was done using a fixation/permeabilization kit (eBioscience). For human cells, we used multiparametric spectral flow cytometry with a 43-color extracellular panel and 42-color intracellular panel that allowed us to study multiple leukocyte populations, CRs, and functional markers simultaneously. In a similar way, for murine cells we used a 36-color extracellular panel and a 35-color intracellular panel to study multiple leukocyte populations, CRs, and other markers. For certain murine experiments traditional flow cytometry was used to identify T cell populations. All flow cytometric analyses were performed using Aurora 5 lasers (Cytek) or Fortessa II (BD) and analyzed using FlowJo software (BD). Cell sorting of CR-defined subsets was performed using an Influx Cell Sorter (BD). Human and murine sorted cells were stained with CellTrace kit followed by stimulation with Phorbol 12-myristate 13-acetate (PMA) for 2 hours to study activation or 2 days for proliferation. See Supplementary Table 3 for a list of antibodies used.

**Competitive homing experiment**

To determine the role of each individual CR in homing and tissue-infiltration of T cells, we performed a competitive short-term homing strategy [47] to identify four groups of cells injected at identical proportions. First, we treated WT, CCR5$^{-/-}$, CXCR6$^{-/-}$, or CXCR3 -/- mice with anti-PD-1 + anti-CTLA-4 (P1C4) or PBS on days 5, 8, and 11. On Day 12, mice were euthanized, and T cells were isolated from the spleen using Pan T Cell Isolation Kit II (Miltenyi Biotec).  After labelling T cells with different CellTrace Kits (except R6$^{-/-}$ mice which express GFP reporter) and combining them at similar proportion, $15 \times 10^6$ T cells were injected into the tail vein of B16 tumor-bearing PepBoy mice (Day 12 after tumor injection), also treated with PBS or P1C4 following the same treatment schedule and respecting the same group (PBS in PBS, P1C4 in P1C4). After 48 hours recipient mice were intravenously injected with anti-CD90.2 antibody for discrimination between vascular (circulating) and tissue lymphocytes [48]. After 3 minutes, mice were sacrificed, and the percentage of WT, CCR5$^{-/-}$, CXCR6$^{-/-}$, or CXCR3$^{-/-}$ in CD45.2$^+$ CD3$^+$ cells in each tissue was divided by the percentage of the same cells before injection.

**Immunohistochemistry of mouse tissues**

For liver immunohistochemistry, rabbit anti-CD3ε antibody (Invitrogen SP7) was diluted 1:100 and incubated at room temperature for 2 hours. Signal amplification was performed using ImmPRESS Horse anti-rabbit IgG antibody-HRP (Vector) and DAB substrate Kit (BD Pharmingen) following the manufacturer's protocol. Slides were counterstained and mounted.

**TCGA RNAseq**

RNAseq data was downloaded from the TCGA Pan-Cancer and TCGA Melanoma (SKCM) cohorts from xenabrowser.net [51] (IDs: EB++AdjustPANCAN_IlluminaHiSeq_RNASeqV2.geneExp.xena and TCGA.SKCM.sampleMap/HiSeqV2; unit: log2(norm_count + 1)), and corresponding phenotype datasets (IDs: Survival_SupplementalTable_S1_20171025_xena_sp, survival/SKCM_survival.txt,

TCGA_phenotype_denseDataOnlyDownload.tsv, TCGA.SKCM.sampleMap/SKCM_clinicalMatrix).
This dataset contained 11060 samples for PANCAN and 464 samples for SKCM, but only primary
and metastatic tumors were analyzed (n=10272 for PANCAN, n=104 for primary SKCM, and n=368
for metastatic SKCM). R value from Pearson correlation between normalized RSEM counts (> 5
log2(x+1)) of CRs/ligands and cell-type-specific genes (Supplementary Table 1; based on Tirosh
et al) was calculated using GraphPad Prism v10. Then, an unsupervised hierarchical clustering
gene tree was generated based on these correlations using Multi Experiment Viewer (MeV) v4.9
[52]. For survival analysis, optimal cutoffs were calculated using Kaplan-Meier Plotter (kmplot.com)
[53], and Kaplan-Meier curves and Log-rank p values were generated using GraphPad Prism v10.
Identification of upregulated differentially expressed genes from TCGA and Liu *et al* primary
melanoma samples was done using SeqGeq (BD), a platform that allows visualization of scRNA
seq data and RNA seq data. STRING database (https://string-db.org/), was used to generate
biological process gene ontology analysis [54].

**Analysis of publicly available single-cell datasets**

To perform an *in-silico* analysis of single-cell RNA seq data from human metastatic melanoma,
the GSE72056_melanoma_single_cell_revised_v2.txt.gz file was downloaded from GEO Series
GSE72056. Using SeqGeq (BD), a characterization of immune and non-immune cells in the TME
was performed with UMAP multidimensional analysis. Subsets were defined in CD8$^+$ T cells by
the co-expression of *CCR7*, *CXCR6*, *CXCR3* and *CCR5* following the gating strategy defined in Fig.
1d left. Finally, gene expression of other functional markers was measured within CR-defined
subsets.

**scRNA-seq analysis of external clinical melanoma cohort immune cells**

Pre-processed scRNA-seq datasets of immune cells (n=49,009) from patients studied in Pozniak
et. al [39] were loaded into Seurat in R and subsequently Scanpy in Python. Immune cell annotation
was guided by automatic annotation using SingleR and its built-in reference dataset of stroma
and immune cells 'BlueprintEncodeData' as previously described [49]. Non-cycling CD8$^+$ T cell
subtypes (n=13,304) were selected for downstream analysis in Scanpy. The top 8,000 highly
variable genes were calculated using the 'Seurat_V3' method implemented in Scanpy. Gene
expression counts were normalized to median total counts and logarithmized in Scanpy. CD8$^+$ T
cells from individual samples were batch corrected using an scVI model trained with two hidden
layers and 30-dimensional latent space [50]. After performing neighborhood graph construction
using 20 nearest neighbors in scVI latent space, leiden clustering with default resolution, and
evaluation of known marker genes and CD8$^+$ T cell gene signatures in Scanpy, a noisy outlier
cluster was removed to prevent distortions in subsequent diffusion maps. Then, after
reconstructing the neighborhood graph using 300 nearest neighbors in scVI space, we computed
diffusion components (DCs) using the Scanpy implementation to examine continuous phenotypic
dynamics. The first and second DCs reflected expected *TCF7*$^+$ to *TOX*$^+$ progression of CD8$^+$ T cells,
and diffusion pseudotime along this trajectory was computed using the Scanpy implementation
to reflect this progression. We then plotted the expression of *CCR7*, *CXCR3*, *CCR5*, and *CXCR6*
across pseudotime using CD8$^+$ T cells ordered by diffusion pseudotime rank. Given the sparsity of
CR expression, which is typical for a clinical scRNA-seq dataset, we used sliding windows of 300
cells to average normalized expression across the CD8$^+$ T cell trajectory. We also computed the

95% confidence interval on the same window of 300 cells to reflect variation in expression across the trajectory. The window size of 300 reduced sparsity while still conserving the expected negative binomial distribution of CR gene expression counts. To compare CR expression patterns over CD8$^+$ T cell trajectory, expression of each CR was scaled and plotted across cell pseudotime rank, reflecting the expected progression of *CCR7*$^+$ to *CXCR6*$^+$ expression in CD8$^+$ T cells. Finally, we examined differential expression of CR genes and the R6 co-expressing signatures (R6R3, R6R5, R6R3R5) across ICB responder/non-responder clinical groups using average gene expression pseudobulked by sample (summation of counts in each sample divided by total number of cells in each sample) to account for CR expression sparsity and intrapatient heterogeneity. Pseudobulked sample gene expression and signature expression were compared across clinical groups using boxplots and significance was assessed using Mann-Whitney U-tests. Code relating to this data analysis is publicly available on GitHub: https://github.com/azizilab/chemokine_receptor_reproducibility.

**Statistical analysis**

One-way ANOVA was used for normally distributed data. Mann–Whitney was used for non-parametric data. Kaplan–Meier log-rank analysis was used to evaluate the survival differences between groups. Statistical analysis was conducted using Prism 10.0 software (GraphPad). Only p values < 0.05 were considered as significant.

**Methods Bibliography.**

51. Goldman, M.J., Craft, B., Hastie, M. et al. Visualizing and interpreting cancer genomics data via the Xena platform. Nat Biotechnol (2020). https://doi.org/10.1038/s41587-020-0546-8

52. Saeed AI, Sharov V, White J, Li J, Liang W, Bhagabati N, Braisted J, Klapa M, Currier T, Thiagarajan M, Sturn A, Snuffin M, Rezantsev A, Popov D, Ryltsov A, Kostukovich E, Borisovsky I, Liu Z, Vinsavich A, Trush V, Quackenbush J. TM4: a free, open-source system for microarray data management and analysis. Biotechniques. 2003 Feb;34(2):374-8

53. Lanczky A, Gyorffy B. Web-Based Survival Analysis Tool Tailored for Medical Research (KMplot): Development and Implementation, J Med Internet Res, 2021; 23(7):e27633, doi:10.2196/27633

54. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, Gable AL, Fang T, Doncheva NT, Pyysalo S, Bork P, Jensen LJ, von Mering C. The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Res. 2023 Jan 6;51(D1):D638-D646. doi: 10.1093/nar/gkac1000.