Cancer pathway connectivity resolved by drug perturbation and RNA sequencing

Thi Huong Lan Do[1]* Caroline Lohoff[2,3,4]*, Ferris Jung[5], Sandra Kummer[1], Marcel Fabian Pohly[1,6,7], Sebastian Scheinost[8], Jennifer Huellein[3,8], Vladimir Benes[5], Wolfgang Huber[3,4,7✉], Junyan Lu[2,3,4✉], Thorsten Zenz[1,7✉]

[1] Department of Medical Oncology and Hematology, University Hospital Zurich and University of Zurich (UZH), Zurich, Switzerland
[2] Institute for Computational Biomedicine, Medical Faculty Heidelberg, Heidelberg University, Heidelberg, Germany
[3] Genome Biology Unit, European Molecular Biology Laboratory (EMBL), Heidelberg, Germany
[4] Molecular Medicine Partnership Unit (MMPU), Heidelberg, Germany
[5] Genomic Core Facility, European Molecular Biology Laboratory (EMBL), Heidelberg, Germany
[6] Department of Biosystems Science and Engineering, ETH Zurich, Basel, Switzerland
[7] INTeRCePT consortium, The LOOP Zurich Medical Research Center, Zurich, Switzerland
[8] Department of Translational Medical Oncology, National Center for Tumor Diseases (NCT) Heidelberg and German Cancer Research Center (DKFZ), Heidelberg, Germany

* These authors contributed equally.
✉ Corresponding authors

Correspondence:

Thorsten Zenz, Department of Medical Oncology and Hematology, University Hospital and University of Zurich, Rämistrasse 100, CH-8091 Zurich, Switzerland. ✉ Thorsten.Zenz@usz.ch

Junyan Lu, Institute for Computational Biomedicine, Medical Faculty Heidelberg, Heidelberg University, Im Neuenheimer Feld 130, D-69120 Heidelberg, Germany. ✉ Junyan.Lu@uni-heidelberg.de

Wolfgang Huber, European Molecular Biology Laboratory, Meyerhofstraße 1, D-69117 Heidelberg, Germany. ✉ Wolfgang.Huber@embl.org

Abstract

Pathway inhibitors are a backbone of cancer treatment. The configuration of pathway dependencies varies from tumour to tumour. Better treatment for individual patients could be designed if the pathway wiring were readily measurable.

Here, we characterise the transcriptional responses of 116 lymphoma patient samples exposed to ten drug perturbations. We used factor analysis to decompose individual and shared drug effects, thereby generating a pathway connectivity map of chronic lymphocytic leukemia (CLL). The expression profiles of the major disease subgroups, defined by IGHV mutation status, became more similar to each other after BTK inhibition, consistent with B-cell receptor (BCR) signalling as a driver of their phenotypic difference. An even stronger convergence was observed with combined IRAK4 and BTK inhibition, indicating cooperation of BCR and toll-like receptor (TLR) signalling in CLL. We identified genetic aberrations (*BRAF*, *TP53*, deletion 17p, deletion 15q, trisomy 12) that modulated drug effects in CLL and constituted specific interaction patterns. IRAK4 inhibition effects depended on the presence of trisomy 12, a finding that suggests that the trisomy 12 driver event in CLL acts by gene dosage-dependent IRAK4 upregulation and amplification of the BCR/TLR cooperation.

Our results highlight the potential of systematic drug perturbation assays with transcriptome readout to map pathway interconnectivity and functionally annotate tumour drivers.

Introduction

Chronic lymphocytic leukemia (CLL) is a common B-cell lymphoma in adults. More than 200 driver genes and chromosomal aberrations are known, which target a broad range of signalling pathways including the B-cell receptor (BCR), DNA damage response and NOTCH pathway[1–4]. A more precise understanding how this broad range of pathways cooperate and contribute to the CLL phenotype should facilitate better treatments. Multi-omics profiling of CLL patient samples has revealed major biological dimensions of CLL, such as the epigenetic cell-of-origin fingerprint[5,6], the clinical and biological impact of IGHV mutations[7,8], autonomous BCR signalling[9] and CLL proliferative drive (CLL-PD)[8,10–13]. Nonetheless, many common mutations and copy number variants (CNVs), e. g. trisomy 12, have not yet been conclusively linked to a function[10].

The ability to measure multi-omics data in large cancer cohorts has refined molecular subgrouping, improved outcome prediction and treatments[12,14,15]. Recently, however, progress has often focused on the discovery of rarer disease variants with the inherent downside that findings only apply to a small proportion of patients[1,3]. Furthermore, clinical precision oncology has largely relied on cancer genome sequencing and the identification of genetic mutations as biomarkers[16,17], while other omics data have remained research tools. Among these, gene expression profiling is particularly suited for the detailed characterisation of drug effects. This concept has been applied in the "Connectivity Map", which represents a landmark study mapping over 80,000 genetic and chemical perturbations to gene expression signatures in 29 cancer cell lines[18,19], and has been widely used to identify drug effects, drug repurposing and construct gene regulatory networks[18,20–22]. By using the perturbation signatures, transcriptional consequences can specifically be linked to the targeted upstream cause and therefore facilitate the understanding of cellular mechanisms[18].

In this study, we applied a set of 10 small-molecule drugs to a collection of 116 primary tumours, to investigate how these drugs work, capture the wiring of targeted pathways and identify modulators (i.e. mutations) of drug response.

Results

Shallow depth transcriptional profiles capture CLL disease dimensions and drug effects

In order to increase feasibility and cost-effectiveness of profiling hundreds to thousands of samples, we first compared gene expression from five CLL samples profiled with whole-transcript deep (TruSeq) and 3'-end-based library preparation with shallow depth RNA sequencing (QuantSeq[23]) (Suppl. Fig. 1a-d). Although the sequencing depth was reduced up to 30-fold (2.2-3.8 vs. 43-73 million reads per sample), QuantSeq recovered 71% of the genes detected by TruSeq with reproducible quantification (R = 0.82-0.85) (Suppl. Fig. 1e-g).

We selected a cohort of 108 chronic lymphocytic leukemia (CLL) and eight other B-cell Non-Hodgkin (B-NHL) lymphoma patient samples with available molecular profiles and clinical data (Fig. 1) and used QuantSeq to generate 1,086 (1,010 for CLL, 76 for B-NHL; 92% successfully prepared libraries) gene expression profiles after 48h of *ex-vivo* treatment with inhibitors used clinically (BTK inhibitor ibrutinib, PI3K inhibitor duvelisib, MEK inhibitor trametinib, mTOR inhibitor everolimus, XPO1 inhibitor selinexor), compounds targeting key signalling pathways (IRAK4 inhibitor compound 26, AKT inhibitor MK2206, MDM2 inhibitor and p53 activator nutlin-3a, BET inhibitor IBET762), and one combinatorial drug treatment (BTK and IRAK4 inhibition) (Fig. 1a, Suppl. Fig. 2a,b, Suppl. Table 1). IGHV mutational status, trisomy 12 and *TP53* mutation influenced viability of cells after drug treatment consistent with our prior findings[11] (Suppl. Fig. 2c-e).

We began our analysis by performing principal component analysis (PCA) to investigate which gene expression programmes, as characterised by factors ($\triangleq$ principal components), were the main contributors of transcriptional variation in our data set. PCA of the DMSO-treated control samples showed that transcriptional variation between patients was predominantly driven by disease type, IGHV, methylation group[5,6] and trisomy 12 status (Suppl. Fig. 2f). This result was highly similar to results from studies using conventional RNA sequencing[13] and shows that we were able to capture the main biological disease dimensions after *ex-vivo* culture and markedly reduced sequencing depth (Suppl. Fig. 2g,h). PCA of all samples including drug treatments

revealed that the first component (PC1) captured a contribution of sample viability to the

transcriptional variation independent of perturbation (Fig. 2a,b, Suppl. Fig. 2i, Suppl. Fig. 3a-

c). This viability signature was enriched in pathways linked to a general upregulation of

hypoxia, p53 pathway, TNFα signalling via NFκB, and downregulation of MYC targets and

OXPHOS preferentially in drug-perturbed samples (Suppl. Fig. 3d). Variation driven by IGHV

status/methylation programme and trisomy 12 were reflected by PC2 and PC3 (Fig. 2c, Suppl.

Fig. 2i,j). We also used *t*-Distributed Stochastic Neighbour Embedding (*t*-SNE) to visualise

samples' gene expression similarities and found patterns that were dominated by inter-patient

differences and inherent molecular disease features (Fig. 2d-e). After subtracting the patient-

specific baseline signatures, the embeddings captured drug specific-effects (Fig. 2f, Suppl.

Fig. 2k).

Individual drug effects

To understand the effect of individual drugs, we extracted drug-specific transcriptional

signatures by performing differential gene expression analysis between drug- and control-

treated samples (Fig. 2g). We first assessed the response to nutlin-3a, a MDM2 inhibitor and

activator of the transcription factor p53[24]. We identified the p53 activation signature in primary

CLL cells, which consisted of a strong upregulation of well-known transcriptional p53 targets

and pathways including *BAX*, *MDM2* and *CDKN1A*/p21 (Fig. 2h-i). This response was blunted

in cases with *TP53* mutation and/or deletion in chromosome 17p (Fig. 2j).

Inhibition of PI3K and BTK elicited the strongest responses, as assessed by the number of

differentially expressed genes, in line with the role of BCR signalling as a key driver of CLL[7–9]

(Fig. 2g). BTK inhibition led to the downregulation of genes associated with B-cell activation

and function (*FCRL5*, *FILIP1L*, *MS4A1*/CD20, *CCL3*/*4*) and phosphatases (*PTPN6*, *DUSP4/5*)

(Fig. 2k,l). We also found downregulation of key B-cell transcription factors (*PAX5*, *IRF4*,

IKZF3) and members of the NFκB pathway (*REL*, *RELB, NFKB1),* suggesting a role of active

BCR signalling in the maintenance of the regulatory B-cell identity programme. This signature

overlapped with reported transcriptional signatures after BCR stimulation and *in-vivo* BTK

inhibitor treatment[25–27] (Suppl. Fig. 4a).

PI3K inhibition led to the downregulation of gene sets enriched in immune cell response,

cytokine signalling, interferon (IFN) response and BCR signalling, similar to the effects upon

BTK inhibition but more pronounced (Fig. 2l,m, Suppl. Fig. 4b,c). We investigated the

similarities and differences between BTK and PI3K inhibition by directly comparing the effects

of the two inhibitors against each other (Fig. 2m, Suppl. Fig. 4d). Genes only modulated by the

BTK inhibitor included components of B-cell development and function, endocytosis and

autophagy, and the IFN response pathway. PI3K inhibition exclusively downregulated genes

associated with IFN and cytokine signalling (Fig. 2n, Suppl. Fig. 4d). To understand whether

these differences were driven by a common upstream mediator, we inferred the transcription

factor activities from the differentially expressed genes[28] in which transcription factor activity of

STAT family members and in particular gene expression of *STAT1* was downregulated by PI3K

but not BTK inhibition (Fig. 2o, Suppl. Fig. 4e-f).

Inhibition of AKT modulated gene sets involved in antigen processing and presentation, IFN

signalling and suppression of MHC class II molecules similar to the PI3K inhibitor signature

(Suppl. Fig. 5a,b). Differentially expressed genes between PI3K and AKT inhibitor treatment

mostly consisted of BTK-dependent targets that were only suppressed upon PI3K inhibition

(Suppl. Fig. 5c). The mTOR inhibitor signature showed overlap with the AKT and PI3K inhibitor

signatures in the context of antigen processing and adaptive immune system (Suppl. Fig.

5b,d), and specifically included suppression of gene sets associated with lysosome,

cholesterol homeostasis and mTORC1 signalling (Suppl. Fig. 5b,g). The predominant

mechanisms of action (MoA) induced by the MEK inhibitor were cytokine signalling and IFN

response (Suppl. Fig. 5b,e-g).

Bromodomain and extra-terminal motif (BET) proteins are readers of acetylated histone marks

and regulate gene expression[29]. Inhibition of BET proteins led to upregulation of genes

associated with translation, ribosome and metabolism of RNA (Suppl. Fig. 5h,i). Selinexor is a

small-molecule that prevents the translocation of proteins from the nucleus to the cytoplasm

by inhibiting the nuclear exporter XPO1[30]. *XPO1* was among the top upregulated genes

together with solute carriers, nucleotide- and protein-binding adapters, and members of the

zinc finger protein family (Suppl. Fig. 5j,k). Gene expression of p53 was also increased after

XPO1 inhibitor treatment without induction of the transcriptional p53 activation signature

(Suppl. Fig. 5l). Inhibition of toll-like receptor (TLR) signalling with the IRAK4 inhibitor did not

lead to significant differential gene expression (Fig. 2g).

Connectivity of signalling and perturbation effects in CLL

The use of multiple drugs allowed us to understand how important signalling axes in CLL are

connected. For this, we constructed a pathway connectivity network as a bipartite graph in

which treatment and gene nodes are connected by edges when the perturbation led to

differential expression of that gene. We found that the BTK and PI3K inhibitors were strongly

linked through their common BCR-inhibiting MoA. The AKT inhibitor formed close connections

to the PI3K and mTOR inhibitors along the PI3K/AKT/mTOR pathway (Fig. 3a). We also

illustrated these relationships by comparing the perturbations directly against each other (Fig.

3b). Because drug perturbations typically do not affect a single gene in isolation, but

simultaneously modulate the expression of multiple related genes, we used a factorisation

technique, namely guided sparse factor analysis[31] (GSFA) to decompose the full set of drug-

induced differentially expressed genes into individual latent factors of co-regulated genes (Fig.

3c-e, Suppl. Fig. 6). We identified drug-specific effects upon inhibition of MDM2 (factor 18),

XPO1 (factor 12), BET (factor 16) and MEK (factor 14), but also found modules affected by

multiple drugs at variable intensities (Fig. 3c,d, Suppl. Fig. 6a,b). These shared axes of

pathway heterogeneity distinguished canonical BCR signalling (factor 8; BTK/PI3K inhibition)

from antigen processing (factor 17; PI3K/AKT inhibition), and the lysosome pathway (factor

20; mTOR/MEK inhibition) (Fig. 3c,e, Suppl. Fig. 6b,c). We also identified modules regulated

in opposing directions, which could indicate compensatory pathway activation, and link them

to IFN (factor 14; MEK/XPO1 vs. mTOR/BTK inhibition) and lysosomal signalling (factor 20;

MEK/mTOR vs. PI3K/BTK inhibition) (Fig. 3c-e). Next, we used linear regression to identify the

impact of the most prevalent disease aberrations and CNVs on individual patient factor

contributions. Mutation or deletion of *TP53* was negatively associated, while trisomy 12 was

positively associated with the response to the MDM2 inhibitor captured in factor 18 (Suppl. Fig.

6d). BTK and MEK inhibitor responses reflected in the factors 17 and 14, respectively, were

positively associated with unmutated IGHV and trisomy 12 (Suppl. Fig. 6e-f).

Pathway dependency of known CLL drivers

We aimed to understand how signalling activities contribute to signatures associated with

major CLL subtypes that affect clinical outcome. The proliferative drive of CLL (CLL-PD), linked

to prognosis and increased mTOR-MYC-OXPHOS activity, was significantly reduced upon

AKT, BET and mTOR inhibition, demonstrating a direct dependency of pro-proliferative

capacities of CLL cells on these pathways[12] (Fig. 3f). The IGHV status is the strongest

modulator of the transcriptional variation in CLL and a key determinant of outcome (Fig. 2c,

Suppl. Fig. 2f-j). Some well-known markers of IGHV status, i. e. *ZAP70* and *CD38*, were stable

after drug treatment, while other signature genes, i. e. *CCL3* and *CD72*, were dependent on

BCR activity, suggesting that active signalling maintains part of the IGHV signature (Suppl.

Fig. 7a,b). To test the impact of the drugs on the overall signature, we quantified the

differentially expressed genes between patients with unmutated and mutated IGHV with each

treatment and found that the signature size was moderately reduced by PI3K, mTOR and BTK

inhibition (Fig. 3g). This partial reversal was driven by downregulation of IGHV signature genes

in unmutated patients which enriched in the cholesterol and glucose metabolism, allograft

rejection, adipogenesis, mTORC1 signalling and IFN signalling gene sets (Suppl. Fig. 7c,d).

When the BTK and IRAK4 inhibitors were combined, we observed the most pronounced

collapse of the IGHV signature, with an enrichment of genes linked to ribosomes and

translation, in line with a more pronounced effect on these downstream consequences when

the BCR and TLR were simultaneously perturbed[32] (Fig. 3g, Suppl. Fig. 7c,e). To understand

to what extent BCR and TLR signalling cooperate in CLL, we assessed how the sum of the

individual BTK and IRAK4 inhibitor effects - with the expectation that there is no interaction -

compared to the observed effect from the BTK/IRAK4 inhibitor combination. We found that

genes affected by BTK, IRAK4 or BTK/IRAK4 inhibition showed stronger effects than what

would be expected from the sum of the individual effects (Suppl. Fig. 7f). Cooperation of TLR

and BCR signalling has been described in other B-NHL, where it is linked to activating gene

mutations targeting the pathway[33,34]. Our finding suggests a cooperation between TLR and

BCR signalling in CLL even in the absence of such a driver mutation.

Trisomy 12 occurs in 15%–20% of CLL patients and is considered an early event[26]. It is

associated with disease progression and a higher incidence of aggressive transformation, but

its molecular function is not well understood[35,36]. We repeated the same analysis with the

trisomy 12 signature. Trisomy 12 signature genes showed partial dependencies on BTK, PI3K

and mTOR (Fig. 3h, Suppl. Fig. 7g-i). Differential expression of signature genes encoded on

chromosome 12 was largely stable and not affected by disruption of signalling activities, in line

with a gene dosage-driven "passenger" signature with no driver role (Fig. 3h). In contrast, we

found more genes to discriminate trisomy 12 after MEK, BTK/IRAK4 and MDM2 inhibitor

treatment. These findings suggest a modulation of these pathways by trisomy 12 (Fig. 3h,

Suppl. Fig. 7j).

Disease drivers influence pathway connectivity in CLL

Epistasis describes a phenomenon of genetic interactions in which the phenotype of a mutation

is modified by the presence or absence of mutations in one or more genes in a non-additive

manner[37]. In this study, we adopted this concept and systematically tested for drug-gene

interactions in linear models and identified modifications in the transcriptional phenotype of a

chemically perturbed gene target in the presence of unmutated IGHV and the common disease

drivers trisomy 12, *BRAF*, deletion 17p, deletion 15q and *TP53* (Fig. 4a, Suppl. Fig. 8a). We

then grouped the observed phenotypes of the epistatic interactions based on the direction of

which perturbations mask, modify, or otherwise influence the expression of genes. We

identified genes upregulated by a given aberration which were downregulated in the presence

of both perturbation and driver to levels below baseline and vice-versa (inversion; n = 247).

Suppression of the driver phenotype in the presence of a perturbation was observed for 112

genes. We also observed synergy (n = 94), where drug effects are amplified and only

discernible in the presence of both aberration and drug treatment. Suppression of drug effects

in the presence of an aberration was found for 60 genes (Fig. 4b-c, Suppl. Fig. 8b-d). We used

these categories to understand the functional implications of disease drivers based on known

drug targets or mechanisms. A clear example for loss of function (LOF) in CLL was observed

with *TP53* mutations or deletion 17p, which showed impaired transcriptional activation of p53

targets, e. g. *MDM2*, *CDKN1A*, in response to MDM2 inhibitor treatment (≙ suppression of

drug effect) (Fig. 4d-f, Suppl. Data Fig. 8d). Consistent with the activation of the MAPK

pathway, *BRAF* mutations showed most interactions with MEK inhibition (synergy and driver

suppression type (Fig. 4d, Suppl. Fig. 8d)). We also identified so far unknown interactions

linking mTOR inhibition to deletion 15q (Fig. 4d, Suppl. Fig. 8d). For unmutated IGHV, we

observed interactions with mTOR predominantly of the driver and drug suppression type (Fig.

4d, Suppl. Fig. 8d). We also found important therapeutic targets in lymphoma, e. g. *CD19* and

MS4A1/CD20, in the synergy group. These findings suggest that expression of these targets

are dependent on a certain pathway in molecular subgroups which could serve as a basis for

designing new treatment strategies (Fig. 4g,h).

Most drug-gene interactions were observed for trisomy 12 (n = 314) characterised by inversion

and synergy with BTK, MEK, IRAK4 and combined BTK/IRAK4 inhibition (Fig. 4d,i-l, Suppl.

Fig. 8d). The effect of IRAK4 inhibition in the presence of trisomy 12 is remarkable as the

inhibitor's effect was undetectable when the complete cohort was analysed (Fig. 2g). We

identified most drug-gene interactions were modulated by trisomy 12 with IRAK4 (n = 114),

combined BTK/IRAK4 (n = 83) and BTK inhibition (n = 39) providing further evidence for

crosstalk and cooperative effects between BCR and TLR signalling in trisomy 12 (Fig. 4d,i-l).

To demonstrate the role of IRAK4 as a mediator of TLR signalling in CLL, we treated patient

samples with either IRAK4 inhibitor alone or in combination with the TLR9 agonist CpG. TLR9

stimulation led to increased viability in line with a pro-tumourigenic role of TLR signalling, which

was abrogated by IRAK4 inhibition (Suppl. Fig. 9a). We searched for focal gains in the CNV

profiles of our CLL cohort (n = 366) which may allow restricting the number of candidate genes

targeted by trisomy 12, and identified a CLL case with a focal gain between chr12:42,538,673

and chr12: 44,229,999. This gain included nine protein-coding genes including *IRAK4* (Fig. 5a-

c, Suppl. Fig. 9b). We clustered gene expression of CLL samples (n = 210) based on the

trisomy 12 baseline signature and restricted the analysis to genes not encoded on

chromosome 12 (642 out of 917 total trisomy 12 signature genes) to prevent clustering due to

gene dosage. The sample with the focal gain of *IRAK4* clustered with the trisomy 12 samples

demonstrating the phenotype resemblance (Fig. 5c).

Our findings suggest that trisomy 12 confers increased TLR signalling capacity driven by the

increased *IRAK4* expression from gene dosage and cooperates with BCR signalling.

Discussion

We performed a functional perturbation screening directly in cancer cells at scale to capture

drug-induced global gene expression changes with low-depth RNA sequencing. Prior studies

investigating perturbation-induced gene expression changes have exploited single-cell

technology (few patients, many cells e. g. MIX-seq[38], RASL-seq[39] or DRUG-seq[40]) and focused

on establishment of technical frameworks, but less on exploiting the signatures to dissect

disease biology and cellular processes. The Connectivity Map focused on the differences and

signatures induced by a large number of drugs in fewer cell lines[18,19]. By design, it therefore

has unique resolution on the drug effects and contributes to the understanding how drugs work.

Our approach complements the Connectivity Map by leveraging many primary tumours from

an individual disease, and resolving the effects of a limited number of clinically used drugs on

a broad and representative sampling of molecularly diverse tumours without immortalisation

artefacts. We exploit the drug effects on transcriptomes to resolve disease mechanisms as

well as inter-patient heterogeneity. This design allowed us to measure downstream effects of

signalling events even if effects are subtle. Our findings suggest that insights and hypotheses

from gene expression studies on primary tumours may be vastly increased by the analysis of

such perturbed samples as the ability to specifically target molecules and pathways grows.

Multiple clinical inhibitors targeting PI3K, BTK, mTOR and MEK, were used in this study as

optimal combination and clinical assignment to patients is a challenge, which cannot be easily

addressed in trials[41].

We used interaction testing to study the effect of gene mutations on the perturbed gene

expression, e.g. by LOF mutations (*TP53*) or GOF (trisomy 12, *BRAF*). A particularly notable

example is trisomy 12, present in approximately 20% of CLL patients. We identified IRAK4 as

a driver in CLL with trisomy 12, highlighting how perturbation transcriptomics can uncover

actionable targets within genetically defined subgroups. In line with our proposed mechanism

where IRAK4 is a key target, trisomy 12 has been found to be strongly associated with

response to TLR stimuli such as resiquimod (TLR7/8) and CpG-ODN (TLR9)[42]. Reid et al.

studied genes targeted by trisomy 12 in a human pluripotent cell line model and identified

IRAK4 inhibition to be potent against human trisomy 12-harboring CLL cells in patient-derived

xenografts[43].

Our experiments enable the deconvolution of drug effects by factor analysis, and facilitate the

transition to patient-specific pathway assembly read-outs by drug perturbation and global

expression analysis (Fig. 6). The identification of compensatory pathways (factor 14, 20) may

provide a basis for optimised combination treatment. While prior drug response perturbation

studies have hinted at some of these findings and structure[11], the use of a large set of tumours

and high-dimensional readout of gene expression gives a more comprehensive and systematic

picture of the connectivity of signalling pathways in CLL, where effects of inhibitors on

individual pathways can be deconvoluted and linked to other drugs. On the route to improved

precision medicine for individual cancer patients, the workflows established in our study

provide a framework to measure pathway connectivity of primary cancer cells using clinically

relevant drugs and adapting the treatment along the lines of measured pathway activity which

can be generated by the projection of the individual patients response compared to a larger
reference set of measurements such as the one presented here. While the precise strategy for
optimal drug combinations will need to be demonstrated, our data provides a roadmap for such
experimentation.

Drug-perturbed transcriptome profiling measures connectivity of cancer pathway
dependencies and functionally annotates mutations. As more specific drugs become available,
tumour pathway wiring could be fully resolved and improve precision oncology, moving closer
to treatments guided by measured pathway activity rather than inferred potential.

Material and Methods

Patient samples and study approval

331 Patient material was acquired in accordance with the Declaration of Helsinki and approval from
332 the local ethics committees (S-206/2011; S-356/2013, 2009-0062; 2019/01744), after written
333 informed consent. Detailed patient information is listed in Suppl. Table 1. DNA mutations,
334 CNVs, DNA methylation and gene expression were performed as described previously[11].

Isolation and cryoconservation of MNCs from whole blood

336 Mononuclear cells from peripheral blood were isolated using Ficoll-Paque (Cytiva) density
337 gradient centrifugation, cryopreserved in freezing medium (72.5% RPMI-1640 (Gibco), 20%
338 fetal bovine serum (FBS, Gibco), 7.5% DMSO (Sigma-Adrich)) and stored at -180°C in liquid
339 nitrogen.

Compounds

Small-molecules (except compound 26 (Calbiochem)) were purchased from Selleckchem and
reconstituted in DMSO at 10 mM.

In a pilot, we treated four CLL samples with concentrations based on results from multiple prior
drug screens[11,12] with average toxicity below 20% (Suppl. Methods Fig. 1a,b) and confirmed
drug-specific gene expression changes in line with the drugs' mechanisms of action (Suppl.

Methods Fig. 1c-h). For the screen, stocks were diluted to 200-fold of the final assay

concentrations with DMSO as "ready-to-use" aliquots and stored at -20°C.

Ex-vivo drug treatment and cell viability assessment

Primary samples were thawed and incubated in RPMI-1640 supplemented with 10% heat-

inactivated human serum from male AB clotted whole blood (Sigma-Aldrich), 1% 10,000 U/mL

penicillin/10,000 µg/mL streptomycin (P/S, Gibco) and 2% 200 mM L-glutamine (Gln, Gibco)

(RPMI-HS) on a roller mixer for 3h at room temperature to allow recovery from the thawing

process and support the removal of DMSO from the freezing medium. The medium was

replaced and 5 x 10^6 cells were treated with each drug (Suppl. Methods Fig. 1c) and cultured

at 37°C and 5% CO_2 for 48h. Cells were harvested, washed twice with cold phosphate-buffered

saline (Gibco), lysed in 600 µL TRI Reagent (Zymo Research) and stored at -80°C until RNA

extraction.

Sample viabilities of each sample (before and after perturbation) were assessed on the flow

cytometer LSR II Fortessa with FACSDiva v8.0.1 software (BD Biosciences) and analysed with

FlowJo v10.10.0 (FlowJo LLC) (Suppl. Method Fig. 2).

TLR9 stimulation and ATP-based viability assessment

For the 5x concentrated working dilution, a 5-point 5-fold serial dilution of IRAK4i was prepared

with a starting concentration of 50 µM (≙ 10 µM–0.016 nM final assay concentration). For the

negative control, 5 µL of 0.5% DMSO working dilution was pre-plated (≙ 0.1% final

concentration).

MNCs from 20 CLL patients were thawed and incubated for 3h on the roller mixer prior to

seeding as described above and reconstituted to 1 x 10^6/mL (1.25X) cell concentration. In

unstimulated conditions, 20 µL (≙ 20,000 cells) were added to each well containing the pre-

plated DMSO or IRAK4i. For the CpG stimulation, cells were supplemented with 1.25 µg/mL

of the TLR9 agonist ODN2006 (InvivoGen) before seeding to reach a final assay concentration

of 1 µg/mL. Cells were incubated at 37°C and 5% CO_2 for 48h. For the ATP-based viability

assessment, plates were spun down briefly before adding 7 µL CellTiter-Glo (Promega) to

each well. The reaction was incubated for 20 min at RT and luminescence measured on a

BioTek Synergy LX multimode microplate reader (Agilent). To obtain normalised viabilities,

intensity values were divided by the median of all DMSO values for each condition and

concentration for each patient.

RNA extraction

Frozen lysates were thawed at 37°C for 3 min. RNA was precipitated by thoroughly mixing 100

µL of 100% ethanol (Sigma-Aldrich) with the lysate. RNA was extracted with the Direct-zol

RNA MiniPrep kit (Zymo Research) including on-column DNase treatment (30U per sample)

for 15 min at room temperature according to the manufacturer's protocol and eluted in 25 µL

nuclease-free water (QIAGEN). The concentration was determined with Quant-it RiboGreen

RNA assay (ThermoFisher Scientific) on a BioTek Synergy LX multimode microplate reader.

RNA was stored at -80°C in 5 µL nuclease-free water until use for library preparation and RNA

sequencing.

Library preparation and 3' RNA sequencing

For the pilot screen (4 patients) 200 ng of total RNA were used for cDNA library generation

with the QuantSeq 3′ mRNA-Seq Library Prep Kit FWD for Illumina (Lexogen). Libraries from

42 samples were pooled and sequenced on the NextSeq500 (Illumina) using 75 bp and single-

end reads yielding 3.9 to 13.5 million reads. For the comparison between the library

preparation methods, baseline profiles of 5 patients generated by whole-transcript deep

sequencing (TruSeq, Illumina)[13], the same RNA material was sequenced in the pilot screen.

For the remaining 112 patients, cDNA libraries were prepared from 100 ng total RNA with the

QuantSeq 3′ mRNA-Seq Library Prep Kit FWD for Illumina (Lexogen). Up to 48 libraries (22-

48) were pooled and sequenced on the HiSeq 4000 (Illumina) using 50 bp and single-end

reads yielding 2.3 to 21.7 million reads.

Processing of the 3'-RNA sequencing data

The same libraries from the first sequencing run were sequenced twice and the resulting reads

were pooled. The demultiplexed RNA sequencing FASTQ files were mapped to the ENSEMBL

human reference genome release 104 (Homo sapiens GRCh38.104) using STAR (version

2.7.9a) with default parameters. The aligned reads were summarised into per-gene counts

using htseq-count (version 0.9.1) with default parameters in union mode, counting only those

reads unambiguously mapped to a single gene. The resulting gene count matrix was imported

into R (version 4.5.1) for further processing. Given the shallow sequencing depth, the gene

count matrix exhibited higher sparsity compared to traditional bulk RNA-seq data. This sparsity

posed challenges for accurately estimating between-sample normalisation factors using

methods designed for deep RNA sequencing, such as DESeq2[44]. To address this,

scTransform[45], a method that applies regularised negative binomial regression to sparse gene

count matrices from single-cell sequencing data, was used to estimate the normalisation

factors. These factors were then supplied to DESeq2 (version 1.44) for variance-stabilising

transformation and differential expression analysis. For all the downstream analyses, only

protein-coding genes with minimum 10 counts in at least one sample were retained.

The baseline profiles were combined with profiles generated by 3'-end shallow depth

sequencing (QuantSeq, Lexogen) and normalised for library size before comparative analysis.

Data analysis

Statistics

Correlation trends were indicated by Pearson correlation (R) and linear regression lines with

95% confidence intervals. Density distributions of single values were visualised with sina plots

and summarised with Tukey-style box and whiskers plots. Statistical differences between

groups were computed with unpaired two-tailed Student's t-test when groups contained

samples from different patients, e. g. mutant vs. wild-type. Paired two-tailed Student's t-tests

were used when conditions of the same patients were compared. Nominal P-values were

corrected for multiple testing using the Benjamini-Hochberg procedure and indicated as false-

discovery rate (FDR).

Dimensionality reduction and clustering analysis

Dimensionality reduction including principal component analysis (PCA), *t*-distributed Stochastic Neighbor Embedding (*t*-SNE) and hierarchical clustering were conducted on gene counts after variance-stabilising transformation (*vst* function from DESeq2 package). PCA (*prcomp* function from base R) and *t*-SNE (*tsne* function from M3C package[46], version 1.26) were performed on the 1,000 most variable genes, unless indicated. Hierarchical clustering was carried out using the Ward.D2 method based on Euclidean distances computed from the centered and scaled expression values of genes selected from differential expression analyses. The selection criteria are detailed in the figure legends. The ComplexHeatmap package[47] (version 2.20) was used for visualisation. To analyse expression patterns after drug treatment, patient-specific variations were adjusted in the gene expression matrix before dimension reduction (*removeBatchEffect* function from limma package[48], version 3.60.6). The application of this adjustment is noted in the respective figure legends.

Differential expression analysis

All differential expression analyses were conducted on the scTransform-adjusted gene count matrix using the R package DESeq2[44] (versions 1.44.0 and 1.48.2).

To compute genes differentially expressed between drug- and DMSO-treated samples, patient IDs were included as a covariate in the design formula to account for patient-specific variation. Log2-fold changes were shrunken with the *lfcShrink* function using the apeglm[49] method. Genes were filtered by significance (*FDR* < 0.5%) and effect size (|fold change| > 1.05).

p53 expression scores were downloaded from the Target Gene Regulation Database 2.0[50].

The CLL-PD score was computed from the Z-score gene count matrix using *CLLPDestimate* from the mofaCLL package[12] (version 0.0.9).

The transcriptional signature of a disease dimension was defined as the set of differentially expressed genes between molecular subgroups individually for each perturbation. To account for the confounding effects of IGHV and trisomy 12 status, these variables were included in the design formula. To compare the number of signature genes between perturbations, the

90% confidence interval was estimated using bootstrapping (1,000 times, sampling 90% of

patients while maintaining the ratio between wild-type and mutated/CNV patients).

Drug-genotype interaction testing and epistasis clusters

The interactions between genetic aberrations and drug effects were modeled by linear

interaction testing with DESeq2 using a design formula that accounts for aberration, patient ID

and treatment condition. Drug-gene interaction genes were filtered by *FDR* (< 10%). Median

and scaled expression was calculated for each gene and group (DMSO/wild-type, drug/wild-

type, DMSO/mut, drug/mut) and normalised against DMSO/wild-type as the reference.

Interaction types were grouped using soft Fuzzy C-Means Clustering (FCM) with the *cmeans*

function (centers = 20, iter.max = 1000, dist = "euclidian", method = "cmeans", m = 1.4) from

the e1071 package[51] (version 1.7.16). FCM was repeated three times for genes with probability

scores below 0.85 until every gene was assigned to a cluster with a probability score of at least

0.85, resulting in 60 pre-clusters. Pre-clusters with similar patterns were summarised to 20

final clusters using harder FCM (m = 1.1) and manual curation.

Pathway connectivity networks

Pathway connectivity networks of significant DEGs are shown as bipartite graphs in which

perturbation and DEGs denote the disjoint and independent sets of nodes and are connected

by the edges. The edges are weighted by the significance level of the drug effects. For the

connectivity network of interactions, nodes were filtered for aberrations with more than 5

mutated/CNV cases. The edges are weighted by the significance level of the interaction. Node

coordinates were computed using the network package[52] (version 1.19.0).

Guided sparse factor analysis

Guided Sparse Factor Analysis (GSFA) was conducted using the GSFA package (version

0.2.8)[31]. The gene count matrix was pre-processed by accounting for the patient-specific

variation and selecting the 5,000 genes with the largest deviance after deviance residual

transformation[53]. DMSO-treated samples were indicated as negative controls, and a

"mixture_normal" distribution was used as the sparse prior. Singular value decomposition was

employed to initialise the factor values, followed by 3,000 iterations of Gibbs sampling to train

the model. Posterior means were calculated using the last 1,000 iterations. All other

parameters were used with default settings.

The model with 20 factors was chosen for further analysis (Suppl. Method Fig. 3). Perturbations

and genes with association effect sizes above 0.25 and posterior inclusion probabilities (*PIP*)

above 0.95 were considered to have a significant effect on a particular factor.

Multiple linear regression was conducted with Lasso penalisation, leave-one-out cross-

validation, mean squared error loss, 200 iterations, mutations/CNVs as predictors and

centered and scaled individual drug effects as response vectors using the glmnet package[54]

(version 4.1-8).

Pathway enrichment analysis

Parametric analysis of gene set enrichment[55] was performed on the DEGs (ranked by *t*-

statistics) or genes associated with GSFA factors (ranked by factor loading). Human gene set

collections (version 2024.1) were downloaded from MSigDB ([https://www.gsea-](https://www.gsea-msigdb.org/gsea/msigdb/human/collections.jsp)

[msigdb.org/gsea/msigdb/human/collections.jsp](https://www.gsea-msigdb.org/gsea/msigdb/human/collections.jsp)). The decoupleR package (version 2.14.0)

was used to infer the impact on transcription factor activity by each drug treatment[28]. The filter

criteria for model input and results are indicated in the respective figure legends.

Data availability

All RNA sequencing data generated in this study will be available at European Genome-

Phenome Archive (EGA) upon publication. The sequencing data generated in our previous

studies and used in the current study, including whole-exome/whole-genome sequencing,

targeted-sequencing, DNA methylation profiling and RNA sequencing of baseline patient

samples, are available at EGA under the accession code EGAS00001001746. Processed

data, including RNA sequencing data, drug sensitivity profiling and sample metadata are

available in our project repository at GitHub (https://github.com/THLDo/CLL_Connectivity).

The RNA drug perturbation dataset can also be explored interactively via the Shiny app

available upon publication.

Code availability

The computational codes for reproducing all figures and results reported in this article will be provided in our project repository at GitHub (https://github.com/THLDo/CLL_Connectivity) under the GNU General Public License v3.0 upon publication.

References

1. Knisbacher, B. A. *et al.* Molecular map of chronic lymphocytic leukemia and its impact on outcome. *Nat. Genet.* **54**, 1664–1674 (2022).

2. Nadeu, F., Diaz-Navarro, A., Delgado, J., Puente, X. S. & Campo, E. Genomic and Epigenomic Alterations in Chronic Lymphocytic Leukemia. *Annu. Rev. Pathol. Mech. Dis.* **15**, 149–177 (2020).

3. Robbe, P. *et al.* Whole-genome sequencing of chronic lymphocytic leukemia identifies subgroups with distinct biological and clinical features. *Nat. Genet.* **54**, 1675–1689 (2022).

4. Burger, J. A. Treatment of Chronic Lymphocytic Leukemia. *N. Engl. J. Med.* **383**, 460–473 (2020).

5. Oakes, C. C. *et al.* DNA methylation dynamics during B cell maturation underlie a continuum of disease phenotypes in chronic lymphocytic leukemia. *Nat. Genet.* **48**, 253–264 (2016).

6. Kulis, M. & Esteller, M. DNA Methylation and Cancer. in *Advances in Genetics* vol. 70 27–56 (Elsevier, 2010).

7. Damle, R. N. *et al.* Ig V Gene Mutation Status and CD38 Expression As Novel Prognostic Indicators in Chronic Lymphocytic Leukemia. *Blood* **94**, 1840–1847 (1999).

8. Hamblin, T. J., Davis, Z., Gardiner, A., Oscier, D. G. & Stevenson, F. K. Unmutated Ig V(H) genes are associated with a more aggressive form of chronic lymphocytic leukemia. *Blood* **94**, 1848–1854 (1999).

9. Minden, M. D. *et al.* Chronic lymphocytic leukaemia is driven by antigen-independent cell-autonomous signalling. *Nature* **489**, 309–312 (2012).

10. Abruzzo, L. V. *et al.* Trisomy 12 chronic lymphocytic leukemia expresses a unique set of

activated and targetable pathways. *Haematologica* **103**, 2069–2078 (2018).

11. Dietrich, S. *et al.* Drug-perturbation-based stratification of blood cancer. *J. Clin. Invest.*

128, 427–445 (2017).

12. Lu, J. *et al.* Multi-omics reveals clinically relevant proliferative drive associated with

mTOR-MYC-OXPHOS activity in chronic lymphocytic leukemia. *Nat. Cancer* **2**, 853–864

(2021).

13. Lütge, A. *et al.* Subgroup-specific gene expression profiles and mixed epistasis in chronic

lymphocytic leukemia. *Haematologica* https://doi.org/10.3324/haematol.2022.281869

(2023) doi:10.3324/haematol.2022.281869.

14. Burd, A. *et al.* Precision medicine treatment in acute myeloid leukemia using prospective

genomic profiling: feasibility and preliminary efficacy of the Beat AML Master Trial. *Nat.*

Med. **26**, 1852–1858 (2020).

15. Irmisch, A. *et al.* The Tumor Profiler Study: integrated, multi-omic, functional tumor

profiling for clinical decision support. *Cancer Cell* **39**, 288–293 (2021).

16. Stratton, M. R., Campbell, P. J. & Futreal, P. A. The cancer genome. *Nature* **458**, 719–

724 (2009).

17. The International Cancer Genome Consortium. International network of cancer genome

projects. *Nature* **464**, 993–998 (2010).

18. Lamb, J. *et al.* The Connectivity Map: Using Gene-Expression Signatures to Connect

Small Molecules, Genes, and Disease. *Science* **313**, 1929–1935 (2006).

19. Subramanian, A. *et al.* A Next Generation Connectivity Map: L1000 Platform and the

First 1,000,000 Profiles. *Cell* **171**, 1437-1452.e17 (2017).

20. Ben Guebila, M. *et al.* GRAND: a database of gene regulatory network models across

human conditions. *Nucleic Acids Res.* **50**, D610–D621 (2022).

21. Cheng, J., Yang, L., Kumar, V. & Agarwal, P. Systematic evaluation of connectivity map

for disease indications. *Genome Med.* **6**, 540 (2014).

22. Musa, A. *et al.* A review of connectivity map and computational approaches in

pharmacogenomics. *Brief. Bioinform.* **19**, 506–523 (2018).

23. Moll, P., Ante, M., Seitz, A. & Reda, T. QuantSeq 3′ mRNA sequencing for RNA

quantification. *Nat. Methods* **11**, i–iii (2014).

24. Oren, M. & Prives, C. p53: A tale of complexity and context. *Cell* **187**, 1569–1573 (2024).

25. Herishanu, Y. *et al.* The lymph node microenvironment promotes B-cell receptor

signaling, NF-κB activation, and tumor proliferation in chronic lymphocytic leukemia.

Blood **117**, 563–574 (2011).

26. Landau, D. A. *et al.* Mutations driving CLL and their evolution in progression and relapse.

Nature **526**, 525–530 (2015).

27. Rendeiro, A. F. *et al.* Chromatin mapping and single-cell immune profiling define the

temporal dynamics of ibrutinib response in CLL. *Nat. Commun.* **11**, 577 (2020).

28. Badia-i-Mompel, P. *et al.* decoupleR: ensemble of computational methods to infer

biological activities from omics data. *Bioinforma. Adv.* **2**, vbac016 (2022).

29. Zaware, N. & Zhou, M.-M. Bromodomain biology and drug discovery. *Nat. Struct. Mol.*

Biol. **26**, 870–879 (2019).

30. Azmi, A. S., Uddin, M. H. & Mohammad, R. M. The nuclear export protein XPO1 — from

biology to targeted therapy. *Nat. Rev. Clin. Oncol.* **18**, 152–169 (2021).

31. Zhou, Y., Luo, K., Liang, L., Chen, M. & He, X. A new Bayesian factor analysis method

improves detection of genes and biological processes affected by perturbations in single-

cell CRISPR screening. *Nat. Methods* **20**, 1693–1703 (2023).

32. Kielbassa, K. *et al.* Ibrutinib sensitizes CLL cells to venetoclax by interrupting TLR9-

induced CD40 upregulation and protein translation. *Leukemia* **37**, 1268–1276 (2023).

33. Mandato, E. *et al.* MYD88L265P augments proximal B-cell receptor signaling in large B-

cell lymphomas via an interaction with DOCK8. *Blood* **142**, 1219–1232 (2023).

34. Phelan, J. D. *et al.* A multiprotein supercomplex controlling oncogenic signalling in

lymphoma. *Nature* **560**, 387–391 (2018).

35. Döhner, H. *et al.* Genomic Aberrations and Survival in Chronic Lymphocytic Leukemia.

N. Engl. J. Med. **343**, 1910–1916 (2000).

36. Fabbri, G. & Dalla-Favera, R. The molecular pathogenesis of chronic lymphocytic

leukaemia. *Nat. Rev. Cancer* **16**, 145–162 (2016).

37. Van De Haar, J. *et al.* Identifying Epistasis in Cancer Genomes: A Delicate Affair. *Cell*

177, 1375–1383 (2019).

38. McFarland, J. M. *et al.* Multiplexed single-cell transcriptional response profiling to define

cancer vulnerabilities and therapeutic mechanism of action. *Nat. Commun.* **11**, 4296

(2020).

39. Li, H., Qiu, J. & Fu, X. RASL-seq for Massively Parallel and Quantitative Analysis of

Gene Expression. *Curr. Protoc. Mol. Biol.* **98**, (2012).

40. Ye, C. *et al.* DRUG-seq for miniaturized high-throughput transcriptome profiling in drug

discovery. *Nat. Commun.* **9**, 4307 (2018).

41. O'Dwyer, P. J. *et al.* The NCI-MATCH trial: lessons for precision oncology. *Nat. Med.* **29**,

1349–1357 (2023).

42. Bruch, P. *et al.* Drug-microenvironment perturbations reveal resistance mechanisms and

prognostic subgroups in CLL. *Mol. Syst. Biol.* **18**, e10855 (2022).

43. Reid, J. C. *et al.* Human pluripotent stem cells identify molecular targets of trisomy 12 in

chronic lymphocytic leukemia patients. *Cell Rep.* **34**, 108845 (2021).

44. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

45. Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-

seq data using regularized negative binomial regression. *Genome Biol.* **20**, 296 (2019).

46. John, C. R. *et al.* M3C: Monte Carlo reference-based consensus clustering. Preprint at

https://doi.org/10.1101/377002 (2018).

47. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

48. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing

and microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).

49. Zhu, A., Ibrahim, J. G. & Love, M. I. Heavy-tailed prior distributions for sequence count

data: removing the noise and preserving large differences. *Bioinformatics* **35**, 2084–2092

(2019).

50. Fischer, M., Schwarz, R., Riege, K., DeCaprio, J. A. & Hoffmann, S. TargetGeneReg 2.0:

a comprehensive web-atlas for p53, p63, and cell cycle-dependent gene regulation. *NAR*

Cancer **4**, zcac009 (2022).

51. Meyer, D., Dimitriadou, E., Hornik, K., Weingessel, A. & Leisch, F. e1071: Misc Functions

of the Department of Statistics, Probability Theory Group (Formerly: E1071), TU Wien.

1.7-16 https://doi.org/10.32614/CRAN.package.e1071 (1999).

52. Butts, C. T. **network** : A Package for Managing Relational Data in *R. J. Stat. Softw.* **24**,

(2008).

53. Townes, F. W., Hicks, S. C., Aryee, M. J. & Irizarry, R. A. Feature selection and

dimension reduction for single-cell RNA-Seq based on a multinomial model. *Genome*

Biol. **20**, 295 (2019).

54. Friedman, J., Hastie, T. & Tibshirani, R. Regularization Paths for Generalized Linear

Models via Coordinate Descent. *J. Stat. Softw.* **33**, (2010).

55. Kim, S.-Y. & Volsky, D. J. PAGE: Parametric Analysis of Gene Set Enrichment. *BMC*

Bioinformatics **6**, 144 (2005).

56. Landau, D. A. *et al.* The evolutionary landscape of chronic lymphocytic leukemia treated

with ibrutinib targeted therapy. *Nat. Commun.* **8**, 2185 (2017).

Acknowledgements

T. Zenz was supported by the CRPP 'Next Generation Drug Response Profiling for

Personalized Cancer Care', the Swiss Cancer Research foundation (KFS-4439-02-2018), the

Monique-Dornonville-de-la-Cour Stiftung and the Helmut Horten Stiftung. M. F. Pohly, W.

Huber were supported by the INTeRCePT, and T. Zenz by the INTeRCePT and POLAR

(Uniscientia Stiftung) projects funded by The LOOP Zurich. J. Lu, C. Lohoff and W. Huber were

supported by the SMART-CARE project funded by the Federal Ministry of Research,

Technology and Space of Germany (161L0213 and 16LW0237). For access to infrastructure,

we thank M. Bawohl, M. Rechsteiner and A. Weber at the Department of Pathology and

Molecular Pathology, and the Functional Genomics Center Zurich. For access to the EMBL

Heidelberg HPC Cluster, we gratefully thank J. Pečar and EMBL IT Services.

Contributions

THLD, JL, WH, TZ conceptualised the study and designed the experiments. THLD, SK, MFP

and FJ conducted experiments. SS, JH, WH, JL and TZ were responsible for data curation.

CL, FJ and VB performed RNA sequencing and data pre-processing. THLD, CL, JL and TZ

performed data analysis and data visualisation. THLD, CL, JL, WH and TZ interpreted the data.

THLD, CL, JL, WH and TZ wrote the original draft of the manuscript. All authors reviewed and

edited the manuscript.

Ethics declarations

Competing interests

656 TZ received consulting fees from Astra-Zeneca, BeiGene, AbbVie, Janssen, Novartis, Lilly,

657 Roche, Bristol-Myers Squibb, and Gilead Sciences, and payment for lectures from Astra-

658 Zeneca, BeiGene, AbbVie, Janssen, Novartis, Lilly, Roche, Bristol-Myers Squibb, and Gilead

659 Sciences. All other authors declare no competing interests.

Figures

Fig. 1: Study overview

a, Study outline (https://BioRender.com/6i76n18). **b**, Genetic landscape of study cohort including recurrent genetic aberrations (> 5 cases), clinical and demographic parameters, and availability of additional molecular omics profiling data. Colours for continuous variables are scaled by minimum value, 25th, 50th, 75th percentile and maximum value. Numbers indicated in brackets show case number and frequency of patients with respective aberration for the CLL cohort.

Fig. 2: Transcriptional landscape of drug effects in CLL

a, First two principal components of PCA based on the 1,000 most variant genes coloured by

perturbation and **b**, absolute viability of each drug-treated sample measured by flow cytometry.

c, PC2 and PC3 of PCA based on the 1,000 most variant genes coloured by IGHV and trisomy

12 status. **d**, t-SNE plot based on the 1,000 most variant genes coloured by patient and **e**, by

methylation groups. Samples from the same patient form tight clusters. **f**, t-SNE plot based on

the 1,000 most variant genes after adjustment for the patient effect coloured by perturbation.

g, Number of differentially expressed genes (DEGs) between drug- and DMSO-treated

samples (0.5% *FDR*, |fold change| > 1.05). Induction or suppression for each treatment are

shown. **h**, Volcano plot summarising the differential gene expression after MDM2 inhibitor

treatment. Genes are annotated according to p53 expression score[50]. **i**, GSEA with DEGs after

MDM2 inhibitor treatment (0.5% *FDR*) using hallmark and canonical pathways gene sets.

Gene sets were filtered by *FDR* (< 1%) and number of enriched genes (> 10). **j**, Heatmap with

hierarchical clustering showing patient effect-adjusted normalised expression of top 40 DEGs

after MDM2 inhibitor treatment. *TP53* mutation and deletion status are annotated. **k**, Volcano

plot and zoom-in summarising the differential gene expression after BTK inhibitor treatment. **l**,

GSEA with DEGs after BTK and PI3K inhibitor treatment (0.5% *FDR*) using hallmark and

canonical pathways gene sets. Gene sets were filtered by *FDR* (< 1%) and number of enriched

genes (> 10). Depicted are the top 10 gene sets for BTK and PI3K inhibition. **m**, Scatter plot

of drug effect sizes for DEGs between PI3K and BTK inhibitor treatment (10% *FDR*). Colours

and shapes depict in which comparisons the gene was differentially expressed (0.5% *FDR*).

n, GSEA with DEGs between PI3K and BTK inhibitor treatment (10% *FDR*) using hallmark and

canonical pathways gene sets. Gene sets were filtered by FDR (< 10%) and number of

enriched genes (> 10). **o**, Inferred transcription factor (TF) activity using all genes. TFs were

filtered by *FDR* (< 10%) and number of targets in TF regulon (> 30 target genes).

Fig. 3: Pathway connectivity in CLL

a, Pathway connectivity network shown as a bipartite graph. Genes and perturbations are connected if the perturbation led to differential expression of that gene. The total number of DEGs for each treatment is depicted by the size of the central nodes. Colour and thickness of edges represent whether genes are induced or suppressed, and significance of drug effect, respectively. **b**, Heatmap with hierarchical clustering showing number of DEGs after pairwise drug-drug comparison (0.5% *FDR*). **c**, Estimated effects of drug perturbations on co-regulated gene modules represented by latent factors. The posterior inclusion probability (*PIP*) denotes the probability that a perturbation affects the factor loading. Factors with significant loadings (*PIP* > 0.95, |effect size| > 0.25) are shown. The sign was inverted for selected factors to reflect the direction of the drug effects. **d**, Heatmap showing the drug effects for 50 genes with the

highest gene weights in each factor. **e**, GSEA with the gene weights for each factor using

hallmark and canonical pathways gene sets. Gene sets were filtered by *FDR* (< 1%) and

number of enriched genes (> 10). The sign of weights was reverted in accordance to (c). **f**,

CLL-PD scores computed from the transcriptomics data for each individual treatment.

Significance levels were calculated using two-tailed paired Student's *t*-test with DMSO as

reference and adjusted for multiple testing. Asterisks denote *FDR* < 10%. **g**, Median number

of DEGs and 90% confidence interval (CI) between samples with unmutated and mutated

IGHV status for each treatment at 10% *FDR* split into genes with lower or higher expression in

U-CLL compared to M-CLL. **h**, Median number of DEGs and 90% CI between samples with

and without trisomy 12 for each treatment at 10% *FDR* split into genes with lower or higher

expression in trisomy 12 compared to non-trisomy 12, and whether genes are encoded on

chromosome 12.

722 **Fig. 4: Interaction of perturbation effects and disease drivers**

723 **a**, Pathway connectivity network of significant drug-genotype interactions (DGIs) shown as a

724 bipartite graph. Genes, drugs and disease aberration are connected if a differential

725 transcriptional drug response was observed in the presence of the genetic driver (*FDR* < 10%).

726 Drug-genotype pairs with at least 10 DGIs are displayed. **b**, Schematic classification of

727 epistasis categories based on median values from all genes in the respective groups. **c**,

728 Heatmap of relative gene expression of DGIs in the presence of either drug, genetic driver or

729 both compared to WT/DMSO samples by epistasis class. **d**, Proportion of aberration

730 contributing to each epistasis class (outer circle) and proportion of perturbations contributing

731 to each aberration within each epistasis class (inner circle) with more than 5 DGIs. **e**, Heatmap

732 visualising the effect sizes of genes with significant interaction between MDM2 inhibition and

del17p or *TP53*^mut (*FDR* < 10%). **f**, Effect of MDM2 inhibition on normalised expression of

MDM2 and *CDKN1A* by deletion 17p status. **g**, Heatmap visualising the effect sizes of genes

with significant interaction between MEK inhibition and trisomy 12 (*FDR* < 10%). **h**, Normalised

expression of MEK inhibition on *HELZ2* and *CD19* by trisomy 12 status. **i**, Heatmap visualising

the effect sizes of genes with significant interaction between IRAK4 inhibition and trisomy 12

(*FDR* < 10%). **j**, Effect of IRAK4 inhibition on normalised expression of *POLR3B*, *BCL2L12*,

DDX19A and *HMGB1* by trisomy 12 status. **k**, Heatmap visualising the effect sizes of genes

with significant interaction between combined BTK/IRAK4 inhibition and trisomy 12 (*FDR* <

10%). **l**, Effect of DMSO (red), IRAK4 (cyan), BTK (green) and BTK/IRAK4 inhibition (pink) on

normalised expression of *PELP1*, *DHX33* and *TRIM39* by trisomy 12 status.

Fig. 5: IRAK4 is a disease driver targeted by trisomy 12

a, CNV plot with ideogram of chromosome 12 for patient P0004. The red line shows the copy

numbers across the chromosome. **b**, Genomic track displaying the protein-coding genes on

the amplified locus chr12:42,550,907-chr12: 44,229,999 (hg19). **c**, Gene expression of IRAK4

by trisomy 12 status for the CLL cohort from Lütge et al.[13]. The significance level was

calculated using two-tailed Student's *t*-test and shown as nominal *P*-value. **d**, Heatmap

visualising the scaled gene expression of trisomy 12 signature genes excluding the genes on

chromosome 12 (n = 642) for the CLL cohort (n = 210) from Lütge et al.[13]

Fig. 6: Pathway diagram and dependency map in CLL

Pathway dependencies modulated by gene mutations/CNVs or IGHV status were identified by drug-genotype interaction analysis and depicted as activating (blue) or inhibitory (red) (https://BioRender.com/svmtjkd).

Supplementary figures

Suppl. Fig. 1: Comparison of shallow 3'-QuantSeq to regular coverage whole-transcript TruSeq sequencing

a, Overview of experimental workflow (https://BioRender.com/u4e6koa). **b**, Schematic comparison of read distribution along transcript between 3'-QuantSeq[23] and whole-transcript TruSeq (https://BioRender.com/sx3vfjw). **c**, Biased read coverage of transcript 3'-end for exemplary samples. **d**, Mapped biotypes, **e**, number of mapped reads, **f**, and number of detected genes by method. Colours depict whether genes were detected by both methods, only by TruSeq or only by QuantSeq. **g**, Correlation of log2 gene expression counts detected by both methods.

769 **Suppl. Fig. 2: Overview viability effect of drugs and transcriptional landscape**

770 **a**, Overview of completeness and **b**, number of available treatment conditions by patient. **c**,

771 DMSO-normalised sample viability after 48h treatment measured by flow cytometry and

772 separated by diagnosis. Significance levels were calculated using two-tailed paired Student's

773 *t*-test and adjusted for multiple testing. Asterisk denotes significant differences between drug-

774 and DMSO-treated samples (*FDR* < 10%). **d**, Genomic association with drug response for CLL

775 samples. Asterisk denotes associations with significant differences in drug response between

776 patients with and without respective genetic aberrations with more than 5 cases (*FDR* < 10%).

777 Significance levels were calculated using two-tailed Student's *t*-test and adjusted for multiple

778 testing. **e**, Boxplots of individual patient data shown for top significant associations (*FDR* <

779 10%). Significance levels were calculated using two-tailed Student's *t*-test and adjusted for

780 multiple testing. **f**, PCA based on 1,000 most variant genes from DMSO-treated CLL and

781 lymphoma samples (n = 107), **g**, only CLL samples (n = 100), and **h**, from overlapping CLL

782 patients from Lütge et al.[13] profiled by deep whole-transcript RNA sequencing (n = 95). **i**, PCA

783 based on 1,000 most variant genes from DMSO- and drug-treated CLL and lymphoma

784 samples (n = 1,086), **j**, only CLL samples (n = 1,010), **k**, and after adjustment for the patient

785 effect (n = 1,010).

Suppl. Fig. 3: Transcriptional signature associated with viability

a, Correlation of cell death (100% - sample viability) with PC1 for all samples and **b**, by treatment. Correlation trends are indicated by Pearson correlation coefficient (*R*) and adjusted for multiple testing. **c**, Heatmap showing normalised expression of top 50 correlated genes with cell death. Correlation trends are indicated by Pearson correlation coefficient (*R*). **d**, GSEA individually for each treatment using genes with significant correlations between expression and cell death (*FDR* < 0.5%) using hallmark gene sets and number of enriched genes (< 10). Gene sets were filtered by *FDR* (< 1%). Enrichments indicate positive (red) or negative (blue) associations with cell death.

Suppl. Fig. 4: Drug effects of BTK and PI3K inhibitors on BCR inhibition

a, Projection of data from Herishanu et al.[25] (*ex-vivo* BCR activation signature), Landau et al.[56] and Rendeiro et al.[27] *(in-vivo* BTK inhibitor signature) into volcano plot of differential gene expression after BTK inhibition. Shapes depict whether the gene is regulated into the same or different direction. **b**, Volcano plot summarising the differential gene expression after PI3K inhibition. **c**, GSEA with DEGs after BTK and PI3K inhibitor treatment (0.5% *FDR*) using hallmark and canonical pathways. Gene sets were then filtered by *FDR* (< 1%) and number of enriched genes (> 10). Depicted are all significant gene sets for PI3K inhibition. **d**, Volcano plot summarising the differential gene expression between PI3K and BTK inhibition. Colours and shapes depict in which comparisons the gene was deregulated (0.5% *FDR*). **e**, Normalised *STAT1* expression in samples treated with DMSO, BTK or PI3K inhibitor. Significance levels

808   were calculated using two-tailed paired Student's *t*-test and shown as nominal *P*-values. **f**,

809   Projection of STAT1, STAT2 and STAT3 transcription factor target genes into volcano plot of

810   differential expression after BTK and PI3K inhibition.

Suppl. Fig. 5: Drug effects of AKT, mTOR, MEK, XPO1 and BET inhibitors

a, Volcano plot summarising gene expression after AKT inhibitor treatment. **b**, GSEA with DEGs after AKT, mTOR and MEK inhibitor treatment (0.5% *FDR*) using hallmark and canonical pathways gene sets. Gene sets were filtered by *FDR* (< 1%) and number of enriched genes (> 10). Depicted are all significant gene sets for AKT and the top 10 gene sets for MEK and mTOR inhibition. **c**, Scatter plot of drug effect sizes for DEGs between AKT and PI3K inhibitor treatment (0.5% *FDR*). Colours and shapes depict in which comparisons the gene was deregulated (0.5% *FDR*). **d**, Volcano plot summarising gene expression after mTOR and **e**, MEK inhibitor treatment. **f**, Scatter plot of drug effect sizes for DEGs between MEK and mTOR inhibitor treatment (0.5% *FDR*). Colours and shapes depict in which comparisons the gene was deregulated (0.5% FDR). **g**, GSEA with genes differentially expressed between MEK and

mTOR inhibition (0.5% *FDR*) using hallmark and canonical pathways gene sets. Gene sets

were filtered by *FDR* (< 1%) and number of enriched genes (> 10). **h**, Volcano plot summarising

gene expression after BET inhibitor treatment. **i**, GSEA with DEGs after BET inhibitor treatment

(0.5% *FDR*) using hallmark and canonical pathways gene sets. Gene sets were filtered by

FDR (< 1%) and number of enriched genes (> 10). **j**, Volcano plot summarising gene

expression after XPO inhibitor treatment. **k**, GSEA with DEGs after XPO1 inhibitor treatment

(0.5% *FDR*) using hallmark and canonical pathways gene sets. Gene sets were filtered by

FDR (< 1%) and number of enriched genes (≥ 3). **l**, Effect of XPO1 (green) and MDM2 inhibition

(rose) on the expression of *TP53* and the p53 target genes *MDM2*, *CDKN1A*, *BBC3*.

Significance levels were calculated using two-tailed paired Student's *t*-test and shown as

nominal *P*-values.

Suppl. Fig. 6: Guided sparse factor analysis of drug effects

a, Heatmap shows drug effects for 50 genes with the highest gene weights for factors with

significant loadings. Top 10 genes with the highest gene weights for each factor are labelled.

b, Estimated effects of drug perturbations on co-regulated gene modules represented by 20

latent factors. The posterior inclusion probability (*PIP*) denotes the probability that a

perturbation affects the factor loading. Factors with significant loadings (*PIP* > 0.95, |effect

size| > 0.25) are shown. Asterisks denote factors for which the sign was inverted to reflect the

direction of the drug effects. **c**, GSEA with the gene weights for each factor using hallmark and

canonical pathways gene sets. Gene sets were filtered by *FDR* (< 1%) and number of enriched

genes (> 10). The sign of weights was reverted in accordance to (b). **d**, Influence of genetic

disease drivers on the contribution of the MDM2 inhibitor effect in factor 18, **e**, BTK inhibitor

846   effect in factor 17, and **f**, MEK inhibitor effect in factor 14 computed by multiple linear

847   regression.

 Suppl. Fig. 7: Pathway dependency of expression signatures and disease drivers

 a, Normalised expression of *ZAP70 and CD38* by IGHV status in DMSO or BTK inhibitor-

 treated samples. **b**, Normalised expression of *CCL3* and *CD72* by IGHV status in DMSO or

 BTK inhibitor-treated samples. Significance levels were calculated using two-tailed Student's

 t-test and shown as nominal *P*-values. **c**, GSEA with genes differentially expressed between

 U- and M-CLL (10% *FDR* in > 90% of bootstrapping performed to increase stringency) for each

 treatment using hallmark and canonical pathways gene sets. Gene sets were filtered by *FDR*

 (< 1%) and number of enriched genes (> 10). **d**, Differential expression of IGHV signature

 genes within the IFN response and **e**, ribosome gene set with BTKi, PI3Ki, mTORi, IRAK4i and

 BTKi/IRAK4i compared to DMSO. **f**, Expected independent additive effect from single BTK and

 IRAK4 inhibition against the observed effect from the BTK/IRAK4 inhibitor combination for

 down- and up-regulated BTK, IRAK4 and BTK/IRAK4 inhibitor-dependent genes (black dots).

 The expected independent effect was calculated using the Bliss independence model

 ($LFC_{expected} = LFC_{BTKi} + LFC_{IRAK4i} - LFC_{BTKi} * LFC_{IRAK4i}$) and compared to the observed effect

 ($LFC_{BTKi/IRAK4i}$). The gray dots depict the background distribution of expected additive effect for

all genes (n_{Down} = 8,821, n_{Up} = 8,040). Shaded areas indicate increased synergistic (blue) or

antagonistic (red) effects for BTK, IRAK4 and BTK/IRAK4 inhibitor-dependent genes

compared to all genes. **g**, Normalised expression of dynamic BCR-dependent genes MTOR

and **h**, CD81, and **i**, mTOR-dependent gene *CSTB* divided by trisomy 12 status in DMSO or

drug-treated samples. **j**, Normalised expression of MEK-induced trisomy 12 signature gene

CD19 divided by trisomy 12 status in DMSO or MEK inhibitor-treated samples. Significance

levels were calculated using two-tailed Student's *t*-test and shown as nominal *P*-values.

872 Suppl. Fig. 8: Interaction of perturbation effects and disease drivers

a, Pathway connectivity network of all DGI genes shown as a bipartite graph. Genes, drugs

and disease drivers are connected if a differential drug response was observed in the presence

of the mutation (*FDR* < 10%). **b**, Cluster assignment of DGIs based on gene expression in the

presence of either drug, mutation/CNV or both compared to WT/DMSO samples by Fuzzy C-

Means Clustering. The thick black line depicts the median LFC of all genes from the respective

cluster. **c**, Heatmap of relative gene expression of DGIs in the presence of either drug,

mutation/CNV or both compared to WT/DMSO samples by epistasis class. **d**, Number of target

genes with significant DGIs for each perturbation, and all recurrent drivers with more than 5

cases by epistasis class (*FDR* < 10%).

Suppl. Fig. 9: IRAK4 is a disease driver targeted by trisomy 12

a, Dose-response curves of 20 patients treated with IRAK4 inhibitor and IRAK4 inhibitor

combined with 1 µg/mL CpG-ODN for TLR9 stimulation. **b**, Gene expression of genes on the

amplified locus chr12:42,538,673-chr12: 44,229,999 (hg19) by trisomy 12 status for the CLL

cohort (n = 210) from Lütge et al.[13]. Significance levels were calculated using two-tailed

Student's *t*-test and shown as nominal *P*-values.

Supplementary methods figures

Suppl. Methods Fig. 1: Design and pilot drug screening using shallow QuantSeq

a, Dose-response curves of drug response in CLL patients (n = 88) (viability drug screen

from Lu et al.[12]). **b**, Significant genetic association with drug response in CLL samples (*FDR*

< 10%) for aberration with more than 5 cases. Significance levels were calculated using two-

tailed Student's *t*-test and adjusted for multiple testing. **c**, Overview of treatments and

concentrations. **d**, Number of DEGs between 4 drug- and DMSO-treated CLL samples. **e**,

Volcano plot summarising the differential gene expression after PI3K, BTK, combined

BTK/IRAK4, MEK, MDM2 and XPO1 inhibitor treatment. **f**, Gene expression of exemplary

genes with treatment-specific down- and upregulation. **g**, GSEA with DEGs after each

treatment (all genes) using hallmark gene sets. Gene sets were filtered by *FDR* (< 1%) and

number of enriched genes (> 10). **h**, Pathway connectivity network shown as a bipartite

graph. Genes and drugs are connected if the drug perturbation leads to differential

expression (10% *FDR*).

Suppl. Methods Fig. 2 Gating strategy for flow cytometric assessment of cell viability

Cells were pre-gated by excluding debris and doublet cells. Viable and dead cells were then

gated based on cell size (FSC-A) and granularity (SSC-A).

Suppl. Methods Fig. 3 Guided sparse factor analysis model evaluation

a, Proportion of variation explained by each GSFA model with 1 to 30 factors. **b**, Percentage

of additional variation explained by each GSFA model compared to the previous model with

the highest explained variation. The model with 20 factors was selected for further analysis.