GenomoBase: an integrated resource for *Genomoviridae* genomics and metagenomics

Yair Cárdenas-Conejo[1,2,3]*

1 Laboratorio de Biología Sintética, Estructural y Molecular, Universidad de Colima,
Coquimatlán, Colima, México.

2 Secretaría de Ciencia, Humanidades, Tecnología e Innovación, Gobierno de México,
Ciudad de México, México

3 Centro Universitario de Investigaciones Biomédicas, Universidad de Colima. Colima,
Colima, México

Corresponding Author:

Yair Cárdenas-Conejo*

ORCID: 0000-0002-0190-244X

email address: ycardenas11@ucol.mx

Abstract

The family *Genomoviridae* (phylum Cressdnaviricota) comprises viruses with circular

single-stranded DNA (ssDNA) genomes that typically encode a replication protein and a

capsid protein. High-throughput sequencing has greatly expanded the number of known

genomoviruses, but sequences and annotations remain dispersed across multiple

repositories and publications, and no family-focused resource has been available. Here I

present GenomoBase (https://www.genomobase.org), a curated web resource that integrates

genomic, ecological and bibliographic data for all 237 *Genomoviridae* species currently

recognized by the International Committee on Taxonomy of Viruses (ICTV). GenomoBase

leverages a PostgreSQL backend to organize data according to standardized taxonomy,

summarizing viral occurrences across ten genera, twelve host categories and nineteen

countries. An "Explore Metagenomes" module incorporates more than 600 Sequence Read

Archive (SRA) runs identified via Serratus, linking taxonomic records to metagenomic

detections and highlighting the global distribution of genomovirus-related reads. To support

research, the platform also provides automated literature updates via PubMed and local

BLAST services for nucleotide and protein queries. GenomoBase offers a user-friendly

entry point for exploring *Genomoviridae* diversity and is designed to facilitate taxonomic

curation, ecological and evolutionary analyses, and the discovery of novel genomoviruses

in both environmental and host-associated niches.

Keywords

Genomoviridae; CRESS-DNA viruses; metagenomics; viral database; Cressdnaviricota;

Introduction

Circular replication-associated protein encoding single-stranded (CRESS) DNA viruses are ubiquitous components of environmental and host-associated viromes and are currently classified within the phylum Cressdnaviricota. Members of this phylum possess small circular ssDNA genomes that replicate by rolling-circle replication mediated by a conserved replication-associated protein (Rep), a hallmark of all CRESS-DNA viruses, and they infect a broad range of eukaryotic hosts, including plants, fungi, invertebrates and vertebrates [1]. The best-known families include *Geminiviridae*, which infect mono- and dicotyledonous plants; *Circoviridae*, which infect pigs, poultry and other vertebrates; and *Nanoviridae*, which infect mainly crop plants [1]. More recently, within Cressdnaviricota, the family *Genomoviridae* was established following the discovery of *Sclerotinia sclerotiorum* hypovirulence-associated DNA virus 1 (SsHADV-1) [2], a circular ssDNA virus infecting the plant-pathogenic fungus *Sclerotinia sclerotiorum* [3]. Subsequent taxonomic work has expanded *Genomoviridae* to ten genera and more than two hundred species recognized by the International Committee on Taxonomy of Viruses (ICTV) [4]. Genomoviruses typically have compact genomes of approximately 1.9–3.9 kb that encode two major proteins in an ambisense arrangement: a rolling-circle replication initiation protein (Rep) and a capsid protein (CP). The Rep protein is homologous to the corresponding Rep of members of the family *Geminiviridae*, sharing conserved domains and sequence motifs required for viral replication [2]. The origin of replication is also similar in both families, consistent with replication via rolling-circle mechanisms [2]. In contrast, the CP forms a non-enveloped, isometric T=1 icosahedral capsid and shows no recognizable amino acid sequence similarity to capsid proteins of other known virus groups. Some members additionally encode small hypothetical open reading frames of unknown function. To date, no canonical movement protein has been identified in

genomoviruses, consistent with the etymology of the family name *Genomoviridae* (Ge- for

"geminivirus-like", nomo- for "no movement protein") [2]. Although genomoviruses were

initially described in fungi, subsequent metagenomic studies have detected them in a wide

range of contexts, including plants, birds, mammals, invertebrates and diverse

environmental matrices such as wastewater, sediments, soils and food-associated

microbiomes. Experimental evidence of infection remains limited to *Sclerotinia*

sclerotiorum [3] and successful replication in the sciarid fly *Lycoriella ingenua* [5], and the

true host range of many lineages remains unclear.

The rapid expansion of genomovirus sequence data has been driven almost entirely by

metagenomic studies of human, animal and environmental viromes and by large-scale

surveillance projects. Gemykibiviruses and related taxa have now been reported in a variety

of human clinical samples, including cerebrospinal fluid from acute encephalitis patients

[6], vaginal secretions [7] and blood donors in the Brazilian Amazon [8]; the same virus has

also been identified in bats [9], suggesting a possible zoonotic link. More recently, a human

gemykibivirus was shown to actively replicate in HEK293T cells [10], further underscoring

the need to better understand the biology and potential medical relevance of these viruses.

Despite the growing interest, genomovirus sequences and annotations remain dispersed

across taxonomic resources, GenBank records and individual metagenomic studies, and no

dedicated, curated database has been available for this rapidly expanding family. To

address this gap, I developed GenomoBase (https://www.genomobase.org), an online

resource focused specifically on the family *Genomoviridae*. GenomoBase integrates ICTV-

based taxonomy with curated genome annotations, standardized host and geographic

metadata, Serratus-filtered metagenomic detections from the Sequence Read Archive

(SRA), local BLAST services and a PubMed-derived literature catalogue. Here we describe

the construction, content and main features of GenomoBase and provide a concise

overview of the patterns that can be readily explored through the portal.

Materials and methods

Reference taxonomy and genome retrieval

ICTV-recognized genomovirus species were retrieved from the Virus Metadata Resource

(VMR) file VMR_MSL40.v2.20251013 (https://ictv.global/vmr), which lists all species

and genera formally assigned to the family *Genomoviridae*. From this file we extracted the

scientific name and the corresponding GenBank accession number for each species. The

curated table underlying GenomoBase contains 239 genome records: one representative

genome for each of the 237 species, plus the three genomic segments (A, B and C) of the

tripartite *Gemytripvirus fugra1*. Using a custom Python script (fetch_genomes.py) built on

Biopython v1.86 [11], we downloaded the corresponding GenBank records via NCBI E-

utilities and parsed the following information: accession, genus, species name, virus name,

sequence definition line, genome length, full genomic sequence and all annotated coding

sequences (CDSs). For each genome we identified the canonical Rep and CP ORFs based

on gene annotations and product descriptions, extracted their nucleotide and amino acid

sequences and stored them as separate fields. Any additional predicted CDS were

concatenated into "other proteins" fields (names, amino acid sequences and nucleotide

sequences).

The resulting dataset, including host and geographic information, was manually inspected

to ensure that all entries had complete circular genomes and coherent annotation of Rep and

CP. Where necessary, minor inconsistencies in the genus or species names were corrected

to match the latest ICTV taxonomy. The final curated table (**Supplementary Table 1**) was

used as the master source for populating the GenomoBase database.

Host, isolation source and country information were also extracted from the GenBank

records. For each genome we collected the host, isolation_source, country and

collection_date qualifiers when present. Host names were standardized and manually

grouped into 11 ecological categories, including mammals, birds, plants, insects, fungi,

molluscs, reptiles, arachnids, actinopterygian fishes, humans and environmental samples.

This manual curation step allowed us to separate human-associated records from other

mammalian hosts and to distinguish non-living matrices (e.g. wastewater or environmental

swabs) from host organisms.

Country strings were cleaned to remove subnational descriptors and harmonized to ISO-

like country names (e.g. "USA: California" to "USA"). When possible, additional locality

information (regions, cities or sampling sites) was kept as free-text descriptors in the

database. These standardized fields are used in the statistics and visualization pages of the

portal to summarize host and geographic patterns.

Metagenomic survey and SRA/BioSample integration

Because most genomoviruses have been discovered by metagenomics, we implemented an

"Explore Metagenomes" module to connect reference genomes with public high-

throughput sequencing projects. As a starting point we used Serratus, a large-scale

alignment platform for viral sequence discovery [12], to identify SRA runs with reads

matching genomovirus references. We queried the Serratus explorer for alignments against

genomovirus targets and retained runs whose best hits satisfied the specific filters on

alignment quality: genome alignment scores and percent identity between 75% and 100%

(query date November 2025). This range was chosen to capture both close relatives and

more divergent genomovirus-like sequences while excluding low-similarity matches that

are likely spurious.

We extracted the unique run identifiers and obtained 613 distinct SRA runs with putative

genomovirus signal. To enrich these with standardized metadata we developed a custom

Python script (fetch_sra_metadata.py) that queries the NCBI SRA and BioSample API for

NCBI-hosted runs and the ENA browser API for European accessions (e.g. SAMEA and

ERS identifiers). For each run, the script retrieves: SRA accession, BioSample accession,

release date, SRA "organism" or "environment" label, BioSample "host" and "isolation

source" attributes, when available location string and, when possible, latitude/longitude

coordinate. All retrieved fields are cleaned to remove control characters and harmonized

into a JSON structure with one dictionary per SRA run. For BioSamples hosted at ENA

(typically SAMEA, ERS or ERP accessions), the script parses XML records to extract host

and geographic information, including coordinates embedded in free-text fields. The final

JSON file (**Supplementary file 1**) is both downloadable from the portal and used internally

to populate the metagenomes table in PostgreSQL. A small fraction of runs lack either

explicit location names or coordinates; in these cases, only the available fields are stored.

The Explore Metagenomes web page uses these curated attributes to generate summary

plots and an interactive map of sampling locations.

Literature retrieval and curation

To provide an up-to-date overview of genomovirus research, we implemented a literature module driven by a dedicated PubMed query. A custom Python script (fetch_pubmed.py) uses NCBI E-utilities to search PubMed for articles containing the terms "Genomoviridae", "genomovirus" and the names of each of the ten ICTV-recognized genera in titles, abstracts or Medical Subject Headings terms. The script retrieves the PubMed IDs of all matching records, then uses esummary to obtain titles, journal names, publication years and DOIs, and efetch to access additional metadata where necessary. Each record is annotated with one or more broad categories (e.g. "environmental metagenomics", "animal viromes", "human-associated viromes", "fungal genomoviruses", "reviews") based on keyword matching in titles and abstracts. The output is stored in a JSON file, which is rendered on the Literature page as an interactive, searchable table with filters by category, year and journal.

BLAST services

To enable sequence similarity searches, GenomoBase provides local BLASTN and BLASTP services based on NCBI BLAST+ v2.17.0 [13]. From the curated genomovirus table we generated FASTA files for complete genomes, Rep proteins and CP proteins, which were converted into dedicated BLAST databases. User-submitted nucleotide or protein queries are run against these databases with a default parameters and E-value threshold of 1×10^{-3}, and the results are returned as sortable tables with basic alignment statistics and direct links from hit accessions to the corresponding GenomoBase virus detail pages.

Database implementation, BLAST services and web interface

GenomoBase is implemented using the Django web framework (Python v3.10+) with a

PostgreSQL backend. A primary database model stores curated genome-level information,

including accession, taxonomy, host and geographic metadata, as well as the nucleotide and

amino acid sequences of Rep, CP and additional CDS. Separate tables store metagenomic

run metadata and literature entries. The application is deployed on an Amazon Web

Services (AWS) EC2 instance, which provides a stable, scalable environment with regular

backups and straightforward future upgrades of both the database and the web front-end.

The web interface is styled using Bootstrap and custom CSS, with dedicated views for the

index page, master species list, species detail, statistics, BLAST search, metagenome

explorer, downloads, literature and external resources. Summary statistics (e.g. numbers of

species per genus, host category counts and country distributions) are computed

dynamically using Django's ORM and rendered as charts (bar plots and maps) using

JavaScript-based visualizations.

Results

Resource content and example analyses

GenomoBase organizes the currently ICTV-recognized *Genomoviridae* diversity into ten

genera, with *Gemycircularvirus* (126 species) and *Gemykibivirus* (50 species) clearly

dominating the dataset, and the remaining genera represented by between 1 and 16 species

each (Fig. 1a). Genome lengths range from ~1.99 to 3.9 kb (mean ≈ 2.19 kb), consistent

with compact CRESS-DNA genomes (Fig. 1b). The upper bound of ~3.9 kb corresponds to

the combined size of the three genomic segments of the tripartite *Gemytripvirus fugra1*.

Beyond taxonomic composition and genome size, host annotations reveal associations with

12 ecological categories, including mammals, birds, plants, insects, fungi, molluscs,

reptiles, arachnids, actinopterygian fishes and non-living matrices. Mammals represent the

largest category (37.66%, excluding humans) in terms of species–host combinations,

followed by plants (14.64%), birds (13.39%) and insects (10.88%) (Fig. 1c). A discrete

subset of seven genomoviruses belonging to *Gemykibivirus* (5 spp.), *Gemykrogvirus* and

Gemyvongvirus genera has been recovered from human-associated samples, highlighting a

small but potentially important group of lineages at the interface between environmental

and medical virology. Geographic metadata indicate that genomoviruses have been

reported from at least ~19 countries, with especially high counts from the USA (97 spp.),

New Zealand (38 spp.), China (26 spp.) and Brazil (21), illustrating their broad global

distribution (Fig. 1d).

Figure 1. Overview of *Genomoviridae* species and associated metadata in GenomoBase. (a) Bar chart showing the distribution of currently classified species across ICTV-recognized genera. (b) Genome length distribution visualized as a raincloud plot combining a density curve, individual genomes plotted as points, and a boxplot. (c) Broad host categories for genomoviruses, summarized as a pie chart based on curated host metadata. (d) Global map indicating the number of genomovirus species reported per country according to available sampling location information.

Based on the harmonized host annotations for the 613 *Genomoviridae*-positive SRA runs, most detections fall into a set of recurrent host categories (Fig. 2a). The largest single class corresponds to "environmental samples", which account for 172/613 runs (28.1%) and encompass marine and freshwater systems, wastewater, air, soils and food-associated microbiomes. Among runs with biological hosts, human-associated samples (human gut, general human metagenomes, oral/skin/vaginal metagenomes) constitute the most frequent category, with 98 runs (16.0%), followed by metagenomes from birds (85 runs, 13.9%). Additional mammalian hosts include giant pandas (*Ailuropoda melanoleuca*; 27 runs, 4.4%), domestic pigs (*Sus scrofa*; 23 runs, 3.8%), mice (*Mus musculus*; 21 runs, 3.4%), bovines (*Bos* spp.; 17 runs, 2.8%) and *Macaca mulatta* (7 runs, 1.1%). There are also smaller contributions from *Xenopus laevis* (11 runs, 1.8%) and samples with unknown host information (50 runs, 8.2%). Beyond these dominant categories, the remaining runs form a long tail of individually represented wildlife and domestic animal hosts. Collectively, these patterns indicate that genomovirus-related reads are detected across a broad spectrum of vertebrate hosts and environmental matrices, while a substantial fraction of SRA runs still lacks precise host information.

Geographic metadata indicate that 462 of 613 runs (75.3%) include a textual location descriptor, corresponding to 180 distinct place names distributed across 47 countries or territories. In addition, 282 runs (46.0%) carry explicit latitude/longitude coordinates,

enabling map-based visualization of sampling sites. Within this set, the largest

contributions come from projects in China (approximately 165 runs) and the USA

(approximately 96 runs), followed by individual efforts in India, Ireland, Singapore,

Norway, South Korea, Canada and several European and Latin American countries (Fig.

2b).

Figure 2. Overview of metagenomic runs with genomovirus detections in GenomoBase. (a) Pie chart
summarizing SRA runs with genomovirus hits, grouped into broad host or sample categories (e.g. birds,
humans, domestic pigs, environmental and others). (b) Global heat map of sampling locations for the 282 runs

with latitude/longitude metadata, illustrating the relative density of genomovirus-positive metagenomes.

Warmer colours indicate higher local density across countries and regions.

Discussion

The rapid expansion of genomovirus sequence data, largely driven by metagenomic studies,

has created a challenge for data integration and standardization. Until now, genomovirus

sequences and annotations have been scattered across general taxonomic resources,

GenBank (https://www.ncbi.nlm.nih.gov/datasets/taxonomy/1910928/), ViralZone

(https://viralzone.expasy.org/7382), ICTV (VMR_MSL40.v2.20251013) and individual

metagenomic surveys, without a family-focused curation framework. GenomoBase

addresses this gap by providing, to our knowledge, the first integrated resource that aligns

Genomoviridae genomes with the official ICTV taxonomy, harmonized host and

geographic metadata and systematically filtered metagenomic detections. By centralizing

these data within a dedicated web interface, the platform facilitates taxonomic curation and

enables ecological and evolutionary analyses that are difficult to perform from isolated

records.

The aggregated view provided by GenomoBase makes current sampling biases in

Genomoviridae immediately apparent. Geographic metadata for formally described species

indicate detections in 19 countries across all inhabited continents, while the metagenomic

module reveals SRA runs potentially containing genomoviruses from around 47 countries

worldwide. This pattern likely reflects sequencing effort rather than the true biogeography

of the family, even as it clearly indicates that members of the family *Genomoviridae* are

widespread across continents and environments. GenomoBase also highlights a broad

apparent host spectrum in both the species and metagenomic datasets, including mammals,

birds, invertebrates, plants and environmental samples; although infection has not been

experimentally confirmed for all of these hosts [4], many of these associations remain

plausible and merit further investigation. Of particular interest, seven formally recognized

species belonging to three genera have been reported in human clinical samples [6,8], and

genomoviruses are detected in at least ~90 human-associated metagenomes, including gut

[14], vaginal [7] and brain [15]. These observations suggest that genomovirus are recurrent

components of the human virome and underline the value of continued surveillance of this

family in a public-health context, especially in light of recent evidence that at least one

gemykibivirus can actively replicate in human cells [10].

A second key contribution of GenomoBase is that it does not merely summarize existing

detections, but explicitly links ICTV-aligned reference genomes with Serratus-derived

metagenomic reads and contigs. Connecting these "orphan" SRA sequences to curated

Genomoviridae genomes shows that members of this family are ubiquitous components of

environmental and host-associated viromes, often detected in studies that were not designed

to target them. Making such hidden detections visible and searchable within a family-

specific context allows researchers to identify under-sampled hosts and environments, and

to prioritize public SRA runs for targeted re-analysis or assembly of novel genomovirus

genomes. In this sense, GenomoBase functions not only as a static catalogue but also as a

hypothesis-generating platform for future work on genomovirus diversity, ecology and

evolution.

Conclusions and future directions

GenomoBase provides a focused, user-friendly platform for the *Genomoviridae* research

community and for researchers interested more broadly in circular ssDNA viruses. By

integrating ICTV-based reference genomes, harmonized host and geographic metadata,

Serratus-filtered SRA detections, local BLAST services and an automatically updated

literature catalogue, the resource enables users to (i) search and download curated

genomovirus genomes and genes, (ii) explore host and geographic patterns of diversity, and

(iii) investigate the occurrence of genomoviruses across public metagenomic datasets. At

the same time, the current implementation makes explicit several limitations of the

available data, such as incomplete host annotations and strong geographic sampling biases,

which can guide future sequencing surveys. GenomoBase is freely available and will be

periodically updated in line with ICTV releases and Serratus-based screenings, and its

underlying framework can be readily extended to other CRESS-DNA virus families and

related viral taxa.

Data and code availability

The online instance of GenomoBase is accessible at https://www.genomobase.org. All

curated genomovirus genome metadata, host and geographic annotations, and the

metagenomic summary used in this study are available for download through the

GenomoBase portal and as archived files on Zenodo (DOI: 10.5281/zenodo.17860246).

The complete species table and metagenome JSON, together with all custom scripts and the

Django source code used to build and update GenomoBase, as well as sample input data

and the corresponding output, are deposited on Zenodo and documented in detail in the

accompanying README file, which provides additional implementation details.

Funding

320   This study was supported by grants from the Secretaría de Ciencia, Humanidades,

321   Tecnología e Innovación (CBF-2025-G-902).

Author contributions

Y.C.-C. was responsible for conceptualization, data curation, software and web

implementation (GenomoBase), formal analysis, visualization and writing of the

manuscript.

Conflict of interest statement

The author declares no competing interests.

References

[1]   M. Krupovic, A. Varsani, D. Kazlauskas, M. Breitbart, E. Delwart, K. Rosario, N. Yutin, Y.I. Wolf, B. Harrach, F.M. Zerbini, V.V. Dolja, J.H. Kuhn, E.V. Koonin, *Cressdnaviricota* : a Virus Phylum Unifying Seven Families of Rep-Encoding Viruses with Single-Stranded, Circular DNA Genomes, J Virol 94 (2020) e00582-20. https://doi.org/10.1128/JVI.00582-20.

[2]   M. Krupovic, S.A. Ghabrial, D. Jiang, A. Varsani, Genomoviridae: a new family of widespread single-stranded DNA viruses, Arch Virol 161 (2016) 2633–2643. https://doi.org/10.1007/s00705-016-2943-3.

[3]   X. Yu, B. Li, Y. Fu, D. Jiang, S.A. Ghabrial, G. Li, Y. Peng, J. Xie, J. Cheng, J. Huang, X. Yi, A geminivirus-related DNA mycovirus that confers hypovirulence to a plant pathogenic fungus, Proc. Natl. Acad. Sci. U.S.A. 107 (2010) 8387–8392. https://doi.org/10.1073/pnas.0913535107.

[4]   A. Varsani, M. Krupovic, Family Genomoviridae: 2021 taxonomy update, Arch Virol 166 (2021) 2911–2926. https://doi.org/10.1007/s00705-021-05183-y.

[5]   S. Liu, J. Xie, J. Cheng, B. Li, T. Chen, Y. Fu, G. Li, M. Wang, H. Jin, H. Wan, D. Jiang, Fungal DNA virus infects a mycophagous insect and utilizes it as a transmission vector, Proc. Natl. Acad. Sci. U.S.A. 113 (2016) 12803–12808. https://doi.org/10.1073/pnas.1608013113.

[6]   E.T. Tuladhar, S. Shrestha, S. Vernon, L. Droit, K.A. Mihindukulasuriya, M. Tamang, L. Karki, A. Elong Ngono, B. Jha, B.K. Awal, B.S. Chalise, R. Jha, S. Shresta, D. Wang, K.D. Manandhar, Gemykibivirus detection in acute encephalitis patients from Nepal, mSphere 9 (2024) e00219-24. https://doi.org/10.1128/msphere.00219-24.

[7]   E.D.S.F. Ramos, T.R. Tozetto-Mendoza, P. Bortoletto, N.E. Ferreira, L. Honorato, E.M.G. Barbosa, A. Luchs, I.M. Linhares, S.D. Spandorfer, E. Leal, A.C. Da Costa, S.S. Witkin, M.C. Mendes-Correa, Characterization of CRESS-DNA viruses in human vaginal secretions: An exploratory metagenomic investigation, Journal of Medical Virology 96 (2024) e29750. https://doi.org/10.1002/jmv.29750.

[8]   R.S. Bezerra, H.T. Bitencourt, D.T. Covas, S. Kashima, S.N. Slavov, Metagenomic identification of human Gemykibivirus-2 (HuGkV-2) in parenterally infected blood donors from the Brazilian Amazon, International Journal of Infectious Diseases 98 (2020) 249–251. https://doi.org/10.1016/j.ijid.2020.06.096.

[9]   B.S. Silvério, R.L. Sanz Duro, L.L.F. De Sousa, M.D. Guilardi, G. Cabral-Miranda, L.M.R. Janini, R. Durães-Carvalho, Detection and Phylogenetic Analysis of Emerging Human-Associated Gemykibivirus-2 in *Molossus molossus* Bat From Brazil, Journal of Medical Virology 97 (2025) e70154. https://doi.org/10.1002/jmv.70154.

[10] R. Yang, H. Yan, Y. Wang, W. Yang, J. Qin, Transcriptome Analysis Reveals Gemykibivirus Infection Induces Mitochondrial DNA Release in HEK293T Cells, Viruses 17 (2025) 1331. https://doi.org/10.3390/v17101331.

[11] P.J.A. Cock, T. Antao, J.T. Chang, B.A. Chapman, C.J. Cox, A. Dalke, I. Friedberg, T. Hamelryck, F. Kauff, B. Wilczynski, M.J.L. De Hoon, Biopython: freely available Python tools for computational molecular biology and bioinformatics, Bioinformatics 25 (2009) 1422–1423. https://doi.org/10.1093/bioinformatics/btp163.

[12] R.C. Edgar, B. Taylor, V. Lin, T. Altman, P. Barbera, D. Meleshko, D. Lohr, G. Novakovsky, B.
Buchfink, B. Al-Shayeb, J.F. Banfield, M. De La Peña, A. Korobeynikov, R. Chikhi, A. Babaian,
Petabase-scale sequence alignment catalyses viral discovery, Nature 602 (2022) 142–147.
https://doi.org/10.1038/s41586-021-04332-2.

[13] C. Camacho, G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, T.L. Madden, BLAST+:
architecture and applications, BMC Bioinformatics 10 (2009) 421. https://doi.org/10.1186/1471-2105-
10-421.

[14] T.K. Dao, T.T.N. Pham, H.D. Nguyen, Q.T. Dam, T.B.T. Phung, T.V.H. Nguyen, T.Q. Nguyen, K.C.
Hoang, T.H. Do, Metagenomic Analysis of the Gastrointestinal Phageome and Incorporated Dysbiosis
in Children with Persistent Diarrhea of Unknown Etiology in Vietnam, Pathogens 14 (2025) 985.
https://doi.org/10.3390/pathogens14100985.

[15] I. Lamberto, K. Gunst, H. Müller, H. Zur Hausen, E.-M. De Villiers, Mycovirus-Like DNA Virus
Sequences from Cattle Serum and Human Brain and Serum Samples from Multiple Sclerosis Patients,
Genome Announc 2 (2014) e00848-14. https://doi.org/10.1128/genomeA.00848-14.

GenomoBase: an integrated resource for Genomoviridae genomics and metagenomics

Global detections from
ICTV Species and metagenomes

237 Genomoviridae spp.
from 10 genera

Gemykolovirus

Gemycircularvirus

Gemykibivirus

GenBank/SRA inputs

Curated genomovirus
genomes and SRA Runs

GenomoBase
implementation