1 Transcriptional Activation of Estrogen Receptor-alpha and Estrogen Receptor-beta

2 from Elephant Shark (*Callorhynchus milii*)

4 Ya Ao[1], Haruka Narita[1], Wataru Takagi[2], Susumu Hyodo[2],

5 Michael E. Baker[4,5], Yoshinao Katsu[1,3]

[1]Graduate School of Life Science, Hokkaido University, Sapporo 060-0810, Japan

[2]Laboratory of Physiology, Atmosphere and Ocean Research Institute, The University of Tokyo,

Chiba 277-8564, Japan

[3]Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan

[4]Division of Nephrology-Hypertension, Department of Medicine, 0693, University of California,

San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0693

[5]Center for Academic Research and Training in Anthropogeny (CARTA), University of Califor-

nia, San Diego, La Jolla, CA 92093

Correspondence:

Yoshinao Katsu: ykatsu@sci.hokudai.ac.jp

Michael E. Baker: mbaker@health.ucsd.edu

[Type here]

Abstract

Humans and other vertebrates contain two estrogen receptors (ERs), ERα and ERβ.   Among

cartilaginous fish (sharks, rays, skates), which are chondrichthyans that evolved about 425 mil-

lion years ago, only activation by steroids of ERβ orthologs has been characterized.   To remedy

this gap in understanding estrogen signaling in chondrichthyans, we studied estrogen activation

of orthologs of human ERα and ERβ from elephant shark (*Callorhynchus milii*).   Unexpectedly,

we found that *C. milii* contained three estrogen-responsive ERα genes: ERα1 (596 amino acids),

ERα2 (600 amino acids), and ERα3 (599 amino acids) with strong sequence similarity to each

other.   We also found an estrogen-unresponsive gene, ERα4 (561 amino acids), with a 39 amino

acid deletion in the DNA-binding domain.   An estrogen-responsive ERβ ortholog (580 amino

acids) also was present in *C. milii*.   The three active *C. milii* ERαs are of similar length to hu-

man ERα (595 amino acids); however, *C. milii* ERβ is longer than human ERβ (530 amino ac-

ids).   We studied transcriptional activation of ERα and ERβ by estradiol (E2), the main repro-

ductive estrogen in humans.   We also studied estrone (E1), the main postmenopausal estrogen,

and estriol (E3), which is synthesized during pregnancy.   We determined the half-maximal re-

sponse (EC50) and fold-activation to E2, E1, and   E3 of *C. milii* ERα1, ERα2, ERα3, and ERβ.

Among these estrogens, E2 had the lowest EC50 for all four ERs.   Fold-activation by E2 and

E3 was similar for ERα1, ERα2, ERα3, and ERβ.   Overall, estrogen activation of *C. milii* ERα

and ERβ was similar to that for human ERα and ERβ, indicating substantial conservation of the

vertebrate ER during the 425 million years since the divergence of cartilaginous fish and humans

from a common ancestor.

Keywords: elephant shark; estrogen receptor evolution; ERα; ERβ.

1. Introduction

Estrogens are a class of steroid hormones that have diverse physiological activities in

females and males in humans and other vertebrates [1-5].   Estrogens act by binding to estrogen

receptors (ERs) [5-6], which are transcription factors that belong to the nuclear receptor family,

which also includes receptors for progestins, androgens, and corticosteroids [7-9].   To date, two

distinct estrogen receptor gene: estrogen receptor alpha (ERα) and estrogen receptor beta (ERβ),

respectively, have been isolated in mammals and other vertebrates [10-12].   ER α and ERβ

have strong sequence similarity (97%) in their DNA binding domains, but only about 55% simi-

larity in their ligand-binding domains.

ERα is expressed in reproductive tissues (uterus, ovary), breast, kidney and bone,

while ERβ is expressed in the ovary and male <u>reproductive organs</u>, (prostate), colon, kidney and

the immune system.   The response of ERα and ERβ in humans and other vertebrates to

17β-estradiol (E2), the main physiological estrogen, as well as to estrone (E1) a postmenopausal

estrogen, and estriol (E3), which is synthesized during pregnancy (Figure 1) has been studied

extensively [5,6,10,13].   Although ERα and ERβ can have opposing actions, both receptors are

activated by binding estrogens, which induce a conformational change that promotes binding to

specific DNA sequences and the initiation of estrogen-dependent gene transcription.   ERα pro-

motes tissue proliferation, while ERβ may act as a suppressor of ERα-mediated proliferation.

75
Figure 1. Structures of estrogen.
Estrone (E1), 17β-estradiol (E2), and estriol (E3) are the natural estrogens.   E1 is a minor fe-
male sex hormone and serves mainly as a precursor of E2.   E3 is produced in the placenta dur-
ing pregnancy.   E2 is a major female sex hormone mainly produced in the ovary [38, 39].

Although ERs with sequence similarity to ERα [14,15] and ERβ [14-16] have been

isolated from elasmobranchs (sharks, rays, and skates), only the response to steroids by ERβ from

the cloudy catshark (*Scyliorhinus torazame*) and whale shark (*Rhincodon typus*) has been studied

[16].   Estrogen activation of a gnathostome ERα has not been reported.   Elasmobranchs, along

with Holocephali, comprise the two extant subclasses of Chondrichthyes, jawed fishes with skel-

etons composed of cartilage rather than calcified bone [17].   Chondrichthyes occupy a key posi-

tion relative to fish and terrestrial vertebrates, having diverged from bony vertebrates about 450

million years ago [14,18].   As the oldest group of extant jawed vertebrates (gnathostomes), the

properties of ERα and ERβ in cartilaginous fishes can provide important insights into the early evo-

lution of ERα and ERβ from an ancestral ER in lamprey [19].   Moreover, elephant shark has the

slowest evolving genome of all known vertebrates, including the slowly evolving coelacanth [14],

which makes elephant shark genes a good system to study the evolution of the ER, including the

early evolution of selectivity of the ER in a vertebrate and the subsequent divergence in steroid
selectivity in human ER, in other vertebrate ERs and in ray-finned fish ERs [20].   Another point
of interest for the ER in elephant shark is that elephant sharks contain three full-length ERαs [14],
raising the question of each how these three ERαs compare with each other and with mammalian
ERαs in their response to estrogens.

To remedy this gap in our understanding of ERα activation in vertebrates, we have
studied the response to estradiol, estrone, and estriol of three full-length *Callorhinchus milii*
ERαs, as well as *C. milii* ERβ.   Here we report that despite the strong sequence similarity
among *C. milii* ERα1, ERα2, ERα3, and ERβ, there are some differences in their half-maximal
response (EC50) and fold-activation to these three physiological estrogens.   We find that E2 had
the lowest EC50 for all four ERs.   E2 and E3 had a similar fold activation for ERα1, ERα2,
ERα3, and ERβ.   Overall, estrogen activation of *C. milii* ERα and ERβ was similar to that for
human ERα and ERβ [5,7], indicating substantial conservation of the vertebrate ER in the 425
million years since the divergence of cartilaginous fish and humans from a common ancestor.
106

2. Materials and methods
2.1. Animals
C. milii elephant sharks of both sexes were collected in Western Port Bay, Victoria, Aus-
tralia, using recreational fishing equipment, and transported to Primary Industries Research Vic-
toria, Queenscliff, using a fish transporter.   The animals were kept in a 10-t round tank with
running seawater (SW) under a natural photoperiod for at least several days before sampling.
Animals were anesthetized in 0.1% (w/v) 3-amino benzoic acid ethyl ester (Sigma-Aldrich, St.
Louis, MO, USA).   After the decapitation of the animal, tissues were dissected out, quickly
frozen in liquid nitrogen, and kept at −80°C.   All animal experiments were conducted according
to the Guideline for Care and Use of Animals approved by the committees of University of To-
kyo as described in [21] [Animal Ethics Committee of the Ocean Research Institute of the Uni-
versity of Tokyo], Approval Code: [17-3], and Approval Date: [March 29, 2005]).
119

120 *2.2. Chemical reagents*
121   Estrone (E1), 17β-estradiol (E2), and estriol (E3) were purchased from Sigma-Aldrich
122 Corp. (St. Louis, MO).   All chemicals were dissolved in dimethyl-sulfoxide (DMSO).   The

123   concentration of DMSO in the culture medium did not exceed 0.1%.

125   *2.3. Cloning of estrogen receptors*

126   When we began this project, the genome of *C. milii* had not yet been sequenced [14].

Thus, we needed to sequence an ER in *C. milii* as a first step in determining the number of ERs

and their sequences in *C. milii*.   For this goal, we used two conserved amino acid regions in the

DNA-binding domain (GYHYGVW) and the ligand-binding domain (NKGM/IEH) of vertebrate

ERs to design the degenerate oligonucleotides to clone *C. milii* ERs.   The second PCR used the

first PCR amplicon, and nested primers that were selected in the DNA binding domain

(CEGCKAF) and the ligand binding domain (NKGM/IEH).   As a template for PCR, the

first-strand cDNA was synthesized using total RNA isolated from the ovary.   The amplified

DNA fragments were subcloned with TA-cloning plasmids pCR2.1-TOPO (Invitrogen, Carlsbad,

CA).   The 5' and 3' ends of the ER cDNAs were amplified by rapid amplification of the cDNA

end (RACE) using a SMART RACE cDNA Amplification kit (BD Biosciences Clontech., Palo

Alto, CA).   To amplify the isoforms of elephant shark ER, we applied a PCR-based cDNA am-

plification technique using the primers et (5'-TGAAGTGTGATCGTCCAGGCGACAG-3' and

5'-CAAGCTGGAGGATAAGACATCGAC -3').   Sequencing was performed using a BigDye

Terminator Cycle Sequencing kit and analyzed on the Applied Biosystems 3730 DNA Analyzer.

141   We used the *C. milii* extracted RNA to clone ERα1, ERα2, ERα3, ERα4, and ERβ,

142   which had sequences corresponding to the sequences of the *C. milii* genome deposited by

143   Venkatesh et al., 2014 [14].

145   *2.4. Database and sequence analyses*

146   All sequences generated were searched for similarity using BLASTn and BLASTp at

the web servers of the National Center of Biotechnology Information (NCBI).   We found four

ERα genes and one ERβ gene [14], which were aligned using ClustalW [22].

2.5. Phylogenetic tree analysis

The evolutionary history was inferred using the Neighbor-Joining method [23].   The optimal

tree is presented in this study.   The percentage of replicate trees in which the associated taxa

clustered together in the bootstrap test (1000 replicates) is shown next to the branches [24].

The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary dis-

tances used to infer the phylogenetic tree.   The evolutionary distances were computed using the

Poisson correction method and are in units of the number of amino acid substitutions per site.

Evolutionary analyses were conducted in MEGA 11 [25].

2.6. Reporter gene assay

Full-length estrogen receptors were amplified using specific forward and reverse primers

designed at start and stop codons and cloned into the mammalian expression vector pcDNA3.1

(Invitrogen).   In addition, ERs with a FLAG-tag added to the N-terminus were also prepared by

PCR.   A reporter construct, pGL4.23-4xERE was produced by subcloning of oligonucleotides

having 4xERE into the *Kpn*I-*Hind*III site of pGL4.23 vector (Promega).   All cloned DNA se-

quences were verified by sequencing.

Reporter gene assays using full-length ERs were performed in Human Embryonic Kidney

293 cells (HEK293 cells).   HEK293 cells were seeded in 24-well plates at 5 x 10^4 cells/well in

phenol-red-free Dulbecco's modified Eagle's medium with 10% charcoal/dextran-treated fetal

bovine serum.   After 24 h, the cells were transfected with 400 ng of reporter construct, 25 ng of

pRL-TK (as an internal control to normalize the variation in transfection efficiency; contains the

Renilla reniformis luciferase gene with the herpes simplex virus thymidine kinase promoter), and

200 ng of pcDNA3.1-estrogen receptor using polyethylenimine (PEI).   After 5 h of incubation,

ligands were applied to the medium at various concentrations.   After an additional 43 h, the

cells were collected, and the luciferase activity of the cells was measured with the Du-

al-Luciferase Reporter Assay System.   Promoter activity was calculated as firefly (*P.*

pyralis)-luciferase activity/sea pansy (*R. reniformis*)-luciferase activity.   The values shown are

mean ± SEM from three separate experiments, and dose-response data, which were used to cal-

culate the half maximal response (EC50) for each steroid, were analyzed using GraphPad Prism

(Graph Pad Software, Inc., San Diego, CA).   All experiments were performed in triplicate.

3. Results

3.1. Cloning and sequence analysis of cDNAs for four elephant shark ERα genes and one ERβ

gene.

Using standard RT-PCR and RACE techniques, we successfully cloned four full-length

ERs, designated as ERα1, ERα2, ERα3 (Figure 2A, B), and ERβ (Figure 2C), from elephant

shark ovary RNA.   The cDNA for elephant shark ERα1 predicted a 596 amino acid protein with

a calculated molecular mass of 66.6 kDa (GenBank accession no. LC068847), and the cDNA for

elephant shark ERβ predicted a 580 amino acid protein with a calculated molecular mass of 64.9

kDa (GenBank accession no. LC068848), Figure 2A, B.   In addition, we cloned the other two

full-length elephant shark ERs: ERα2 and ERα3, which are found in GenBank (GenBank acces-

sion no. XM_007894403 for ERα2, XM_007894404 for ERα3).   ERα4 (GenBank accession no.

XM_007894406) with a 39 amino acid deletion in the DNA-binding domain was also cloned

(Figure 2A, B).

The elephant shark ERα and ERβ sequences contain the five recognizable steroid

hormone receptor sub-domains in the expected order: the N-terminal region (A/B domain), DNA

binding domain, DBD (C domain), hinge region (D domain), ligand binding domain, LBD (E

domain), and C-terminal extension (F domain) (Figure 2A).   In Figure 2A we compare the

functional domains of ERα1, ERα2, ERα3 and ERα4.   Figure 2A shows that there is excellent

conservation among these ERs of the A/B, C, D, E and F domains.   This strong sequence con-

servation also is seen in the amino acid alignment of ERα1, ERα2, ERα3 and ERα4 shown in

Figure 2B.   Figure 2B shows the deletion in the DBD of ERα4, which probably explains the

lack of transcriptional activation by estrogens of *C. milii* ERα4.   The amino acid sequence of

ERα4 is closest to ERα3.   Overall, Figures 2A and 2B show the strong sequence conservation

in the four *C. milii* ERα genes.

In Figure 2C, we show a sequence alignment of *C. milii* ERα1 and ERβ.   The DNA

response element recognition motif (P-box, CEGCKA) in the DBD and the AF-2 motif (core

LLLEML region) in the LBD, which is required for interaction with transcriptional co-activators,

are conserved in all four -elephant shark ERα sequences and in ERβ (Figure 2B and 2C).

217
Figure 2. Amino acid sequences of the elephant shark ERs.
(A) Comparison of functional domains of four elephant shark ERα genes.
Elephant Shark ERα1, ERα2, and ERα3 show strong sequence conservation of their A/B, C, D,
and E domains.   ERα4 has a deletion in the DNA-binding domain [C domain], which we pro-
pose explains its lack of transcriptional activation by estradiol.
(B) Sequence alignment of Elephant Shark ERα1, ERα2, ERα3, and ERα4.   There is excellent
conservation of the four-elephant shark ERα genes.
(C) Aligned sequences of the elephant shark ERα1 and ERβ.   The DBD and LBD are indicated
by open boxes.   Residues are important for DNA response element recognition (the P-box), and
the AF-2 region, which mediates contact of the LBD with transcriptional coactivators, is shaded
in gray.   Numbers to the right indicate amino acid position.   Asterisks indicate residues con-
served in both ERα1 and ERβ.

3.2. Comparison of functional domains on human ERα and ERβ with functional domains in ele-

phant shark ERα and ERβ.

To gain additional insights into the evolution of the *C. milii* ERs, we compared the

functional domains of elephant shark ERα1to elephant shark ERβ (Figure 3A), of elephant shark

ERα1 to human ERα and human ERβ (Figure 3C), and of elephant shark ERβ to human ERα

and human ERβ (Figure 3D).   We also compared human ERα and ERβ to each other (Figure

3B).

238
Figure 3. Comparison of the functional domains of elephant shark ERs with human ERs.
The functional domains A/B through F are shown schematically with the number of amino acid residues
indicated.   The percentage of amino acid identity is shown.   (A) Comparison of elephant shark ERα1
with elephant shark ERβ. The N-terminal (A/B) domains of elephant shark ERα1 and ERβ show strong
divergence.   There is also some divergence between the ligand (E) binding domain on elephant shark
ERα1 and ERβ.   (B) Comparison of human ERα (accession no. NM000125) with human ERβ (acces-
sion no. AB006590).   The domains of human ERα with human ERβ have similar sequence similarity to
each other as elephant shark ERα1 and ERβ.   (C) Comparison of elephant shark ERα1 with human ERα
and human ERβ.   Elephant shark ERα1 is closer to human ERα than to human ERβ.   (D) Comparison
of elephant shark ERβ with human ERα and human ERβ.   Elephant shark ERβ is closer to human ERβ
than to human ERα.

Figure 3A shows the excellent conservation of the DNA-binding domains [C domains]

(94% identity) and ligand-binding domains [E domains] (66% identity) in elephant shark ERα1

and ERβ, in contrast to the weak conservation of the A/B domains (13% identity) at their

N-terminus.   Comparison of Figures 3A and B reveals that the DBD in human ERα and *C. milii*

ERα1 and ERβ have similar sequence conservation, while the LBD sequence in *C. milii* ERα1

and *C. milii* ERβ are more conserved (66% identity) than the LBD in human ERα and ERβ (58%

identity).

Figure 3C shows the strong sequence conservation of the DBD in elephant shark ERα1

and human ERα and ERβ.   Figure 3C also shows that there is stronger conservation (76%) of

the LBD in elephant shark ERα1 and human ERα, compared to the similarity (64%) of the cor-

responding LBD in elephant shark ERα1 to human ERβ.   Figure 3D shows the strong sequence

conservation of the DBD in elephant shark ERα1 and human ERα and ERβ.   Figure 3D also

shows the stronger conservation (72%) of the LBD in elephant shark ERβ and human ERβ,

compared to the similarity (64%) to the corresponding LBD in human ERα.

3.3. Comparison of functional domains on elephant shark ERα and ERβ with functional domains

in zebrafish, and whale shark.

To gain an additional insight into the evolution of elephant shark ERα and ERβ, we

compared their sequences to ERα and ERβ sequences in zebrafish and whale shark (Figure 4).

270
Figure 4.   Comparison of the functional domains of elephant shark ERs with ERs from zebrafish
and whale shark.
The functional domains A/B through F are shown schematically with the number of amino acid residues
indicated.   The percentage of amino acid identity is shown.   (A) Comparison of elephant shark ERα1

with zebrafish and whale shark ERs.   (B) Comparison of elephant shark ERβ with zebrafish and whale
shark ERs.   Genbank accession no. NM_152959 (zebrafish ERα), AF516874 (zebrafish ERβ1),
AF349413 (zebrafish ERβ2), XM_048600891 (whale shark ERα), and AB551716 (whale shark ERβ).
278

The sequence comparisons (Figure 4) between elephant shark ERα and ERβ domains

with ERα and ERβ domains in whale shark revealed an unexpected sequence similarity (87%

identity) between the A/B domain on elephant shark ERα and whale shark ERα.   The A/B do-

main in whale shark ERβ and elephant shark ERβ has 40% identity.   Interestingly, the A/B do-

mains on elephant shark ERα and ERβ domains have 39% and 35% sequence identity, respec-

tively, with the A/B domains on human ERα and human ERβ, respectively.

The stronger conservation of the LBD in elephant shark ERβ and human ERβ (72%

identity), compared to the similarity to the corresponding LBD in human ERα (63% identity) is

shown in Figure 3C.   The LBD of elephant shark ERα1 is closer to the LBD in human ERα

than the LBD in human ERβ, and the LBD of elephant shark ERβ is closer to the LBD in human

ERβ than the LBD in human ERα (Figure 3C).

3.4. Phylogenetic analysis of estrogen receptors in elephant sharks, humans and other verte-

brates.

To better understand the relationship between elephant shark ERs and other vertebrates

ERs, we constructed a phylogeny of ERs from sharks, humans and other vertebrates [Figure 5]

using the Neighbor-Joining method [23].   The phylogeny in shows the early divergence of shark

ERs from other ERs.

Figure 5. Phylogenetic analysis of estrogen receptors in elephant sharks, humans and other verte-
brates.
The phylogenetic tree was constructed using the maximum likelihood analysis [24]. The two elephant
shark ERs are marked with open boxes. The scale bar indicates 0.1 expected amino acid substitutions
per site. The GenBank accession numbers are NM_000125 (human ERα), AB006590 (human ERβ),
NM_203535 (tropical clawed frog ERα), NM_001040012 (tropical clawed frog ERβ), XM_048600891
(whale shark ERα), AB551716 (whale shark ERβ), XM_068044449 (horn shark ERα), XM_068038484
(horn shark ERβ), XM_060831156 (epaulette shark ERα), XM_060829508 (epaulette shark ERβ),
NM_152959 (zebrafish ERα), AF516874 (zebrafish ERβ1), AF349413 (zebrafish ERβ2).

3.5. Transcriptional activities of elephant shark ERα and ERβ.

A transactivation assay was used to examine the response to the physiological estrogens

E1, E2, and E3 of the three active isoforms of elephant shark ERα. The three isoforms, ERα

showed ligand-dependent transactivation (Figure 6A, B, C). The EC50s for transcriptional ac-

tivation of ERα1, ERα2, and ERα3 were 0.49 nM, 0.29 nM, and 0.25 nM (E1), 0.0059 nM,

0.016 nM, and 0.0093 nM (E2), and 0.2 nM, 0.31 nM, and 0.29 nM (E3), respectively (Table 1).

We also examined the induction of elephant shark ERβ transcriptional activity by dif-

ferent concentrations of estrogens (E1, E2, and E3) (Figure 6). All three estrogens activated

transcription of elephant shark ERβ in a dose-dependent manner (Figure 6D). The

half-maximal response (EC50) for transcriptional activation of elephant shark ERβ was 0.14 nM

for E1, 0.001 nM for E2, and 0.02 nM for E3 (Table 1). Although there was no significant dif-

ference in EC50 values for activation by an estrogen of ERα1 and ERβ, the fold activation val-

ues for E2 were higher than those for E1 and E2 (Table 1). Thus, the elephant shark estrogen

receptors exhibited a ligand responsiveness observed human estrogen receptors [1-10].

[Type here]

324
325 **Figure 6.   Transcriptional activation of elephant shark ERs.**
326 Elephant shark ERs were transfected into HEK293 cells with an ERE-driven reporter gene.   Concentra-
327 tion-response profile for ERα1 (A), ERα2 (B), ERα3 (C), and ERβ (D) for E1, E2, and E3 (10^{-13}M to
328 10^{-6}M).   Data are expressed as a ratio of a vehicle (DMSO) to other test chemicals.   Each point repre-
329 sents the mean of triplicate determinations, and vertical bars represent the mean ± SEM.

[Type here]

Table 1. Gene transcriptional activities of elephant shark ERs by estrogens in HEK293

		E1	E2	E3
ERα1	EC50 (nM)	0.49	0.0059	0.2
	Fold-Activation (± SEM) *	1.62 (± 0.16)	1.66 (± 0.04)	1.57 (± 0.09)
	Ratio**	0.98	1.00	0.95
ERα2	EC50 (nM)	0.29	0.016	0.31
	Fold-Activation (± SEM) *	1.51 (± 0.12)	1.79 (± 0.12)	1.82 (± 0.13)
	Ratio**	0.84	1.00	1.02
ERα3	EC50 (nM)	0.25	0.0093	0.29
	Fold-Activation (± SEM) *	1.55 (± 0.11)	1.75 (± 0.08)	1.87 (± 0.15)
	Ratio**	0.89	1.00	1.07
ERβ	EC50 (nM)	0.14	0.001	0.02
	Fold-Activation (± SEM) *	2.87 (± 0.16)	3.23 (± 0.26)	3.49 (± 0.26)
	Ratio**	0.89	1.00	1.08

*Fold activation values were calculated based on the 10 nM values.
**Ratio values were calculated by dividing the fold-activation value of estrogens by the value of E2.

4. Discussion

Significant advances have been made in elucidating the evolution of estrogen signaling in animals [1, 2, 9, 11, 19, 26, 27]. Amphioxus, is a chordate that contains an ER and a steroid receptor (SR) that is an ortholog of receptors for 3-keto-steroids, including testosterone, proges- terone, cortisol, and aldosterone [11, 28]. Contrary to expectations, the amphioxus ER does not bind E2 or other steroids [11, 27, 28], while E2 and E1 are transcriptional activators of the SR [11, 28]. Atlantic sea lamprey contains an ER [19] that is activated by E2 and E1 [11, 27].

The estrogen receptors in Chondrichthyes, cartilaginous fishes with jaws, are basal to lobe-finned fishes and are ancestors of terrestrial vertebrates [14]. These ERs are still not fully characterized [15, 29]. The identification of ERα1, ERα2, ERα3, ERα4, and ERβ from the sequencing of the elephant shark [14] provided an opportunity to investigate estrogen signaling in a basal jawless vertebrate that evolved about 425 million years ago. We investigated the re- sponse to E1, E2, and E3 of ERα1, ERα2, ERα3, and ERβ from elephant shark expressed in HEK293 cells (Figure 5). These studies revealed that elephant shark ERα1, ERα2, ERα3, and

ERβ are activated by E1, E2, and E3.   We note that we found that ERα4 did not respond to an
estrogen (data not shown), which we propose is due to a sequence deletion in the DBD.   Thus,
we did not continue studies of estrogen activation of ERα4.
Our data with estrogen activation of ERα1, ERα2, ERα3, and ERβ align with the established
ligand binding characteristics of ERs in other vertebrates, thereby substantiating the conserved
nature of estrogen signaling pathways [28, 30-32].   Estrogen-dependent transcriptional activity
of the ERs from various species, including other sharks, revealed that the ERs from all species
have a stronger response to E2 than to E3 [16, 33-37].
Instead of only one ERα gene, elephant shark contains three ERα genes with very similar
sequences, which is unexpected.   The strong sequence conservation of ERα4, which is not lig-
and-activated, also is unexpected.   Further research is needed to gain a more comprehensive
understanding of the physiological function of estrogen and the multiple estrogen receptors in
elephant sharks.   This data should provide clues for understanding the physiological function(s)
of each isoform.
In summary, this is the first report with evidence for the conservation of elephant shark
ERα and ERβ compared to human ERα and ERβ, which provides a valuable perspective on the
early evolutionary history of these critical nuclear receptors in vertebrates.   The findings from
this study not only confirm the conserved nature of ER-mediated estrogen signaling across ver-
tebrates but also open new avenues for research into the functional diversification of these re-
ceptors across vertebrates. These findings furnish researchers with critical molecular data to ex-
amine the role of ERs in future studies, such as those examining gonadal development, reproduc-
tive biology, and developmental and evolutionary endocrinology.
394

Author Contribution
Y.A.: Writing – review & editing, Writing – original draft, Methodology, Investigation, Formal
analysis, and Data curation.   **H.N.:** Writing – original draft, Methodology, Investigation, Formal
analysis, Data curation.   **W. T.:** Resources**.   S.H.:** Resources**.   M.E.B.:** Writing – review &
editing, Project administration, Validation, Supervision, Conceptualization.   **Y.K.:** Writing –
review & editing, Project administration, Visualization, Validation, Funding acquisition, Super-
vision, Conceptualization.
402

403 **Funding**

404 This research was supported by Grants-in-Aid for Scientific Research from the Ministry of Edu-
405 cation, Culture, Sports, Science and Technology of Japan (23K05839) to Y.K., and the Takeda
406 Science Foundation to Y.K.

Declaration of competing interest

408
409 The authors declare that they have no known competing financial interests or personal relation-
410 ships that could have appeared to influence the work reported in this paper.

Data availability

412
413 All data is freely available on request to Dr. Katsu or to Dr. Baker.

Acknowledgment

415
416 The authors are grateful to Dr. Kaori Oka for her suggestions and helpful advice.   We also thank
417 colleagues in our laboratories.

5. References

419
[1] Bondesson M, Hao R, Lin CY, Williams C, Gustafsson JÅ. 2015. Estrogen receptor signaling
during vertebrate development. Biochim Biophys Acta 1849(2), 142-151. doi:
10.1016/j.bbagrm.2014.06.005.

[2] Gao H, Dahlman-Wright K. 2011. The gene regulatory networks controlled by estrogens. Mol
Cell Endocrinol. 334(1-2), 83-90. doi: 10.1016/j.mce.2010.09.002.

[3] Deroo BJ, Korach KS. Estrogen receptors and human disease. J Clin Invest. 2006
Mar;116(3):561-70. doi: 10.1172/JCI27987.

[4] Hamilton KJ, Hewitt SC, Arao Y & Korach KS 2017 Estrogen Hormone Biology. *Curr Top
Dev Biol* **125** 109-146.

[5] Paterni I, Granchi C, Katzenellenbogen JA & Minutolo F 2014 Estrogen receptors alpha
(ERalpha) and beta (ERbeta): subtype-selective ligands and clinical potential. *Steroids* **90**
13-29

[6] Heldring N, Pawson T, McDonnell D, Treuter E, Gustafsson JA, Pike AC. Structural insights
into corepressor recognition by antagonist-bound estrogen receptors. J Biol Chem. 2007
Apr 6;282(14):10449-55. doi: 10.1074/jbc.M611424200.

[7] Dahlman-Wright K, Cavailles V, Fuqua SA, Jordan VC, Katzenellenbogen JA, Korach KS,
Maggi A, Muramatsu M, Parker MG, Gustafsson JA. 2006. International Union of Phar-
macology. LXIV. Estrogen receptors. Pharmacol Rev. 58(4), 773-781. doi:
10.1124/pr.58.4.8.

[8] Markov GV, Tavares, R, Dauphin-Villemant C, Demeneix BA, Baker ME, Laudet V. 2009.
Independent elaboration of steroid hormone signaling pathways in metazoans. Proc Natl
Acad Sci USA. 106(29), 11913-11918. doi: 10.1073/pnas.0812138106.

[9] Baker ME. 2011. Origin and diversification of steroids: co-evolution of enzymes and nuclear
receptors. Mol Cell Endocrinol. 334(1-2), 14-20. doi: 10.1016/j.mce.2010.07.013.

[10] Kuiper GG, Enmark E, Pelto-Huikko M, Nilsson S, Gustafsson JA. Cloning of a novel re-
ceptor expressed in rat prostate and ovary. Proc Natl Acad Sci U S A. 1996 Jun
11;93(12):5925-30. doi: 10.1073/pnas.93.12.5925.

[11] Bridgham JT, Brown JE, Rodriguez-Mari A, Catchen JM, Thornton JW. 2008. Evolution of
a new function by degenerative mutation in cephalochordate steroid receptors. PLoS
Genet. 4(9), e1000191. doi: 10.1371/journal.pgen.1000191.

[12] Baker ME, Nelson DR, Studer RA. 2015. Origin of the response to adrenal and sex steroids:
Roles of promiscuity and co-evolution of enzymes and steroid receptors. J. Steroid
Biochem. Mol Biol 151, 12-24. doi: 10.1016/j.jsbmb.2014.10.020.

[13] Escande A, Servant N, Rabenoelina F, Auzou G, Kloosterboer H, Cavaillès V, Balaguer P,
Maudelonde T. 2009. Regulation of activities of steroid hormone receptors by tibolone
and its primary metabolites. J Steroid Biochem Mol Biol. 116(1-2):8-14. doi:

10.1016/j.jsbmb.2009.03.008.

[14] Venkatesh B, Lee AP, Ravi V, Maurya AK, Lian MM, Swann JB, Ohta Y, Flajnik MF,

Sutoh Y, Kasahara M, Hoon S, Gangu V, Roy SW, Irimia M, Korzh V, Kondrychyn I,

Lim ZW, Tay BH, Tohari S, Kong KW, Ho S, Lorente-Galdos B, Quilez J,

Marques-Bonet T, Raney BJ, Ingham PW, Tay A, Hillier LW, Minx P, Boehm T, Wilson

RK, Brenner S, Warren WC. 2014. Elephant shark genome provides unique insights into

gnathostome evolution. Nature 505(7482), 174-179. doi: 10.1038/nature12826.

[15] Filowitz GL, Rajakumar R, O'Shaughnessy KL, Cohn MJ. 2018. Cartilaginous Fishes Pro-

vide Insights into the Origin, Diversification, and Sexually Dimorphic Expression of Ver-

tebrate Estrogen Receptor Genes. Mol Biol Evol. 35(11), 2695-2701. doi:

10.1093/molbev/msy165.

[16] Katsu Y, Kohno S, Narita,H, Urushitani H, Yamane K, Hara A, Clauss TM, Walsh MT,

Miyagawa S, Guillette LJ, Iguchi T. 2010a. Cloning and functional characterization of

Chondrichtyes, cloudy catshark, Scyliorhinus torazame and whale shark, Rhincodon

typus estrogen receptors. Gen Comp Endocrinol. 168(3), 496-504. doi:

10.1016/j.ygcen.2010.06.010.

[17] Inoue JG, Miya M, Lam K, Tay BH, Danks JA, Bell J, Walker TI, Venkatesh B.

2010.Evolutionary origin and phylogeny of the modern holocephalans (Chondrichthyes:

Chimaeriformes): a mitogenomic perspective. Mol Biol Evol. 27(11):2576-2586. doi:

10.1093/molbev/msq147.

[18] Yu WP, Rajasegaran V, Yew K, Loh WL, Tay BH, Amemiya CT, Brenner S, Venkatesh B.

2008. Elephant shark sequence reveals unique insights into the evolutionary history of

verte-brate genes: A comparative analysis of the protocadherin cluster. Proc Natl Acad Sci

U S A. 105(10):3819-3824. doi: 10.1073/pnas.0800398105.

[19] Thornton JW. 2001. Evolution of vertebrate steroid receptors from an ancestral estrogen re-

ceptor by ligand exploitation and serial genome expansions. Proc Natl Acad Sci USA.

98(10), 5671-5676. doi: 10.1073/pnas.091553298.

[20] Baker ME. 2019. Steroid receptors and vertebrate evolution. Mol Cell Endocrinol.

496:110526. doi: 10.1016/j.mce.2019.110526.

[21] Kakumura K, Watanabe S, Bell JD, Donald JA, Toop T, Kaneko T, Hyodo S. 2009. Multi-

ple urea transporter proteins in the kidney of holocephalan elephant fish (*Callorhinchus*

milii). Comp Biochem Physiol B Biochem Mol Biol. 154(2), 239-247. doi:

10.1016/j.cbpb.2009.06.009.

[22] Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of

progressive multiple sequence alignment through sequence weighting, position-specific

gap penalties and weight matrix choice. Nucleic Acids Res. 22(22), 4673-4680. doi:

10.1093/nar/22.22.4673.

[23] Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phyloge-

netic trees. Mol Biol Evol. 1987 Jul;4(4):406-25. doi:

10.1093/oxfordjournals.molbev.a040454.

[24] Felsenstein J. CONFIDENCE LIMITS ON PHYLOGENIES: AN APPROACH USING

THE BOOTSTRAP. Evolution. 1985 Jul;39(4):783-791. doi:

10.1111/j.1558-5646.1985.tb00420.x.

[25] Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis Ver-

sion 11. Mol Biol Evol. 2021 Jun 25;38(7):3022-3027. doi: 10.1093/molbev/msab120.

[26] Cotnoir-White D, Laperriere D, Mader S. 2011. Evolution of the repertoire of nuclear

receptor binding sites in genomes. Mol Cell Endocrinol. 334(1-2), 76-82. doi:

10.1016/j.mce.2010.10.021.

[27] Paris M, Pettersson K, Schubert M, Bertrand S, Pongratz I, Escriva H, Laudet V. 2008. An

amphioxus orthologue of the estrogen receptor that does not bind estradiol: insights into

532   estrogen receptor evolution. BMC Evol Biol. 8, 219. doi: 10.1186/1471-2148-8-219.

534  [28] Katsu Y, Kubokawa K, Urushitani H, Iguchi T. 2010. Estrogen-dependent transactivation of

535   amphioxus steroid hormone receptor via both estrogen and androgen response elements.

536   Endocrinology 151(2), 639-648. doi: 10.1210/en.2009-0766.

538  [29] Awruch CA. 2013. Reproductive endocrinology in chondrichtyans: the present and the future.

539   Gen Comp Endocrinol. 192, 60-70. doi: 10.1016/j.ygcen.2013.05.021.

541  [30] Green S, Walter P, Kumar V, Krust A, Bornert JM, Argos P, Chambon P . 1986. Human oes-

542   trogen receptor cDNA: sequence, expression and homology to v-erb-A. Nature.

543   320(6058), 134-139. doi: 10.1038/320134a0.

545  [31] Matthews J, Gustafsson JA. 2003. Estrogen signaling: a subtle balance between ER alpha

546   and ER beta. Mol Interv. 3(5), 281-292. doi: 10.1124/mi.3.5.281.

548  [32] Katzenellenbogen BS, Katzenellenbogen JA. 2000. Estrogen receptor transcription and

549   transactivation: Estrogen receptor alpha and estrogen receptor beta: regulation by selec-

550   tive estrogen receptor modulators and importance in breast cancer. Breast Cancer Res.

551   2(5), 335-344. doi: 10.1186/bcr78.

553  [33] Katsu Y, Lange A, Urushitani H, Ichikawa R, Paul GC, Cahill LL, Jobling S, Tyler CR,

554   Iguchi T. 2007. Functional associations between two estrogen receptors, environmental

555   estrogens and sexual disruption in the roach (Rutilus rutilus). Environ Sci Tech. 41(9),

556   3368-3374. doi: 10.1021/es062797I

558  [34] Katsu Y, Kohno S, Hyodo S, Ijiri S, Adachi S, Hara A, Guillette LJ, Iguchi T. 2008. Molec-

559   ular cloning, characterization and evolutionary analysis of estrogen receptors from

560   phylogenetically ancient fish. Endocrinology 149(12), 6300-6310. doi:

561   10.1210/en.2008-0670

563  [35] Katsu Y, Taniguchi E, Urushitani H, Miyagawa S, Takase M, Kubokawa K, Tooi O, Oka T,

Santo N, Myburgh J, Matsuno A, Iguchi T. 2010c. Molecular cloning and characterization

of ligand- and species-specificity of amphibian estrogen receptors. Gen Comp Endocrinol.

168(2), 220-230. doi: 10.1016/j.ygcen.2010.01.002

[36] Lange A, Katsu Y, Miyagawa S, Ogino Y, Urushitani H, Kobayashi T, Hirai T, Shears JA,

Nagae M, Yamamoto J, Ohnishi Y, Oka T, Tatarazako N, Ohta Y, Tyler CR, Iguchi T.

2012. Comparative responsiveness to natural and synthetic estrogens of fish species

commonly used in the laboratory and field monitoring. Aquat Toxicol 109, 250-258. doi:

10.1016/j.aquatox.2011.09.004.

[37] Yatsu R, Katsu Y, Kohno S, Mizutani T, Ogino Y, Ohta Y, Myburgh J, van Wyk JH, Guillette

LJ Jr, Miyagawa S, Iguchi T. 2016. Characterization of evolutionary trend in squamate

estrogen receptor sensitivity. Gen Comp Endocrinol. 238, 88-95. doi:

10.1016/j.ygcen.2016.04.005.

[38] Kuhl H. Pharmacology of estrogens and progestogens: influence of different routes of ad-

min-istration. Climacteric. 2005 Aug;8 Suppl 1:3-63. doi: 10.1080/13697130500148875.

[39] Stanczyk FZ. 2024. Metabolism of endogenous and exogenous estrogens in women. J Ster-

oid Biochem Mol Biol. 2024 Sep;242:106539. doi: 10.1016/j.jsbmb.2024.106539.