1 Transcriptional Activation of Estrogen Receptor-alpha and Estrogen Receptor-beta
2 from Elephant Shark (*Callorhynchus milii*)
3

4 Ya Ao[1], Haruka Narita[1], Wataru Takagi[2], Susumu Hyodo[2],
5 Michael E. Baker[4,5], Yoshinao Katsu[1,3]
6
7
8
9
[1]Graduate School of Life Science, Hokkaido University, Sapporo 060-0810, Japan
[2]Laboratory of Physiology, Atmosphere and Ocean Research Institute, The University of Tokyo,
Chiba 277-8564, Japan
[3]Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan
[4]Division of Nephrology-Hypertension, Department of Medicine, 0693, University of California,
San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0693
[5]Center for Academic Research and Training in Anthropogeny (CARTA), University of Califor-
nia, San Diego, La Jolla, CA 92093
18
19 Correspondence:
20 Yoshinao Katsu: ykatsu@sci.hokudai.ac.jp
21 Michael E. Baker: mbaker@health.ucsd.edu

22
23 **Short title:** Transcriptional activation of elephant shark estrogen receptors

[Type here]

Abstract

Humans and other terrestrial vertebrates contain two estrogen receptors (ERs), ER-alpha and

ERbeta.   Among cartilaginous fish (sharks, rays, skates), which are jawless vertebrates that

evolved about 525 million years ago, only activation by steroids of ER-beta orthologs has been

characterized.   To remedy this gap in understanding estrogen signaling in jawless vertebrates,

we studied estrogen activation of orthologs of human ER-alpha and ER-beta.   Unexpectedly, we

found that *C. milii* contained three estrogen-responsive ER-alpha genes: ER-alpha1 (596 amino

acids), ER-alpha-2 (600 amino acids), and ER-alpha-3 (599 amino acids) with strong sequence

similarity to each other.   We also found an estrogen-unresponsive gene, ER-alpha4 (561 amino

acids), with a 39 amino acid deletion in the DNA-binding domain.   An estrogen-responsive

ER-beta ortholog (580 amino acids) also was present in *C. milii*.   The three active *C. milii*

ER-alphas have a similar length to human ER-alpha (595 amino acids); however, *C. milii*

ER-beta is longer than human ER-beta (530 amino acids).   We determined the half-maximal

response (EC50) and fold-activation to estradiol (E2), estrone (E1), and estriol (E3) of *C. milii*

ER-alpha1, ER-alpha2, ER-alpha3, and ER-beta.   Among these estrogens, E2 had the lowest

EC50 for all four ERs.   Fold-activation by E2 and E3 was similar for ER-alpha1, ER-alpha2,

ER-alpha3, and ER-beta.   Overall, estrogen activation of *C. milii* ER-alpha and ER-beta was

similar to that for human ER-alpha and ER-beta indicating substantial conservation of the verte-

brate ER in the 525 million years since the divergence of cartilaginous fish and humans from a

common ancestor.

Keywords: elephant shark; estrogen receptor evolution; ER-alpha; ER-beta.

1. Introduction

Estrogens are a class of steroid hormones that have diverse physiological activities in

females and males in humans and other vertebrates [1-5].   Estrogens act by binding to estrogen

receptors (ERs) [5-6], which are transcription factors that belong to the nuclear receptor family,

which also includes receptors for progestins, androgens, and corticosteroids [7-9].   To date, two

distinct estrogen receptors: estrogen receptor alpha (ERα) and estrogen receptor beta (ERβ), re-

spectively, have been isolated in mammals and other vertebrates [10-12].   The response of ERα

and ERβ in humans and other vertebrates to 17β-estradiol (E2), the main physiological estrogen,

as well as to estrone (E1) and estriol (E3) (Figure 1) has been studied extensively [5,6,10,13].

[Type here]

Figure 1.   Structures of estrogen.
Estrone (E1), 17β-estradiol (E2), and estriol (E3) are the natural estrogens.   E1 is a minor fe-
male sex hormone and serves mainly as a precursor of E2.   E3 is produced in the placenta dur-
ing pregnancy.   E2 is a major female sex hormone mainly produced in the ovary [35, 36].

Although ERs with sequence similarity to ERα [14,15] and ERβ [14-16] have been

isolated from elasmobranchs (sharks, rays, and skates), only the response to steroids by ERβ from

the cloudy catshark (*Scyliorhinus torazame*) and whale shark (*Rhincodon typus*) has been studied

[16].   Estrogen activation of a gnathostome ERα has not been reported.   Elasmobranchs, along

with Holocephali, comprise the two extant subclasses of Chondrichthyes, jawed fishes with skel-

etons composed of cartilage rather than calcified bone [17].   Chondrichthyes occupy a key posi-

tion relative to fish and terrestrial vertebrates, having diverged from bony vertebrates about 450

million years ago [14,18].   As the oldest group of extant jawed vertebrates (gnathostomes), the

properties of ERα and ERβ in cartilaginous fishes can provide important insights into the early evo-

lution of ERα and ERβ from an ancestral ER in lamprey [19].   Moreover, elephant shark has the

slowest evolving genome of all known vertebrates, including the slowly evolving coelacanth [14],

which makes elephant shark genes windows into the past, and, thus, a good system to study the

evolution of the ER, including selectivity of the ER in a basal vertebrate for steroids and the

subsequent divergence in steroid selectivity in human ER, in other terrestrial vertebrate ERs and in

ray-finned fish ERs [20].   Another point of interest for the ER in elephant shark is that elephant

sharks contain three full-length ERαs [14], raising the question of each how these three ERαs

compare with each other and with mammalian ERαs in their response to estrogens.

To remedy this gap in our understanding of ERα activation in a basal jawless verte-

brate, we have studied the response to estradiol, estrone, and estriol of three full-length

Callorhinchus milii ERαs, as well as *C. milii* ERβ.   Here we report that despite the strong se-

quence similarity among *C. milii* ERα1, ERα2, ERα3, and ERβ, there are some differences in

their half-maximal response (EC50) and fold-activation to these three physiological estrogens.

We find that E2 had the lowest EC50 for all four ERs.   E2 and E3 had a similar fold activation

for ERα1, ERα2, ERα3, and ERβ.   Overall, estrogen activation of *C. milii* ERα and ERβ was

similar to that for human ERα and ERβ [5,7], indicating substantial conservation of the verte-

brate ER in the 525 million years since the divergence of cartilaginous fish and humans from a

common ancestor.

2. Materials and methods

2.1. Animals

C. milii elephant sharks of both sexes were collected in Western Port Bay, Victoria, Aus-

tralia, using recreational fishing equipment, and transported to Primary Industries Research Vic-

toria, Queenscliff, using a fish transporter.   The animals were kept in a 10-t round tank with

running seawater (SW) under a natural photoperiod for at least several days before sampling.

Animals were anesthetized in 0.1% (w/v) 3-amino benzoic acid ethyl ester (Sigma-Aldrich, St.

Louis, MO, USA).   After the decapitation of the animal, tissues were dissected out, quickly

frozen in liquid nitrogen, and kept at −80°C.   All animal experiments were conducted according

to the Guideline for Care and Use of Animals approved by the committees of University of To-

kyo [21].

2.2. Chemical reagents

Estrone (E1), 17β-estradiol (E2), and estriol (E3) were purchased from Sigma-Aldrich

Corp. (St. Louis, MO).   All chemicals were dissolved in dimethyl-sulfoxide (DMSO).   The

concentration of DMSO in the culture medium did not exceed 0.1%.

2.3. Cloning of estrogen receptors

When we began this project, the genome of *C. milii* had not yet been sequenced [14].

Thus, we needed to sequence an ER in *C. milii* as a first step in determining the number of ERs

and their sequences in *C. milii*.   For this goal, we used two conserved amino acid regions in the

DNA-binding domain (GYHYGVW) and the ligand-binding domain (NKGM/IEH) of vertebrate

ERs to design the degenerate oligonucleotides to clone *C. milii* ERs.   The second PCR used the

first PCR amplicon, and nested primers that were selected in the DNA binding domain

(CEGCKAF) and the ligand binding domain (NKGM/IEH).   As a template for PCR, the

first-strand cDNA was synthesized using total RNA isolated from the ovary.   The amplified

DNA fragments were subcloned with TA-cloning plasmids pCR2.1-TOPO (Invitrogen, Carlsbad,

CA).   The 5' and 3' ends of the ER cDNAs were amplified by rapid amplification of the cDNA

end (RACE) using a SMART RACE cDNA Amplification kit (BD Biosciences Clontech., Palo

Alto, CA).   To amplify the isoforms of elephant shark ER, we applied a PCR-based cDNA am-

plification technique using the primers et (5'-TGAAGTGTGATCGTCCAGGCGACAG-3' and

5'-CAAGCTGGAGGATAAGACATCGAC -3').   Sequencing was performed using a BigDye

Terminator Cycle Sequencing kit and analyzed on the Applied Biosystems 3730 DNA Analyzer.

We used the *C. milii* extracted RNA to clone ERα1, ERα2, ERα3, ERα4, and ERβ,

which had sequences corresponding to the sequences of the *C. milii* genome deposited by

Venkatesh et al., 2014 [14].

2.4. Database and sequence analysis

All sequences generated were searched for similarity using BLASTn and BLASTp at

the web servers of the National Center of Biotechnology Information (NCBI).   We found four

ERα genes and one ERβ gene [14], which were aligned using ClustalW [22].

2.5. Reporter gene assay

Full-length estrogen receptors were amplified using specific forward and reverse primers

designed at start and stop codons and cloned into the mammalian expression vector pcDNA3.1

(Invitrogen).   In addition, ERs with a FLAG-tag added to the N-terminus were also prepared by

PCR.   A reporter construct, pGL4.23-4xERE was produced by subcloning of oligonucleotides

having 4xERE into the *Kpn*I-*Hind*III site of pGL4.23 vector (Promega).   All cloned DNA se-

quences were verified by sequencing.

Reporter gene assays using full-length ERs were performed in Human Embryonic Kidney

293 cells (HEK293 cells).   HEK293 cells were seeded in 24-well plates at 5×10^4 cells/well in

phenol-red-free Dulbecco's modified Eagle's medium with 10% charcoal/dextran-treated fetal

bovine serum.   After 24 h, the cells were transfected with 400 ng of reporter construct, 25 ng of

pRL-TK (as an internal control to normalize the variation in transfection efficiency; contains the

Renilla reniformis luciferase gene with the herpes simplex virus thymidine kinase promoter), and

200 ng of pcDNA3.1-estrogen receptor using polyethylenimine (PEI).   After 5 h of incubation,

ligands were applied to the medium at various concentrations.   After an additional 43 h, the

cells were collected, and the luciferase activity of the cells was measured with the Du-

al-Luciferase Reporter Assay System.   Promoter activity was calculated as firefly (*P.*

pyralis)-luciferase activity/sea pansy (*R. reniformis*)-luciferase activity.   The values shown are

mean ± SEM from three separate experiments, and dose-response data, which were used to cal-

culate the half maximal response (EC50) for each steroid, were analyzed using GraphPad Prism

(Graph Pad Software, Inc., San Diego, CA).   All experiments were performed in triplicate.

3. Results

3.1. Cloning and sequence analysis of cDNAs for four elephant shark ERα genes and one ERβ

gene.

Using standard RT-PCR and RACE techniques, we successfully cloned four full-length

ERs, designated as ERα1, ERα2, ERα3 (Figure 2A, B), and ERβ (Figure 2C), from elephant

shark ovary RNA.   The cDNA for elephant shark ERα1 predicted a 596 amino acid protein with

a calculated molecular mass of 66.6 kDa (GenBank accession no. LC068847), and the cDNA for

elephant shark ERβ predicted a 580 amino acid protein with a calculated molecular mass of 64.9

kDa (GenBank accession no. LC068848), Figure 2A, B.   In addition, we cloned the other two

full-length elephant shark ERs: ERα2 and ERα3, which are found in GenBank (GenBank acces-

sion no. XM_007894403 for ERα2, XM_007894404 for ERα3).   ERα4 (GenBank accession no.

XM_007894406) with a 39 amino acid deletion in the DNA-binding domain was also cloned

(Figure 2A, B).

The elephant shark ERα and ERβ sequences contain the five recognizable steroid

hormone receptor sub-domains in the expected order: the N-terminal region (A/B domain), DNA

binding domain, DBD (C domain), hinge region (D domain), ligand binding domain, LBD (E

domain), and C-terminal extension (F domain) (Figure 2A).   In Figure 2A we compare the

functional domains of ERα1, ERα2, ERα3 and ERα4.   Figure 2A shows that there is excellent

conservation among these ERs of the A/B, C, D, E and F domains.   This strong sequence con-

servation also is seen in the amino acid alignment of ERα1, ERα2, ERα3 and ERα4 shown in

Figure 2B.   Figure 2B shows the deletion in the DBD of ERα4, which probably explains the

lack of binding of estrogens by *C. milii* ERα4.   The amino acid sequence of ERα4 is closest to

ERα3.   Overall, Figures 2A and 2B show the strong sequence conservation in the four *C. milii*

ERα genes.

In Figure 2C, we show a sequence alignment of *C. milii* ERα1 and ERβ.   The DNA

response element recognition motif (P-box, CEGCKA) in the DBD and the AF-2 motif (core

LLLEML region) in the LBD, which is required for interaction with transcriptional co-activators,

are conserved in all four -elephant shark ERα sequences and in ERβ (Figure 2B and 2C).

Figure 2.   Amino acid sequences of the elephant shark ERs.
(A) Comparison of functional domains of four elephant shark ERα genes.
Elephant Shark ERα1, ERα2, and ERα3 show strong sequence conservation of their A/B, C, D,
and E domains.   ERα4 has a deletion in the DNA-binding domain [C domain], which explains
its lack of transcriptional activation by estradiol.   (B) Sequence alignment of Elephant Shark
ERα1, ERα2, ERα3, and ERα4.   There is excellent conservation of the four-elephant shark
ERα genes.   (C) Aligned sequences of the elephant shark ERα1 and ERβ.   The DBD and LBD
are indicated by open boxes.   Residues are important for DNA response element recognition
(the P-box), and the AF-2 region, which mediates contact of the LBD with transcriptional
coactivators, is shaded gray.   Numbers to the right indicate amino acid position.   Asterisks in-
dicate residues conserved in both ERα1 and ERβ.

3.2 Comparison of functional domains on human ERα and ERβ with functional domains in ele-

phant shark ERα and ERβ.

204   To gain additional insights into the evolution of the *C. milii* ERs, we compared the

205   functional domains of elephant shark ERα1 to elephant shark ERβ (Figure 3A), of elephant shark

206   ERα1 to human ERα and human ERβ (Figure 3C), and of elephant shark ERβ to human ERα

207   and human ERβ (Figure 3D).   We also compared human ERα and ERβ to each other (Figure

208   3B).

209
Figure 3.   Comparison of the functional domains of elephant shark ERs with human ERs.
The functional domains A/B through F are shown schematically with the number of amino acid residues
indicated.   The percentage of amino acid identity is shown.   (A) Comparison of elephant shark ERα1
with elephant shark ERβ. The N-terminal (A/B) domains of elephant shark ERα1 and ERβ show strong
divergence.   There is also some divergence between the ligand (E) binding domain on elephant shark
ERα1 and ERβ.   (B) Comparison of human ERα (accession no. NM000125) with human ERβ (acces-
sion no. AB006590).   The domains of human ERα with human ERβ have similar sequence similarity to
each other as elephant shark ERα1 and ERβ.   (C) Comparison of elephant shark ERα1 with human ERα
and human ERβ.   Elephant shark ERα1 is closer to human ERα than to human ERβ.   (D) Comparison
of elephant shark ERβ with human ERα and human ERβ.   Elephant shark ERβ is closer to human ERβ
than to human ERα.

222   Figure 3A shows the excellent conservation of the DNA-binding domains [C domains]

223   (94% identity) and ligand-binding domains [E domains] (66% identity) in elephant shark ERα1

[Type here]

and ERβ, in contrast to the weak conservation of the A/B domains (13% identity) at their

N-terminus.   Comparison of Figures 3A and B reveals that the DBD in human ERα and *C. milii*

ERα1 and ERβ have similar sequence conservation, while the LBD sequence in *C. milii* ERα1

and *C. milii* ERβ are more conserved (66% identity) than the LBD in human ERα and ERβ (58%

identity).

Figure 3C shows the strong sequence conservation of the DBD in elephant shark ERα1

and human ERα and ERβ.   Figure 3C also shows that there is stronger conservation (76%) of

the LBD in elephant shark ERα1 and human ERα, compared to the similarity (64%) of the cor-

responding LBD in elephant shark ERα1 to human ERβ.   Figure 3D shows the strong sequence

conservation of the DBD in elephant shark ERα1 and human ERα and ERβ.   Figure 3D also

shows the stronger conservation (72%) of the LBD in elephant shark ERβ and human ERβ,

compared to the similarity (64%) to the corresponding LBD in human ERα.

3.3 Comparison of functional domains on elephant shark ERα and ERβ with functional domains

in zebrafish, and whale shark.

To gain an additional insight into the evolution of elephant shark ERα and ERβ, we

compared their sequences to ERα and ERβ sequences in zebrafish and whale shark (Figure 4).

241
Figure 4.   Comparison of the functional domains of elephant shark ERs with ERs from zebrafish
and whale shark.
The functional domains A/B through F are shown schematically with the number of amino acid residues
indicated.   The percentage of amino acid identity is shown.   (A) Comparison of elephant shark ERα1
with zebrafish and whale shark ERs.   (B) Comparison of elephant shark ERβ with zebrafish and whale
shark ERs.   Genbank accession no. NM_152959 (zebrafish ERα), AF516874 (zebrafish ERβ1),
AF349413 (zebrafish ERβ2), XM_048600891 (whale shark ERα), and AB551716 (whale shark ERβ).

The sequence comparisons between elephant shark ERα and ERβ domains with ERα

and ERβ domains in whale shark revealed an unexpected sequence similarity (87% identity) be-

tween the A/B domain on elephant shark ERα and whale shark ERα.   The A/B domain in whale

shark ERβ and elephant shark ERβ has 40% identity.   Interestingly, the A/B domains on ele-

phant shark ERα and ERβ domains have 39% and 35% sequence identity, respectively, with the

A/B domains on human ERα and human ERβ, respectively.

Figure 3C shows the stronger conservation of the LBD in elephant shark ERβ and hu-

man ERβ (72% identity), compared to the similarity to the corresponding LBD in human ERα

(63% identity).   The LBD of elephant shark ERα1 is closer to the LBD in human ERα than the

LBD in human ERβ, and the LBD of elephant shark ERβ is closer to the LBD in human ERβ

than the LBD in human ERα.

260
261   **Figure 5.   Transcriptional activation of elephant shark ERs.**
262   Elephant shark ERs were transfected into HEK293 cells with an ERE-driven reporter gene.   Concentra-
263   tion-response profile for ERα1 (A), ERα2 (B), ERα3 (C), and ERβ (D) for E1, E2, and E3 (10^{-13}M to
264   10^{-6}M).   Data are expressed as a ratio of a vehicle (DMSO) to other test chemicals.   Each point repre-
265   sents the mean of triplicate determinations, and vertical bars represent the mean ± SEM.
266

3.4. Transcriptional activities of elephant shark ERα and ERβ.

A transactivation assay was used to examine the response to the physiological estrogens

E1, E2, and E3 of the three active isoforms of elephant shark ERα.   The three isoforms, ERα

showed ligand-dependent transactivation (Figure 5A, B, C).   The EC50s for transcriptional ac-

tivation of ERα1, ERα2, and ERα3 were 0.49 nM, 0.29 nM, and 0.25 nM (E1), 0.0059 nM,

0.016 nM, and 0.0093 nM (E2), and 0.2 nM, 0.31 nM, and 0.29 nM (E3), respectively (Table 1).

We also examined the induction of elephant shark ERβ transcriptional activity by differ-

ent concentrations of estrogens (E1, E2, and E3).   All three estrogens activated transcription of

elephant shark ERβ in a dose-dependent manner (Figure 5D).   The half-maximal response

(EC50) for transcriptional activation of elephant shark ERβ was 0.14 nM for E1, 0.001 nM for

E2, and 0.02 nM for E3 (Table 1).   Although there was no significant difference in EC50 values

for activation by an estrogen of ERα1 and ERβ, the fold activation values for E2 were higher

than those for E1 and E2 (Table 1).   Thus, the elephant shark estrogen receptors exhibited a

ligand responsiveness observed human estrogen receptors [1-10].

281
282
283 **Table 1. Gene transcriptional activities of elephant shark ERs by estrogens in HEK293**
284

		E1	E2	E3
ERα1	EC50 (nM)	0.49	0.0059	0.2
	Fold-Activation (± SEM) *	1.62 (± 0.16)	1.66 (± 0.04)	1.57 (± 0.09)
	Ratio**	0.98	1.00	0.95
ERα2	EC50 (nM)	0.29	0.016	0.31
	Fold-Activation (± SEM) *	1.51 (± 0.12)	1.79 (± 0.12)	1.82 (± 0.13)
	Ratio**	0.84	1.00	1.02
ERα3	EC50 (nM)	0.25	0.0093	0.29
	Fold-Activation (± SEM) *	1.55 (± 0.11)	1.75 (± 0.08)	1.87 (± 0.15)
	Ratio**	0.89	1.00	1.07
ERβ	EC50 (nM)	0.14	0.001	0.02
	Fold-Activation (± SEM) *	2.87 (± 0.16)	3.23 (± 0.26)	3.49 (± 0.26)
	Ratio**	0.89	1.00	1.08

285
*Fold activation values were calculated based on the 10 nM values.
**Ratio values were calculated by dividing the fold-activation value of estrogens by the value of E2.

4. Discussion

Significant advances have been made in elucidating the evolution of estrogen signaling in

animals [1, 2, 9, 11, 19, 23, 24].   Amphioxus, a chordate that is a close ancestor of vertebrates,

contains an ER and a steroid receptor (SR) that is an ortholog of receptors for 3-keto-steroids,

including testosterone, progesterone, cortisol, and aldosterone [11, 25].   Contrary to expecta-

tions, the amphioxus ER does not bind E2 or other steroids [11, 24, 25], while E2 and E1 are

transcriptional activators of the SR [11, 25].   Atlantic sea lamprey, a jawless fish, contains an

ER [19] that is activated by E2 and E1 [11, 24].

The estrogen receptors in Chondrichthyes, cartilaginous fishes with jaws, are basal to

lobe-finned fishes and are ancestors of terrestrial vertebrates [14].   These ERs are still not fully

characterized [15, 26].   The identification of ERα1, ERα2, ERα3, ERα4, and ERβ from the

sequencing of the elephant shark [14] provided an opportunity to investigate estrogen signaling

in a basal jawless vertebrate that evolved about 525 million years ago.   We investigated the re-

sponse to E1, E2, and E3 of ERα1, ERα2, ERα3, and ERβ from elephant shark expressed in

HEK293 cells (Figure 5).   These studies revealed that elephant shark ERα1, ERα2, ERα3, and

ERβ are activated by E1, E2, and E3.   We note that we found that ERα4 did not respond to an

estrogen, which we propose is due to a sequence deletion in the DBD (data not shown).   Thus,

we did not continue studies of estrogen activation of ERα4.

Our data with estrogen activation of ERα1, ERα2, ERα3, and ERβ align with the established

ligand binding characteristics of ERs in other vertebrates, thereby substantiating the conserved

nature of estrogen signaling pathways [25, 27-29].   Estrogen-dependent transcriptional activity

of the ERs from various species, including other sharks, revealed that the ERs from all species

have a stronger response to E2 than to E3 [16, 30-34].

Instead of only one ERα gene, elephant shark contains three ERα genes with very similar

sequences, which is unexpected.   The strong conservation of ERα4, which is not lig-

and-activated, also is unexpected.   Further research is needed to gain a more comprehensive

understanding of the physiological function of estrogen and the multiple estrogen receptors in

elephant sharks.   This data should provide clues for understanding the physiological function(s)

of each isoform.

In summary, this is the first report with evidence for the conservation of elephant shark

ERα and ERβ compared to human ERα and ERβ, which provides a valuable perspective on the

early evolutionary history of these critical nuclear receptors in vertebrates.   The findings from

this study not only confirm the conserved nature of ER-mediated estrogen signaling across ver-

tebrates but also open new avenues for research into the functional diversification of these re-

ceptors across vertebrates. These findings furnish researchers with critical molecular data to ex-

amine the role of ERs in future studies, such as those examining gonadal development, reproduc-

tive biology, and developmental and evolutionary endocrinology.

Author Contribution

Y.A.: Writing – review & editing, Writing – original draft, Methodology, Investigation, Formal

analysis, and Data curation.   **H.N.:** Writing – original draft, Methodology, Investigation, Formal

analysis, Data curation.   **W. T.:** Resources**.**   **S.H.:** Resources**.**   **M.E.B.:** Writing – review &

editing, Project administration, Validation, Supervision, Conceptualization.   **Y.K.:** Writing –

review & editing, Project administration, Visualization, Validation, Funding acquisition, Super-

vision, Conceptualization.

Funding

This research was supported by Grants-in-Aid for Scientific Research from the Ministry of Edu-

cation, Culture, Sports, Science and Technology of Japan (23K05839) to Y.K., and the Takeda

Science Foundation to Y.K.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relation-

ships that could have appeared to influence the work reported in this paper.

Data availability

All data is freely available on request to Dr. Katsu or to Dr. Baker.

Acknowledgment

The authors are grateful to Dr. Kaori Oka for her suggestions and helpful advice.   We also thank

colleagues in our laboratories.

5. References

[1] Bondesson M, Hao R, Lin CY, Williams C, Gustafsson JÅ. 2015. Estrogen receptor signaling

during vertebrate development. Biochim Biophys Acta 1849(2), 142-151. doi:
10.1016/j.bbagrm.2014.06.005.

[2] Gao H, Dahlman-Wright K. 2011. The gene regulatory networks controlled by estrogens. Mol
Cell Endocrinol. 334(1-2), 83-90. doi: 10.1016/j.mce.2010.09.002.

[3] Deroo BJ, Korach KS. Estrogen receptors and human disease. J Clin Invest. 2006
Mar;116(3):561-70. doi: 10.1172/JCI27987.

[4] Hamilton KJ, Hewitt SC, Arao Y & Korach KS 2017 Estrogen Hormone Biology. *Curr Top*
Dev Biol **125** 109-146.

[5] Paterni I, Granchi C, Katzenellenbogen JA & Minutolo F 2014 Estrogen receptors alpha
(ERalpha) and beta (ERbeta): subtype-selective ligands and clinical potential. *Steroids* **90**
13-29

[6] Heldring N, Pawson T, McDonnell D, Treuter E, Gustafsson JA, Pike AC. Structural insights
into corepressor recognition by antagonist-bound estrogen receptors. J Biol Chem. 2007
Apr 6;282(14):10449-55. doi: 10.1074/jbc.M611424200.

[7] Dahlman-Wright K, Cavailles V, Fuqua SA, Jordan VC, Katzenellenbogen JA, Korach KS,
Maggi A, Muramatsu M, Parker MG, Gustafsson JA. 2006. International Union of Phar-
macology. LXIV. Estrogen receptors. Pharmacol Rev. 58(4), 773-781. doi:
10.1124/pr.58.4.8.

[8] Markov GV, Tavares, R, Dauphin-Villemant C, Demeneix BA, Baker ME, Laudet V. 2009.
Independent elaboration of steroid hormone signaling pathways in metazoans. Proc Natl
Acad Sci USA. 106(29), 11913-11918. doi: 10.1073/pnas.0812138106.

[9] Baker ME. 2011. Origin and diversification of steroids: co-evolution of enzymes and nuclear
receptors. Mol Cell Endocrinol. 334(1-2), 14-20. doi: 10.1016/j.mce.2010.07.013.

[10] Kuiper GG, Enmark E, Pelto-Huikko M, Nilsson S, Gustafsson JA. Cloning of a novel re-
ceptor expressed in rat prostate and ovary. Proc Natl Acad Sci U S A. 1996 Jun
11;93(12):5925-30. doi: 10.1073/pnas.93.12.5925.

[11] Bridgham JT, Brown JE, Rodriguez-Mari A, Catchen JM, Thornton JW. 2008. Evolution of
a new function by degenerative mutation in cephalochordate steroid receptors. PLoS
Genet. 4(9), e1000191. doi: 10.1371/journal.pgen.1000191.

[12] Baker ME, Nelson DR, Studer RA. 2015. Origin of the response to adrenal and sex steroids:
Roles of promiscuity and co-evolution of enzymes and steroid receptors. J. Steroid
Biochem. Mol Biol 151, 12-24. doi: 10.1016/j.jsbmb.2014.10.020.

[13] Escande A, Servant N, Rabenoelina F, Auzou G, Kloosterboer H, Cavaillès V, Balaguer P,
Maudelonde T. 2009. Regulation of activities of steroid hormone receptors by tibolone
and its primary metabolites. J Steroid Biochem Mol Biol. 116(1-2):8-14. doi:
10.1016/j.jsbmb.2009.03.008.

[14] Venkatesh B, Lee AP, Ravi V, Maurya AK, Lian MM, Swann JB, Ohta Y, Flajnik MF,
Sutoh Y, Kasahara M, Hoon S, Gangu V, Roy SW, Irimia M, Korzh V, Kondrychyn I,
Lim ZW, Tay BH, Tohari S, Kong KW, Ho S, Lorente-Galdos B, Quilez J,
Marques-Bonet T, Raney BJ, Ingham PW, Tay A, Hillier LW, Minx P, Boehm T, Wilson
RK, Brenner S, Warren WC. 2014. Elephant shark genome provides unique insights into
gnathostome evolution. Nature 505(7482), 174-179. doi: 10.1038/nature12826.

[15] Filowitz GL, Rajakumar R, O'Shaughnessy KL, Cohn MJ. 2018. Cartilaginous Fishes Pro-
vide Insights into the Origin, Diversification, and Sexually Dimorphic Expression of Ver-
tebrate Estrogen Receptor Genes. Mol Biol Evol. 35(11), 2695-2701. doi:
10.1093/molbev/msy165.

[16] Katsu Y, Kohno S, Narita,H, Urushitani H, Yamane K, Hara A, Clauss TM, Walsh MT,
Miyagawa S, Guillette LJ, Iguchi T. 2010a. Cloning and functional characterization of
Chondrichtyes, cloudy catshark, Scyliorhinus torazame and whale shark, Rhincodon

typus estrogen receptors. Gen Comp Endocrinol. 168(3), 496-504. doi:
10.1016/j.ygcen.2010.06.010.
419

420 [17] Inoue JG, Miya M, Lam K, Tay BH, Danks JA, Bell J, Walker TI, Venkatesh B.
421   2010.Evolutionary origin and phylogeny of the modern holocephalans (Chondrichthyes:
422   Chimaeriformes): a mitogenomic perspective. Mol Biol Evol. 27(11):2576-2586. doi:
423   10.1093/molbev/msq147.
424

425 [18] Yu WP, Rajasegaran V, Yew K, Loh WL, Tay BH, Amemiya CT, Brenner S, Venkatesh B.
426   2008. Elephant shark sequence reveals unique insights into the evolutionary history of
427   verte-brate genes: A comparative analysis of the protocadherin cluster. Proc Natl Acad Sci
428   U S A. 105(10):3819-3824. doi: 10.1073/pnas.0800398105.
429

430 [19] Thornton JW. 2001. Evolution of vertebrate steroid receptors from an ancestral estrogen re-
431   ceptor by ligand exploitation and serial genome expansions. Proc Natl Acad Sci USA.
432   98(10), 5671-5676. doi: 10.1073/pnas.091553298.
433

434 [20] Baker ME. 2019. Steroid receptors and vertebrate evolution. Mol Cell Endocrinol.
435   496:110526. doi: 10.1016/j.mce.2019.110526.
436

437 [21] Kakumura K, Watanabe S, Bell JD, Donald JA, Toop T, Kaneko T, Hyodo S. 2009. Multi-
438   ple urea transporter proteins in the kidney of holocephalan elephant fish (*Callorhinchus*
439   *milii*). Comp Biochem Physiol B Biochem Mol Biol. 154(2), 239-247. doi:
440   10.1016/j.cbpb.2009.06.009.
441

442 [22] Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of
443   progressive multiple sequence alignment through sequence weighting, position-specific
444   gap penalties and weight matrix choice. Nucleic Acids Res. 22(22), 4673-4680. doi:
445   10.1093/nar/22.22.4673.
446

[23] Cotnoir-White D, Laperriere D, Mader S. 2011. Evolution of the repertoire of nuclear
receptor binding sites in genomes. Mol Cell Endocrinol. 334(1-2), 76-82. doi:

10.1016/j.mce.2010.10.021.

[24] Paris M, Pettersson K, Schubert M, Bertrand S, Pongratz I, Escriva H, Laudet V. 2008. An
amphioxus orthologue of the estrogen receptor that does not bind estradiol: insights into
estrogen receptor evolution. BMC Evol Biol. 8, 219. doi: 10.1186/1471-2148-8-219.

[25] Katsu Y, Kubokawa K, Urushitani H, Iguchi T. 2010. Estrogen-dependent transactivation of
amphioxus steroid hormone receptor via both estrogen and androgen response elements.
Endocrinology 151(2), 639-648. doi: 10.1210/en.2009-0766.

[26] Awruch CA. 2013. Reproductive endocrinology in chondrichtyans: the present and the future.
Gen Comp Endocrinol. 192, 60-70. doi: 10.1016/j.ygcen.2013.05.021.

[27] Green S, Walter P, Kumar V, Krust A, Bornert JM, Argos P, Chambon P . 1986. Human oes-
trogen receptor cDNA: sequence, expression and homology to v-erb-A. Nature.
320(6058), 134-139. doi: 10.1038/320134a0.

[28] Matthews J, Gustafsson JA. 2003. Estrogen signaling: a subtle balance between ER alpha
and ER beta. Mol Interv. 3(5), 281-292. doi: 10.1124/mi.3.5.281.

[29] Katzenellenbogen BS, Katzenellenbogen JA. 2000. Estrogen receptor transcription and
transactivation: Estrogen receptor alpha and estrogen receptor beta: regulation by selec-
tive estrogen receptor modulators and importance in breast cancer. Breast Cancer Res.
2(5), 335-344. doi: 10.1186/bcr78.

[30] Katsu Y, Lange A, Urushitani H, Ichikawa R, Paul GC, Cahill LL, Jobling S, Tyler CR,
Iguchi T. 2007. Functional associations between two estrogen receptors, environmental
estrogens and sexual disruption in the roach (Rutilus rutilus). Environ Sci Tech. 41(9),
3368-3374. doi: 10.1021/es062797I

[31] Katsu Y, Kohno S, Hyodo S, Ijiri S, Adachi S, Hara A, Guillette LJ, Iguchi T. 2008. Molec-
ular cloning, characterization and evolutionary analysis of estrogen receptors from

phylogenetically   ancient   fish.   Endocrinology   149(12),   6300-6310.   doi:
10.1210/en.2008-0670

[32] Katsu Y, Taniguchi E, Urushitani H, Miyagawa S, Takase M, Kubokawa K, Tooi O, Oka T,
Santo N, Myburgh J, Matsuno A, Iguchi T. 2010c. Molecular cloning and characterization
of ligand- and species-specificity of amphibian estrogen receptors. Gen Comp Endocrinol.
168(2), 220-230. doi: 10.1016/j.ygcen.2010.01.002

[33] Lange A, Katsu Y, Miyagawa S, Ogino Y, Urushitani H, Kobayashi T, Hirai T, Shears JA,
Nagae M, Yamamoto J, Ohnishi Y, Oka T, Tatarazako N, Ohta Y, Tyler CR, Iguchi T.
2012. Comparative responsiveness to natural and synthetic estrogens of fish species
commonly used in the laboratory and field monitoring. Aquat Toxicol 109, 250-258. doi:
10.1016/j.aquatox.2011.09.004.

[34] Yatsu R, Katsu Y, Kohno S, Mizutani T, Ogino Y, Ohta Y, Myburgh J, van Wyk JH, Guillette
LJ Jr, Miyagawa S, Iguchi T. 2016. Characterization of evolutionary trend in squamate
estrogen   receptor   sensitivity.   Gen   Comp   Endocrinol.   238,   88-95.   doi:
10.1016/j.ygcen.2016.04.005.

[35] Kuhl H. Pharmacology of estrogens and progestogens: influence of different routes of ad-
min-istration. Climacteric. 2005 Aug;8 Suppl 1:3-63. doi: 10.1080/13697130500148875.

[36] Stanczyk FZ. 2024. Metabolism of endogenous and exogenous estrogens in women. J Ster-
oid Biochem Mol Biol. 2024 Sep;242:106539. doi: 10.1016/j.jsbmb.2024.106539.