Immune-mediated Engagement of T Regulatory Cells with Tumor Cells Results in Trogocytosis and Tumor Cell Killing

Amy K Erbe[1]*, Arika S. Feils[1], Anqi Gao[2], Sabrina VandenHeuvel[1], Simon Boyenga[1], Alina Hampton[1], Mackenzie Heck[1], Jen Zaborek[3], Daniel Spiegelman[1], Noah Tsarovsky[1], Anna Hoefges[1], Alexander A. Pieper[1], Peter M. Carlson[1], Mildred Felder[1], Ravi B Patel[4], Stephen D. Gillies[5], Alexa Heaton[1,6,7], Zachary Morris[1], Huy Q. Dinh[2,3], Alexander L. Rakhmilevich[1], Paul M. Sondel[1,8]

[1]Department of Human Oncology, University of Wisconsin, Madison WI
[2]Department of Oncology, University of Wisconsin, Madison WI
[3]Department of Biostatistics and Medical Informatics, University of Wisconsin, Madison WI
[4]University of Pittsburgh, Pittsburgh PA
[5]PineTree Therapeutics Inc. Carlisle, MA
[6]Morgridge Institute for Research, Madison WI
[7]Department of Biomedical Engineering, University of Wisconsin, Madison WI
[8]Department of Pediatrics, University of Wisconsin, Madison, WI

Corresponding Author:
Amy Erbe PhD
Research Asst. Professor
University of Wisconsin
UW Carbone Cancer Center
aerbe@wisc.edu

Summary

Despite advances in treatment, >50% of patients with advanced melanoma are unresponsive to current therapies. Using the B78-D14 melanoma model (GD2$^+$/MHC-I$^-$/MHC-II$^+$), we can cure mice with a regimen that includes radiotherapy (RT) in combination with immunocytokine (IC; anti-GD2 monoclonal antibody linked to IL-2) while establishing immunological memory. We interrogated the role of T cells in the antitumor and memory responses following RT+IC. We show a requirement for CD4, but not CD8 T cells, to achieve both the initial and memory responses. Upon IC-induced cell-cell contact, subsets of CD4 T cells, including Foxp3$^+$ T regulatory cells, trogocytose GD2 from tumor cells, acquire cytotoxic granules, and kill tumor cells. These results were confirmed using human tumor cell lines. These findings reveal that CD4$^+$ T regulatory cells, upon immunologically-induced binding to tumor cells, can trogocytose tumor antigens and directly kill tumor cells, redefining their potential role in antitumor immunity.

Keywords

Immunocytokine, Effector Tregs, Trogocytosis, CD4s, Radiation, Treg, Melanoma, MHC-I, MHC-II

Introduction

The incidence of melanoma in the U.S continues to rise. Among patients with advanced disease, the current standard of care—immune checkpoint inhibition (ICI)—achieves a 5-year progression-free survival (PFS) rate of less than 30%.[1,2] Tumors that fail to respond to ICI often harbor features that promote growth while suppressing or evading immune recognition and elimination.[3-5] Although ICI has markedly improved outcomes across several cancers, including melanoma, its benefit is not universal.[4,6] Melanomas with a high tumor mutation burden (TMB) are considered "hot" and have actionable mutation-driven neoantigens, prominent Major Histocompatibility Complex class I (MHC-I) expression, and substantial T cell infiltration prior to therapy. These characteristics indicate an active endogenous immune response that is being dampened by checkpoint signaling. Patients with "hot" melanomas are more likely to benefit from ICI;[7-9] however >70% of advanced melanoma patients do not show prolonged PFS in response to ICI. Most of these non-responsive tumors are considered "cold", characterized by a low actionable TMB, poor immune infiltration, and/or loss of MHC-I.[10-15]

T cells can be primed to specifically recognize, remember, and eliminate melanoma by engaging melanoma-derived peptides presented in the cleft of MHC-I and/or MHC-II molecules on tumor cells. The resulting immune activation and memory enable ongoing surveillance, through which T cells identify and eradicate refractory, recurrent or disseminated melanoma cells unless their activity is suppressed by immune checkpoints—circumstances under which ICI therapy can restore function. However, downregulation or loss of MHC-I is common in melanoma, preventing recognition by tumor-specific CD8 T cells (CD8s). Thus, MHC-I loss is a common mechanism of immune evasion and resistance to therapy.[5,16,17] Separately, MHC-II molecules can present antigen to CD4 T cells (CD4s), initiating a cascade of immune responses. While not commonly expressed on most solid tumors and usually restricted to expression on immune cells, MHC-II is expressed on 50-60% of melanomas, but the implications of MHC-II expression on melanoma are not well understood.[16,18-21] Whether MHC-II+ tumors directly, or indirectly, engage CD4s likely depends on the tumor microenvironment (TME) and treatment.[22] CD4 effector, helper, and T regulatory (Treg) cells can be activated through MHC-II. Within the TME, Tregs may expand preferentially in MHC-II+ tumors, and stimulation of Tregs through MHC-II interactions can enhance their immunosuppressive activity.[23] Yet paradoxically, patients with MHC-II+ melanomas tend to respond more favorably to ICI therapy.[21] Thus, it is possible the antitumor response to ICI for patients with MHC-II+ tumors may reflect a balance of Treg versus T effector (T_{eff}) activation prior to ICI treatment, though additional data are needed to clarify these mechanisms.

To investigate how MHC-I/II expression influences T cell responses induced by immunotherapy, we used the "immunologically cold" B78-D14 murine melanoma tumor model, which is poorly infiltrated by immune cells and resistant to ICI alone.[24-27] B78-D14 cells (B78-D14s) express GD2, a tumor-antigen expressed on many melanomas, but expressed minimally on most normal tissues. Using this translationally relevant model, combining radiation therapy (RT) with an immunocytokine [IC; anti-GD2 monoclonal antibody (mAb) linked to IL-2] cures approximately 50% of B78-D14 tumor-bearing mice, inducing durable, T cell-dependent, tumor-specific immune memory.[25] Notably, B78-D14s lack MHC-I expression, even after IFN-γ stimulation, but can express MHC-II following IFN-γ treatment.[28,29] We therefore examined the mechanisms underlying curative immune responses in this MHC-II+ B78-D14 tumor model to identify TME factors that support adaptive immunity and that mediate tumor killing. Our studies revealed a critical role for CD4s in response to the RT+IC regimen. We observed that CD4s, including Tregs, engaged in tumor cell trogocytosis, which was accompanied by degranulation, indicating cytotoxic activity. These findings demonstrate that, upon IC treatment, Tregs can trogocytose

tumor antigens, acquire effector-like properties, and contribute directly to tumor killing in MHC-II[+] melanoma.

Results

CD4 T cells are required for tumor control

We have previously established a radio-immunotherapy regimen in which radiation therapy (RT) is followed by five consecutive doses of immunocytokine (IC) (Fig. 1A), showing that RT+IC elicits durable, T cell dependent antitumor responses in the B78-D14 melanoma model.[25] B78-D14s have aberrant expression of MHC-I and MHC-II, with discordant results reported and we observe that co-culture of B78-D14 with IFN-γ for 48h induces expression of MHC-II but not MHC-I (**Fig S1A**).[28-30] Although IFN-γ treatment can upregulate MHC-II on B78-D14 cells, the resulting expression is heterogeneous, with a subset of tumor cells exhibiting little to no detectable MHC-II by flow cytometry (**Fig. S1A**). Tumor cell expression of MHC-II can augment the response to immunotherapy, as MHC-II[+] tumor cells can function as antigen-presenting cells (APCs), stimulating T-helper (Th) and T-effector (T_eff) responses to promote adaptive antitumor immunity.[21,31,32] *Ex vivo* analyses of B78-D14 tumors following RT+IC revealed increased MHC-II expression in tumors from "responding" mice (based on tumor regression) compared to tumors from non-responders (**Fig. 1B-C, Fig. S1B-D**), suggesting that MHC-II expression may contribute to effective antitumor responses. Because MHC-II interacts with CD4s, these findings raise the possibility that CD4-mediated mechanisms underlie part of the therapeutic efficacy in this model. Indeed, immune cell depletion studies confirmed that CD4s are required for the initial antitumor response: mice depleted of CD4s during RT+IC therapy failed to respond, whereas depletion of CD8 or NK cells did not impair therapeutic efficacy (**Fig. 1D, Fig. S2A-2B**). Conversely, in two other tumor models (NXS2 and MC38) that express MHC-I (**Fig S1A**), antitumor efficacy to radio-immunotherapy (RT+IC in NXS2 or RT+IL-2+anti-CTLA4 for MC38) exhibited a distinct dependency on CD8s, but not CD4s (**Fig. S2B-2D**).[33] These findings suggest that, in contrast to the MHC-II[+] B78-D14 model, which depends on CD4s, expression of MHC-I in NXS2 and MC38 promotes CD8-driven antitumor responses. Taken together, these findings are consistent with a model in which MHC-II expression favors CD4-driven antitumor responses, whereas MHC-I expression is associated with CD8-driven responses, although additional factors within the TME likely also influence these dynamics.[34]

We next examined the efficacy of T cell priming within the tumor-draining lymph node (TDLN) following RT+IC treatment to determine whether CD4s and CD8s were sufficiently activated to attack the tumor. In the TDLN, RT+IC treatment increased proliferation of both CD4s and CD8s (**Fig. 1E**). Single cell secretory cytokine analyses showed that CD4s and CD8s from RT+IC-treated mice were similarly activated (**Fig. 1F**), with t-SNE plots showing clustering of both treated subsets (CD4s and CD8s) together, distinct from their respective no treatment (NT) counterparts (**Fig. 1G**). This indicates that RT+IC induces a shared cytokine-production phenotype in CD4s and CD8s, different from those in NT conditions. Both CD4 and CD8 T cell types from RT+IC groups also exhibited elevated production of effector cytokines compared with NT controls (**Fig. 1F-H**).

To understand the contribution CD4s play during the antitumor response, we depleted CD4s during the course of RT+IC treatment and compared immune activity within the TME to mice that had CD4s. Within the TME, on D8 after RT+IC treatment initiation, immune cells from RT+IC-treated mice exhibited distinct immunophenotypes with enriched effector subsets which were lost if CD4s were depleted during treatment, suggesting CD4 depletion abrogates RT+IC-induced TME reprogramming (**Fig. 1I-M**). RT+IC significantly increased IFN-γ and TNF-α in CD4s, which could explain the parallel

IFN-γ and TNF-α enhancements observed in CD8s and NK cells (**Fig. 1I-J**). However, when CD4s were depleted during RT+IC, IFN-γ and TNF-α production by CD8s and NK cells was significantly reduced, indicating impaired activation. We also examined cytotoxic potential, defined by lytic granule release of perforin (Prf) and CD107a expression. RT+IC increased cytotoxic CD4s in the TME (**Fig. 1K-L**). Although CD8s and NK cells were not required for tumor control against B78-D14s, we observed an increase in Prf-expressing CD8s and NK cells following RT+IC, with NK cells having the highest levels (**Fig. 1K**). Notably, despite increased Prf expression, CD8s and NKs lacked CD107a surface expression, suggesting limited degranulation and cytotoxic activity. In contrast, CD4s were the only subset with increased CD107a$^+$ and Prf$^+$ cells following RT+IC, implicating cytotoxic CD4s as key effector cells (**Fig. 1M**).

Figure 1: CD4$^+$ T cells are Required for Response to RT+IC in B78-D14 Tumors.
(A) Treatment and harvest schema for mice bearing B78-D14 tumors treated with RT and IC. **(B)** In three independent studies, mice treated with RT+IC were categorized as responders (R) vs. non-responders (NR) based on tumor size at time of harvest (top) prior to harvest. Tumors were then analyzed for MHC-II expression (bottom); see additional details in **Fig. S1**. **(C)** Increased tumor MHC-II expression significantly correlated with treatment response. n = 7-11/group; 3 independent experiments combined. **(D)** Immune subset depletion demonstrated that CD4s were required for tumor rejection, while CD8s and NK cells were dispensable. n=11-18/group; 3 independent experiments combined. **(E–H)** From TDLNs harvested on day 7: **(E)** RT+IC increased Ki67 expression in both CD4$^+$ and CD8$^+$ T cells. **(F)** Isoplexis analyses showed RT+IC enhanced production of effector cytokines in both T cell subsets, along with increased regulatory cytokines**. (G)** IsoPlexis tSNE plots showed that RT+IC-CD4s and RT+IC-CD8s clustered together with distinct cytokine profiles compared to NT counterparts. **(H)** Polyfunctional Activation Topology Principal Component Analysis displays cytokine secretion for CD4s and CD8s based on treatment. Each grouping/circle of datapoints represents the same functional group that are randomly offset, but remain within a radius proportional to the secretion frequency of the corresponding group (i.e., large groups = large circles, small groups = small circles). The principal components are labeled according to their correlation with specific cytokines. Here again, both RT+IC-CD8s and RT+IC-CD4s are highly polyfunctional with expression combinations of MIP-1α and IL-27 secreting cells, as well as IFN-γ, CCL-11, IL-5 and sCD137 polyfunctional cells. NT samples pooled from n=3 mice; RT+IC samples pooled from n=3 mice. **(I–J)** From tumors excised on day 8: RT+IC increased IFN-γ (**I**) and TNF-α (**J**) in NK, CD4s, and CD8s, but CD4 depletion abrogated this effect in CD8s and NKs. Perforin was increased in all subsets with RT+IC and remained elevated in CD8s and NKs following CD4 depletion (**K**). However, CD107a expression was restricted to CD4s (**L**), which uniquely co-expressed perforin and CD107a (**M**), indicating direct cytotoxic activity. n = 18/group; 4 independent experiments combined. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001; Mean +/- SEM.

CD4 T cells mediate both primary and memory responses

In addition to a requirement for CD4s in the initial antitumor response against B78-D14, we interrogated whether CD4s played a role in long-term immune memory. After initial tumor clearance, tumor-free mice were rechallenged with B78-D14 tumors to evaluate immune memory (**Fig 2A**). Mice depleted of CD4s during the rechallenge were unable to reject B78-D14 tumors, whereas depletion of CD8s or NK cells did not impair rejection (**Fig. 2B-C**). These findings indicate that CD4s not only drive the primary antitumor response but are also essential for mediating rejection of the rechallenged tumor, indicating sustained immune memory.

To further characterize T cell responses in the memory setting, we analyzed TDLNs from mice cured of B78-D14 tumors with RT+IC 7d following rechallenge with B78-D14 ("memory-mice"). Flow cytometry t-SNE analyses showed marked phenotypic changes in both CD4s and CD8s compared with naïve or tumor-bearing (NT) controls (**Fig. 2D**). Upon rechallenge, memory-mice displayed decreased CD4 T central memory (T_{cm}) but increased CD4 T effector memory (T_{em}) (**Fig. 2E**), with no major shifts in the CD8 memory subsets (**Fig. 2F**). Both CD4s and CD8s expressed elevated CD38 during rechallenge, but only CD4s showed significantly increased CD69, suggesting preferential CD4 activation (**Fig. 2G-2H**).[35,36]

We examined LNs from memory-mice in comparison to tumor-bearing control or naïve mice. In the LNs of naïve mice and the TDLN of tumor-bearing (NT) mice, the T_{em} phenotype was more prominent among CD8s than CD4s. In contrast, in the TDLNs of memory-mice, CD4s exhibited a more pronounced T_{em} phenotype compared to CD8s (**Fig. 2I**). Within tumors, memory-mice likewise showed a higher proportion of CD4 T_{em} relative to tumor-bearing controls (**Fig. 2I**). Notably, among CD45$^+$ cells in the memory-mice, CD4 and CD8 subset distributions were comparable in TDLNs, whereas the tumor compartment was dominated by CD4s (**Fig. 2J**). Functionally, only CD4s—not CD8s—contained significantly increased cytotoxic granule content (GzmB and/or Prf; **Fig. 2K**) and degranulation (CD107a; **Fig. 2L**) in memory-mice vs. tumor-bearing mice. Furthermore, CD4s from memory-mice had increase CD107a expression compared to CD8s from the same mice (**Fig. 2L**). Together, these results indicate that CD4s are the primary mediators of the memory response to B78-D14 tumors, characterized by a dominant effector memory and cytotoxic phenotype driving long-term protection.

Figure 2. CD4$^+$ T cells are Required for the Memory Response in mice cured by RT+IC.
A) Treatment and harvest schematic for tumor-free (TF) mice after RT+IC that were rechallenged with B78-D14s ~30 days post-clearance. **B-C)** Immune cell depletion during rechallenge revealed that CD4s, but not CD8s or NK cells, were required for rejection, as CD4-depleted mice exhibited tumor growth comparable to naïve controls. n=15-30/group; 2 independent experiments combined. **D)** Flow cytometry of LNs from: age matched naïve mice (i.e., never injected with B78-D14s); or the TDLN 7d after previously naïve mice were injected with B78-D14s ("tumor bearing"), or TDLN from "memory" mice using mice cured B78-D14 tumors by RT+IC and rechallenged with B78-D14s, with TDLN harvested 7d after B78-D14 injection, showed shifts in T cell subsets (T_{cm} and T_{em}) based on tSNE analysis of CD4, CD8, CD62L and CD44). **E-H)** Flow cytometry of TDLNs identified memory phenotypes (Naïve, CD44$^-$CD62L$^+$; T_{cm}, CD44$^+$CD62L$^+$; $T_{em,}$ CD44$^+$CD62L$^-$; terminally differentiated effector memory cells (Temra, CD44$^-$CD62L$^-$). Memory-mice exhibited significantly reduced CD4$^+$ T_{cm} and significantly increased CD4$^+$ T_{em} (**E**), with no differences found in CD8$^+$ memory subsets (**F**). Memory-mice also showed enhanced expression of activation markers—CD69 on CD4s (**G**) and CD38 on both CD4s and CD8s (**H**). **(I)** In TDLNs and TME, CD4$^+$ Tem were enriched in memory-mice compared to naïve or tumor-bearing controls. (**J**) CD4s also constituted a greater fraction of the CD45$^+$ cells than the CD8s in the TME (but not TDLNs) in memory versus tumor-bearing mice. **(K-L)** Within the TME of memory-mice, CD4s expressed higher levels of granzyme B, perforin, and CD107a, consistent with cytotoxic function, whereas CD8s did not. n=5/group; 2 independent experiments. *p < 0.05, ***p < 0.001; Mean +/- SEM.

scRNA-seq identifies phenotypic shifts with functional adaptation of CD4 subsets

To define how RT+IC reshapes CD4$^+$ T cell states, we performed scRNA-seq on tumors from NT and RT+IC-treated mice (**Fig. 1A**). RT+IC increased NK and T cell abundance and substantially reduced tumor cells, consistent with treatment-induced tumor clearance (**Fig. 3A**). Within CD4s, we identified six transcriptionally distinct clusters: naïve, T effector cells (T$_{eff}$), T memory cells (T$_{em}$), pTregs (proliferative Tregs), eTregs (effector *GzmB_high* Tregs), and oTregs (other Tregs) (**Fig. 3B)** based on their respective gene expression patterns (**Fig. 3C**). Of these six clusters, oTregs were decreased following RT+IC treatment, whereas both T$_{em}$ and eTregs were increased (**Fig. 3D**).

Given that CD4s are required for both primary B78-D14 tumor control and memory rechallenge (**Fig. 1C** and **Fig. 2B**), and the association between tumor MHC-II expression and antitumor response (**Fig. 1B**), we first focused on transcriptional programs relevant to CD4 TCR signaling (**Fig. 3E**). T_{eff} cells (from both NT and RT+IC mice) expressed high levels of TCR activation genes (*Nr4a1*[37] and *Erg1*[38]), while T_{em} cells (from both NT and RT+IC mice) expressed proximal signaling *Lck* and *Lat*, critical for TCR signaling[39] (**Fig. 3E**). pTregs also showed high levels of expression of these genes after RT+IC, including *Nr4a1* and *Egr1*, which are linked to Treg differentiation.[40,41] Core Treg-identity genes, *Ctla4* and *Foxp3*, were decreased with RT+IC treatment in pTregs, while *IL-2Ra* was increased across all Treg subsets with RT+IC (**Fig. 3F**). This increased *IL-2Ra* (IL-2Rα/CD25) expression in the RT+IC group may be due to elevated IL-2 availability in the TME following IC treatment, acting on IL-2Ra expressing Tregs.[42] Despite increased *IL-2Ra* expression with RT +IC, none of the Treg subsets showed increased expression of classical inhibitory or suppressive pathway genes such as *Lag3*[43], *Pcdc1*[44], *Tgfb1*[45] with treatment (**Fig. 3G**).

We then evaluated transcriptional programs related to Treg stability and effector adaptation (**Fig. 3H-I**).[46] Helios (*Ikzf2*), a key regulator of Treg stability[47], was expressed in all Treg subsets, but was decreased in pTregs following RT+IC (**Fig. 3H**). Conversely, Eos (*Ikzf4*)[48], a major regulator of Treg stability was minimally expressed in each Treg subset. *Foxo1*[49,50], required for Treg development and function, was decreased with RT+IC in all 3 Treg populations (**Fig. 3H**). Together, these features suggest that RT+IC promotes Tregs plasticity[51], and may promote Tregs to adopt effector-like characteristics upon RT+IC. Supporting this, *Socs2,* an essential regulator of Treg plasticity[52], was increased in all 3 Treg clusters with RT+IC (**Fig. 3I**). Genes linked to effector Treg adaptation, including GITR (*Tnfrsf18*)[53], were upregulated in all three Treg clusters with treatment, with increased *IFN-γ* expression in pTregs and substantially elevated *Gzmb* in all Treg subsets, particularly eTregs (**Fig. 3I**). These features suggest that RT+IC promotes a shift toward effector-like, cytolytic Treg states within the TME.

Figure 3. scRNAseq Reveals CD4 Treg Cells are Engaged in the Antitumor Response
Mice bearing B78-D14 tumors, either untreated (NT) or treated with RT+IC, were assessed by scRNA-seq on day 8. **A)** RT+IC treatment reduced tumor cell abundance and reshaped the immune landscape, with redistribution of CD4$^+$ subsets (non-Tregs and Tregs) and increased infiltration of NK cells and CD8s within the TME. Myeloid and dendritic cells remained minimal in both NT and RT+IC tumors. **B)** Focusing on CD4s, six distinct populations were identified: naïve, Teff, Tem, eTregs, oTregs and pTregs. **C)** Subset-specific RNA expression patterns revealed distinct transcriptional profiles for each CD4 grouping. **D)** With RT+IC, the frequency of Tem and eTregs increased, while naïve, Teff, and oTregs decreased. (**E-H**) Key gene expression levels were examined in pathways that contribute to: (**E**) TCR signaling; (**F**) Treg phenotype; (**G**) Treg inhibition; (**H**) Treg stability; and (**I**) Treg effector function in Treg and other CD4 subsets following RT+IC. n=2/group; 1 independent experiment (the color scheme corresponds to that in **Fig. 3B**).

Tumor trogocytosis by CD4 T cells correlates with effector function

Following RT+IC-treatment, a subset of CD45$^+$ immune cells became GD2$^+$ (**Fig. 4A**). GD2, the target antigen of the anti-GD2 IC component of this therapy, is a disialoganglioside that is not often expressed on immune subsets.[54] Among the CD45$^+$ cells, only CD4s, but not CD8s or NK cells, were GD2$^+$ (**Fig. 4B**). There was no difference in the amount of tumor-infiltrating CD4s based on treatment, but with RT+IC there was a significant in increase in the amount of GD2$^+$CD4s compared to NT (**Fig. 4C**). RT+IC increased CD107a expression on CD4s, with GD2$^+$ CD4s expressing significantly more CD107a than GD2$^-$ CD4s (**Fig. 4D**). Various methods were applied to validate this observed GD2 expression on T cells as opposed to the presence of doublets (CD4s in direct contact with GD2$^+$ tumor cells) or GD2$^+$ tumor cells aberrantly expressing CD4. Image Stream flow cytometry revealed that while some CD4s were detected in direct contact with GD2$^+$ tumor cells as doublets (**Fig. 4E**), the majority of CD4$^+$/GD2$^+$ signals were instead isolated CD4$^+$ cells that displayed punctate GD2 staining on their membranes (**Fig. 4E**, **Fig. S3A**). *In vitro* coculture of T cells isolated from the TDLN of a naïve mouse with B78-D14s, followed by sorting and cytospin of GD2$^+$CD4$^+$ cells, further confirmed that the double-positive population represented CD4s that had acquired GD2, rather than doublets of CD4 cells bound to GD2$^+$ tumor cells or tumor cells that aberrantly expressed CD4 (**Fig. S3B-C**). These data indicated that CD4s acquired GD2 expression via trogocytosis of GD2$^+$ tumor membranes.

Trogocytosis involves direct membrane transfer from a donor cell (here, GD2$^+$ tumor cells) to a recipient cell (CD4$^+$ T cells) during close contact.[55-58] Trogocytosis by CD4s is primarily thought to be mediated via interaction of the CD4 TCR with MHC-II, most commonly observed through CD4 interactions with APCs. Because B78-D14s express MHC-II (upregulated in responding mice, **Fig. 1B–C**), we hypothesized that CD4 trogocytosis was mediated via TCR:MHC-II interactions. To test this, we incubated TDLNs from RT+IC-treated mice with various tumor lines differing in MHC and GD2 expression (**Fig. 4F**). Significant GD2 acquisition by CD4s was observed only after co-culture with MHC-II$^+$GD2$^+$ B78-D14 cells (**Fig. 4G**), suggesting that tumor co-expression of both GD2 and MHC-II were needed to facilitate or enhance trogocytosis by CD4s in this setting. Similarly, CD4s from naïve and memory-mice became GD2$^+$ only after exposure to B78-D14, with significantly higher acquisition in memory CD4s, consistent with TCR recognition facilitating trogocytosis (**Fig. 4H**).

To determine how trogocytosis was associated with CD4 effector function, CD4s were isolated from TDLNs of control (NT) and treated (RT+IC) B78-D14-bearing mice and co-incubated with B78-D14s (**Fig. 4F**). In contrast to tumor-infiltrating CD4s (**Fig 4C**), TDLN-CD4s were GD2$^-$ at baseline but became GD2$^+$ following co-culture (**Fig. 4I**). TDLN-CD4s from RT+IC-treated mice showed high GzmB expression prior to co-culture (pre-incubation), which was lost after incubation (**Fig. 4J**), consistent with degranulation by cells that had released their cytotoxic granules. In line with this, the degranulation marker, CD107a, a separate indicator of recent cytotoxic activity, was increased with co-culture specifically in GD2$^+$CD4s (**Fig. 4K**). The amount of GD2 expression on CD4s correlated with CD107a levels (**Fig. S4**), with GD2$^+$ CD4s from RT+IC-treated TDLNs expressing significantly higher CD107a than those from NT mice after co-culture, further supporting the link between GD2 acquisition and enhanced effector function (**Fig. 4K**). Together, the loss of intracellular GzmB, the presence of CD107a, and the acquisition of GD2 post-incubation, indicate that these GD2$^+$CD4s had degranulated upon target (GD2$^+$ tumor cell) engagement and acquired GD2 via trogocytosis.

This phenomenon persisted within the TME of rechallenged memory-mice. Mice cured of primary B78-D14 tumors were subsequently rechallenged with a second B78-D14 injection and TME cells were harvested 7 days later for analyses of CD45$^+$GD2$^+$ cells (**Fig. 4L**). Similar to trends observed in the initial antitumor response, most GD2$^+$ cells in the rechallenge TME were CD4s with punctate GD2 staining, consistent with trogocytosis rather than tumor:T cell doublets (**Fig. 4M-N, Fig. S3D**). CD4$^+$ cells

remained the predominant CD45$^+$GD2$^+$ population, with more GD2$^+$CD4$^+$ cells in rechallenge memory-mice than GD2$^+$CD8$^+$ cells (**Fig. 4O**). Notably, GzmB expression was significantly increased on GD2$^+$CD4s from the TME of rechallenged memory-mice compared to GD2$^+$CD8s (**Fig. 4P**). GD2$^+$CD4s from rechallenged memory mice exhibited significantly higher CD107a expression compared with GD2$^+$CD8s from the same mice, indicating enhanced degranulation by CD4s during the memory response (**Fig. 4Q**). Collectively, these results demonstrate that CD4$^+$ T cells undergo tumor-directed trogocytosis during both primary and memory responses. Acquisition of GD2 correlates with cytotoxic degranulation (CD107a), suggesting that trogocytosis is linked to the effector function of CD4s in this tumor model.

Figure 4. Trogocytosis of B78-D14 by CD4s Influences Effector Cell Function
A) Within the TME of B78-D14 tumor–bearing mice, a subset of CD45$^+$ cells became GD2$^+$ by day 8 after RT+IC (Fig. 1A). **B)** Among immune subsets, CD4s—rather than CD8s or NK cells—displayed the highest GD2 expression following RT+IC. **C)** RT+IC did not alter the overall proportion of CD4s within the TME (left plot) but significantly increased the frequency of GD2$^+$CD4s (right plot). **D)** RT+IC increased the % of CD107a$^+$GD2$^+$CD4s (**D**) compared to NT. No CD107a$^+$ cells were detected among GD2$^-$CD4s under either treatment condition. **E)** Image Stream analysis of tumors excised on day 8 revealed (top panel) uniform GD2 expression on tumor cells (anti-GD2 mAb, green), while CD4s were smaller and stained homogeneously with anti-CD4 (red). Some CD4$^+$/GD2$^+$ signals reflect doublets of green tumor cells adhering to red CD4+ cells (top panel), while most reflected a subset of isolated CD4s that displayed bright CD4 signal and punctate GD2 staining (bottom panel), consistent with trogocytosis of GD2 from tumor cells. n=8/group; 2 independent experiments combined. **F)** Schema for TDLNs from B78-D14-bearing mice cocultured with MC38 (GD2$^-$MHC-I$^+$/MHC-II^{lo}), B16 (GD2$^-$MHC-I$^+$/MHC-II$^+$), B78-H1 cells (GD2$^-$/MHC-I$^-$/MHC-II$^-$), 9464D-GD2 (GD2$^+$MHC-II$^-$/MHC-I$^-$/) or B78-D14 (GD2$^+$/MHC-I$^-$/MHC-II$^+$) cells. **G)** Pre-incubation, CD4s expressed minimal GD2. Post-incubation, GD2 acquisition occurred only with the GD2$^+$ B78-D14 coculture, not with other lines. **H)** In similar cocultures, CD4s from TDLNs of tumor-free RT+IC-treated "memory" mice showed greater GD2 acquisition than CD4s from naïve controls. **I)** Some CD4s acquire GD2 post-incubation with B78-D14s regardless of *in vivo* treatment status. **J)** Prior to coculture, CD4s from RT+IC-treated mice expressed elevated GzmB compared to NT; post-incubation with B78-D14, GzmB levels returned to baseline. **K)** CD107a was absent from GD2$^-$CD4s but selectively upregulated in GD2$^+$CD4s after B78-D14 coculture, with significantly higher GD2$^+$CD107a$^+$ frequencies in RT+IC-treated mice. n=3-18/group; 2-5 independent experiments combined. **L)** In the TME of rechallenged memory-mice, a subset of CD45$^+$ cells became GD2$^+$ by day 7 post-rechallenge. **M-N)** Image Stream confirmed some CD4/tumor doublets with GD2$^+$ tumor cells interacting with CD4s (**M**), and isolated CD4s displaying punctate GD2 staining (**N**). **O)** Within the TME post-rechallenge, GD2 expression on CD45$^+$ cells was predominantly restricted to CD4s, with higher frequencies in memory versus tumor-bearing mice. **P-Q)** Among GD2$^+$CD45$^+$ cells, CD4s from memory mice expressed significantly more GzmB+/-Prf (**P**) and CD107a (**Q**) compared to CD8s, highlighting cytotoxic activity unique to CD4s. n=8/group; 2 independent experiments combined. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001; Mean +/- SEM.

Treg subsets mediate trogocytosis and acquire cytotoxic potential

Trogocytosis, the direct acquisition of membrane proteins from other cells, can influence CD4 T cell activation and function.[56,59,60] Because CD4s undergoing trogocytosis with RT+IC treatment (**Fig. 4C&E**) correlated with degranulation, and that Tregs exhibited an effector phenotype during treatment (**Fig. 3B–D**), we examined how GD2 acquisition related to the cytotoxic phenotype of CD4 Tregs. Within the TME, treatment-dependent shifts were observed in GD2 acquisition, cytotoxic markers (CD107a, GzmB), and Treg markers (Foxp3, CD25) (**Fig. 5A, Fig. S5**). Following RT+IC, GD2$^+$Foxp3$^+$ cells were detected in the TME, including both CD25$^+$ and CD25$^-$ subsets (**Fig. 5B**), with a significant increase in GD2$^+$/CD25$^-$/Foxp3$^+$ cells compared to NT tumors (**Fig. 5B**). GzmB expression was absent in CD4 subsets from NT tumors, regardless of CD25/Foxp3/GD2 expression, but was present in multiple populations following RT+IC—including GD2$^{-/+}$/CD25$^-$/Foxp3$^-$ CD4s, GD2$^-$/CD25$^-$/Foxp3$^+$ CD4s and GD2$^+$/Foxp3$^+$ CD4s (**Fig. 5C and Fig. S5D**). Following RT+IC, with the exception of GD2$^+$/CD25$^+$/Foxp3$^-$ cells, all GD2$^+$CD4 subsets (CD25$^+$Foxp3$^+$, CD25$^-$Foxp3$^+$, and CD25$^-$Foxp3$^-$) displayed significantly increased CD107a expression, indicating degranulation (**Fig. 5D, Fig. S5E**).

To determine which components of the treatment regimen contributed to these phenotypes, we evaluated each element alone or in combination: NT, RT alone, IC (anti-GD2 genetically linked to IL-2), RT+IL-2, RT+anti-GD2mAb, RT+anti-GD2mAb+IL-2, or RT+IC. Treatments including RT significantly increased Foxp3$^+$CD4s compared to NT, whereas IC alone did not (**Fig. 5E**). Only treatments containing IC (IC alone or RT+IC) generated GD2$^+$Foxp3$^+$CD4s (**Fig. 5F**), with these groups showing the highest expression of GzmB (**Fig. 5G**) and CD107a (**Fig. 5H**) within the GD2$^+$Foxp3$^+$CD4s. Separately, *in vitro* observations (**Fig. 4J**) indicated that some CD4$^+$ cells from RT+IC–treated mice expressed GzmB prior to co-culture and acquired GD2 during IC-mediated interactions with tumor cells. GD2$^+$Foxp3$^+$CD4s isolated directly from the TME exhibited both GD2 and GzmB at harvest, consistent with dynamic regulation of cytotoxic granules during tumor engagement.[61]

Figure 5. Effector Tregs Trogocytose Tumor Cells and Degranulate
A) Flow cytometry of TMEs harvested on day 8 post-treatment revealed treatment-dependent shifts in cytotoxicity markers (GzmB, CD107a) and Treg markers (Foxp3, CD25), aligned with GD2 expression patterns (UMAP analysis, **Fig. S5**). **B)** Following RT+IC, GD2 expression was detected on Foxp3$^+$ cells (CD25$^{+/-}$) as well as other CD4 subsets. **C)** RT+IC significantly increased GzmB expression across CD4 subsets, including Foxp3$^+$CD25$^{+/-}$ Tregs. **D)** A subset of Foxp3$^+$CD25$^{+/-}$ Tregs that became GD2$^+$ with RT+IC also showed significantly higher CD107a co-expression compared to GD2$^-$ Tregs, with elevated CD107a expression from RT+IC group restricted to GD2$^+$CD4s (**Fig. S6**). n=8/group; 2 independent experiments combined. **E)** Treatments including RT (RT alone, RT+anti-GD2 ("aGD2"), RT+IL2, RT+aGD2+IL2, or RT+IC) increased Foxp3$^+$CD4s compared to NT, whereas IC alone did not. **F)** Only regimens containing IC (IC alone or RT+IC) induced GD2$^+$CD4s in the TME. **G–H)** Similarly, IC-containing treatments (IC alone or RT+IC) drove increased GzmB$^+$Foxp3$^+$GD2$^+$ (**G**) and CD107a$^+$Foxp3$^+$GD2$^+$ (**H**) subsets within CD4s. n=6-12/group; 3 independent experiments combined. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001; Mean +/- SEM.

Immunocytokine treatment enables Murine Treg-Mediated Tumor Killing

We then considered whether $CD4^+$ T_{eff} cells were being converted to Tregs or whether Tregs were acquiring effector-like properties during RT+IC treatment. To address this, TDLNs from tumor-bearing mice were sorted into Treg cells ($CD4^+CD25^+CD127^-$) or "non-Tregs" ($CD4^+CD25^-CD127^{+/-}$) (**Fig. S6A**).[62] Analysis of these sorted populations confirmed that the Treg fraction was $Foxp3^+$, whereas non-Tregs were $Foxp3^-$ (**Fig. S6A**) as CD127 expression inversely correlates with Foxp3.[63,64] Sorted Tregs were then incubated with B78-D14 tumor cells in media for 24hrs and analyzed by flow cytometry (**Fig. 6A**). Consistent with observations in CD4s (**Fig. 4I**), $CD25^+Foxp3^+CD4s$ did not express GD2 prior to co-culture with B78-D14s (pre-incubation; **Fig. 6B**), but GD2 expression was significantly increased on $CD25^+Foxp3^+CD4s$ post-24hr incubation (**Fig. 6B**). Furthermore, CD107a expression was significantly increased on $GD2^+CD25^+Foxp3^+CD4s$ compared to $GD2^-CD25^+Foxp3^+CD4s$ (**Fig. 6C**). Importantly,

after co-culture, Tregs maintained a CD25⁺Foxp3⁺ phenotype, and non-Tregs did not convert into Tregs (**Fig. S6B**).

We also monitored the interaction of the sorted Tregs with B78-D14s with or without IC treatment using IncuCyte live-cell imaging (**Fig. 6A**). Over the course of 18hrs following co-culture initiation, Tregs trafficked to and surrounded the tumor cells, leading to tumor cell death by 24hrs (**Fig. 6D-E; Video S1-2**). With IC included during treatment, we observed significant induction of B78-D14 killing, with an increased rate of apoptosis of B78-D14s observed within 2hrs post-coculture (**Fig. 6E**). Together with the *in vivo* observation that IC treatment increased expression of GD2 and CD107a on Tregs (**Fig 5B-D**), these results suggest that IC (anti-GD2 linked to IL-2) may promote close interactions between Tregs [via high-affinity IL-2 receptors (IL-2R) binding][65-68] and GD2⁺ tumor cells (via anti-GD2), enabling both tumor killing and acquisition of tumor-derived GD2 through trogocytosis.

Since we observed rapid tumor-cell killing by Tregs (<4hrs) in the presence of IC and noted that GD2 acquisition correlates with cytotoxicity, we next asked whether IC directly increased GD2 acquisition on the surface of Tregs. To test this, B78-D14 (GD2⁺) or B78-H1 (GD2⁻) tumor cells were co-incubated with T cells in the presence or absence of IC, and GD2 acquisition was assessed after 4hrs. T cells did not acquire GD2 in the absence of IC or when cultured with GD2⁻ B78-H1 cells. In contrast, IC treatment induced detectable GD2 acquisition when T cells were cultured with GD2⁺ B78-D14 cells (**Fig. 6F**).

Figure 6. Treg Killing of B78-D14 Tumors is Mediated via IC Treatment
A) Schematic of immune cells collected from the TDLN of untreated tumor-bearing mice; Tregs were sorted based on CD4⁺/CD127⁻/CD25⁺ and then incubated with B78-D14+mKate2 cells in a live cell imaging assay. **B)** Prior to incubation, Foxp3⁺CD25⁺ cells were GD2⁻, but some become GD2⁺ post-incubation with B78-D14s. **C)** The Foxp3⁺CD25⁺ cells that were GD2⁺ had increased CD107a expression as compared to GD2⁻ Foxp3⁺CD25⁺ cells after incubation with B78-D14. **D)** Over the course of 22hrs culture with IC, IncuCyte imaging showed that the sorted Tregs (small gray cells) interact with B78-D14+mKate2 cells (red signal) and cause apoptosis (increased green signal) on the B78-D14s. **E)** Sorted Tregs incubated with B78-D14+mKate2 cells showed increased killing (as measured by increased green signal of red tumor cells) when co-incubated with IC as compared to Tregs without IC, IC alone, or media alone. **F)** CD4s isolated from naïve mouse spleens and co-cultured with B78-D14 tumor cells showed increased in the percentage of GD2⁺Foxp3⁺CD4s after 4h of IC treatment compared with untreated controls, whereas CD4s co-cultured with B78-H1 (GD2⁻) tumor cells failed to acquire GD2 even in the presence of IC. n=3 mice/experiment; 2 independent experiments. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001; Mean +/- SEM.

Immunocytokine treatment enables human Treg-mediated tumor killing

To determine whether these findings translate to the human setting, T cells isolated from healthy donor human PBMCs were incubated *in vitro* with the GD2$^+$ human melanoma cell line, M21—which does expresses MHC-I and MHC-II with IFN-γ stimulation (**Fig S1A**)—for 4hrs under various treatments (**Fig. 7A**). No differences in the percentages of Foxp3$^+$CD4s were found across treatment groups (**Fig. S8A**). Single-agent treatment with anti-GD2 mAb or IL-2, or the combination of anti-GD2 mAb + IL-2, did not induce GD2 expression on CD4s (**Fig. 7B**), GD2$^+$Foxp3$^+$CD4s (**Fig. 7C**) or GD2$^+$CD107a$^+$ on Tregs (**Fig. 7D**). Conversely, IC treatment led to significant induction in GD2 expression on all CD4s (**Fig. 7B**), including Foxp3$^+$CD4s (**Fig. 7C**), as well as a marked increase in CD107a expression on GD2$^+$Foxp3$^+$CD4s (**Fig. 7D**). These effects were abrogated by IL-2R blockade with anti-CD25, confirming that the IC directly links Tregs to tumor cells via IL-2R engagement (**Fig. 7B-7D**). To further confirm that these effects occur through direct linkage of Tregs to tumor cells, we evaluated trogocytosis using a GD2/CD3-engager (anti-GD2 × anti-CD3 bispecific antibody, "aGD2×aCD3"). Similar to IC, aGD2×aCD3 treatment resulted in significantly increased expression of GD2 on CD4s and Foxp3$^+$CD4s (**Fig. 7E-F**) and enhanced CD107a expression on GD2$^+$Foxp3$^+$CD4s (**Fig. 7G**). These effects were not seen with the antibodies (anti-GD2mAb or anti-CD3mAb) given individually and were reduced by anti-CD3 blockade prior to coculture (**Fig. 7E–G, Fig S7B-E**).

Finally, to test whether human Tregs are cytotoxic to human GD2$^+$ tumors, we co-cultured LA-N-1 human neuroblastoma cells—which expresses MHC-I, but not MHC-II, with IFN-γ stimulation (**Fig S1A**)—with human Tregs in the presence of IC (**Fig. 7H**). Over a 10hr co-culture, only Tregs treated with IC were capable of killing LA-N-1 cells (**Fig. 7I**). Collectively, these results demonstrate that IC co-culture with GD2$^+$ tumor cells enables mouse and human Foxp3$^+$ Tregs to mediate effector-like cytotoxic functions—including GD2 trogocytosis and degranulation—while maintaining their regulatory phenotype, highlighting their capacity for functional adaptation in response to IC-mediated tumor engagement. Because LA-N-1 cells lack MHC-II expression, we next tested whether IC can bypass the requirement for tumor-cell MHC-II expression for GD2 acquisition on Tregs. After a 4hr co-culture of LA-N-1 cells with healthy donor CD4s treated with or without IC, IC treatment induced detectable GD2 acquisition on Tregs (**Fig. 7J**).

Figure 7. Treg Trogocytosis and Killing of Human Tumors
A) Schematic of T cells isolated from PBMCs were cocultured with human GD2$^+$ M21 melanoma cells in the presence of aGD2-mAb, IL2, aGD2-mAb+IL2, IC, IC+aCD25-mAb (to block the CD25 IL2 receptor), aCD3-mAb+aGD2-mAb, aCD3xaGD2 bispecific Ab, or aCD3xaGD2 bispecific Ab + aCD3-mAb (to block the CD3 receptor) for 4 hours and then analyzed via flow cytometry. **B-C)** Coculture with IC significantly increased the expression of GD2 on CD4s (**B**) as well as on Foxp3$^+$CD4s (**C**). **D)** Coculture with IC also significantly increased CD107a expression on GD2$^+$Foxp3$^+$CD4s. aCD25-mAb pretreatment of CD4s, prior to culturing with IC significantly reduced the IC induced GD2 expression on CD4s and Foxp3$^+$CD4s, as well as decreased CD107a on GD2$^+$Foxp3$^+$CD4s (**B,C,D**). **E-F)** Treatment with aCD3xaGD2 bispecific Ab significantly increased the expression of GD2 on CD4s (**E**) as well as on Foxp3$^+$CD4s (**F**); and aCD3xaGD2 bispecific Ab treatment also significantly increased CD107a expression on GD2$^+$Foxp3$^+$CD4s (**G**). These effects were significantly diminished by pretreatment with aCD3-mAb with aCD3xaGD2 bispecific Ab treatment, confirming dependence on CD3 engagement (**E-G**). (A compilation of these data graphed together can be found in **Fig. S7B-D**). 2 independent experiments. **H)** Schematic of PBMCs isolated from a healthy donor were sorted for Tregs by magnetic separation and incubated with human LA-N-1+NLSmkate2 neuroblastoma cells with or without IC and monitored via live cell imaging. **Fig. S7E** shows efficiency of Treg isolation. **I)** LA-N-1+NLSmkate2 cells express red fluorescent signal within their nuclei; an increase in red signal over time indicates proliferation, as seen in cocultures with Tregs+media, IC or media alone. In contrast, coculture with Tregs in the presence of IC led to a progressive loss of red fluorescence, indicating Treg-mediated killing of LA-N-1$^+$NLS-mKate2 cells. **J)** Healthy donor human CD4s co-cultured with LA-N-1 tumor cells showed increased in the percentage of GD2$^+$Foxp3$^+$CD4s after 4h of IC treatment compared with untreated cells.1 independent experiment. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001; Mean +/- SEM.

HUMAN T CELL ISOLATION and COCULTURE SCHEMA

A

GD2 TROGOCYTOSIS WITH IMMUNOCYTOKINE: anti-GD2 x IL-2

B   **C**   **D**

GD2 TROGOCYTOSIS WITH BISPECIFIC: anti-GD2 x anti-CD3

E   **F**   **G**

TREGS MEDIATE KILLING OF GD2+ NEUROBLASTOMA WITH IC

H   **I**   **J**

Discussion

External beam RT followed by IC induces curative responses in mice bearing established B78-D14 tumors.[25] RT not only triggers DNA damage and tumor cell death but also promotes immune activation within the TME. IC amplifies this response by stimulating infiltrating immune cells, including CD4$^+$ T cells. In the B78-D14 model—MHC-I–deficient but MHC-II–inducible—CD4s are the primary mediators of RT+IC responses. Direct tumor engagement by CD4s was evident via GD2 trogocytosis, with Foxp3$^+$ subsets (Tregs and CD25$^-$/Foxp3$^+$ cells) comprising ~1/3 of GD2$^+$CD4s and displaying antitumor effector activity. These findings demonstrate that regulatory populations can participate directly in tumor cytotoxicity.

CD4$^+$ T cells are functionally diverse, and their plasticity can be harnessed to drive antitumor responses following immunotherapy.[69-73] Prior work has demonstrated MHC-II dependent cytotoxic activity of CD4s against B16 melanoma.[72,74] Like B16, B78-D14 also express MHC-II under cytokine stimulation (e.g., IFN-γ). Unlike B16, B78-D14 do not express MHC-I, even after IFN-γ. In this context, CD4s take a lead role in the initial antitumor response and demonstrates the essential role of CD4s in memory (rechallenge) responses. Upon treatment, CD4s exhibited increased activation markers, degranulation molecules, and cytolytic granules. While CD8 and NK cells showed elevated perforin, they did not degranulate, and their activation depended on CD4-mediated signaling, as CD4 depletion abrogated TNF-α and IFN-γ production. Loss of tumor rejection and memory responses following CD4—but not CD8 or NK—depletion underscores CD4s as central orchestrators of RT+IC–mediated immunity.

Trogocytosis—the transfer of membrane components and associated antigens between cells—has been described as a means by which T cells dynamically acquire information from antigen-presenting or target cells.[58,75] GD2 acquisition by CD4s occurs, in part, through an MHC-II–dependent process. CD4s acquired GD2 after *in vitro* co-culture only when tumor cell lines expressed MHC-II, and GD2 acquisition was enhanced in memory CD4s. These findings are consistent with a model in which memory CD4s undergo enhanced trogocytosis upon re-encountering MHC-II–presented tumor antigens. This process likely reflects a dynamic mode of antigen engagement that enables CD4s to sustain cytokine production and cytolytic potential.[55,58,60,75] Beyond MHC-II–dependent processes, IC accelerated GD2 acquisition and cytotoxicity even when target cells lacked MHC-II, indicating an MHC-independent mechanism. *In vitro*, IC accelerated GD2 trogocytosis by CD4s and enhanced tumor killing, even when target cells lacked MHC-II. *In vivo*, during the antitumor response with RT+IC, a subset of Foxp3$^+$ Tregs infiltrating the TME acquired GD2, indicating direct contact with GD2$^+$ tumor cells. Given that IL-2 preferentially binds to and supports cells expressing the high-affinity IL-2 receptor—including Tregs[55,60]—the IC likely supports Treg persistence and engagement in the TME through IL-2Rα/βγ (or high-affinity IL-2R).[58,75]

The capacity of Tregs to mediate cytotoxicity has been debated. Choi *et al.* used anti-CD3 x anti-EGFRvIII bispecific antibody to report the cytotoxic activity of human Tregs against glioblastoma cells.[76] However, Koristka *et al.* reported that purified human Tregs did not exhibit detectable cytotoxic activity against tumor cells when engaged with several bispecific antibodies.[77] Our data clarify this by showing that Tregs demonstrate potent cytotoxic function when bridged to GD2$^+$ tumor cells via either anti-GD2×IL-2 IC (mouse and human Tregs) or an anti-CD3×anti-GD2 (human Tregs) bispecific engager. Under these conditions, Tregs acquired GD2 through trogocytosis and upregulated CD107a, indicating active degranulation, representing a previously unrecognized mechanism by which Tregs can exert direct tumoricidal activity.[76] Importantly, these effects were abrogated by IL-2Rα (for IC) or CD3 blockade (for the anti-CD3×anti-GD2 bispecific engager), confirming that direct physical linkage

between Tregs and tumor cells via high-affinity IL-2R or CD3 engagement is required. The prominence of Treg engagement in this process may reflect their constitutive expression of the high-affinity IL-2R, which we previously showed is required for IC-mediated bridging of immune cells to tumor cells.[65-68] In both mice and humans, resting Tregs uniquely express this receptor complex at high levels,[78] providing a mechanistic explanation for the robust GD2 acquisition, trogocytosis, and cytolytic activity we observed in Tregs relative to other CD4+ subsets.

Trogocytosis by Tregs has been described in the context of acquiring the molecules from antigen-presenting cells such as dendritic cells. [45, 79] In tumors, Treg cytotoxicity has primarily been associated with suppression of effector lymphocytes,[80] yet direct tumor cell killing by Tregs has remained largely anecdotal.[76,77] Here, we identify a previously unrecognized role for Tregs in antitumor immunity: following immune engagement, such as that induced by IC, Tregs can trogocytose tumor antigens and directly kill tumor cells. In our model of MHC-I–deficient, MHC-II-inducible tumor cells, a subset of CD4s mediate trogocytosis *in vivo* during the rechallenge-memory response, even without IC, supporting a mechanism in which CD4+ T cells directly kill tumor cells that upregulate MHC-II via MHC-II–restricted TCR recognition. Together, these findings demonstrate that RT+IC therapy reshapes the tumor immune landscape by engaging CD4+ T cells as both direct effectors and central coordinators of antitumor immunity. Through cytokine signaling, IL-2-driven activation, trogocytosis, and IC-mediated tumor engagement, Tregs and other CD4 subsets gain cytolytic competence and contribute to tumor clearance. These results highlight the functional plasticity of CD4+ T cells, illustrating how targeted cytokine delivery can repurpose classically suppressive populations toward productive antitumor immunity.[81]

Methods

Tumor cells:
Murine Cell Lines: B78-D14 melanoma cells were obtained from Ralph Reisfeld (Scripps Research Institute), and are derived from B78-H1 cells (obtained from Dr. Matthew Krummel, University of California San Francisco)[82], which were originally derived from B16 melanoma.[83-85] The B16F10 (B16) melanoma cell line was originally obtained from Dr. William Ershler (University of Wisconsin-Madison).[86] MC38 is a mouse colon adenocarcinoma cell line that was obtained from Brad St. Croix (NIH). NXS2 is a mouse neuroblastoma cell line that was obtained from Ralph Reisfeld (Scripps Research Institute). B78 cell lines, B16 cells and MC38 cells were grown in RPMI 1640 (Mediatech) supplemented with 10% heat-inactivated FBS, 2 mmol/L l-glutamine, 100 U/mL penicillin, and 100 µg/mL streptomycin. B78-D14 cells lack melanin, but were transfected with functional GD2/GD3 synthase to GD2.[83,84] Periodic treatment with Hygromycin B (50 µg/mL) and G418 (400 µg/mL) was used to maintain GD2 expression of B78-D14 cells. 9464D-GD2 is a GD2-expressing cell line derived from 9464D as previously described.[87] The parental 9464D cell line is a MYCN-driven neuroblastoma cell line derived from TH-MYCN transgenic mice[88], and a subclone of the parental 9464D cell line selected for that does not express MHCI even when stimulated with interferon-γ.[87] 9464D-GD2 and NXS2 cells were grown in DMEM supplemented with 10% fetal bovine serum, 2 µM L-glutamine, 1 mM sodium pyruvate, 1X MEM non-essential amino acids, 100 U/mL penicillin, and 100 µg/mL streptomycin. Periodic treatment with puromycin (6 µg/mL) and blasticidin (7.5 µg/mL) was used to maintain GD2 expression of 9464D-GD2 cells.
Human Cell Lines: M21 melanoma cells were obtained from Ralph Reisfeld (Scripps Research Institute). LA-N-1 neuroblastoma cells were obtained from Patrick Reynolds (Texas Tech University, Lubbock TX). M21 and LA-N-1 cells were grown in RPMI 1640 (Mediatech) supplemented with 10% heat-inactivated FBS, 2 mmol/L l-glutamine, 100 U/mL penicillin, and 100 µg/mL streptomycin. Mycoplasma testing via PCR was routinely performed.

Immune Cell/Tumor Cell Coincubation and cytotoxicity evaluation:
Tumor Draining Lymph Node/B78-D14 Cocultures
B78-D14 melanoma cells transduced to express NLS-mKate2, B78-D14+NLS-mKate2, were plated at 1000 cells/well in 20 µL/well in 384-well plates (Corning, #353961) and incubated at 37 °C for 24 h. The following day, lymph nodes (LNs) were harvested from mice that had flank B78-D14 tumors that had not received treatment. Immune cells were isolated from the tumor draining LNs by pressing the LNs through a 70µm cell strainer into a 50 mL conical tube using the plunger from a 3ml syringe. Tregs (CD4+CD25+CD127−) or non-Tregs (CD4+CD25+/−CD127+/−) were isolated using BD FACSAria II cell sorters. Sorted cells were pelleted, resuspended in RPMI 1640 supplemented with 10% heat-inactivated FBS and 2 mmol/L L-glutamine, and plated at 20,000 cells/well in 20 µL together with the B78-D14+NLS-mKate2 cells. To assess cytotoxicity, Annexin V–CF®488A Conjugate (Biotium, #29005R) was added to each well according to the manufacturer's instructions (final volume, 20 µL). Wells were then brought to a total volume of 80 µL by adding either 20 µL of media alone or media containing immunocytokine (IC, 50 ng/mL final concentration). Plates were incubated for 24 h at 37 °C in an IncuCyte SX5 live-cell imaging system.

Human Immune Cell/Melanoma Cocultures
LA-N-1 human neuroblastoma transduced to express nuclear-localized mKate2 (NLS-mKate2), LA-N-1 +NLS-mKate2 were plated at 1,000 cells/well in 20 µL/well in 384-well plates (Corning, #353961) and incubated at 37 °C for 24 h. The following day, peripheral blood mononuclear cells (PBMCs) were

isolated from healthy donor whole blood by Ficoll density gradient centrifugation. PBMCs were resuspended in RPMI 1640 supplemented with 10% heat-inactivated FBS and 2 mmol/L L-glutamine. Treg cells were isolated using the EasySep Human Treg Cell Isolation kit (Stemcell, #100-1136) per the manufacturers protocol and plated at 50,000 cells/well in 20 µL together with the LA-N-1 +NLS-mKate2 cells. Wells were then brought to a total volume of 60 µL by adding either 20 µL of media alone or media containing immune complex (IC, 50 ng/mL final concentration). Cells were monitored on an IncuCyte SX5 live cell imager.

T Cell Coincubation for Flow Cytometry Analysis:
Human Melanoma Cocultures
M21 cells were plated at 10,000 cells/well in 50µL/well in a 96-well plate and incubated 24 h at 37 °C. After 24h, PBMCs were isolated from healthy donor whole blood by Ficoll density gradient centrifugation. PBMCs were resuspended in RPMI 1640 supplemented with 10% heat-inactivated FBS and 2 mmol/L L-glutamine. T cells were isolated using the EasySep Human T Cell Isolation kit (Stemcell, #17951) per the manufacturers protocol and plated at 50,000 cells/well in 50 µL together with the M21 cells. After 4 h, PBMCs were collected and assessed by flow cytometry.

Tumor Draining Lymph Node Cocultures for Flow Cytometry
Tumor cell lines (B78-D14, B78-H1, MC38, B16 or 9464D+GD2) cells were plated in 12 well plates at 200,000 cells/well in 1 ml standard culture media/cell line with 200U of IFN-γ and incubated 24 h at 37 °C. After 24h, media was replaced with fresh media without IFN-γ. T cells were isolated as above from TDLN from naïve mice, untreated tumor-bearing mice, tumor-bearing mice with RT+IC, or rechallenge memory-mice. T cells were suspended in RPMI 1640 supplemented with 10% heat-inactivated FBS + 0.05 mM β-Mercaptoethanol (final concentration) and co-incubated for 24hrs with tumor cell lines. After 24hrs, T cells were removed from plate, washed with PBS, and processed for flow cytometry.

T Cell Coincubation for Cytospin Analysis:
B78-D14+NLSmKate2 cells were plated in 12 well plates at 200,000 cells/well in 1 ml standard culture media/cell line with 200U of IFN-γ and incubated 24 h at 37 °C. After 24h, media was replaced with fresh media without IFN-γ. T cells were isolated as above from TDLN from naïve mice and suspended in RPMI 1640 supplemented with 10% heat-inactivated FBS + 0.05 mM β-Mercaptoethanol (final concentration) and co-incubated for 24hrs with B78-D14+NLSmKate2 cells. Following co-incubation, mkate2⁻/GD2⁺/CD4⁺ T cells were sorted and plated onto glass chamber slides for an additional 24 h at 37 °C. As controls, B78-D14+NLSmKate2 cells alone and T cells alone (not co-incubated) were also plated on glass chamber slides for 24 h at 37 °C. After incubation, slides were stained with Wright–Giemsa stain and imaged by light microscopy.

Isoplexis Single-Cell Cytokine Profiling:
Single-cell multiplex cytokine profiling was performed using the Isoplexis platform on LNs isolated from naïve mice or B78-D14-tumor bearing mice treated with NT or RT+IC treated mice on day 8 after the start of treatment. Immune cells were isolated from LNs by pressing the LNs through a 70µm cell strainer into a 50 mL conical tube using the plunger from a 3ml syringe. Isolated immune cells were washed with PBS and immediately processed for single-cell profiling using the IsoCode Single-Cell Adaptive Immune: Mouse T-cells kit (Isoplexis, ISOCODE-1004-4) per the manufacturers protocol. Data were analyzed using IsoSpeak software.

Animal Tumor Growth Studies:

C57BL/6NTac mice were obtained from Taconic Biosciences between ages of 5-7 weeks old. A/J mice were obtained from Jackson laboratories between ages of 5-7 weeks old. After arrival, mice were allowed to acclimate to the facility for 1 to 2 weeks prior to tumor cell injection.

Tumor Implantation and Tumor Growth Monitoring

2×10^6 B78 cells were injected intradermally into the right flank of C57BL/6NTac mice in a volume of 0.1 mL PBS.[89] Tumor growth was monitored via twice weekly caliper measurements. Tumor volume was determined using the calculation: width^2 x length/2. When tumors reached ~50-100 mm^3, mice were enrolled in studies for treatment and randomized into treatment groups. Investigators measuring tumor growth once enrolled in treatment groups were blinded to the treatment mice received. Mice were euthanized when either the length or the width of a tumor reached 20mm and tracked for survival.

Treatment Administration

Each randomized group in any single experiment had equivalent starting tumor volumes. Mice receiving RT therapy were immobilized using custom lead jigs that exposed only the dorsal right flank. External beam radiation therapy (single dose of 12Gy) was delivered to the right flank tumor surface on day 0 using an X-RAD 320 cabinet irradiator system (Precision X-Ray, North Branford, Connecticut, United States). Mice receiving IC (AnYxis Immuno-Oncology GmbH, Vienna, Austria) were injected intratumorally (IT) on days 5-9 with 50µg in 100µL volume. Mice receiving anti-GD2 were given dinutuximab (Unituxin) at 50mg/dose with or without IL-2 (Proleukin) at 75,000 U/dose in 100µL volume.

For immune cell depletion studies, CD8 T cells were depleted via anti-CD8 antibody (clone 2.43, BioXcell, # BE0061) at 50 µg in 0.5 mL PBS per dose on days -2, 2, 5, 8, 12, 15, 18, 22. CD4 T cells were depleted via anti-CD4 antibody (clone GK1.5, BioXcell, #BE0003-1) at 200 µg in 0.5 mL PBS per dose on days -2, 5, 12, 18. NK cells were depleted via anti-NK1.1 antibody (clone PK136, BioXcell, # BE0036) 200 µg in 0.5 mL PBS per dose on days -2, 2, 5, 8, 12, 15, 18, 22. Control mice received normal rat IgG antibody (Sigma-Aldrich) 200 µg in 0.5 mL PBS per dose on days -2, 2, 5, 8, 12, 15, 18, 22. For Treg depletion studies, mice were treated with anti-CTLA4 (clone 9D9, BRMS) to deplete Tregs at 200µg in 200ml PBS per dose. As a positive control, mice were depleted of CD4 T cells using anti-CD4 antibody (clone GK1.5, BioXcell, #BE0003-1) at 200 µg in 0.2 mL PBS per dose. For these studies depletion antibodies were administered on days -1, 2, 5, 8, 11. Flow cytometry of peripheral blood was performed to confirm antibody depleted was successful.

Tumor Rechallenge Studies

At least two months after treated mice were cured, tumor rechallenge studies were performed. For the B78 model, 2×10^6 B78-D14 cells were injected intradermally into the right abdomen of both cured / memory-mice and age-matched naïve mice. All animal work was performed under an animal protocol approved by the Institutional Animal Care and Use Committee at the University of Wisconsin, Madison. A maximum of 5 mice were housed per cage. Standard chow and water were provided *ad libitum*. Enrichment (nesting material and igloo) was provided to all cages. Male and female mice were used for experiments.

Single Cell RNA sequencing Studies:

scRNAseq sample preparation and acquisition

C57BL/6NTac mice bearing B78-D14 tumors were treated with no treatment or external beam radiation therapy (12 Gy, day 0) plus IC (50 µg dose, days 5-8). On day 8, 1-hour post-injection of IC, mice were euthanized via CO_2 and tumors were excised. Two tumors per treatment group were finely chopped and suspended in 2.35 mL of RPMI-1640 + Enzyme D (100 µL), R (50 µL) and A (12.5 µL) (Tumor Dissociation Kit, Miltenyi), transferred to a gentleMACS C Tube (Miltenyi) and placed on a

gentleMACS Octo Dissociator with Heaters. On the gentleMACs dissociator, program *m_imp* Tumor_02 was used, after which the tubes were then inverted and placed in a shaking incubator at 37°C for 40 minutes, and then returned to the gentleMACS dissociator and run on program *m_imp* Tumor_03. Dissociated tumors were filtered through a 40 μm cell strainer (Corning), washed with 15 mL of RPMI-1640, and centrifuged at 300xg for 10 minutes. After removal of the supernatant, live cells were purified from the tumor cell pellet using a Dead Cell Removal Kit (Miltenyi) according to the manufacturer's protocol. Cell viability was determined and confirmed to be >90%. The Chromium Next GEM Single Cell 5' Reagent Kits v2 (Dual Index) protocol from 10X Genomics was followed for scRNAseq processing. Quantity and quality of the RNA were assessed using High Sensitivity D5000 ScreenTape on an Agilent 4200 TapeStation System (UW Biotech Center). Around 10,000 cells per sample were captured for library preparation, and sequenced on an Illumina NovaSeq6000 (Novogene, Sacramento, CA). Two biological replicates were included in a single experiment.

scRNAseq data analysis
Raw reads were aligned to the mm10 reference genome together with UMI (unique molecular identifier) counting using the Cell Ranger pipeline (v3 10X Genomics). Data were filtered using DoubletFinder[101] to remove potential doublets. Further filtering includes only the cells with low mitochondria contents (≤10%) and more than 200 genes covered by the mapping. To integrate the scRNAseq, we used a fuzzy clustering-based integration method (Harmony method)[102] to account for potential technical variance across samples. Downstream analyses were based on Seurat single-cell analysis package[103] including: principal component analysis with standard deviation saturation elbow plot to select the optimal number of principal components, graph-based clustering based on 20 Principal Components using *FindCluster* with different resolution from 0.1 to 2 to justify the number of clusters based on representative markers overlaid in the hierarchical tree across different resolution (Clustree R package), differentially expression analysis using MAST[104] implemented in Seurat with the cutoff average log2FC 0.25, adjusted *p*-value cutoff of 0.01, and at least 20% of cell expressed the markers. Differentially expressed genes for each subset (major cell types in Fig 3A and CD4 T cell in Fig 3B) were used to annotate the cell types, including *Sox10* to identify tumor cells, *Cd4* for CD4+ T cells, *Foxp3* for Tregs, *Cd8a* for CD8+ T cells, *Ncr1* for NK cells, *Fn1* for fibroblasts, *Pecam1* for endothelial cells, *Csf3r* for neutrophils, *Csf1r* for macrophages, and *Itgax* as well as *Flt3* for dendritic cells. Visualization with dot plots and violin plots were done using Seurat in R (v4.4.2) and Bioconductor platform.

Single-cell RNA-seq data have been deposited at GEO and are publicly available (GEO code: GSE310495). Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Flow Cytometry Sample Preparation and Acquisition:
Tumor Disaggregation and Flow Cytometry
On day 8 post-treatment, tumors were collected from mice. Tumors were manually disaggregated in petri dishes by chopping into ~1 mm pieces using a scalpel. Tumors were transferred to 50 mL conical tubes containing 5 mL of complete RPMI (Gibco), 2.5 mg of collagenase type IV, 250 μg of DNase, and 1X protein transport inhibitor (PTI, eBioscience). Samples were further disaggregated by placing on a heated shaker at 37°C for 45 minutes. Final single cell suspensions of tumor samples were obtained by passing through a 70 μm filter. Separately, a spleen was collected from a mouse as a control sample for staining. Immune cells were isolated from a spleen by pressing the spleen through a 70μm cell strainer into a 50 mL conical tube using the plunger from a 3ml syringe. Splenocytes were washed with 10ml of PBS, followed by red blood cell lysis. For red blood cell lysis, splenocytes were resuspended in 1ml of sterile water with gentle vortexing and incubated for 10 sec. After 30 seconds, 49ml of PBS was added and

cells were centrifuged to remove wash. Tumor samples and splenocytes were resuspended in flow buffer (PBS + 2% fetal bovine serum) plus 1X PTI, kept on ice, and protected from light for 4-6 hours. 3×10^6 cells from each individual tumor sample were added to individual 5 mL flow tubes. Leftover tumor cells were pooled and mixed with splenocytes and used for fluorescence minus one control. For the live/dead control, 1.5×10^6 cells were placed in a 1.5ml tube, frozen at -20C, thawed and then mixed with 1.5×10^6 live cells. Samples were resuspended in PBS plus PTI and stained with GhostRed780 Viability Dye (Tonbo Biosciences) for 30 minutes at 4°C, protected from light. After being washed with flow buffer plus PTI, samples were incubated with Fc block (TruStain FcX, BioLegend, #101319) for 5 minutes at room temperature to reduce nonspecific binding. Samples were stained with surface antibodies in brilliant stain buffer (BD Biosciences) for 30 minutes at 4°C, protected from light. After being washed with flow buffer plus PTI, samples were fixed with fixation buffer (eBioscience, #00-5523-00) for 30 minutes at 4°C, protected from light. After being washed with 1X permeabilization buffer, samples were stained with intracellular antibodies in permeabilization buffer overnight at 4°C, protected from light. The following day, samples were washed once with permeabilization buffer and once with flow buffer. Samples were acquired on an Attune NxT flow cytometer (Thermo Fisher). Antibody panels can be found in supplemental materials.

FlowJo Analysis
Data were standardized between timepoints using rainbow beads (Spherotech). Data were analyzed using FlowJo (v10.7.1) and gates were determined using fluorescent minus one control.

Statistical Analyses:
All details on statistical tests, replicates, etc. can be found within the figure captions.

For statistical analysis of tumor volume, killed tumor cells, and red object count, the time-weighted average (area under the volume-time curve, calculated using trapezoidal method) was calculated for each mouse tumor. Time-weighted averages using ANOVA of the time-weighted average of these values, then pairwise t-tests if the overall test was significant. The time-weighted averages were aggregated using the trapezoidal method. No p-value adjustments were made to adjust for inflated type 1 error rate. Survival curves were compared with pairwise log rank tests and response was evaluated by two-sample tests of proportions. Tumor response rate (i.e., tumor free vs. tumor bearing) was compared using a chi-square test then pairwise two-sample tests of proportions if the overall test was significant. Significance was assessed at the alpha = 0.05 level and no adjustments were made to account for inflated type 1 error rate. Analysis was conducted using R version 4.3.1 (2023-06-16). Differences in primary and rechallenge cures were calculated via Fisher exact test. Mouse analyses were carried out in R v4.1.2.

Flow cytometry results are represented as scatter dot plots showing mean (bar) ± standard deviation. Each dot represents a single mouse tumor. A one-way ANOVA was calculated with Šídák's multiple comparisons test, α = 0.05 (GraphPad Prism v10.4).

Single-cell RNA sequencing results are represented as UMAPs, heatmaps (relative gene expression), dot plots (average gene expression), and violin plots showing mean ± standard deviation.

Data Analysis and Schematic Design Software:
Data were graphed using GraphPad Prism version 10.01 (GraphPad Software, Inc.). Schematic diagrams of study designs were created in BioRender.com.

Acknowledgements

This work was supported by the Midwest Athletes Against Childhood Cancer, the Cancer Research Institute, the University of Wisconsin Carbone Cancer Center, the Hyundai Hope on Wheels Foundation, the End Kids Cancer Foundation, Stand Up 2 Cancer, The St. Baldrick's Foundation, Alex's Lemonade Stand Foundation, The Band of Parents Foundation, The CURE Childhood Cancer Foundation and by public health service grants R35-CA197078, and P01 CA250972 from the National Cancer Institute. NIH Shared Instrumentation Grants (1S10RR025483-01).

The authors thank the University of Wisconsin Carbone Cancer Center Flow Cytometry Laboratory, supported by P30 CA014520, for use of its facilities and services. We additionally thank Dr. Johannes Schoeneberg for his valuable assistance with video stabilization.

Conflicts of Interest

SDG is an employee of PineTree Therapeutics Inc. Cambridge, MA. There are no other conflicts of interest related to the work included in this manuscript.

Inclusion and Ethics Statement

All of the coauthors in this study agree that the research conducted is relevant, and they have fulfilled the criteria for authorship required by Nature Portfolio journals, as their participation was essential for the design and implementation of the study. The roles and responsibilities were agreed among collaborators ahead of the research. All of the research conducted was approved under approved Biological Safety, University of Wisconsin Institutional Review Board and Institutional Care and Animal Use Committee for the University of Wisconsin, Madison. This research was not severely restricted or prohibited in the setting of the researchers, and does not result in stigmatization, incrimination, discrimination or personal risk to participants. Local, regional and international research relevant to our study was taken into account in citations.

Declaration of generative AI and AI-assisted technologies

No AI or AI assisted technologies were used for any component of this research.

References

1. Siegel, R.L., Giaquinto, A.N., and Jemal, A. (2024). Cancer statistics, 2024. CA Cancer J Clin *74*, 12–49. 10.3322/caac.21820.
2. van Not, O.J., van den Eertwegh, A.J.M., Jalving, H., Bloem, M., Haanen, J.B., van Rijn, R.S., Aarts, M.J.B., van den Berkmortel, F., Blank, C.U., Boers-Sonderen, M.J., et al. (2024). Long-Term Survival in Patients With Advanced Melanoma. JAMA Netw Open *7*, e2426641. 10.1001/jamanetworkopen.2024.26641.
3. Ouwerkerk, W., van den Berg, M., van der Niet, S., Limpens, J., and Luiten, R.M. (2019). Biomarkers, measured during therapy, for response of melanoma patients to immune checkpoint inhibitors: a systematic review. Melanoma Res *29*, 453–464. 10.1097/CMR.0000000000000589.
4. Olbryt, M., Rajczykowski, M., and Widlak, W. (2020). Biological Factors behind Melanoma Response to Immune Checkpoint Inhibitors. Int J Mol Sci *21*. 10.3390/ijms21114071.
5. Cornel, A.M., Mimpen, I.L., and Nierkens, S. (2020). MHC Class I Downregulation in Cancer: Underlying Mechanisms and Potential Targets for Cancer Immunotherapy. Cancers (Basel) *12*. 10.3390/cancers12071760.
6. Versluis, J.M., Hoefsmit, E.P., Shehwana, H., Dimitriadis, P., Sanders, J., Broeks, A., and Blank, C.U. (2024). Tumor characteristics of dissociated response to immune checkpoint inhibition in advanced melanoma. Cancer Immunology, Immunotherapy *73*. 10.1007/s00262-023-03581-6.
7. Bai, R., Lv, Z., Xu, D., and Cui, J. (2020). Predictive biomarkers for cancer immunotherapy with immune checkpoint inhibitors. Biomark Res *8*, 34. 10.1186/s40364-020-00209-0.
8. Camelliti, S., Le Noci, V., Bianchi, F., Moscheni, C., Arnaboldi, F., Gagliano, N., Balsari, A., Garassino, M.C., Tagliabue, E., Sfondrini, L., and Sommariva, M. (2020). Mechanisms of hyperprogressive disease after immune checkpoint inhibitor therapy: what we (don't) know. Journal of Experimental & Clinical Cancer Research *39*. 10.1186/s13046-020-01721-9.
9. Krebs, F.K., Trzeciak, E.R., Zimmer, S., Ozistanbullu, D., Mitzel-Rink, H., Meissner, M., Grabbe, S., Loquai, C., and Tuettenberg, A. (2021). Immune signature as predictive marker for response to checkpoint inhibitor immunotherapy and overall survival in melanoma. Cancer Med *10*, 1562–1575. 10.1002/cam4.3710.
10. Callahan, M.K., Kluger, H., Postow, M.A., Segal, N.H., Lesokhin, A., Atkins, M.B., Kirkwood, J.M., Krishnan, S., Bhore, R., Horak, C., et al. (2018). Nivolumab Plus Ipilimumab in Patients With Advanced Melanoma: Updated Survival, Response, and Safety Data in a Phase I Dose-Escalation Study. J Clin Oncol *36*, 391–398. 10.1200/JCO.2017.72.2850.
11. Tsai, K.K., and Daud, A.I. (2015). Nivolumab plus ipilimumab in the treatment of advanced melanoma. J Hematol Oncol *8*, 123. 10.1186/s13045-015-0219-0.
12. Wolchok, J.D., Kluger, H., Callahan, M.K., Postow, M.A., Rizvi, N.A., Lesokhin, A.M., Segal, N.H., Ariyan, C.E., Gordon, R.A., Reed, K., et al. (2013). Nivolumab plus ipilimumab in advanced melanoma. N Engl J Med *369*, 122–133. 10.1056/NEJMoa1302369.
13. Maleki Vareki, S. (2018). High and low mutational burden tumors versus immunologically hot and cold tumors and response to immune checkpoint inhibitors. J Immunother Cancer *6*, 157. 10.1186/s40425-018-0479-7.
14. Maes, K., Mondino, A., Lasarte, J.J., Agirre, X., Vanderkerken, K., Prosper, F., and Breckpot, K. (2021). Epigenetic Modifiers: Anti-Neoplastic Drugs With Immunomodulating Potential. Front Immunol *12*, 652160. 10.3389/fimmu.2021.652160.
15. Bonaventura, P., Shekarian, T., Alcazer, V., Valladeau-Guilemond, J., Valsesia-Wittmann, S., Amigorena, S., Caux, C., and Depil, S. (2019). Cold Tumors: A Therapeutic Challenge for Immunotherapy. Front Immunol *10*, 168. 10.3389/fimmu.2019.00168.

16. Mendez, R., Aptsiauri, N., Del Campo, A., Maleno, I., Cabrera, T., Ruiz-Cabello, F., Garrido, F., and Garcia-Lora, A. (2009). HLA and melanoma: multiple alterations in HLA class I and II expression in human melanoma cell lines from ESTDAB cell bank. Cancer Immunol Immunother *58*, 1507–1515. 10.1007/s00262-009-0701-z.

17. Mendez, R., Rodriguez, T., Del Campo, A., Monge, E., Maleno, I., Aptsiauri, N., Jimenez, P., Pedrinaci, S., Pawelec, G., Ruiz-Cabello, F., and Garrido, F. (2008). Characterization of HLA class I altered phenotypes in a panel of human melanoma cell lines. Cancer Immunol Immunother *57*, 719–729. 10.1007/s00262-007-0411-3.

18. Donia, M., Andersen, R., Kjeldsen, J.W., Fagone, P., Munir, S., Nicoletti, F., Andersen, M.H., Thor Straten, P., and Svane, I.M. (2015). Aberrant Expression of MHC Class II in Melanoma Attracts Inflammatory Tumor-Specific CD4+ T- Cells, Which Dampen CD8+ T-cell Antitumor Reactivity. Cancer Res *75*, 3747–3759. 10.1158/0008-5472.CAN-14-2956.

19. Erbe, A., Feils, A., Felder, M., Heck, M., Hoefges, A., Carlson, P., Kravtsov, D., Morris, Z., Patel, R., Hank, J., et al. (2020). MHC-I and -II Expression Levels on Tumor cells and T cell Response to Immunotherapy in B78 Murine Melanoma Model. held in Virtual, (JITC).

20. Deffrennes, V., Vedrenne, J., Stolzenberg, M.C., Piskurich, J., Barbieri, G., Ting, J.P., Charron, D., and Alcaide-Loridan, C. (2001). Constitutive expression of MHC class II genes in melanoma cell lines results from the transcription of class II transactivator abnormally initiated from its B cell-specific promoter. J Immunol *167*, 98–106. 10.4049/jimmunol.167.1.98.

21. Johnson, D.B., Estrada, M.V., Salgado, R., Sanchez, V., Doxie, D.B., Opalenik, S.R., Vilgelm, A.E., Feld, E., Johnson, A.S., Greenplate, A.R., et al. (2016). Melanoma-specific MHC-II expression represents a tumour-autonomous phenotype and predicts response to anti-PD-1/PD-L1 therapy. Nat Commun *7*, 10582. 10.1038/ncomms10582.

22. Haabeth, O.A.W., Fauskanger, M., Manzke, M., Lundin, K.U., Corthay, A., Bogen, B., and Tveita, A.A. (2018). CD4(+) T-cell-Mediated Rejection of MHC Class II-Positive Tumor Cells Is Dependent on Antigen Secretion and Indirect Presentation on Host APCs. Cancer Res *78*, 4573–4585. 10.1158/0008-5472.CAN-17-2426.

23. Oliveira, G., Stromhaug, K., Cieri, N., Iorgulescu, J.B., Klaeger, S., Wolff, J.O., Rachimi, S., Chea, V., Krause, K., Freeman, S.S., et al. (2022). Landscape of helper and regulatory antitumour CD4(+) T cells in melanoma. Nature *605*, 532–538. 10.1038/s41586-022-04682-5.

24. Rakhmilevich, A.L., Felder, M., Lever, L., Slowinski, J., Rasmussen, K., Hoefges, A., Van De Voort, T.J., Loibner, H., Korman, A.J., Gillies, S.D., and Sondel, P.M. (2017). Effective Combination of Innate and Adaptive Immunotherapeutic Approaches in a Mouse Melanoma Model. J Immunol *198*, 1575–1584. 10.4049/jimmunol.1601255.

25. Morris, Z.S., Guy, E.I., Francis, D.M., Gressett, M.M., Werner, L.R., Carmichael, L.L., Yang, R.K., Armstrong, E.A., Huang, S., Navid, F., et al. (2016). In Situ Tumor Vaccination by Combining Local Radiation and Tumor-Specific Antibody or Immunocytokine Treatments. Cancer Res *76*, 3929–3941. 10.1158/0008-5472.CAN-15-2644.

26. Aiken, T.J., Komjathy, D., Rodriguez, M., Stuckwisch, A., Feils, A., Subbotin, V., Birstler, J., Gillies, S.D., Rakhmilevich, A.L., Erbe, A.K., and Sondel, P.M. (2022). Short-course neoadjuvant in situ vaccination for murine melanoma. J Immunother Cancer *10*. 10.1136/jitc-2021-003586.

27. Campbell, B.B., Light, N., Fabrizio, D., Zatzman, M., Fuligni, F., de Borja, R., Davidson, S., Edwards, M., Elvin, J.A., Hodel, K.P., et al. (2017). Comprehensive Analysis of Hypermutation in Human Cancer. Cell *171*, 1042–1056 e1010. 10.1016/j.cell.2017.09.048.

28. Lode, H.N., Xiang, R., Pertl, U., Forster, E., Schoenberger, S.P., Gillies, S.D., and Reisfeld, R.A. (2000). Melanoma immunotherapy by targeted IL-2 depends on CD4(+) T-cell help mediated by CD40/CD40L interaction. J Clin Invest *105*, 1623–1630. 10.1172/JCI9177.

29. Lasek, W., Mackiewicz, A., Czajka, A., Switaj, T., Gol b, J., Wiznerowicz, M., Korczak-Kowalska, G., Bakowiec-Iskra, E.Z., Gryska, K., Izycki, D., and Jakobisiak, M. (2000). Antitumor effects of the combination therapy with TNF-alpha gene-modified tumor cells and interleukin 12 in a melanoma model in mice. Cancer Gene Ther *7*, 1581–1590. 10.1038/sj.cgt.7700263.

30. da Silva, I.L., Montero-Montero, L., Ferreira, E., and Quintanilla, M. (2018). New Insights Into the Role of Qa-2 and HLA-G Non-classical MHC-I Complexes in Malignancy. Front Immunol *9*, 2894. 10.3389/fimmu.2018.02894.

31. Accolla, R.S., Ramia, E., Tedeschi, A., and Forlani, G. (2019). CIITA-Driven MHC Class II Expressing Tumor Cells as Antigen Presenting Cell Performers: Toward the Construction of an Optimal Anti-tumor Vaccine. Front Immunol *10*, 1806. 10.3389/fimmu.2019.01806.

32. Johnson, A.M., Bullock, B.L., Neuwelt, A.J., Poczobutt, J.M., Kaspar, R.E., Li, H.Y., Kwak, J.W., Hopp, K., Weiser-Evans, M.C.M., Heasley, L.E., et al. (2020). Cancer Cell-Intrinsic Expression of MHC Class II Regulates the Immune Microenvironment and Response to Anti-PD-1 Therapy in Lung Adenocarcinoma. J Immunol *204*, 2295–2307. 10.4049/jimmunol.1900778.

33. Heaton, A.R., Burkard, N.J., Gao, A., Hoefges, A., Feils, A.S., Burkard, S.K., Felder, M.A., Tsarovsky, N.W., Spiegelman, D.V., Lublin, G.M., et al. (2025). Divergent role of CD8 T cells with distinct metabolic phenotypes during curative radio-immunotherapy in hot versus cold tumors. bioRxiv. 10.1101/2025.10.24.684274.

34. Levitsky, H.I., Lazenby, A., Hayashi, R.J., and Pardoll, D.M. (1994). In vivo priming of two distinct antitumor effector populations: the role of MHC class I expression. J Exp Med *179*, 1215–1224. 10.1084/jem.179.4.1215.

35. Piedra-Quintero, Z.L., Wilson, Z., Nava, P., and Guerau-de-Arellano, M. (2020). CD38: An Immunomodulatory Molecule in Inflammation and Autoimmunity. Front Immunol *11*, 597959. 10.3389/fimmu.2020.597959.

36. Cibrian, D., and Sanchez-Madrid, F. (2017). CD69: from activation marker to metabolic gatekeeper. Eur J Immunol *47*, 946–953. 10.1002/eji.201646837.

37. Jennings, E., Elliot, T.A.E., Thawait, N., Kanabar, S., Yam-Puc, J.C., Ono, M., Toellner, K.M., Wraith, D.C., Anderson, G., and Bending, D. (2020). Nr4a1 and Nr4a3 Reporter Mice Are Differentially Sensitive to T Cell Receptor Signal Strength and Duration. Cell Rep *33*, 108328. 10.1016/j.celrep.2020.108328.

38. Carleton, M., Haks, M.l.C., Smeele, S.A.A., Jones, A., Belkowski, S.M., Berger, M.A., Linsley, P., Kruisbeek, A.M., and Wiest, D.L. (2002). Early Growth Response Transcription Factors Are Required for Development of CD4−CD8− Thymocytes to the CD4+CD8+ Stage. The Journal of Immunology *168*, 1649–1658. 10.4049/jimmunol.168.4.1649.

39. Bosselut, R., Zhang, W., Ashe, J.M., Kopacz, J.L., Samelson, L.E., and Singer, A. (1999). Association of the adaptor molecule LAT with CD4 and CD8 coreceptors identifies a new coreceptor function in T cell receptor signal transduction. J Exp Med *190*, 1517–1526. 10.1084/jem.190.10.1517.

40. Fassett, M.S., Jiang, W., D'Alise, A.M., Mathis, D., and Benoist, C. (2012). Nuclear receptor Nr4a1 modulates both regulatory T-cell (Treg) differentiation and clonal deletion. Proceedings of the National Academy of Sciences *109*, 3891–3896. 10.1073/pnas.1200090109.

41. Yang, L., Han, X., Wang, M., Zhang, X., Wang, L., Xu, N., Wu, H., Shi, H., Pan, W., Huang, F., and Wu, X. (2025). Early Growth Response Gene 1 Benefits Autoimmune Disease by Promoting Regulatory T Cell Differentiation as a Regulator of Foxp3. Research *8*. 10.34133/research.0662.

42. Wang, Y., Li, J., Nakahata, S., and Iha, H. (2024). Complex Role of Regulatory T Cells (Tregs) in the Tumor Microenvironment: Their Molecular Mechanisms and Bidirectional Effects on Cancer Progression. Int J Mol Sci *25*. 10.3390/ijms25137346.

43. Zhang, Q., Chikina, M., Szymczak-Workman, A.L., Horne, W., Kolls, J.K., Vignali, K.M., Normolle, D., Bettini, M., Workman, C.J., and Vignali, D.A.A. (2017). LAG3 limits regulatory T cell proliferation and function in autoimmune diabetes. Sci Immunol *2*. 10.1126/sciimmunol.aah4569.

44. Asano, T., Meguri, Y., Yoshioka, T., Kishi, Y., Iwamoto, M., Nakamura, M., Sando, Y., Yagita, H., Koreth, J., Kim, H.T., et al. (2017). PD-1 modulates regulatory T-cell homeostasis during low-dose interleukin-2 therapy. Blood *129*, 2186–2197. 10.1182/blood-2016-09-741629.

45. Moreau, J.M., Velegraki, M., Bolyard, C., Rosenblum, M.D., and Li, Z. (2022). Transforming growth factor-beta1 in regulatory T cell biology. Sci Immunol *7*, eabi4613. 10.1126/sciimmunol.abi4613.

46. Shevyrev, D., and Tereshchenko, V. (2020). Treg Heterogeneity, Function, and Homeostasis. Frontiers in Immunology *10*. 10.3389/fimmu.2019.03100.

47. Chougnet, C., and Hildeman, D. (2016). Helios-controller of Treg stability and function. Transl Cancer Res *5*, S338–S341. 10.21037/tcr.2016.07.37.

48. Sharma, M.D., Huang, L., Choi, J.H., Lee, E.J., Wilson, J.M., Lemos, H., Pan, F., Blazar, B.R., Pardoll, D.M., Mellor, A.L., et al. (2013). An inherently bifunctional subset of Foxp3+ T helper cells is controlled by the transcription factor eos. Immunity *38*, 998–1012. 10.1016/j.immuni.2013.01.013.

49. Kerdiles, Y.M., Stone, E.L., Beisner, D.L., McGargill, M.A., Ch'en, I.L., Stockmann, C., Katayama, C.D., and Hedrick, S.M. (2010). Foxo Transcription Factors Control Regulatory T Cell Development and Function. Immunity *33*, 890–904. 10.1016/j.immuni.2010.12.002.

50. Ouyang, W., Liao, W., Luo, C.T., Yin, N., Huse, M., Kim, M.V., Peng, M., Chan, P., Ma, Q., Mo, Y., et al. (2012). Novel Foxo1-dependent transcriptional programs control T(reg) cell function. Nature *491*, 554–559. 10.1038/nature11581.

51. Dixon, M.L., Leavenworth, J.D., and Leavenworth, J.W. (2021). Lineage Reprogramming of Effector Regulatory T Cells in Cancer. Front Immunol *12*, 717421. 10.3389/fimmu.2021.717421.

52. Knosp, C.A., Schiering, C., Spence, S., Carroll, H.P., Nel, H.J., Osbourn, M., Jackson, R., Lyubomska, O., Malissen, B., Ingram, R., et al. (2013). Regulation of Foxp3+ inducible regulatory T cell stability by SOCS2. J Immunol *190*, 3235–3245. 10.4049/jimmunol.1201396.

53. Ronchetti, S., Ricci, E., Petrillo, M.G., Cari, L., Migliorati, G., Nocentini, G., and Riccardi, C. (2015). Glucocorticoid-induced tumour necrosis factor receptor-related protein: a key marker of functional regulatory T cells. J Immunol Res *2015*, 171520. 10.1155/2015/171520.

54. Grosser, B., Kohlruss, M., Slotta-Huspenina, J., Jesinghaus, M., Pfarr, N., Steiger, K., Novotny, A., Gaida, M.M., Schmidt, T., Hapfelmeier, A., et al. (2020). Impact of Tumor Localization and Molecular Subtypes on the Prognostic and Predictive Significance of p53 Expression in Gastric Cancer. Cancers (Basel) *12*. 10.3390/cancers12061689.

55. Zhou, G., Ding, Z.C., Fu, J., and Levitsky, H.I. (2011). Presentation of acquired peptide-MHC class II ligands by CD4+ regulatory T cells or helper cells differentially regulates antigen-specific CD4+ T cell response. J Immunol *186*, 2148–2155. 10.4049/jimmunol.1002917.

56. Reed, J., Reichelt, M., and Wetzel, S.A. (2021). Lymphocytes and Trogocytosis-Mediated Signaling. Cells-Basel *10*. 10.3390/cells10061478.

57. Nakayama, M., Hori, A., Toyoura, S., and Yamaguchi, S.I. (2021). Shaping of T Cell Functions by Trogocytosis. Cells-Basel *10*. 10.3390/cells10051155.

58. Kim, J., Park, S., Kim, J., Kim, Y., Yoon, H.M., Rayhan, B.R., Jeong, J., Bothwell, A.L.M., and Shin, J.H. (2025). Trogocytosis-mediated immune evasion in the tumor microenvironment. Exp Mol Med *57*, 1–12. 10.1038/s12276-024-01364-2.

59. Osborne, D.G., and Wetzel, S.A. (2012). Trogocytosis results in sustained intracellular signaling in CD4(+) T cells. J Immunol *189*, 4728–4739. 10.4049/jimmunol.1201507.

60. Reed, J., and Wetzel, S.A. (2019). Trogocytosis-Mediated Intracellular Signaling in CD4(+) T Cells Drives T(H)2-Associated Effector Cytokine Production and Differentiation. J Immunol *202*, 2873–2887. 10.4049/jimmunol.1801577.

61. Hay, Z.L.Z., and Slansky, J.E. (2022). Granzymes: The Molecular Executors of Immune-Mediated Cytotoxicity. Int J Mol Sci *23*. 10.3390/ijms23031833.

62. Nettenstrom, L., Alderson, K., Raschke, E.E., Evans, M.D., Sondel, P.M., Olek, S., and Seroogy, C.M. (2013). An optimized multi-parameter flow cytometry protocol for human T regulatory cell analysis on fresh and viably frozen cells, correlation with epigenetic analysis, and comparison of cord and adult blood. J Immunol Methods *387*, 81–88. 10.1016/j.jim.2012.09.014.

63. Liu, W., Putnam, A.L., Xu-Yu, Z., Szot, G.L., Lee, M.R., Zhu, S., Gottlieb, P.A., Kapranov, P., Gingeras, T.R., Fazekas de St Groth, B., et al. (2006). CD127 expression inversely correlates with FoxP3 and suppressive function of human CD4+ T reg cells. J Exp Med *203*, 1701–1711. 10.1084/jem.20060772.

64. Seddiki, N., Santner-Nanan, B., Martinson, J., Zaunders, J., Sasson, S., Landay, A., Solomon, M., Selby, W., Alexander, S.I., Nanan, R., et al. (2006). Expression of interleukin (IL)-2 and IL-7 receptors discriminates between human regulatory and activated T cells. J Exp Med *203*, 1693–1700. 10.1084/jem.20060468.

65. Buhtoiarov, I.N., Neal, Z.C., Gan, J., Buhtoiarova, T.N., Patankar, M.S., Gubbels, J.A., Hank, J.A., Yamane, B., Rakhmilevich, A.L., Reisfeld, R.A., et al. (2011). Differential internalization of hu14.18-IL2 immunocytokine by NK and tumor cell: impact on conjugation, cytotoxicity, and targeting. J Leukoc Biol *89*, 625–638. 10.1189/jlb.0710422.

66. Gubbels, J.A., Gadbaw, B., Buhtoiarov, I.N., Horibata, S., Kapur, A.K., Patel, D., Hank, J.A., Gillies, S.D., Sondel, P.M., Patankar, M.S., and Connor, J. (2011). Ab-IL2 fusion proteins mediate NK cell immune synapse formation by polarizing CD25 to the target cell-effector cell interface. Cancer Immunol Immunother *60*, 1789–1800. 10.1007/s00262-011-1072-9.

67. Perez Horta, Z., Saseedhar, S., Rakhmilevich, A.L., Carmichael, L., Hank, J.A., Boyden, M., Gillies, S.D., and Sondel, P.M. (2019). Human and murine IL2 receptors differentially respond to the human-IL2 component of immunocytokines. Oncoimmunology *8*, e1238538. 10.1080/2162402X.2016.1238538.

68. Perez Horta, Z. (2016). Evaluation of Novel Immunocytokines That Preferentially Target High or Intermediate Affinity Interleukin-2 Receptors. Ph.D. (University of Wisconsin).

69. Montauti, E., Oh, D.Y., and Fong, L. (2024). CD4(+) T cells in antitumor immunity. Trends Cancer *10*, 969–985. 10.1016/j.trecan.2024.07.009.

70. Kravtsov, D.S., Erbe, A.K., Sondel, P.M., and Rakhmilevich, A.L. (2022). Roles of CD4+ T cells as mediators of antitumor immunity. Front Immunol *13*, 972021. 10.3389/fimmu.2022.972021.

71. Oh, D.Y., and Fong, L. (2021). Cytotoxic CD4(+) T cells in cancer: Expanding the immune effector toolbox. Immunity *54*, 2701–2711. 10.1016/j.immuni.2021.11.015.

72. Quezada, S.A., Simpson, T.R., Peggs, K.S., Merghoub, T., Vider, J., Fan, X., Blasberg, R., Yagita, H., Muranski, P., Antony, P.A., et al. (2010). Tumor-reactive CD4(+) T cells develop cytotoxic activity and eradicate large established melanoma after transfer into lymphopenic hosts. J Exp Med *207*, 637–650. 10.1084/jem.20091918.

73. Kim, S.I., Haerr, M.E., Al-Ghezi, M., Chen, C., Zhang, Y., Phipps, J.L., Arora, C., Gribbin, K.P., Kim, G., Markosyan, N., et al. (2025). CD4+ T cells mediate MHC-deficient tumor rejection and endothelial cell reprogramming. Cancer Immunol Res. 10.1158/2326-6066.CIR-24-1342.

74. Xie, Y., Akpinarli, A., Maris, C., Hipkiss, E.L., Lane, M., Kwon, E.K., Muranski, P., Restifo, N.P., and Antony, P.A. (2010). Naive tumor-specific CD4(+) T cells differentiated in vivo eradicate established melanoma. J Exp Med *207*, 651–667. 10.1084/jem.20091921.

75. Eisenberg, G., Uzana, R., Pato, A., Frankenburg, S., Merims, S., Yefenof, E., Ferrone, S., Peretz, T., Machlenkin, A., and Lotem, M. (2013). Imprinting of lymphocytes with melanoma antigens

acquired by trogocytosis facilitates identification of tumor-reactive T cells. J Immunol *190*, 5856–5865. 10.4049/jimmunol.1202879.

76. Choi, B.D., Gedeon, P.C., Herndon, J.E., 2nd, Archer, G.E., Reap, E.A., Sanchez-Perez, L., Mitchell, D.A., Bigner, D.D., and Sampson, J.H. (2013). Human regulatory T cells kill tumor cells through granzyme-dependent cytotoxicity upon retargeting with a bispecific antibody. Cancer Immunol Res *1*, 163. 10.1158/2326-6066.CIR-13-0049.

77. Koristka, S., Cartellieri, M., Arndt, C., Feldmann, A., Seliger, B., Ehninger, G., and Bachmann, M.P. (2015). Tregs activated by bispecific antibodies: Killers or suppressors? Oncoimmunology *4*, e994441. 10.4161/2162402X.2014.994441.

78. Mitchell, D.A., Cui, X., Schmittling, R.J., Sanchez-Perez, L., Snyder, D.J., Congdon, K.L., Archer, G.E., Desjardins, A., Friedman, A.H., Friedman, H.S., et al. (2011). Monoclonal antibody blockade of IL-2 receptor alpha during lymphopenia selectively depletes regulatory T cells in mice and humans. Blood *118*, 3003–3012. 10.1182/blood-2011-02-334565.

79. Tekguc, M., Wing, J.B., Osaki, M., Long, J., and Sakaguchi, S. (2021). Treg-expressed CTLA-4 depletes CD80/CD86 by trogocytosis, releasing free PD-L1 on antigen-presenting cells. Proc Natl Acad Sci U S A *118*. 10.1073/pnas.2023739118.

80. Cao, X., Cai, S.F., Fehniger, T.A., Song, J., Collins, L.I., Piwnica-Worms, D.R., and Ley, T.J. (2007). Granzyme B and perforin are important for regulatory T cell-mediated suppression of tumor clearance. Immunity *27*, 635–646. 10.1016/j.immuni.2007.08.014.

81. Ward, N.C., Yu, A., Moro, A., Ban, Y., Chen, X., Hsiung, S., Keegan, J., Arbanas, J.M., Loubeau, M., Thankappan, A., et al. (2018). IL-2/CD25: A Long-Acting Fusion Protein That Promotes Immune Tolerance by Selectively Targeting the IL-2 Receptor on Regulatory T Cells. J Immunol *201*, 2579–2592. 10.4049/jimmunol.1800907.

82. Broz, M.L., Binnewies, M., Boldajipour, B., Nelson, A.E., Pollack, J.L., Erle, D.J., Barczak, A., Rosenblum, M.D., Daud, A., Barber, D.L., et al. (2014). Dissecting the tumor myeloid compartment reveals rare activating antigen-presenting cells critical for T cell immunity. Cancer Cell *26*, 638–652. 10.1016/j.ccell.2014.09.007.

83. Becker, J.C., Varki, N., Gillies, S.D., Furukawa, K., and Reisfeld, R.A. (1996). An antibody-interleukin 2 fusion protein overcomes tumor heterogeneity by induction of a cellular immune response. Proc Natl Acad Sci U S A *93*, 7826–7831. 10.1073/pnas.93.15.7826.

84. Haraguchi, M., Yamashiro, S., Yamamoto, A., Furukawa, K., Takamiya, K., Lloyd, K.O., Shiku, H., and Furukawa, K. (1994). Isolation of GD3 synthase gene by expression cloning of GM3 alpha-2,8-sialyltransferase cDNA using anti-GD2 monoclonal antibody. Proc Natl Acad Sci U S A *91*, 10455–10459. 10.1073/pnas.91.22.10455.

85. Silagi, S. (1969). Control of pigment production in mouse melanoma cells in vitro. Evocation and maintenance. J Cell Biol *43*, 263–274. 10.1083/jcb.43.2.263.

86. Rakhmilevich, A.L., Imboden, M., Hao, Z., Macklin, M.D., Roberts, T., Wright, K.M., Albertini, M.R., Yang, N.S., and Sondel, P.M. (2001). Effective particle-mediated vaccination against mouse melanoma by coadministration of plasmid DNA encoding Gp100 and granulocyte-macrophage colony-stimulating factor. Clin Cancer Res *7*, 952–961.

87. Aiken, T.J., Erbe, A.K., Zebertavage, L., Komjathy, D., Feils, A.S., Rodriguez, M., Stuckwisch, A., Gillies, S.D., Morris, Z.S., Birstler, J., et al. (2022). Mechanism of effective combination radio-immunotherapy against 9464D-GD2, an immunologically cold murine neuroblastoma. J Immunother Cancer *10*. 10.1136/jitc-2022-004834.

88. Norris, M.D., Burkhart, C.A., Marshall, G.M., Weiss, W.A., and Haber, M. (2000). Expression of N-myc and MRP genes and their relationship to N-myc gene dosage and tumor formation in a murine neuroblastoma model. Med Pediatr Oncol *35*, 585–589. 10.1002/1096-911x(20001201)35:6<585::aid-mpo20>3.0.co;2-p.

89.   Carlson, P.M., Mohan, M., Rodriguez, M., Subbotin, V., Sun, C.X., Patel, R.B., Birstler, J., Hank, J.A., Rakhmilevich, A.L., Morris, Z.S., et al. (2021). Depth of tumor implantation affects response to in situ vaccination in a syngeneic murine melanoma model. J Immunother Cancer *9*. 10.1136/jitc-2020-002107.

Supplementary Figures

Supplemental Figure 1. MHC-II expression on B78-D14 tumors correlates with response. A) Expression of MHC-I and MHC-II following 48-hour IFN-γ stimulation on murine tumor cell lines (B78-D14, MC38, NXS2, and 9464D+GD2) and human tumor cell lines (M21 and LA-N-1) used in this study. **B)** Tumor-bearing mice were treated with RT+IC. Tumors were collected for flow cytometry from 3 different experiments at time points in which tumor growth differentiation between mice responding (tumors continued to shrink) vs non-responding (tumors began to grow) was observed (D29 for Expt. 1 & 2 and D53 for Expt.3). **C)** MHC-II expression was assessed on tumor cells by gating on CD45⁻ followed by gating on GD2⁺ for MHC-II expression based on FMO gating. **D)** MHC-II expression graphs for Responders (R) and Non-responders (NR) for each individual experiment (GD2 on X-axis, and. MHC-II on Y axis). **E)** When B78-D14 cells are cultured for 48h *in vitro* with IFN-γ, some of the cells show substantial upregulation of MHC-II, but none show any expression of MHC-I.

Supplemental Figure 2. Antitumor response to depletion of CD4 or CD8 T cells depends in part on MHC molecule expression on the tumor cells. A) Mice with B78 tumors depleted of CD8 T cells or NK cells still respond to treatment with RT+IC, but mice don't respond if depleted of CD4 T cells. **B**) A/J mice bearing NXS2 tumors respond to RT+IC similarly if not depleted or if depleted of CD4 or NK cells; if depleted of CD8 T cells these mice are less likely to respond. **C**) C57Bl/6 mice bearing MC38 tumors respond to RT+IL2+anti-CTLA4, but if depleted of CD8 T cells they do not. TF = Tumor Free; TB = Tumor Bearing.[33]

Supplemental Figure 3. GD2/CD4 co-expressing cells are CD4 T cells that have GD2 expression. A)
Of the GD2[+]/CD4[+] cells identified in the RT+IC-treated TME following treatment with RT+IC (**Fig 4E**),
the majority were identified by imagestream flow cytometry as CD4s containing fragments of GD2 rather
than a GD2 cell in direct contact with a CD4 T cell (i.e., a doublet). **B**) Following a 24h *in vitro* coculture
of B78-D14s with T cells isolated from the TDLN of a naïve mouse, sorted GD2[+]CD4s were plated on glass
slides, and 24 hrs later stained with Geimsa (top photo) and found to be 6-7 μm in diameter. Separately,
pure B78-D14s cells obtained from culture were plated on glass slides and stained with Geimsa (bottom
photo), and found to be 18-21 μm in diameter. **C**) Sorted GD2+CD4s all appeared to be the same size
(~7μm) as lymphocytes, and were smaller than the diameter of isolated pure B78-D14 tumor cells (~19
μm). **D**) Of the GD2[+]/CD4[+] cells identified from the TME during memory-rechallenge response, the
majority were identified as CD4s with punctate GD2 as compared to doublets of a GD2[+] cell in direct
contact with a CD4 T cell.

**Supplemental Figure 4. Expression of CD107a on CD4 T cells correlates with the amount of GD2
expression on CD4 T cells.** As shown in Fig 4I&K, TDLN-CD4s from RT+IC-treated or untreated tumor-
bearing mice cocultured with B78-D14 cells for 24h showed acquistion of GD2 on CD4 cells after co-
culture and became CD107a[+]. The relationship of GD2 and CD107 expression levels were analyzed. **A**)
Increased expression of GD2 has a significant positive correlation with CD107a on CD4s, with RT+IC
treated samples having higher expression of both markers. **B**) Non-expression of GD2 on CD4s has a
significant negative correlation with the amount of expression of CD107a on CD4s. Each point represents
data from a separate mouse.

Supplemental Figure 5. Unbiased FlowSOM Clustering Identifies Treatment-Driven CD4[+] T-Cell Subsets and Cytotoxic Marker Expression After RT+IC. A) Unbiased cluster analysis identifies 8 populations based on expression levels of GD2, CD107a, GzmB, Foxp3 and CD25 combining data from untreated (NT) and treated (RT+IC) CD4s in the TME as plotted on UMAP_4LZK using FlowSOM clustering technique. The intensity of expression in the heat map in the right panel indicates the level of expression of each marker when all samples are combined. **B**) Each individual population is plotted on UMAP_4LZK. **C**) Each UMAP_4LZK shows where Not Treated CD4 T cells (top panel) and RT+IC CD4s (bottom panel) cluster based on expression levels of GD2, CD107a, GzmB, Foxp3 and CD25.**D**) GzmB is expressed on CD4 T cell CD25$^{+/-}$/Foxp3/GD2$^{+/-}$ subsets taken from the TME d7 after treatment with RT+IC. **E**) CD107a is expressed on CD4 T cell CD25$^{+/-}$/Foxp3/GD2$^{+/-}$ subsets taken from the TME d7 after treatment with RT+IC. 2way ANOVA with Tukey's multiple comparison test. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001; Mean +/- SEM.

Supplemental Figure 6. Phenotype of Tregs sorted for the *in vitro* assay. A) Tregs were sorted from TDLN from tumor-bearing (but untreated) mice, based on CD4$^+$CD25$^+$CD127$^-$, "NOT" Tregs were sorted based on CD4$^+$CD25$^+$CD127$^+$ or CD4$^+$CD25$^-$. Post-sort, ~92% of cells sorted as "NOT" Tregs were CD25$^-$Foxp3$^-$ with ~5% of sorted cells having CD25$^+$Foxp3$^-$ (right panel), and ~91% of cells sorted on as "Tregs" were CD25$^+$Foxp3$^+$ (bottom panel) with 9% of sorted cells being CD25-Foxp3+. **B**) Post-coculture with B78-D14 cells, CD25/Foxp3 expression phenotype of cells sorted on as "Tregs" maintains expression of CD25 and Foxp3. Cells sorted on as "Not Tregs" remain largely CD25$^-$Foxp3$^-$ with a small subset of cells expressing CD25$^+$ without co-expression of Foxp3.

Sup Fig 7. Coculture of human Tregs with tumor cells and IC or aGD2xaCD3 bispecific Ab influences Treg trogocytosis and CD107a expression. A-D) T cells isolated from human PBMCs cocultured with M21 human melanoma cells were treated in the presence of aGD2, IL2, aGD2 + IL2, IC, IC + aCD25 (to block the CD25 IL2 receptor), aCD3 + aGD2, aGD2xaCD3 bispecific Ab, or a aGD2xaCD3 bispecific Ab + aCD3 (to block the CD3 receptor) for 4 hours and analyzed via flow cytometry. **A**) The frequency of Foxp3+CD4s remained unchanged across treatment conditions. **B**) Only coculture with antitumor antibodies linked to immune cell–targeting domains (IC or aGD2×aCD3) increased the proportion of GD2+CD4s; this effect of IC was abrogated by anti-CD25 blockade and for aGD2xaCD3 was partially reduced by aCD3 blockade. The aGD2×aCD3 treatment produced the most GD2+CD4s. **C**) Similarly, coculture with IC and aGD2×aCD3 increased GD2 expression on Foxp3+CD4s, with aCD25 abrogating this effect for IC and aCD3 blockade slightly reducing this effect for aGD2xaCD3. IC treatment resulted in the highest GD2 expression on Foxp3+CD4s. **D**) Coculture with IC and aGD2×aCD3 also enhanced CD107a expression on GD2+Foxp3+CD4s, indicative of degranulation and cytotoxic potential; this IC effect was abrogated by aCD25 and this aGD2xaCD3 effect was modestly reduced by aCD3 blockade. **E**) Efficiency of Treg isolation from human PBMC by magnetic separation shown in **Fig. 7H&I**.