The protein turnover and trafficking of Chlamyopsin6 is regulated by IFT88 and IFT52 in the *Chlamydomonas reinhardtii*

Short running title: Intraflagellar transport-mediated trafficking of modular rhodopsins in *Chlamydomonas reinhardtii*

Kumari Sushmita[1#], Sunita Sharma[1,2#], Rajani Singh[1], Suneel Kateriya[1*]

[1]Laboratory of Optobiotechnology, School of Biotechnology, Jawaharlal Nehru University, New Delhi 110067, India.

[2]Department of Cellular Biology, University of Georgia, Athens 30602, Georgia, USA.

Kumarisushmita.sbt@gmail.com; sssharma100595@gmail.com; singh.rajani07@gmail.com

*To whom correspondence should be addressed: Suneel Kateriya, skateriya@jnu.ac.in

#The authors contributed equally to this work.

Orcid ID: Kumari Sushmita (0000-0001-9649-1446), Sunita Sharma (0000-0002-6531-7522), Rajani Singh (R.S. 0000-0002-1520-2794), Suneel Kateriya (0000-0001-5428-4297)

Highlights

- Chlamyopsin6 localizes in eyespot and flagella of *Chlamydomonas reinhardtii* and its trafficking is IFT-mediated.

- Chlamyopsin5 localizes in the eyespot and its trafficking is independent of IFT machinery.

- Disruption of IFT52 and IFT88 results in reduced Chlamyopsin6 protein content within the cell.

- The protein interactome suggests the crosstalk among photoreceptors to mediate light-triggered cellular responses.

Abstract

Microbial rhodopsin-based optogenetics has been widely applied to diverse mammalian and plant cell types for controlling membrane potential mediated responses. However, trafficking of optogenetically active protein to the desired subcellular organelle is still a major concern in optogenetic field. This could be resolved by studying the trafficking mechanism of optogenetically active protein in the native system. Current study is focused on the trafficking of two of the microbial rhodopsins named Chalmyopsin5 and Chlamyopsin6 in a green alga, *Chlamydomonas reinhardtii*. Chlamyopsin5 and Chlamyopsin6 are modular in nature and possess rhodopsin, histidine kinase, response regulator and cyclase domain in tandem. Immunolocalization of Chlamyopsin5 and Chlamyopsin6 in wild strain suggests their different subcellular localization; Chlamyopsin5 in eyespot and Chlamyopsin6 in flagella. Extensive immunocytochemistry of Chlamyopsin5 and Chlamyopsin6 was performed in different intraflagellar transport (IFT) components-defective strains of *Chlamydomonas* to dissect their trafficking mode to the destined subcellular compartment. Our results indicated the trafficking of Chlamyopsin5 to the eyespot to be independent of IFT machinery while Chlamyopsin6 to the flagella to be IFT dependent. Further, we demonstrate that IFT88 and IFT52 stabilizes Chlamyopsin6 and IFT20 interacts with Chlamyopsin6 in *Chlamydomonas*. Protein interactome of Chlamyopsin5 and Chlamyopsin6 indicate their role in nitrogen assimilation, gametogenesis and photoprotection in co-ordination with other photoreceptors. Collectively, our study enabled us to understand the targeting of Chlamyopsins to the subcellular compartment (eyespot and flagella). Further research in this direction is required to resolve the current challenge of targeting of optogenetic protein to desired subcellular compartment.

Keywords: Chlamyopsin5, Chlamyopsin6, *Chlamydomonas reinhardtii*, Intraflagellar transport, Immunolocalization

1. Introduction

Rhodopsins are membrane-embedded light-sensing proteins widely distributed in all the domains of life. The protein moiety in rhodopsin is bound to the light-sensing chromophore (retinal) via Schiff base linkage with the conserved lysine in the seventh helix (Ernst *et al.*, 2014). The isomeric form of retinal in the ground state broadly divides rhodopsin into two categories- animal-type rhodopsin and microbial-type rhodopsin. Animal-type rhodopsins are further categorized as vertebrate and invertebrate rhodopsins present in photoreceptor cells in vertebrate eyes and also in the brain of invertebrates. Animal-type rhodopsins have a major role in vision and circadian clock regulation (Ernst *et al.*, 2014). Microbial-type rhodopsins are also further categorized based on their functions as light-driven pumps, channels, sensory, or enzyme rhodopsins (Ernst *et al.*, 2014). While optogenetic application of animal-type rhodopsin is limited due to involvement of multiple components in signaling, however, microbial-type rhodopsins are widely used for controlling the membrane potential-driven responses in diverse cell types, including mammalian and plant cells (Govorunova *et al.*, 2017; Kojima, Shibukawa, and Sudo, 2020; Ochoa-Fernandez *et al.*, 2020; Reyer *et al.*, 2020; Zhou, Ding, Duan, *et al.*, 2021; Zhou, Ding, Gao, *et al.*, 2021). Despite of immense potential of rhodopsins as an optogenetic tool, the major challenges are their targeting to the required subcellular organelle or membrane surface (Zalocusky, Fenno and Deisseroth, 2013). This could be resolved to a greater extent by studying the trafficking mechanism of microbial-type rhodopsins. In this respect, trafficking of animal type rhodopsins is well studied (Wang and Deretic, 2014). Within the photoreceptor cells (rod and cone), rhodopsins are synthesized in the

endoplasmic reticulum (ER) of the inner segment and trafficked to the Golgi complex via a COP (coatomer) mediated pathway (Schmidt *et al.*, 2013). From the Golgi complex, rhodopsins are loaded on the Golgi-derived vesicles that integrates with membrane in the vicinity of the ciliary base (Wang and Deretic, 2014). From the ciliary base, rhodopsins are taken by intraflagellar transport (IFT) proteins along the connecting cilium to the outer segment, and from where they get integrated into the membrane where signaling takes place (Bhowmick *et al.*, 2009; Khanna, 2015).

IFT is the motor-driven bidirectional movement of proteins within the cilia, defects in which lead to ciliopathies (Rosenbaum and Witman, 2002; Cole, 2003; Webb, Mukhopadhyay, and Roberts, 2020). IFT is composed of two complexes- complex A and complex B (Rosenbaum and Witman, 2002; Cole, 2003; Webb, Mukhopadhyay, and Roberts, 2020). Complex A consists of 5-6 proteins along which cargoes are driven by dynein motor from the tip to the base of cilia/flagella (retrograde movement) (Rosenbaum and Witman, 2002; Cole, 2003; Webb, Mukhopadhyay and Roberts, 2020). Complex B consists of 12-16 proteins and with accompanying cargoes, is driven by kinesin motor from the base to the tip of cilia/flagella (Rosenbaum and Witman, 2002; Cole, 2003; Webb, Mukhopadhyay and Roberts, 2020). Among the IFT components, IFT52 and IFT88 are of prime importance since they form the core component of complex B required for assembly and stabilization of complex (Richey and Qin, 2012; Taschner *et al.*, 2014). IFT172 of complex B mediates the switch from anterograde to retrograde movement and reassembly at the ciliary tip (Pedersen *et al.*, 2005). IFT20, one of the anterograde IFT components is unique in terms of its localization and function. IFT20 localizes at the basal body as well as Golgi complex and is associated with post-Golgi trafficking of animal type rhodopsin and other membrane proteins (Follit *et al.*, 2006; Keady, Le and Pazour, 2011; Wang and Deretic, 2014). Though the signaling

and trafficking of animal-type rhodopsin is well studied, the trafficking of microbial-type rhodopsin is lacking.

We have selected two of the microbial type rhodopsins (Chamyopsin5 and Chlamyopsin6) (Awasthi *et al.*, 2020) belonging to the enzymerhodopsin category for studying the trafficking mechanism in *Chlamydomonas*. Chlamyopsin5 and Chlamyopsin6 (would be referred to as Cop5 and Cop6 hereafter) possess histidine kinase (HK), response regulator (RR), and cyclase domain in tandem and are therefore named as histidine kinase rhodopsin (HKR) or Cyclopsin (Cyclop). Biochemical characterization purified rhodopsin domain of Cop5 (HKR1) suggests its uniqueness in terms to spectral properties. Cop5 exists in two interconvertible forms: UV and blue light absorbing state. Upon blue light exposure, it converts to the UV-absorbing state and vice versa. This interconvertible state results from the isomerization of retinal (Luck *et al.*, 2012, 2015; Luck and Hegemann, 2017). Sequence analysis of Cop5 suggests that the cyclase domain lacks a few crucial residues responsible for its function and, therefore, is suspected to interact with the functional subunit, forming a heterodimer in the native system. In-vitro characterization of full-length Cop6 in *Xenopus leavis* suggests it to be a light-inhibited guanylate cyclase with supplementation of exogenous ATP (Tian *et al.*, 2018). Also, the signal proceeds through HK and RR since the replacement of critical residues led to the inhibition of cyclase activity (Tian *et al.*, 2018). However, in the native system, the signal transduction and function might be different. We studied the trafficking and function of Cop5 and Cop6 in *Chlamydomonas reinhardtii*. Extensive immunocytochemistry of Cop5 and Cop6 was performed in wild-type and mutant strains defective in IFT proteins (kinesin, dynein, IFT52, IFT88, and IFT172). Co-immunoprecipitation and protein interactome prediction via web-based software were performed to get insight into the interacting partners of Cop5 and Cop6. Insight into the trafficking mechanism of microbial-type rhodopsin

would help us to understand their subcellular targeting and overcome the current challenge of membrane or subcellular targeting of optogenetic proteins.

2. Materials and Methods:

2.1 *Chlamydomonas* strains and culture: All strains of *Chlamydomonas* were obtained from the Chlamydomonas resource center (University of Minnesota). All wild-type strains were maintained on TAP (Tris-Acetate-Phosphate) media and mutants on TAP supplemented with 0.1g/L arginine under 14hr light and 10hr dark cycles at 22˚C. All temperature-sensitive IFT mutants were grown at 22˚C (permissive temperature), and experiments were carried out by incubation at 33˚C (non-permissive temperature).

2.2 Immunoblotting: Proteins were first resolved on 6% SDS-PAGE and transferred on a nitrocellulose membrane. The protein blotted membrane was blocked with 5% skimmed milk in PBST (PBS supplemented with 0.1% Tween 20) at room temperature, incubated for 1 hr. The blocked membrane was incubated overnight with the primary antibody. The membrane was washed five times, each for five minutes, with PBST. After washing, the membrane was incubated with HRP-conjugated anti-goat IgG (secondary antibody) for 1hr followed by washing five times, each for five minutes with PBST. Membrane was developed by enhanced chemiluminescence (ECL) using solution A (100mM Tris-Cl, pH 8.5 + luminol + coumaric acid) and solution B (100mM Tris-Cl, pH 8.5 + hydrogen peroxide), and protein bands were captured on photographic films in the dark.

2.3 Immunolocalization: Coverslips were washed and coated with 0.01mg/ml poly-L-lysine. *Chlamydomonas* cells were seeded on poly-L-lysine-coated coverslips and kept for 15 minutes at room temperature. Algal cells seeded coverslips were then treated with freshly prepared 3.7%

paraformaldehyde solution in PBS for approximately 4 min. Coverslips were then dipped in ice-cold absolute ethanol solution and incubated at − 20 °C for 20 min with brief agitation. Cells were rehydrated with PBS containing 250mM NaCl, followed by four washings with PBST (PBS supplemented with 0.5% Triton X-100). The permeabilized *Chlamydomonas* cells were incubated overnight with freshly diluted primary antibody at 4°C. Then washed four times, each for 5mins with PBST, followed by incubation with FITC or Alexa Fluor conjugated secondary antibody (Invitrogen, USA) for 2 h at room temperature. After washing with PBST followed by PBS, coverslips were mounted onto the acid-washed glass slides using SlowFade® Gold antifade reagent (Molecular Probes, Invitrogen, USA) in an inverted manner (Awasthi *et al.*, 2016). Slides were visualized by the Olympus Fluoview 1000 confocal microscope.

2.4 Co-immunoprecipitation: *Chlamydomonas* cells grown for 3-4 days with 14hr light and 10hr dark cycles were harvested by centrifugation at 6000rpm for 10mins. Cells were lysed in IP-lysis buffer (RIPA) supplemented with 1mM PMSF, 1mM Sodium orthovanadate, and protease inhibitor cocktail (Sigma). Cells were fractionated by sonication, and cell debris was clarified by centrifugation at 13000rpm for 15 min. Supernatant obtained was pre-cleared by incubation with the pre-immune serum of the host organism, in which respective antibody was raised, and protein A Sepharose beads (Invitrogen) for 2hrs at 4°C on a gyratory shaker. 1ml of pre-cleared supernatant was incubated separately with the pre-immune serum and respective antibody and kept overnight at 4°C on gyration. Protein A Sepharose beads were added to it and kept for an additional 2hr at 4°C on gyration. Beads with antigen antigen-antibody complex were obtained by centrifugation at low speed for 10 min. Bead pellet was washed thrice with 1X PBS. Antigen-antibody complex was eluted from the beads by resuspending the beads in 0.1M glycine, pH-2.8, equal to the volume of bead pellet, followed by centrifugation at 2000 rpm for 4min. Supernatant was mixed with 2X

Laemmli dye and heated at 65°C for 30 mins. Proteins were resolved on SDS-PAGE and immunoblotted with the antibody of interest.

3. Results and Discussion

3.1. Spatial distribution of Chlamyopsin5 and Chlamyopsin6 differ within *Chlamydomonas* cell

The conserved domain architecture of Chlamyopsin5 and Chlamyopsin6 is mentioned in **Figure S1**. To study the localization and trafficking of Cop5 and Cop6 in *Chlamydomonas reinhardtii*, the antibodies against the peptides of the rhodopsin domains of Cop5 and Cop6 were used **(Figure S1)**. The antibodies were specific towards their antigens, and full-length Cop5 and Cop6 were detected in the cellular fractions of the *Chlamydomonas* cells **(Figure S1).** The spatial distribution of Cop5 and Cop6 differs within the *Chlamydomonas* cell. Though Cop5 possesses domains similar to Cop6, its localization was found to be restricted to the eyespot **(Figure 1)**. Cop6 behaves differently from Cop5 and localizes predominantly in flagella, with a fraction of it in the eyespot **(Figure 1)**. Dissimilar cellular destinations might be accountable for the different cellular functions of Cop5 and Cop6 within the *Chlamydomonas* cells. Since Cop6 is destined for flagella in *Chlamydomonas* cells, we sought to dissect the role of IFT machinery in the trafficking of Cop6 to the flagella.

3.2. Defects in the kinesin-2 motor protein in *Chlamydomonas* perturb the anterograde movement of Chlamyopsin6

Heterotrimeric motor protein kinesin-2 encoded by *fla8* in *C. reinhardtii* drives the cargoes loaded IFT complex within the cilia/flagella. Previous reports elucidated the dependency of ChR1 on kinesin-2 motor protein for its trafficking to the eyespot (Awasthi *et al.*, 2016). Therefore, we

investigated the involvement of kinesin-2 motor in the trafficking of Cop5 and Cop6 to the destined organelle (eyespot or flagella), via immunolocalization of Cop5 and Cop6 in a kinesin-2 motor subunit defective strain of *Chlamydomonas* (CC-1396). CC-1396 is a temperature-sensitive strain with a defect in the assembly and transport of anterograde particles; however, retrograde movement is comparable to the wild type. The temperature-sensitive mutant strains have phenotypes similar to wild-type at permissive temperature (22˚C), whereas at non-permissive temperature (33˚C), they show characteristic features of mutations. Cop5 localization to the eyespot is unperturbed in the kinesin motor mutant strain at non-permissive temperature **(Figure S2)** while Cop6 remained concentrated near the basal body and the flagellar tip **(Figure 2)**. Thus, mislocalization of Cop6 in the absence of functional kinesin-2 indicates the dependency of Cop6 on kinesin-2 motor protein for its trafficking to flagella. On the other hand, trafficking of Cop5 to the eyespot is independent of the kinesin-2 motor.

3.3. IFT88 and IFT52 stabilize the Chlamyopsin6 in *Chlamydomonas*

Interaction of IFT52 and IFT46 via their C-terminus serves as the docking site for other IFT-B proteins, IFT70, IFT88, and subcomplex (IFT81/74/27/25) (Taschner *et al.*, 2014). IFT52 is crucial for the assembly of the IFT-B core complex; IFT52 mutants fail to assemble the IFT-B core complex (Richey and Qin, 2012). IFT88 directly interacts with IFT52 and stabilizes the IFT-B core complex (Richey and Qin, 2012; Taschner *et al.*, 2014). Thus, IFT52 and IFT88 are essential for the assembly and stabilization of anterograde IFT complex. CC-477 and CC-3943 *Chlamydomonas* strains defective in IFT52 and IFT88, respectively, were used to study the localization of Cop5 and Cop6. CC-477 (IFT52) and CC-3943 (IFT88) mutant strains are devoid of flagella even at permissive temperature (22°C). Localization of Cop5 to the eyespot is unrestrained in IFT52 and IFT88 mutants of *Chlamydomonas,* implicating that Cop5 trafficking remains unaffected by IFT

disassembly and destabilization **(Figure S3)**. Surprisingly, no fluorescence for Cop6 was detected in IFT52 and IFT88 mutant strains, while in wild-type cells, fluorescence for Cop6 was detected in flagella **(Figure 3)**. In addition, a signal for acetylated tubulin was detected in both IFT52 and IFT88 mutants **(Figure 3)**. No detection of fluorescence for Cop6 in IFT52 and IFT88 led to the speculation of Cop6 degradation in the absence of IFT52 and IFT88 core components. Therefore, the protein amount of Cop6 was compared in CC-477 (IFT52 mutant) and CC-3943 (IFT88 mutant) with CC-124 (wild type) via immunoblotting. The protein amount was comparatively low in IFT52 and IFT88 mutant cell lysate than in the wild type **(Figure 4)**. The total protein amount was loaded equally in all wells. Reduced protein amount in IFT52 and IFT88 mutants with respect to wild type strengthens our speculation of Cop6 degradation due to the instability of IFT core components and the essential role of the complex for Cop6 trafficking to the destined organelle (flagella). However, further experiments would be required for the confirmation of the Cop6 degradation and the associated degradation pathway.

Taken together, we deduce that IFT core components IFT52 and IFT88 stabilize their cargo protein Cop6 and are crucial for its trafficking to flagella, whereas Cop5 stabilization and trafficking to the eyespot is independent of IFT52 and IFT88.

3.4. IFT172 regulates the transition of Chlamyopsin6 from anterograde to retrograde movement at the flagellar tip

IFT172 is a component of the anterograde IFT complex encoded by *fla11* (Pedersen *et al.*, 2005). IFT172 mediates the switch from anterograde to retrograde movement at the flagellar tip in *Chlamydomonas* (Pedersen *et al.*, 2005). Temperature-sensitive, flagellar assembly mutant strain of IFT172 (CC-1920) was used to study the localization of Cop6 to dissect the involvement of

IFT172 in the turnover of Cop6 at the flagellar tip. CC-1920 is a temperature-sensitive, flagellar assembly mutant of IFT172 encoded by *fla11* with replaced Leucine with Proline ($L_{1615}P$) at position 1615. The mutant strain shows wild-type-like phenotypes at 22°C, but the flagella get reabsorbed with time when incubated at 33°C. Cop6 sticks at the flagellar tip in CC-1920 strain at incubation of 30, 60, and 90 min at a non-permissive temperature (33°C) while evenly distributed within the flagella at a permissive temperature (22°C) **(Figure 5)**. Accumulation of Cop6 at the flagellar tip indicates the role of IFT172 in turnover at the flagellar tip (switch of anterograde to retrograde movement).

3.5. Chlamyopsin6 interacts with IFT20 in *Chlamydomonas*

IFT20 is a part of complex B (anterograde IFT components) and is unique among all IFT components regarding its distribution within the cell. All IFT components show their presence at the basal body, while IFT20 was found associated with the Golgi complex as well as the basal body. IFT20 mediates the trafficking of ATR from the Golgi to the basal body and within the cilia/flagella (Follit *et al.*, 2006; Keady, Le and Pazour, 2011). It interacts with Rabin8, a GEF (guanosine exchange factor) for GTPase (Rab8) involved in post-Golgi trafficking of ATR in vertebrates (Wang and Deretic, 2014). To dissect its role in the trafficking of modular sensory-type rhodopsin (Cop5 and Cop6) in *Chlamydomonas,* the antibody was raised against the full-length recombinant IFT20. Specificity of anti-IFT20 was confirmed by immunoblotting varying concentrations (50ng to 2000ng) of recombinant IFT20 protein. Anti-IFT20 was able to detect 50ng of recombinant IFT20. After confirmation of the specificity of the IFT20 antibody, it was used to detect the full-length IFT20 from the cell lysate of *Chlamydomonas.* Anti-IFT20 was able to detect the specific full-length IFT20 from the total cell lysate of *Chlamydomonas,* and no band

was detected with pre-immune serum. IFT20 was seen to be localized in the flagella of *Chlamydomonas* cells **(Figure S4)**.

Interactome prediction of IFT20 in *Chlamydomonas* suggests its interaction with other IFT components and, therefore, is an important component of IFT machinery. In the mammalian system, IFT20 was reported to form a link between IFT components and kinesin-2 by interacting with the motor protein (KIF3B) of kinesin-2 motor (Baker *et al.*, 2003). To assess the role of IFT20 in the trafficking of Cop6, the interaction of IFT20 and Cop6 was checked via co-immunoprecipitation. Anti-IFT20 was used as bait to precipitate Cop6 from the total cell lysate of *Chlamydomonas* cells. Cop6 precipitated with anti-IFT20, suggesting the interaction between IFT20 and Cop6 in the cellular system **(Figure 6)**. The interaction of IFT20 with Cop6 indicates that IFT20 might assist in the trafficking of Cop6. However, the assistance of IFT20 in trafficking Cop6 from the Golgi complex or the basal body is still a question.

3.6. *Dynein motor drives the retrograde movement of Chlamyopsin6 in Chlamydomonas*

The dynein motor is composed of heavy, light, and intermediate chains. Dynein motors drive the retrograde movement of IFT components and their cargo proteins. However, some proteins return to the cell body via diffusion, including kinesin-2 (Chien *et al.*, 2017). To explore the retrograde movement of Cop6 via the dynein motor, *Chlamydomonas* strain (CC-4424) with a defect in the kinesin motor and dynein heavy chain was used to study immunolocalization. This mutation affects the retrograde movement of cargo proteins at a non-permissive temperature (33°C), resulting in the accumulation of cargo proteins at the tip of cilia/flagella. Accumulation of Cop6 at the flagellar tip in dynein-defective *Chlamydomonas* mutants implicates that Cop6 requires the dynein motor for its movement back to the cell body from the flagella **(Figure 7)**. Localization of Cop5 in the

CC-4424 mutant strain was unperturbed at non-permissive temperature **(Figure S5)**; it localized to the eyespot, indicating the retrograde movement of Cop5 is independent of dynein motors.

3.7. Speculation of Chlamyopsin5 and Chlamyopsin6 function via protein-protein interaction

To gain insight into the pathways associated with Cop5 and Cop6 and their physiological role in *Chlamydomonas*, the interacting partners of Cop5 and Cop6 were predicted through the STRING database (Szklarczyk *et al.*, 2019). Addition of the sequence of regulatory protein (14-3-3) to the interactome established the link between Cop5 and phototropin and in turn with other photoreceptors (ChR1, ChR2, and cryptochrome), suggesting the complex cross-talk among photoreceptors might regulate signaling pathways (photosynthesis, photoprotection, eyespot development, sexual cycle or circadian rhythm). Therefore, a combined interactome was built based on the information from STRING software, literature, nano-LC-MS/MS lab data, and sequence analysis for the presence of a 14-3-3 binding motif **(Figure 8A)**. cGK2 and adenylyl cyclases obtained in the interactome are associated with gametogenesis (Wang, Pan and Snell, 2006). In interactome analysis, Cop5, Cop6, 14-3-3, cGK2, and adenylyl cyclase clustered together, indicating Cop5, Cop6, and 14-3-3 might have some role during gametogenesis. 14-3-3 may have other roles such as abiotic stress, role in trafficking, or proteasomal degradation (**Figure 8B**). Further, cytohubba analysis was also performed to identify the hub protein governing the network **(Figure S6, Table S1).**

4. **Conclusions**

In this study, we intended to study the IFT-mediated trafficking of two of the modular rhodopsins (Chlamyopsin5 and Chlamyopsin6) from *Chlamydomonas reinhardtii*. Despite being similar in domain organization, the spatial distribution of Cop5 and Cop6 in wild wild-type strain of *Chlamydomonas* differs. Cop5 localizes in the eyespot, whereas Cop6 localizes in the flagella of

the *Chlamydomonas* cells. Further, an *in vivo* localization study of Cop5 and Cop6 in temperature-sensitive IFT motor (kinesin and dynein) mutants of *Chlamydomonas* suggested that Cop6 is dependent on IFT motors for its transport to the flagella, while transport of Cop5 to the eyespot is independent of IFT motors. The combined result of immunofluorescence and immunoblotting of Cop6 in IFT88 and IFT52 mutant strains suggested the dependency of Cop6 on IFT88 and IFT52 for stability and trafficking to flagella in the *Chlamydomonas* cells. IFT172 mediates the reassembly of Cop6 for retrograde movement at the flagellar tip. Cop6 co-precipitated with IFT20 and might serve as a link between Cop6 and IFT proteins. Protein-protein interaction suggested the role of Cop5 and Cop6 in nitrogen assimilation, gametogenesis, phototaxis/photophobic response, and photoprotection via direct or indirect interaction with other photoreceptors (ChR1, ChR2, and phototropin), 14-3-3, and cGMP-dependent protein kinases (CGK2). Interactions should be further confirmed by pull-down assays. IFT-mediated trafficking and functional analysis of Cop5 and Cop6 would enable us to understand the selectivity mechanism of IFT machinery for microbial-type rhodopsin and their subcellular targeting. In addition to enhancing the optogenetic applicability of rhodopsin, such a study would also enable us to understand ciliopathies. In conclusion, our study indicates that trafficking of all microbial-type rhodopsins is not IFT-mediated. Some of the IFT components (e.g-IFT52 and IFT88) not only mediate the trafficking of Cop6 but also help in its stabilization. IFT20, directing the movement of animal photoreceptor from the Golgi complex, also interacts with Cop6. Protein interactome of Cop5 and Cop6 suggests the crosstalk among the photoreceptors, and would enable us to establish its link with many light-mediated biological pathways in *Chlamydomonas reinhardtii*.

Conflicts of Interest

The authors declare no competing interests.

Author Contributions

S.K. conceived the project. K.S. designed experiments under the guidance of S.K. and drafted the manuscript with contributions from S.K., S.S., and R.S. K.S. conducted immunofluorescence, immunoblotting, Co-immunoprecipitation experiments, and PPI interactome publication. S.S. contributed to immunoblotting, immunofluorescence, and image processing. R.S. contributed to immunoblotting, PPI network prediction, and image processing. All authors reviewed, revised, and approved the final manuscript.

Acknowledgement

SK is thankful to SERB ANRF for providing financial support (CRG/2021/00315) for the research project. Dr. Peeyush Ranjan and Dr. Mayanka Awasthi are acknowledged for the gift of Cop5 and Cop6 antibodies. DBT and CSIR are highly acknowledged for providing financial assistance to KS and SS, respectively. RS acknowledges the DBT-RA program (DBT-RA/2022/July/N/2560) for financial assistance.

References:

Awasthi, M. *et al.* (2016) 'The trafficking of bacterial type rhodopsins into the Chlamydomonas eyespot and flagella is IFT mediated', *Scientific Reports*, 6(1), p. 34646. doi: 10.1038/srep34646.

Awasthi, M. *et al.* (2020) 'Novel Modular Rhodopsins from Green Algae Hold Great Potential for Cellular Optogenetic Modulation Across the Biological Model Systems', *Life*, 10(11), p. 259. doi: 10.3390/life10110259.

Baker, S. A. *et al.* (2003) 'IFT20 Links Kinesin II with a Mammalian Intraflagellar Transport Complex that Is Conserved in Motile Flagella and Sensory Cilia', *Journal of Biological*

Chemistry, 278(36), pp. 34211–34218. doi: 10.1074/jbc.M300156200.

Bhowmick, R. *et al.* (2009) 'Photoreceptor IFT complexes containing chaperones, guanylyl cyclase 1 and rhodopsin', *Traffic*, 10(6), pp. 648–663. doi: 10.1111/j.1600-0854.2009.00896.x.

Chien, A. *et al.* (2017) 'Dynamics of the IFT machinery at the ciliary tip', *eLife*, 6, pp. 1–25. doi: 10.7554/eLife.28606.

Cole, D. G. (2003) 'The Intraflagellar Transport Machinery of Chlamydomonas reinhardtii', *traffic*, 4(13), pp. 435–442.

Engel, B. D. *et al.* (2012) 'The role of retrograde intraflagellar transport in flagellar assembly, maintenance, and function', *Journal of Cell Biology*, 199(1), pp. 151–167. doi: 10.1083/jcb.201206068.

Ernst, O. P. *et al.* (2014) 'Microbial and animal rhodopsins: Structures, functions, and molecular mechanisms', *Chemical Reviews*, 114(1), pp. 126–163. doi: 10.1021/cr4003769.

Follit, J. A. *et al.* (2006) 'The Intraflagellar Transport Protein IFT20 Is Associated with the Golgi Complex and Is Required for Cilia Assembly □', 17(September), pp. 3781–3792. doi: 10.1091/mbc.E06.

Govorunova, E. G. *et al.* (2017) 'Microbial Rhodopsins: Diversity, Mechanisms, and Optogenetic Applications', *Annual Review of Biochemistry*, 86(1), pp. 845–872. doi: 10.1146/annurev-biochem-101910-144233.

Keady, B. T., Le, Y. Z. and Pazour, G. J. (2011) 'IFT20 is required for opsin trafficking and photoreceptor outer segment development', *Molecular Biology of the Cell*. Edited by T. Stearns, 22(7), pp. 921–930. doi: 10.1091/mbc.e10-09-0792.

Khanna, H. (2015) 'Photoreceptor Sensory Cilium: Traversing the Ciliary Gate', *Cells*, 4(4), pp. 674–686. doi: 10.3390/cells4040674.

Kojima, K., Shibukawa, A. and Sudo, Y. (2020) 'The Unlimited Potential of Microbial Rhodopsins as Optical Tools.', *Biochemistry*, 59(3), pp. 218–229. doi: 10.1021/acs.biochem.9b00768.

Luck, M. *et al.* (2012) 'A Photochromic Histidine Kinase Rhodopsin (HKR1) That Is Bimodally Switched by Ultraviolet and Blue Light', *Journal of Biological chemistry*, 287(47), pp. 40083–40090. doi: 10.1074/jbc.M112.401604.

Luck, M. *et al.* (2015) 'Photochemical chromophore isomerization in histidine kinase rhodopsin', *FEBS Letters*, 589(10), pp. 1067–1071. doi: 10.1016/j.febslet.2015.03.024.

Luck, M. and Hegemann, P. (2017) 'The two parallel photocycles of the Chlamydomonas sensory photoreceptor histidine kinase rhodopsin 1☆', *Journal of Plant Physiology*, 217, pp. 77–84. doi: 10.1016/j.jplph.2017.07.008.

Ochoa-Fernandez, R. *et al.* (2020) 'Optogenetic control of gene expression in plants in the presence of ambient white light', *Nature Methods*, 17(7), pp. 717–725. doi: 10.1038/s41592-020-0868-y.

Pedersen, L. B. *et al.* (2005) 'Chlamydomonas IFT172 Is Encoded by FLA11, Interacts with CrEB1, and Regulates IFT at the Flagellar Tip', *Current Biology*, 15(3), pp. 262–266. doi: 10.1016/j.cub.2005.01.037.

Reyer, A. *et al.* (2020) 'Channelrhodopsin-mediated optogenetics highlights a central role of depolarization-dependent plant proton pumps', *Proceedings of the National Academy of*

Sciences, 117(34), pp. 20920–20925. doi: 10.1073/pnas.2005626117.

Richey, E. A. and Qin, H. (2012) 'Dissecting the Sequential Assembly and Localization of Intraflagellar Transport Particle Complex B in Chlamydomonas', *PLoS ONE*. Edited by D. Zhou, 7(8), p. e43118. doi: 10.1371/journal.pone.0043118.

Rosenbaum, J. L. and Witman, G. B. (2002) 'Intraflagellar transport', *nature reviews Molecular cell biology*, 3(11), pp. 813–825. doi: 10.1038/nrm952.

Schmidt, K. *et al.* (2013) 'Early stages of retinal development depend on Sec13 function', *Biology Open*, 2(3), pp. 256–266. doi: 10.1242/bio.20133251.

Szklarczyk, D. *et al.* (2019) 'STRING v11: protein – protein association networks with increased coverage , supporting functional discovery in genome-wide experimental datasets', *Nucleic Acids Research*, 47, pp. D607–D613. doi: 10.1093/nar/gky1131.

Taschner, M. *et al.* (2014) 'Crystal structures of IFT70/52 and IFT52/46 provide insight into intraflagellar transport B core complex assembly', *Journal of Cell Biology*, 207(2), pp. 269–282. doi: 10.1083/jcb.201408002.

Tian, Y. *et al.* (2018) 'Two-component cyclase opsins of green algae are ATP-dependent and light-inhibited guanylyl cyclases', *BMC Biology*, 16(1), p. 144.

Wang, J. and Deretic, D. (2014) 'Molecular complexes that direct rhodopsin transport to primary cilia', *Progress in Retinal and Eye Research*, 38(2), pp. 1–19. doi: 10.1016/j.preteyeres.2013.08.004.

Wang, Q., Pan, J. and Snell, W. J. (2006) 'Intraflagellar Transport Particles Participate Directly in Cilium- Generated Signaling in Chlamydomonas', *Cell*, 125(3), pp. 549–562. doi:

10.1016/j.cell.2006.02.044.

Webb, S., Mukhopadhyay, A. G. and Roberts, A. J. (2020) 'Intraflagellar transport trains and motors: Insights from structure', *Seminars in Cell & Developmental Biology*, 107(February), pp. 82–90. doi: 10.1016/j.semcdb.2020.05.021.

Zalocusky, K. A., Fenno, L. E. and Deisseroth, K. (2013) '2 Current challenges in optogenetics', in *Optogenetics*. DE GRUYTER, pp. 23–34. doi: 10.1515/9783110270723.23.

Zhou, Y., Ding, M., Duan, X., *et al.* (2021) 'Extending the Anion Channelrhodopsin-Based Toolbox for Plant Optogenetics', *Membranes*, 11(4), p. 287. doi: 10.3390/membranes11040287.

Zhou, Y., Ding, M., Gao, S., *et al.* (2021) 'Optogenetic control of plant growth by a microbial rhodopsin', *Nature Plants*, 7(2), pp. 144–151. doi: 10.1038/s41477-021-00853-w.

Figures:

(A)

Figure 1: Immunolocalization of Cop5 and Cop6 in *C. reinhardtii* cells. (A) Immunostaining of cells

was done using anti-acetylated tubulin (green), anti-Cop5 (Red). The secondary antibody for acetylated

tubulin and Cop5 is Alexa 488 and Alexa 546 conjugated anti-mouse IgG and anti-rabbit IgG, respectively.
(B) Immunostaining of cells was done using anti-acetylated tubulin (Red). Fluorescence of Cop6 (Green)
was obtained with anti-Cop6. The secondary antibody for acetylated tubulin and Cop6 is Alexa 647 and
Alexa 488 conjugated anti-mouse IgG and anti-goat IgG, respectively. Scale bar=2 μm in single cell and
scale bar = 5 μm in group of cells.

Figure 2: Immunolocalization of Cop6 in kinesin motor mutant strain (CC-1396) of *C. reinhardtii*
cells. Immunostaining of Cop6 was done using Anti-Cop6 (Green). The secondary antibody for Cop6 is
Alexa 488-conjugated anti-goat IgG. The first channel depicts the fluorescence specific for Cop6, second

for acetylated tubulin and the third panel depicts overlay of the first two panels with DIC. Scale bar=2 μm

in single cell and scale bar = 5 μm in group of cells.

Figure 3: Immunolocalization of Cop6 in IFT52, IFT88, and CEP290 mutant strain of *C. reinhardtii*

cells. The first panel represents the signal specific for Cop6 with the primary antibody anti-Cop6. Secondary

labelling was done with Alexa 488 conjugated anti-goat IgG. The second panel represents the signal for

acetylated tubulin, secondary labelling was done with Alexa 647 conjugated anti-mouse IgG. The third panel represents the overlay of the first two panels with DIC. Scale bar=2 μm.

Figure 4: Comparison of the expression profile of Cop5 and Cop6 in wild type versus IFT52 and IFT88 mutants. (A) Immunoblot of cell lysate of the wild-type strain, IFT52, and the IFT88 mutant strain with anti-Cop6. (B) Represents the relative band intensity of Cop6 in wild-type strain, IFT52, and IFT88 mutant strain. Value represents mean S±3 *P < 0.05, **P < 0.001.

Figure 5: Immunolocalization of Cop6 in IFT172 mutant strain (CC-1920) of *C. reinhardtii* cells.

Immunostaining of cells was performed using anti-acetylated tubulin (Red) and anti-Cop6 (Green). The secondary antibody for acetylated tubulin and Cop6 was Alexa 647 and Alexa 488 conjugated anti-mouse IgG and anti-goat IgG, respectively. Scale bar=2 µm.

Figure 6: Co-immunoprecipitation of Cop6 with IFT20 antibody. Co-immunoprecipitation of Cop6 with IFT20. Anti-IFT20 (IP: IFT20) and pre-immune serum (IP: PIS) of a goat in which antibody against IFT20 was raised were incubated with the cell lysate of *Chlamydomonas*, followed by an immunoprecipitation protocol. Immunoblotting was performed with anti-Cop6. The band at the size of 250 kDa was detected in IP with anti-IFT20. The band at the corresponding position was also observed in the cell lysate utilized for IP. No band was detected in IP with pre-immune serum, reflecting the specificity of the band in IP with anti-IFT20. An equal amount of heavy chain in IP with anti-IFT20 and pre-immune serum reflects the loading control.

Figure 7: Immunolocalization of Cop6 in kinesin and dynein heavy chain motor double mutant strain (CC-4424) of *C. reinhardtii* cells. Immunostaining of cells was performed using anti-acetylated tubulin (Red) and anti-Cop6 (Green). The secondary antibody for acetylated tubulin and Cop6 is Alexa 647 and Alexa 488 conjugated anti-mouse IgG and anti-goat IgG, respectively. Scale bar=2 μm in single cell and scale bar=5μm in group of cells.

(A)

Chlamyopsin5

A0A2K3E007

Protein degradation

A0A2K3CPY7
A0A2K3CPZ6
A0A2K3DIH3
NIT1
A0A2K3DEH6
A0A2K3DMV5
UBC3
A0A2K3DQP2
STM2
A8J3A7_CHLRE
A8J3A8_CHLRE
A0A2K3CVV1

Intraflagellar transport machinery

CPKG1
FAP259
FAP116
IFT80
FAP22
DYF13
IFT74
BLD1
FLA10
IFT81
IFT172
DHC1b
IFT57
FLA8
IDA5
IFT20
IFT88
EB1
ARL2

CAM1

Phosphatases

FTT1
PKL1
PPA1

A0A2K3D7N7

Chlamyopsin6

Photoreceptors

CPH1
PHOT
cop4
COP3
UVR3
MKP1

Clusters description

● Intraciliary transport
● Protein sumoylation
● Protein phosphatase 2A homologues
● Entrainment of circadian clock
  Phototransduction

(B)

Might assist in the sexual life cycle

Role in sexual life cycle-

Chlamyopsin5

CGK2 (cGMP dependent protein kinase)

Chlamyopsin6

Adenylyl cyclase associated protein

NIT1 (Nitrate reductase)

PKL1 (Serine/threonine protein phosphatase) PP2Ac

MKP1 (MAP Kinase Phosphatase 1)

Phototropin
Cryptochrome
ChR2
ChR1

-Role in eyespot development, sexual life cycle, photosynthesis and circadian clock

IFT172
Kinesin-2 motor protein

-Role in trafficking

EB1

Actin/ IDA5

ARLA1A/B

ARL11

14-3-3

VDAC
UBC3
PPA1 (Protein Phosphatase)
CaM1 Calmodulin

Role in abiotic stress

—— Nano-LC-MS/MS
—— Binding motif
—— Literature
—— STRING database

Figure 8: Predicted interactome of Cop5, Cop6, and 14-3-3.

(A) A combined interacting network was constructed based on the interactions reported in literature, interactions derived from the STRING database, proteins possessing an interacting motif for 14-3-3, and proteins detected in Nano-LC-MS/MS of lab data. The red dotted line denotes interaction is based on our results.

(B) Depicts the major biological processes involving different components of the predicted interactome in *C. reinhardtii* based on literature, proteomics data (available in the database and our lab data), and experimental data (from our study).

Graphical abstract

Cop6 is re-assembled for retrograde movement at flagellar tip with the help of IFT172

Cop6 are taken back to cell body by dynein motor

Cop6 are taken to the flagella by kinesin motor

IFT52 and IFT88 are essential for the assembly of Cop6 onto the IFT complex for trafficking to flagella

IFT20 interacts with Cop6

Retrograde

Anterograde

Basal body

IFT52   IFT88

IFT20

Nucleus

Golgi stacks

Eyespot

Cop5

Cop5 localizes in the eyespot

Chloroplast

A   Complex A

B   Complex B

K   Kinesin-2

D   Dynein

IFT52   IFT52

IFT88   IFT88

172   IFT172

Cop5

Cop6

IFT20   IFT20