Evidence of a distinct *Blumeria graminis* f. sp. *tritici* pathotype structure in

Australian wheat powdery mildew: implications for resistance breeding

Ancy Tony[1], Daniel Mullan[2], Huyen T.T. Phan[1], Francisco J. Lopez-Ruiz[1], Kejal N.

Dodhia[1], Benjamin J. Saunders[3], Cao (Grace) Fang[1], Simon Ellwood[1],

Kar-Chun Tan[1*]

[1]The Centre for Crop and Disease Management, School of Molecular and Life

Sciences, Curtin University, Bentley 6102, Australia

[2]Intergrain Pty Ltd, Bibra Lake 6163, Australia

[3]School of Molecular and Life Sciences, Curtin University, Bentley 6102, Australia.

*Corresponding author: Kar-Chun.Tan@curtin.edu.au

Abstract

The obligate biotrophic fungus *Blumeria graminis* f. sp. *tritici* (*Bgt*) is the causal agent

of wheat powdery mildew. To investigate virulence diversity, 30 Australian *Bgt* isolates

collected between 2020 and 2024 were screened against a panel of 24 wheat lines

carrying defined resistance (*R*) genes. Pathotyping analysis indicated that the

Australian *Bgt* population is diverse in its virulence profile. Non-metric

multidimensional scaling analysis of disease phenotyping revealed evidence of six

distinct *Bgt* pathotype groups. In addition, we identified wheat varieties carrying

defined *R* genes and their combinations that remain effective against the

contemporary Australian *Bgt*. Notably, wheat cvs Khapli and Tabasco, possessing

Pm4a and *Pm2,* respectively, in addition to unknown *R* gene(s), were highly resistant

to all the isolates tested. Furthermore, elite European varieties that possess durable

resistance, such as Alchemy and Cordiale, display complete to moderate resistance

against all selected isolates representing the Australian pathotype groups.

Key words: *Blumeria graminis* f. sp. *tritici,* powdery mildew resistance, pathotype

structure, virulence diversity, resistance breeding.

The obligate fungal pathogen *Blumeria graminis* f. sp. *tritici* (*Bgt*) causes wheat

powdery mildew (WPM) that results in typical yield losses of 10–15% and up to 50%

under favourable conditions (Rana et al. 2022). In Australia, around 12.6 million

hectares are sown with wheat annually, providing a large breeding ground for *Bgt*

(Department of Agriculture 2025). The fungus develops haustoria in epidermal cells to

extract nutrients and reproduces asexually through airborne conidia, while

chasmothecia are formed sexually under stress or at the end of the season (Spanu

2014; Wicker et al. 2013). The deployment of genetic resistance in wheat is an

environmentally sustainable strategy to minimize fungicide dependence and mitigate

associated ecological impacts. Resistance to WPM comprises race-specific

resistance, mediated by major WPM resistance (*R/Pm*) genes that induce

hypersensitive responses, and adult-plant or partial resistance, mostly a slow-

mildewing form associated with broad-spectrum and durable protection (Alam et al.

2011; Ge et al. 2021). To date, over 100 *Pm* genes and allelic variants (*Pm1-*

Pm71) conferring major and minor resistance to *Bgt* have been characterized in

wheat, but only a small subset of these have been cloned (Kunz et al. 2025; Li et al.

2025). To achieve durable resistance, resistance-associated quantitative trait loci must

exhibit consistent effects across multiple years and diverse environmental conditions,

while maintaining stable expression throughout plant development (Kang et al. 2020).

Most wheat cultivars currently grown in Australia exhibit partial to complete

susceptibility to *Bgt* (Shackley et al. 2025). The limited presence of widespread

resistance to WPM in current germplasm highlights the urgent need for improvement,

especially in light of proliferating fungicide resistance (Ireland et al. 2022).

Demethylation inhibitor (DMI) and quinone-outside inhibitor resistance have recently

become widespread in Eastern Australia, and DMI-resistance mutations have also

been detected in Western Australia, as reported by the Australian Fungicide

Resistance Extension Network (2025). This underscores declining fungicide efficacy

against WPM (GRDC 2025; Lopez-Ruiz et al. 2023).

Race-specific resistance underlies the existence of pathotypes, or variants with a

distinct virulence pattern on host genotypes (Thrall et al. 2001). Monitoring WPM

pathotype structure across Australian agro-ecological zones is crucial for tracking

virulence variation and guiding effective, region-specific resistance deployment in

geographically isolated growing regions, as different commercial wheat varieties

adopted by Australian regions affect *Bgt* adaption to varietal differences (Brown and

Harris 2020; Shackley et al. 2023). The aims of this study were to determine the

pathotypes present in Australia, analyse virulence differences among geographic

locations, and determine the resistance status of selected sources of *Pm* genes

against the current Australian *Bgt* population.

Thirty *Bgt* isolates were collected from major wheat-growing regions across Australia

between 2020 and 2024 (Supplementary Table 1 and Supplementary Figure 1).

Monoconidial isolates were established and maintained on the susceptible wheat cv.

Trojan using a high-throughput detached leaf assay (DLA) method for infection studies

as previously described (Lopez-Ruiz et al. 2023; Tucker et al. 2013). To assess

pathogenicity, a panel of 24 wheat varieties carrying different combinations of defined

Pm genes sourced from the Australian Grains Genebank (AGG) was evaluated to

identify effective *Pm* genes and determine any evidence of local *Bgt* pathotypes

(Supplementary Table 2). Selected Australian commercial wheat lines were also

evaluated against WPM. These varieties are Brumby (rated as resistant except to a

pathotype), DS Pascal (resistant to moderately resistant), Rockstar (Moderately

susceptible to susceptible) and Trojan (susceptible) (Shackley 2022). All wheat

varieties were sown in individual pots with three replicates and grown in an

environment-controlled growth chamber. Seven-day-old seedling leaf sections were

then excised and placed onto benzimidazole agar plates. Isolate inoculation was

performed using an inoculation tower and plates were incubated at 20°C under a

12:12 h light:dark photoperiod as previously described (Tucker et al. 2013). Disease

severity was scored at seven days past infection based on a five-point (0-4) infection

type (IT) scale (Supplementary Figure 2). 0 and 1 were considered highly resistant

(R), 2 moderately resistant (MR), 3 moderately susceptible (MS), and 4 highly

susceptible(S) (Li et al. 2019).

We used a heatmap approach to visualise the virulence profile of all 30 *Bgt* isolates

and observed significant variations in the virulence profiles (Figure 1). Isolates 2023-

060 (SA), 22PMWR15 (WA) and 20PMFRG01 (WA) were highly virulent on the wheat

panel, whereas 2024PM121 (WA), 2023-046 (SA), 2024PM111 FT168 (QLD) and

2024PM108 (WA) were the least. Wheat cvs Khapli and Tabasco, both carrying an

unknown *Pm* gene as well as *Pm4a* and *Pm2,* respectively, exhibited complete

resistance to all the isolates tested. The Australian wheat cv Brumby possesses the

highest disease resistance rating against WPM. However, resistance was overcome

by the highly virulent isolate 2023-060 from a field in Malinong (SA), as was previously

observed in another field location elsewhere in SA (Shackley 2022). Despite this, our

study indicates that wheat cvs Khapli and Tabasco possess highly effective resistance

genes against isolate 2023-060 (Supplementary Table 3). Wheat cvs Mich.Amber/8Cc,

Seri 82, and Sonora/8Cc possess *Pm3f*, *Pm8* and *Pm3c,* respectively, and displayed

the lowest levels of resistance. Thus, these *Pm* genes are ineffective against the

current Australian *Bgt* population. It was observed that cv Ulka/8Cc, possessing *Pm2*

(Briggle 1969), shows a mixed resistance response to the isolate panel, whereas

Tabasco, also possessing *Pm2* (Wu et al. 2023), is resistant to all isolates. This

strongly suggests that cv Tabasco carries an unknown *Pm* gene in addition to *Pm2*.

Likewise, although cvs Kavkaz and Seri 82 both carry *Pm8* (Hurni et al. 2013; Villareal

et al. 1998), their contrasting responses suggest the presence of additional resistance

in Kavkaz.

* WA = Western Australia, SA = South Australia, VIC = Victoria, QLD = Queensland

Figure 1: Virulence profiling of Australian *Bgt* isolates on differential and commercial

wheat lines. The heatmap represents the virulence spectrum of *Bgt* isolates (*x*-axis)

tested on a set of differential wheat lines (*y*-axis). The heatmap was generated using

ComplexHeatmap package (ver.2.22.0) in R (ver.4.4.1) (Gu et al. 2016). Compiled

using Adobe Photoshop (ver. 25.12.3). Detailed disease scores are provided in

Supplementary Table 3.

A two-dimensional non-metric multidimensional scaling (2D nMDS) analysis was

performed to assess whether virulence variability across the panel reflected any

underlying population structure. The disease severity score data were exported and

analysed in PRIMER 7 (ver. 7.0.24), and a 2D nMDS ordination plot was constructed

based on the Euclidean distance. A similarity profile then used to visualise the *Bgt*

clustering pattern that is indicative of pathotypes (M. J. Anderson 2008) (PRIMER 7

PERMANOVA+ add-on, PRIMER-E Ltd). 2D nMDS ordination identified six distinct

pathotype groups (Figure 2). Correlation vectors demonstrate the contribution of

specific host *R* genes to the observed variation (Figure 2A; Supplementary Figure 3).

Thus, the longer the vector, the greater its influence in differentiating isolates and their

orientation reflects the direction of increasing susceptibility to those isolates. cv Khapli,

which exhibited an IT of 0 against all isolates, is positioned near the centre of the

correlation vectors due to the lack of variation in its effectiveness. Wheat cvs Virest,

Thew, Normandie and Rockstar exhibited similar susceptibility patterns and were the

major contributors to pathotype IV formation. Brumby was susceptible only to isolate

2023-060, and it aligns with pathotype V, which carries the corresponding virulence.

Wheat cvs Khapli/8Cc and Armada displayed related *Bgt* susceptibility profiles, with

both showing pronounced susceptibility to QLD-derived isolates that predominantly

define pathotype I. Wheat cvs TD 1656, Timgalen and Coker 9803 exhibited similar

Bgt susceptibility profiles and contributed to the clustering of isolate 22PMWR26 (VIC),

together with the variety Brigand, which showed complete susceptibility exclusively to

this isolate. Out of these pathotypes, III and V, each carrying a single isolate, were

collected in 2023 from South Australia. However, III contains the second least virulent

isolate (2023-046), and the most virulent isolate (2023-060) is assigned to pathotype

V. To assess the geographical variation of these pathotypes, a 2D nMDS plot was

generated to determine if there is any correlation between state collection location and

pathotype. Clustering of isolates by location is difficult to conclude due to insufficient

isolates collected from VIC and among pathotype groups (Figure 2B). We then

examined for temporal variation (Figure 2C). This indicated pathotypes I and II were

comprised solely of isolates from 2024. Some isolates collected from 2024 are also

found in pathotype IV, which consists of isolates collected in 2020, 2021, 2022 and

2024 (Figure 2C). However, more isolates are needed to conclusively determine

whether there is a spatial-temporal effect on virulence.

Figure 2: 2D nMDS scaling ordination plot reveals distinct Australian *Bgt* pathotypes.

A Euclidean distance similarity matrix was calculated based on the pairwise virulence

profile of 30 *Bgt* isolates and used to generate nMDS coordinates of each sample and

was overlaid with the hierarchical clustering analysis of these isolates, denoted as

green ellipses or as groups in roman numerals. **A**, Each isolate, denoted by black dots

and is overlaid with the correlation vectors calculated based on the Pearson

correlation, denoted as blue lines with wheat varieties. **B**, Isolate plotted based on

collection location. **C**, Isolate plotted based on the year of collection.

Single major *R* genes are often rapidly overcome by new virulent pathogen races due

to strong selection pressure, whereas durable resistance remains effective over time

and across environments. Durable polygenic resistance provides complete to partial

protection against a wider range of pathotypes by lowering the selection pressure

(Burdon et al. 2014). In the United Kingdom, Alchemy, Cordiale, and Einstein

demonstrated long-lasting resistance against the local *Bgt* population over a 10-year

disease trial survey (Brown 2015). These varieties were screened to assess their

effectiveness against the Australian *Bgt* population by selecting representative isolates

from each pathotyping group and screened using seedling DLAs. Pathotype VI was

excluded due to loss of isolates. Wheat cvs Alchemy and Cordiale were resistant to all

pathotypes, whereas Einstein displayed moderate resistance across all pathotypes

(Table 1). Even though pathotype V contains the most virulent isolate, all three elite

European cultivars showed moderate to strong resistance. This suggests the potential

effectiveness of these varieties in providing long-term broad-spectrum WPM

resistance in Australia. However, further testing over several years across diverse

geographical and environmental conditions is required.

180   **Table 1 here**

Knowledge on Australian *Bgt* aggressiveness is limited, with only two existing

published data focused exclusively on WA isolates. For instance, Golzar et al.(2016)

pathotyped cumulative pools of WA *Bgt* isolates collected between 2011 and 2014,

whereas Kloppe et al. (2022) characterised 15 WA isolates collected in 2016. We then

compared these earlier findings with those from the contemporary Australian isolates

analysed in this study (Table 2). For statistical analysis, the disease scores from both

studies were converted to a 0–4 numerical scale and assigned infection types

following the method described by Li et al. (2019). Comparisons of WA isolates

revealed that *Pm3b*, *Pm3e*, *Pm1+2+9*, and *Pm2+6* exhibited a loss of effective

resistance over time. In contrast, *Pm4b* and *Pm17* demonstrated renewed

effectiveness against the WA *Bgt* isolates examined in this study. *Pm3a, Pm4a* and

the *Pm4a+unknown R* gene (from cvs. Asosan/8Cc, Khapli/8Cc and Khapli,

respectively) maintained their effectiveness over time in WA, with *Pm3a* and

Pm4a+unknown also remaining effective against *Bgt* isolates from other Australian

states. The effectiveness of these identified *Pm* genes require further experimental

validation to ensure that WPM resistance in varieties that carry them is not due to

unknown background resistance. However, Golzar et al.(2016) observed that Khapli

demonstrated adult plant WPM resistance.

199   **Table 2 here**

Since 2020, severe WPM outbreaks have occurred across the eastern states of

Australia, causing yield losses of up to 50% (Ellwood et al. 2024). Strategic use of

resistance genes is only effective when supported by detailed characterisation of *Bgt*

virulence and pathotype diversity (Czembor et al. 2014; El-Shamy and Mohamed

2022; Lalošević et al. 2021; Parks et al. 2008). Since the deployment of a major *R*

gene often leads to boom-and-bust disease dynamics driven by the emergence of new

pathogen races (Pink and Puddephat 1999). This underscores the need for continued

monitoring of pathogen diversity and resistance gene efficacy to guide region-specific

disease management (Zou et al. 2023).

This study represents the first reported simultaneous pathotyping of *Bgt* across a

broad coverage of major wheat-growing regions in Australia, resulting in the

identification of multiple virulences and six distinct pathotype groups. As many of the

most popular wheat varieties in Australia are susceptible to WPM, this suggests a

history of exposing *R*-genes in the field, many of which have lost their effectiveness.

Despite this, wheat cvs Tabasco and Khapli exhibit complete resistance to the tested

Australian *Bgt* population and represent valuable sources of resistance that can be

deployed by the cereal industry into existing parental varieties. In addition to major *R*

genes, the use of durable broad-spectrum resistance from Alchemy and Cordiale to

minimise the impact of WPM in Australia highlights the potential of tapping into

international germplasm as a resource for WPM resistance breeding in Australia.

Findings from this study support targeted deployment and stacking of effective *R*

genes and, ideally, implementing broad-spectrum WPM resistance. Materials

collected for the current research can be used to study WPM evolution, reveal their

origins, and identify their virulence factors. Furthermore, we hypothesise novel *Pm*

genes effective against Australian *Bgt* are present and remain undiscovered in the

differential wheat panel. Efforts are underway to locate these unknown genes and to

assess their efficacy in disease control.

Acknowledgements

We thank the AGG (Horsham, Australia) for the provision of wheat germplasm used in

this study. Sam Trengove (Trengove Consulting), Hossein Golzar (DPIRD), Andrea

Hills (DPIRD) and Ciara Beard (DPIRD) for the provision of *Bgt* samples. Julie

Lawrence (Curtin University) for valuable technical assistance. Kul Adhikari (Curtin

University), Steven Chang (Curtin University), Marta Lopez-Ruiz (Curtin University),

and Nirmala Sharma (Curtin University) for isolate maintenance.

Funding: Support was provided by Intergrain Pty Ltd (Grant CTR18532) and the

Centre for Crop and Disease Management, a co-investment between the Grains

Research and Development Corporation and Curtin University (Grant CUR1403-

002BLX).

Literature Cited

Alam, M. A., Xue, F., Wang, C., and Ji, W. 2011. Powdery mildew resistance genes in wheat: identification and genetic analysis. J. Mol. Biol. Res. 1:20.

Australian Fungicide Resistance Extension Network, A. 2025. Fungicide resistance. Grains Research and Development Corporation (GRDC).

Briggle, L. 1969. Near-isogenic lines of wheat with genes for resistance to *Erysiphe graminis* f. sp. *tritici* 1. Crop Sci. 9:70-72.

Brown, J. K. 2015. Durable resistance of crops to disease: a Darwinian perspective. Annual Review of Phytopathology 53:513-539.

Brown, S., and Harris, K. 2020. 2021 Victorian crop sowing guide. Grains Research and Development Corporation (GRDC) Barton, Australia.

Burdon, J., Barrett, L., Rebetzke, G., and Thrall, P. 2014. Guiding deployment of resistance in cereals using evolutionary principles. . Evol. Appl. 7(6):609-624.

Czembor, H. J., Domeradzka, O., Czembor, J. H., and Mańkowski, D. R. 2014. Virulence structure of the powdery mildew (*Blumeria graminis*) population occurring on triticale (x *Triticosecale*) in Poland. Phytopathology 162:499-512.

Department of Agriculture, F. a. F. D. 2025. Australian Crop Report June 2025. DAFF.

El-Shamy, M. M., and Mohamed, M. E. 2022. Virulence analysis of wheat powdery mildew races during 2019–2020 seasons in Egypt. Cereal Res. Commun. 50:67-73.

Ellwood, S. R., Lopez-Ruiz, F. J., and Tan, K. C. 2024. Barley powdery mildew control in Western Australia and beyond. Plant Pathol. 73:1666-1674.

Ge, C., Wentzel, E., D'Souza, N., Chen, K., Oliver, R. P., and Ellwood, S. R. 2021. Adult resistance genes to barley powdery mildew confer basal penetration resistance associated with broad-spectrum resistance. The Plant Genome 14:e20129.

Golzar, H., Shankar, M., and D'Antuono, M. 2016. Responses of commercial wheat varieties and differential lines to western Australian powdery mildew (*Blumeria graminis* f. Sp. *tritici*) populations. Australas. Pl. Pathol. 45:347-355.

GRDC. 2025. Paddock practices: Potential for fungicide resistance detected in Esperance wheat powdery mildew. Grains Research and Development Corporation(GRDC).

Gu, Z., Eils, R., and Schlesner, M. 2016. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. Bioinformatics 32:2847-2849.

Hurni, S., Brunner, S., Buchmann, G., Herren, G., Jordan, T., Krukowski, P., Wicker, T., Yahiaoui, N., Mago, R., and Keller, B. 2013. Rye *Pm8* and wheat *Pm3* are orthologous genes and show evolutionary conservation of resistance function against powdery mildew. Plant J. 76:957-969.

Ireland, K. B., Beard, C., Cameron, J., Chang, S., Davidson, J., Dodhia, K., Garrard, T., Hills, A., Holloway, G., and Jayasena, K. 2022. Fungicide resistance management in Australian grain crops. Grains Research and Development Corporation.

Kang, Y., Zhou, M., Merry, A., and Barry, K. 2020. Mechanisms of powdery mildew resistance of wheat–a review of molecular breeding. Plant Pathol. 69:601-617.

Kloppe, T., Boshoff, W., Pretorius, Z., Lesch, D., Akin, B., Morgounov, A., Shamanin, V., Kuhnem, P., Murphy, P., and Cowger, C. 2022. Virulence of *Blumeria graminis* f. sp. *tritici* in Brazil, South Africa, Turkey, Russia, and Australia. Front. Plant Sci. 13:954958.

Kunz, L., Jigisha, J., Menardo, F., Sotiropoulos, A. G., Zbinden, H., Zou, S., Tang, D., Hückelhoven, R., Keller, B., and Müller, M. C. 2025. Avirulence depletion assay: Combining *R* gene-mediated selection with bulk sequencing for rapid avirulence gene identification in wheat powdery mildew. PLoS pathog. 21:e1012799.

Lalošević, M., Jevtić, R., Župunski, V., Maširević, S., and Orbović, B. 2021. Virulence structure of the wheat powdery mildew population in Serbia. Agronomy 12:45.

289   Li, G., Xu, X., Tan, C., Carver, B. F., Bai, G., Wang, X., Bonman, J. M., Wu, Y., Hunger, R., and
290   Cowger, C. 2019. Identification of powdery mildew resistance loci in wheat by
291   integrating genome-wide association study (GWAS) and linkage mapping. Crop J. 7:294-
292   306.
293   Li, J., Yu, N., Sun, N., Geng, L., Qie, Y., Zhai, D., Wang, Y., Li, L., Liu, X., and Sun, X. 2025.
294   Identification and characterization of the powdery mildew resistance in cultivated
295   emmer wheat accession Lxd-682 via bulked segregant RNA sequencing. BMC Plant Biol.
296   25:583.
297   Lopez-Ruiz, F., Dodhia, K., Chang, S., Simpfendorfer, S., and Trengove, S. 2023. Fungicide
298   resistance in wheat powdery mildew. Pages 29-34 in: Grains research update 2023
299   M. J. Anderson, R. N. G., and K. R. Clarke. 2008. PERMANOVA+ for PRIMER: guide to software
300   and statistical methods. Primer-E Limited.
301   Parks, R., Carbone, I., Murphy, J. P., Marshall, D., and Cowger, C. 2008. Virulence structure of the
302   eastern US wheat powdery mildew population. Plant Dis. 92:1074-1082.
303   Pink, D., and Puddephat, I. 1999. Deployment of disease resistance genes by plant
304   transformation–a 'mix and match'approach. Trends Plant Sci. 4:71-75.
305   Rana, V., Batheja, A., Sharma, R., Rana, A., and Priyanka. 2022. Powdery mildew of wheat:
306   research progress, opportunities, and challenges.
307   Shackley, B., Power, S., Paynter, B., Dhammu, H., Wackett, B., Nicol, D., Seymour, M., and Shea,
308   G. 2025. 2026 Western Australian crop sowing guide. State of Western Australia
309   (Department of Primary Industries and Regional Development).
310   Shackley, B., Paynter, B, Bucat, J, Seymour, M, and Dhammu, H. 2022. 2023 Western Australian
311   crop sowing guide. Page 209 Department of Primary Industries and Regional
312   Development, Western Australia.
313   Shackley, B. J., Power, S., Paynter, B., Troup, G., Seymour, M., and Dhammu, H. S. 2023. 2024
314   Western Australian crop sowing guide. DPIRD.
315   Spanu, P. D. 2014. The Genomes of the Cereal Powdery Mildew Fungi, Blumeria graminis. Pages
316   161-175 in: Genomics of Plant-Associated Fungi: Monocot Pathogens. R. A. Dean, A.
317   Lichens-Park and C. Kole, eds. Springer Berlin Heidelberg, Berlin, Heidelberg.
318   Thrall, P. H., Burdon, J. J., and Young, A. 2001. Variation in resistance and virulence among
319   demes of a plant host-pathogen metapopulation. J. Ecol. 89:736-748.
320   Tucker, M., Jayasena, K., Ellwood, S., and Oliver, R. 2013. Pathotype variation of barley powdery
321   mildew in Western Australia. Australas. Plant Pathol. 42:617-623.
322   Villareal, R. L., Bañuelos, O., Mujeeb-Kazi, A., and Rajaram, S. 1998. Agronomic performance of
323   chromosomes 1B and T1BL. 1RS near-isolines in the spring bread wheat Seri M82.
324   Euphytica 103:195-202.
325   Wicker, T., Oberhaensli, S., Parlange, F., Buchmann, J. P., Shatalina, M., Roffler, S., Ben-David,
326   R., Doležel, J., Šimková, H., and Schulze-Lefert, P. 2013. The wheat powdery mildew
327   genome shows the unique evolution of an obligate biotroph. Nat. Genet. 45:1092-1096.
328   Wu, J., Jia, H., Qiao, L., Fu, B., Brown-Guedira, G., Nagarajan, R., and Yan, L. 2023. Genetic basis
329   of resistance against powdery mildew in the wheat cultivar "Tabasco". Mol. Breed.
330   43:56.
331   Zou, S., Xu, Y., Li, Q., Wei, Y., Zhang, Y., and Tang, D. 2023. Wheat powdery mildew resistance:
332   from gene identification to immunity deployment. Front. Plant Sci. 14:1269498.

335 **Table 1:** Disease phenotypes of European durable WPM resistant wheat lines to

336 Australian *Bgt.* Each column contains representative isolates of different pathotypes,

337 and each row represents wheat lines. Wheat cv. Khapli (resistant) and Trojan

338 (susceptible) were used as controls and scored/rated based on 0-4 numerical scale

339 as described above (Li et al. 2019).

	Pathotype I	Pathotype II	Pathotype III	Pathotype IV	Pathotype V
	2024PM110172	2024PM126	2023-046	20PMFRG01	2023-060
Alchemy	R	R	R	R	R
Cordiale	R	R	R	R	R
Einstein	MR	MR	R	MR	MR
Khapli	R	R	R	R	R
Trojan	S	S	S	S	S

341   **Table 2:** Comparative analysis of previous pathotyping studies on Australian *Bgt* with

342   this study. 'NA' indicates *Pm* genes not examined. Seedling infection scores from these

343   studies were converted to a 0 to 4 numeric scale for statistical comparisons (Li et al.

344   2019). * Indicates that the wheat varieties used for the particular *Pm* genes differ

345   among studies.

	(Golzar et al. 2016)	(Kloppe et al. 2022)	This study			
Region	WA	WA	WA	SA	QLD	VIC
R gene/sampling year	2011-2014	2016		2020-2024		
*Pm1a**	S	S	S	S	R	R
Pm2	NA	R	MR	MR	R	S
Pm3a	R	R	R	R	R	R
Pm3b	NA	MR	MS	MR	R	R
Pm3c	S	NA	S	MS	MS	MS
*Pm3e**	R	NA	MR	R	R	R
Pm4a	R	R	R	R	S	R
Pm4a+unknown	R	NA	R	R	R	R
*Pm4b**	S	R	R	R	S	R
Pm5a	S	NA	MS	MS	S	S
*Pm6**	S	S	MR	MS	MR	S
Pm7	MS	NA	MS	MR	MR	R
*Pm8**	S	MR	MR	MR	S	MS

Pm17	S	R	R	R	MS	S
Pm1+2+9	R	NA	S	MS	R	R
Pm2+6	R	NA	MS	MS	MR	S

347