Multi-Omics Reprogramming Drives a Counterintuitive Reversal of Disease Susceptibility During Ageing

Alejandro Valdivieso[1], Léo Duperret[2], Bruno Petton[3], Gaelle Courtay[1], Océane Romatif[1], Juliette Pouzadoux[1], Sylvain Henry[4], Andrei Turtoi[4,5] Eve Toulza[2], Arnaud Lagorce[2], Lionel Degremont[4], Benjamin Morga[6], Celine Cosseau[2], Fabrice Pernet[3], Guillaume Mitta[7,*], Jeremie Vidal-Dupiol[1,*]

[1] IHPE, Univ Montpellier, CNRS, IFREMER, Univ Perpignan Via Domitia, Montpellier, France

[2] IHPE, Univ Perpignan Via Domitia, CNRS, IFREMER, Univ Montpellier, Perpignan, France

[3] Ifremer, LEMAR UMR 6539, UBO/CNRS/IRD/Ifremer, Argenton-en-Landunvez, France

[4] Platform for Translational Oncometabolomics, Biocampus, CNRS, INSERM, Université de Montpellier, Montpellier, France

[4] Tumor Microenvironment and Resistance to Therapy Lab, Institut de Recherche en Cancérologie de Montpellier, INSERM U1194, Montpellier, France

[6] Ifremer, ASIM, Adaptation Santé des Invertébrés Marins, La Tremblade, France

[7] Ifremer, ILM, IRD, Université Polynesie Francaise, UMR SECOPOL, Tahiti, French Polynesia, France

*Correspondence: guillaume.mitta@ifremer.fr and jeremie.vidal.dupiol@ifremer.fr

Abstract

Ageing is a progressive and irreversible biological process characterized by the deterioration of physiological functions and increased vulnerability to mortality. Although extensively studied in vertebrates, ageing in long-lived invertebrates remains comparatively unexplored. While ageing typically leads to greater susceptibility to infectious diseases, a striking and unexpected reversal was identified in oysters: older oysters exhibit markedly increased tolerance to the Pacific Oyster Mortality Syndrome (POMS), a panzootic disease primarily driven by the OsHV-1 herpesvirus and responsible for severe losses in global aquaculture. To investigate this counterintuitive pattern, we challenged oysters aged 4, 16, and 28 months from four biparental families and conducted an integrative multi-omics analysis, including epigenomics, transcriptomics, and metabolomics on the two families showing the strongest age-related increase in survival. Our results reveal that ageing in *Magallana gigas* is characterized by coordinated epigenetic, transcriptional, and metabolic reprogramming that reduces host permissiveness to POMS. We show that the epigenetic remodeling of key immune regulators (e.g., Toll-like receptors, MyD88) aligns with transcriptional rewiring of NF-κB and ubiquitin pathways, producing a finely tuned innate immune state marked by enhanced antiviral activity but reduced antibacterial responsiveness. We also identify age-related repression of mTOR signaling, likely promoting autophagy and improving viral control. These regulatory changes are tightly linked to metabolic adjustments, including reduced TCA cycle flux, remodeled nitrogen metabolism, and altered glutathione dynamics, which collectively support a stress-tolerant, energy-conserving phenotype. Together, our findings reveal a fundamental evolutionary trade-off: juveniles prioritize growth at the cost of viral susceptibility, whereas adults invest in cellular maintenance and antiviral preparedness.

Keywords

POMS, Host-pathogen interaction, Multi-omics integration, Ageing, Pacific oyster

Introduction

Ageing is a progressive and irreversible biological process marked by the gradual deterioration of physiological functions, which over time reduces organismal performance and increases susceptibility to mortality (López-Otín et al., 2013). It is driven by a set of conserved molecular and cellular alterations, including DNA damage, telomere shortening, epigenetic drift, loss of proteostasis, and mitochondrial dysfunction (Mahmoudi & Brunet, 2012). These hallmarks disrupt key processes such as immune regulation, metabolic balance, oxidative stress detoxification, and cell-to-cell communication (López-Otín et al., 2023).

Ageing was mostly studied in vertebrates, but mollusk bivalves are increasingly used as a model given the inter-species variability of maximum life span, ranging from a couple of years for some vagile species (*Argopecten*

irradians irradians) to almost 400 years for sessile species (*Artica islandica*) (Abele & Philipp, 2013). In general
ageing in bivalves was shown to induce, at the functional level a lowering of the energetic metabolism, a
diminishing of the antioxidant defense, an induction of apoptosis, autophagy, and DNA repair mechanisms, and
an attenuated stress response, including immune gene expression (Philipp & Abele, 2010; Husmann et al., 2014).
The latter predicts that ageing in mollusks should be accompanied by an increasing vulnerability to diseases.
For *Magallana gigas*, increased susceptibility with age to the bacterial pathogen *Vibrio aestuarianus* was well
documented (Azéma et al., 2017). Surprisingly, this trend is totally reversed in face of the Pacific Oyster
Mortality Syndrome (POMS), where individuals show lower susceptibility through time. Indeed, spat and
juvenile oysters exhibit remarkably higher mortality rates when compared to older individuals (Petton et al.,
2015; Green et al., 2016; Carrasco et al., 2017; Dégremont, 2013).
63
POMS is a polymicrobial disease initiated by primary infection with the Ostreid herpesvirus type 1 microvariant
(OsHV-1 µVar) targeting and disrupting hemocyte physiology, thus impairing host immunity, and facilitating a
secondary bacterial infection, including *Vibrio* species (de Lorgeril et al., 2018; Rubio et al., 2019; Delisle et al.,
2022; Clerissi et al., 2023; Oyanedel et al., 2023; Kunselman et al., 2024). Given its panzootic status and the
huge pressure it exerts on the oyster industry, a better understanding of the factors and associated molecular
mechanisms influencing POMS pathogenesis is necessary (Abbadi et al., 2018; Gittenberger et al., 2016; Hwang
et al., 2013; Keeling et al., 2014; Martenot et al., 2011; Mortensen et al., 2016; Peeler et al., 2012; Petton et al.,
2021; Roque et al., 2012; Segarra et al., 2010). Permissiveness to POMS is defined as the extent to which an
oyster allows the infection process initiated by OsHV-1 µVar, including subsequent dysbiosis and opportunistic
bacterial proliferation, to progress to mortality. Survival of POMS depends on a complex interplay between host
genetic and epigenetic inherited components (Dégremont et al., 2015; Azéma et al., 2017; de Lorgeril et al.,
2018; Gutierrez et al., 2020; Divilov et al., 2019, 2021; Gawra et al., 2023; Valdivieso et al., 2025), the influence
of temperature (Duperret et al., 2025), and food availability (Pernet et al., 2019). Although age strongly
influences permissiveness to POMS in oysters (Dégremont, 2013; Petton et al., 2015; Green et al., 2016;
Carrasco et al., 2017), the molecular mechanisms underlying age-dependent reduced susceptibility to POMS
remain totally unknown.
80
Given both the complexity of POMS pathogenesis in oysters and the profound physiological changes occurring
during ageing, single-omic approaches would be insufficient to fully capture the regulatory changes underlying
the age-dependent shift in disease susceptibility. Multi-omics integration offers an opportunity to simultaneously
assess upstream regulatory layers (e.g., epigenetic modifications such as DNA methylation), intermediate
molecular responses (e.g., gene expression changes), and downstream functional responses (e.g., metabolic
reprogramming). By integrating these complementary biological layers, it becomes possible to identify
coordinated molecular signatures and cross-layer interactions that would remain hidden in isolated datasets,
providing a more comprehensive and mechanistic understanding of how ageing reshapes the oyster's
susceptibility to POMS. We induced POMS challenge to oysters aged 4, 16, and 28 months, using four
biparental families. Two of them displaying the highest survival to POMS gain in function of age were subjected
to an integrative multi-omics analysis combining comparative epigenetics, transcriptomics, and metabolomics.
Our findings reveal both shared and family-specific molecular trajectories across age, involving key biological
processes such as cellular differentiation and antiviral defense. Metabolomic profiling further uncovered an age-
related accumulation of tricarboxylic acid (TCA) cycle intermediates, indicative of enhanced energy
mobilization in older oysters. Altogether, our multi-layered data highlight molecular signatures underlying the
age-driven transition from a disease-permissive to a non-permissive state, offering novel insights into the
improved survival of oysters during POMS outbreaks.
98

Materials and Methods
Production of full-sibling families
Four biparental oyster (H2D, F14R, F11N, and F14V) families (*Magallana gigas*) were produced at the Ifremer
hatchery in Argenton (France) in 2020. The Spat (F_1) were transferred and maintained under biosecured
conditions at the Ifremer nursery (Bouin, France). Further details about hatchery and nursery steps are described
in (de Lorgeril et al., 2018, 2020). At 4-, 16-, and 28-month-old, the offspring were relocated to the Ifremer

facilities in Argenton. Oysters were kept at 23°C and fed *ad libitum* (50/50 mixture of *Tisochrysis lutea* and
Skeletonema costatum, 1,500 µm³/µL) for 15 days for acclimation (Rico-Villa et al., 2006). The pathogen-free
status was confirmed by testing for the absence of OsHV-1 µVar DNA and ensuring that *Vibrio sp.*
concentrations were below ten colony-forming units per milligram of tissue, as described in (Petton et al., 2015).
109

Production of the viral suspension for the experimental infection

A viral suspension of Ostreid herpesvirus 1 µVar (OsHV-1 µVar) was prepared following the protocol from a
mix of several variants described by (Schikorski et al., 2011). From moribund oysters, the tissues were dissected,
pooled, and homogenized in sterile artificial seawater using a blender under chilled conditions. The homogenate
was subjected to sequential centrifugation steps to remove cellular debris (3,000 × g for 10 min followed by
10,000 g for 20 min at 4°C). The resulting supernatant was filtered through 0.45 µm and 0.22 µm pore-size
filters to eliminate bacteria and larger particles, yielding a clarified viral suspension. The quantification of
genomics units (viral load) of OsHV-1 µVar was performed by real-time quantitative Polymerase Chain
Reaction (qPCR) using a Stratagene Mx3005P Real-Time thermocycler. The qPCR reaction volume (20 µL)
was: 5 µL DNA (5 ng/µL), 2 µL of each primer at the final concentration of 550 nM (Eurogentec SA), 1 µL of
distilled water, and 10 µL of Brilliant III Ultra-Fast SYBR®Green PCR Master Mix (Agilent). The virus-specific
primer pairs targeted a region of the OsHV-1 µVar genome predicted to a gene that encodes a DNA polymerase
catalytic subunit (open reading frame, ORF) 100AY509253: Forward: 5'-
ATTGATGATGTGGATAATCTGTG-3' and Reverse: 5'-GGTAAATACCATTGGTCTTGTTCC-3' (Webb et
al., 2007; Pepin, 2013). The amplification qPCR program consisted of 3 min at 95°C, followed by 40 cycles at
95°C for 5 s and 60°C for 20 s, with a melting temperature curve of the amplicon to verify the specificity of the
amplification. Quantification of viral DNA copies was estimated by comparing the observed Cq values to a
standard curve of the dimer primer amplification product cloned into the pCR4-TOPO vector. The final
concentration was adjusted to 2.2×10 genome copies/µL and stored at –80°C.
129

Experimental infection

131 To investigate the effects of ageing on oysters' permissiveness to POMS, the four families (called recipients)
132 were exposed to the disease through a cohabitation experiment with OsHV-1 µVar-injected donor oysters at ages
133 4-, 16-, and 28 months (see below), mimicking the natural transmission of POMS (**Figure 1A** and **Table S1**).
134

Donor oysters came from two stocks of "pathogen-free" oysters to maximize the production of OsHV-1 µVar
during cohabitation: one stock, called Dellec, showing a highly susceptible to POMS (> 90%) (de Lorgeril et al.,
2018), and one stock, called NSI, with an expected 50–60% mortality (Petton et al., 2013). Twenty hours before
the onset of cohabitation, 4-month-old donor oysters were anaesthetized with hexahydrate MgCl$_2$ (ACROS, 50
g/L) for 2 hours (Suquet et al., 2009). Then, 100 µL of OsHV-1 µVar suspension was injected into their adductor
muscle with a 26-gauge needle using a multi-dispensing pipette (Schikorski et al., 2011). The injected donor
oysters were maintained at 23°C in tanks (Volume = 400 L) for 18 hours, an optimal temperature for the disease
(Petton et al., 2021). Temperature, pH, salinity, and dissolved oxygen were monitored twice a day with a WTW
Multiparameter MultiLine 3630 IDS and a Dotsman P750, equipped with a PT100 sensor and an iButton
DS1922L-F5 thermochron.
145

Cohabitation infection was performed by placing the injected donor and recipient oysters together at a 1:1
biomass ratio. The 4-, 16-, and 28-month-old oysters were acclimated in three, six, and fifteen tanks,
respectively, to maintain a constant biomass per liter, as the individual weight increased throughout the age. To
increase the transmission of OsHV-1 µVar to the recipient oysters, water renewal was halted for the first 125
hours post-cohabitation (hpc). At the end of this period, all surviving donor oysters were removed from the
tanks. Once cohabitation began, six live recipient oysters were collected at 0 (basal conditions), 3, 6, 12, and 24
hours hpc for each family in each experiment. All tissues were removed from the shells, flash-frozen in liquid
nitrogen, and stored at –80°C. Mortality was monitored twice daily, and dead oysters were removed as soon as
they were observed. The experiment ended when no deaths were recorded in any of the tanks for a continuous 48
hours. The infection experiments were conducted over three consecutive years, each dedicated to a specific age
group (4-, 16-, and 28-month-old recipient oysters), under the same conditions and within the same facility with

the same donor oysters. To validate the mortality across the three cohabitation experiments, an internal control
consisting of 4-month-old NSI recipient oysters was included in each trial.
159

160
Figure 1. Age-dependent survival and viral load dynamics following OsHV-1 infection for the four oyster
biparental families
A) Graphical representation of the experimental design of the four biparental Magallana gigas families (H2D,
F14R, F11N, F14V) exposed to OsHV-1 µVar through cohabitation with infected donor oysters. The families
were challenged at three different ages (4-, 16-, and 28-month-old) under controlled and permissive to the

disease tank conditions (23°C, O☐-saturated, ad libitum feeding, 400 L volume). **B)** Kaplan–Meier survival
curves of oyster families H2D, F14R, F11N, and F14V at 4-, 16-, and 28-month-old. Statistical differences were
assessed by Mantel-Cox log-rank tests using the 4-month-old group as reference within each family (*** =
P☐<☐0.01). **C)** Positive correlations between age and survival rates in all families. **D)** The OsHV-1 µVar DNA
load and RNA viral activity kinetics were measured during the first 24 hours post-cohabitation (hpc) in H2D and
F14R. Letters denote significant differences among the three age groups at 24 hpc, with the 4-month-old group
serving as the reference (One-way ANOVA followed by a Tukey's post hoc test). Data are presented as the mean
± standard deviation in each group.
174

DNA and RNA extractions from the same individual oyster
Samples were ground in 50 mL stainless steel bowls with 20 mm grinding balls continuously cooled with liquid
nitrogen as described (de Lorgeril et al., 2018) using the Retsch MM400 MILLa device. Grounded samples were
stored at –80°C. Following the manufacturer's protocol, genomic DNA (gDNA) was extracted using the
Genomic DNA Tissue Kit (Macherey-Nagel, 740952.250) and stored at –20°C. The gDNA concentration,
purity, and quality were assessed using the NanoDrop One spectrophotometer (Thermo Scientific) and 1%
agarose gel electrophoresis. Total RNA was extracted using the Direct-zol RNA Miniprep Kit (Zymo Research)
according to the manufacturer's protocol. The RNA concentration was measured using a NanoDrop One
spectrophotometer, and integrity was assessed with a BioAnalyzer 2100 (Agilent). All RNA samples were stored
at –80°C. To quantify the viral load present in the gDNA samples we followed the same protocol as described
above.
186

Genome references
For this study, we used the *M. gigas* reference genome (GCA_902806645.1) (Peñaloza et al., 2021; G. Zhang et
al., 2012). The *M. gigas* genome comprises a total of 30,418 coding genes, and a description of each gene is
provided in **Table S2**. For the virus, we used the OsHV-1 µVar genome (KY242785.1) (Burioli et al., 2017). For
this study, we used a previously constructed Gene Ontology (GO-term) gene annotation database (Duperret et
al., 2025), needed for the GO_MWU software (see below). Briefly, the coding sequences were extracted with
TransDecoder, annotated using ORSON (BLASTX against Uniprot-SwissProt; InterProScan domain prediction),
validated with the Omicsbox software (v. 3.4.5) (Gotz et al., 2008), and further enriched with immune-related
terms.
196

Epigenetic analysis
Preparation and library sequencing of samples
DNA methylation libraries were prepared and sequenced by IntegraGen SA (Evry, France) using the NEBNext
Enzymatic Methyl-sequencing (EM-seq) protocol. Briefly, 100 ng of extracted genomic DNA, supplemented
with unmethylated lambda DNA and methylated puc19 DNA as internal spike-in controls, was used for library
preparation. Sequencing was performed on an Illumina NovaSeq platform with 150 bp paired-end reads.
203

Processing and analysis of the samples
The raw paired read quality control for each sample was analysed using FastQC (v.0.53) software (Andrews,
2010), and the adapters were trimmed using TrimGalore! (v.0.6.10) software (Krueger et al., 2023) with: -q 30 --
paired --clip_R1 5 and --clip_R2 5 --Illumina. Any remaining adapters were removed in a second trimming
round with default parameters. We used the *'Bismark'* (v0.24.2) software (Krueger & Andrews, 2011),
employing the *'bismark_genome_preparation'* function to bisulfite convert the *M. gigas* genome reference. Then,
the trimmed reads were aligned to the bisulfite converted *M. gigas* genome using the *'bismark'* function: -q -N 0
-- score_min L,0, –0.4. The duplicated reads were removed using the *'deduplicate_bismark'* function, and the
methylation extraction calling was accomplished using the *'bismark_methylation_extractor'* function: --
no_overlap --cutoff 8. We applied the same methodology to bacteriophage lambda to assess enzymatic
conversion efficiency. We discarded the samples with < 98.0% of conversion efficiency. We used the *'methylKit'*
package (v.1.24.0) for data treatment (Akalin et al., 2012), and we retained only the common CpGs present in all
samples with a minimum coverage of 8X. For normalization, we used the "median" method. Because the
methylated fraction in bivalves primarily occurs within gene body regions (Gavery & Roberts, 2014; Männer et
al., 2021; Venkataraman et al., 2020, 2022), we intersect the CpGs located within gene body regions. First, we

selected the genic coordinates (chromosome, start, and end positions) of all coding genes in *M. gigas* using the
'biomaRt' package (v.2.54.1) (Durinck et al., 2009). We then employed the *'foverlaps'* function from the
'data.table' package (v.1.14.8) (Dowle et al., 2019) to overlap these gene coordinates with the identified CpG
coordinates to compile a list of genes containing CpGs within their boundaries. Finally, we calculated the
average methylation values for those genes that contained multiple CpG sites within the same gene body.
224
225 *Transcriptomic analysis*
226 *Preparation and library sequencing*
227 RNA-seq libraries were prepared using the NEBNext Ultra II Directional RNA Library Prep Kit. Paired-End
228 (PE) reads sequencing was performed with 100 base pair reads on the Illumina NovaSeq sequencer. Library
229 construction and sequencing were carried out by IntegraGen SA (Evry, France)
230
Processing and analysis of samples
The raw paired reads were trimmed using Trim Galore! software (v.0.6.7) (Martin, 2011; Krueger et al., 2023)
with parameters: -q 30 --illumina --stringency 1 -e 0.1 --length 35. The quality of the sequences was assessed
before and after trimming using FastQC software (v.0.11.9) (Andrews, 2010). We mapped the trimmed PE reads
to the *M. gigas* genome using the following parameters: --runMode alignReads, --outSAMmapqUnique 60, and -
-outSAMattributes All with the STAR software (v.2.7.10b) (Dobin et al., 2013). Finally, count reads were
obtained using the HTSeq software (v.0.9.1) (Putri et al., 2022) with: --stranded reverse -t gene -i ID -m union -q
--minequal 10. The trimmed reads were also mapped to the OsHV-1 µVar with Bowtie2 software (v2.3.4.3)
(Langmead et al., 2009; Langmead & Salzberg, 2012) to analyze the viral transcriptional activity. The HTSeq-
count software was used to obtain the number of counts per ORF with the same parameters as above, except for
the feature type as CDS and the ID attribute as product. The viral raw counts were normalized based on the
formula: $C = \left(\frac{C_{ORF}}{S_{ORF}}\right) \times \left(\frac{\bar{L}}{L_i}\right)$. While the first ratio corrects for the size of the ORF by dividing the raw counts
(C_{ORF}) with the size of it (S_{ORF}), the second one corrects the sequencing depth with a ratio of the average reads
for all the time points ($\bar{L}$) and the number of total reads for the specific time points (L_i), as described (de Lorgeril
et al., 2018). We used the *'DESeq2'* package (v.1.42.1) for data treatment (Love et al., 2014), and raw count gene
matrix was filtered (counts ≤ 15 across all the samples), and library normalization was applied, followed by a
variance-stabilizing transformation with the *'vst'* function.
248
Metabolomic analysis
The Montpellier Alliance for Metabolomics and Metabolism Analysis (MAMMA) platform (BioCampus,
Montpellier, France) carried out targeted metabolites on global metabolites (GM), nucleotide metabolites, and
metabolites involved in the tricarboxylic acid (TCA) cycle. Briefly, 20 mg of powder samples were
supplemented with 980 µL of ice-cold methanol (0013684102BS, Biosolve), 5 µL of 2-(N-Morpholino)-
ethanesulfonic acid (M8250, Sigma-Aldrich) and 20 µL of mixed internal standards along with 0.5 g of glass
beads. Samples were purified via SPE column and dried in a Speedvac (Turtoi, Jeudy, Henry, et al., 2023).
256
Processing and analysis of the samples
For the GM analysis (Turtoi, Jeudy, Valette, et al., 2023), samples were reconstituted in 200 µL of UPLC water
and transferred to LC-MS polypropylene vials. UHPLC 1290 Infinity II with an HPLC Discovery® HS F5, 3 µm
particle size, L × I.D. 15 cm × 2.1 mm (567503-U, Sigma-Aldrich) was used for chromatographic analysis. GM
profiling analysis was performed on a 6495 LC/TQ mass system and the Agilent MassHunter workstation
(Agilent, Santa Clara, USA) with the dynamic MRM scan mode. The nucleotide metabolites analysis was
conducted using chromatography on a UHPLC 1290 Infinity II system equipped with an InfinityLab Poroshell
HPH-C18 column (2.7 µm particle size; L × I.D. 100 cm × 3.1 mm; 695975-502, Agilent). The mobile phase A
consisted of 10 mM ammonium acetate (5.33004; Sigma-Aldrich), 2.5 µM medronic acid (5191-3940; Agilent),
and 0.1% formic acid in UPLC-grade water (0023214102BS; Biosolve). The mobile phase B consisted of 90%
UPLC-grade methanol (0013684102BS; Biosolve), 10% UPLC-grade water, 2.5 µM medronic acid, and 0.1%
formic acid. The gradient for the analysis was as follows: 2% B at 0 min; 2% B at 1 min; 42% B at 4 min; 100%
B at 5 min; 100% B at 6 min; 2% B at 6.5 min; and 0% B at 8 min. The column temperature was maintained at
30°C, with a flow rate of 0.4 mL/min and an injection volume of 3 µL per run. Nucleotide metabolite profiling

was performed using a 6495 LC/TQ mass spectrometer utilizing the dynamic MRM scan mode. Source parameter settings for both positive and negative ion modes were as follows: capillary voltage at 3.5 kV, gas temperature at 120°C, gas flow at 11 L/min, nebulizer pressure at 40 psi, sheath gas temperature at 400°C, and sheath gas flow at 12 L/min. The Krebs cycle metabolites analysis was performed using gas chromatography coupled with mass spectrometry (GC-MS). Purified and dried samples were reconstituted in 200 µL of an 80:20 UPLC methanol/UPLC water solution, and 10 µL of d27-Myristic acid was added as an internal standard (75 µg/mL; 400505, Agilent). The samples were then transferred into brown crimp vials with inserts and dried using a SpeedVac. Derivatization was performed by a PAL autosampler as follows: 90 µL of methoxyamine hydrochloride was added to the samples (20 g/L; 89803, Sigma-Aldrich), which were incubated at 30°C for 90 min. Subsequently, 10 µL of MSTFA (TS-48915; Thermo Scientific) was added, followed by incubation at 37°C for 30 min. Chromatographic analysis was performed on an Agilent 8890 GC system equipped with a DB-MS + DG column (0.25 µm film thickness, L × I.D. 30 m × 0.250 mm; 122-5532G, Agilent). One microliter of the prepared sample was injected in split mode (10:1), with the inlet temperature maintained at 250°C. Helium was used as the carrier gas at a flow rate of 1.1 mL/min. The GC temperature program started at 60°C for 1 min, followed by a ramp of 60°C/min to 120°C, then 30°C/min to 250°C, and finally 60°C/min to 325°C, with a 5-min hold at the final temperature. Krebs cycle metabolite profiling was conducted using a 5977B MSD mass spectrometer with dynamic MRM scan mode. The temperature was set to 250°C, and the electron energy was maintained at 70 eV. The peak area counts were normalized based on their respective internal standard control compounds for each detected metabolite from the GM, Nucleotides, and Krebs cycle. The *'MetaboAnalystR'* package (v.4.0.0) (Chong & Xia, 2018; Chong et al., 2019; Pang et al., 2020) was used to filter compounds based on an interquartile range (IQR) cutoff of < 25, removing 10% of the data. Subsequently, quantile normalization was applied, followed by a logarithmic transformation.

Statistical analyses
Statistical analyses were performed using R software (v.4.4.1). Plots were primarily generated using the *'ggplot2'* package (v.3.5.1) (Wickham, 2009) unless otherwise. To ensure data quality, samples falling outside the 95% confidence ellipse in PCA were considered outliers and subsequently removed.

Survival curves
Survival curves (Kaplan-Meier model) were generated using the *'survfit'* and plotted using the *'ggsurvplot'* functions from the *'survival'* (v.3.2-11) (Therneau & Lumley, 2015) and the *'survminer'* (v.0.4.9) packages (Kassambara et al., 2017), respectively. The Cox proportional hazards model was performed using the *'coxph'* function from the *'survival'* package.

Viral DNA and RNA load in oyster
Normality and homogeneity of variance for viral DNA loads and RNA transcription were evaluated using the functions of *'shapiro.test'* from the *'stats'* (v.4.4.1) and *'levenetest'* from the *'car'* packages (Fox & Weisberg, 2019), respectively. A one-way ANOVA was conducted with the *'aov'* function to compare viral DNA and RNA loads across age groups, followed by Tukey's Honest Significant Difference (HSD) test using the *'Tukeyhsd'* function from *Stats'* package.

Weighted Correlation Network Analysis
To identify biologically relevant co-expression modules associated with age, we applied a Weighted Correlation Network Analysis (WGCNA) using the *'WGCNA'* package (v.1.73) (Langfelder & Horvath, 2008). This analysis was performed across the three omics layers per family: epigenetics (DNA methylation), transcriptomics (gene expression), and metabolomics (metabolites). For all the omics, the trait of interest was the age, and samples were re-coded as follows: 4 months =1, 16 months = 2, and 28 months = 3. The associations between the network modules and age were assessed using the Pearson (r) correlation ($P < 0.05$). For each dataset, an unsupervised co-expression network was constructed using the following parameters: maxBlockSize = 30,000, networkType = signed, TOMtype = signed, mergeCutHeight = 0.25, and minModuleSize = 100. A soft power threshold was selected to achieve a scale-free topology, optimizing the identification of biologically meaningful network structures. Modules showing either significant positive or negative correlations with age were retained,

and when multiple modules shared the same direction, they were merged to generate a unified list of age-associated positive and negative features (i.e., gene methylation, gene expression and metabolites). To evaluate the biological significance of age-associated modules in epigenetic and transcriptomic analyses, we retrieved the kME values (eigengene connectivity) of all genes present in the significant modules, considering both positive and negative associations. The analysis included all 30,418 coding genes, assigning kME values to the genes retrieved from the significant modules, while giving a value of 0 for the genes outside the modules. We use the GO_MWU package (github.com/z0on/GO_MWU) (Wright et al., 2015), focusing on the enrichment analysis of Biological Process (BP) GO-terms with the following parameters: largest = 0.5, smallest = 10, clusterCutHeight = 0.25, Module = TRUE, and Alternative = g. The BP GO-terms were significant at *P.adj* < 0.05. To assess the impact of age on metabolomics, we used the normalized metabolite data from the *'MetaboAnalystR'* package as input for WGCNA analysis, identifying significant modules associated with age. To simplify and summarize the list of significant BP GO-terms obtained in epigenetics and transcriptomics from GO_MWU, we categorized the BP GO-terms into broader groups based on the Gene Ontology database (**Table S3**). This grouping, based on the current understanding of oyster–POMS interactions, further supports the rationale for organizing these categories according to established knowledge of POMS pathogenesis (de Lorgeril et al., 2018, 2020; Duperret et al., 2025).

The metabolites within the significant modules associated with age were further used as an input list for pathway reconstruction and enrichment analysis using the *'FELLA'* package (v.1.22.0) (Picart-Armada et al., 2018), based on a diffusion algorithm. Specifically, each input metabolite introduced a unitary flow model that propagated through intermediate KEGG entities (including reactions, enzymes, and modules) to identify biological pathways in which the metabolite is involved. Once identified, we extracted the genes annotated to participate in the significantly enriched KEGG pathways by querying KEGG through the 'KEGGREST' package (v.1.46.0) (Tenenbaum, 2017) with database *M. gigas* (T03920), thereby enabling the integration of metabolomic findings with the corresponding genes.

Results

Ageing reduces viral replication in POMS-susceptible oyster families

Following the three infection challenges (one for each age: 4-, 16-, and 28-month-old), mortality decreased with ageing for the four families (**Figures 1B–C**). For each infection challenge, both the 4-month-old NSI donors and the 4-month-old NSI recipient oysters (positive control) deployed in the experimental tanks showed similar survival curves across the three experiments (**Figure S1A–B**), confirming that the mortality patterns observed in the four families tested were not biased by differences in experimental infections (virus load or infectious efficiency). Only the F14R and H2D were selected for further analysis due to their contrasting age‐related gains of survival (**Figure 1B**). Key infection indicators revealed that 28-month-old oysters () in H2D showed significantly lower viral DNA loads (One-way ANOVA: $F_{\square,\square\square} = 5.9$, $P < 0.05$) and reduced viral transcriptional activity ($F_{\square,\square\square} = 8.2$, $P < 0.01$) compared with their 4‐month‐old siblings (**Figure 1D**). Whereas age-related differences are not significant in F14R (viral DNA: $F_{\square,\square\square} = 0.74$, $P = 0.50$; and viral transcription: $F_{\square,\square\square} = 1.4$, $P = 0.28$), although a trend toward a decrease in viral load and transcriptional activity was apparent in the oldest oysters (**Figure 1D**). Overall, our findings demonstrated that ageing limits OsHV-1 µVar replication associated with increased survival. To investigate the molecular mechanisms underlying age-related changes influencing permissiveness to POMS, we analyzed the methylome, transcriptome, and metabolome of 4-, 16-, and 28-month-old oysters just before the infection challenge (0 hpc),

In the epigenetic dataset, each sample yielded an average of 84,991,198 ± 8,765,131 trimmed reads, with mapping of 42.87 ± 2.52% and a duplicate read rate of 15.20 ± 3.35%. The enzymatic conversion efficiency estimated by the lambda genome spike-in positive control was 99.18% ± 3.62% (**Table S4**). We identified 2,164,482 CpG sites in F14R and 2,300,773 CpG sites in H2D that were shared across the three age groups in each family. Both families exhibited age-related methylation patterns (**Figures S2A** and **S2B**). For further analysis, we averaged the methylation values at the gene level (within each gene's body region) for each sample and age group, resulting in two gene-level methylation lists (26,104 genes for F14R and 26,534 genes for H2D, **Table S5** and **Table S6**, respectively). As for the transcriptomic data, each sample yielded an average of 86.65%

±⬜1.32% alignment rate (**Table S4**) and showed a clear age-dependent pattern of gene expression in both
families (**Figures S2C** and **S2D**). For the metabolomic data, based on 76 filtered out of the initial 114
metabolites, a strong separation between 4-month-old and older oysters (16- and 28-month-old) was observed,
highlighting a remarkable age-related metabolic shift (**Figures S2E** and **S2F**). Our results revealed that ageing
induced profound changes in methylome, gene expression, and metabolism, which could govern physiological
changes influencing permissiveness to POMS. Outlier removal resulted in the exclusion of two epigenetic, one
transcriptomic, and four metabolomic samples (**Table S4**).
382
Ageing drives a shift from a permissive to a non-permissive state to POMS, reshaping critical
pathways that enhance the host's ability to cope with POMS
Ageing drives a shift from a permissive to a non-permissive state to POMS, reshaping critical pathways that
enhance the host's ability to cope with infection. To investigate the molecular basis of this age-dependent shift,
we sought to integrate epigenetic, transcriptomic, and metabolomic layers. Since changes in DNA methylation,
gene expression, and metabolite profiles are expected to be coordinated and functionally linked, we applied a
Weighted Gene Co-methylation/Co-expression/Co-metabolite Network Analysis. This systems-level approach
allowed us to identify age-associated molecular modules significantly correlated (positively or negatively) with
age. To further characterize the biological meaning of these modules, we performed enrichment analyses of BP
GO-terms, revealing functional pathways that either converge or diverge between families.
393
Epigenetics reflects age-dependent changes in immunity
At the epigenetic level for F14R, we identified one positively correlated module (693 methylation genes; r =
0.73; *P* < 0.001) and two negatively correlated modules (574 and 425 methylation genes; r = –0.78 and –0.50; *P*
< 0.001 and 0.05, respectively) with age (**Figure S3A** and **Table S7**). For H2D, we identified two positively
correlated modules (660 and 163 methylation genes; r = 0.55 and 0.63; both *P* < 0.05) and one negatively
correlated module (2,085 methylation genes; r = –0.86; *P* < 0.001) with age (**Figure S3B** and **Table S7**). From
these gene methylation sets, 10 were overrepresented and 9 underrepresented GO-terms for the F14R, while it
was 10 and 23 GO terms for the H2D, respectively (**Table S8**). The BP GO-term analysis revealed biological
processes that are either common or specific to each family. For both families, genes with increased DNA
methylation levels were enriched in biological regulation, developmental processes, responses to stimuli, and
immune system processes. Among the genes showing decreased DNA methylation levels with age, the BP GO-
terms were mostly related to cellular processes and localization. Divergent epigenetic signatures mainly
concerned the H2D, where several biological processes linked to biological regulation, developmental processes,
and reproduction were enriched among genes with decreased DNA methylation levels, while processes related to
localization were enriched among genes where DNA methylation levels increased with age (**Figure 2A–2B**).
Overall, these results aligned with biological expectations of ageing in bivalves, showing a shift from cellular
proliferation and structural modification in younger oysters to a remodeling of metabolism, stress responses, and
immune functions in the older ones.
412
Given the pivotal role of immune and stress responses in host-pathogen interactions in oysters, we focused on
the genes belonging to the Response to stimulus and Immune system process GO-terms. Between the 693 and
823 positively, and the 999 and 2,085 negatively age-associated methylation genes in F14R and H2D,
respectively (**Figures S3A–3B**), a total of 19 (13 + 6) immune- and stress-related genes were shared (**Figure**
S4A). These shared genes were involved in pathogen recognitions, such as the C1q domain-containing protein
(G18427), complement C1q tumor necrosis factor-related protein 2 (G3352), and ficolin-2 (G22089); in immune
signaling as the inhibitor of nuclear factor kappa-B kinase subunit α (IKKα, G1756), serine/threonine-protein
kinase H1 homolog (G20959), interferon regulatory factor 8 (IRF8, G23210), TNF receptor-associated factor 2
(TRAF2, G25366), serine/threonine-protein kinase TBK1 (G30487); and in immune regulation as the E3
ubiquitin ligases TRIM63 (G10197), TRIM45 (G22126), TRIM56 (G25021), tripartite motif-containing protein
2 (G15281), RNF146 (G2646), FMRFamide receptor (G3282), interleukin-6 receptor subunit β (G628), G-
protein coupled receptor family 1 domain-containing protein (G6283) B-cell lymphoma/leukemia 11A BCL11A
(**Figure S4A**). Among the shared genes, we examined the IRF8 (G23210), TBK1 (G30487), RNF146 (G2646),
G-protein coupled receptor (G6283), IKKα (G1756), and ficolin-2 (G22089), and all genes displayed aligned

age-associated DNA methylation trends (increases or decreases) in both families (**Figure 2C**). Additionally, we
detected Toll-like receptors (TLRs), critical components of innate immune recognition, which also exhibited
notable age-associated methylation patterns, such as the toll-like receptor 4 (G26336) in F14R and the toll-like
receptor 4 (G9660 and G9701) and toll-like receptor 13 (G21066) in H2D. This family-specific modulation of
TLR at the methylation level may reflect nuanced changes in pathogen sensing and immune responsiveness
across the lifespan, underscoring the complexity of immune epigenetic regulation in oyster host-pathogen
interactions. Overall, these results indicated that age-associated DNA methylation changes also occur in
immune- and stress-related genes, suggesting that epigenetic regulation at the methylome level during ageing
contributes to the modulation of the immune landscape in oysters. Such epigenetic regulation may also influence
the capacity of oysters to respond to OsHV-1 µVar infection, potentially explaining why the older oysters
displayed reduced permissiveness to POMS compared to the younger ones. To investigate whether these age-
related epigenetic modifications are also reflected at the transcriptional level, either gene-to-gene or biological
functions, we analyzed the gene expression patterns across the same age groups.

Figure 2. Age-associated DNA methylation modules and Gene Ontology enrichment (Epigenetics)
Heatmaps of age-associated DNA methylation (Epigenetics)
Heatmaps displaying genes with positive (red) or negative (blue) associations between DNA methylation and age, alongside a summary of enriched Gene Ontology (GO) terms of the Biological Process category for **A)** F14R and **B)** H2D. Complete GO-term lists are available in Table S6. **C)** Examples of immune/stress genes showing concordant age-associated methylation trends in both families.

Ageing-associated transcriptomic shifts reflect metabolic and immunity remodeling
At the transcriptomic level in F14R, we identified 14,546 genes grouped into six modules correlated with age: two positively (6,468 and 1,420 genes; r = 0.96 and 0.50; $P < 0.001$ and 0.05, respectively) and four negatively (4,517; 1,380; 463; and 298 genes; r = –0.96, –0.70, –0.80, and –0.55; $P < 0.001$, 0.01, 0.001, and 0.05, respectively) modules (**Figure S3C** and **Table S9**). In the H2D, we identified 11,533 genes grouped into nine modules correlated with age: five positive (2,253; 1,975; 1,558; 313; and 214 genes; r = 0.93, 0.75, 0.76, 0.63, and 0.51; $P < 0.001$, 0.001, 0.001, 0.01, and 0.05, respectively) and four negative modules (3,184; 1,771; 133; and 105 genes; r = –0.91, –0.91, –0.49, and –0.53; $P < 0.001$, 0.001, 0.05, and 0.05, respectively) (**Figure S3D** and **Table S9**). The enrichment analysis revealed 167 positively and 175 negatively BP GO-terms in F14R, and 95 positively and 195 negatively BP GO-terms in H2D correlated with age (**Table S10**). Among these, 181 BP GO-terms (77 positive and 104 negative) were in common (**Table S11**). To synthesize the important number of BP GO-terms, as we did in epigenetics, we classified the BP GO-terms from transcriptomic analysis into broader functional categories. Among the significant BP GO-terms, those most directly involved between ageing and susceptibility to POMS included over-representation pathways in the Immune system process (e.g., Cell surface receptor signaling pathway and Defense response to symbiont) and antiviral immunity (e.g., Defense response to virus), and to the under-representation of Toll-like receptors and Myd88 (e.g., Toll signaling, and MyD88-dependent toll-like receptor signaling pathways), along with regulation of canonical NF-kappaB signal transduction. DNA repair mechanisms (e.g., DNA damage response and Regulation of DNA repair) were exclusively positively associated with age, suggesting a conserved activation of these protective pathways across both families. The response to stimulus and homeostatic processes was in both positive and negative sets of genes; however, the results were quite nonspecific among the genes positively associated with age (e.g., Cellular response to stimulus). A strong and specific under-expression of genes in the response to oxidative stress and the maintenance of homeostasis processes was also evidenced (e.g., Response to oxidative stress, the Wnt signaling, and Cellular homeostasis). Additionally, the enrichment analysis underscored the regulation of a broad spectrum of metabolic processes. Among the genes positively associated with ageing, those involved in protein catabolism (e.g., Proteolysis involved in protein catabolic process) and nucleotide metabolism (e.g., Nucleobase-containing compound metabolic process) were evident. For the genes negatively associated with age, the metabolism of protein processes was shown to be under-represented (e.g., Proteolysis) as well as the metabolism of lipids (e.g., Lipid metabolic process). Between families, differences in metabolic processes were more pronounced in H2D, which showed a marked under-expression of genes involved in energy production (e.g., ATP biosynthetic process, Generation of precursor metabolites and energy, and Energy derivation by oxidation of organic compounds) (**Figure 3A–3B**).

We also extended our analysis to the genes exhibiting transcriptional regulation involved in Response to stimulus and Immune system process GO-terms in the context of POMS. Between the 7,888 and 6,340 positively, and the 6,658 and 5,193 genes negatively age-associated in F14R and H2D (**Figures S3C–D**), 174 (150 + 24) immune- and stress-related genes were shared (**Figure S4B** and **Table S12**). Specifically, we found age-associated increases in the expression of 150 genes, including the interferon regulatory factor 1 (IRF8, G748), DNA damage-regulated autophagy modulator protein 2 (G6254), a set involving E3 ubiquitin-protein ligases genes such as (TRIM33, G4565), (UBR2, G12475), and (TRIP12, G16081) which are key regulators of protein turnover and immune signaling, and ATP-dependent RNA and DNA helicases genes including (DDX42, G34539), (DHX8, G31806) and (DDX31, G27896) genes, which are essential for RNA metabolism, DNA repair, and antiviral defense (**Figure 3C**). On the other hand, we observed an age-associated decrease in the expression of 24 genes in innate immune signaling, including multiple genes of myeloid differentiation primary response protein MyD88 (e.g., G6951 and G5576) and related to Toll-like receptor (TLR) pathways such as TLR 4 (G28) and TLR 6 (G26605) (**Figure 3C**). Overall, these results indicated that ageing in oysters orchestrates a

complex and coordinated biological program, marked by enhanced DNA repair capacity, a reshaped immune landscape with heightened antiviral response genes, and reduced NF-κB signaling. In the context of POMS, such gene expression reprogramming may promote the improved survival of older oysters by strengthening intracellular antiviral defenses and repair mechanisms, while tempering the excessive inflammatory stress responses. Consistent with previous observations in ageing in bivalves (Abele et al., 2009), we also detected a decline in energetic metabolism and antioxidant defenses. Given the critical role of energy metabolism in determining susceptibility to POMS in oysters, we next performed a metabolomic analysis across the same age groups.

Figure 3. Age-associated gene-expression modules and GO enrichment (Transcriptomics) Heatmaps of age-associated gene expression (Transcriptomics).

Heatmaps displaying genes with positive (red) or negative (blue) associations between gene expression and age, alongside a summary of enriched Gene Ontology (GO) terms of the Biological Process category for **A)** F14R and **B)** H2D. Complete GO-term lists are available in Table S7. **C)** Examples of immune/stress genes showing concordant age-associated gene expression trends in both families.

Global metabolomic analysis reveals age-dependent decline in mitochondrial function and central metabolism

At the metabolomic level in F14R, 21 metabolites were clustered into a module positively correlated ($r = 0.94$, $P < 0.001$), while 35 metabolites were in two modules negatively correlated with age ($r = -0.90$ and -0.76, $P < 0.001$) (**Figure S3E**). In H2D, 23 and 26 metabolites were grouped into one positive and one negative module ($r = 0.85$ and -0.92, $P < 0.001$, respectively) correlated with age (**Figure S3F**). The most pronounced differences were between the 4-month-old oysters and the older age groups (16- and 28-month-old) (**Figure 4**), where a metabolic shift was evident in key amino acids (e.g., alanine, creatine, cysteine, argininosuccinic acid) and nucleotides (e.g., AMP, GMP, UMP), as well as in energy-related metabolites (e.g., NAD☐). Notably, a progressive decline was observed in several TCA cycle intermediates, such as fumarate, malate, and citrate, indicating a reduction in mitochondrial oxidative metabolism with age. Consistent with transcriptomic data, this metabolic shift was more pronounced in H2D. A set of 37 (15 + 22) metabolites was shared between the two families, including central metabolic intermediates involved in ATP generation, redox homeostasis (e.g., glutathione derivatives), nucleotide turnover, and amino acid metabolism (**Figure S5**). These patterns revealed a conserved metabolic aging signature characterized by mitochondrial downregulation, altered redox balance, and remodeling of amino acid and nucleotide metabolism. To gain deeper functional insights into the metabolic reprogramming associated with ageing in oysters, we performed enrichment analysis of the positively and negatively age-associated metabolites using the *'FELLA'* package. This approach integrates metabolomic profiles with the *M. gigas* KEGG database, enabling the identification of enriched KEGG pathways. Results revealed family-specific KEGG pathways, such as the Toll-like receptor signaling pathway in F14R (**Figure S6A**) and Pyruvate metabolism in H2D (**Figure S6B**). To obtain a broader view of metabolism and ageing in oysters, we analyzed the 37 common age-associated metabolites shared by both families. This analysis identified KEGG pathways, including those involved in amino acid metabolism, carbohydrate metabolism (e.g., the TCA cycle), and nucleotide metabolism. Pathways related to genetic information processing, such as RNA degradation, protein export, and mRNA surveillance, were also enriched. Notably, we observed enrichment of cellular processes previously linked to POMS outcomes, including autophagy, mitophagy, peroxisome activity, and the mTOR signaling pathway, all of which are central to antiviral defense, oxidative stress control, and energy homeostasis in oysters. Regulation of neuroactive ligand–receptor interaction and related signaling pathways further suggests an interplay between metabolic processes and neuroendocrine regulation, which may modulate immune and stress response (**Figure S6C**). Together, these findings highlighted a core set of metabolic and regulatory networks potentially underpinning age-dependent permissiveness to POMS.

Figure 4. Age-associated metabolites (Metabolomics).

Heatmaps displaying metabolites with positive (red) or negative (blue) associations with age for **A)** F14R and **B)** H2D.

Integration of multi-omics: linking epigenomics, transcriptomics, and metabolomics with signals of ageing influencing permissiveness to POMS

Epigenetic and transcriptomic relationship in age-dependent modulation of immune and stress responses

To explore the relationship between epigenomic and transcriptomic layers, we overlapped the list of genes showing age-related methylation changes with those exhibiting age-associated expression patterns for both positive and negative associations. This gene-to-gene overlap between epigenomic and transcriptomic lists revealed 855 genes in F14R and 1,282 genes in H2D, but only 74 were in common between the two families. Although we identified the same set of genes associated with age at both the epigenetic and transcriptomic levels, it has been consistently shown in invertebrates that DNA methylation levels do not directly correlate with gene expression levels (Dixon & Matz, 2022). It seems more plausible that DNA methylation interacts with other epigenetic modifications, such as chromatin accessibility, and there exists a more complex interrelationship with epigenetics and transcription regulation in invertebrates (Bogan et al., 2023; G. Wang & Mai, 2025). Moreover, genetic backgrounds may differ between families, and regulatory mechanisms can vary accordingly; nevertheless, these differences often converge on similar biological functions by mobilizing different members of multigenic families (de Lorgeril et al., 2020). To characterize the relationship between age-associated changes in DNA methylation and the gene expression at the biological function and the context of POMS, we focused on the significant BP GO-terms belonging to Response to stimulus and Immune system process in both omics. This targeted approach provided an integrated view of immune- and stress-related processes potentially modulating the permissiveness to POMS in oysters in ageing. Notably, several biological processes, including Defense response to virus, Response to symbiont, Response to oxidative stress, and Chemokine-mediated signaling, displayed consistent directionality for both omics layers in both families, indicating the existence of a core of

567 age-modulated pathways (**Figure 5**). The recurrence of the same biological functions in two independent genetic
568 backgrounds reinforces their potential roles as fundamental components of the immune landscape in oysters.
569 Together, these observations supported the hypothesis that age-related epigenetic modifications may act as
570 upstream regulatory layers, either promoting or repressing transcriptional programs that shape the organism's
571 permissiveness to POMS.
572

573
574 **Figure 5. Gene Ontology (GO-terms) enrichment of age-associated immune and stimulus-response genes.**
575 Biological Process GO terms related to Response to stimulus and Immune system process, derived from genes
576 with positive (red) or negative (blue) associations with age. Results are presented separately for DNA
577 methylation (epigenomic) and gene expression (transcriptomic) datasets in each family (F14R and H2D).
578
Coordinated age-related metabolic shifts highlight the regulation of oxidative stress and redox imbalance
To validate the functional relevance of the enriched KEGG pathways identified through metabolomics, we
compiled all genes annotated to these pathways and cross-referenced them with our transcriptomic datasets to
determine age-dependent expression dynamics. Given the large number of KEGG pathways, we focused our
validation on seven key pathways: Carbohydrate metabolism (citrate cycle, TCA cycle; crg00020), Nucleotide
metabolism (purine metabolism, crg00230; and pyrimidine metabolism, crg00240), Amino acid metabolism
(alanine, aspartate, and glutamate metabolism, crg00250; and arginine and proline metabolism, crg00330),
Metabolism of other amino acids (taurine and hypotaurine metabolism, crg00430), and Glutathione metabolism
(crg00480). These pathways have been linked to POMS permissiveness in relation to temperature, the major
driver of the disease (Duperret et al., 2025) (**Figures 6**, **S7,** and **S8**).
589

590
591 **Figure 6. Metabolic map integrating transcriptomic and metabolomic data associated with age in F14R**
592 **and H2D.**
593 Schematic representation of metabolite and gene expression changes associated with ageing. The map illustrates
594 metabolites and corresponding gene expression changes linked to age. Metabolites that accumulate (higher
595 abundance) in 28-month-old oysters are shown in orange, whereas those without accumulation (low abundance)
596 are shown in black. Metabolites detected in both families are highlighted in bold. Genes are color-coded
597 according to fold change (FC) at a false discovery rate (FDR) < 0.05, using 4-month-old oysters as the reference
598 group F14R (Blue) and H2D (Green). Arrows represent direct metabolic conversions. Family-specific results are
599 shown separately in Figure S7 (F14R) and Figure S8 (H2D).
600

The concordance between metabolite levels and corresponding gene expression patterns associated with younger and older conditions strongly supports our integrative multi-omics approach and underscores the biological relevance of the identified pathways in age-dependent metabolic modulation within the context of POMS in oysters (**Figures 6**, **S7**, and **S8**).

Integrating transcriptomic and metabolomic data revealed a coherent age-associated reprogramming of central carbon and amino acid metabolism, reflecting altered mitochondrial function during ageing in oysters. The accumulation of TCA cycle intermediates with age (metabolomics: citrate, malate, and fumarate) coincided with the down-expression of multiple downstream genes, such as the succinate-CoA ligase (ADP-forming) subunit β mitochondrial (G16455), the isocitrate dehydrogenase (NAD) subunit β mitochondrial (G24230), the isocitrate dehydrogenase (NADP) mitochondrial-like (G5092), and the isocitrate dehydrogenase (NADP) cytoplasmic (G4888). These enzymes are involved in mitochondrial oxidative phosphorylation, suggesting a slowdown of TCA cycle flux of energy and reduced metabolic turnover in older oysters. In parallel, increased glutamate and glutamine levels in older oysters suggested a shift toward nitrogen storage, likely reflecting reduced amino acid turnover due to diminished TCA cycle activity. This interpretation is supported by the under-expression of the mitochondrial glutamate dehydrogenase gene (G3271) identified in both families, a key link between amino acid metabolism and the TCA cycle. Interestingly, the accumulation of glutamate and glutamine did not lead to enhanced antioxidant capacity, as levels of both reduced (GSH) and oxidized (GSSG) glutathione were lower in older oysters compared to younger ones, indicating regulated antioxidant defense. In both families, we also observed accumulation of L-Proline, accompanied by reduced L-Hydroxyproline levels and strong over-expression of the trans-L-3-hydroxyproline dehydratase gene (G10696) in older oysters. This enzyme catalyzes the dehydration of hydroxyproline, a metabolite derived from proline hydroxylation, which may serve as an adaptive strategy to buffer oxidative stress and preserve cellular integrity under diminished metabolic capacity. Finally, we detected accumulation of L-Asparagine alongside marked under-expression of the isoaspartyl peptidase/L-asparaginase gene (G13495). This was associated with reduced levels of L-Arginosuccinate and L-Arginine, two essential intermediates in arginine biosynthesis. Notably, at least in the F14R, over-expression of the nitric oxide synthase gene (G23422), homologous to salivary gland NOS, suggests an increased demand for L-Arginine as a substrate for the nitric oxide (NO) production. NO is a key mediator of innate immune responses and redox signaling. The observed metabolic imbalance, characterized by excessive L-Asparagine retention, limited arginine biosynthesis, and regulation of NO-generating pathways, suggests a complex age-associated reprogramming of nitrogen metabolism.

Taken together, the integration of epigenomic, transcriptomic, and metabolomic datasets reveals a coherent, multi-layered regulatory program associated with ageing in oysters. Epigenetic modifications appear to act as upstream regulators that shape transcriptional landscapes, particularly in immune- and stress-related pathways, thereby influencing the organism's ability to cope with OsHV-1 µVar. These transcriptional programs are further mirrored in metabolic reorganization, where shifts in central carbon, amino acid, and redox metabolism reflect the functional consequences of gene expression regulation. The convergence across omics layers demonstrates that age-dependent changes are tightly interconnected, together driving a progressive transition toward reduced viral and altered physiological state, influencing the permissiveness to POMS in oysters.

Discussion

This study aims to elucidate the molecular mechanisms by which ageing in *Magallana gigas* influences the permissiveness to Pacific Oyster Mortality Syndrome (POMS). The older oysters consistently displayed improved survival and markedly reduced OsHV-1 µVar replication compared with younger individuals of the same genetic background. This age-dependent protective effect, observed across all four families, aligns with previous reports of reduced POMS-related mortality in adult oysters (Arzul et al., 2002; Peeler et al., 2012; Pernet et al., 2012; Dégremont, 2013). To investigate the molecular basis of age-dependent differences in permissiveness to POMS, we performed an integrative multi-omics analysis (epigenetics, transcriptomics, and metabolomics) on two genetically distinct oyster families with contrasted phenotypes regarding permissiveness to POMS (F14R and H2D).

Epigenetic and transcriptomic remodeling drives age-related immune trade-offs, influencing POMS permissiveness in oysters

The role of epigenetic mechanisms in innate immunity remains poorly understood. However, growing evidence suggests that epigenetic regulation acts as a key modulator of immune responses in invertebrates, including oysters (Gómez-Díaz et al., 2012; Fallet et al., 2020; Gu et al., 2022). Our analysis revealed age-associated DNA methylation changes in genes involved in cellular processes, metabolism, environmental stress responses, and immune functions, consistent with a previous study highlighting the contribution of epigenetic modulation to POMS pathogenesis in oysters (Fallet et al., 2022). We found that ageing progressively reshapes the oyster DNA methylation landscape, with pronounced changes in immune-related genes. Notably, several Toll-like receptor (TLR) genes, key pattern recognition receptors, and the adaptor gene Myd88 all exhibited increased DNA methylation levels with age. This finding supports previous findings linking epigenetic modulation of the TLR–NF-κB–Myd88 signaling to resistant and susceptible phenotypes in the POMS context (Gawra et al., 2023; Valdivieso et al., 2025).

Moreover, genome-wide DNA methylation profiling has shown that the NF-κB pathway is epigenetically regulated during *Vibrio alginolyticus* infections in oysters (J. Li et al., 2024), supporting the idea that age-dependent methylation shapes innate immune signaling and contributes to the improved outcomes observed in adult oysters during POMS outbreaks (Tang et al., 2020). DNA methylation is already recognized as essential during early oyster development (Riviere et al., 2013), and its gradual remodeling during ageing is likely to be not only a marker of age but also a driver of transcriptional reprogramming (Saint-Carlier & Riviere, 2015). While direct correlations between methylation and gene expression in invertebrates are often weak or absent (X. Wang et al., 2014; Dixon & Matz, 2022), evidence suggests that DNA methylation interacts with other epigenetic mechanisms to regulate chromatin accessibility (Badeaux & Shi, 2013). Such interactions may profoundly shape the relationship between epigenetic and transcriptional regulation (Bogan et al., 2023), ultimately influencing oyster defense capacity against POMS, as supported by our findings. The lack of gene-to-gene biological correlations between DNA methylation and gene expression regulation is a recurrent observation in oysters and other invertebrates (Gavery & Roberts, 2010; Johnson et al., 2022). Previous studies have shown that different genetic backgrounds in oysters can strongly influence the expression of specific genes within multigenic families, leading to different gene-level signatures but ultimately converging on similar biological functions (de Lorgeril et al., 2020). Together, these findings support the view that integrative multi-omics approaches are more informative when interpreted at the functional pathway or process level, rather than focusing solely on one-to-one gene relationships.

Ageing was also accompanied by extensive transcriptional reprogramming, affecting key biological processes that influence permissiveness to POMS in oysters. Specifically, these transcriptomic changes included an enhanced DNA repair capacity (Zhao et al., 2023) and a reshaped immune landscape characterized by the over-expression of antiviral defense genes (e.g., Defense/Response to virus, Defense response to symbiont) and the under-expression of the NF-κB signaling components, including genes in the Toll signaling pathway, MyD88-dependent Toll-like receptor signaling, and cytokinesis-related processes. These processes have been widely reported in oyster immunity (Huang et al., 2019) and in POMS pathogenesis in oysters (Green & Speck, 2018; de Lorgeril et al., 2018, 2020; Duperret et al., 2025). The attenuation of gene expression involved in the NF-κB signaling in adults suggests a dampened inflammatory response (Rothschild et al., 2018), potentially reflecting a shift from broad pro-inflammatory programs in young oysters toward more targeted antiviral defenses in older oysters. In *M. gigas*, this pathway plays roles in developmental, cellular, and immune response processes (Yu et al., 2018), and it is mostly mediated by the TLR (L. Zhang et al., 2011). This immune reprogramming, together with the over-expression of virus recognition and response genes, may represent a strategy that enhances viral control while limiting inflammatory damage (de Lorgeril et al., 2020). Indeed, host immune gene expression dynamics are key determinants of infection outcome, influencing both pathogen proliferation and disease progression (Alizon et al., 2011). These dynamics are also observed from our results, where age-associated epigenetic and transcriptional remodeling shifts, enabling older oysters to exert stronger control over OsHV-1 μVar, whereas younger individuals remain highly susceptible, as reflected by higher mortality rates and viral replication. However, this ontogenetic shift in immune prioritization may involve a life-stage trade-off. Young

704 oysters appear biased toward antibacterial defenses, while older ones may favor antiviral responses, as depicted
705 with epidemiological observations that juveniles were more susceptible to OsHV-1 μVar, whereas adults were
706 predominantly affected by *Vibrio aestuarianus* (Azéma et al., 2017; Green et al., 2016).
707

Age-dependent redox and metabolic reprogramming shape immune responses and susceptibility to
POMS

In bivalves, the immune defense mainly relies on phagocytosis by haemocytes by enzymatic or oxidative
degradation (Canesi et al., 2002). Upon activation, the oyster immune cells generate reactive oxygen species
(ROS), a highly reactive molecule produced as byproducts of mitochondrial respiration and enzymatic activity
(Sussarellu et al., 2013). While ROS production is crucial for pathogen elimination, their excessive accumulation
can damage host tissues (C. Li & Jackson, 2002), making their tight regulation essential. This balance is
achieved through the action of antioxidant enzymes, which sustain the cellular redox state. In *M. gigas*, redox
homeostasis has been identified as a central integrator of stress adaptation, linking antioxidant defenses with
energy metabolism, immunity, and development (Trevisan & Mello, 2024). Our findings are consistent with this
view, highlighting that redox regulation not only safeguards cellular integrity, but also shapes the metabolic and
immune responses during oyster aging, therefore influencing permissiveness to POMS. Understanding the
intricate relationship between immune, metabolic pathways, and redox balance is pivotal for environmental
stress induced by pathogens in oysters, especially in the context of some redox-mediated immuno-metabolic
adaptations that may alter the susceptibility.
723

The integration of transcriptomic and metabolomic data revealed age-associated regulation of glutathione
metabolism, underscoring its central role in shaping the cellular redox environment and influencing
susceptibility to POMS (Samain, 2011). Glutathione (GSH), a tripeptide composed of glutamate, cysteine, and
glycine, is a key antioxidant that neutralizes reactive oxygen species (ROS) through reversible redox cycling
between its reduced (GSH) and oxidized (GSSG) forms. An optimal GSH/GSSG ratio is critical for maintaining
redox homeostasis, protecting macromolecules from oxidative damage, and supporting cellular detoxification
(Trevisan & Mello, 2024). However, excessive reduction (reductive stress) can also be deleterious, disrupting
redox-sensitive signaling and protein folding, and causing DNA damage (Xiao & Loscalzo, 2020). In our
dataset, younger oysters exhibited higher concentrations of both GSH and GSSG, reflecting a more dynamic
redox buffering system and greater antioxidant capacity at early life stages compared to older oysters. This redox
flexibility may enhance protection against oxidative stress and support immune activation during OsHV-1
infection. However, at these early stages, the organisms remain largely unable to control the infection, resulting
in higher mortalities. These findings are consistent with the observation that ageing is associated with the over-
expression of genes involved in the DNA repair process, likely reflecting a compensatory mechanism to
counteract cumulative oxidative damage and maintain genome integrity. Older oysters displayed lower
GSH/GSSG levels, consistent with a less reducing intracellular environment and altered immune signaling, as
reflected in their transcriptomic profiles in genes involved in response to virus. The metabolic shifts in glycine
and glutamate in old oysters, which are necessary precursors of GSH, further indicate age-related remodeling of
antioxidant defenses (Rebrin & Sohal, 2008). These adaptations likely fine-tune the nuclear factor kappa-light-
chain-enhancer of activated B cells (NF-κB) signaling (Asehnoune et al., 2004; Deponte, 2013; Lingappan,
2018), a process also observed in the transcriptomic data, thereby enhancing immune responsiveness. The Nitric
oxide (NO) metabolism also emerged as a redox-linked pathway modulated by age. NO is synthesized by nitric
oxide synthases from L-arginine and serves as both a potent antimicrobial effector and a sentinel of
environmental stress (Nakayama & Maruyama, 1998; Ottaviani et al., 1993; Rivero, 2006; Tafalla et al., 2003).
In young oysters, we showed elevated L-arginine levels and higher nitric oxide synthase expression, suggesting
enhanced NO production and immune responsiveness. Conversely, older oysters exhibited reduced L-arginine
availability, accumulation of L-asparagine, and down-expression of isoaspartyl peptidase/L-asparaginase, an
enzyme driving L-asparagine degradation into L-aspartate in the argininosuccinate–arginine pathway, indicating
a reduced flux toward NO synthesis. Transcriptomic analyses also revealed that multiple biosynthetic and
metabolic pathways, including lipid metabolism, catabolic processes, the TCA cycle, and ATP metabolism, were
negatively associated with age. The metabolomic data further illustrated a rewiring of the TCA cycle in oysters,
with fumarate, malate, citrate, and α-ketoglutarate accumulated in older oysters, whereas succinate was more

abundant in younger ones. Malate, derived from aspartate, succinate or glucose, succinate pathways, enters the mitochondria and can be processed through two complementary routes: an oxidative branch, generating NADH and ATP via substrate-level phosphorylation and complex I–mediated proton pumping, or a reductive branch, which regenerates succinate and uses fumarate as an alternative electron acceptor under low oxygen conditions (Müller et al., 2012; Trevisan & Mello, 2024). These metabolic routes highlight the flexibility of oyster mitochondrial metabolism, which can shift between oxidative and reductive modes in response to changing energy demands. The observed metabolite patterns may suggest a reduced oxidative flux and energy-conserving metabolism in older oysters, contrasted with a more active aerobic profile in younger ones.

mTOR signaling as a central hub linking metabolism, autophagy, and immune defense in oyster aging

Our integrated omics analyses point to the mechanistic target of rapamycin (mTOR) pathway as a central regulator of age-associated changes in the oyster physiology. The mTOR pathway is a well-established master regulator of growth, metabolism, and cellular senescence, integrating environmental and cellular signals to coordinate anabolic processes, mitochondrial activity, and stress responses (Weichhart, 2018; Papadopoli et al., 2019). The observed age-dependent decline in energy metabolism and altered redox regulation in old oysters may reflect changes in the mTOR signaling activity, as cellular energy and nutritional state directly regulate mTOR to control associated signaling pathways downstream (Q. Zhang et al., 2023) and autophagy in bivalves (Kim et al., 2011; Levine & Deretic, 2007). In the context of POMS, starved oysters enhanced autophagy, improving their capacity to cope with the OsHV-1 µVar infection by limiting viral replication (Pernet et al., 2019). In oysters, autophagy is functional and modulates OsHV-1 µVar proliferation (Leprêtre et al., 2021; Moreau et al., 2015; Picot et al., 2019, 2020, 2022). In this study, we identified several genes involved in the autophagy/mTOR pathway displaying age-related changes at the epigenetic and transcriptional level, such as the inhibitor of nuclear factor kappa-B kinase subunit alpha (IKKα, G1756) and rapamycin-insensitive companion of mTOR (Rictor, G270) in both families. In addition, we found 28 and 34 genes downregulated with age in F14R and H2D, respectively, that are implicated in the KEGG mTOR signaling pathway, underscoring widespread transcriptional repression of this central regulatory network with age. Together, these findings highlight that mTOR-associated signaling components are broadly modulated during oyster ageing, potentially altering metabolic regulation, immune signaling, and autophagy, and providing a molecular basis for age-dependent variation in permissiveness to POMS.

Conclusion

This multi-omics study demonstrates that ageing in *Magallana gigas* is characterized by adaptive reprogramming that reduces permissiveness to Pacific Oyster Mortality Syndrome (POMS). By integrating epigenomic, transcriptomic, and metabolomic data, we show that epigenetic remodeling of key immune regulators, such as Toll-like receptors and MyD88, aligns with transcriptional rewiring of NF-κB and ubiquitin pathways, indicating fine-tuned control of innate immunity. Our data also provide strong evidence that mTOR signaling is repressed with age, likely promoting autophagy and enhancing viral control, thereby improving outcomes during OsHV-1 infection. This regulatory shift is tightly interconnected with metabolic reprogramming, including reduced tricarboxylic acid (TCA) cycle flux, remodeling of nitrogen metabolism, and altered glutathione dynamics, collectively supporting a stress-tolerant, energy-conserving phenotype. Together, these results reveal an evolutionary trade-off between growth and immune defense: juveniles' metabolic activity increases their susceptibility to viral proliferation, whereas adults shift their resources toward cellular maintenance and antiviral preparedness, resulting in improved survival during POMS outbreaks.

Data availability

RNA-seq data and EM-seq analyses have been made available through the ENA database under Bioproject DECICOMP: PRJEB86647.

EM-seq data were submitted under accession numbers PRJEB105019.

RNA-seq data were submitted under accession numbers PRJEB105132.

Metabolomic data have been made available through Metabolight database under the accession number MTBLS13463

808 Complementary information is available from the corresponding authors on reasonable request.
809

Credit author statement

Funding acquisition: G.M.
Conceptualization: B.P., J.V.D, and G.M.
Methodology: B.P., J.V.D, and G.M
Investigation: A.V., L.D., E.T., A.L., J.V.D., and G.M.
Supervision: E.T., A.L., J.V.D., and G.M
Writing original draft: A.V., J.V.D.
Writing review and editing: all authors
818

Declaration of competing interests

820 The authors declare that they have no known financial conflicts of interest or personal relationships that could
821 have influenced the work reported in this paper.
822

Funding

824 The present study was supported by the ANR project DECICOMP (ANR-19-CE20-0004) and within the
825 framework of the "Laboratoire d'Excellence (LABEX)" TULIP (ANR-10-LABX-41).
826

Acknowledgments

We would like to thank the staff of the Ifremer experimental platforms at Argenton and Bouin for providing
experimental facilities and for the production of the biological material used in this study. We thank Bruno
Chollet, Louis Boismorand, Charlotte Corporeau, Mathias Huber, Marianne Alunno-Bruscia and Moussa Diagne
for assistance in the experimental design and procedure. We thank the Montpellier Alliance for Metabolomics
and Metabolism Analysis (MAMMA) platform (BioCampus Montpellier) for technical support in the
metabolomic analysis.
834

References

Abbadi, M., Zamperin, G., Gastaldelli, M., Pascoli, F., Rosani, U., Milani, A., Schivo, A., Rossetti, E., Turolla,
E., Gennari, L., Toffan, A., Arcangeli, G., & Venier, P. (2018). Identification of a newly described OsHV-1 µvar
from the North Adriatic Sea (Italy). Journal of General Virology, 99(5), 693–703.
https://doi.org/10.1099/jgv.0.001042
Abele, D., Brey, T., & Philipp, E. (2009). Bivalve models of aging and the determination of molluscan lifespans.
Experimental Gerontology, 44(5), 307–315. https://doi.org/10.1016/j.exger.2009.02.012
Abele, D., & Philipp, E. (2013). Environmental control and control of the environment: The basis of longevity in
bivalves. Gerontology, 59(3), 261–266. https://doi.org/10.1159/000345331
Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick, A., & Mason, C. E.
(2012). MethylKit: A comprehensive R package for the analysis of genome-wide DNA methylation profiles.
Genome Biology, 13(10), R87. https://doi.org/10.1186/gb-2012-13-10-r87
Alizon, S., Luciani, F., & Regoes, R. R. (2011). Epidemiological and clinical consequences of within-host
evolution. Trends in Microbiology, 19(1), 24–32. https://doi.org/10.1016/j.tim.2010.09.005
Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data [Computer software].
Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom.
Arzul, I., Renault, T., Thébault, A., & Gérard, A. (2002). Detection of oyster herpesvirus DNA and proteins in
asymptomatic Crassostrea gigas adults. Virus Research, 84(1–2), 151–160. https://doi.org/10.1016/S0168-
1702(02)00007-2
Asehnoune, K., Strassheim, D., Mitra, S., Kim, J. Y., & Abraham, E. (2004). Involvement of Reactive Oxygen
Species in Toll-Like Receptor 4-Dependent Activation of NF-κB. The Journal of Immunology, 172(4), 2522–
2529. https://doi.org/10.4049/jimmunol.172.4.2522

Azéma, P., Lamy, J.-B., Boudry, P., Renault, T., Travers, M.-A., & Dégremont, L. (2017). Genetic parameters of
resistance to Vibrio aestuarianus, and OsHV-1 infections in the Pacific oyster, Crassostrea gigas, at three
different life stages. Genetics Selection Evolution, 49(1), 23. https://doi.org/10.1186/s12711-017-0297-2
Badeaux, A. I., & Shi, Y. (2013). Emerging roles for chromatin as a signal integration and storage platform.
Nature Reviews Molecular Cell Biology, 14(4), 211–224. https://doi.org/10.1038/nrm3545
Bogan, S. N., Strader, M. E., & Hofmann, G. E. (2023). Associations between DNA methylation and gene
regulation depend on chromatin accessibility during transgenerational plasticity. BMC Biology, 21(1), 149.
https://doi.org/10.1186/s12915-023-01645-8
Burioli, E. A. V., Prearo, M., & Houssin, M. (2017). Complete genome sequence of Ostreid herpesvirus type 1
μVar isolated during mortality events in the Pacific oyster Crassostrea gigas in France and Ireland. Virology,
509, 239–251. https://doi.org/10.1016/j.virol.2017.06.027
Canesi, L., Gallo, G., Gavioli, M., & Pruzzo, C. (2002). Bacteria–hemocyte interactions and phagocytosis in
marine bivalves. Microscopy Research and Technique, 57(6), 469–476. https://doi.org/10.1002/jemt.10100
Carrasco, N., Gairin, I., Pérez, J., Andree, K. B., Roque, A., Fernández-Tejedor, M., Rodgers, C. J., Aguilera, C.,
& Furones, M. D. (2017). A production calendar based on water temperature, spat size, and husbandry practices
reduce OsHV-1 μvar impact on cultured Pacific oyster Crassostrea gigas in the Ebro delta (Catalonia),
Mediterranean coast of Spain. Frontiers in Physiology, 8. https://doi.org/10.3389/fphys.2017.00125
Chong, J., & Xia, J. (2018). MetaboAnalystR: An R package for flexible and reproducible analysis of
metabolomics data. Bioinformatics, 34(24), 4313–4314. https://doi.org/10.1093/bioinformatics/bty528
Chong, J., Yamamoto, M., & Xia, J. (2019). MetaboAnalystR 2.0: From Raw Spectra to Biological Insights.
Metabolites, 9(3), 57. https://doi.org/10.3390/metabo9030057
Clerissi, C., Luo, X., Lucasson, A., Mortaza, S., De Lorgeril, J., Toulza, E., Petton, B., Escoubas, J.-M.,
Dégremont, L., Gueguen, Y., Destoumieux-Garzón, D., Jacq, A., & Mitta, G. (2023). A core of functional
complementary bacteria infects oysters in Pacific Oyster Mortality Syndrome. Animal Microbiome, 5(1), 26.
https://doi.org/10.1186/s42523-023-00246-8
de Lorgeril, J., Lucasson, A., Petton, B., Toulza, E., Montagnani, C., Clerissi, C., Vidal-Dupiol, J., Chaparro, C.,
Galinier, R., Escoubas, J.-M., Haffner, P., Dégremont, L., Charrière, G. M., Lafont, M., Delort, A., Vergnes, A.,
Chiarello, M., Faury, N., Rubio, T., … Mitta, G. (2018). Immune-suppression by OsHV-1 viral infection causes
fatal bacteraemia in Pacific oysters. Nature Communications, 9(1), 4215. https://doi.org/10.1038/s41467-018-
06659-3
de Lorgeril, J., Petton, B., Lucasson, A., Perez, V., Stenger, P.-L., Dégremont, L., Montagnani, C., Escoubas, J.-
M., Haffner, P., Allienne, J.-F., Leroy, M., Lagarde, F., Vidal-Dupiol, J., Gueguen, Y., & Mitta, G. (2020).
Differential basal expression of immune genes confers Crassostrea gigas resistance to Pacific oyster mortality
syndrome. BMC Genomics, 21(1), 63. https://doi.org/10.1186/s12864-020-6471-x
Dégremont, L. (2013). Size and genotype affect resistance to mortality caused by OsHV-1 in Crassostrea gigas.
Aquaculture, 416–417, 129–134. https://doi.org/10.1016/j.aquaculture.2013.09.011
Dégremont, L., Nourry, M., & Maurouard, E. (2015). Mass selection for survival and resistance to OsHV-1
infection in Crassostrea gigas spat in field conditions: Response to selection after four generations. Aquaculture,
446, 111–121. https://doi.org/10.1016/j.aquaculture.2015.04.029
Delisle, L., Laroche, O., Hilton, Z., Burguin, J.-F., Rolton, A., Berry, J., Pochon, X., Boudry, P., & Vignier, J.
(2022). Understanding the dynamic of POMS infection and the role of microbiota composition in the survival of
Pacific Oysters, Crassostrea gigas. Microbiology Spectrum, 10(6), e01959-22.
https://doi.org/10.1128/spectrum.01959-22
Deponte, M. (2013). Glutathione catalysis and the reaction mechanisms of glutathione-dependent enzymes.
Biochimica et Biophysica Acta (BBA) - General Subjects, 1830(5), 3217–3266.
https://doi.org/10.1016/j.bbagen.2012.09.018
Divilov, K., Schoolfield, B., Mancilla Cortez, D., Wang, X., Fleener, G. B., Jin, L., Dumbauld, B. R., &
Langdon, C. (2021). Genetic improvement of survival in Pacific oysters to the Tomales Bay strain of OsHV-1
over two cycles of selection. Aquaculture, 543, 737020. https://doi.org/10.1016/j.aquaculture.2021.737020
Divilov, K., Schoolfield, B., Morga, B., Dégremont, L., Burge, C. A., Mancilla Cortez, D., Friedman, C. S.,
Fleener, G. B., Dumbauld, B. R., & Langdon, C. (2019). First evaluation of resistance to both a California
OsHV-1 variant and a French OsHV-1 microvariant in Pacific oysters. BMC Genetics, 20(1), 96.
https://doi.org/10.1186/s12863-019-0791-3

Dixon, G., & Matz, M. (2022). Changes in gene body methylation do not correlate with changes in gene expression in Anthozoa or Hexapoda. BMC Genomics, 23(1), 234. https://doi.org/10.1186/s12864-022-08474-z

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. Bioinformatics, 29(1), 15–21. https://doi.org/10.1093/bioinformatics/bts635

Dowle, M., Srinivasan, A., Gorecki, J., Chirico, M., Stetsenko, P., Short, T., Lianoglou, S., Antonyan, E., Bonsch, M., & Parsonage, H. (2019). Package 'data. Table'. Extension of 'data. Frame, 596.

Duperret, L., Valdivieso, A., Kunselman, E., Petton, B., Morga, B., De Lorgeril, J., Pernet, F., Degremont, L., Faury, N., Allienne, J.-F., Pouzadoux, J., Romatif, O., Courtay, G., Monaco, C. J., Toulza, E., Delisle, L., Vidal-Dupiol, J., Lagorce, A., & Mitta, G. (2025). Deciphering the molecular mechanisms of oyster resistance to Pacific Oyster Mortality Syndrome (POMS) disease induced by high temperatures. Science of The Total Environment, 994, 180026. https://doi.org/10.1016/j.scitotenv.2025.180026

Durinck, S., Spellman, P. T., Birney, E., & Huber, W. (2009). Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt. Nature Protocols, 4(8), 1184–1191. https://doi.org/10.1038/nprot.2009.97

Fallet, M., Luquet, E., David, P., & Cosseau, C. (2020). Epigenetic inheritance and intergenerational effects in mollusks. Gene, 729, 144166. https://doi.org/10.1016/j.gene.2019.144166

Fallet, M., Montagnani, C., Petton, B., Dantan, L., De Lorgeril, J., Comarmond, S., Chaparro, C., Toulza, E., Boitard, S., Escoubas, J.-M., Vergnes, A., Le Grand, J., Bulla, I., Gueguen, Y., Vidal-Dupiol, J., Grunau, C., Mitta, G., & Cosseau, C. (2022). Early life microbial exposures shape the Crassostrea gigas immune system for lifelong and intergenerational disease protection. Microbiome, 10(1), 85. https://doi.org/10.1186/s40168-022-01280-5

Fox, J., & Weisberg, S. (2019). An R companion to applied regression. Sage publications.

Gavery, M. R., & Roberts, S. B. (2010). DNA methylation patterns provide insight into epigenetic regulation in the Pacific oyster (Crassostrea gigas). BMC Genomics, 11(1), 483. https://doi.org/10.1186/1471-2164-11-483

Gavery, M. R., & Roberts, S. B. (2014). A context dependent role for DNA methylation in bivalves. Briefings in Functional Genomics, 13(3), 217–222. https://doi.org/10.1093/bfgp/elt054

Gawra, J., Valdivieso, A., Roux, F., Laporte, M., De Lorgeril, J., Gueguen, Y., Saccas, M., Escoubas, J.-M., Montagnani, C., Destoumieux-Garzón, D., Lagarde, F., Leroy, M. A., Haffner, P., Petton, B., Cosseau, C., Morga, B., Dégremont, L., Mitta, G., Grunau, C., & Vidal-Dupiol, J. (2023). Epigenetic variations are more substantial than genetic variations in rapid adaptation of oyster to Pacific oyster mortality syndrome. Science Advances, 9(36), eadh8990. https://doi.org/10.1126/sciadv.adh8990

Gittenberger, A., Voorbergen☐Laarman, M. A., & Engelsma, M. Y. (2016). Ostreid herpesvirus Os HV ☐1 μVar in Pacific oysters Crassostrea gigas (Thunberg 1793) of the Wadden Sea, a UNESCO world heritage site. Journal of Fish Diseases, 39(1), 105–109. https://doi.org/10.1111/jfd.12332

Gómez-Díaz, E., Jordà, M., Peinado, M. A., & Rivero, A. (2012). Epigenetics of host–pathogen interactions: The road ahead and the road behind. PLoS Pathogens, 8(11), e1003007. https://doi.org/10.1371/journal.ppat.1003007

Gotz, S., Garcia-Gomez, J. M., Terol, J., Williams, T. D., Nagaraj, S. H., Nueda, M. J., Robles, M., Talon, M., Dopazo, J., & Conesa, A. (2008). High-throughput functional annotation and data mining with the Blast2GO suite. Nucleic Acids Research, 36(10), 3420–3435. https://doi.org/10.1093/nar/gkn176

Green, T., & Speck, P. (2018). Antiviral defense and innate immune memory in the oyster. Viruses, 10(3), 133. https://doi.org/10.3390/v10030133

Green, T., Vergnes, A., Montagnani, C., & De Lorgeril, J. (2016). Distinct immune responses of juvenile and adult oysters (Crassostrea gigas) to viral and bacterial infections. Veterinary Research, 47(1), 72. https://doi.org/10.1186/s13567-016-0356-7

Gu, X., Qiao, X., Yu, S., Song, X., Wang, L., & Song, L. (2022). Histone lysine-specific demethylase 1 regulates the proliferation of hemocytes in the oyster Crassostrea gigas. Frontiers in Immunology, 13, 1088149. https://doi.org/10.3389/fimmu.2022.1088149

Gutierrez, A. P., Symonds, J., King, N., Steiner, K., Bean, T. P., & Houston, R. D. (2020). Potential of genomic selection for improvement of resistance to ostreid herpesvirus in Pacific oyster (Crassostrea gigas). Animal Genetics, 51(2), 249–257. https://doi.org/10.1111/age.12909

Huang, B., Zhang, L., Xu, F., Tang, X., Li, L., Wang, W., Liu, M., & Zhang, G. (2019). Oyster versatile IKKα/βs are involved in toll-like receptor and RIG-I-like receptor signaling for innate immune response. Frontiers in Immunology, 10, 1826. https://doi.org/10.3389/fimmu.2019.01826

Husmann, G., Abele, D., Rosenstiel, P., Clark, M. S., Kraemer, L., & Philipp, E. E. R. (2014). Age-dependent expression of stress and antimicrobial genes in the hemocytes and siphon tissue of the Antarctic bivalve, Laternula elliptica, exposed to injury and starvation. Cell Stress and Chaperones, 19(1), 15–32. https://doi.org/10.1007/s12192-013-0431-1

Hwang, J. Y., Park, J. J., Yu, H. J., Hur, Y. B., Arzul, I., Couraleau, Y., & Park, M. A. (2013). Ostreid herpesvirus 1 infection in farmed Pacific oyster larvae (Crassostrea virginica) (Thunberg) in Korea. Journal of Fish Diseases, 36(11), 969–972. https://doi.org/10.1111/jfd.12093

Johnson, K. M., Sirovy, K. A., & Kelly, M. W. (2022). Differential DNA methylation across environments has no effect on gene expression in the eastern oyster. Journal of Animal Ecology, 91(6), 1135–1147. https://doi.org/10.1111/1365-2656.13645

Kassambara, A., Kosinski, M., Biecek, P., & Fabian, S. (2017). Package 'survminer'. Drawing Survival Curves Using 'Ggplot2'(R Package Version 03 1).

Keeling, S., Brosnahan, C., Williams, R., Gias, E., Hannah, M., Bueno, R., McDonald, W., & Johnston, C. (2014). New Zealand juvenile oyster mortality associated with ostreid herpesvirus 1—An opportunistic longitudinal study. Diseases of Aquatic Organisms, 109(3), 231–239. https://doi.org/10.3354/dao02735

Kim, J., Kundu, M., Viollet, B., & Guan, K.-L. (2011). AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. Nature Cell Biology, 13(2), 132–141. https://doi.org/10.1038/ncb2152

Krueger, F., & Andrews, S. R. (2011). Bismark: A flexible aligner and methylation caller for Bisulfite-seq applications. Bioinformatics, 27(11), 1571–1572. https://doi.org/10.1093/bioinformatics/btr167

Krueger, F., James, F., Ewels, P., Afyounian, E., Weinstein, M., Schuster-Boeckler, B., Hulselmans, G., & Sclamons. (2023). FelixKrueger/TrimGalore: V0.6.10 - add default decompression path (Version 0.6.10) [Computer software]. Zenodo. https://doi.org/10.5281/ZENODO.5127898

Kunselman, E., Manrique, D., Burge, C. A., Allard, S., Daniel, Z., Mitta, G., Petton, B., & Gilbert, J. A. (2024). Temperature and microbe mediated impacts of the San Diego Bay ostreid herpesvirus (OsHV-1) microvariant on juvenile Pacific oysters. Sustainable Microbiology, 1(1), qvae014. https://doi.org/10.1093/sumbio/qvae014

Langfelder, P., & Horvath, S. (2008). WGCNA: An R package for weighted correlation network analysis. BMC Bioinformatics, 9(1), 559. https://doi.org/10.1186/1471-2105-9-559

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. Nature Methods, 9(4), 357–359. https://doi.org/10.1038/nmeth.1923

Langmead, B., Trapnell, C., Pop, M., & Salzberg, S. L. (2009). Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biology, 10(3), R25. https://doi.org/10.1186/gb-2009-10-3-r25

Leprêtre, M., Faury, N., Segarra, A., Claverol, S., Degremont, L., Palos-Ladeiro, M., Armengaud, J., Renault, T., & Morga, B. (2021). Comparative proteomics of ostreid herpesvirus 1 and Pacific oyster interactions with two families exhibiting contrasted susceptibility to viral infection. Frontiers in Immunology, 11, 621994. https://doi.org/10.3389/fimmu.2020.621994

Levine, B., & Deretic, V. (2007). Unveiling the roles of autophagy in innate and adaptive immunity. Nature Reviews Immunology, 7(10), 767–777. https://doi.org/10.1038/nri2161

Li, C., & Jackson, R. M. (2002). Reactive species mechanisms of cellular hypoxia-reoxygenation injury. American Journal of Physiology-Cell Physiology, 282(2), C227–C241. https://doi.org/10.1152/ajpcell.00112.2001

Li, J., He, Y., Yang, B., Mokrani, A., Li, Y., Tan, C., Li, Q., & Liu, S. (2024). Whole-genome DNA methylation profiling revealed epigenetic regulation of NF-κB signaling pathway involved in response to Vibrio alginolyticus infection in the Pacific oyster, Crassostrea gigas. Fish & Shellfish Immunology, 151, 109705. https://doi.org/10.1016/j.fsi.2024.109705

Lingappan, K. (2018). NF-κB in oxidative stress. Current Opinion in Toxicology, 7, 81–86. https://doi.org/10.1016/j.cotox.2017.11.002

López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M., & Kroemer, G. (2013). The hallmarks of aging. Cell, 153(6), 1194–1217. https://doi.org/10.1016/j.cell.2013.05.039

López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M., & Kroemer, G. (2023). Hallmarks of aging: An
expanding universe. Cell, 186(2), 243–278. https://doi.org/10.1016/j.cell.2022.11.001

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. Genome Biology, 15(12), 550. https://doi.org/10.1186/s13059-014-0550-8

Mahmoudi, S., & Brunet, A. (2012). Aging and reprogramming: A two-way street. Current Opinion in Cell
Biology, 24(6), 744–756. https://doi.org/10.1016/j.ceb.2012.10.004

Männer, L., Schell, T., Provataris, P., Haase, M., & Greve, C. (2021). Inference of DNA methylation patterns in
molluscs. Philosophical Transactions of the Royal Society B: Biological Sciences, 376(1825), 20200166.
https://doi.org/10.1098/rstb.2020.0166

Martenot, C., Oden, E., Travaillé, E., Malas, J.-P., & Houssin, M. (2011). Detection of different variants of
Ostreid Herpesvirus 1 in the Pacific oyster, Crassostrea gigas between 2008 and 2010. Virus Research, 160(1–2),
25–31. https://doi.org/10.1016/j.virusres.2011.04.012

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.Journal, 17(1), Article 1. https://doi.org/10.14806/ej.17.1.200

Moreau, P., Moreau, K., Segarra, A., Tourbiez, D., Travers, M.-A., Rubinsztein, D. C., & Renault, T. (2015).
Autophagy plays an important role in protecting Pacific oysters from OsHV-1 and Vibrio aestuarianus
infections. Autophagy, 11(3), 516–526. https://doi.org/10.1080/15548627.2015.1017188

Mortensen, S., Strand, Å., Bodvin, T., Alfjorden, A., Skår, C., Jelmert, A., Aspán, A., Sælemyr, L., Naustvoll,
L., & Albretsen, J. (2016). Summer mortalities and detection of ostreid herpesvirus microvariant in Pacific
oyster Crassostrea gigas in Sweden and Norway. Diseases of Aquatic Organisms, 117(3), 171–176.
https://doi.org/10.3354/dao02944

Müller, M., Mentel, M., Van Hellemond, J. J., Henze, K., Woehle, C., Gould, S. B., Yu, R.-Y., Van Der Giezen,
M., Tielens, A. G. M., & Martin, W. F. (2012). Biochemistry and evolution of anaerobic energy metabolism in
eukaryotes. Microbiology and Molecular Biology Reviews, 76(2), 444–495.
https://doi.org/10.1128/MMBR.05024-11

Nakayama, K., & Maruyama, T. (1998). Differential production of active oxygen species in photo-symbiotic and
non-symbiotic bivalves. Developmental & Comparative Immunology, 22(2), 151–159.
https://doi.org/10.1016/s0145-305x(97)00060-8

Ottaviani, E., Paeman, L. R., Cadet, P., & Stefano, G. B. (1993). Evidence for nitric oxide production and
utilization as a bacteriocidal agent by invertebrate immunocytes. European Journal of Pharmacology:
Environmental Toxicology and Pharmacology, 248(4), 319–324. https://doi.org/10.1016/0926-6917(93)90006-c

Oyanedel, D., Lagorce, A., Bruto, M., Haffner, P., Morot, A., Labreuche, Y., Dorant, Y., De La Forest Divonne,
S., Delavat, F., Inguimbert, N., Montagnani, C., Morga, B., Toulza, E., Chaparro, C., Escoubas, J.-M., Gueguen,
Y., Vidal-Dupiol, J., De Lorgeril, J., Petton, B., … Destoumieux-Garzón, D. (2023). Cooperation and cheating
orchestrate Vibrio assemblages and polymicrobial synergy in oysters infected with OsHV-1 virus. Proceedings
of the National Academy of Sciences, 120(40), e2305195120. https://doi.org/10.1073/pnas.2305195120

Pang, Z., Chong, J., Li, S., & Xia, J. (2020). MetaboAnalystR 3.0: Toward an optimized workflow for global
metabolomics. Metabolites, 10(5), 186. https://doi.org/10.3390/metabo10050186

Papadopoli, D., Boulay, K., Kazak, L., Pollak, M., Mallette, F., Topisirovic, I., & Hulea, L. (2019). mTOR as a
central regulator of lifespan and aging. F1000Research, 8, 998. https://doi.org/10.12688/f1000research.17196.1

Peeler, E. J., Allan Reese, R., Cheslett, D. L., Geoghegan, F., Power, A., & Thrush, M. A. (2012). Investigation
of mortality in Pacific oysters associated with Ostreid herpesvirus-1μVar in the Republic of Ireland in 2009.
Preventive Veterinary Medicine, 105(1–2), 136–143. https://doi.org/10.1016/j.prevetmed.2012.02.001

Peñaloza, C., Gutierrez, A. P., Eöry, L., Wang, S., Guo, X., Archibald, A. L., Bean, T. P., & Houston, R. D.
(2021). A chromosome-level genome assembly for the Pacific oyster Crassostrea gigas. GigaScience, 10(3),
giab020. https://doi.org/10.1093/gigascience/giab020

Pepin, J. F. (2013). European Union Reference Laboratory for Molluscs Diseases, IFREMER,. https://www.eurl-
mollusc.eu/content/download/57054/795216/file/

Pernet, F., Barret, J., Le Gall, P., Corporeau, C., Dégremont, L., Lagarde, F., Pépin, J., & Keck, N. (2012). Mass
mortalities of Pacific oysters Crassostrea gigas reflect infectious diseases and vary with farming practices in the
Mediterranean Thau lagoon, France. Aquaculture Environment Interactions, 2(3), 215–237.
https://doi.org/10.3354/aei00041

Pernet, F., Tamayo, D., Fuhrmann, M., & Petton, B. (2019). Deciphering the effect of food availability, growth and host condition on disease susceptibility in a marine invertebrate. Journal of Experimental Biology, jeb.210534. https://doi.org/10.1242/jeb.210534

Petton, B., Bruto, M., James, A., Labreuche, Y., Alunno-Bruscia, M., & Le Roux, F. (2015). Crassostrea gigas mortality in France: The usual suspect, a herpes virus, may not be the killer in this polymicrobial opportunistic disease. Frontiers in Microbiology, 6. https://doi.org/10.3389/fmicb.2015.00686

Petton, B., Destoumieux-Garzón, D., Pernet, F., Toulza, E., De Lorgeril, J., Degremont, L., & Mitta, G. (2021). The Pacific oyster mortality syndrome, a polymicrobial and multifactorial disease: State of knowledge and future directions. Frontiers in Immunology, 12, 630343. https://doi.org/10.3389/fimmu.2021.630343

Petton, B., Pernet, F., Robert, R., & Boudry, P. (2013). Temperature influence on pathogen transmission and subsequent mortalities in juvenile Pacific oysters Crassostrea gigas . Aquaculture Environment Interactions, 3(3), 257–273. https://doi.org/10.3354/aei00070

Philipp, E. E. R., & Abele, D. (2010). Masters of longevity: Lessons from long-lived bivalves – a mini-review. Gerontology, 56(1), 55–65. https://doi.org/10.1159/000221004

Picart-Armada, S., Fernández-Albert, F., Vinaixa, M., Yanes, O., & Perera-Lluna, A. (2018). FELLA: An R package to enrich metabolomics data. BMC Bioinformatics, 19(1), 538. https://doi.org/10.1186/s12859-018-2487-5

Picot, S., Faury, N., Arzul, I., Chollet, B., Renault, T., & Morga, B. (2020). Identification of the autophagy pathway in a mollusk bivalve, Crassostrea gigas. Autophagy, 16(11), 2017–2035. https://doi.org/10.1080/15548627.2020.1713643

Picot, S., Faury, N., Pelletier, C., Arzul, I., Chollet, B., Dégremont, L., Renault, T., & Morga, B. (2022). Monitoring autophagy at cellular and molecular level in Crassostrea gigas during an experimental ostreid herpesvirus 1 (OsHV-1) infection. Frontiers in Cellular and Infection Microbiology, 12, 858311. https://doi.org/10.3389/fcimb.2022.858311

Picot, S., Morga, B., Faury, N., Chollet, B., Dégremont, L., Travers, M.-A., Renault, T., & Arzul, I. (2019). A study of autophagy in hemocytes of the Pacific oyster, Crassostrea gigas. Autophagy, 15(10), 1801–1809. https://doi.org/10.1080/15548627.2019.1596490

Putri, G. H., Anders, S., Pyl, P. T., Pimanda, J. E., & Zanini, F. (2022). Analysing high-throughput sequencing data in Python with HTSeq 2.0. Bioinformatics, 38(10), 2943–2945. https://doi.org/10.1093/bioinformatics/btac166

Rebrin, I., & Sohal, R. S. (2008). Pro-oxidant shift in glutathione redox state during aging. Advanced Drug Delivery Reviews, 60(13–14), 1545–1552. https://doi.org/10.1016/j.addr.2008.06.001

Rico-Villa, B., Le Coz, J. R., Mingant, C., & Robert, R. (2006). Influence of phytoplankton diet mixtures on microalgae consumption, larval development and settlement of the Pacific oyster Crassostrea gigas (Thunberg). Aquaculture, 256(1–4), 377–388. https://doi.org/10.1016/j.aquaculture.2006.02.015

Rivero, A. (2006). Nitric oxide: An antiparasitic molecule of invertebrates. Trends in Parasitology, 22(5), 219–225. https://doi.org/10.1016/j.pt.2006.02.014

Riviere, G., Wu, G.-C., Fellous, A., Goux, D., Sourdaine, P., & Favrel, P. (2013). DNA methylation is crucial for the early development in the oyster C. gigas. Marine Biotechnology, 15(6), 739–753. https://doi.org/10.1007/s10126-013-9523-2

Roque, A., Carrasco, N., Andree, K. B., Lacuesta, B., Elandaloussi, L., Gairin, I., Rodgers, C. J., & Furones, M. D. (2012). First report of OsHV-1 microvar in Pacific oyster (Crassostrea gigas) cultured in Spain. Aquaculture, 324–325, 303–306. https://doi.org/10.1016/j.aquaculture.2011.10.018

Rothschild, D. E., McDaniel, D. K., Ringel-Scaia, V. M., & Allen, I. C. (2018). Modulating inflammation through the negative regulation of NF-κB signaling. Journal of Leukocyte Biology, 103(6), 1131–1150. https://doi.org/10.1002/JLB.3MIR0817-346RRR

Rubio, T., Oyanedel, D., Labreuche, Y., Toulza, E., Luo, X., Bruto, M., Chaparro, C., Torres, M., De Lorgeril, J., Haffner, P., Vidal-Dupiol, J., Lagorce, A., Petton, B., Mitta, G., Jacq, A., Le Roux, F., Charrière, G. M., & Destoumieux-Garzón, D. (2019). Species-specific mechanisms of cytotoxicity toward immune cells determine the successful outcome of Vibrio infections. Proceedings of the National Academy of Sciences, 116(28), 14238–14247. https://doi.org/10.1073/pnas.1905747116

Saint-Carlier, E., & Riviere, G. (2015). Regulation of Hox orthologues in the oyster Crassostrea gigas evidences a functional role for promoter DNA methylation in an invertebrate. FEBS Letters, 589(13), 1459–1466. https://doi.org/10.1016/j.febslet.2015.04.043

Samain, J.-F. (2011). Review and perspectives of physiological mechanisms underlying genetically-based resistance of the Pacific oyster Crassostrea gigas to summer mortality. Aquatic Living Resources, 24(3), 227–236. https://doi.org/10.1051/alr/2011144

Schikorski, D., Renault, T., Saulnier, D., Faury, N., Moreau, P., & Pépin, J.-F. (2011). Experimental infection of Pacific oyster Crassostrea gigas spat by ostreid herpesvirus 1: Demonstration of oyster spat susceptibility. Veterinary Research, 42(1), 27. https://doi.org/10.1186/1297-9716-42-27

Segarra, A., Pépin, J. F., Arzul, I., Morga, B., Faury, N., & Renault, T. (2010). Detection and description of a particular Ostreid herpesvirus 1 genotype associated with massive mortality outbreaks of Pacific oysters, (Crassostrea gigas), in France in 2008. Virus Research, 153(1), 92–99. https://doi.org/10.1016/j.virusres.2010.07.011

Suquet, M., de Kermoysan, G., Araya, R. G., Queau, I., Lebrun, L., Le Souchu, P., & Mingant, C. (2009). Anesthesia in Pacific oyster, Crassostrea gigas. Aquatic Living Resources, 22(1), 29–34. https://doi.org/10.1051/alr/2009006

Sussarellu, R., Dudognon, T., Fabioux, C., Soudant, P., Moraga, D., & Kraffe, E. (2013). Rapid mitochondrial adjustments in response to short-term hypoxia and re-oxygenation in the Pacific oyster Crassostrea gigas. Journal of Experimental Biology, jeb.075879. https://doi.org/10.1242/jeb.075879

Tafalla, C., Gómez-León, J., Novoa, B., & Figueras, A. (2003). Nitric oxide production by carpet shell clam (Ruditapes decussatus) hemocytes. Developmental & Comparative Immunology, 27(3), 197–205. https://doi.org/10.1016/s0145-305x(02)00098-8

Tang, X., Huang, B., Lin, S., Wang, W., Zhang, G., & Li, L. (2020). CgMyD88s serves as an innate immune system plug during Ostreid Herpesvirus 1 infection in the Pacific oyster (Crassostrea gigas). Frontiers in Immunology, 11, 1247. https://doi.org/10.3389/fimmu.2020.01247

Tenenbaum, D. (2017). KEGGREST [Computer software]. Bioconductor. https://doi.org/10.18129/B9.BIOC.KEGGREST

Therneau, T. M., & Lumley, T. (2015). Package 'survival'. R Top Doc, 128(10), 28–33.

Trevisan, R., & Mello, D. F. (2024). Redox control of antioxidants, metabolism, immunity, and development at the core of stress adaptation of the oyster Crassostrea gigas to the dynamic intertidal environment. Free Radical Biology and Medicine, 210, 85–106. https://doi.org/10.1016/j.freeradbiomed.2023.11.003

Turtoi, E., Jeudy, J., Henry, S., Dadi, I., Valette, G., Enjalbal, C., & Turtoi, A. (2023). Analysis of polar primary metabolites in biological samples using targeted metabolomics and LC-MS. STAR Protocols, 4(3), 102400. https://doi.org/10.1016/j.xpro.2023.102400

Turtoi, E., Jeudy, J., Valette, G., Enjalbal, C., Vila, I. K., Laguette, N., & Turtoi, A. (2023). Multiplexed targeted analysis of polyunsaturated fatty acids and oxylipins using liquid chromatography-tandem mass spectrometry. STAR Protocols, 4(3), 102226. https://doi.org/10.1016/j.xpro.2023.102226

Valdivieso, A., Morga, B., Degremont, L., Mege, M., Courtay, G., Dorant, Y., Escoubas, J.-M., Gawra, J., De Lorgeril, J., Mitta, G., Cosseau, C., & Vidal-Dupiol, J. (2025). DNA methylation landscapes before and after Pacific Oyster Mortality Syndrome are different within and between resistant and susceptible Magallana gigas. Science of The Total Environment, 962, 178385. https://doi.org/10.1016/j.scitotenv.2025.178385

Venkataraman, Y. R., Downey-Wall, A. M., Ries, J., Westfield, I., White, S. J., Roberts, S. B., & Lotterhos, K. E. (2020). General DNA methylation patterns and environmentally-induced differential methylation in the Eastern oyster (Crassostrea virginica). Frontiers in Marine Science, 7, 225. https://doi.org/10.3389/fmars.2020.00225

Venkataraman, Y. R., White, S. J., & Roberts, S. B. (2022). Differential DNA methylation in Pacific oyster reproductive tissue in response to ocean acidification. BMC Genomics, 23(1), 556. https://doi.org/10.1186/s12864-022-08781-5

Wang, G., & Mai, R. (2025). Epigenetic regulation of growth and stress response in oysters. International Journal of Aquaculture. https://doi.org/10.5376/ija.2025.15.0019

Wang, X., Li, Q., Lian, J., Li, L., Jin, L., Cai, H., Xu, F., Qi, H., Zhang, L., Wu, F., Meng, J., Que, H., Fang, X., Guo, X., & Zhang, G. (2014). Genome-wide and single-base resolution DNA methylomes of the Pacific oyster

Crassostrea gigas provide insight into the evolution of invertebrate CpG methylation. BMC Genomics, 15(1), 1119. https://doi.org/10.1186/1471-2164-15-1119

Webb, S. C., Fidler, A., & Renault, T. (2007). Primers for PCR-based detection of ostreid herpes virus-1 (OsHV-1): Application in a survey of New Zealand molluscs. Aquaculture, 272(1–4), 126–139. https://doi.org/10.1016/j.aquaculture.2007.07.224

Weichhart, T. (2018). mTOR as regulator of lifespan, aging, and cellular senescence: A mini-review. Gerontology, 64(2), 127–134. https://doi.org/10.1159/000484629

Wickham, H. (2009). Elegant graphics for data analysis. Media, 35(211), 10–1007.

Wright, R. M., Aglyamova, G. V., Meyer, E., & Matz, M. V. (2015). Gene expression associated with white syndromes in a reef building coral, Acropora hyacinthus. BMC Genomics, 16(1), 371. https://doi.org/10.1186/s12864-015-1540-2

Xiao, W., & Loscalzo, J. (2020). Metabolic Responses to Reductive Stress. Antioxidants & Redox Signaling, 32(18), 1330–1347. https://doi.org/10.1089/ars.2019.7803

Yu, M., Chen, J., Bao, Y., & Li, J. (2018). Genomic analysis of NF-κB signaling pathway reveals its complexity in Crassostrea gigas. Fish & Shellfish Immunology, 72, 510–518. https://doi.org/10.1016/j.fsi.2017.11.034

Zhang, G., Fang, X., Guo, X., Li, L., Luo, R., Xu, F., Yang, P., Zhang, L., Wang, X., Qi, H., Xiong, Z., Que, H., Xie, Y., Holland, P. W. H., Paps, J., Zhu, Y., Wu, F., Chen, Y., Wang, J., … Wang, J. (2012). The oyster genome reveals stress adaptation and complexity of shell formation. Nature, 490(7418), 49–54. https://doi.org/10.1038/nature11413

Zhang, L., Li, L., & Zhang, G. (2011). A Crassostrea gigas Toll-like receptor and comparative analysis of TLR pathway in invertebrates. Fish & Shellfish Immunology, 30(2), 653–660. https://doi.org/10.1016/j.fsi.2010.12.023

Zhang, Q., Li, Y., Liao, K., Chen, D., Qiu, Y., Yan, X., & Xu, J. (2023). mTOR plays a conserved role in regulation of nutritional metabolism in bivalve Sinonovacula constricta. Journal of Marine Science and Engineering, 11(5), 1040. https://doi.org/10.3390/jmse11051040

Zhao, Y., Simon, M., Seluanov, A., & Gorbunova, V. (2023). DNA damage and repair in age-related inflammation. Nature Reviews Immunology, 23(2), 75–89. https://doi.org/10.1038/s41577-022-00751-y