In situ chromatin dynamics and HIV-1 nuclear trafficking

Zhen Hou[#1], Yao Shen[#1], Long Chen[1,3], Juan Shen[1], David I. Stuart[1,3], Peijun Zhang[1,2,3]*

[1]Division of Structural Biology, Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK
[2]Diamond Light Source, Harwell Science and Innovation Campus, Didcot, UK
[3]Chinese Academy of Medical Sciences Oxford Institute, University of Oxford, Oxford, UK

[#]Equal contribution
*Correspondence to: peijun.zhang@strubi.ox.ac.uk

Keywords
Chromatin, nucleosome, HIV-1, nuclear trafficking, *in situ*, cryo-CLEM, cryo-ET, cryo-FIB, subtomogram averaging

Abstract
The organization and dynamics of chromatin within intact nuclei remain poorly defined, limiting understanding of how nuclear architecture influences macromolecular transport. HIV-1 provides a striking example of a large complex that must traverse this dynamic landscape to access integration sites within transcriptionally active regions. Here, we combine a functional nuclear import system with correlative cryo-electron tomography to visualize chromatin architecture and HIV-1 cores *in situ* during nuclear penetration. Native nucleosomes resolved at 5.6 Å reveal four structural classes, from compact conformations at the nuclear periphery to open, flexible forms in the interior, defining a spatially heterogeneous and dynamic chromatin landscape. Within this environment, HIV-1 cores decorated with nuclear factors exclude nearby nucleosomes and preferentially associate with open chromatin. Disruption of CPSF6–capsid interactions abolishes these associations, confining viral cores to peripheral chromatin. These findings establish an *in situ* framework linking chromatin structure and dynamics to HIV-1 nuclear trafficking and provide a structural basis for viral navigation within the dynamic nuclear landscape.

Introduction

In human cells, the genome is organized within the nucleus as chromatin, a dynamic nucleoprotein polymer that integrates DNA packaging with gene regulation[1]. Within the confined approximately 10 μm nucleus, roughly two meters of genomic DNA are compacted through hierarchical folding of chromatin. At the molecular level, chromatin is composed of repeating arrays of nucleosomes, each consisting of ~147 base pairs of DNA wrapped around a histone octamer containing two copies each of H2A, H2B, H3, and H4[2, 3, 4]. Linker DNA connecting adjacent nucleosomes, when bound by histone H1, forms chromatosomes that serve as the building blocks of higher-order chromatin structures. Through nucleosome-nucleosome interactions and progressive folding, chromatin establishes domains of distinct accessibility and functionally within the nucleus, balancing genome compaction with the need for molecular accessibility during various DNA metabolic processes[5, 6]. Compact, transcriptionally repressive heterochromatin is typically enriched at the nuclear periphery and around nucleoli, whereas open, transcriptionally active euchromatin predominates in the nuclear interior[7]. Interspersed among these chromatin domains are membraneless nuclear compartments, such as nuclear speckles enriched in splicing and transcriptional regulators[8, 9]. Together, chromatin and associated nuclear bodies create a densely packed yet dynamic landscape that governs genome function and constrains the movement of macromolecular assemblies within the nucleus.

Within this crowded and heterogeneous chromatin environment, HIV-1 must penetrate and navigate the nuclear interior to access the host chromatin for integration[10, 11]. To establish infection in non-dividing cells[12, 13, 14, 15], the HIV-1 core, a conical capsid enclosing the viral genome and enzymes, must cross the nuclear envelope through the nuclear pore complex (NPC)[16, 17, 18, 19]. The HIV-1 capsid recruits host factors to mediate nuclear transport, while simultaneously supporting the completion of reverse transcription of the viral RNA genome for subsequent integration[19, 20, 21, 22, 23, 24, 25].

Among these host factors, cleavage and polyadenylation specificity factor 6 (CPSF6) plays a pivotal role in guiding viral cores through the nuclear environment[26, 27, 28, 29, 30]. CPSF6, a component of the cleavage factor I complex involved in mRNA 3′ end processing, binds directly to the viral capsid, facilitating nuclear entry and steering viral cores away from the heterochromatin-rich nuclear periphery toward gene-dense, transcriptionally active regions within the nuclear interior[26, 27, 28, 29, 30]. Within these subnuclear compartments, reverse transcription is thought to be completed[29, 31, 32], and viral integration is subsequently mediated by the transcriptional coactivator lens epithelium-derived growth factor p75 (LEDGF/p75)[33, 34, 35]. LEDGF binds HIV-1 integrase and tethers the pre-integration complex to nucleosomes enriched

in histone H3K36me3[33, 34, 35], thereby promoting integration within transcriptionally active

chromatin domains known as speckle-associated domains (SPADs)[29, 31, 32] and ensuring efficient

proviral transcription[33, 34, 35]. Despite these insights, how viral cores physically navigate through

the dynamic chromatin environment and achieve integration site targeting remains unclear.

To address this question, we combined a functional HIV-1 nuclear import system coupled with

cryo-focused ion beam (cryo-FIB) milling and correlative cryo-electron tomography (cryo-ET)

to directly visualize chromatin architecture and HIV-1 cores in intact T-cell nucleus during

nuclear trafficking. We resolved the *in situ* structure of nucleosomes at 5.6 Å and identified four

distinct classes of nucleosome conformations, with compact nucleosome conformations

preferentially enriched at the nuclear periphery and open and flexible conformations

predominating within the interior, defining a dynamic and heterogeneous chromatin landscape

that HIV-1 must traverse. Within this nuclear environment, HIV-1 cores decorated with nuclear

factors locally exclude nucleosomes within approximately 35 nm of the capsid surface, creating

nucleosome-depleted zones. Surrounding these regions, nucleosomes in open conformations are

enriched, whereas compact conformations are diminished, suggesting that HIV-1 traffics through

locally accessible chromatin regions. Furthermore, the N74D capsid mutation abolishes the

association of viral cores with nuclear factors, thereby confining them to peripheral, compact

chromatin regions. Together, these findings establish a structural framework connecting

chromatin dynamics to HIV-1 nuclear trafficking, laying a foundation for future studies on

chromatin remodelling, viral integration, and transcription control.

Result

Correlative cryo-ET of HIV-1 nuclear trafficking

While the process of HIV-1 nuclear import through NPCs has been investigated[19, 23, 24, 25], how

viral cores traverse the nuclear interior and interact with host chromatin after nuclear entry

remains poorly understood. To directly visualize this process, we combined our previously

established HIV-1 nuclear import assay with a correlative cryo-ET imaging approach[19, 24].

We performed the HIV-1 nuclear import assay using isolated viral cores with permeabilized

CEM cells in the presence of rabbit reticulocyte lysate and an ATP-regenerating system (RRL-

ATP) (Supplementary Fig. 1a) and imaged the samples by confocal fluorescent microscopy

(FM). Briefly, HIV-1 cores were isolated from virions carrying near full-length genomes (Env-

deficient R9 derivatives)[36] that expressed mNeonGreen-tagged integrase (mNeonGreen-IN).

Sucrose gradient centrifugation followed by Triton X-100 "spin-through" treatment produced a

discrete fluorescent band corresponding to intact cores[37] (Supplementary Fig. 1b, inset), which

was confirmed by cryo-EM imaging (Supplementary Fig. 1b). To allow cytoplasmic access while
preserving nuclear integrity, CEM cells were gently permeabilized with a low concentration of
digitonin (0.018%) and supplemented with RRL-ATP[38, 39] (Supplementary Fig. 1a). Mixing the
isolated HIV-1 cores with these permeabilized cells resulted in robust nuclear import, as
demonstrated by confocal imaging showing abundant mNeonGreen-IN puncta within nuclei
(Supplementary Fig. 1c).
113
We next employed a correlative cryo-FIB and cryo-ET workflow to image imported HIV-1 cores
within intact nuclei[19, 24]. Cryo-FIB milling generated cell lamellae 100–150 nm thick that
contained mNeonGreen-IN signals marking HIV-1 cores and SiR-DNA signals delineating
nuclei. Cryo-fluorescence microscopy (cryo-FM) of the post-milled lamellae confirmed the
presence of mNeonGreen-IN signals within the nucleus (Fig. 1a), guiding cryo-ET data
acquisition from the corresponding regions (Fig. 1b, Supplementary Table 1-2).
120
From 253 tomograms, we identified 55 HIV-1 cores within the nucleus (Fig. 1c and
Supplementary Figure 2) exhibiting both cone- and tube-shaped morphologies (21 cone-shaped
and 34 tube-shaped), consistant with our previous findings[19]. Based on their distance from the
nuclear envelope, we classified the cores into three categories: just-imported (<75 nm; still
associated with the NPC), en-route (75 nm–1.5 µm; detached from the NPC but within 1.5 µm of
the nuclear envelope), and deep-inside (>1.5 µm from the nuclear envelope) (Fig. 1d–i and
Supplementary Video 1-3). Notably, the number of detectable cores decreased with increasing
depth of nuclear penetration (Fig. 1c), consistent with previous fluorescence microscopy studies
of HIV-1–infected cells[29, 30].
130
Nearly all nuclear HIV-1 cores displayed distinct densities on their capsid surfaces, likely
corresponding to nuclear factors such as CPSF6 that mediate HIV-1 nuclear import and
trafficking (Fig. 1d-i, Supplementary Fig. 2). Chromatin fibers were clearly visible beneath the
nuclear envelope (Fig. 1d), where short nucleosome clutches exhibited a zigzag organization
(Fig. 1d–i). Nucleosomes appeared more loosely packed in the nuclear interior (Fig. 1h) than at
the periphery (Fig. 1d, f). Notably, the regions immediately surrounding HIV-1 cores were
markedly depleted of nucleosomes (Fig. 1d–i, Supplementary Fig. 2), suggesting that the
decorated cores may locally displace chromatin during nuclear trafficking. The mechanism
underlying this chromatin exclusion during HIV-1 core transport through the nucleus remains to
be elucidated.
141
***In situ* structure and dynamics of nucleosomes**

It is worth noting that in this HIV-1 nuclear import system, imported viral cores were relatively

rare (a total of 55 cores observed across 253 tomograms, with 9 located deep within the nucleus).

The majority of nucleosomes were far from HIV-1 cores and are therefore expected to remain in

a largely native state. Because the HIV-1 cores were introduced exogenously for 30 minutes, the

overall chromatin landscape was likely only minimally perturbed. Thus, the chromatin structures

described here represent an approximately native configuration in the absence of direct

association with HIV-1 cores.

Using template matching and subtomogram averaging (STA) of individual nucleosomes, we

resolved four distinct nucleosome classes within the nucleus at up to 5.6 Å resolution

(Supplementary Fig. 3-4). The majority of nucleosomes (Class 1, 69%) corresponded to H1-

bound nucleosomes, representing canonical nucleosome core particles with histone H1 bound to

linker DNA (chromatosomes). The 5.6 Å STA map revealed well-defined histone octamers with

clearly resolved secondary structural features and discernible major and minor grooves of the

DNA (Fig. 2a, Supplementary Fig. 3-4). This map closely fits the atomic model of *in vitro*

reconstituted human nucleosomes with H1 bound on-dyad (Fig. 2b)[40]. The C-terminal tail of

histone H2A was clearly resolved, extending outward from the nucleosome core to engage the

linker DNA and contribute to higher-order chromatin organization[41]. Similarly, the N-terminal

tails of histone H3 projected from the DNA groove, consistent with a previous high-resolution

structural study[6]. These findings demonstrate the capability of *in situ* structural determination to

resolve molecular details of small protein–DNA complexes (~250 kDa) within the crowded

nuclear environment.

The second class (Class 2, 8%) revealed two parallelly stacked-nucleosomes, each associated

with a histone H1 molecule (stacked H1-bound nucleosomes) (Fig. 2c, Supplementary Fig. 3-4).

Focused refinement of a single nucleosome within this stack yielded a 9.9 Å map, showing that

the H4 N-terminal tails extended toward adjacent nucleosomes, contacting the acidic patches on

H2A–H2B dimers of neighboring nucleosomes to stabilize inter-nucleosome packing

(Supplementary Fig. 5)[2, 42]. This configuration contrasts with the Class 1 H1-bound nucleosome

(Supplementary Fig. 5b), where the H4 N-terminal tails fold back to form intra-nucleosomal

contacts with the H3 α2 helix, as previously described[40, 43]. Notably, several histone regions at

the inter-nucleosome interface, including H3 α3 and the C-terminal β3 strands of H4 and H2A,

appeared disordered in the focused-refined H1-bound nucleosome within the stacked pair

(Supplementary Fig. 5b). In contrast, in Class 1 nucleosomes, H4-β3 and H2A-β3 form parallel

β-strands that stabilize the histone octamer and maintain nucleosome integrity[40] (Supplementary

Fig. 5b). The absence of these β-strand densities in the stacked configuration suggests that the

H4-β3 tail fails to fold back, instead adopting a flexible conformation that may in turn destabilize
H3-α3. Similar deviations from the folded-back H4-β3 conformation have been observed in
complexes with histone chaperones during histone deposition or eviction[44, 45].
182
Two additional nucleosome classes were associated with open or flexible linker DNA. Class 3
nucleosomes (core nucleosomes, 11%; 9.3 Å resolution) lacked detectable densities for H1 and
linker DNA, whereas Class 4 nucleosomes (open-linker H1-bound nucleosomes, 15%; 12.1 Å
resolution) featured parallel linker DNA arms, with histone H1 engaging asymmetrically with
one of the linkers (Fig. 2c-d, Supplementary Figs. 3-4). Although linker histone H1 is known to
stabilize linker DNA, the weaker density observed for the H1-bound linker in this class likely
reflects partial H1 occupancy and conformational heterogeneity under native nuclear conditions
(Fig. 2d, Supplementary Fig. 4d), reinforcing the role of H1 in regulating chromatin compaction
and dynamic remodeling[6, 46, 47].
192
Chromatin architecture and nucleosome arrangement in the nucleus
To investigate how nucleosome organization contributes to higher-order chromatin architecture,
we mapped nucleosome particles back into their original tomograms based on their determined
positions and orientations. Neighboring nucleosomes were then connected by linker DNA
segments of varying lengths (Fig. 3a), enabling analysis of their spatial relationships and overall
distributions.
199
Consistent with our previous observations[48], chromatin was not organized into regular 30-nm
fibers, but instead exhibited a heterogeneous architecture, in which compact chromatin regions
coexisted with relaxed, irregular zigzag arrangements of nucleosomes (Fig. 3a, Supplementary
Fig. 6, 7a, Supplementary Video 4). We also observed chromatin loops and locally zig-zagged
nucleosome clutches connected by visible linker DNA segments (Supplementary Fig. 6, 7a). To
infer multi-nucleosome connectivity and linker geometry directly from tomograms, we
developed a physics-based probabilistic linker-assignment framework, inspired by Kreysing et
al[49]. Nucleosome coordinates and orientations derived from subtomogram refinement defined
two DNA exit arms per particle. Candidate inter-nucleosome linkages were iteratively evaluated
for geometric compatibility and assigned using a probability-ranked greedy algorithm, ensuring
that each DNA arm participated in only one connection and that no closed cycles formed. The
resulting connectivity graph revealed irregularly packed chromatin trajectories spanning kilobase
scales, from local, short-range zigzag configurations to the longest continuous chains comprising
up to 12 nucleosomes (~2 kb of DNA) (Fig. 3b–c).
214

To examine whether specific nucleosome classes exhibit spatial preferences within the nucleus,

we mapped all four classes of nucleosomes back into their original tomograms (Fig. 3d,

Supplementary Fig. 7b) and quantified their global and local distributions relative to the nuclear

envelope (≤1.5 μm vs. >1.5 μm from the periphery) (Fig. 3e, f). Pairwise nearest-neighbor

distance analysis revealed a dominant inter-nucleosome spacing of ~120–140 Å, accompanied by

a secondary shorter-distance peak at ~55 Å (Fig. 3e). These distances correspond to separations

between i and i + 1 nucleosomes within zigzag chromatin fibers and between i and i + 2

nucleosomes in stacked H1-bound configurations, respectively. H1-bound nucleosomes (Class

1) were significantly enriched near the nuclear periphery ($p = 1.8 \times 10^{-11}$), whereas core

nucleosomes (Class 3) were preferentially localized to the nuclear interior (1.19-fold, $p < 1 \times$

10^{-15}). In contrast, both stacked H1-bound nucleosomes and open-linker H1-bound nucleosomes

(Classes 2 and 4) exhibited relatively uniform distributions across peripheral and interior nuclear

regions (Fig. 3f).

We next investigated whether different nucleosome classes exhibited preferential association

patterns with themselves or with other classes. All four classes showed some degree of self-

association, though to varying extents (Fig. 3g). Core nucleosomes (Class 3) displayed the

strongest self-association with >2-fold enrichment, followed by stacked H1-bound nucleosomes

(Class 2) with a 1.63-fold enrichment and open-linker H1-bound nucleosomes (Class 4) with a

1.43-fold enrichment. In contrast, H1-bound nucleosomes (Class 1) showed minimal preference

among the four classes (Fig. 3g). These findings indicate that nucleosome organization within the

nucleus is non-uniform yet non-random, suggesting the presence of domains enriched in specific

chromatin states that collectively contribute to a dynamic and heterogeneous chromatin

architecture. Further correlative cryo-ET analyses will be instrumental in elucidating which

nuclear factors and histone/DNA modifications underlie the formation and regulation of these

distinct chromatin domains, insights that are not readily captured by conventional bulk

measurements.

HIV-1 cores locally exclude nucleosomes and prefer open chromatin

To investigate the chromatin organization encountered by HIV-1 during nuclear traversal, we

analyzed nucleosome distributions surrounding HIV-1 cores in tomograms. Nucleosome

densities were projected through 130 nm thick lamellae to generate 2D heatmaps (Fig. 4a). These

maps revealed that chromatin is less condensed in deeper nuclear regions compared to areas

adjacent to the nuclear envelope, as expected. Consistent with previous reports[49, 50], the regions

directly beneath the NPC basket were consistently devoid of nucleosomes (Fig. 1e).

Strikingly, we observed a persistent nucleosome-free zone in the immediate vicinity of HIV-1

cores, regardless of their position within the nucleus (Fig. 4a, n = 55 cores). Density profiling

around the cores showed that while nuclear factors occupied regions approximately 7 nm from

the capsid surface, the nucleosome-depleted zone extended up to ~35 nm (Supplementary Fig.

9). Moreover, pairwise nearest-neighbor distance analysis revealed that typical inter-nucleosome

distances were 120–140 Å, whereas capsid–nucleosome separations peaked around 350 Å (Fig.

4b). Together, these observations support a model in which HIV-1 cores locally exclude

nucleosomes, potentially favoring access to open chromatin, although the mechanism of

exclusion remains to be determined.

We next analyzed the spatial distribution of the four nucleosome classes surrounding HIV-1

cores by quantifying their local enrichment, defined as the ratio of each class's proportion within

the region of HIV-1 cores to its global nuclear proportion (Fig. 4c, d). This analysis revealed that

HIV-1 cores strongly prefer open chromatin regions containing core nucleosomes (Class 3) and

open-linker H1-bound nucleosomes (Class 4) (Fig. 4d). In contrast, stacked H1-bound

nucleosomes (Class 2) were the least favored by HIV-1 cores (Fig. 4d). These findings suggest

that HIV-1 cores selectively associate with accessible chromatin environments, potentially

facilitating their trafficking toward nuclear speckle–associated domains (SPADs), where viral

integration preferentially occurs.

CPSF6 facilitates nuclear trafficking of HIV-1 cores

CPSF6 is a key nuclear host factor that dictates integration site selection[28, 51, 52]. It is thought to

direct HIV-1 cores away from heterochromatin-rich regions toward the nuclear interior, thereby

promoting integration into gene-dense, transcriptionally active chromatin within SPADs[29, 31, 53].

CPSF6 contains a single phenylalanine–glycine (FG) motif, and mutation of this motif abolishes

its association with the HIV-1 capsid[53], resulting in a loss of CPSF6-dependent targeting of

integration into euchromatin and transcriptionally active genes, as demonstrated by integration-

site mapping and fluorescence microscopy studies[27, 28]. Conversely, HIV-1 capsids carrying the

N74D mutation fail to bind CPSF6[53] and are defective in CPSF6-dependent trafficking to nuclear

speckles and integration into SPADs[31, 54].

We analyzed the nuclear trafficking of HIV-1 cores carrying the N74D mutation and compared

them with WT cores. Although N74D cores exhibited similar morphologies and nuclear envelope

binding as WT cores, they failed to accumulate efficiently within nuclei (Supplementary Fig. 1b,

c). Instead, mNeonGreen-IN puncta were predominantly localized near the nuclear periphery and

were rarely observed in the nuclear interior. Correlative cryo-ET imaging further revealed that

most N74D nuclear cores remained tethered to NPCs, with only a single detached core detected
near the nuclear envelope and none penetrating deeper into the nucleus (Fig. 5a-c,
Supplementary Fig. 10, Supplementary Video 5). Notably, these N74D cores lacked the
prominent surface densities observed coating imported WT cores (Fig. 5a-b). Density profiling
confirmed the presence of distinct density layer at a distance of ~7 nm from the capsid layer only
around imported WT cores, but absent in both N74D cores within the nucleus and WT cores
outside the nucleus (Fig. 5d), consistent with our previous observations[19].
294
To further characterize capsid-nuclear factor interactions, we performed STA of CA lattices from
both WT and imported N74D cores which had not been previously resolved[19]. This analysis
yielded three CA hexamer maps from WT cores outside the nucleus, WT cores inside the
nucleus, and N74D cores inside the nucleus, resolved at 10.1 Å, 9.6 Å, and 12.8 Å, respectively
(Fig. 5e, Supplementary Fig. 11, Supplementary Table 3). Only imported WT cores, but not
unimported WT or N74D cores, exhibited additional densities ~65 Å above the capsid surface
(Fig. 5e-f). The **location** of these densities is consistent with structural studies showing the prion-
like low-complexity regions (LCRs) of CPSF6 bound to CA tubes[55], providing direct *in situ*
evidence that CPSF6 interacts with the viral capsid, likely guiding HIV-1 core trafficking from
peripheral heterochromatin toward the transcriptionally active nuclear interior.
305
Discussion
For decades, determining the *in situ* structures of chromatin and nucleosomes within intact nuclei
has remained a major challenge. Recent advances in cryo-FIB milling and cryo-ET have made it
possible to visualize native chromatin fibers and nucleosomes at moderate resolutions (> 10 Å)[48,
56, 57, 58, 59]. In this study, we resolved the native nucleosome structure at 5.6 Å resolution,
providing an unprecedented level of structural detail for a ~250 kDa complex within the crowded
environment of the cell nucleus. Our analyses further revealed multiple nucleosome classes
corresponding to compact and open chromatin conformations, and distinct histone tail
organizations critical for epigenetic regulation and other DNA metabolic processes. *In situ*
visualization of off-dyad H1 binding uncovered asymmetric linker DNA engagement, providing
structural insight into the flexibility and dynamic behavior of chromatin. These findings mark a
step change in the direct visualization of chromatin architecture and heterogeneity within intact
nuclei.
319
HIV-1 post-nuclear events have been extensively investigated through cell biology approaches[28,
32, 51, 60], fluorescence imaging[61, 62, 63, 64, 65], and more recently, cryo-ET[19, 23, 24, 25]. These studies
established that HIV-1 enters the nucleus with a largely intact capsid, which interacts with

CPSF6 on the nuclear side and is subsequently directed toward SPADs, where integration

preferentially occurs within transcriptionally active chromatin. However, how viral cores

navigate the crowded nuclear environment and complex chromatin landscape to reach these

integration sites has remained unclear. By combining a functional nuclear import system with

high-resolution *in situ* structural analysis, we directly imaged HIV-1 cores at varying depths of

nuclear penetration and characterized their interactions with chromatin inside intact nuclei.

A key finding is that HIV-1 cores exhibit distinct preferences for specific local chromatin

environments. Viral cores excluded nucleosomes from within ~35 nm of their capsid surface,

creating local nucleosome-depleted zones. Similar exclusion zones were observed beneath NPCs,

both in this study and by others[49, 50], and are thought to facilitate macromolecular transport

across the nuclear envelope. Within the surrounding chromatin, open nucleosome states (Classes

3 and 4) were enriched, whereas compact nucleosome states (Classes 1 and 2) were

underrepresented. This pattern suggests that nuclear factor–decorated HIV-1 cores preferentially

traverse open, accessible chromatin regions, potentially displacing nucleosomes as they progress

toward transcriptionally active integration sites.

We further show that CPSF6 decorates the HIV-1 core upon nuclear entry, enabling the viral core

to traverse compact peripheral chromatin and access the more dispersed chromatin of the nuclear

interior. Disruption of the CA–CPSF6 interaction (as in the N74D mutant) confined viral cores to

the nuclear periphery, consistent with previous observations that loss of CPSF6 association

redirects HIV-1 integration toward lamina-associated heterochromatin[28, 30, 31, 65, 66, 67, 68]. The C-

terminal arginine/serine-rich domain of CPSF6 is intrinsically disordered and capable of

mediating phase separation and nuclear speckles association[60, 69]. Whether this condensation

property contributes to the nuclear trafficking of HIV-1 cores toward nuclear speckle remains to

be determined. Future studies should also explore whether additional nuclear factors cooperate

with CPSF6 to regulate HIV-1 trafficking and integration site selection.

Based on these findings, we propose a model in which CPSF6 binding and chromatin

accessibility act in concert to guide HIV-1 cores toward transcriptionally active regions (Fig. 6).

Upon nuclear entry, HIV-1 cores rapidly recruit CPSF6 and potentially other nuclear cofactors,

forming a proteinaceous coat on the viral capsid that facilitates passage through compact

peripheral chromatin into the more open and accessible chromatin of the nuclear interior. During

this process, HIV-1 cores locally exclude nucleosomes through a yet-unknown mechanism.

Collectively, these findings establish a structural framework directly linking HIV-1 nuclear

trafficking to chromatin organization, providing a foundation for future studies of integration site

selection and viral latency. More broadly, this work highlights the potential of *in situ* structural

biology to illuminate the 3D organization of the genome, as well as the mechanisms of

replication, repair, and transcriptional regulation across physiological and disease contexts.

Methods

Mammalian cell culture

All cell cultures were maintained at 37 °C in a humidified incubator with 5% CO_2. Human

embryonic kidney (HEK) 293T Lenti-X cells (632180; Takara/Clontech) were obtained directly

from the vendor, who performed authentication by short tandem repeat (STR) profiling and

confirmed the absence of mycoplasma contamination. HEK293T Lenti-X cells were grown in

Dulbecco's modified Eagle medium (DMEM; Gibco) supplemented with 10% fetal bovine

serum (FBS), 2 mM L-glutamine (Gibco), and 1% MEM non-essential amino acids (Gibco).

$CD4^+$ T lymphocyte CEM cells (NIH HIV Reagent Program, ARP-117) were maintained in

RPMI-1640 medium (Sigma-Aldrich) containing 10% FBS and 2 mM L-glutamine (Gibco).

These cells were not independently authenticated following acquisition.

Production of mNeonGreen-IN-labelled HIV-1 cores containing near-full-length HIV-1

RNA

mNeonGreen-IN–labelled HIV-1 cores carrying near–full-length viral RNA were generated as

previously described[19]. Briefly, HEK293T Lenti-X cells (~80% confluence, four T175 flasks)

were transfected using polyethyleneimine (PEI; branched, MW ~25,000; Sigma-Aldrich). Each

transfection contained 24 µg of Env-defective HIV-1 plasmid (R9ΔE-CA-WT or R9ΔE-CA-

N74D) and 6 µg of Vpr–mNeonGreen–IN plasmid (a gift from Prof. Zandrea Ambrose,

University of Pittsburgh) mixed with 120 µg PEI in Opti-MEM. After incubation at room

temperature for 20 min, the mixture was added to the cells. 16 h post-transfection, the medium

was replaced with fresh DMEM supplemented with 10% FBS, 1% nonessential amino acids, and

1% L-glutamine. Supernatants containing HIV-1 particles were collected after 24 h.

Virus-containing supernatants (120 ml) were filtered through a 0.45 µm filter (Sarstedt) and

layered onto 5 ml of 20% sucrose in 1× STE buffer (10 mM Tris-HCl pH 7.4, 100 mM NaCl, 1

mM EDTA) in four 38.5 ml ultracentrifuge tubes (Beckman Coulter). Virions were pelleted using

a SW32Ti rotor at 85,500 × g for 3 h at 4 °C (Optima XPN-90, Beckman Coulter). Each pellet

was resuspended in 400 µl 1× STE supplemented with 1 mM IP6.

For core purification, a continuous sucrose gradient (30–85%) was prepared in 13.2 ml

ultracentrifuge tubes (Beckman Coulter) in 1× STE with 1 mM IP6. The gradient was overlaid

with 250 µl of 15% sucrose containing 1% Triton X-100, followed by 7.5% sucrose to create a

protective barrier. Approximately 800 µl of concentrated HIV-1 particles was layered on top, and

gradients were centrifuged at 182,600 × g for 16 h at 4 °C using a SW41Ti rotor.

After centrifugation, the fluorescent mNeonGreen band corresponding to HIV-1 cores was
visualized under blue light and collected by side puncture with a 25 G needle (BD Microlance 3).
Core fractions were quantified by p24 ELISA and either snap-frozen in single-use aliquots at
−80 °C or dialyzed overnight at 4 °C against 500 ml of 1× SHE buffer (10 mM HEPES-NaOH
pH 7.4, 100 mM NaCl, 1 mM EDTA) containing 0.8 mM IP6 using a 7 kDa cut-off Slide-A-
Lyzer Dialysis Cassette (Thermo Fisher Scientific) before mixing with permeabilized T cells.
405

T cell membrane permeabilization
Exponentially growing CEM T cells (~1 × 10^6 cells/ml) were collected by centrifugation at 500 ×
g for 5 min at 4 °C, washed twice with 10 ml of ice-cold PBS, and pelleted under the same
conditions. The cell pellet was resuspended in lysis buffer (20 mM HEPES pH 7.5, 25 mM KCl,
5 mM MgCl$_2$, 1 mM DTT, and 1× protease inhibitor) containing 0.018% digitonin. Cells were
gently rotated at room temperature for 10 min to permeabilize the plasma membrane, following
previously established procedures[19]. After incubation, the cells were briefly centrifuged at 200 ×
g for 1 min, and the supernatant was discarded. The pellet was resuspended in fresh lysis buffer
without digitonin to remove residual detergent. To verify nuclear integrity, digitonin-treated
CEM cells were incubated with 0.2 mg/ml FITC–dextran (500 kDa; Sigma-Aldrich, 46947) in
lysis buffer for 15 min at room temperature. Images were acquired using a Leica TCS SP8
confocal microscope.
418

Nuclear import of mNeonGreen-IN-labelled HIV-1 cores in permeabilized T cells
Nuclear import of mNeonGreen-IN-labelled HIV-1 cores in permeabilized T cells was carried
out as previously described[19]. Nuclei of digitonin-permeabilized CEM cells were stained on ice
with 1 μM SiR-DNA (Spirochrome) for 30 min. Approximately 4 × 10^5 permeabilized cells were
then incubated with HIV-1 cores (WT or N74D) at a final CA concentration of 20 μg/ ml in a
40 μl reaction mixture containing 0.2 mM IP6, 20 mM HEPES (pH 7.5), 18.8 mM KCl, 20 mM
NaCl, 0.25 mM EDTA, 3.8 mM MgCl$_2$, 1 mM DTT, and 1× protease inhibitor cocktail. After a
30 min incubation on ice, 10 μl of supplement mix (rabbit reticulocyte lysate + ATP + GTP +
creatine phosphate + creatine kinase) was added, resulting in final concentrations of 16% (v/v)
rabbit reticulocyte lysate (RRL; Promega, L4151), 0.6 mM ATP, 0.06 mM GTP, 5 mM creatine
phosphate, and 10 U μl^{-1} creatine kinase. The reaction was incubated at 37 °C for 30 min to allow
nuclear import. For confocal microscopy, samples were transferred to μ-Slide 18-well glass-
bottom chambers (Ibidi) and imaged using a Leica TCS SP8 confocal microscope. For cryo-FIB
milling and cryo-ET, samples were fixed with 5 mM ethylene glycol bis(succinimidyl succinate)
(EGS) for 30 min at room temperature, quenched with 50 mM Tris-HCl (pH 7.5), and kept on ice
until grid preparation.

Confocal microscopy and fluorescence image processing

All fluorescence imaging was performed using a Leica TCS SP8 confocal microscope equipped with a HC PL Apo 63×/1.2 NA MotCORR water-immersion CS2 objective and a HyD GaAsP detector, operated through LAS X software (Leica Microsystems). Fluorescent signals from GFP and SiR-DNA were acquired using 488 nm and 633 nm excitation laser lines, respectively, with optimized spectral filters to minimize channel overlap. Z-stacks were acquired with a step size of 0.3–0.5 µm using a 1 a.u. pinhole and an image resolution of 1,024 × 1,024 pixels. Image stacks were processed and analysed using Leica Application Suite X (LAS X) and Fiji (ImageJ)[72].

Plunge-freezing vitrification

Sample vitrification was performed using a Leica EM GP2 automated plunge freezer under controlled conditions (20 °C, 95% relative humidity). The mixture of HIV-1 cores and permeabilized CEM cells was incubated with 10% glycerol for 2 min prior to freezing. A 3–6 µL aliquot of the mixture (containing approximately 4,000 cells per µL) was applied to the carbon side of glow-discharged Quantifoil R2/1 Cu 300 grids, while 1 µL of PBS was placed on the reverse side. Grids were back-blotted for 6 s using Whatman filter paper and then rapidly plunge-frozen in liquid ethane. For HIV-1 cores, 3.5 µL of WT or N74D cores were applied to the carbon side of glow-discharged lacey carbon film grids (Cu 300 mesh, Electron Microscopy Sciences), back-blotted for 3 s, and plunge-frozen in liquid ethane under identical environmental conditions. Cryo-EM data were collected on a Glacios transmission electron microscope (Thermo Fisher Scientific) equipped with a Falcon III direct electron detector at a nominal magnification of 73,000×.

Correlative cryo-FIB milling

Vitrified mixtures of permeabilized CEM cells supplemented with RRL-ATP and HIV-1 cores were thinned by cryo-FIB milling to produce lamellae, guided by cryo-CLEM using two distinct systems. Two grids of WT cores were loaded via an Autoloader onto a dual-beam FIB/SEM microscope (Arctis, Thermo Fisher Scientific), equipped with a cryogenic stage cooled to −191 °C, a wide-field integrated fluorescence microscope (iFLM; 100×, NA 0.75), and a plasma multi-ion source (argon, xenon, and oxygen). Argon was used as the FIB source in this study.

Prior to milling, a platinum layer was sputter-coated for 30 s, followed by deposition of an organometallic platinum layer via the GIS system (Thermo Fisher Scientific) for 50 s. Three-dimensional correlative milling was performed using the embedded protocol in WebUI v1.3 (Thermo Fisher Scientific) with a milling angle of 10°. For each position, a 15 µm z-stack of

fluorescence (GFP) and reflection images was acquired at 500 nm step size after rough milling
under default parameters. Target fluorescence coordinates were registered using discernible ice
chunks as fiducial markers in both SEM and FIB images to guide lamella placement. Lamellae
were produced stepwise: (i) opening at 2 nA, (ii) rough milling at 0.74 nA and 0.2 nA, and (iii)
polishing at 60 pA, yielding a final lamella thickness of 140 nm. To enhance fluorescence
detection on polished lamellae, the step size was reduced to 100 nm, generating a 15 µm stack
(101 images) in both GFP and far-red channels.
478
Four grids of WT cores and one grid of N74D cores were milled on a dual-beam FIB/SEM
microscope (Aquilos 2, Thermo Fisher Scientific) equipped with a cryo-transfer system and a
rotatable cryo-stage cooled to −191 °C. The instrument was modified to incorporate a
fluorescence module (METEOR, Delmic Cryo BV; 50×, NA 0.8). Grids were mounted on a
METEOR shuttle with a pre-tilt of 26° and coated with an organometallic platinum layer for 30 s
using the GIS system. Milling was performed at 10° using AutoTEM 5 (Thermo Fisher
Scientific). Cells located near grid centers were selected for lamella preparation. Before milling,
fluorescence stacks were acquired for each site in GFP and far-red channels at 200 nm steps (± 6
µm), processed in ImageJ to improve signal-to-noise ratio (SNR), and correlated with FIB/SEM
images using the 3D Correlation Toolbox[73], with ice features as fiducial markers. Sequential
milling was performed from 0.5 nA (rough) to 0.3 nA (medium), 0.1 nA (fine), 60 pA (first
polishing), and 30 pA (final polishing), yielding lamellae ~120 nm thick. Final fluorescence
stacks were collected (GFP and far-red) at 100 nm steps (± 2 µm) using a 400 mW laser and 300
ms exposure. In total, 15 lamellae (Arctis) and 21 lamellae (Aquilos 2) containing detectable
fluorescence were generated for WT cores, and 9 lamellae were produced in Aquilos 2 for N74D
cores.
495
Cryo-electron tomography data collection
Lamellae were transferred to two Titan Krios G3 (Thermo Fisher Scientific) microscopes
operating at 300 kV, each equipped with a Falcon 4i detector and Selectris X energy filter.
Objective apertures (100 µm) were inserted. Overview images were collected at 8100×
magnification and stitched to create lamella maps, which were aligned with corresponding
fluorescence images in ImageJ. Tilt series were then collected at the correlated sites and
neighboring regions (without overlap) using Tomography 5 (Thermo Fisher Scientific) at 64k×
magnification.
504
For WT cores, 253 tilt series were recorded at 1.94 Å/pixel, using lamella pre-tilts of ± 12°, and a
dose-symmetric scheme from −42° to +66° in 2° increments. Each series comprised 55 tilts with

5 movie frames per tilt at 2.5 e$^-$/Å²/tilt (total dose 137.5 e$^-$/Å²) and −2 to −5 μm defocus. For

N74D cores, 45 tilt series were collected at 1.903 Å/pixel using identical imaging parameters.

Alignment of tilt series

Movie frames were aligned with MotionCor2[74]. Tilt-series alignment and tomogram

reconstruction were performed in IMOD v4.11[75] by patch tracking (200×200 pixel patches, 0.45

overlap). Alignments were inspected manually and poor frames removed. Reconstructed

tomograms were binned 6× (11.64 Å/pixel for WT, 11.418 Å/pixel for N74D) and SIRT-like

filtered (8 iterations) for visualization.

Density profiling

To measure the intensity of surround densities of HIV-1 cores, grey value measurements were

taken to assess the change in density as a function of distance from the surface of HIV-1 cores.

For each core, lines were derived from the capsid surface including partial density of the CA (~ 3

nm). Black density was assigned a value of 0, and white was assigned a value of 1, with higher

densities corresponding to lower values. In total, 233 lines were derived from 12 imported WT

cores, 155 lines were derived from 12 outside WT cores, and 101 lines were derived from 10

imported N74D cores. For chromatin, 227 lines were derived and profiled in 12 tomograms.

Template matching

To localize nucleosomes, template matching was performed using emClarity version 1.5.0.2[76]

with non-CTF-corrected tomograms binned at 6×. The procedure utilized a template derived

from EMD-3947[77], which was low-pass filtered to 40 Å (Supplementary Fig. 3a). An exhaustive

search was applied by giving a peak threshold ranging from 1,500 to 3,500 based on the size of

nuclear region in the tomogram. Matched coordinates were then imported into Chimera using the

plug-in Place Object[78, 79] for the initial cleaning based on the cross-correlation coefficient (ccc)

value, the cut-off was set to 0.35. Particles residing outside nuclei or with a ccc value lower than

0.35 were treated as false positives and removed (Supplementary Fig. 3a).

To localize individual CA hexamers, template matching was performed in emClarity version

1.5.0.2[76] with non-CTF-corrected tomograms binned at 6× using a low-pass filtered (30 Å)

template derived from EMD-12452[80]. An exhaustive search was applied by giving a peak

threshold of 2,000 in the small cropped-out tomogram region (150 nm³) containing the HIV-1

core. HIV-1 CA hexamer peaks were selected with the MagpiEM tool (available at

https://github.com/fnight128/MagpiEM), and particle selection was identified based on the

geometric constraints of the capsid lattice. Any hexamers that did not conform to the expected

hexagonal lattice geometry were automatically excluded, followed by manual inspection to
ensure the selection precision (Supplementary Fig. 11a).
545
546 NPCs and 80S ribosomes were located adopting the same procedure, using 1/8 of the original
547 density map EMD-11967[23] for NPCs, and EMD-1780[81] for ribosomes, respectively.
548
Subtomogram averaging
The 3D alignment and averaging of nucleosomes were conducted in RELION version 4.0[82] with
an initial particle number of 551,565. Prior to the iterative refinement, iterative classification was
performed to further clean the particles until no prominent bad class was generated, resulting in
230,106 particles for the final classification (Supplementary Fig. 3a). In the final classification,
four major classes were generated: class 1 (H1-bound nucleosome) with 158,436 particles, class
2 (stacked H1-bound nucleosomes) with 18,563 particles, class 3 (core nucleosome) with 25,153
particles, and class 4 (open-linker H1-bound nucleosome) with 27,954 particles. Iterative
refinement was then carried out for each class from bin6 to bin1. The resolution of the
reconstructed structure was calculated using a gold-standard Fourier shell correlation (FSC) cut-
off of 0.143. For nucleosomes, the final resolution was determined at 5.6 Å, 20.0 Å, 9.9 Å, 9.3
Å, and 12.1 Å for class 1, class 2, mono H1-bound nucleosome in class 2, class 3, and class 4,
respectively (Supplementary Fig. 3b). For HIV-1 CA hexamers, a similar procedure was
employed (Supplementary Fig. 11a), and the resolution was determined at 10.1 Å, 9.6 Å, and
12.8 Å for outside WT CA (6,404 particles), imported WT CA (4,445 particles), and imported
N74D CA (988 particles) (Supplementary Fig. 11b). Structural fitting was conducted in
ChimeraX[83].
566
Segmentation
To enhance the segmentation, reconstructed bin6 tomograms were corrected for the missing
wedge and denoised using IsoNet version 0.2[84], applying 35 iterations with sequential noise cut-
off levels of 0.05, 0.1, 0.15, 0.2, and 0.25 at iterations 10, 15, 20, 25, and 30, respectively.
Membranes and nuclear envelopes in all tomograms were initially segmented using MemBrain-
seg[85], then imported into ChimeraX for manual cleaning and polishing. Nucleosomes mapped
back to the tomograms with segmented membranes using ChimeraX and ArtiaX[86], based on their
positions and orientations after final refinement. For better visualization, nucleosomes depicted
in the segmented volume were low-pass filtered structures from this study. HIV-1 cores were
mapped back based on the refined coordinates and orientations of CA hexamers from
subtomogram averaging, with missing CA hexamers manually placed back using the *in situ* CA
hexamer structures resolved in this study. CA pentamers were only placed back on cores with

most CA hexamers matched and the five surrounding hexamers identified and then modelled

using the low-pass filtered EMD-13422[80]. NPCs and 80S ribosomes were mapped back based on

their coordinates after the template matching using the low-pass filtered model generated from

EMD-11967 and EMD-1780, respectively. Movies were made using ChimeraX.

Definition of linker arms and geometric parameters

Each nucleosome was represented by two exit points ("linker arms") and corresponding tangent

vectors indicating the directions of DNA emerging from the dyad axis. The spatial positions and

orientations of these linker arms were derived from subtomogram-averaged nucleosome densities

and guided by the canonical chromatosome structure (PDB 7K5Y)[6].

Candidate pair identification

For each nucleosome, potential partners were identified based on the spatial proximity of linker-

arm exit points. Neighbours located within 30 nm of either arm were considered candidate

partners. For each valid pair (i, j), the four possible arm-to-arm combinations (arm$_1$–arm$_1$, arm$_1$–

arm$_2$, arm$_2$–arm$_1$, arm$_2$–arm$_2$) were evaluated.

DNA bending-energy model and connection probability

For each possible connection, the linker bending angle (θ) and arc length (L) were derived from

the positions and orientations of the two arms, where θ is defined as the angle between the unit

tangent vectors of the two arms, t_1 and t_2:

$$\theta = \cos^{-1}(-\mathbf{t}_1 \cdot \mathbf{t}_2),$$

and L was computed from the distance between the two DNA exit points ($\|r_2 - r_1\|$) and the

bending angle using circular-arc geometry:

$$L = \frac{\theta}{2\sin(\theta/2)} \|r_2 - r_1\|.$$

Where r_1 and r_2 are the coordinates of the two arms.

The DNA linker bending energy was then estimated using the worm-like chain (WLC) model,

which describes the entropic elasticity of double-stranded DNA[87, 88, 89]:

$$U_{\text{bend}}(\theta, L) = \frac{2l_p}{L}\left(\frac{\theta}{2}\right)^2 k_B T,$$

Where L is the linker length, $l_p = 50$ nm, the persistence length of B-form DNA under

physiological ionic conditions [90], and $k_B T$ is the thermal energy. The total connection probability

was defined as:

$$P(L, \theta) \propto \exp\left(-\frac{L}{L_0}\right) \exp\left[-\frac{U_{\text{bend}}(\theta, L)}{k_B T}\right]$$

with $L_0 = 25$ nm, the characteristic decay length. This formulation penalizes both excessively

long and sharply bent linkers, thereby favouring geometrically plausible DNA trajectories.

Probabilistic connection assignment and graph construction

A three-dimensional probability matrix P_{ijk} was computed, representing the likelihood of each

possible arm-to-arm connection between nucleosome pairs (i, j), where k indexes the four

possible arm combinations. Candidate connections were incorporated iteratively in descending

order of probability.

At each iteration, the highest-probability entry was examined; if $P_{ijk} > 0.1$, the corresponding

connection was accepted. A union–find (disjoint-set) data structure was used to track connected

components, ensuring that each arm was used only once and that no cyclic connections occurred.

After each accepted connection, all entries involving the used arms were excluded from further

consideration. The process continued until no remaining probabilities exceeded the threshold.

The resulting adjacency graph $G(V, E)$ contains nucleosomes as vertices V and their most

probable linker connections as edges E. Each connected component of G represents a continuous

chromatin fragment of variable length and serves as the basis for further statistical and structural

analyses.

Statistical analysis

To assess the significance of distribution differences across nucleosomes, two-sided Chi-square

test and Fisher's exact test were applied. The calculation and plots were carried out using Prism

v10.

Data availability

All data needed to evaluate the conclusions in the paper are present in the paper and/or the

Supplementary Information. The subtomogram-averaged maps for nucleosomes subunits have

been deposited in the public database EMDB under the following accession codes: EMD-55446

(H1-bound nucleosome), EMD-55447 (stacked H1-bound nucleosomes), EMD-55448 (H1-

bound nucleosome in stacking nucleosomes), EMD-55449 (core nucleosome), and EMD-55450

(open-linker H1-bound nucleosome). The subtomogram-averaged maps for the CA hexamers

from both unimported and imported WT cores and from imported N74D cores have been

deposited in the public database EMDB under the following accession codes: EMD-55441,

EMD-55443, and EMD-55445, respectively. Source data are provided with this paper.

Code availability

The scripts used in this study and relevant codes are deposited in GitHub

(https://github.com/fnight128/MagpiEM).

Acknowledgements

We thank L. Carrique, H. Duyvesteyn, D. Owen, J. Bancroft, E. Drydale and J. Gilchrist for their

support in data collection. We thank J. Shi and C. Aiken (Vanderbilt University Medical Center)

for kindly providing plasmids to produce HIV-1 cores. We thank F. Nightingale for the MagpiEM

software. Y.S. is further supported by a CIHR fellowship from the Canadian Institutes of Health

Research (194032), and an EMBO fellowship (ALTF 96-2024). We acknowledge the Oxford

Particle Imaging Centre (OPIC) for access to cryo-FIB/SEM instruments (Arctis and Aquilos 2)

and the cryo-EM instrument (Krios). We acknowledge the Oxford Cellular Imaging Core Facility

(CICF) for access to fluorescence microscopes and imaging analysis software. We acknowledge

Diamond Light Source for access to and support of the cryo-EM facilities at the UK national

electron Bio-Imaging Centre (eBIC), proposal NT29812. Computation was performed at the

Diamond Light Source and Oxford Biomedical Research Computing (BMRC) facility supported

by the Wellcome Trust Core Award grant number 203141/Z/16/Z with additional support from

the NIHR Oxford BRC. This work was supported by US National Institutes of Health grants

U54AI170791 (P.Z.), R21AI184080 (P.Z.); the UK Wellcome Investigator Award 206422/Z/17/Z

(P.Z.); the UK Wellcome Discovery Award 311427/Z/24/Z (P.Z.); the UK Biotechnology and

Biological Sciences Research Council grant BB/S003339/1 (P.Z.); ERC AdG grant 101021133

(P.Z.); and the Chinese Academy of Medical Sciences (CAMS) Innovation Fund for Medical

Science (CIFMS) (grant no. 2018-I2M-2-002; P.Z., D.I.S.). The funders had no role in study

design, data collection and analysis, decision to publish, or preparation of the paper.

Contributions

P.Z. conceived the research. Z.H., Y.S. and P.Z. designed the experiments. Y.S. prepared samples.

Y.S. conducted the fluorescence microscopy. J.S. helped with HIV-1 core preparation. Z.H.

performed the cryo-CLEM and cryo-FIB. Z.H. and Y.S. collected the cryo-ET data and

performed reconstruction of tomograms. Y.S. collected the micrographs of input cores. Z.H.

performed STA and the segmentation. L.C. performed the linker DNA analysis under the

supervision of D. I. S.. Z.H. performed the statistical analyses. Z.H. and Y.S. prepared figures.

Z.H., Y.S., and P.Z. wrote the paper with help from other co-authors.

Reference

1. Delvaux de Fenffe CM, Govers J, Mattiroli F. Always on the Move: Overview on Chromatin Dynamics within Nuclear Processes. *Biochemistry* **64**, 2138-2153 (2025).

2. Luger K, Mader AW, Richmond RK, Sargent DF, Richmond TJ. Crystal structure of the nucleosome core particle at 2.8 A resolution. *Nature* **389**, 251-260 (1997).

3. Kornberg RD. Chromatin structure: a repeating unit of histones and DNA. *Science* **184**, 868-871 (1974).

4. Oudet P, Gross-Bellard M, Chambon P. Electron microscopic and biochemical evidence that chromatin structure is a repeating unit. *Cell* **4**, 281-300 (1975).

5. Woodcock CL, Frado LL, Rattner JB. The higher-order structure of chromatin: evidence for a helical ribbon arrangement. *J Cell Biol* **99**, 42-52 (1984).

6. Zhou BR*, et al.* Distinct Structures and Dynamics of Chromatosomes with Different Human Linker Histone Isoforms. *Mol Cell* **81**, 166-182 e166 (2021).

7. Bickmore WA, van Steensel B. Genome architecture: domain organization of interphase chromosomes. *Cell* **152**, 1270-1284 (2013).

8. Mao YS, Zhang B, Spector DL. Biogenesis and function of nuclear bodies. *Trends Genet* **27**, 295-306 (2011).

9. Sawyer IA, Sturgill D, Dundr M. Membraneless nuclear organelles and the search for phases within phases. *Wiley Interdiscip Rev RNA* **10**, e1514 (2019).

10. Jang S, Engelman AN. Capsid-host interactions for HIV-1 ingress. *Microbiol Mol Biol Rev* **87**, e0004822 (2023).

11. Bedwell GJ, Engelman AN. Factors that mold the nuclear landscape of HIV-1 integration. *Nucleic Acids Res* **49**, 621-635 (2021).

12. Katz RA, Greger JG, Boimel P, Skalka AM. Human immunodeficiency virus type 1 DNA nuclear import and integration are mitosis independent in cycling cells. *J Virol* **77**, 13412-13417 (2003).

13. Mannioui A*, et al.* Cell cycle regulation of human immunodeficiency virus type 1 integration in T cells: antagonistic effects of nuclear envelope breakdown and chromatin condensation. *Virology* **329**, 77-88 (2004).

14. Lusic M, Siliciano RF. Nuclear landscape of HIV-1 infection and integration. *Nat Rev Microbiol* **15**, 69-82 (2017).

729   15.   Fassati A. HIV infection of non-dividing cells: a divisive problem. *Retrovirology* **3**,
730   74 (2006).
731
732   16.   Schuller AP*, et al.* The cellular environment shapes the nuclear pore complex
733   architecture. *Nature* **598**, 667-671 (2021).
734
735   17.   Zimmerli CE*, et al.* Nuclear pores dilate and constrict in cellulo. *Science* **374**,
736   eabd9776 (2021).
737
738   18.   Singh D*, et al.* The molecular architecture of the nuclear basket. *Cell* **187**, 5267-
739   5281 e5213 (2024).
740
741   19.   Hou Z*, et al.* HIV-1 nuclear import is selective and depends on both capsid
742   elasticity and nuclear pore adaptability. *Nat Microbiol* **10**, 1868-1885 (2025).
743
744   20.   Campbell EM, Hope TJ. HIV-1 capsid: the multifaceted key player in HIV-1
745   infection. *Nat Rev Microbiol* **13**, 471-483 (2015).
746
747   21.   Shen Q, Wu C, Freniere C, Tripler TN, Xiong Y. Nuclear Import of HIV-1. *Viruses*
748   **13**, (2021).
749
750   22.   Wilbourne M, Zhang P. Visualizing HIV-1 Capsid and Its Interactions with
751   Antivirals and Host Factors. *Viruses* **13**, (2021).
752
753   23.   Zila V*, et al.* Cone-shaped HIV-1 capsids are transported through intact nuclear
754   pores. *Cell* **184**, 1032-1046 e1018 (2021).
755
756   24.   Hou Z*, et al.* Direct visualization of HIV-1 core nuclear import and its interplay
757   with the nuclear pore. *EMBO Rep*, (2025).
758
759   25.   Kreysing JP*, et al.* Passage of the HIV capsid cracks the nuclear pore. *Cell* **188**,
760   930-943 e921 (2025).
761
762   26.   Bialas K, Diaz-Griffero F. HIV-1-induced translocation of CPSF6 to biomolecular
763   condensates. *Trends Microbiol* **32**, 781-790 (2024).
764
765   27.   Sowd GA*, et al.* A critical role for alternative polyadenylation factor CPSF6 in
766   targeting HIV-1 integration to transcriptionally active chromatin. *Proc Natl Acad
767   Sci U S A* **113**, E1054-1063 (2016).
768
769   28.   Achuthan V*, et al.* Capsid-CPSF6 Interaction Licenses Nuclear HIV-1 Trafficking
770   to Sites of Viral DNA Integration. *Cell Host Microbe* **24**, 392-404 e398 (2018).
771
772   29.   Francis AC*, et al.* HIV-1 replication complexes accumulate in nuclear speckles
773   and integrate into speckle-associated genomic domains. *Nat Commun* **11**, 3505
774   (2020).

776   30.   Chin CR*, et al.* Direct Visualization of HIV-1 Replication Intermediates Shows that
777   Capsid and CPSF6 Modulate HIV-1 Intra-nuclear Invasion and Integration. *Cell*
778   *Rep* **13**, 1717-1731 (2015).
779

780   31.   Li W*, et al.* CPSF6-Dependent Targeting of Speckle-Associated Domains
781   Distinguishes Primate from Nonprimate Lentiviral Integration. *mBio* **11**, (2020).
782

783   32.   Singh PK, Bedwell GJ, Engelman AN. Spatial and Genomic Correlates of HIV-1
784   Integration Site Targeting. *Cells* **11**, (2022).
785

786   33.   Ciuffi A, Bushman FD. Retroviral DNA integration: HIV and the role of
787   LEDGF/p75. *Trends Genet* **22**, 388-395 (2006).
788

789   34.   Llano M*, et al.* An essential role for LEDGF/p75 in HIV integration. *Science* **314**,
790   461-464 (2006).
791

792   35.   Engelman A, Cherepanov P. The lentiviral integrase binding protein LEDGF/p75
793   and HIV-1 replication. *PLoS Pathog* **4**, e1000046 (2008).
794

795   36.   Deshpande A*, et al.* Elasticity of the HIV-1 core facilitates nuclear entry and
796   infection. *PLoS Pathog* **20**, e1012537 (2024).
797

798   37.   Aiken C. Cell-free assays for HIV-1 uncoating. *Methods Mol Biol* **485**, 41-53
799   (2009).
800

801   38.   Adam SA, Sterne-Marr R, Gerace L. Nuclear protein import using digitonin-
802   permeabilized cells. *Methods Enzymol* **219**, 97-110 (1992).
803

804   39.   Wilson GL, Dean BS, Wang G, Dean DA. Nuclear import of plasmid DNA in
805   digitonin-permeabilized cells requires both cytoplasmic factors and specific DNA
806   sequences. *J Biol Chem* **274**, 22025-22032 (1999).
807

808   40.   Dombrowski M, Engeholm M, Dienemann C, Dodonova S, Cramer P. Histone H1
809   binding to nucleosome arrays depends on linker DNA length and trajectory. *Nat*
810   *Struct Mol Biol* **29**, 493-501 (2022).
811

812   41.   Vogler C*, et al.* Histone H2A C-terminus regulates chromatin dynamics,
813   remodeling, and histone H1 binding. *PLoS Genet* **6**, e1001234 (2010).
814

815   42.   Dorigo B, Schalch T, Bystricky K, Richmond TJ. Chromatin fiber folding:
816   requirement for the histone H4 N-terminal tail. *J Mol Biol* **327**, 85-96 (2003).
817

818   43.   Kan PY, Caterino TL, Hayes JJ. The H4 tail domain participates in intra- and
819   internucleosome interactions with protein and DNA during folding and
820   oligomerization of nucleosome arrays. *Mol Cell Biol* **29**, 538-546 (2009).

822 44. English CM, Adkins MW, Carson JJ, Churchill ME, Tyler JK. Structural basis for
823   the histone chaperone activity of Asf1. *Cell* **127**, 495-508 (2006).
824

825 45. Chavez MS, Scorgie JK, Dennehey BK, Noone S, Tyler JK, Churchill ME. The
826   conformational flexibility of the C-terminus of histone H4 promotes histone
827   octamer and nucleosome stability and yeast viability. *Epigenetics Chromatin* **5**, 5
828   (2012).
829

830 46. Wu H, Dalal Y, Papoian GA. Binding Dynamics of Disordered Linker Histone H1
831   with a Nucleosomal Particle. *J Mol Biol* **433**, 166881 (2021).
832

833 47. Zhou BR, Jiang J, Feng H, Ghirlando R, Xiao TS, Bai Y. Structural Mechanisms
834   of Nucleosome Recognition by Linker Histones. *Mol Cell* **59**, 628-638 (2015).
835

836 48. Hou Z, Nightingale F, Zhu Y, MacGregor-Chatwin C, Zhang P. Structure of native
837   chromatin fibres revealed by Cryo-ET in situ. *Nat Commun* **14**, 6324 (2023).
838

839 49. Kreysing JP, *et al.* Molecular architecture of heterochromatin at the nuclear
840   periphery of primary human cells. *bioRxiv*, 2025.2004.2009.647790 (2025).
841

842 50. Wang B, *et al.* The molecular basis of lamin-specific chromatin interactions. *Nat
843   Struct Mol Biol*, (2025).
844

845 51. Rohlfes N, *et al.* The nuclear localization signal of CPSF6 governs post-nuclear
846   import steps of HIV-1 infection. *PLoS Pathog* **21**, e1012354 (2025).
847

848 52. Ingram Z, *et al.* Spatiotemporal binding of cyclophilin A and CPSF6 to capsid
849   regulates HIV-1 nuclear entry and integration. *mBio* **16**, e0016925 (2025).
850

851 53. Price AJ, *et al.* CPSF6 defines a conserved capsid interface that modulates HIV-
852   1 replication. *PLoS Pathog* **8**, e1002896 (2012).
853

854 54. Lee K, *et al.* Flexible use of nuclear import pathways by HIV-1. *Cell Host Microbe*
855   **7**, 221-233 (2010).
856

857 55. Wei G, *et al.* Prion-like low complexity regions enable avid virus-host interactions
858   during HIV-1 infection. *Nat Commun* **13**, 5879 (2022).
859

860 56. Zhou H, *et al.* Quantitative spatial analysis of chromatin biomolecular
861   condensates using cryoelectron tomography. *Proc Natl Acad Sci U S A* **122**,
862   e2426449122 (2025).
863

864 57. Li Y, Zhang H, Li X, Wu W, Zhu P. Cryo-ET study from in vitro to in vivo revealed
865   a general folding mode of chromatin with two-start helical architecture. *Cell Rep*
866   **42**, 113134 (2023).

868 58. Tan ZY, Cai S, Noble AJ, Chen JK, Shi J, Gan L. Heterogeneous non-canonical
869   nucleosomes predominate in yeast cells in situ. *Elife* **12**, (2023).
870

871 59. Chen JK*, et al.* Nanoscale analysis of human G1 and metaphase chromatin in
872   situ. *EMBO J* **44**, 2658-2694 (2025).
873

874 60. Jang S*, et al.* HIV-1 usurps mixed-charge domain-dependent CPSF6 phase
875   separation for higher-order capsid binding, nuclear entry and viral DNA
876   integration. *Nucleic Acids Res* **52**, 11060-11082 (2024).
877

878 61. Burdick RC, Morse M, Rouzina I, Williams MC, Hu WS, Pathak VK. HIV-1
879   uncoating requires long double-stranded reverse transcription products. *Sci Adv*
880   **10**, eadn7033 (2024).
881

882 62. Li C, Burdick RC, Nagashima K, Hu WS, Pathak VK. HIV-1 cores retain their
883   integrity until minutes before uncoating in the nucleus. *Proc Natl Acad Sci U S A*
884   **118**, (2021).
885

886 63. Burdick RC*, et al.* Dynamics and regulation of nuclear import and nuclear
887   movements of HIV-1 complexes. *PLoS Pathog* **13**, e1006570 (2017).
888

889 64. Francis AC, Marin M, Prellberg MJ, Palermino-Rowland K, Melikyan GB. HIV-1
890   Uncoating and Nuclear Import Precede the Completion of Reverse Transcription
891   in Cell Lines and in Primary Macrophages. *Viruses* **12**, (2020).
892

893 65. Bejarano DA*, et al.* HIV-1 nuclear import in macrophages is regulated by CPSF6-
894   capsid interactions at the nuclear pore complex. *Elife* **8**, (2019).
895

896 66. Francis AC, Melikyan GB. Single HIV-1 Imaging Reveals Progression of Infection
897   through CA-Dependent Steps of Docking at the Nuclear Pore, Uncoating, and
898   Nuclear Transport. *Cell Host Microbe* **23**, 536-548 e536 (2018).
899

900 67. Balasubramaniam M*, et al.* PF74 Inhibits HIV-1 Integration by Altering the
901   Composition of the Preintegration Complex. *J Virol* **93**, (2019).
902

903 68. Müller TG*, et al.* Lenacapavir-induced capsid damage exposes HIV-1 genomes
904   from nuclear speckles. *bioRxiv*, 2025.2007.2007.663551 (2025).
905

906 69. Liu S*, et al.* CPSF6 regulates alternative polyadenylation and proliferation of
907   cancer cells through phase separation. *Cell Rep* **42**, 113197 (2023).
908

909 70. Boettiger AN*, et al.* Super-resolution imaging reveals distinct chromatin folding for
910   different epigenetic states. *Nature* **529**, 418-422 (2016).
911

71.   Strom AR, Emelyanov AV, Mir M, Fyodorov DV, Darzacq X, Karpen GH. Phase separation drives heterochromatin domain formation. *Nature* **547**, 241-245 (2017).

72.   Schindelin J*, et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012).

73.   Arnold J*, et al.* Site-Specific Cryo-focused Ion Beam Sample Preparation Guided by 3D Correlative Microscopy. *Biophys J* **110**, 860-869 (2016).

74.   Zheng SQ, Palovcak E, Armache JP, Verba KA, Cheng Y, Agard DA. MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017).

75.   Kremer JR, Mastronarde DN, McIntosh JR. Computer visualization of three-dimensional image data using IMOD. *J Struct Biol* **116**, 71-76 (1996).

76.   Himes BA, Zhang P. emClarity: software for high-resolution cryo-electron tomography and subtomogram averaging. *Nat Methods* **15**, 955-961 (2018).

77.   Bilokapic S, Strauss M, Halic M. Histone octamer rearranges to adapt to DNA unwrapping. *Nat Struct Mol Biol* **25**, 101-108 (2018).

78.   Qu K*, et al.* Structure and architecture of immature and mature murine leukemia virus capsids. *Proc Natl Acad Sci U S A* **115**, E11751-E11760 (2018).

79.   Pettersen EF*, et al.* UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* **25**, 1605-1612 (2004).

80.   Ni T*, et al.* Structure of native HIV-1 cores and their interactions with IP6 and CypA. *Sci Adv* **7**, eabj5715 (2021).

81.   Armache JP*, et al.* Localization of eukaryote-specific ribosomal proteins in a 5.5-A cryo-EM map of the 80S eukaryotic ribosome. *Proc Natl Acad Sci U S A* **107**, 19754-19759 (2010).

82.   Zivanov J*, et al.* A Bayesian approach to single-particle electron cryo-tomography in RELION-4.0. *Elife* **11**, (2022).

83.   Goddard TD*, et al.* UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci* **27**, 14-25 (2018).

84.   Liu YT, Zhang H, Wang H, Tao CL, Bi GQ, Zhou ZH. Isotropic reconstruction for electron tomography with deep learning. *Nat Commun* **13**, 6482 (2022).

957 85. Isensee F, Jaeger PF, Kohl SAA, Petersen J, Maier-Hein KH. nnU-Net: a self-
958   configuring method for deep learning-based biomedical image segmentation. *Nat*
959   *Methods* **18**, 203-211 (2021).
960

961 86. Ermel UH, Arghittu SM, Frangakis AS. ArtiaX: An electron tomography toolbox for
962   the interactive handling of sub-tomograms in UCSF ChimeraX. *Protein Sci* **31**,
963   e4472 (2022).
964

965 87. Marko JF, Siggia ED. Bending and twisting elasticity of DNA. *Macromolecules* **27**,
966   981-988 (1994).
967

968 88. Bustamante C, Marko JF, Siggia ED, Smith S. Entropic elasticity of lambda-
969   phage DNA. *Science* **265**, 1599-1600 (1994).
970

971 89. Hagerman PJ. Flexibility of DNA. *Annual review of biophysics and biophysical*
972   *chemistry* **17**, 265-286 (1988).
973

974 90. Marko JF, Siggia ED. Stretching dna. *Macromolecules* **28**, 8759-8770 (1995).
975
976

a mNeongreen-IN SiR-DNA

c

Core number

Status of imported cores: Just-imported, En-route, Deep-in
Values: 25, 21, 9

d Ribosome NPC NE NE NPC Chromatin fibre Nuclear granule Nuclear factors WT HIV-1 core Nucleosome Nuclear granule Nucleus **Just-imported**

e Ribosome | NE | NPC | CA hexamer | CA pentamer | Nuclear factors | DNA | Core histones | H1

f Nucleosome Nuclear factors NE WT HIV-1 core Nucleus Mito ER Ribosomes **En-route**

g Ribosome | Mito & ER | CA hexamer | CA pentamer | Nuclear factors | DNA | Core histones | H1
Top bound of the tomogram

h Nuclear factors WT HIV-1 core Nuclear granule Nucleosome Nucleus **Deep-in**

i CA hexamer | CA pentamer | Nuclear factors | DNA | Core histones | H1

Figure 1. Correlative cryo-ET of nuclear trafficking of HIV-1. a-b, A representative cryo-fluorescence image of a lamella (**a**) and overlapped with a low magnification of cryo-EM image of the same lamella (**b**). HIV-1 cores are indicated by white arrowheads. Nuclei are outlined by white dashed lines. Inset in (**a**) is an enlarged view of the area containing HIV-1 cores. Scale bars = 5 μm. **c,** A bar chart showing the distribution of HIV-1 cores captured in nuclei at the stage of "Just-imported (<75 nm from NPC), "En-route" (between 75 nm to 1.5 μm from NPC), and "Deep-in" (> 1.5 μm from NPC). **d,** A representative tomographic slice of a "Just-imported" cone-shaped WT HIV-1 core. The NPC, ribosomes, nucleosomes, prominent chromatin fibres, nuclear granules, and nuclear factors coating the core are labelled. The nucleus and nuclear envelope (NE) are annotated accordingly. **e,** The segmented volume of (**d**), shown as an overview (left) and zoomed-in views of the "Just-imported" WT core (upper right) and zig-zag organized nucleosomes connected by black dashed lines (right lower). HIV-1 CA hexamers and pentamers, NPC, DNA, core histones and histone H1 of nucleosomes, ribosomes, nuclear factors, and NE are segmented and mapped back with indicated colours. Of note, two subunits of the right NPC exceeds the upper bound of the tomogram, thus are not depicted. **f,** A representative tomographic slice of an "En-route" tube-shaped WT core. Labelling of the subjects is the same as in (**d**). NE is indicated by the white dashed lines. **g,** The segmented volume of (**f**), shown as an overview (left) and zoomed-in views of the "En-route" WT core (upper right) and zig-zag organized nucleosomes connected by black dashed lines (right lower). Of note, one end of the "En-route" core exceeds the top bound (red dashed line) of the tomogram, thus is not fully depicted. HIV-1 CA hexamers and pentamers, mitochondria/ER, DNA, core histones and histone H1 of nucleosomes, ribosomes, nuclear factors are segmented and mapped back with indicated colours. NE is indicated by the black dashed lines. **h,** A representative tomographic slice of a "Deep-in" tube-shaped WT core. Labelling of the subjects is the same as in (**d**). **i,** The segmented volume of (**h**), shown as an overview (left) and zoomed-in views of the "Deep-in" WT core (upper right) and zig-zag organized nucleosomes connected by black dashed lines (right lower). HIV-1 CA hexamers and pentamers, DNA, core histones and histone H1 of nucleosomes, and nuclear factors are mapped back and shown with indicated colours. Scale bars = 100 nm in **d, f, h**.

a

Linker DNA

Linker DNA

DNA
H3
H4
H2A
H2B
H1

Class 1 (68.85%)

b

30°

H2A

30°

H3

c

Class 2 (8.07%)   Class 3 (10.93%)   Class 4 (12.15%)

d

Class 4   Class 1

H1

Linker DNA

H1

Linker DNA

Figure 2. *In situ* structure and dynamics of nucleosomes. a, Cryo-ET STA structure of the major class (class 1) of *in situ* H1-bound nucleosome at 5.6 Å resolution. Histones and DNAs are colored as indicated. Two orthogonal views are displayed with core histones shown at 6σ, and H1 and linker DNA shown at 3σ. Linker DNA indicated by black arrows. **b**, The histone proteins are fitted with atomic model (PDB 7PEX). DNA densities are removed for clarity. Two orthogonal views are displayed with regions containing H2A and H3 tails enlarged on the right with model coloured red. **c**, Cryo-ET STA structures of class 2 (stacking H1-bound nucleosomes), class 3 (core nucleosome), and class 4 (open-linker H1-bound nucleosome). Two orthogonal views are depicted, class 2 is fitted with the atomic model (PDB 7PEX), class 3 and class 4 are fitted with the atomic model (PDB 2CV5). Class 2 is contoured at 3σ, class3 is contoured at 5σ, class 4 is contoured at 2.5σ with linker DNA indicated by black arrows. The deviation of the linker DNA density from the model is indicated by the dashed curve. **d**, Comparison between H1-bound nucleosome (class1) and open-linker H1-bound nucleosome (class 4). Two orthogonal views are depicted. The trajectory of the linker DNA is indicated by dashed arrows and H1 is coloured pink.

Figure 3. *In situ* **chromatin architecture in the nucleus. a**, A map-back representation of nucleosomes linked by linear DNA strand of various lengths in a tomogram collected from a heterochromatin region without HIV-1 cores. Only zig-zag linked nucleosomes (see methods section) are highlighted for clarity. **b**, A histogram showing the distribution of nucleosomes in various zig-zag connected oligomers (n = 253 tomograms). **c**, Representative zig-zag linked di-, tri-, tetra-, penta-, and hexa-nucleosomes. **d**, A map-back representation of four characterized nucleosome classes in a tomogram collected from a heterochromatin region without HIV-1 core. Nucleosome classes are colored accordingly. **e**, Pair-wise nucleosome distance distribution (n = 253 tomograms), the peak is indicated. **f**, The distribution of nucleosome classes in periphery (distance to NE $\leqq$ 1.5 µm) (n = 235 tomograms) and interior (distance to NE > 1.5 µm) regions of nuclei (n = 18 tomograms). The enrichment factor is calculated by the radio of regional nucleosome proportion over the global proportion of each class. Two-sided Fisher's exact test is applied for each class: for class 1, p = 1.8372 × 10^{-11} < 0.0001, n = 146,656 ($\leqq$ 1.5 µm), n = 11,780 (> 1.5 µm); for class 2, p = 0.3654 > 0.05, no significance, n = 17,168 ($\leqq$ 1.5 µm), n = 1,395 (> 1.5 µm); for class 3, p < 1 × 10^{-15}, n = 22,844 ($\leqq$ 1.5 µm), n = 2,309 (> 1.5 µm); for class 4, p = 0.1721 > 0.05, no significance, n = 25,748 ($\leqq$ 1.5 µm), n = 2,206 (> 1.5 µm). **g**, The distribution of nearest neighbour classes (coloured) for each nucleosome class. The enrichment factor is calculated by the ratio of local (the nearest neighbor) nucleosome proportion over the global proportion of each class. Two-sided Chi-square test is applied for each class: for class 1, p < 1 × 10^{-15}, n = 158,436; for class 2, p < 1 × 10^{-15}, n = 18,563; for class 3, p < 1 × 10^{-15}, n = 25,153; for class 4, p < 1 × 10^{-15}, n = 27,954.

Figure 4. Chromatin exclusion by HIV-1 cores during nuclear trafficking. a, Heat maps from the three tomograms in **Figure 1d-i**, showing the map-backed nucleosome density (projected over consistent 130 nm thick tomogram) during HIV-1 nuclear traversal. Areas occupied by HIV-1 cores are indicated by purple arrowheads. NPC occupation is indicated by the white arrowhead. NE is indicated by the black dashed lines. **b**, The distributions of pair-wise inter-nucleosome distance and distance between HIV-1 CA and nucleosomes. **c**, Map-back of nucleosomes by classes in interior region of the nucleus containing a HIV-1 core, with all classes included and coloured accordingly. The HIV-1 core and nuclear factors are mapped back as in **Figure 1i**. **d**, The distribution of nucleosome classes in the nearest neighbours of HIV-1 cores compared to the global distribution. Two-sided Fisher's exact test is applied: for class 1, $p < 1 \times 10^{-15}$, n = 2,787; for class 2, $p = 1.14523030 \times 10^{-7} < 0.0001$, n = 265; for class 3, $p < 1 \times 10^{-15}$, n = 772 ; for class 4, $p = 3.02791446817 \times 10^{-4} < 0.001$, n = 621.

a

Nucleosome

Chromatin fibre

Nuclear granule

NE

Nucleus

N74D HIV-1 core

Nuclear granule

NPC

b

Ribosome | NE | ER | NPC | N74D CA | DNA | Core histones | H1

c

WT
N74D

Percentage (%)

Just-imported   En-route   Deep-in

d

Normalised grey value

7 nm

WT outside
WT imported
N74D imported

Distance (nm)

e

WT outside

WT imported

N74D imported

f

90°

90°

65 Å
Nuclear factors

90°

Figure 5. CPSF6 facilitates the nuclear traversal of HIV-1. a-b, A representative tomographic slice a "Just-imported" cone-shaped N74D core **(a)** and corresponding segmented volume **(b)**. No discernable core-surrounding density is observed (enlarged in the inset). The NPC, ribosomes, nucleosomes, nuclear granules, and chromatin fibres are labelled. The nucleus, NE, and ER are annotated accordingly. Ribosomes, NE, ER, N74D CA hexamers, DNA, core histones and histone H1 of nucleosomes are segmented and mapped back with the indicated colours. Of note, three subunits of the NPC exceeds the upper bound of tomogram, thus are not depicted. Scale bar, 100 nm. **c,** Distributions of WT and N74D cores in the nucleus. Fisher's exact test is applied, p = 0.0424 < 0.05, n of WT cores = 55, n of N74D cores = 10. **d,** Density profiling of the vicinity of cores. The same strategy illustrated in **Supplementary Figure 9** is applied with shorter lines. For WT outside cores, n of lines = 155, n of cores = 12. For WT imported cores, n of lines = 111, n of cores = 10. For N74D imported cores, n of lines = 101, n of cores = 10. The central lines represent the mean, and whisks represent the S.D. The distance from the surface of imported WT HIV-1 cores to the density center of nuclear factors approximates 7 nm. **e,** Cryo-ET STA structures of the HIV-1 CA hexamers in capsid lattices from WT cores outside the nucleus (top), WT cores inside the nucleus (middle), and N74D cores inside the nucleus (bottom). The left panel: top views of the STA map (contoured at 2.5σ) with the central hexamer coloured; the middle panel: top views of the STA map (WT contoured at 2.5σ, N74D contoured at 1.5σ) fitted with a CA hexamer model (PDB 6SKK); the right panel: A slice of CA hexamer density maps (position shown in the upper left corner) fitted with atomic model (PDB 6SKK). **f,** CPSF6 density shown in the imported WT CA STA map (middle panel with an arrowhead), are absent in the STA maps of imported N74D cores (bottom) and outside WT cores (top). The left panel: CA hexamer density maps contoured at 0.25σ, coloured according to the height, from white (bottom) to purple (top); the right panel: the side views of CA hexamer density maps (major hexamer density contoured at 2.5σ for WT and 1.5σ for N74D, density of nuclear factors contoured at 0.25σ), the distance between CA hexamer and center of nuclear factors measures approximately 65 Å for the imported WT core.

Figure 6. Schematic model of HIV-1 nuclear trafficking. This schematic summarizes the distinct behaviors of WT and CPSF6-binding-deficient (N74D) HIV-1 cores during the nuclear trafficking. Upon nuclear import, WT HIV-1 cores rapidly engage nuclear factors such as CPSF6, which assemble into a phase-separation-like layer that excludes chromatin along the trajectory of core traversal. This CPSF6-associated environment facilitates HIV-1 core movement through the condensed heterochromatin at the nuclear periphery. As the cores traverse into the nuclear interior, they encounter regions of less-compact chromatin enriched in dynamically positioned nucleosomes, likely corresponding to transcriptionally active domains. In contrast, N74D HIV-1 cores, which fail to interact with CPSF6 and related nuclear factors, are unable to penetrate the dense peripheral heterochromatin and consequently remain trapped at the nuclear periphery.