Evolutionary and ecological determinants of the phenology of births in wild large herbivores, a systematic review

Lucie Thel[1], Christophe Bonenfant[2], Simon Chamaillé-Jammes[3,4]

Authors details

[1] Department of Conservation Management, Faculty of Science, George Campus, Nelson Mandela University, George, South Africa.

[2] Laboratoire de Biométrie et Biologie Évolutive, Unité Mixte de Recherche 5558, Université Lyon I, Bâtiment 711, 43 Boulevard du 11 novembre 1918, 69622 Villeurbanne Cedex, France.

[3] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France.

[4] Mammal Research Institute, Department of Zoology & Entomology, University of Pretoria, South Africa.

Corresponding author: Lucie Thel, lucie.mcv.thel@gmail.com.

Abstract

1. Introduction: Timing of birth is a major determinant of individuals' reproductive success and survival, and is driven by multiple internal and external factors. However, there is currently no quantitative assessment of the extent and support received by the drivers of the phenology of births in mammals. Due to their diverse phenology of births, large herbivores represent an ideal case study to quantify this.

2. Aims: our objective was to identify the different drivers tested for their effect on the phenology of births in large herbivores and how much empirical support they received, as well as determine thematic, taxonomic and spatial gaps in the literature.

3; Methods: We conducted a systematic review focusing on the drivers of the phenology of births in wild large herbivores. We extracted the outcomes of 124 relevant articles and assessed the reliability of their findings using a general index based on the quality of the data and methods used.

4. Results and Discussion: As expected, the two dominant hypotheses (seasonality and predation) received strong support regarding their effect on birth timing and synchrony, respectively. The effect of female, offspring and population characteristics remained marginally accounted for, even though there was evidence of their importance in the determination of the phenology of births. We identified a major research gap in Asian and South and Central American species.

5. Conclusions: We encourage research in these understudied domains, as well as large scale multi-species comparative analyses to fill in the gaps and improve our understanding of reproductive phenology.

Keywords: food resource, population ecology, predation, reproduction, seasonality, synchrony, timing.

Introduction

The timing of birth is a major determinant of an individual's reproductive success, as it may affect both the short- and long-term survival of the offspring (e.g., via early growth, Côté and Festa-Bianchet 2001; via density-dependent phenomena, Williams et al. 2013), the survival of the parent, and possibly the later reproductive success of both the parent and the offspring (Green and Rothstein 1993a). The timing of birth can be affected by numerous drivers, such as the seasonality of the environment (e.g., food availability, Bunnell 1982; temperature variations, McNutt et al. 2019; date of snowmelt, Laforge et al. 2023), interacting with life history traits (e.g. diet, Sinclair et al. 2000; mating system, Boness et al. 1995; sociality, Bertram 1975) and predation (O'Donoghue and Boutin 1995). These drivers can induce both plastic and evolutionary responses in the phenology of births (Radchuk et al. 2019). For instance, most mammal species in the northern hemisphere, such as the roe deer (*Capreolus capreolus*) and the Eurasian lynx (*Lynx lynx*), have evolved to give birth in spring when the temperature becomes milder, reducing the risks of hypothermia for the newborn, and the vegetation blooms, providing sufficient food of high quality to cope with the energetic costs of gestation and lactation (Plard et al. 2014, Mattisson et al. 2022). But there is evidence that individuals can also actively tune their parturition date by adjusting their gestation length in response to variations of temperature and snow conditions during the ongoing year (e.g., in the reindeer (*Rangifer tarandus*), Paoli et al. 2020). The phenology of births is thus the result of a complex interaction between multiple drivers acting at different spatio-temporal scales.

The study of the phenology of births consists mainly in describing the timing and synchrony of births in a given population and understanding their drivers (Thel et al. 2022). Although some main patterns of animal phenology have been recognized since long, timing and synchrony of births are highly variable in time, space, but also between and within species and even populations (Bronson 1989). The phenology of births in mammals ranges

from year-round breeding (e.g., the chacma baboon (*Papio ursinus*) in Tsaobis Nature Park,

Namibia, Dezeure et al. 2021) to a concentration of births during a very short period of time

(e.g., 80% of births happen in less than 30 days in a French population of roe deer, Gaillard et

al. 1993). The phenology of births can vary from one species to the other in the same location

(e.g., in the guild of East African antelopes in the Serengeti, Tanzania, Sinclair et al. 2000),

but also from one population to the other for the same species across its geographical range.

In Africa, the impala (*Aepyceros melampus*) gives birth year round in Kenya and Tanzania,

but seasonally in Zambia, Botswana and Rwanda (Ansell 1960, Dasmann and Mossman

1962, Leuthold and Leuthold 1975, Sinclair et al. 2000, Moe et al. 2007).

The study of the drivers of the phenology of births has been a major research avenue,

producing numerous studies going back a long way in time (e.g., in large herbivores: Ansell

1960, Spinage 1973, Bunnell 1980, Gaillard et al. 1993, Post et al. 2003, Ryan et al. 2007,

Ogutu et al. 2010, Paoli et al. 2018, Neumann et al. 2020) and generating several qualitative

and quantitative reviews and book chapters (e.g., Rutberg 1987, Bronson 1989, Post 2003,

Brockman and van Schaik 2005, Zerbe et al. 2012, Heldstab et al. 2018). Two main

hypotheses emerged from this abundant literature: the seasonality hypothesis stating that the

timing of births is primarily set by the seasonality of climate and food resources (Bunnell

1980, Bunnell 1982, Festa-Bianchet 1988, Post 2003), and in the case of prey species such as

large herbivores and rodents, the predation hypothesis stating that the synchrony of births is

shaped, in addition to environmental drivers, by the type of predation and the antipredator

behaviour of the juveniles (Estes 1976, Rutberg 1987, Ims 1990a, Ims 1990b, O'Donoghue

and Boutin 1995). Although intrinsically related (a population displaying asynchronous births

does not show a set timing of births), timing and synchrony do not necessarily respond to the

same selection pressures. The seasonality hypothesis is usually accepted as affecting the

timing of birth at the individual scale, whereas the predation hypothesis emerges from

population-scale properties and affects the synchrony of births. Over the last century, a growing number of more specific hypotheses have been put forward to account for the deviation of birth phenology from the general matching with environmental seasonality. For instance, it is generally accepted that grazer species should have a shorter birth period than browser species because the availability of their food resource, mainly composed of grasses, is more concentrated in time than tree-based resources such as leaves, preferred by browsers (Leuthold and Leuthold 1975).

Due to their great diversity in their phenology of births as well as their pivotal position in the ecological processes as a primary consumer and a prey, large herbivores provide an ideal framework for analysing and understanding the regulatory processes of the phenology of births at the mammal scale. Large herbivores are present at almost all latitudes from the poles (e.g., the caribou (*Rangifer tarandus*), Post et al. 2003) to the equator (e.g., the Bornean yellow muntjac (*Muntiacus atherodes*), Giman et al. 2007), and occur in habitats ranging from sea level (e.g., the moose, Singh et al. 2012) to mountainous landscapes as elevated as 5000 m a.s.l. (e.g., the alpine musk deer (*Moschus chrysogaster*), Buzzard et al. 2018). Those species present a large variety of life history traits in terms of their diet (Hofmann 1989), sociality (Szemán et al. 2021), and reproductive traits (e.g., gestation length, Owen-Smith and Ogutu 2013; mating system, Bowyer et al. 2020), amongst others. Additionally, they are exposed to a wide range of predation pressures, from the complete absence of predators, as it has been the case for almost two centuries in Rum Island, Scotland (Shi et al. 2010), to the full guild of medium and large predators such as in the Serengeti ecosystem, Tanzania (Sinclair and Northon-Griffiths 1995), including recovering systems such as Yellowstone National Park, United States of America (Ripple et al. 2010). Such a diversity of conditions and traits gathered in one clade has generated a vast range of research and participated in the emergence and development of major ecological theories revolving

around reproductive biology (e.g., density-dependence effects, Bonenfant et al. 2009; capital-income breeding strategies, Apollonio et al. 2020), some of them directly related to the phenology of births (e.g., Gosling 1969, Post 2003).

In this study, we conducted a systematic review of the scientific literature focusing on the drivers of the phenology of births in large herbivores living in natural conditions. Current reviews on the phenology of births are either dated and do not include the latest publications on the subject (e.g., Rutberg 1987), they focus on a limited geographical range (e.g., Bunnell 1982), or they are based mainly on captive individuals (e.g., Zerbe et al. 2012). Additionally, empirical tests conducted on some populations led to inconclusive outcomes or results contradicting theoretical expectations. Currently, there is no quantitative assessment of the reliability and empirical validity of the hypotheses tested in terms of phenology of births in wild large herbivores. A comprehensive summary of the methodological approaches used to explore these different hypotheses is also lacking. Using a systematic search on the Web of Science and complementing it with our personal list of references, we identified 124 relevant scientific articles published since the early 1960's, from which we summarised the different hypotheses and how much empirical support they have received so far. We highlighted the hypotheses, the geographic areas and the taxonomic groups that are currently understudied and we suggested new avenues of research for a better understanding of the drivers of the phenology of births in large herbivores, and in mammals in general.

Methods

Focus of the literature search

The "ungulates" consist in a polyphyletic group generally used to describe herbivorous hoofed mammals with a body mass > 2 kg. It commonly includes the orders Artiodactyla (i.e., even number of fingers) and Perissodactyla (i.e., uneven number of fingers), but can be

extended to some large marsupials and to the Proboscidea (i.e., characterised by a trunk)

according to the definition selected (Danell et al. 2006). As part of the Artiodactyla, the Suina

(i.e., pigs and peccaries) and the Cetacea (i.e., aquatic mammals), which are not strictly

herbivorous, can also be included in the ungulates *lato sensu*. Based on their similarities in

terms of eco-evolutionary traits, we retained in this systematic review the Artiodactyla

(including the Suina but not the Cetacea), the Perissodactyla and the Proboscidea, and refer to

them as "large herbivores".

Systematic literature search

We conducted an initial literature search (16th of July, 2025) on the Web of Science

(www.webofscience.com, WoS hereafter) using all databases available (Web of Science Core

Collection, Grants Index, Inspec, KCI-Korean Journal Database, Medline, Preprint Citation

Index, ProQuest Dissertations & Theses Citation Index, SciELO Citation Index). We

searched all scientific articles, review articles, books and data papers published before 2025

(i.e., 2025 excluded) and written in English, included in the following catalogues (inbuilt in

the WoS search engine): "reproductive biology", "environmental science ecology",

"zoology", "behavioural sciences", "biodiversity conservation", "demography",

"evolutionary biology" and "remote sensing". To conduct the search, we used the following

list of keywords: "phenolog* OR *synchron* OR timing* OR period* OR *season* OR

date* OR pattern*" ("OR" allowed us to search for records containing any of the terms

separated by the operator; "*" allowed us to search for words with variant zero to many

characters, e.g., the use of "period*" allowed us to search for words such as "period",

"periods", "periodic", "periodicity"). We combined these terms with the Boolean connector

"AND" ("AND" finds records containing both terms separated by the operator) with the

following list of keywords: "breed* OR birth* OR parturition* OR reproduc* OR calv* OR

fawn* OR lamb* OR foal* OR farrow* OR litter* OR offspring* OR newborn*", and looked

for this combination of keywords in the title of the articles referenced in the WoS.

We obtained n = 15,919 initial results (see PRISMA diagram in Supplementary

Material 1, Page et al. 2021). We excluded n = 28 duplicates based on identical DOI and/or

identical combination of title and list of authors. Based on a quick screening of the title (and

abstract, when necessary), we excluded n = 15,822 irrelevant articles that were either not

dealing with large herbivores reproduction, or when dealing with large herbivores

reproduction, that were focusing on: i) populations subject to experimental designs or

veterinary treatments; ii) populations in captive, farming or zoo conditions; iii) non-endemic

or feral populations; iv) subjects not dealing with phenology of births in general (e.g., male

reproduction, mating instead of parturition phenology). We also excluded purely

methodological articles, which presented methods to estimate the date of birth but did not

explore the effect of any driver on the phenology of births. We extracted n = 69 articles

dealing explicitly with the phenology of births in large herbivores in wild conditions.

Although the WoS references publications since 1637, we mostly had access to publications

after 1993 for some databases due to academic subscription restrictions, and we acknowledge

that due to probable imperfect referencing of the oldest studies (Say-Sallaz et al. 2019), the

first half of the twentieth century and before was likely less well covered than the more recent

period in our literature review. We thus added relevant articles from our personal

bibliography that did not come up in the systematic search (n = 76 articles).

We conducted a detailed analysis based on the reading of n = 145 full articles. From

this final screening, we only included the articles directly exploring the phenology of births in

large herbivores in wild conditions and their potential drivers (i.e., excluding purely

descriptive articles). Eventually, we included a total of n = 124 articles in the review (WoS =

63, personal bibliography = 61, Supplementary Material 2).

Data extraction from the articles

We extracted manually the following information from each article included in the review:

references of the article (e.g., year of publication and journal), references of each population

studied (e.g., species vernacular and scientific names, sample size), references of the study

site and study period (e.g., country and duration of the study; see Supplementary Material 1

for a detailed description of each variable extracted). Each variable explored in a given

population was recorded as one unique observation, which means that an article could contain

more than one observation even if a single hypothesis was tested. For instance, an article

studying the effect of vegetation phenology on two different populations of caribou consisted

in two observations.

For each variable, we extracted the citation of the hypothesis when available (i.e., the

formal description of the expectations as formulated by the authors, usually in the

Introduction section of the article). In several instances, a variable was tested while the

authors did not explicitly state their expectations. For example, we encountered this situation

when the authors tested the effect of the sex of the offspring on the timing of birth as a

control variable, but did not initially mention if they expected an effect or not, or the direction

of the effect. We then classified each variable tested according to 39 factors (e.g., female age,

latitude, population density; see Results section for the detailed list), themselves grouped into

seven main themes (e.g., the factors mentioned above belong to the themes: female

characteristics, environmental characteristics and population characteristics, respectively; see

Results section for the detailed list). We identified the parameter of phenology addressed by

the variable, i.e., the timing or synchrony of births. We used the codes "supported", "partially

supported" or "unsupported" to report on the outcome of the study. We considered the effect

of a variable as "supported" if the associated significance value reported by the authors was

below the level alpha = 0.05 (statistical approach). In the absence of a formal statistical test

(descriptive approach), we followed the conclusions of the authors, which could be reported

either in the Results or in the Discussion section. In the case of a significant effect detected

only on a subset of the population (e.g., when considering male but not female offsprings, or

when considering multiparous but not primiparous females), we recorded that the hypothesis

was "partially supported".

Quality index

To assess the reliability of the hypothesis testing procedure in the articles retrieved for this

review, we calculated a general quality index based on the data collection method, the sample

size, the study period length and the type of test (statistical or descriptive) used in the articles.

For each variable tested in an article, we attributed a value between 1 (lowest quality) and 3

(highest quality) to the data collection method, the sample size, the study period length and

the type of test conducted (see Supplementary Material 3 for details). We then summed these

four values to obtain an index for each variable tested.

We explored the change in quality across time using a linear regression with a

quadratic effect (assuming a potential stabilisation of quality over time) to model the

variation of the quality index of the articles (mean of the quality indexes of all the variables

tested in the article) according to the year of publication. We also explored the variation of

quality according to the continent and taxonomic family of the populations studied in the

articles using ANOVAs. Due to small sample sizes for some taxonomic families (sample size

for Antilocapridae, Elephantidae, Hippopotamidae, Moschidae, Rhinocerotidae and

Tayassuidae ≤ 3), we proceed to the following groupments: Tayassuidae were grouped with

Suidae, Moschidae were grouped with Cervidae, Antilocapridae were grouped with Bovidae,

and finally, Elephantidae, Hippopotamidae and Rhinocerotidae were grouped together. All

statistical analyses were conducted using the R software (version 4.5.1, R Core Development

Team 2025). The data used in the analyses is available from the Zenodo Repository

(Anonymised reference).

Results

Temporal, geographical and taxonomic focus

We identified a total of n = 124 articles published between 1962 and 2023 in 52 different peer

reviewed journals (Fig. 1a). The number of articles published increased quadratically since

the 1960's (Fig. 1b). A total of 82 species (Fig. 2) distributed across 29 countries were

included in the articles. One article did not directly mention the species names (English et al.

2012) and two articles did not directly mention the country of origin of the populations

included in the study (English et al. 2012, Owen-Smith and Ogutu 2013). Almost half (46%)

of the species were studied in only one article (e.g., the chital (*Axis axis*), Ahrestani et al.

2011), and a very limited number of species appeared in more than five articles (n = 16, e.g.,

the roe deer appeared in 13 articles, the reindeer and bighorn sheep (*Ovis canadensis*) in 12

articles). Several populations characterised by continuous long-term monitoring were subject

to multiple publications (e.g., the red deer (*Cervus elaphus*) of Rum Island, Scotland; the roe

deer of Trois Fontaines, France; the guild of large herbivores of Serengeti National Park,

Tanzania). Most of the articles focused on one or two populations (n = 96, from the same or

different species), and 11 articles included more than 10 populations (range = [1; 80]

populations studied in a given article, all the articles including more than 25 populations

corresponding to datasets collected from literature reviews). For the articles providing the

duration of the study or the starting and ending dates of data collection for the populations

studied (n = 170), the average study length (± standard deviation) was 11 ± 11 years (median

[Q1; Q3] = 6 [3; 14] years, Fig. 3).

At the time of this study, the continent most represented amongst the articles retrieved

was North America with 46 articles, while we found only four articles in Central and South

America (Fig. 4). Three articles included study sites on several continents ("worldwide"

category in Table 1). The two main data collection methods, mostly used in North America

and Europe, were neonate captures and direct observations, the latter also mainly used in

Africa (Fig. 5). Studies conducted in North America also frequently used movement tracking

devices (i.e., identification of parturition date from the mother's movements and/or

identification of juvenile survival from the juvenile's movements) and less frequently vaginal

implant transmitters, while birth date prediction from foetuses was a fairly common method

in Africa. Four other methods were infrequently used to estimate birth date and consisted in

aerial surveys, camera trapping, hormonal assay of reproductive stage of the female based on

faecal samples, and aged culled individuals to identify the date of birth.

Hypotheses and analytical approach

We identified three main ways to approach the analyses in the articles retrieved. The most

common one (n = 311) corresponded to what we defined here as a "closed hypothesis". In

this case, the expectation was clearly stated by the authors (e.g., "*As grass is more seasonal*

than browse, grazers are expected to have a shorter birth season length than mixed feeders

or browsers", English et al. 2012). In what we defined as an "open hypothesis" approach (n =

161), an hypothesis was stated by the authors, but there was no specific expectation regarding

the effect of the variable tested (e.g., "*We tested relationships between the timing of*

parturition and maternal age", Adams and Dale 1998). In some instances, we could not

identify any statement announcing that the effect of a given variable would be tested even

though the effect or absence of effect of this variable would be reported in the Results section

(n = 98). We recorded these cases as the "absence of clear hypothesis" (e.g., Berger 1992

mentioned their hypotheses and expectations after presenting their results: "*Hence the data*

support the hypothesis that (1) late-breeding females in good body condition shorten

gestation, synchronizing births with other females"). In most cases, the effect of a variable

was tested with a statistical approach (n = 391) rather than a descriptive approach with no

statistical support to draw conclusions, but rather a visual correlation observed by the authors,

such as a concentration of births during the rainy season (n = 181). However, 61% of the

variables were tested descriptively rather than statistically in articles published on African

populations, reaching 67%, on Central and South American populations.

Quality index

There was a significant relationship between the year of publication and quality index

of the articles (n = 103 articles included in the analysis; β_{year} = 5.28 ± 2.25, p = 0.02; $\beta_{year*year}$ =

-1.30 ± 0.00, p = 0.02; adj. R^2 = 0.27), with quality increasing over time but progressively

reaching a plateau after 2005 (Supplementary Material 3). The quality index varied

depending on the continent of the study (p < 0.01), and was significantly larger in Europe and

North America than in Africa, Asia and Central and South America (Supplementary Material

3). The quality index also varied depending on the taxonomic family of the population

studied (n = 42 species included in the analysis; p < 0.01; Supplementary Material 3). It was

significantly larger for Cervidae-Moschidae than for any other taxonomic family (all p <

0.05), except Girafidae (p = 0.07). All other taxonomic families did not show significant

differences (all p > 0.05).

Variables tested

We identified seven themes: i) environmental characteristics, ii) plant characteristics, iii) life

history traits, iv) female characteristics, v) offspring characteristics, vi) population

characteristics and vii) predation exposure. In each theme we found articles focusing on the

timing but also the synchrony of births (Table 2). In some themes, articles were however

mainly focused on one or the other (e.g., female, juvenile and population characteristics often

tested for their effect on timing of births; predation exposure mostly tested for its effect on

synchrony of births). These themes encompassed 39 factors (Table 3), themselves

encompassing numerous variables. For instance, the factor "rainfall" was investigated via

variables such as "monthly rainfall", "cumulative rainfall", "total rainfall during the wet

season", "autumn rainfall", "onset of the rains". In total, we identified 408 variables whose

effect on the timing of birth was tested, whereas we identified 162 variables whose effect on

the synchrony of births was tested.

When investigating the timing of births, rainfall was the most common factor tested

(environmental characteristics theme, in n = 28 articles). Female age was the most

investigated factor among female characteristics (in n = 19 articles), while it was plant

phenology in plant characteristics (in n = 14 articles), offspring sex in offspring

characteristics (in n = 13 articles), and population density in population characteristics (in n =

12 articles, Supplementary Material 4). When investigating the synchrony of births, predation

exposure was the most common factor tested (in n = 16 articles). Latitude was the most

investigated factor among environmental characteristics (in n = 9 articles), while it was the

anti-predator behaviour of the juvenile in the life history traits (in n = 7 articles).

The effects of plant characteristics as well as environmental characteristics were

commonly explored in all continents (Fig. 6). In Africa, they were mainly explored via

rainfall and annual climatic seasonality (mostly based on a broad description of the temporal

succession of wet and dry seasons), while these themes were often described via local

environmental conditions (i.e., difference of seasonality and climate based on spatial

differences) in Europe, and plant phenology as well as temperature in North America. In

addition, the effects of female and population characteristics were also commonly explored in Europe and North America, mostly via female quality and identity as well as population density in Europe, and via female age and previous reproductive status in North America. The effects of predation exposure as well as offspring characteristics were also commonly explored in North America.

When considering the effect of a variable on the timing of births, 50% or more tests received partial or full support in all themes, except from the theme offspring characteristics, where more than 75% received no support (Fig. 7a). When considering the effect of a variable on the synchrony of births, life history traits, female characteristics and offspring characteristics (but n = 1 in the latter) received largely more than 50% support (Fig. 7b). The other themes received between 25 and 50% support overall (see details in Supplementary Material 5).

Discussion

The dominant hypotheses confirmed: seasonality and predation

Most of the articles that explored the potential effect of a climatic or a vegetation-related variable (environmental and plant characteristics) found support for the existence of an effect on timing, and to a lower extent, synchrony of births (e.g., Bunnell 1982, Sinclair et al. 2000, Moyes et al. 2011, Renaud et al. 2022). This conclusion is strengthened by the diversity of approaches identified in this literature review. Almost 20 factors were used (e.g., rainfall, temperature, greenness indexes, forage biomass, plant evapotranspiration). The effect of rainfall was mainly studied in Africa, and the one of temperature mainly in the northern hemisphere (Europe and North America), in accordance with the region-specific limiting factor of vegetation growth, the southern hemisphere seasonality being predominantly

determined by rainfall (wet versus dry seasons) and the northern hemisphere seasonality by

temperature (hot versus cold seasons, Owen-Smith and Ogutu 2013).

The articles retrieved in the present literature review explored the effect of predation

pressure predominantly via two approaches, the direct impact of predation on juvenile

mortality and the antipredator behaviour of the juvenile, and found moderate support (e.g.,

Michel et al. 2020). The articles using the antipredator behaviour of the juvenile were often

descriptive (but see Rutberg 1987 for a statistical approach), which can affect the strength of

the conclusions (e.g., Ogutu et al. 2013). Additionally, the antipredator behaviour of the

juvenile was generally classified according to the hider/follower dichotomy (Lent 1974).

However, this trait correlates with other life history traits such as group size and sociality,

diet and body size of the species, which might themselves affect birth phenology (Sinclair et

al. 2000). The classification of certain species at one end or the other of the gradient is

therefore debated (e.g., African buffalo (*Syncerus caffer*) and plains bison (*Bison bison*),

Green and Rothstein 1993b, Ryan et al. 2007), making it more difficult to establish general

and universal conclusions based on this approach. From the present literature, we conclude

that predation pressure might have a significant effect on the synchrony of births in African

species (effect supported in 63% of the cases). African ecosystems are less subject to food

resource and climate seasonality than, for instance, northern temperate ecosystems, in which

predation seems to have less effect than climate or vegetation phenology on the distribution

of births (effect partially or fully supported in only 42% of the cases). Such a finding

corroborates the results of previous comparative studies (Zerbe et al. 2012).

Emerging research theme avenues

Although fairly frequently explored in European and North American populations, the effects

of female, offspring and population characteristics remain marginally accounted for in the

study of the phenology of births in large herbivores. When investigated, evidence of their effect is mixed. It remains unclear if this lack of consistency in the results comes from biological reasons, these drivers potentially acting as a secondary driver after environmental conditions and mostly at the individual level might have a limited effect on the phenology of births, or from methodological reasons. The investigation of such drivers requires a detailed understanding in the long-term of the population studied and individually-centred monitoring, which can be challenging to implement and is thus infrequently done or might lead to imperfect datasets (e.g., data collected for a small sample of the population or biased toward certain individuals, data collected on a short period of time, Festa-Bianchet et al. 2017). Accordingly, we found that the articles focusing on European and North American populations (more particularly Cervidae populations with long-term monitoring) extensively exploring these drivers were also characterised by higher quality indexes than the other articles. Due to the difficulty to collect accurate data as well as their expected secondary role in the determination of the timing of births, these drivers are often included as control or secondary variables. Although these characteristics are often related to environmental conditions (e.g., the reproductive status of the female during the previous reproductive season can be related to forage availability and climatic conditions), intrinsic characteristics such as female age, social rank and experience (e.g., primiparous or multiparous) have also been identified as relevant drivers of the phenology of births (e.g., Berger and Cain 1999, Bonnet et al. 2019). Even if these variables do not actively influence the phenology of births, they nevertheless mechanically affect the distribution (e.g., in a population of mainly young females, one can expect the timing of births to be delayed), and failing to take these processes into account can affect conclusions when evolutionary or plastic drivers such as environmental seasonality are explored. Consequently, population structure should not be disregarded when studying the phenology of births.

Besides, despite directly affecting the phenology of birth, food resources and

environmental characteristics can also affect female's ability to breed and the timing of

breeding, and eventually the timing of parturition (Hogg et al. 2017). However, articles

dealing exclusively with the phenology of mating were not considered in this review (e.g.,

Dauphiné and McClure 1974). Gestation length is adjustable within some limits (Clements et

al. 2011, Zerbe et al. 2012), so the phenology of mating cannot be regarded as an absolute

reflection of the phenology of birth. Additionally, mating is more difficult to observe in the

field than birth and does not necessarily lead to a birth (i.e., abortion). We trust the number of

articles in this field is limited and their exclusion should not affect the general conclusions of

our study.

Eventually, the effect of diseases (e.g., Berger and Cain 1999) and inbreeding (e.g.,

Bonnet et al. 2019) prevalence on the phenology of births are currently under-explored.

Several pathogens can cause abortion in large herbivores (e.g., brucellosis (*Brucella abortus*),

Cross et al. 2015), thus leading to the possibility for an altered phenology of births in infected

populations, depending on the prevalence of the disease in the population and the timing of

abortion. Similarly under-explored, human interventions such as hunting (e.g., Gamelon et al.

2011) or active management (Kaze et al. 2016) might affect the evolutionary and/or plastic

determination of the synchrony of births in a population. Gamelon et al. (2011) showed that

hunting acts as a selection pressure for earlier dates of birth in a population of wild boar (*Sus*

scrofa), as being born earlier increases development time and allows females to reproduce

earlier, an evolutionary advantage when mortality is high. Kaze et al. (2016) hypothesised

that a better management of bison natural habitat via environmental rehabilitation and

reduction of competition with livestock for resource access enhanced female condition and

increased birth synchrony. In the current context of climate change, modification of land use

and development of human activities across the globe, there is a need for a better

understanding of the effect of such anthropogenic and epidemiological factors on the

phenology of births in large herbivores, and more broadly in mammals.

Research challenges

As illustrated in this study, the phenology of births is driven by a combination of drivers,

which might be difficult to separate and describe. Although most of the articles used "direct

tests", when the variable of interest is directly used in the test (e.g., effect of the age of the

female on the date of parturition, Bon et al. 1993), the use of "tests via proxy" was common

(when the variable of interest is not available or difficult to measure in the field leading to the

use of a proxy instead, e.g., effect of food availability via the variability of rainfall,

Ndibalema 2009). Additionally, some articles also used "logical tests", when the variable of

interest is not available or difficult to measure in the field, and a theoretical reasoning is used

to draw conclusions from the observations (e.g., "*if synchrony in births is primarily an*

antipredator adaptation by ungulates, then episodic and unpredictable droughts and floods

should not significantly shift the timing and synchrony of births in seasonally breeding

ungulates", Ogutu et al. 2010). However, in such cases, the effect of the driver tested cannot

be proven conclusively.

The importance of interacting and cascading effects is frequently highlighted in the

Discussion section or even tested in the articles retrieved for the present literature review

(e.g., Stopher et al. 2008, Zerbe et al. 2012, Aikens et al. 2021). The variability of our

findings in terms of the support received by each factor tested also illustrates this complexity.

To address this complexity, the use of aggregated indices (i.e., one index combining several

variables) such as latitude (which combines factors such as temperature, day length,

vegetation type, seasonality of the environment, e.g., English et al. 2012) and antipredator

behaviour of the juvenile (which refers to the level of precocity of the juvenile but also

correlates with other life history traits, e.g., Sinclair et al. 2000), as well as site-specific

comparisons (which uses spatial variability to describe different levels of exposure to

predation for example, but also includes variable local environmental conditions, e.g., Post et

al. 2003) is frequent. Such an approach can provide a general understanding of a set of factors

on the phenology of births, but limits our understanding of individual effects. Although the

possibilities are limited when studying populations in natural conditions (Clauss et al. 2021),

the study of the phenology of births in systems where one or several factors are constant or

fairly stable can help in this regard.

Most of the factors tested in the articles retrieved correspond to proximal factors

inducing a short-term response fairly easy to monitor. However, monitoring the effect of

ultimate factors generating slow evolutionary responses is more challenging. Such

investigations require comparative analyses involving multi-species comparisons controlling

for genetic relatedness (Zerbe et al. 2012, Bonnet et al. 2019). The majority of the articles

retrieved for the present literature review focused on a single species, and the rare multi-

species studies usually used descriptive comparisons between species based on life history

traits and/or did not account for phylogenetic correlations (e.g., Gosling 1969, Rutberg 1987).

Large scale comparative analyses can be of particular interest to empirically test theoretical

hypotheses about life history traits and evolutionary responses in reproductive phenology

(Rutberg 1987, Zerbe et al. 2012).

Geographic and taxonomic gap

Our literature review identified a large geographical and taxonomic gap in the study of the

phenology of births in large herbivores, likely to be generalized to other mammals too. More

than 90% of the articles retrieved concern Africa, Europe and North America, leaving Asia

and Central and South America mostly unexplored. At a finer geographical scale, West and

Central Africa as well as Eastern Europe also have almost no data available, or are concerned

by purely descriptive studies (e.g., in Honduras, Klein 1982; in Nepal, Bhusal et al. 2020).

Most of these ecosystems are characterised by dense vegetation, difficult conditions and/or

remote and secluded wilderness, explaining the overall gap in the scientific literature (Kelt

and Meserve 2014). Additionally, the large herbivores occupying these environments are

often elusive, mostly nocturnal and/or occurring at low densities (e.g., Cephalophus and

Philantomba genera in Central Africa, Zerbe et al. 2012), which makes their study

challenging, as illustrated by the lower quality index of these papers.

Moreover, although seasonals in terms of food availability, the absence of major

environmental seasonality in equatorial ecosystems (e.g., photoperiod, temperature) might

also have played a role in the lack of studies conducted under these latitudes (Bronson 2009),

and led to the assumption that no reproductive seasonality should be expected, thus reducing

the interest for the exploration of the phenology of births. However, there is evidence for

reproductive seasonality around the equator, as illustrated in other taxa such as Chiroptera

(e.g., Heideman and Utzurrum 2003) and in rare examples in the present literature review

(e.g., the phenology of births of the collared peccary (*Pecari tajacu*) is correlated with

fruiting phenology in French Guiana, Dubost and Henry 2017). Additionally, the

investigation of the phenology of births in species under stable environmental conditions

(compared to temperate systems characterised by dramatic seasonal changes) opens the door

to the exploration of other drivers such as predator pressure and social characteristics (e.g., in

this study, Sinclair et al. 2000; in Chiroptera, Porter and Wilkinson 2001) by providing

natural experimental conditions where the dominant driver regulating the phenology of births

in large herbivores, namely environmental seasonality, is controlled. Such a framework can

help better understand the role of secondary drivers when environmental seasonality is not a

strong constraint (Zerbe et al. 2012). Eventually, even in the absence of reproductive

seasonality, investigating the phenology of births is critical for the understanding of its

cascading effects on population dynamics, in particular in the context of climate change (e.g.,

the increase of winter births due to earlier onset of plant growth is negatively correlated with

juvenile recruitment in feral cattle (*Bos taurus*), Burthe et al. 2011).

Data availability statement

Data available from the Zenodo Repository: https://doi.org/10.5281/zenodo.17483679

(Anonymised reference).

Acknowledgments

We thank Elise Huchard for her suggestions during the early-stage conceptualization of the project and Hermanus Swanepoel for his help in data collection. This work was supported by a grant from the "Ministère Français de l'Enseignement Supérieur, de la Recherche et de l'Innovation" through the "Ecole Doctorale E2M2" of the "Université Claude Bernard Lyon 1" as well as Postgraduate Research Scholarship funding from Nelson Mandela University.

Conflict of interest statement

The authors declare that they have no conflict of interest.

Authors' contributions

Lucie Thel: Conceptualization - Methodology - Formal analysis - Investigation - Data Curation - Writing: Original Draft - Writing: Review & Editing - Visualisation - Funding acquisition - Validation

Christophe Bonenfant: Conceptualization - Methodology - Writing: Review & Editing - Supervision - Funding acquisition - Validation

Simon Chamaillé-Jammes: Conceptualization - Methodology - Writing: Review & Editing - Supervision - Funding acquisition - Validation

References

References

Anonymised reference (dataset access, for revision process)

Adams L. G. and Dale B. W. 1998. "Timing and synchrony of parturition in Alaskan

caribou." *Journal of Mammalogy* 79: 287-294.

Ahrestani F. S., Van Langevelde F., Heitkönig I. M. and Prins H. H. 2011.

"Contrasting timing of parturition of chital *Axis axis* and gaur *Bos gaurus* in tropical South

India–the role of body mass and seasonal forage quality." *Oikos* 121: 1300-1310.

Aikens E. O., Dwinnell S. P., LaSharr T. N., Jakopak R. P., Fralick G. L., Randall J.,

Kaiser R., Thonhoff M., Kauffman M. J. and Monteith K. L. 2021. "Migration distance and

maternal resource allocation determine timing of birth in a large herbivore." *Ecology* 102:

e03334.

Ansell W. 1960. "The breeding of some larger mammals in Northern Rhodesia."

Proceedings of the Zoological Society of London 134: 251-274.

Apollonio M., Merli E., Chirichella R., Pokorny B., Alagić A., Flajšman K. and

Stephens P. A. 2020. "Capital-income breeding in male ungulates: Causes and consequences

of strategy differences among species." *Frontiers in Ecology and Evolution* 8: 521767.

Berger J. 1992. "Facilitation of reproductive synchrony by gestation adjustment in

gregarious mammals: a new hypothesis." *Ecology* 73: 323-329.

Berger J. and Cain S. L. 1999. "Reproductive synchrony in brucellosis-exposed bison

in the southern Greater Yellowstone Ecosystem and in noninfected populations."

Conservation Biology 13: 357-366.

Bertram B. C. 1975. "Social factors influencing reproduction in wild lions." *Journal*

of Zoology 177: 463-482.

Bhusal A., Neupane B., Bhattarai S., Kapali A., Bhatta S., Bajagain S. 2020.

"Breeding seasonality of Chital (*Axis axis*) in the Hetauda Valley of Nepal." *Forestry Journal*

of Institute of Forestry 17: 174-183.

Bon R., Dardaillon M. and Estevez I. 1993. "Mating and lambing periods as related to

age of female mouflon." *Journal of Mammalogy* 74: 752-757.

Bonenfant C., Gaillard J.-M., Coulson T., Festa-Bianchet M., Loison A., Garel M.,

Loe L. E., Blanchard P., Pettorelli N., Owen-Smith N., Du Toit J. and Duncan P. 2009.

"Empirical evidence of density-dependence in populations of large herbivores." *Advances in*

Ecological Research 41: 313-357.

Boness D. J., Bowen W. D. and Iverson S. J. 1995. "Does male harassment of females

contribute to reproductive synchrony in the grey seal by affecting maternal performance?"

Behavioral Ecology and Sociobiology 36: 1-10.

Bonnet T., Morrissey M. B., Morris A., Morris S., Clutton-Brock T. H., Pemberton J.

M. and Kruuk L. E. 2019. "The role of selection and evolution in changing parturition date in

a red deer population." *PLoS biology* 17: e3000493.

Bowyer R. T., McCullough D. R., Rachlow J. L., Ciuti S. and Whiting J. C. 2020.

"Evolution of ungulate mating systems: integrating social and environmental factors."

Ecology and Evolution 10: 5160-5178.

Brockman D. K. and van Schaik C. P. 2005. "Seasonality and reproductive function".

In *Seasonality in primates: Studies of living and extinct human and non-human primates*,

Cambridge University Press.

Bronson F. 2009. "Climate change and seasonal reproduction in mammals."

Philosophical Transactions of the Royal Society B: Biological Sciences 364: 3331-3340.

Bronson F. H. 1989. *Mammalian reproductive biology*. University of Chicago Press.

Bunnell F. L. 1980. "Factors controlling lambing period of Dall's sheep." *Canadian*

Journal of Zoology 58: 1027-1031.

Bunnell F. L. 1982. "The lambing period of mountain sheep: synthesis, hypotheses,

and tests." *Canadian Journal of Zoology* 60: 1-14.

Burthe S., Butler A., Searle K. R., Hall S. J., Thackeray S. J. and Wanless S. 2011.

"Demographic consequences of increased winter births in a large aseasonally breeding

mammal (*Bos taurus*) in response to climate change." *Journal of Animal Ecology* 80: 1134-

1144.

Buzzard P. J., Li X. and Bleisch W. V. 2018. "High altitude ungulate communities in

southwest China." *Mammalia* 82: 415-422.

Clauss M., Zerbe P., Bingaman Lackey L., Codron D., Müller D. W. H. 2021. "Basic

considerations on seasonal breeding in mammals including their testing by comparing natural

habitats and zoos." *Mammalian Biology* 101: 373-386.

Clements M. N., Clutton-Brock T. H., Albon S. D., Pemberton J. M. and Kruuk L. E.

2011. "Gestation length variation in a wild ungulate." *Functional Ecology* 25: 691-703.

Côté S. D. and Festa-Bianchet M. 2001. "Birthdate, mass and survival in mountain

goat kids: effects of maternal characteristics and forage quality." *Oecologia* 127: 230-238.

Cross P. C., Maichak E. J., Rogerson J. D., Irvine K. M., Jones J. D., Heisey D. M.,

Edwards W. H. and Scurlock B. M. 2015. "Estimating the phenology of elk brucellosis

transmission with hierarchical models of cause-specific and baseline hazards." *The Journal of*

Wildlife Management 79: 739-748.

Danell K., Bergström R., Duncan P. and Pastor J. 2006. *Large herbivore ecology,*

ecosystem dynamics and conservation. Cambridge University Press.

Dasmann R. F. and Mossman A. S. 1962. "Reproduction in some ungulates in

Southern Rhodesia." *Journal of Mammalogy* 43: 533-537.

Dauphiné Jr T. C. and McClure R. L. 1974. "Synchronous mating in Canadian barren-

ground caribou." *The Journal of Wildlife Management* 38: 54-66.

Dezeure J., Baniel A., Carter A., Cowlishaw G., Godelle B. and Huchard E. 2021.

"Birth timing generates reproductive trade-offs in a non-seasonal breeding primate."

Proceedings of the Royal Society B 288: 20210286.

Dubost G. and Henry O. 2017. "Seasonal reproduction in neotropical rainforest

mammals." *Zoological studies* 56: e2.

English A. K., Chauvenet A. L., Safi K. and Pettorelli N. 2012. "Reassessing the

determinants of breeding synchrony in ungulates." *PLoS ONE* 7: e41444.

Estes R. D. 1976. "The significance of breeding synchrony in the wildebeest." *African

Journal of Ecology* 14: 135-152.

Festa-Bianchet M., Douhard M., Gaillard J.-M. and Pelletier F. 2017. "Successes and

challenges of long-term field studies of marked ungulates." *Journal of Mammalogy* 98: 612-

620.

Gaillard J.-M., Delorme D., Jullien J. and Tatin D. 1993. "Timing and synchrony of

births in roe deer." *Journal of Mammalogy* 74: 738-744.

Gamelon M., Besnard A., Gaillard J.-M., Servanty S., Baubet E., Brandt S. and

Gimenez O. 2011. "High hunting pressure selects for earlier birth date: wild boar as a case

study." *Evolution* 65: 3100-3112.

Giman B., Stuebing R., Megum N., Mcshea W. J. and Stewart C. M. 2007. "A camera

trapping inventory for mammals in a mixed use planted forest in Sarawak." *Raffles Bulletin of

Zoology* 55: 209-2015.

Gosling L. 1969. "Parturition and related behaviour in Coke's hartebeest, *Alcelaphus

buselaphus cokei* Günther." *Journal of Reproduction and Fertility Supplement* 6: 265-286.

Green W. C. and Rothstein A. 1993b. "Asynchronous parturition in bison:

implications for the hider-follower dichotomy." *Journal of Mammalogy* 74: 920-925.

Green W. C. and Rothstein A. 1993a. "Persistent influences of birth date on

dominance, growth and reproductive success in bison." *Journal of Zoology* 230: 177-186.

Heideman P. D. and Utzurrum R. C. 2003. "Seasonality and synchrony of

reproduction in three species of nectarivorous Philippines bats." *BMC Ecology* 3: 11.

Heldstab S. A., Müller D. W. H., Graber S. M., Bingaman Lackey L., Rensch E., Hatt

J.-M., Zerbe P. and Clauss M. 2018. "Geographical Origin, Delayed Implantation, and

Induced Ovulation Explain Reproductive Seasonality in the Carnivora." *Journal of Biological*

Rhythms 33: 402-419.

Hofmann R. R. 1989. "Evolutionary steps of ecophysiological adaptation and

diversification of ruminants: a comparative view of their digestive system." *Oecologia* 78:

443-457.

Hogg J. T., Dunn S. J., Poissant J., Pelletier F. and Byers J. A. 2017. "Capital vs.

income-dependent optimal birth date in two North American ungulates." *Ecosphere* 8:

e01766.

Ims R. A. 1990b. "On the adaptive value of reproductive synchrony as a predator-

swamping strategy." *The American Naturalist* 136: 485-498.

Ims R. A. 1990a. "The ecology and evolution of reproductive synchrony." *Trends in*

Ecology & Evolution 5: 135-140.

Kaze J., Whiting J. C., Freeman E. D., Bates S. B. and Larsen R. T. 2016. "Birth-site

selection and timing of births in American bison: effects of habitat and proximity to

anthropogenic features." *Wildlife Research* 43: 418-428.

Kelt D. A. and Meserve P. L. 2014. "Status and challenges for conservation of small

mammal assemblages in South America." *Biological Reviews* 89: 705-722.

Klein E. H. 1982. "Phenology of breeding and antler growth in white-tailed deer in

Honduras." *The Journal of Wildlife Management* 46: 826-829.

Laforge M. P., Webber Q. M. and Vander Wal E. 2023. "Plasticity and repeatability

in spring migration and parturition dates with implications for annual reproductive success."

Journal of Animal Ecology 92: 1042-1054.

Lent P. C. 1974. "Mother-infant relationships in ungulates." In *The Behaviour of*

Ungulates and its relation to management, 14-55. International Union for Conservation of

Nature and Natural Resources.

Leuthold W. and Leuthold B. 1975. "Temporal patterns of reproduction in ungulates

of Tsavo East National Park." *East African Wildlife Journal* 13: 159-169.

Mattisson J., Linnell J. D. C., Anders O., Belotti E., Breitenmoser-Würsten C., Bufka

L., Fuxjäger C., Heurich M., Ivanov G., Jędrzejewski W., Kont R., Kowalczyk R., Krofel M.,

Melovski D., Mengüllüoğlu D., Middelhoff T. L., Molinari-Jobin A., Odden J., Ozoliņš J.,

Okarma H., Persson J., Schmidt K., Vogt K., Zimmermann F. and Andrén H. 2022. "Timing

and synchrony of birth in Eurasian lynx across Europe." *Ecology and Evolution* 12: e9147.

McNutt J., Groom R. and Woodroffe R. 2019. "Ambient temperature provides an

adaptive explanation for seasonal reproduction in a tropical mammal." *Journal of Zoology*

309: 153-160.

Michel E. S., Strickland B. K., Demarais S., Belant J. L., Kautz T. M., Duquette J. F.,

Beyer Jr D. E., Chamberlain M. J., Miller K. V., Shuman R. M., Kilgo J. C., Diefenbach D.

R., Wallingford B. D., Vreeland J. K., Ditchkoff S. S., DePerno C. S., Moorman C. E.,

Chitwood M. C. and Lashley M. A. 2020. "Relative reproductive phenology and synchrony

affect neonate survival in a nonprecocial ungulate." *Functional Ecology* 34: 2536-2547.

Moe S., Rutina L. and Du Toit J. 2007. "Trade-off between resource seasonality and

predation risk explains reproductive chronology in impala." *Journal of Zoology* 273: 237-

243.

Moyes K., Nussey D. H., Clements M. N., Guinness F. E., Morris A., Morris S.,

Pemberton J. M., Kruuk L. E. and Clutton-Brock T. H. 2011. "Advancing breeding

phenology in response to environmental change in a wild red deer population." *Global*

Change Biology 17: 2455-2469.

Ndibalema V. G. 2009. "A comparison of sex ratio, birth periods and calf survival

among Serengeti wildebeest sub-populations, Tanzania." *African Journal of Ecology* 47: 574-

582.

Neumann W., Singh N. J., Stenbacka F., Malmsten J., Wallin K., Ball J. P. and

Ericsson G. 2020. "Divergence in parturition timing and vegetation onset in a large herbivore

—differences along a latitudinal gradient." *Biology Letters* 16: 20200044.

O'Donoghue M. and Boutin S. 1995. "Does reproductive synchrony affect juvenile

survival rates of northern mammals?" *Oikos* 74: 115-121.

Ogutu J. O., Piepho H.-P. and Dublin H. T. 2013. "Responses of phenology,

synchrony and fecundity of breeding by African ungulates to interannual variation in

rainfall." *Wildlife Research* 40: 698-717.

Ogutu J. O., Piepho H.-P., Dublin H. T., Bhola N. and Reid R. S. 2010. "Rainfall

extremes explain interannual shifts in timing and synchrony of calving in topi and warthog."

Population Ecology 52: 89-102.

Owen-Smith N. and Ogutu J. O. 2013. "Controls over reproductive phenology among

ungulates: allometry and tropical-temperate contrasts." *Ecography* 36: 256-263.

Page M. J., Moher D., Bossuyt P. M., Boutron I., Hoffmann T. C., Mulrow C. D.,

Shamseer L., Tetzlaff J. M., Akl E. A., Brennan S. E., Chou R., Glanville J., Grimshaw J. M.,

Hróbjartsson A., Lalu M. M., Li T., Loder E. W., Mayo-Wilson E., McDonald S.,

McGuinness L. A., Stewart L. A., Thomas J., Tricco A. C., Welch V. A., Whiting P. and

McKenzie J. E. 2021. "PRISMA 2020 explanation and elaboration: updated guidance and

exemplars for reporting systematic reviews." *BMJ* 372: n160.

Paoli A., Weladji R. B., Holand Ø. and Kumpula J. 2018. "Winter and spring climatic

conditions influence timing and synchrony of calving in reindeer." *PLoS ONE* 13: e0195603.

Paoli A., Weladji R. B., Holand Ø. and Kumpula J. 2020. "Early-life conditions

determine the between-individual heterogeneity in plasticity of calving date in reindeer."

Journal of Animal Ecology 89: 370-383.

Plard F., Gaillard J.-M., Coulson T., Hewison A. M., Delorme D., Warnant C. and

Bonenfant C. 2014. "Mismatch between birth date and vegetation phenology slows the

demography of roe deer." *PLoS Biology* 12: e1001828.

Porter T. and Wilkinson G. 2001. "Birth synchrony in greater spear-nosed bats

(*Phyllostomus hastatus*)." *Journal of Zoology* 253: 383-390.

Post E. 2003. "Timing of reproduction in large mammals". In *Phenology: an

integrative environmental science*, Springer.

Post E., Bøving P. S., Pedersen C. and MacArthur M. A. 2003. "Synchrony between

caribou calving and plant phenology in depredated and non-depredated populations."

Canadian Journal of Zoology 81: 1709-1714.

Radchuk V., Reed T., Teplitsky C., van de Pol M., Charmantier A., Hassall C., ... and

Kramer-Schadt S. 2019. "Adaptive responses of animals to climate change are most likely

insufficient." *Nature communications* 10: 3109.

R Core Team. 2025. "R: A language and environment for statistical computing". R

Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Renaud L. A., Festa-Bianchet M. and Pelletier F. 2022. "Testing the match–mismatch

hypothesis in bighorn sheep in the context of climate change." *Global Change Biology* 28:

21-32.

Ripple W. J., Painter L. E., Beschta R. L. and Gates C. 2010. "Wolves, elk, bison, and

secondary trophic cascades in Yellowstone National Park." *Open Ecology Journal* 3: 31-37.

Rutberg A. T. 1987. "Adaptive hypotheses of birth synchrony in ruminants: an

interspecific test." *The American Naturalist* 130: 692-710.

Ryan S., Knechtel C. and Getz W. 2007. "Ecological cues, gestation length, and birth

timing in African buffalo (*Syncerus caffer*)." *Behavioral Ecology* 18: 635-644.

Say-Sallaz E., Chamaillé-Jammes S., Fritz H. and Valeix M. 2019. "Non-consumptive

effects of predation in large terrestrial mammals: Mapping our knowledge and revealing the

tip of the iceberg." *Biological Conservation* 235: 36-52.

Shi J., Beauchamp G. and Dunbar R. I. 2010. "Group-size effect on vigilance and

foraging in a predator-free population of feral goats (*Capra hircus*) on the Isle of Rum, NW

Scotland." *Ethology* 116: 329-337.

Sinclair A., Mduma S. A. and Arcese P. 2000. "What determines phenology and

synchrony of ungulate breeding in Serengeti?" *Ecology* 81: 2100-2111.

Sinclair A. R. E. and Norton-Griffiths M. 1995. *Serengeti: dynamics of an ecosystem*.

University of Chicago Press.

Singh N. J., Börger L., Dettki H., Bunnefeld N. and Ericsson G. 2012. "From

migration to nomadism: movement variability in a northern ungulate across its latitudinal

range." *Ecological Applications* 22: 2007-2020.

Spinage C. 1973. "The role of photoperiodism in the seasonal breeding of tropical

African ungulates." *Mammal Review* 3: 71-83.

759   Stopher K. V., Pemberton J. M., Clutton-Brock T. H. and Coulson T. 2008.

760 "Individual differences, density dependence and offspring birth traits in a population of red

761 deer." *Proceedings of the Royal Society B: Biological Sciences* 275: 2137-2145.

762   Szemán K., Liker A. and Székely T. 2021. "Social organization in ungulates:

763 revisiting Jarman's hypotheses." *Journal of Evolutionary Biology* 34: 604-613.

764   Thel L., Chamaillé-Jammes S. and Bonenfant C. 2022. "How to describe and measure

765 phenology? An investigation on the diversity of metrics using phenology of births in large

766 herbivores." *Oikos*: e08917.

767   Williams C. T., Lane J. E., Humphries M. M., McAdam A. G. and Boutin S. 2013.

768 "Reproductive phenology of a food-hoarding mast-seed consumer: resource-and density-

769 dependent benefits of early breeding in red squirrels." *Oecologia* 174: 777-788.

770   Zerbe P., Clauss M., Codron D., Bingaman Lackey L., Rensch E., Streich J. W., Hatt

771 J.-M. and Müller D. W. 2012. "Reproductive seasonality in captive wild ruminants:

772 implications for biogeographical adaptation, photoperiodic control, and life history."

773 *Biological Reviews* 87: 965-990.

Figure legends

Figure 1: number of articles investigating the drivers of the phenology of birth in large herbivores (n = 124): a) published every year between 1960 and 2024, b) cumulated number of articles published every year (dots: cumulated numbers, line: predictions of the quadratic regression, shaded area: 95% confidence interval).

Figure 2: Species studied in the articles investigating the drivers of the phenology of birth in large herbivores (n = 82). The size of the dots is proportional to the number of articles studying a given species. The colour of the branches corresponds to taxonomic families: Antilocapridae, Bovidae, Camelidae, Cervidae, Elephantidae, Equidae, Giraffidae, Hippopotamidae, Moschidae, Rhinocerotidae, Suidae, Tayassuidae.

Figure 3: Duration of the studies presented in the articles investigating the drivers of the phenology of births in large herbivores between 1960 and 2024 (n = 170 populations).

Figure 4: Distribution across countries of the total number of articles investigating the drivers of the phenology of births in large herbivores (n = 124). Some articles cover several countries.

Figure 5: Data collection methods used in the articles investigating the drivers of the phenology of births in large herbivores between 1960 and 2024 (n = 170 populations).

Figure 6: Frequency of investigation of each theme (based on the number of articles exploring at least once a factor of the theme considered) in the articles investigating the drivers of the phenology of births in large herbivores between 1960 and 2024: environmental

characteristics; plant characteristics; life history traits; female characteristics; offspring

characteristics; population characteristics; predation exposure.

Figure 7: Proportion of tests receiving support (green), partial support (orange) or no support

(red) in the articles investigating the drivers of the phenology of birth in large herbivores: a)

timing and b) synchrony, for each theme (n = 7; environmental characteristics; plant

characteristics; life history traits; female characteristics; offspring characteristics; population

characteristics; predation exposure).

Tables

Table 1: Number of articles per continent investigating the effect of at least one variable on

the timing or on the synchrony of births in large herbivores (the same article can appear in

both). The "worldwide" category represents articles studying populations from more than one

continent.

Continent	Number of articles exploring the drivers of the timing of births	Number of articles exploring the drivers of the synchrony of births	Total number of articles
Africa	35	9	36
Asia	4	3	5
Central and South America	4	0	4
Europe	27	8	30
North America	39	24	46
Worldwide	1	3	3

813 Table 2: Number of articles investigating at least once the effect of a factor belonging to one

814 of the seven themes identified in the study on the timing or on the synchrony of births in large

815 herbivores (n = 124).

Theme	Timing	Synchrony
Environmental characteristics	79	23
Plant characteristics	33	10
Female characteristics	32	3
Offspring characteristics	18	1
Population characteristics	18	6
Life history traits	9	9
Predation exposure	6	16

817　Table 3: Factors (n = 35) tested for their effect on the timing and/or on the synchrony of

818　births in large herbivores, and their corresponding theme (n = 7) as identified in the study.

Theme	Factor
Environmental characteristics	altitude, annual climatic seasonality, cloud, day length, inter-annual climatic variability, latitude, local environmental conditions, rainfall, snow, temperature
Plant characteristics	faecal or forage protein content, forage biomass, forage quality, general forage quality, general plant phenology, NDVI, plant evapotranspiration
Life history traits	anti-predator behaviour of the juvenile, body size, diet, gregariousness, heritability and plasticity, phylogeny
Female characteristics	female age, female body mass, female condition, female identity, female migratory strategies, female previous reproductive status, female quality, female social rank
Offspring characteristics	offspring inbreeding level, offspring number, offspring sex
Population characteristics	population age structure, population density, population sanitary state, population sex-ratio
Predation exposure	predation exposure

Figures

Figure 1

Figure 2

Figure 4

Figure 6

Figure 7

Supplementary Material

Supplementary Material 1: Data collection process for the systematic review

Supplementary Material 2: References of the articles included in the systematic review (n = 124)

Supplementary Material 3: Quality index according to time, continent and taxonomic family

Supplementary Material 4: Distribution of factors per theme

Supplementary Material 5: Number of times each factor tested received support, partial support and no support

Graphical abstract legend

This systematic review confirmed the strong support received by the two dominant drivers of births phenology, the seasonality and predation hypotheses. Even though there is evidence of their importance, the effects of female, offspring and population characteristics remain marginally accounted for. Asian and South/Central American species are currently understudied.

 Graphical abstract