Differential phoretic vector use among sympatric *Caenorhabditis* nematodes
and an association with invasive nitidulid beetles in southwestern Germany

Ryan Greenway[1]*, Loel Dalan[1], Christian Braendle[2], Marie-Anne Félix[3], and Siyu
Serena Ding[1,4]*

[1]Research Group Genes and Behavior, Max Planck Institute of Animal Behavior,
78464 Konstanz, Germany
[2]Université Côte d'Azur, CNRS, Inserm, IBV, 06108 Nice, France
[3]Institut de Biologie de l'École Normale Supérieure, Centre National de la Recherche
Scientifique, 75005 Paris, France
[4]Centre for the Advanced Study of Collective Behaviour, 78464 Konstanz, Germany
*Corresponding author. Correspondences to:
rgreenway@ab.mpg.de
serena.ding@ab.mpg.de

Abstract

Little is known about the natural history of many *Caenorhabditis* nematodes, despite
their relationship to the model species *C. elegans*. While these nematodes rely on
invertebrate vectors to disperse to new habitats (phoresy), vector use for most species
has not been characterized. We surveyed the invertebrate community of a habitat
containing three sympatric *Caenorhabditis* in southwestern Germany, determining
differential and specific vector use for each of these co-occurring species. We
documented a specific association between *C. apta* n. sp. and two species of invasive
nitidulid beetles, and a particularly strong association of the nematode with the beetle
Stelidota geminata, where we recovered more nematodes per individual beetle and a
higher proportion of beetles carrying nematodes compared to the co-occurring
Epuraea ocularis. Our results provide evidence for group dispersal using beetles in *C.*
apta, supporting previous observations of collective dispersal behavior in this species,
and establish a starting point for further dissecting the evolutionary and mechanistic
causes and consequences of interactions between *Caenorhabditis* nematodes and
their vector species in ecologically relevant conditions.

Keywords

Dauer, invasive species, interspecific interactions, coexistence, Coleoptera

Introduction

Free-living nematodes are some of the most abundant and diverse animals on Earth, and have reached their ubiquitous status despite their small size and limited modes of locomotion (Schratzberger *et al.* 2019). Many free-living nematodes achieve relatively wide distributions for their body size as the result of phoresy, hitchhiking to new habitats by temporarily attaching to a larger and more mobile interspecific vector (Bartlow & Agosta 2021). Phoresy in nematodes is often facilitated by an ability to enter diapaused or quiescent states, during which they can endure starvation and resist environmental extremes over extended time periods, allowing them to safely attach to vectors and disperse over greater distances than possible by crawling alone (Vlaar *et al.* 2021). The specificity of the relationship between phoretic nematodes and vector species varies widely across taxa, and is likely an important driver of nematode diversification and ecological success (Giblin-Davis *et al.* 2013; Bubrig & Fierst 2021; Vlaar *et al.* 2021).

Caenorhabditis nematodes are a key resource used to fuel a wide range of biological discovery over the last century, particularly the laboratory model organism *C. elegans* (Corsi *et al.* 2015; Meneely *et al.* 2019; Ambros *et al.* 2025). Despite their importance, the study of ecology and evolution in natural *Caenorhabditis* populations has only recently received significant attention thanks to the discovery of the natural habitat of these nematodes in the 2010s (Kiontke *et al.* 2011; Cutter 2015; Frézal & Félix 2015). We now know that in the wild *Caenorhabditis* are microbivorous nematodes with a boom-bust life cycle on ephemeral, microbe-rich resource patches, mostly decaying plant material (Kiontke *et al.* 2011; Cutter 2015). In the presence of stressors (*e.g.* starvation, high population density), *Caenorhabditis* nematodes develop a stress-resistant dauer larval form that facilitates dispersal and colonization of new patches after the rapid exhaustion of microbial resources (Barrière & Félix 2007; Félix & Duveau 2012; Schulenburg & Félix 2017; Vlaar *et al.* 2021). Experiments have suggested that *Caenorhabditis* appear to be reliant on phoretic associations with invertebrate carriers to colonize new resource patches (Li *et al.* 2014; Petersen *et al.* 2015; Sloat *et al.* 2022). Although phoretic dispersal is a key component of the life cycle and ecology of *Caenorhabditis*, interactions with potential dispersal vectors remain unknown for the vast majority of *Caenorhabditis* species.

A variety of invertebrate species have been documented carrying *Caenorhabditis* nematodes, with some species found in specialized associations with particular invertebrate taxa and others appearing to be opportunistic or generalists. The best characterized interaction between *Caenorhabditis* and vector is that of *C. japonica* with the shield bug *Parastrachia japonensis,* where the nematode is found exclusively in association with this insect throughout its life-cycle, and has not been found on other invertebrates despite examination of numerous co-occurring taxa (Yoshiga *et al.* 2013). Though not as well characterized as the relationship between *C. japonica* and its vector, *C. niphades* and *C. inopinata* also have specific relationships with particular insects (*Niphades variegatus* weevils and *Ceratosolen spp.* fig pollinating wasps, respectively; (Woodruff & Phillips 2018; Sun *et al.* 2022). Other *Caenorhabditis* may have similarly specific relationships with vector species or taxonomic groups, having been originally isolated and described in association with particular insects (Figure 1), such as *C. drosophilae* with the fruit fly *Drosophila nigrospiracula* (Kiontke 1997), *C. angaria* with the weevils *Metamasius hemipterus* and *Rhynchophorus palmarum* (Sudhaus *et al.* 2011), *C. plicata* with carrion beetles (Volk 1950)*, C. auriculariae* with fungus-feeding *Platydema spp.* beetles (Dayi *et al.* 2021), and *C. monodelphis* with the ciid beetle *Cis castaneus* (Slos *et al.* 2017). However, for these species there does not appear to have been systematic examination of other co-occurring organisms that could also act as vectors.

In contrast to the specific interactions mentioned above, several *Caenorhabditis* species in the Elegans species group (Figure 1) have regularly been found in association with various species of terrestrial invertebrates, including *C. latens* with isopods (Dey *et al.* 2012)*, C brenneri* with gastropods (Diano *et al.* 2022)*, C. briggsae* (Barrière & Félix 2005; Félix & Duveau 2012; Félix *et al.* 2013; Petersen *et al.* 2015) and *C. sinica* (Wang *et al.* 2010) with isopods and gastropods, and *C. remanei* (Fitch *et al.* 1994; Baird 1999; Petersen *et al.* 2015) and *C. elegans* (Barrière & Félix 2005, 2007; Caswell-Chen *et al.* 2005; Félix & Duveau 2012; Ross *et al.* 2012; Petersen *et al.* 2015) with isopods, gastropods, and myriapods. Both *C. briggsae* and *C. elegans* have also been occasionally isolated from insects (Kiontke & Sudhaus 2006; Félix & Duveau 2012), and *C. elegans* dauers will attach to and disperse on various invertebrate groups, as well as inanimate materials, in laboratory experiments (Lee *et al.* 2011, 2017; Chiba *et al.* 2023; Williams *et al.* 2023; Perez *et al.* 2025).

Consequently, it has been hypothesized that these species may be largely
opportunistic in their use of phoretic vectors, exhibiting little specificity or preference
for different taxonomic groups (Schulenburg & Félix 2017; Petersen *et al.* 2023). Under
this scenario, it would be expected that opportunistic *Caenorhabditis* species could be
recovered from many potential vectors frequently encountered in and around their
habitats. Only one study has surveyed the broader invertebrate communities
alongside *Caenorhabditis*-carrying gastropods, isopods, and myriapods but did not
observe any additional invertebrate associations with *Caenorhabditis*, indicating there
may be some degree of vector specificity for these nematodes (Petersen *et al.* 2015).
Despite these described cases, vector use has not been characterized in any detail
for the majority of the >80 known *Caenorhabditis* species.

In southwestern Germany, three species of *Caenorhabditis* can be found co-occurring
at small spatial scales, even being found together on the same rotting fruits (Perez *et*
al. 2025): *C. briggsae*, *C. remanei*, and *C.* sp. 8 described here as *C. apta* n. sp. (see
Appendix 1), a little-studied species in the Angaria group of *Caenorhabditis* nematodes
(Figure 1). Repeated sampling of the orchard meadows around the University of
Konstanz campus has revealed that *C. apta* is the most abundant species in the area
across multiple years; however, it remains unknown what vector(s) *C. apta* uses for
dispersal. To determine whether these three *Caenorhabditis* species rely on the same
or different vectors for dispersal, we conducted a broad survey of invertebrates found
around rotting fruits. If any species were truly opportunistic in vector use, we would
expect to recover them from a broad range of invertebrate groups surveyed from their
habitat. Instead, we found that each *Caenorhabditis* species was isolated only from a
subset of the sampled invertebrate taxa. Based on the results from this general survey,
we conducted a more detailed survey of two locally abundant species of nitidulid
beetles that served as vectors for *C. apta*. We also assessed the life stage, number of
individuals, and sex ratio of *C. apta* on these two beetle species across multiple sites
and across a three-month period at a single site to further characterize interactions
between *C. apta* and its vectors.

Materials and Methods
General invertebrate collection and nematode extraction

To determine which invertebrate taxa serve as vectors for sympatric *C. briggsae, C.*
remanei, and *C. apta*, we opportunistically collected invertebrates found on and
around rotting fruits in orchard meadows near the University of Konstanz campus in
Baden-Württemberg, Germany from June 20, 2025 to October 24, 2025 (Table 1).
Gastropods and large arthropods were individually collected into Whirl-pak ® bags
(The Artemis Corporation, USA) or 2 mL microcentrifuge tubes, while smaller
arthropods (*e.g.* dipterans, coleopterans, hymenopterans) were collected using an
aspirator and deposited into 2 mL microcentrifuge tubes. During this general survey,
invertebrates were identified to order or higher taxonomic levels, with the exception of
nitidulid beetles. Due to their frequent occurrence relative to other taxa, we identified
nitidulid beetles to the species level based on keys for the beetle fauna of Europe
(Lompe 2002; Benisch 2023). Invertebrates were brought back to the lab within two
hours for processing. Invertebrates were individually added to either 6 or 9 cm petri
dishes containing nematode growth media (NGM) seeded with a 200 µL spot of *E. coli*
OP50. Petri dishes were sealed with Parafilm ® (Bemis Company, USA), left at 20º C
on the benchtop in a climate-controlled laboratory, and monitored for 48 hours to
determine if any nematodes emerged onto the bacterial lawn. Emerged nematodes
were identified as *Caenorhabditis* based on morphology using a dissecting microscope
(Stemi 508, Zeiss, Germany) fitted with 2.0x front optics. Following Barrière & Félix
(2006), we identified *Caenorhabditis* based on the presence of both middle and basal
pharyngeal bulbs, a clear grinder in the basal pharyngeal bulb, relatively light brown
and homogeneously colored gut, a relatively short rectum, centrally located vulva (in
females/hermaphrodites), a long and pointy tail in females/hermaphrodites, and a
blunt tail with a fan-shaped bursa in males. We were also able to identify some
additional nematode groups to the genus level, in particular *Pristionchus* based on the
presence of well-developed teeth, the lack of a grinder in the terminal bulb of the
pharynx, large and "dumpy" body shape, and clearly visible longitudinal striations in
the cuticle (Barrière & Félix 2006; Ragsdale *et al.* 2015). *Sheraphelenchus* were
distinguishable based on their diagnostic tail morphology, possessing a digitiform
spike with a finely pointed tip, particularly evident in males (Hunt 1993; Kanzaki &
Tanaka 2013). Lastly, a few individuals belonging to family Panagrolaimidae were able
to be identified based on their long cylindrical pharynx, terminal pharyngeal bulb, and
conical tail (Van Driessche *et al.* 2003). The identity of *Caenorhabditis* that emerged
over the 48-hour period was determined (see below) and recorded for each sample.

Identification of Caenorhabditis *species*

We identified *Caenorhabditis* to the species level based on morphology, behavior, and

reproductive mode using a Stemi 508 dissecting microscope (Zeiss, Germany) fitted

with 2.0x front optics. Previous sampling of this area confirmed the presence of *C.*

remanei, *C. briggsae*, and *C. apta* (Perez *et al.* 2025), which can be distinguished from

each other based on established diagnostic criteria visible under a dissecting

microscope, summarized in Appendix 2 (Barrière & Félix 2006; Kiontke *et al.* 2011;

Sloat *et al.* 2022). After disembarking the invertebrates, nematodes were allowed to

exit the dauer stage and resume development in order to facilitate identification. Upon

reaching sexual maturity, nematodes were examined under the dissecting microscope

and monitored for reproduction. We identified *C. briggsae* based on its self-fertilizing

hermaphroditic reproduction and an absence of males, in contrast to the gonochoristic

reproduction of the other two species. Of the remaining two species, we identified *C.*

apta by its relatively short and "dumpy" body shape, spiral mating behavior, a narrow

fan on the male tail, and stout, amber-colored male spicules. In contrast, *C. remanei*

was identified based on its longer and slimmer body shape, parallel mating behavior,

wide male tail fan, and less pronounced male spicules. In addition to these established

criteria, we also observed that *C. remanei* tends to form large aggregations on

bacterial lawns and readily burrow into the NGM compared to the other two species,

while *C. briggsae* forage individually and tend to remain on the agar surface in the

presence of food. Any uncertain species identifications were confirmed with mating

tests, where isolated males or offspring generated from isolated mothers were

individually transferred to a new 35 mm petri dish containing nematode NGM seeded

with a 100 µL spot of *E. coli* OP50 and the opposite sex of known species identity to

be assessed for reproduction over 72 hours, as each of these species is incapable of

interbreeding with the other two (Greenway *unpublished data*, (Ting *et al.* 2014). In the

case of putative hermaphrodites, virgin worms were plated individually and assessed

for reproduction over 72 hours. Likewise, in cases where only a single female

Caenorhabditis individual emerged from a given sample and did not reproduce 24

hours post-maturation, these individuals were transferred to new plates and provided

with adult males of either *C. remanei* or *C. apta* (pairing based on body shape of the

isolated individual) and monitored for reproduction over 72 hours to confirm species

identity.

Nitidulid beetle dissection and quantitative assessment of Caenorhabditis *occupancy*

Once we had identified beetles in the family Nitidulidae as the only invertebrates associated with *C. apta* (Table 1), we conducted more intense sampling of *Stelidota geminata* and *Epuraea ocularis*, two nitidulid species frequently observed and collected on rotting fruits in our prior sampling. To determine the species, life stage, number of individuals, sex ratio, and locations of nematodes on both beetle species, we performed dissections of adult *E. ocularis* and *S. geminata* collected from 6 locations in Baden-Württemberg, Germany (Table 2). Beetles were collected from rotting fruits using an aspirator, individually placed in 2 mL microcentrifuge tubes, and stored on ice for a minimum of one hour prior to sacrifice by removing the head from the thorax. The beetles were dissected under a stereomicroscope on a 6 cm petri dish containing NGM seeded with 200 µL *E. coli* OP50. We first examined the external surface of the beetle for the presence of nematodes before removing both elytra and transferring them together to a new NGM petri dish, leaving the rest of the corpse on the original dish. We then examined the internal surface of the elytra and the exposed abdomen separately, washing 20 µL of M9 buffer over each sample to loosen attached nematodes, at which point the life stage (larvae, dauer larvae, or adult) of visible nematodes was recorded. Petri dishes were sealed with Parafilm ® and left on the benchtop in a climate-controlled laboratory at 20º C. After 24 hours, the species, total number, and, for a subset of samples, the proportion of males present in the detached nematode population was separately recorded for the elytra and corpse of each beetle. The proportion of males on the beetle could not be determined in all cases, for example when some individuals remained as dauers for several days while others exited dauer rapidly and began to reproduce within 24 hours, resulting in mixed generation larval populations. We further monitored the corpse of each beetle for an additional 48 hours (72 hours total) to qualitatively determine the presence of additional nematode species that emerged following beetle death. To assess seasonal patterns of nematode occupancy and abundance on the beetles, we also collected each beetle species from a single location near the University of Konstanz (Egg Orchard; Figure 2A) two to three times per month from September 4, 2025 to November 10, 2025. Beetles were no longer found on or around fruits after the first freeze in mid-November.

Global distribution patterns of C. apta *and its nitidulid vectors*

Both nitidulid species we found carrying *C. apta* have invaded the European continent
and spread rapidly since the 2000s (Konzelmann 2001; Callot 2007; Jelínek *et al.*
2016; Bibin 2017; Stan 2019; Grujić *et al.* 2020; Lasoń *et al.* 2023), leading us to ask
whether this nematode may have been introduced to Europe alongside either of these
species. To determine whether there is evidence that *C. apta* originated within the
native range of either species, we examined the known distribution of *C. apta*, taking
into account collection dates and geographic context. Specifically, we assessed
whether *C. apta* had been collected in Europe, the Americas (the native range of *S.
geminata*), or Southeast Asia and the western Pacific (the native range of *E. ocularis*).
To this end, we compiled available sampling records from 1990–2024, including both
published and new data, for *C. apta* (N = 55 strains) and other members of the Angaria
clade (*C. quiockensis*, *C. castelli*, *C. angaria*, *C.* sp. 56, and *C. dolens*; N = 50 strains).
252
If *C. apta* was also recently introduced to Europe, we would expect to see collections
of *C. apta* from the native range of one of these beetles, as well as detections in
Europe starting in the mid to late 2000s, coinciding with the spread of the beetles.
Alternatively, if the nematode is native to Europe, we would expect to see earlier
detections of this species in Europe, prior to the spread of the beetles, but not
necessarily overlapping distributions in the native range of either beetle species.
Alternatively, *C. apta* may be a cosmopolitan species, in which case, we would expect
to see a widespread distribution and no temporal pattern to this species detection in
different geographic contexts. To provide a phylogeographic context to the distribution
of *C. apta*, we also compared the known distributions of its closest relatives, species
in the Angaria group. If many or all of the other species in the Angaria group are
concentrated in a particular geographic area, this could indicate that *C. apta* also
originates from the area. On the other hand, if the Angaria group species are
cosmopolitan in distribution, it may indicate a similar natural range for *C. apta*. We also
downloaded occurrence data for *S. geminata* and *E. ocularis* from GBIF (2025) to
compare the geographic and temporal distributions of these two species with collection
data for *C. apta*.
270
Statistics
We analyzed all data in R v4.3.2 (R Core Team 2023). To test whether the proportion
of individuals carrying *C. apta* differs across beetle species and sites, we used a

generalized linear model (GLM) as implemented in the stats package (R Core Team

2023), with presence/absence of *C. apta* as a binomial response variable and a logit

link function, including beetle species and site as factors. To determine whether the

number of extracted *C. apta* individuals differed between the sampled body parts of

the two beetle species across sites, we used a generalized linear mixed model

(GLMM) assuming a negative binomial response variable (nematode counts) and log

link function with the glmmTMB package (Brooks et al. 2017), using beetle species,

body part, and site as factors, and the interaction between species and body part. We

also included a random effect of the individual ID nested within site to account for non-

independence among body parts from the same individual. To test if the sex ratio of

C. apta individuals on beetles differed between species or sites, we modeled the

proportion of male nematodes to total nematodes per beetle using a binomial GLM

with a logit link function in the *stats* package, including beetle species and site as

factors. The interaction between species and site was non-significant for the GLMs

investigating differences in the number of beetles carrying *C. apta* and sex ratio

differences, so were excluded from the final models. For these among site analyses,

beetles collected from the Egg Orchard site were subsampled to better match the

collection dates and sample sizes from other sites, including only individuals collected

from September 15 to November 1, 2025. Additionally, the Niederhofen site was

excluded from all analyses due to the low number of beetles sampled from this

location.

We also used GLMs to test whether the number of individual beetles carrying *C. apta*

and the number of *C. apta* individuals extracted from the two beetle species differed

over time at the Egg Orchard site. We used a GLM modeling *C. apta*

presence/absence on each beetle as a binomial response variable and a logit link

function with the *stats* package, including beetle species as a factor and sampling date

as a number (binning samples collected within 5 days of each other). We used a GLM

with *C. apta* counts per beetle modeled as a negative binomial response with a log

link function in the glmmTMB package, including beetle species and sampling date as

factors. The non-significant interaction between species and date was excluded from

both models.

Results

General invertebrate survey for Caenorhabditis

We first surveyed a broad sample of invertebrates found on and around rotting fruits

near the University of Konstanz, noting the presence or absence of the three

Caenorhabditis species on each individual to determine which taxa serve as vectors

for *Caenorhabditis* nematodes in southern Baden-Württemberg. In total, 478 individual

invertebrates from various taxonomic groups were included in this general survey

(Table 1). Confirming the results of previous studies (Fitch *et al.* 1994; Baird 1999;

Dey *et al.* 2012; Félix & Duveau 2012; Félix *et al.* 2013; Petersen *et al.* 2015), *C.*

briggsae and *C. remanei* were both found on gastropods, and *C. remanei* on isopods.

C. briggsae was isolated from two slugs (3.1% of gastropods), while *C. remanei* was

isolated from four slugs (6.2% of gastropods) and 9 isopods (10.2% of isopods). The

two species were never found occurring on the same vector individual in this survey.

In contrast, *C. apta* was never found on gastropods or isopods, but instead was only

found in association with beetles in the family Nitidulidae, being isolated from 9 nitidulid

beetles (4.6% of all nitidulids) in the general survey. This, however, likely represents

an underestimation of the total occupancy rate of *C. apta* on nitidulids based on closer

inspection of these beetles (see detailed survey below) and laboratory observations.

We found *C. apta* occurring in several independent nitidulid breeding cultures started

from beetles that were deemed "nematode free" after initial screening on agar plates.

Several days after placing these beetles on sterilized apples, many of the cultures

were covered in *C. apta* nematodes, indicating that *C. apta* may be reluctant to

disembark vectors when presented only with *E. coli* OP50 as a food source. In light of

this, we conducted closer inspection of isopods and gastropods to confirm that these

taxa did not also carry *C. apta* or other *Caenorhabditis* that were reluctant to depart

onto the provided media. A subset of the isopods and gastropods from the general

survey (n = 34 and 15, respectively) were killed with a scalpel and washed in water,

plated on NGM petri dishes with the washing liquid, and observed for approximately

one week. We did not find any additional *Caenorhabditis* emerging from these

individuals, confirming the absence of *C. apta* on these taxa. Additionally, *C. briggsae*

and *C. remanei* were never found on nitidulid beetles, and no *Caenorhabditis* species

were found from any other invertebrate group in the survey (Table 1).

Quantitative assessment of C. apta *from nitidulid beetles*

As an association between *Caenorhabditis* and nitidulids has not been previously described, we further studied this relationship by conducting additional sampling of *E. ocularis* and *S. geminata* (Figure 2B and C), the two nitidulid beetles most often encountered in our study area, for quantitative characterization of *C. apta* from these beetles. We dissected an additional 177 *E. ocularis* and 197 *S. geminata* collected between September 15 and November 1, 2025 across six sites in Baden-Württemberg (Figure 2A; Table 2) to determine the lifestage, prevalence, and sex ratio of *C. apta* on individual beetles, as well as where on the beetles the nematodes could be found. We found that the abundance of *C. apta* significantly differed among body parts, beetle species, and sites. Inspection of both the elytra and remainder of the beetle corpse revealed that only dauer larvae were found on the beetles (Figure 2D). The number of *C. apta* observed was higher on the elytra than on the rest of the beetle body (β = 2.31 ± 0.33, z = 7.10, p < 0.001), with *S. geminata* having significantly higher abundances than *E. ocularis* on both the elytra and body (interaction β = 0.97 ± 0.40, z = 2.40, p = 0.016; species β = 1.14 ± 0.37, z = 3.07, p = 0.002; Table 2). Dauers were mostly found on the anterior portion of the elytra, stuck together into clusters in or near a cavity created by the epipleura (the portion of the elytron that folds back towards the abdomen (Goczał & Beutel 2023); Figure 2D and E). Attached dauers were largely motionless until prodded with a wire pick or wetted with M9 solution, at which point they began to wriggle and detach from each other. Despite concentration on the elytra, low numbers of *C. apta* dauers were also found having emerged from the rest of the beetle body in some cases (Table 2). The number of *C. apta* observed on beetles was significantly lower at Möggingen (β = -1.22 ± 0.34, z = -3.55, p < 0.001) and Tübingen (β = -0.99 ± 0.35, z = -2.84, p = 0.004) compared to the Egg Orchard site (Table 2).

We also examined the sex ratio of *C. apta* individuals emerged from each beetle (combined across body parts), finding that the proportion of males on each beetle was significantly lower from an expected 1:1 ratio (Huang *et al.* 2023) for *E. ocularis* (Intercept β = -0.68 ± 0.23, z = -2.93, p = 0.003) and was significantly different between species, closer to an even 1:1 for *S. geminata* (β = 0.53 ± 0.22, z = 2.46, p = 0.014; Table 2), but did not differ among sites (all z ≤ 1.59, all p > 0.110). Differences in the sex ratio of attached nematodes between the two species are likely driven by the low number of nematodes recovered per beetle for *E. ocularis* (Table 2).

In addition to carrying significantly more dauers per beetle, a higher proportion of *S.*
geminata individuals were carrying *C. apta* dauers than were *E. ocularis* (β = 1.57 ±
0.24, z = 6.59, p < 0.001), with *C. apta* found on 68.5% of *S. geminata* individuals
compared to 35.6% of *E. ocularis* across all sites (Table 2). However, there were also
differences in the proportion of all beetle individuals carrying *C. apta* dauers among
sites, with lower overall rates of nematode recovery in both Möggingen (β = -1.03 ±
0.33, z = -3.09, p = 0.002) and Tübingen (β = -1.20 ± 0.35, z = -3.40, p < 0.001)
compared to the Egg Orchard site (Table 2). As these sites were only sampled a single
time later in the season, it is impossible to distinguish if the lower dauer abundance
and proportion of beetle vectors are the result of site-specific or temporal variation in
this interspecific interaction.

Temporal variation in nematode-beetle interactions
We examined temporal variation in the number of beetles carrying *C. apta* and the
number of individual nematodes per beetle at a single site (Egg Orchard, Konstanz;
105 *E. ocularis*, 123 *S. geminata*) from September to November 2025 (Figure 3),
spanning the season when *C. apta* is abundant and active in late summer until its
disappearance in mid-autumn. Dauers of *C. apta* were found on a significantly higher
number of *S. geminata* than *E. ocularis* over the entire period (β = 1.92 ± 0.31, z =6.25,
p < 0.001), though the overall proportion of nitidulids carrying *C. apta* decreased over
time (β = -0.43 ± 0.16, z =-2.73, p = 0.006). Likewise, *S. geminata* carried significantly
more *C. apta* dauers over the whole season than did *E. ocularis* (β = 2.13 ± 0.24, z
=8.97, p < 0.001), and the number of dauers on both species decreased over time (β
= -0.50 ± 0.14, z =-3.71, p < 0.001). Though this survey misses early season and over-
winter dynamics, we recover a consistent pattern of higher *C. apta* dauer occupancy
on *S. geminata* than on *E. ocularis* across time, and expect a qualitatively similar
pattern exists across the year, particularly since nitidulid beetles overwinter as adults.

Other nematodes isolated from nitidulid beetles
While *C. apta* was the nematode species most often observed in our survey of these
two nitidulid species, several additional nematode species were encountered on
occasion. The next most frequently recovered was *Sheraphelenchus* (likely *S. sucus*
based on frequent occurrence on fruits in the area [(Perez *et al.* 2025)]), a

mycetophagous nematode in the family Aphelenchoididae (Kanzaki & Tanaka 2013).
Sheraphelenchus were found on 62 *E. ocularis* (28.1% of sampled beetles) and 14 *S.*
geminata (5.4% of sampled beetles), co-occurring with *C. apta* on 15 and 9 individual
beetles, respectively. As seen with *C. apta*, *Sheraphelenchus* were also frequently
observed clustered into large groups on the underside of the elytra of *E. ocularis*
(Figure 2E). In addition to these commonly observed species, we also recovered a
hermaphroditic *Pristionchus* species from 1 *E. ocularis* (0.45% of sampled beetles)
and 2 *S. geminata* (0.77% of sampled beetles), and a hermaphroditic Panagrolaimid
from 2 *E. ocularis* individuals (0.90% of sampled beetles), all co-occurring on beetles
with *C. apta* with the exception of one of the Panagrolaimids.
419
Global distribution patterns of C. apta *and its nitidulid vectors*
Beginning in 2007, *C. apta* was first isolated from sites across North America. In 2011,
C. apta was recorded for the first time from European collections. With the exception
of *C. apta,* all other known species in the Angaria group of *Caenorhabditis* nematodes
have only been collected from the Americas (Kiontke *et al.* 2011; Sudhaus *et al.* 2011;
Ferrari *et al.* 2017; Stevens *et al.* 2019; Sloat *et al.* 2022). North America is also the
native of range of *S. geminata* (Weber & Connell 1975), where it overlaps with the few
available collections of *C. apta* (Figure 4). *E. ocularis*, on the other hand, is native to
Southeast Asia and the western Pacific (Cline & Audisio 2011; Rittner & Nir 2013),
where *C. apta* has never been detected. Furthermore, *C. apta* has a wide and
overlapping distribution with the invasive range of *S. geminata* in Europe (Figure 4).
431
Discussion
A better understanding of the relationships between *Caenorhabditis* nematodes and
their phoretic vectors is necessary for characterizing the factors shaping the ecology
and evolution of these model organisms. However, studies of variation in vector use
among sympatric *Caenorhabditis* species, as well as basic characterization of vector
use for most *Caenorhabditis* species, have remained limited. In this study, we
quantified associations between three sympatric *Caenorhabditis* species and a variety
of invertebrate taxonomic groups, finding variation in vector use among species, as
well as a novel vector association for *C. apta*.
441
Variation in vector use among sympatric Caenorhabditis *species*

Despite all three *Caenorhabditis* species co-occurring at small spatial scales, even on
the same rotting fruits (Perez *et al.* 2025), we found variation in vector use among
species. In our general survey of 478 invertebrates from the orchard meadows near
the University of Konstanz, we found vector associations between *C. briggsae* and *C.*
remanei with gastropods, as well as *C. remanei* with isopods, but none of the other
surveyed taxa, consistent with vector associations previously reported for these
species (Fitch *et al.* 1994; Baird 1999; Félix & Duveau 2012; Petersen *et al.* 2015).
Caenorhabditis apta was never recovered from gastropods or isopods, instead only
being found in association with nitidulid beetles, and particularly with *S. geminata*. It
cannot be entirely ruled out that other beetle or insect species could act as vectors for
C. apta, particularly given the low sample size for some taxonomic groups; however,
based on the occurrence rates of the sampled invertebrates on and around rotting fruit
in our study area, we are confident that nitidulid beetles are frequently used vectors in
this area. The high occupancy of *C. apta* on nitidulids that we recorded across multiple
sites in Baden-Württemberg may indicate a higher propensity or ability for dispersal in
this species compared with the other two, perhaps mediated by underlying differences
in dispersal-related behaviors as has been observed among *C. elegans* isolates (Lee
et al. 2017). In line with this, the occurrence rates of *C. briggsae* and *C. remanei* on
isopods and gastropods documented in our survey roughly corresponds with the
results of a previous survey of vector-nematode associations from northern Germany
(Petersen *et al.* 2015), potentially indicating species-specific dispersal strategies.
Further work is necessary to determine the relationship between vector use, dispersal
ability, and community dynamics.

Our findings also provide evidence that *Caenorhabditis*-invertebrate interactions are
governed by mechanisms facilitating some degree of vector specificity. If Elegans
group nematodes were entirely opportunistic in their vector use, we would expect to
occasionally recover either species from the extremely common nitidulid beetles found
on most fruits in the study area. However, neither *C. briggsae* nor *C. remanei* was
found on any of the >450 nitidulid beetles sampled in this study. Likewise, *C. apta* was
not recovered from any gastropods or isopods, despite frequent occurrence of these
invertebrates on and around fruit containing *C. apta* dauers. Laboratory experiments
dissecting vector preference for different invertebrate species among these
Caenorhabditis could provide additional insight into the true degree of vector specificity

for each species and the mechanisms governing how they are able to differentiate
among co-occurring vectors. Preference for vector species has been experimentally
demonstrated for the highly specialized *C. japonica*, mediated by chemotactic
attraction to vector cues (Okumura *et al.* 2013; Okumura & Yoshiga 2014), while
experiments with *C. elegans* have failed to find any evidence of attraction to the
chemical profiles of known vector species (Archer *et al.* 2020; Petersen *et al.* 2023).
Determining how chemosensory or other cues aid in vector detection and response
will be of great value for better understanding the ecology, genetics, and evolution of
Caenorhabditis nematodes.

C. apta *dispersal on nitidulid beetles*
Detailed analysis of the association between *C. apta* and the two nitidulid beetles in
our study area highlighted that *C. apta* dauers frequently disperse in mixed sex groups
on the underside of beetles' elytra. Several beetle-dispersed nematodes form similar
clumps or clusters of individual worms on the inner surfaces of elytra (Penas *et al.*
2006; Shimizu *et al.* 2013; Kanzaki *et al.* 2019; Polyanina *et al.* 2019; Goczał & Beutel
2023). It remains unclear whether these elytral clusters are the direct result of group
dispersal behaviors or form once multiple nematodes have individually attached to the
vector. We have previously observed *C. apta* forming dauer towers in nature, as well
as towers reacting to and briefly attaching to fruit flies in the laboratory (Perez *et al.*
2025); however, it remains to be tested how dauer towers directly interact with
nitidulids, the apparent vectors of this species. Given the coordinated locomotion of
dauers in the tower structure on rotten fruits, it is plausible that the elytra clumps we
observe here are the direct results of towers, having collectively migrated underneath
the elytra once attached to the beetle. However, our recovery of both individual dauers
and groups of dauers from each beetle species indicates *C. apta* is can disperse on
nitidulid beetles both as individuals and in groups, providing an excellent system for
disentangling the benefits and costs of collective versus individual dispersal.

In addition to dispersing with members of the same species, we also observed co-
occurrence of more than one nematode species on the same beetle. As the
interactions between *C. apta* and co-occurring nematodes are yet unknown, it is
unclear what the causes or consequences of multi-species phoresy may be.
Sheraphelenchus and *Caenorhabditis* likely are not competing for resources, either

feeding on fungal mycelia or free-living microbes, respectively. On the other hand,

Pristionchus nematodes can be predatory, feeding on the larvae of other nematode

species (Hiramatsu & Lightfoot 2023; Sommer 2025), and may take advantage of

dispersing with other species in order to secure a meal upon arrival to a new habitat

patch. Testing whether there are benefits to dispersing in mixed-species groups

versus alone for any of these species would help shed light on whether co-occurrence

is simply due to chance or if one species may seek out or avoid beetles carrying the

others (Gupta & Borges 2021).

It is an open question whether either of these invasive nitidulid species is the natural

vector of *C. apta*, with which it has expanded its range to Europe by hitchhiking

together, or instead both of these beetle species are novel vectors *C. apta* has recently

begun to use. Given the known distribution of all other Angaria group species, the co-

distribution of *C.* apta and *S. geminata,* and the stronger association we detected

between *C. apta* and *S. geminata* compared to *E. ocularis*, it is possible that *C. apta*

is in fact a recent colonizer of Europe as a result of the expansion of its natural beetle

vector from North America. Co-invasion on insect vectors has been documented for

other phoretic nematodes (D'Anna & Sommer 2011), sometimes with severe

ecological and economic consequences (Giblin-Davis *et al.* 2013)*.* However, it is also

possible that, while initially undetected in Europe due to a patchy distribution or the

lack of targeted sampling, *C. apta* is indeed a native species that has recently been

able to increase in abundance or locally expand its range following a switch to the use

of invasive vector species, as has been observed in *Pristionchus* (D'Anna & Sommer

2011). Currently, the poor sampling of *C. apta* in North America, as well as a lack of

ecological information for this species, hinders further study of the history of these

interactions. Additional support for the invasion hypothesis will require documentation

of a similar phoretic interaction in the native range of *S. geminata*, as well as population

genetic analyses of *C. apta* from populations sampled from across the known range.

If *C. apta* is an invasive species in Europe, this system would provide an excellent

opportunity for studying the ecological and evolutionary consequences of range

expansion in *Caenorhabditis* nematodes, such as how native *Caenorhabditis* species

respond to a novel competitor.

Conclusion

Invertebrate vector associations for *Caenorhabditis* nematodes in southwestern
Germany correspond with current understanding of vector use in the genus. *C.*
briggsae and *C. remanei* associate with terrestrial invertebrates, though the
mechanisms governing this association remain unknown. The sympatric *C. apta*,
however, was exclusively recovered from nitidulid beetles, exhibiting a strong
association with *S. geminata* across both time and space. Further characterization of
the association between *C. apta* and these invasive nitidulid species in Europe and
the native ranges of the beetles will provide a better understanding of the ecology and
evolution of this little-studied nematode species.
554

555 **Data Accessibility Statement**
556 All data and code necessary to replicate the analysis and figures presented in this
557 manuscript are available at
558 https://github.com/SerenaDingLab/Greenway_et_al_Caen-Invert-Intxns.
559

560 **Competing Interests Statement**
561 The authors have no competing interests to declare.
562

563 **Author Contributions**
564 Conceptualization, data curation, formal analysis, visualization, and writing - original
565 draft preparation: R.G. Investigation: R.G., L.D., C.B., and M.A.F. Writing - review
566 and editing: R.G., L.D., C.B., M.A.F., and S.S.D. Supervision, project administration,
567 resources, and funding acquisition: S.S.D.
568

Acknowledgements
We would like to thank Anna-Lena Bucher, Narcís Font-Massot, Matthias Hermann,
Berke Güner, and Christian Weiler for their help accessing and traveling to field sites.
We thank all members of the Ding Lab at the MPI of Animal Behavior for constructive
discussions of the results and feedback on the manuscript draft. This study was funded
through the Max Planck Institute of Animal Behavior.

Appendix 1. Description of *Caenorhabditis apta* sp. n. (Nematoda: Rhabditidae)

Rationale for raising a new species

Most closely related *Caenorhabditis* species are highly similar morphologically; therefore, following Félix *et al.* (2014) we use the amenability of *Caenorhabditis* species to culturing and test crosses to delineate species based on the biological species concept. Following Félix *et al.* (2014), crosses were prioritized using genetic proximity. Building a consensus tree based on sequence alignments of 2,955 genes, Rockman *et al.* (2025) found the closest described species of *C. apta* QX1182 to be *C. agridulce* (Sloat *et al.* 2022). Sloat *et al.* (2022) reported that crosses between *C. apta* QX1182 males and *C. agridulce* QG555 females produced arrested embryos and a small number of L1 larvae that died, while the reciprocal cross only produced arrested embryos. *Caenorhabditis apta* thus defines a new biological species.

Methods

We used scanning electron microscopy for documenting and comparing morphological features of *C. apta* with closely related species. QX1182 nematodes on NGM agar culture plates were resuspended and washed twice in M9 solution, and fixed overnight at 4°C in glutaraldehyde 2.5%, in M9 or 50 mM phosphate pH 7.0 depending on the batch. The fixed animals were rinsed twice in M9 and dehydrated through an ethanol series, pelleting them at each step at 1 g in a tube. The samples were processed through critical point drying and coating with 20 nm of Au/Pd, and observed with a JEOL 6700F microscope at the Ultrastructural Microscopy Platform of the Pasteur Institute.

Species declaration

The electronic edition of this article conforms to the requirements of the amended International Code of Zoological Nomenclature, and hence the new names contained herein are available under that Code from the electronic edition of this article. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix "http://zoobank.org/". The LSID for this publication is: LSID urn:lsid:zoobank.org:pub:E5238355-9049-4697-

9422-97C9F2EA5616. The electronic edition of this work was published in a journal
with an ISSN. We provide pictures in Appendix Figs S1 and S2 to conform with the
ICZN guidelines.

Caenorhabditis apta sp. n. Greenway, Dalan, Braendle, Félix, & Ding
Zoobank identifier
urn:lsid:zoobank.org:act:E679D6EE-FDB9-476C-A274-6712AF93AA26
= *Caenorhabditis sp. 8* in (Kiontke *et al.* 2011; Félix *et al.* 2014; Vielle *et al.* 2016; Sloat
et al. 2022; Rockman *et al.* 2025)

The type isolate by present designation is QX1182, deposited at the *Caenorhabditis*
Genetics Center (https://cgc. umn.edu/). The species is delineated and diagnosed by
the fertile cross with the type isolate QX1182 in both cross directions, yielding highly
fertile hybrid females and males that are interfertile and cross-fertile with their parent
strains. The species reproduces through males and females. The ribosomal DNA
sequence of *C. apta* QX1182 including ITS2 is accessible at NCBI with accession
number JN636143 (Kiontke *et al.* 2011). This species differs by SSU, LSU and ITS2
DNA sequences (Kiontke *et al.* 2011) from all species listed in Tables 1 and 2 of (Félix
et al. 2014; Huang *et al.* 2014; Ferrari *et al.* 2017; Slos *et al.* 2017; Kanzaki *et al.* 2018;
Crombie *et al.* 2019; Stevens *et al.* 2019, 2020; Dayi *et al.* 2021; Sloat *et al.* 2022;
Devi *et al.* 2025; Rockman *et al.* 2025). Note that these ribosomal DNA sequences
may vary within the species (Kiontke *et al.* 2011). The type isolate was collected by
Matthew Rockman from rotting tomatoes in New Jersey, USA, in July 2007 (Kiontke
et al. 2011). Other isolates were found in decomposing fruits in USA and Germany
(Kiontke *et al.* 2011; Perez *et al.* 2025), as well as on decomposing fruits in
Switzerland, Austria, Czech Republic and Hungary and on nitidulid beetles in Germany
in the present study. The mouth is endowed with the usual set of sensory organs
disposed in a concentric manner, namely six labial sensillae, two amphids and six
male-specific cephalic sensillae (Fig. A1A-C). The radial symmetry of the labial
sensillae appears broken by the more external position of the two sensillae closest to
the amphids (Fig. A1A-B). An inner flap is present on each of the six lips, as in *C.*
angaria (Sudhaus *et al.* 2011) and *C. agridulce* (Sloat *et al.* 2022) (Fig. A1A-C). The
male tail displays an anteriorly open and narrow fan with a smooth edge (Fig. A2B-D).
The dorsal rays are in antero-posterior positions 4 and 7 (Fig. A2A-D). Papilliform

phasmids are visible (Kiontke *et al.* 2011) (Fig. A2A). The spicule tips are thick and

curved internally (Fig. A2E-F). The pre-cloacal sensillum is semi-circular and appears

bordered anteriorly by an additional cuticular lip (Fig. A2G-H). In addition, a drawing

of the male tail can be found in Kiontke *et al.* 2011. A Nomarski microscopy picture of

the buccal cavity and a drawing of a spicule can be found in (Sloat *et al.* 2022). The

males mate in a spiral position. The species is named after their ability to be connected

and attached together in masses.

 Appendix 2.

 Table A1. Summary of characters used to distinguish among sympatric
 Caenorhabditis species.

Character	*C. apta*	*C. briggsae*	*C. remanei*
Body shape	Short, dumpy	Long, slender	Long, slender
Reproductive mode	Gonochoristic	Hermaphroditic and selfing	Gonochoristic
Mating behavior	Spiral	-	Parallel
Male morphology	Stout, amber colored spicules Narrow bursal fan	-	Smaller spicules Wide bursal fan
Foraging behavior	Forms small aggregations on food No observed burrowing on food	Solitary on food No observed burrowing on food	Forms large aggregations on food Burrows into NGM when food present

References

Ambros, V.R., Chalfie, M., Daul, A.L., Fire, A.Z., Hall, D.H., Horvitz, H.R., *et al.* (2025). From nematode to Nobel: How community-shared resources fueled the rise of Caenorhabditis elegans as a research organism. *Proc. Natl. Acad. Sci. U. S. A.*, 122, e2522808122.

Archer, H., Deiparine, S. & Andersen, E.C. (2020). The nematode Caenorhabditis elegans and the terrestrial isopod Porcellio scaber likely interact opportunistically. *PLoS One*, 15, e0235000.

Baird, S.E. (1999). Natural and experimental associations of Caenorhabditis remanei with Trachelipus rathkii and other terrestrial isopods. *Nematology*, 1, 471–475.

Barrière, A. & Félix, M.-A. (2005). High local genetic diversity and low outcrossing rate in Caenorhabditis elegans natural populations. *Curr. Biol.*, 15, 1176–1184.

Barrière, A. & Félix, M.A. (2006). Isolation of C. elegans and related nematodes. *WormBook*, 17, 1–9.

Barrière, A. & Félix, M.-A. (2007). Temporal dynamics and linkage disequilibrium in natural Caenorhabditis elegans populations. *Genetics*, 176, 999–1011.

Bartlow, A.W. & Agosta, S.J. (2021). Phoresy in animals: review and synthesis of a common but understudied mode of dispersal. *Biol. Rev. Camb. Philos. Soc.*, 96, 223–246.

Benisch, C. (2023). *kerbtier.de – Beetle Fauna of Germany*. Available at: https://www.kerbtier.de/enindex.html. Last accessed 11 November 2024.

Bibin, A.R. (2017). Invasive sap beetles Epuraea ocularis and Stelidota geminata (Coleoptera, Nitidulidae) in the Russian Black Sea region. *Russian Journal of Biological Invasions*. https://doi.org/10.1134/S2075111717040014

Bubrig, L.T. & Fierst, J. (2021). Review of the dauer hypothesis: What non-parasitic species can tell us about the evolution of parasitism. *J. Parasitol.*, 107, 717–725.

Callot, H. (2007). Stelidota geminata (Say, 1825), espèce envahissante americaine, est bien implantée en Alsace, come ailleurs dans la haut valée du Rhin (Coleoptera …. *Entomologiste (Paris)*.

Chiba, T., Okumura, E., Nishigami, Y., Nakagaki, T., Sugi, T. & Sato, K. (2023). Caenorhabditis elegans transfers across a gap under an electric field as dispersal behavior. *Curr. Biol.*, 33, 2668-2677.e3.

Cline, A.R. & Audisio, P. (2011). Epuraea (Haptoncus) Ocularis Fairmaire (Coleoptera: Nitidulidae) Recently Found in the U.S.A., with Comments on Nearctic Members of Epuraea Erichson. *The Coleopterists Bulletin*, 65, 24–26.

Corsi, A.K., Wightman, B. & Chalfie, M. (2015). A Transparent Window into Biology: A Primer on Caenorhabditis elegans. *Genetics*, 200, 387–407.

Crombie, T.A., Zdraljevic, S., Cook, D.E., Tanny, R.E., Brady, S.C., Wang, Y., *et al.* (2019). Deep sampling of Hawaiian Caenorhabditis elegans reveals high genetic diversity and admixture with global populations. *Elife*, 8, e50465.

Cutter, A.D. (2015). Caenorhabditis evolution in the wild. *Bioessays*, 37, 983–995.

D'Anna, I. & Sommer, R.J. (2011). Pristionchus uniformis, should I stay or should I go? Recent host range expansion in a European nematode: Pristionchus uniformisHost Range Expansion. *Ecol. Evol.*, 1, 468–478.

Dayi, M., Kanzaki, N., Sun, S., Ide, T., Tanaka, R., Masuya, H., *et al.* (2021). Additional description and genome analyses of Caenorhabditis auriculariae representing the basal lineage of genus Caenorhabditis. *Sci. Rep.*, 11, 6720.

Devi, M.P., Haryoso, E., Rais, E.I., Karuniawan, A., Yahya, M.Q., Richaud, A., *et al.* (2025). Five new Caenorhabditis species from Indonesia provide exceptions to Haldane's rule and partial fertility of interspecific hybrids. *G3 (Bethesda)*, 15. https://doi.org/10.1093/g3journal/jkaf134

Dey, A., Jeon, Y., Wang, G.-X. & Cutter, A.D. (2012). Global population genetic structure of Caenorhabditis remanei reveals incipient speciation. *Genetics*, 191, 1257–1269.

Diano, M.A., Dalan, L., Singh, P.R. & Sumaya, N.H. (2022). First report, morphological and molecular characterization of Caenorhabditis brenneri (Nematoda: Rhabditidae)

isolated from the giant African land snail Achatina fulica (Gastropoda: Achatinidae). *Biologia (Bratisl.)*, 77, 469–478.

Félix, M.-A., Braendle, C. & Cutter, A.D. (2014). A streamlined system for species diagnosis in Caenorhabditis (Nematoda: Rhabditidae) with name designations for 15 distinct biological species. *PLoS One*, 9, e94723.

Félix, M.-A. & Duveau, F. (2012). Population dynamics and habitat sharing of natural populations of Caenorhabditis elegans and C. briggsae. *BMC Biol.*, 10, 59.

Félix, M.-A., Jovelin, R., Ferrari, C., Han, S., Cho, Y.R., Andersen, E.C., *et al.* (2013). Species richness, distribution and genetic diversity of Caenorhabditis nematodes in a remote tropical rainforest. *BMC Evol. Biol.*, 13, 10.

Ferrari, C., Salle, R., Callemeyn-Torre, N., Jovelin, R., Cutter, A.D. & Braendle, C. (2017). Ephemeral-habitat colonization and neotropical species richness of Caenorhabditis nematodes. *BMC Ecol.*, 17, 43.

Fitch, D.H.A., Emmons, S.W. & Baird, S.E. (1994). Caenorhabditis Vulgaris Sp.N. (Nematoda: Rhabditidae): a Necromenic Associate of Pill Bugs and Snails. *Nematologica*, 40, 1–11.

Frézal, L. & Félix, M.-A. (2015). C. elegans outside the Petri dish. *Elife*, 4. https://doi.org/10.7554/eLife.05849

Fusca, D.D., Dall'Acqua, M.N., Sánchez-Ramírez, S. & Cutter, A.D. (2025). Phylogenomic timetree-calibrated speciation clocks for Caenorhabditis nematodes reveal slow but disproportionate accumulation of post-zygotic reproductive isolation. *PLoS Genet.*, 21, e1011852.

Giblin-Davis, R.M., Kanzaki, N. & Davies, K.A. (2013). Nematodes that ride insects: Unforeseen consequences of arriving species[§]. *Fla. Entomol.*, 96, 770–780.

Goczał, J. & Beutel, R.G. (2023). Beetle elytra: evolution, modifications and biological functions. *Biol. Lett.*, 19, 20220559.

Grujić, N., Nježić, B., Anifantis, A.S. & Tarasco, E. (2020). Biocontrol potential of some entomopathogenic nematodes against *Stelidota geminata* (Say). *Redia (Firenze)*, 103, 35–39.

Gupta, S. & Borges, R.M. (2021). Hopping on: Conspecific traveller density within a vehicle regulates parasitic hitchhiking between ephemeral microcosms. *J. Anim. Ecol.*, 90, 899–908.

Hiramatsu, F. & Lightfoot, J.W. (2023). Kin-recognition and predation shape collective behaviors in the cannibalistic nematode Pristionchus pacificus. *PLoS Genet.*, 19, e1011056.

Huang, R.-E., Ren, X., Qiu, Y. & Zhao, Z. (2014). Description of Caenorhabditis sinica sp. n. (Nematoda: Rhabditidae), a nematode species used in comparative biology for C. elegans. *PLoS One*, 9, e110957.

Huang, Y., Lo, Y.-H., Hsu, J.-C., Le, T.S., Yang, F.-J., Chang, T., *et al.* (2023). Widespread sex ratio polymorphism in Caenorhabditis nematodes. *R. Soc. Open Sci.*, 10, 221636.

Hunt, D.J. (1993). *Aphelenchida, Longidoridae and Trichodoridae: their systematics and bionomics*. cabidigitallibrary.org.

Jelínek, J., Audisio, P., Hajek, J., Baviera, C., Moncourtier, B., Barnouin, T., *et al.* (2016). Epuraea imperialis (Reitter, 1877). New invasive species of Nitidulidae (Coleoptera) in Europe, with a checklist of sap beetles introduced to Europe and Mediterranean areas. *AAPP| Physical, Mathematical, and Natural Sciences*, 94, 1–24.

Kanzaki, N., Liang, W.-R., Chiu, C.-I., Yang, C.-T., Hsueh, Y.-P. & Li, H.-F. (2019). Nematode-free agricultural system of a fungus-growing termite. *Sci. Rep.*, 9, 8917.

Kanzaki, N. & Tanaka, R. (2013). Sheraphelenchus sucus n. sp. (Tylenchina: Aphelenchoididae) isolated from sap flow of Quercus serrata in Japan. *Nematology*, 15, 975–990.

Kanzaki, N., Tsai, I.J., Tanaka, R., Hunt, V.L., Liu, D., Tsuyama, K., *et al.* (2018). Biology and genome of a newly discovered sibling species of Caenorhabditis elegans. *Nat. Commun.*, 9, 3216.

Kiontke, K. (1997). Description of Rhabditis (Caenorhabditis) drosophilae n. sp. and R. (C.) sonorae n. sp. (Nematoda: Rhabditida) from saguaro cactus rot in Arizona. *Fundamental and applied nematology*, 20, 305–315.

Kiontke, K. & Sudhaus, W. (2006). Ecology of Caenorhabditis species. *WormBook*, 1–14.

Kiontke, K.C., Félix, M.-A., Ailion, M., Rockman, M.V., Braendle, C., Pénigault, J.-B., *et al.* (2011). A phylogeny and molecular barcodes for Caenorhabditis, with numerous new species from rotting fruits. *BMC Evol. Biol.*, 11, 339.

Konzelmann, E. (2001). Epuraea (Haptoncus) ocularis Fairmaire an faulendem Kernobst in Baden, Württemberg und in der Pfalz (Coleoptera: Nitidulidae). *Mitteilungen Entomologischer Verein Stuttgart*, 36, 35–43.

Lasoń, A., Dobosz, R., Greń, C. & Grzywocz, J. (2023). Epuraea (Haptoncus) ocularis Fairmaire, 1849 (Coleoptera: Nitidulidae): the first record in Poland. *Annals of the Upper Silesian Museum in Bytom, Entomology*, 32, 1–4.

Lee, D., Yang, H., Kim, J., Brady, S., Zdraljevic, S., Zamanian, M., *et al.* (2017). The genetic basis of natural variation in a phoretic behavior. *Nat. Commun.*, 8, 273.

Lee, H., Choi, M.-K., Lee, D., Kim, H.-S., Hwang, H., Kim, H., *et al.* (2011). Nictation, a dispersal behavior of the nematode Caenorhabditis elegans, is regulated by IL2 neurons. *Nat. Neurosci.*, 15, 107–112.

Li, S., Jovelin, R., Yoshiga, T., Tanaka, R. & Cutter, A.D. (2014). Specialist versus generalist life histories and nucleotide diversity in Caenorhabditis nematodes. *Proc. Biol. Sci.*, 281, 20132858.

Lompe, A. (2002). *Käfer Europas*. Available at: https://www.coleonet.de/coleo/html/start.htm. Last accessed 1 November 2024.

Meneely, P.M., Dahlberg, C.L. & Rose, J.K. (2019). Working with Worms: *Caenorhabditis elegans* as a Model Organism. *Curr. Protoc. Essent. Lab. Tech.*, 19, e35.

Okumura, E., Tanaka, R. & Yoshiga, T. (2013). Species-specific recognition of the carrier insect by dauer larvae of the nematode Caenorhabditis japonica. *J. Exp. Biol.*, 216, 568–572.

Okumura, E. & Yoshiga, T. (2014). Host orientation using volatiles in the phoretic nematode Caenorhabditis japonica. *J. Exp. Biol.*, 217, 3197–3199.

Penas, A.C., Bravo, M.A., Naves, P., Bonifácio, L., Sousa, E. & Mota, M. (2006). Species of *Bursaphelenchus* Fuchs, 1937 (Nematoda: Parasitaphelenchidae) and other nematode genera associated with insects from *Pinus pinaster* in Portugal. *Ann. Appl. Biol.*, 148, 121–131.

Perez, D.M., Greenway, R., Stier, T., Font-Massot, N., Pertzelan, A. & Ding, S.S. (2025). Towering behavior and collective dispersal in Caenorhabditis nematodes. *Curr. Biol.*, 35, 2980-2986.e4.

Petersen, C., Hermann, R.J., Barg, M.-C., Schalkowski, R., Dirksen, P., Barbosa, C., *et al.* (2015). Travelling at a slug's pace: possible invertebrate vectors of Caenorhabditis nematodes. *BMC Ecol.*, 15, 19.

Petersen, C., Krahn, A. & Leippe, M. (2023). The nematode Caenorhabditis elegans and diverse potential invertebrate vectors predominantly interact opportunistically. *Front. Ecol. Evol.*, 11. https://doi.org/10.3389/fevo.2023.1069056

Polyanina, K.S., Mandelshtam, M.Y. & Ryss, A.Y. (2019). Brief Review of the Associations of Xylobiont Nematodes with Bark Beetles (Coleoptera, Curculionidae: Scolytinae). *Entomological Review*, 99, 598–614.

R Core Team. (2023). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.

Ragsdale, E.J., Kanzaki, N. & Herrmann, M. (2015). Taxonomy and natural history: the genus Pristionchus. In: *Pristionchus pacificus*. BRILL, pp. 77–120.

Rittner, O. & Nir, A. (2013). First record of Myrrha octodecimguttata (Coleoptera: Coccinellidae) and Epuraea ocularis (Coleoptera: Nitidulidae) from Israel. *Zootaxa*, 3609, 349–350.

Rockman, M.V., Tintori, S.C., Nguyen, T.H.M. & Harmony Yomai, V.M. (2025). Caenorhabditis diversity on Pohnpei, Micronesia, provides evidence that the Elegans

820   Supergroup has its roots in the Americas and diversified in the Pacific en route to
821   Asia. *bioRxivorg*. https://doi.org/10.1101/2025.09.22.677770
822 Schratzberger, M., Holterman, M., van Oevelen, D. & Helder, J. (2019). A worm's world:
823   Ecological flexibility pays off for free-living nematodes in sediments and soils.
824   *Bioscience*, 69, 867–876.
825 Schulenburg, H. & Félix, M.-A. (2017). The Natural Biotic Environment of Caenorhabditis
826   elegans. *Genetics*, 206, 55–86.
827 Shimizu, A., Tanaka, R., Akiba, M., Masuya, H., Iwata, R., Fukuda, K., *et al.* (2013).
828   Nematodes associated with Dryocoetes uniseriatus (Coleoptera: Scolytidae).
829   *Environ. Entomol.*, 42, 79–88.
830 Sloat, S.A., Noble, L.M., Paaby, A.B., Bernstein, M., Chang, A., Kaur, T., *et al.* (2022).
831   Caenorhabditis nematodes colonize ephemeral resource patches in neotropical
832   forests. *Ecol. Evol.*, 12, e9124.
833 Slos, D., Sudhaus, W., Stevens, L., Bert, W. & Blaxter, M. (2017). Caenorhabditis
834   monodelphis sp. n.: defining the stem morphology and genomics of the genus
835   Caenorhabditis. *BMC Zool.*, 2, 1–15.
836 Sommer, R.J. (2025). Pristionchus - Beetle associations: Towards a new natural history. *J.*
837   *Invertebr. Pathol.*, 209, 108243.
838 Stan, M. (2019). The first record of Stelidota geminata (Say, 1825) (Coleoptera, Nitidulidae)
839   in Romania. *Trav. Mus. Natl. Hist. Nat. Grigore Antipa*, 62, 57–60.
840 Stevens, L., Félix, M.-A., Beltran, T., Braendle, C., Caurcel, C., Fausett, S., *et al.* (2019).
841   Comparative genomics of 10 new Caenorhabditis species. *Evol Lett*, 3, 217–236.
842 Stevens, L., Rooke, S., Falzon, L.C., Machuka, E.M., Momanyi, K., Murungi, M.K., *et al.*
843   (2020). The genome of Caenorhabditis bovis. *Curr. Biol.*, 30, 1023-1031.e4.
844 Sudhaus, W., Giblin-Davis, R. & Kiontke, K. (2011). Description of Caenorhabditis angaria n.
845   sp. (Nematoda: Rhabditidae), an associate of sugarcane and palm weevils
846   (Coleoptera: Curculionidae). *Nematology*, 13, 61–78.
847 Sun, S., Kanzaki, N., Dayi, M., Maeda, Y., Yoshida, A., Tanaka, R., *et al.* (2022). The
848   compact genome of Caenorhabditis niphades n. sp., isolated from a wood-boring
849   weevil, Niphades variegatus. *BMC Genomics*, 23, 765.
850 Ting, J.J., Woodruff, G.C., Leung, G., Shin, N.-R., Cutter, A.D. & Haag, E.S. (2014). Intense
851   sperm-mediated sexual conflict promotes reproductive isolation in Caenorhabditis
852   nematodes. *PLoS Biol.*, 12, e1001915.
853 Van Driessche, R., De Ley, I.T., De Ley, P., Segers, H. & Bert, W. (2003). Baujardia mirabilis
854   gen. n., sp. n. from pitcher plants and its phylogenetic position within
855   Panagrolaimidae (Nematoda: Rhabditida). *Nematology*, 5, 405–420.
856 Vielle, A., Callemeyn-Torre, N., Gimond, C., Poullet, N., Gray, J.C., Cutter, A.D., *et al.*
857   (2016). Convergent evolution of sperm gigantism and the developmental origins of
858   sperm size variability in Caenorhabditis nematodes. *Evolution*, 70, 2485–2503.
859 Vlaar, L.E., Bertran, A., Rahimi, M., Dong, L., Kammenga, J.E., Helder, J., *et al.* (2021). On
860   the role of dauer in the adaptation of nematodes to a parasitic lifestyle. *Parasit.*
861   *Vectors*, 14, 554.
862 Volk, J. (1950). Nematodes associated with earthworms and carrion beetles, 79, 1–70.
863 Wang, G.-X., Ren, S., Ren, Y., Ai, H. & Cutter, A.D. (2010). Extremely high molecular
864   diversity within the east Asian nematode *Caenorhabditis* Sp. 5. *Mol. Ecol.*, 19, 5022–
865   5029.
866 Weber, R.G. & Connell, W.A. (1975). Stelidota geminata (Say): Studies of Its Biology
867   (Coleoptera: Nitidulidae1). *Ann. Entomol. Soc. Am.*, 68, 649–653.
868 *What is GBIF?* (2025). . Available at: https://www.gbif.org/what-is-gbif. Last accessed 17
869   November 2025.
870 Williams, B., Nelson, M., McRobert, S. & Fingerut, J. (2023). An easy and inexpensive
871   method for determining the rate of individual phoretic events of nematodes.
872   *MicroPubl. Biol.*, 2023. https://doi.org/10.17912/micropub.biology.000942
873 Woodruff, G.C. & Phillips, P.C. (2018). Field studies reveal a close relative of C. elegans
874   thrives in the fresh figs of Ficus septica and disperses on its Ceratosolen pollinating

875   wasps. *BMC Ecol.*, 18, 26.

876   Yoshiga, T., Ishikawa, Y., Tanaka, R., Hironaka, M. & Okumura, E. (2013). Species-specific

877   and female host-biased ectophoresy in the roundworm Caenorhabditis japonica. *Sci.*

878   *Nat.*, 100, 205–208.

Figure 1. Phylogeny of *Caenorhabditis* nematodes, modified from Fusca *et al.* (2025), with the species groups and supergroups mentioned in this study indicated. Branches are colored by vector taxonomic group for species with documented associations, and the three species included in this study are indicated by bold red text.

Figure 2. Collection sites and focal beetle species from this study. (**A**) *Stelidota geminata* and *Epuraea ocularis* were collected from six locations in Baden-Württemberg, Germany. The inset map depicts sites around the University of Konstanz campus. (**B**) *Stelidota geminata*, with the right elytron outlined. (**C**) *Epuraea ocularis*. (**D,E**) Clusters of dauer larvae (circled area) attached to the inner surface of the right elytron of *S. geminata* (>20 *C. apta,* **D**) or *E. ocularis* (>30 *Sheraphelenchus* sp., **E**). The epipleuron is indicated by an arrow in both images.

Figure 3. Temporal variation in (**A**) the proportion of beetles carrying *C. apta* and (**B**) the number of *C. apta* dauer larvae extracted per beetle (square root scaled) for two nitidulid beetle species. In **B**, large points represent the mean number of nematodes per beetle for each species with error bars depicting the standard error, small points represent the number of nematodes extracted from individual beetles.

Figure 4. Global distribution of *S. geminata* and known occurrences of *Caenorhabditis* species in the Angaria group, including *C. apta* (blue triangles). *S. geminata* has spread from its native range in the Americas to Europe since the 2000s (orange versus red points; more intense coloration corresponds to multiple overlapping occurrences). As most Angaria group species have been described and sampled from French Guiana, data points heavily overlap in this region.

Table 1. Summary of *Caenorhabditis* species isolated from different invertebrate groups around the University of Konstanz orchard meadows.

Invertebrate Group	No. Sampled Individuals	*C. briggsae*	*C. remanei*	*C. apta*
Annelida	3	0	0	0
Araneae	2	0	0	0
Blattodea	1	0	0	0
Coleoptera (Nitidulidae)	196	0	0	9 (4.6%)
Coleoptera (Other)	14	0	0	0
Collembola	21	0	0	0
Diptera	80	0	0	0
Gastropoda	65	2 (3.1%)	4 (6.2%)	0
Hemiptera	1	0	0	0
Hymenoptera	3	0	0	0
Isopoda	88	0	9 (10.2%)	0
Myriapoda	4	0	0	0

Table 2. Summary of among site variation in the number of beetles carrying *C. apta* dauers, the location of dauers on the beetles, and the sex ratio of *C. apta* from beetles.

Site	Species	No. beetles	No. positive	Dauers/beetle Mean ± SE	Dauers on elytra Mean ± SE	Dauers on body Mean ± SE	No. beetles (Sex Ratio)	Sex Ratio Mean ± SE
Egg Orchard	*Epuraea ocularis*	61	25 (41.0%)	1.87 ± 0.53	1.84 ± 0.51	0.03 ± 0.02	13	0.34 ± 0.13
	Stelidota geminata	61	47 (77.0%)	10.52 ± 3.37	9.62 ± 2.85	0.90 ± 0.58	30	0.47 ± 0.06
Hockgraben	*Epuraea ocularis*	30	10 (33.3%)	1.37 ± 0.60	0.77 ± 0.23	0.60 ± 0.47	9	0.38 ± 0.14
	Stelidota geminata	30	24 (80.0%)	10.83 ± 2.68	9.23 ± 2.45	1.60 ± 0.72	22	0.52 ± 0.05
Möggingen	*Epuraea ocularis*	35	8 (22.9%)	0.51 ± 0.22	0.46 ± 0.21	0.06 ± 0.04	8	0.17 ± 0.09
	Stelidota geminata	35	18 (51.4%)	2.34 ± 0.65	2.29 ± 0.65	0.06 ± 0.06	17	0.54 ± 0.08
Niederhofen[‡]	*Epuraea ocularis*	1	1 (100.0%)	1.00				
	Stelidota geminata	3	3 (100.0%)	7.67 ± 5.24				
St. Katarina	*Epuraea ocularis*	30	16 (53.3%)	2.00 ± 0.59	1.70 ± 0.57	0.30 ± 0.09	12	0.49 ± 0.14
	Stelidota geminata	29	23 (79.3%)	8.24 ± 1.91	7.59 ± 1.76	0.66 ± 0.38	13	0.48 ± 0.12
Tübingen	*Epuraea ocularis*	20	3 (15.0%)	0.35 ± 0.22	0.35 ± 0.22	0	3	0.42 ± 0.30
	Stelidota geminata	39	20 (51.3%)	4.56 ± 1.10	4.56 ± 1.10	0	20	0.54 ± 0.05
All Sites[‡]	*Epuraea ocularis*	177	63 (35.6%)	1.36 ± 0.24	1.19 ± 0.22	0.18 ± 0.08	45	0.36 ± 0.06
	Stelidota geminata	197	135 (68.5%)	7.56 ± 1.20	6.92 ± 1.05	0.64 ± 0.22	102	0.51 ± 0.03

[‡]Beetles from Niederhofen were excluded from analysis due to low sample sizes
SE = standard error

Appendix Figure A1. Scanning electron microscopy of heads and female vulva. (**A**): Head of adult male in anterior view. (**B**): Head of adult male in antero-lateral view. (**C**) Head of adult female in antero-lateral view (**D,E**): Female vulval openings in ventral views. In (**E**), alae are visible on the top of the micrograph. a: amphid; ls: labial sensillum; cem: male cephalic sensillum; f: flap on lips; v: vulva. Bars: (**A-C**) 1 µm; (**D,E**) 10 µm.

Appendix Figure A2. Scanning electron microscopy of male tails.
(**A**) Dorsal view. (**B-D**) Ventro-lateral views, also from the posterior direction in (**C,D**). (**E,F**) Lateral views with extruded spicules. (**G,H**) Ventro-lateral views of the cloacal opening. Anterior is to the left in (**A,G,H**); to the right in (**B-F**). The male rays are numbered r1 to r9 according to antero-posterior position. The anterior dorsal ray is r4, and the posterior dorsal ray is r7. ph: phasmid. sp: spicule: pcs: precloacal sensillum. Bars: (**A-F**) 10 µm; (**G-H**) 1 µm.