1 **A survey of bacterial and fungal communities of table olives.**
2

3 Eugenio Parente[1], Rocchina Pietrafesa[1], Francesca De Filippis[2], Alessandra De Vivo[2], Maria
4 Grazia Labella[1], Marina Hidalgo[3], Emanuela Lavanga[1], Annamaria Ricciardi[1]
5
6 **Affiliations**
7 1 Dipartimento di Scienze Agrarie, Forestali, Alimentari ed Ambientali, Università degli Studi
8 della Basilicata, Potenza, Italy
9 2 Dipartimento di Agraria, Università degli Studi di Napoli "Federico II", Portici (NA), Italy
10 3 Departamento de Ciencias de la Salud, Universidad de Jaén, Jaén, Spain
11
12
Corresponding author
Prof. Eugenio PARENTE
Dipartimento di Scienze Agrarie, Forestali, Alimentari ed Ambientali
Università degli Studi della Basilicata
Via dell'Ateneo Lucano, 10
85100 Potenza, Italy
19
20

Abstract

Table olives are produced from a large number of olive varieties subjected to different trade preparations, resulting in a highly heterogeneous family of fermented foods. To characterise the diversity of bacterial and fungal communities and its relationship with variety, ripeness, and trade preparation, we surveyed 363 samples from 40 producers across 6 countries, combining physicochemical measurements, viable counts, and amplicon-based metagenomics. This is the largest survey of table olive microbial communities to date and includes the first culture-independent characterisation of microbial communities for several Italian PDO and non-PDO varieties, most notably Oliva di Gaeta. The contrast between alkali-treated and naturally fermented olives was the dominant structuring factor, with HALAB (Halophilic and Alkalophilic Lactic Acid Bacteria) and other halophiles enriched in alkali-treated varieties and a diverse array of *Lactobacillaceae* and *Pseudomonadota* characterising naturally fermented olives. Despite these consistent signals, striking variability was observed within the same variety and even within the same producer, driven by stochastic colonization events, house microbiota, and the widespread use of small fermentation vessels. This variability obscured variety-specific microbial signatures and prevented reliable discrimination of Italian PDO varieties from similar non-PDO counterparts using amplicon-based approaches. The ecological and taxonomic complexity documented here, encompassing bacterial and fungal genera with largely untapped starter and flavour potential, provides the foundation for the development of variety-specific microbiome-based starter cultures.

Keywords

Table olives; amplicon targeted metagenomics; Oliva di Gaeta; olive fermentation; microbiota dynamics.

Highlights

- We report a metataxonomic survey of microbial communities in 363 table olive samples
- Alkali treatment and natural fermentation drive distinct microbial community structures
- House microbiota and stochastic colonization generate high within-variety variability
- Microbiome data provide ecological foundations for olive microbiome-based starters

1. Introduction

Fermented foods have been part of the human diet for millennia and their diversity is huge (Gänzle, 2022; Hernández-Velázquez et al., 2024; Tamang et al., 2020). Table olives are among the most ancient fermented vegetables from the Mediterranean basin (Langgut and Garfinkel, 2022) and this has resulted in the evolution of a great variety of products (Anagnostopoulos and Tsaltas, 2022; Perpetuini et al., 2020). Fermentation of table olives has multiple functions: removing the bitter glucoside oleuropein, enhancing safety and shelf life through acid production and inhibition of spoilage microorganisms, and improving sensory properties. Five main trade preparations of table olives are recognized by the International Olive Oil Council (https://www.internationaloliveoil.org/wp-content/uploads/2019/11/COI-OT-NC1-2004-Eng.pdf). Treated olives, also known as Spanish style olives, are treated with sodium hydroxide to remove oleuropein before being washed and fermented in brine. Green specialties, like Picholine style olives, on the other hand, are

stored at low temperature in brine after the alkali treatment, thus limiting fermentation.
Natural olives (Greek style) include both green and ripe olives directly fermented in brines.
Dried/shrivelled olives and olives darkened by oxidation are preserved by methods other
than fermentation.
Because of their long history, table olives production is strongly associated with local
varieties and practices. In Italy four varieties (Oliva di Gaeta, Nocellara del Belice, Bella di
Daunia, Oliva Ascolana del Piceno) are included in the Protected Denomination of Origin
(PDO) food list and Oliva Taggiasca Ligure has recently received Protected Geographical
Indication status (https://ec.europa.eu/agriculture/eambrosia/geographical-indications-
register/). Many more are included in the Italian List of Traditional Agricultural Products
(https://www.gazzettaufficiale.it/eli/gu/2024/03/13/61/sg/pdf). Spain, Greece, France,
Portugal and Turkey also produce a wide variety of table olives, including PDO products.
Sensory quality, safety and stability of table olives are affected by several factors, including
olive ripeness, trade preparation, and by the growth of microorganisms, either present as
natural contaminants on the fruits, salt, brines, and contact surfaces or added as starter
cultures (Anagnostopoulos and Tsaltas, 2022; Campus et al., 2018). Geographic origin is also
believed to be a contributing factor for initial drupe contamination (Argyri et al., 2020;
Ferrocino et al., 2025; Kamilari et al., 2023). Current trends in table olive production aim at
controlling the microbiota involved in fermentation and spoilage by sanitizing the raw
materials, by adding starter and functional cultures, and/or by reducing salt content
(Campus et al., 2018). Addition of starters has been successfully tested for several varieties
and, typically, involves the inoculation of lactic acid bacteria (*Lactiplantibacillus pentosus* or
Lactipl. plantarum, Lacticaseibacillus paracasei: Alfonzo et al., 2023b; Comunian et al., 2017;
Lanza et al., 2020, 2021; Paba et al., 2020; Pino et al., 2018, 2019; Randazzo et al., 2017;
Rodríguez-Gómez et al., 2023; Ruiz-Barba and Jimenéz-Diaz, 2012; Servili et al., 2006;
Tzamourani et al., 2024; Vougiouklaki et al., 2024; Zago et al., 2013), yeasts (several species
belonging to the genera *Candida, Wyckerhamomyces, Pichia, Cyteromyces* have been tested:
Ciafardini and Zullo, 2019; Montaño et al., 2021; Ruiz-Barba et al., 2024; Tufariello et al.,
2019) or their combinations (Benítez-Cabello et al., 2020; Garrido-Fernández et al., 2021).
Back-slopping practices (Martorana et al., 2015) have also been described.
The composition and dynamics of microbial communities in table olives have been
extensively reviewed (Anagnostopoulos and Tsaltas, 2022; Campus et al., 2018; Portilha-
Cunha et al., 2020; Tsoungos et al., 2023; Vaccalluzzo et al., 2020). The microbiota of table
olives is complex and variable, usually due to the lack of control of unit operations affecting
dispersal and selection, and includes microbial groups which are considered beneficial, like
lactic acid bacteria, yeasts, and other halophilic microorganisms (Halophilic and Alkalophilic
Lactic Acid Bacteria, HALAB, and halophilic *Pseudomonadota*), but also microorganisms
associated with spoilage, like some halophiles (*Alkalibacterium, Marinilactobacillus,*
Celerinatantimonas, some *Vibrio*), propionibacteria, clostridia and yeasts (Arroyo-López et
al., 2012, 2021; Ballester et al., 2021; de Castro et al., 2018; Penland et al., 2021). We have
recently reviewed metataxonomic studies on table olives and found that a core microbiota
of approximately 30 genera can be identified, with wide variations in the composition
among trade preparations and varieties (Ricciardi et al., 2025).
With a few exceptions (Penland et al., 2020, 2021, 2022) studies on table olives microbiota
are relatively small (<50 samples) and often focus on a limited number of varieties (see
reviews by Ricciardi et al., 2025; Tsoungos et al., 2023 for details). This, and the variety of
sampling plans and experimental approaches used, complicates the interpretation and the

integration of these data to obtain a comprehensive view of the factors affecting table olive
microbial community assembly and dynamics. The effect of trade preparation on the
structure and dynamics of bacterial and/or fungal communities has been evaluated for a few
varieties (Cocolin et al., 2013; Maoloni et al., 2022; Ruiz-Barba et al., 2023a, 2023b; Zinno et
al., 2017), and there is evidence that identical varieties produced in different geographic
locations (Kamilari et al., 2023; Kazou et al., 2020) or even within the same geographical
region (Lucena-Padrós and Ruiz-Barba, 2019), or in different years by the same producer
(Penland et al. 2020) have measurable differences in microbial community composition,
although subsets of core taxa can be identified. This information, together with source
tracking studies may assist in understanding the relative importance of dispersal and
selection in the assembly of table olive microbial communities and provide new avenues for
the selection of microbiome – based starters (Nikoloudaki et al., 2024).
Even with all the complexity and variability in table olive microbiomes, a recent study
(López-García et al., 2025), leveraging on a large combined dataset and on machine learning
approaches, has been successful in discriminating table olives bacterial profiles based on
olive processing type, cultivar, country of origin, and isolation matrix. This opens the
possibility that the composition of microbiota can be used in table olive authenticity studies
and help discriminating PDO varieties from their non-PDO counterparts.
This motivated us to carry out a large survey of table olives belonging to a wide array of
varieties and trade preparations from 6 countries (Italy, Greece, Spain, Cyprus, Peru, Egypt)
with a focus on Italian PDO and non-PDO varieties, and including both sound and spoiled
samples with the following objectives:
a. obtaining a comprehensive view of the bacterial and fungal microbial communities of
table olives, and understanding the variability due to systematic effects, like olive
variety, ripeness and trade preparation, and the extent of random variability
between and within producing companies.
b. using black natural Itrana olives as a case study to evaluate if detectable differences
between PDO and non-PDO products exist
143
144 **2. Materials and methods**
145
2.1 Sampling.
Table olive samples (n = 340) were obtained from 40 producers across 6 countries (Italy,
Greece, Spain, Cyprus, Peru, Egypt). Producers were required to deliver olives ready for
packaging in their fermentation brine, , and to fill a questionnaire reporting essential
metadata (location of the company, olive ripeness and variety, the techniques used for
transformation, addition of starter cultures, if any; spoilage; duration of fermentation and of
storage post fermentation, if applicable). None of the samples used in this study had
undergone a heat treatment. Twenty sample, for which the fermentation brine was not
available, were dispatched in the final packages, with the storage brine. Furthermore, 23
metagenomic DNA from table olives were obtained from Instituto de la Grasa, CSIC (Seville,
Spain) and used for amplicon targeted metagenomics, in order to increase coverage of
Spanish varieties. Summary data are shown in Table 1, while detailed data are shown in
Supplementary Table 1. All samples were dispatched in insulated containers with ice packs
and reached our laboratory within 48 (samples from Italy) to 72 h. They were stored at 4°C
for no more than 24 h before being used for microbiological analyses and extraction of DNA.

Supernatants (12,000 x g) of the brine were stored at -20°C for 7-30 gg before the
measurement of pH, titratable acidity and chlorides.

Table 1. The samples used in this project, by olive ripeness and trade preparation.

olive ripeness	olive trade preparation	n
Green olives	Alkali treated olives	64
Green olives	Natural olives	61
Green olives	Olives darkened by oxidation	4
Green olives	Specialties (Picholine, etc.)	30
Olives turning color	Natural olives	41
Olives turning color	Olives darkened by oxidation	7
Black olives	Dehydrated olives	4
Black olives	Natural olives	152

163
2.2 Chemical and physicochemical analyses.
pH and chlorides were measured in brines except for samples of dehydrated olives; the
latter were homogenized (1:4) with deionized water in a Stomacher Lab Blender for 1 min
using a filter bag, and pH and chlorides were measured in the filtered liquid phase.
pH was measured using a digital pH meter (CyberScan pH 110, Eutech Instrument Pte Ltd.,
Thermo Fisher Scientific, Monza, Italy) with a combination electrode (Double Pore Slim
electrode, Hamilton Company, Reno, Nevada, USA). NaCl concentration was estimated from
the measurement of chlorides on diluted (D = 10-50) brines using a selective electrode
(Crison ISE Chlorid 51340400) with a reference electrode (Crison 5241, KNO_3 1 mol/L as a
reference electrolyte), connected to an Orion 420 A plus Ion-analyzer. 5 mol/L $NaNO_3$ was
used as ionic strength adjuster (1 ml / 50 mL diluted brine) and a calibration curve (0.001-0.1
mol/L NaCl). Titratable acidity was only measured in brines using phenophthalein (Sigma) as
an indicator using NaOH 0.1 mol/L and a digital burette (Brand Titrette, Brand, Wertheim,
Germany).
Two technical replicates were used for each measurement.
179
180

2.3 Enumeration of microbial groups, isolation and identification.
Counts were performed on a 1:1 mixture of olives (pits were aseptically removed) and brine
after homogenization (2 min) in a filtered bag (Nasco Whirlpak, Pleasant Prairie, WI) using a
Stomacher Lab Blender 400 (Cole-Parmer, now Antylia Scientific, Vernon Hills, IL, USA) in
peptone water (Bacteriological peptone 0.1% w/v, NaCl 0.85% w/v).
Serial ten-fold dilutions were performed in peptone water. Lactic acid bacteria were
enumerated in modified MRS agar (Lee and Lee, 2008; Ricciardi et al., 2014) with 0.02%
sodium azide (mMRS+NA, 25°C, 72 h), yeasts and moulds on Glucose Yeast Extract agar with
100 mg/L chloramphenicol (GYEAC, 25°C, 5 days), halophilic microorganisms on Plate Count
Agar Standard with 5% NaCl (w/v) (PCA + NaCl, 30°C, 48 h), contaminants on *Pseudomonas*
agar base (PAB, 25°C, 48 h), and *Enterobacteriaceae* in Violet Red Bile Glucose Agar (VRBGA,
30°C, 24 h). All microbiological media were inoculated using a spiral plater (WASP Spiral
Plater, bioMérieux Italia SpA, Bagno a Ripoli, Italy) with the exception of VRBGA, for which
conventional pour plating was used. Colony counts were performed using a digital colony
counter (EasyCount 2, bioMérieux Italia).
Colonies (1-2 from at least one counting plate) were isolated at random and streaked for
purification on the same medium and, after control of cell morphology by light microscopy,
identification was carried out with MALDI-ToF Biotyper Sirius (Bruker Co, Mallerica, MA,
USA) using the "Extended Direct Transfer" SOP provided by the producer.
200
201 *2.4 Microbiological media, ingredients and reagents.*
202 All microbiological media and ingredients were obtained from Oxoid (Basingstoke, United
203 Kingdom) and reagents from Sigma-Aldrich (Merck KGaA, Darmstadt, Germany).
204
2.5 Amplicon targeted metagenomics.
Metagenomic DNA was extracted from both olives and brines (when applicable) in an
attempt to recover both planktonic microorganisms and microorganisms attached to olive
surfaces, while minimizing contamination with fruit DNA. Olives (40 g) were mixed with an
equal volume of brine (for dehydrated olives, sterile distilled water) to which sterile Tween
80 (final concentration 0.1% vol/vol) was aseptically added. After shaking for 10 min on a
rotary shaker (100 rpm) in ice, the liquid was centrifuged (8,000xg for 10 min) and the pellet
was washed twice in sterile NaCl 0.85% and immediately frozen at -80°C. DNA extraction and
purification were performed using DNeasy PowerFood Microbial Kit (QIAGEN Italia, Milan,
Italy) followed by quantitation using NanoDrop One spectrophotometer (Thermo-Fisher
Scientific, Milan, Italia). Library preparation and sequencing was performed by Novogene
(Cambridge, UK). The V3-V4 region of the 16S rRNA gene was amplified using Novogene's
proprietary primers Bakt_341F_Nov (5'-CCTAYGGGRBGCASCAG-3') and Bakt_805R_Nov (5'-
GGACTACNNGGGTATCTAAT-3'). The ITS2 region was amplified using primers ITS3F (5'-
GCATCGATGAAGAACGCAGC-3') and ITS4R (5'-TCCTCCGCTTATTGATATGC-3') for Fungi (White
et al., 1990). Paired end sequencing (2x250 bp) was performed on Illumina NovaSeq X.
Raw sequences were processed using a pipeline based on DADA2 (Callahan et al., 2016)
using R (R Core Team, 2025). Model scripts are available at
https://github.com/ep142/FoodMicrobionet/tree/master/dada2_pipeline. Taxonomic
assignment was carried out using functions assignTaxonomy() and addSpecies(), which
implement the naïve Bayeasian classifier (Wang et al., 2007), with SILVA 138.2 (Quast et al.,
2013) for bacteria and UNITE v10 general release 2024-04-04 for fungi (Abarenkov et al.,
2023) as taxonomic reference databases. The results were assembled in a phyloseq object

(McMurdie and Holmes, 2013) which was then used for further analysis. Extraction blanks
and a mock community (ZymoBIOMICS Microbial Community DNA Standard, Zymo
Research, Invine, CA, USA) were also included. Contaminating ASVs were removed using
package "decontam" (Davis et al., 2018). Non-target sequences were removed prior to
further statistical processing using phyloseq::subset_taxa().
233
2.6 Statistical analysis
Statistical analyses and graphs were produced with scripts combining the functions of the
packages stats (R Core Team, 2025), PCAtools (Blighe and Lun, 2025), phyloseq (McMurdie
and Holmes, 2013), ggplot2 (Wickham, 2016), ggpubr (Kassambara, 2025) and rstatix
(Kassambara, 2023). Non-parametric tests (Wilcoxon-Signed Rank Sum test) were used for
comparing distributions of physico-chemical and chemical variables and counts (as log(cfu/g
brine and olives)). Holm correction for multiple testing was used. Alpha diversity analyses,
ordinations and inferential analyses on microbial community data were performed using
functions of the packages phyloseq (McMurdie and Holmes, 2013), DESeq2 (Love et al.,
2014) and vegan (Oksanen et al., 2025). Heatmaps were generated using package
ComplexHeatmap (Gu, 2022).
To account for unequal sequencing depth, rarefaction was performed before calculation of
alpha-diversity. The threshold for rarefaction (15,000 and 20,000 sequences for bacteria and
fungi, respectively) was chosen based on rarefaction curves (data not shown). Total Sum
Scaling was performed before ordinations and PERMANOVA.
249
250 3. **Results**.
251
3.1 Sampling.
A summary of the samples grouped according to ripeness and trade preparation is shown in
Supplementary Table 1. Most of the samples (natural olives, 44%, Spanish-style, alkali
treated Spanish-style olives, 19%) had undergone a fermentation step, while 27% had
undergone an alkali treatment. The latter included both Spanish-style olives and specialties
like Nocellara del Belice produced with the "Castelvetrano" method (Perpetuini et al., 2020).
The remaining samples were produced with other methods, including drying. Spoiled
samples were also made available by some producers (Supplementary Table 3).
260
3.2 pH, titratable acidity and salt content.
The pH of brines and olives in our study ranged from 3.7 to 8.1 (see Supplementary Table 4,
Supplementary Table 5 for summary statistics). Even within a single variety, the pH was
highly variable (see Supplementary Figure 1 and Supplementary Table 5 for ranges and
standard deviation values). The null hypothesis of no differences in pH distribution among
groups defined by combination of ripeness and trade preparation was rejected (Wilcoxon
rank-sum test, WSRT, with multiple-testing correction; $p < 0.001$). Black dehydrated olives
and specialties showed a higher median pH.
Results for titratable acidity (TA) of brines reflect those of pH with the exception of olives
darkened by oxidation, to which gluconate was added. A box plot of the distribution of TA
for the main varieties is shown in Supplementary Figure 2, while summary statistics are
reported in Supplementary Table 6: the titratable acidity ranged from 0.3 to 24 g lactic
acid/L, with significantly higher median values (WSRT) for natural olives and significantly
lower values for alkali treated olives and specialties in particular. Again, high variability was

observed even within the same variety as shown by the spread of the points in
Supplementary Figure 2. There was a weak (-0.58) but highly significant (p<0.001)
correlation between pH and the $\log_{10}$(TA).
Summary statistics for salt content in brines and olives are shown in Supplementary Tables 7
and 8 while a box plot for the main varieties is shown in Supplementary Figure 3. NaCl
concentrations ranged from 2.2 g/L to near saturation (246 g/L), whereas median values in
groups of samples sharing the same ripeness stage and trade preparation varied between 23
and 82 g/L. High variability was observed. For Oliva di Gaeta samples, which were the most
numerous, NaCl content of brines varied between 30 and 116 g/L. Dehydrated olives and
olives darkened by oxidation had significantly lower median NaCl values (WSRT).
285
3.3 Microbial counts.
Viable microbial counts were obtained using selective or elective media for undesirable
microorganisms (*Enterobacteriaceae*, contaminants on PAB) and potentially beneficial
microorganisms. Based on MALDI-ToF or molecular identification of isolates (Supplementary
Table 9) we found that both mMRS+NA and GYEAC were likely to be selective (all isolates
were LAB or yeasts, respectively), while PAB (Supplementary Figure 5) supported the growth
of yeasts (which formed pinpoint colonies) and PCA+NaCl supported the growth of both
yeast and lactic acid bacteria together with staphylococci and other halophiles. The latter is
therefore of limited use when counts of yeasts and LAB are high.

295
296
297 **Figure 1**. Distribution of viable counts of lactic acid bacteria in table olives used in this study.
298

Counts of *Enterobacteriaceae* are shown in Supplementary Figure 4. For most samples
counts were below the detection limit (<10 cfu/mL), while values > 100 cfu/g (brine + olives)
were found in several samples of naturally fermented or alkali treated olives.
Counts of LAB are shown in Figure 1 while summary statistics are shown in Supplementary
Tables 10 and 11. Several samples (30%) had viable counts below the detection limit of our
method (200 cfu/g), with 10^9 cfu/g detected as maximum counts. As for other variables,
variability within olive varieties was very high but some varieties showed significantly
(p<0.05) higher (WSRT) counts compared to others (for example, Bella di Daunia vs Halkidiki
for Green alkali treated olives; Kalamata and Konservolea vs Oliva di Gaeta for black
naturally fermented olives). High counts of LAB were also found in alkali treated olives which
are stored at low temperature after the alkali treatment, like Nocellara del Belice produced
with the Castelvetrano method (Alfonzo et al., 2024).
Summary results for yeast counts (moulds were only occasionally detected) are reported in
Supplementary Tables 12 and 13, and their distribution is shown in Figure 2: median values
for the groups defined by combinations of ripeness and trade preparation ranged between
5.02 and 6.85 log(cfu/g) and only 6% of the samples had counts below the detection limit of
the method while counts exceeding 10^8 cfu/g were observed thus confirming that yeasts are
among the main microbial groups in table olives. In fact, counts exceeding 10^5 cfu/g were
observed even in varieties which do not undergo fermentation, like Nocellara del Belice,
dried olives and olives darkened by oxidation.

319
320 **Figure 2.** Distribution of viable counts of yeast and moulds in table olives used in this study.
321
To explore the correlation structure among counts and chemical and chemico-physical
parameters, Pearson's correlation matrix was estimated and a correlation test with Holm's

correction for multiple testing was carried out to test the null hypothesis that correlation
coefficients were not significantly different from 0. All coefficients were <|0.5|, with the
highest absolute values for the correlation between counts of contaminants and halophilic
microorganisms (0.45) and pH and titratable acidity (-0.37) although many coefficients were
significantly different from 0. As a result of the general lack of correlation structure, the first
two components of a Principal Component Analysis carried out on the correlation matrix
explained only 43% of the variance, with no clear separation between different olive groups
(Supplementary Figure 6). This again, might be the result of the very high variability within
groups.
333
3.4 The composition of microbial communities of table olives.
3.4.1 Alpha-diversity of microbial communities.
Extraction of DNA proved to be challenging for some low biomass samples, but we were
able to carry out sequencing of the V3-V4 region of the 16S rRNA gene (336 samples) and of
the ITS2 region (337 samples) of most of the samples. Mock communities (2) and blanks (3)
were included for quality control. The number of sequences after QC filtering, ASV inference
and chimera removal was 17.2×10^6 for bacteria and 17.0×10^6 for fungi. 90% of the samples
had 29.3×10^3 sequences or more for bacteria, while the 90th percentile for fungi was 34.9
$\times 10^3$ sequences. The ASV abundance tables for both bacteria and fungi, after removal of
contaminants and non-target sequences, were highly sparse (99.3 and 99.6% for bacteria
and fungi, respectively). To avoid massive loss of rare taxa during prevalence and abundance
filtering and loss of power in inferential tests we decide to perform taxonomic
agglomeration prior to statistical analysis rather than operating on ASVs.
Our pipeline for 16S rRNA sequences theoretically allowed taxonomic assignment down to
the species level. However, this is not always possible for *Lactobacillaceae*, even when using
a custom database (Parente et al., 2023). Therefore, alpha and beta-diversity analysis for
bacterial communities was performed after agglomeration at the genus level. Chao1 values
calculated on the bacterial communities are shown in Figure 3, while summary statistics are
reported in Supplementary Tables 14 and 15.
Chao1, estimated using the estimate_richness() function of the phyloseq package after
rarefaction, ranged from 3 to 21 (with median values for different groups ranging from 16 to
45). A large variability existed within each group and even within the same variety but some
trends were evident. There was a trend in the decrease of diversity from Alkali treated
olives, to natural olives, to olives darkened by oxidation, to specialties. We decided to
investigate if between-firm and within-firm variability was a contributing factor. Distribution
of Chao1 is shown for Itrana, Bella di Daunia and Halkidiki in Supplementary Figures 7, 8 and
9. Within the same variety the diversity of bacterial communities varied between different
producers. We used Spearman's ρ to evaluate correlations between diversity, physical-
chemical parameters, and microbial counts for specific ripeness and trade preparation
combinations (Supplementary Table 16). With the exception of a strong negative correlation
between pH and diversity in Specialties - where diversity increased as pH decreased, likely
due to activity of LAB - all other correlations were low (|ρ| < 0.40). While significantly
different from 0, the remaining correlations often differed in sign across different trade
preparations, indicating that the effect of factors like acidification or NaCl concentration on
Chao1 differed in different groups.

369
370 **Figure 3.** Alpha-diversity (Chao1) of bacterial communities in table olives analysed in this
371 study.
372
Since our pipeline allowed assignment at the species level for the vast majority of sequences
(69.4% of the ASVs, 97.7% of the sequences), diversity indices for fungi were estimated
before agglomeration at the genus level. The distribution of Chao1 values is shown in Figure
4, while summary statistics are in Supplementary Tables 17 and 18.

Figure 4. Alpha-diversity (Chao1) of fungal communities in table olives used in this study.

Chao1 ranged from 1 to 209 (with median values for different groups ranging from 9 to 41,
lower than those for bacteria for most groups). A large variability existed within each group
and even within the same variety, but some trends were evident: the median value for
Chao1 decreased from Specialties to alkali treated olives, to natural olives (Figure 4).
Significant differences among the distributions were found using WSRT for all the
comparisons between alkali treated olives (including specialties) and natural olives (p<0.02).
We investigated if between-firm and within-firm variability was a contributing factor. As for
bacterial communities, distribution of Chao1 values within the same variety (Itrana olives,
Bella di Daunia and Halkidiki olives: Supplementary Figures 10, 11 and 12) varied among
different producers.
We also evaluated the correlation (Spearman ρ) between Chao1 and selected variables
within groups of ripeness and trade preparation (Supplementary Table 19): a rather strong
negative correlation was found between Chao1 and yeasts and moulds counts and duration
of fermentation for most groups, suggesting that, as the fermentation duration increases,
diversity of fungi decreases. Again, differences exist among varieties, possibly because of
systematic combinations of specific factors affecting dispersal and selection: in fact, in Oliva
di Gaeta, there was a weak but significant positive correlation (0.35) between pH and Chao1;
as for bacteria, higher pH is associated with higher diversity.

Figure 5. Distribution of the relative abundance of LAB and HALAB in table olives. Genera are ordered by alphabetical order within family.

After prevalence and abundance filtering (> 1% maximum abundance and >1% prevalence), >99.8% of the sequences were preserved for bacteria (Supplementary Figure 13). The large number of genera retained highlights the high microbial diversity found in table olives. The top 24 bacterial taxa on the basis of abundance and prevalence and are shown in Supplementary Table 20. The distribution of the abundance of LAB and HALAB is shown in Figure 5. Although abundances of the main genera are highly variable within each olive group, *Alkalibacterium* and *Marinilactibacillus* are abundant and prevalent in alkali treated olives, while a diverse set of *Lactobacillaceae*, among which *Lactiplantibacillus* is clearly both prevalent and abundant, dominate naturally fermented olives communities. Several heterofermentative LAB are often (*Lentilactobacillus*) or occasionally *(Limosilactobacillus, Levilactobacillus, Paucilactobacillus* and *Leuconostoc*) dominant or subdominant. It is interesting to note that, even in samples where *Lactiplantibacillus* was not dominant, *Lactiplantibacillus pentosus* was the most frequently isolated species (Supplementary Table 9).

The distribution of non-LAB halophilic genera is shown in Supplementary Figure 14. *Celerinatantimonas* was the most prevalent and abundant halophilic *Pseudomonadota*, but other Gram-negatives, such as *Halomonas, Vibrio, Salinicola* and *Idiomarina* were prevalent and occasionally abundant.

In a recent review we have found that DNA of genera including pathogenic bacteria can be found, albeit at very low abundance, in table olives (Ricciardi et al., 2025b). We repeated the

analysis for this survey and found that only in very few samples DNA sequences assigned to
genera *Salmonella, Yersinia* and *Listeria* were detectable, usually at very low abundance,
with one notable exception (Supplementary Figure 15).
Bar plots of the relative abundances of the 24 most abundant and prevalent genera are
shown in Supplementary Figures 16 (pooled data for all varieties, by ripeness and trade
preparation group) and 17 to 21 (main varieties within ripeness and trade preparation
groups). Some general patterns are evident: HALAB were more frequent and abundant in
alkali treated olives, while LAB, mainly *Lentilactobacillus, Lactiplantibacillus* and *Pediococcus*
were dominating members in naturally fermented olives. However, variety-specific patterns
appear. Two examples, for Bella di Cerignola (used for different types of Bella di Daunia PDO)
and Itrana (used for the production of two non-PDO types and of Oliva di Gaeta PDO) are
shown in Supplementary Figures 22 and 23. In both cases, while there is some similarity in
the composition of the bacterial community within a given variety, differences between
firms are visually evident.
Abundance and prevalence filtering was performed on fungal communities using
approaches similar to those described above. Our pipeline allowed the assignment at the
species level for the vast majority of ASVs, but given the debate on the reliability of species
assignment using regions of the ribosomal RNA operon in fungi (Kauserud, 2023; Rué et al.,
2023; Tedersoo et al., 2022) we decided to perform this analysis both at the species and
genus levels. Abundance and prevalence filtering at the species level retained 195 taxa out
of 795, with 99.6% of sequences, mostly belonging to phylum Ascomycota (Supplementary
Figure 24). Twenty-nine taxa had a prevalence > 0.1 and a maximum abundance > 0.01 and
may be considered as the core mycobiota of table olives (Supplementary Table 10). Six
species (*Pichia membranifaciens, P. manshurica, Candida boidinii, Dekkera custersiana,*
Wickerhamomyces anomalus, Saccharomyces bayanus) have a prevalence >0.5 and a
maximum relative abundance >0.90, but with one exception, their median abundance is
<0.01: in fact, they may dominate some samples but may have a low abundance in others
(see Supplementary Figure 25 for a boxplot of the distribution of core species).
The most abundant species were all cultivable, and were in fact isolated from counting
plates (Supplementary Table 9).
To simplify further descriptive analysis, we performed aggregation at the genus level. Core
fungal genera are shown in Supplementary Figure 26 and, remarkably, they largely overlap
with those found in a recent review (Ricciardi et al., 2025). To avoid losing genera which are
important only in one group of olives we used the same approach used for bacterial
communities (the union of the sets of taxa which passed the filter in each group was
retained): 107 genera were retained. Figure 6 shows a box plot of the most important
families (*Pichiaceae, Saccharomycetales_fam_Incertae_sedis* and *Phaffomycetaceae*), while
other genera are shown in Supplementary Figures 27 and 28.
As for bacterial genera there is a very wide variability within each ripeness and trade
preparation group and sometimes even within the same variety. However, some patterns
appear to emerge: *Pichia* is by far the most abundant and prevalent genus in all types of
olives which have undergone a fermentation (including green alkali treated olives) while the
distribution of other important genera is more variable and often clearly multimodal, with
very high abundance in some samples and very low in others. Specialties clearly have the
largest variety of fungal genera, possibly due to the lower duration of fermentation/storage
and the less selective conditions.
469

Figure 6. Distribution of fungal genera belonging to families *Pichiaceae, Saccharomycetales fam Incertae sedis* and *Phaffomycetaceae* in table olives used in this study. The main varieties are shown (see Supplementary Table 2 for abbreviations).

A summary bar plot with the relative abundance of the main fungal genera in ripeness/trade preparation groups is shown in Supplementary Figure 29. Overall, *Pichia* dominated most of the groups, and the sum of relative abundance of *Pichia, Candida, Wickerhamomyces, Saccharomyces, Zygotorulaspora* and *Dekkera* usually exceeded 75%. On the other hand, there are notable differences among varieties within each group (Supplementary Figures 30-34). The high fungal diversity in specialties, even within the same firm (most samples of Nocellara del Belice were provided by one firm over two years) is evident from Supplementary Figure 33. While the composition of fungal communities of samples of the same combination of olive variety and trade preparation is remarkably similar, samples from some firms have a substantially different composition (Supplementary Figures 35-36).

3.4.2 Beta-diversity of bacterial and fungal communities.
In order to compare the microbiota across samples and varieties, we used a four-step analytical approach. First, we calculated a Bray-Curtis distance matrix, which is a statistical measure used to quantify how much two microbial communities differ in both the types and the abundance of taxa present. Second, we visualized these relationships using non-monotonic Multidimensional Scaling (nMDS). This technique creates a spatial map where the distance between points represents their biological similarity; samples that cluster together have more similar microbiota. Third, we evaluated the homogeneity of group

variances (beta-dispersion) and applied Permutational Multivariate Analysis of Variance
(PERMANOVA) to test whether the differences between ripeness stages and trade
preparations were statistically significant or simply due to random variation.
Finally, we carried out differential abundance analysis with DESeq2 to identify specific taxa
whose abundance was significantly different is pairwise comparisons of ripeness/trade
preparation groups.

500
501   **Figure 7.** Configuration plot for the first two dimensions of a non-monotonic
502   Multidimensional Scaling of the Bray-Curtis distance matrix of the composition of bacterial
503   communities of table olives. See Supplementary Table 2 for abbreviations of olive varieties.
504
nMDS of bacterial communities in three dimensions resulted in a low stress (0.13), thus
explaining a high proportion of the variance, and the configuration for the first two
dimensions is shown in Figure 7. The variability within each ripeness/trade preparation
group is evident: ripeness/trade preparation groups tend to occupy a specific area of the
graph but there was an overlap between cluster of samples belonging to each group. No
clear pattern for the clustering of spoiled samples (which were often, but not always, far
from non-spoiled samples) is evident, but this may be due to the variety of spoilage types in
our dataset. The between-firms and within-firms variability even within a single olive variety
was substantial: selected examples are shown in Supplementary Figures 37-39.
However, inferential methods (PERMANOVA with adonis2) showed that ripeness and trade
preparation significantly (0.001) affected the composition of the bacterial communities. We
further examined if varieties within a specific ripeness/trade preparation group or producers
within a specific olive variety influenced the bacterial composition. Using PERMANOVA, we
found that variety within groups significantly affected the composition of bacterial
communities, except for 'turning color' olives that were naturally fermented (Supplementary

Table 22). The producing firm was a significant factor shaping the microbiota for all varieties
tested, including Itrana nera, Kalamata, Bella di Daunia and Halkidiki (Supplementary Table
23). These results indicate that house microbiota or a producer's specific practices can
create distinct microbial profiles even within the same variety. Statistical testing confirmed
that the observed microbial shifts were primarily caused by true differences in community
composition (group centroids) rather than simply reflecting higher variability within certain
groups (beta-dispersion), with the exception of the comparison for varieties within green
natural olives and firms within Kalamata olives. We also used PERMANOVA to test the null
hypothesis of lack of differences in the structure of bacterial communities for Itrana nera
and Oliva di Gaeta (its PDO counterpart). Olives of the two varieties were provided by 6
different firms, one of which, the largest in the PDO area, F23, provided 90% of the samples
for these varieties. When only samples from F23 were used (n = 80) the null hypothesis was
rejected with p = 0.025. When all firms were included, significant differences were still found
but, due to unbalanced sampling and between and within firm variability, further sampling
and the use supervised machine learning approaches are needed to confirm that differences
in microbiota can indeed be used to differentiate PDO from non PDO Itrana olives.
Finally, we used differential abundance analysis (DESeq2 with alpha = 0.01) to pinpoint the
specific taxa driving the differences between groups. The results are shown in
Supplementary Figures 40 to 47 and in Supplementary Table 24 and summarized in the
heatmap presented in Figure 8. This approach allowed us to detect significant microbial
signatures regardless of their overall rank in the community; indeed, many taxa with low
total abundance were identified as key biological indicators because they exhibited large
effect sizes (high log2 fold-changes) between preparations.
The analysis revealed distinct microbial profiles for selected trade preparation:
Alkali-treated olives were primarily distinguished by a significant enrichment of HALAB and
other salt-tolerant and alkaliphilic microorganisms (*Enterococcus, Alkalibacterium,*
Marinilactobacillus, Halolactibacillus, Natronobacter, Alkalibacterium). Some LAB
(*Lentilactobacillus, Pediococcus, Secundilactobacillus*) were characteristically less abundant
specialties and enriched in black natural olives. The latter showed a distinct prevalence of
Pseudomonadota, including deleterious microorganisms like *Enterobacteriaceae*, and a
diverse array of *Lactobacillaceae*. Specifically, heterofermentative genera—including
Levilactobacillus, Leuconostoc, and *Weissella*—were significantly more abundant in black
natural olives compared to all green varieties.
We observed a clear split in the *Lactobacillaceae* family based on fruit ripeness;
Lactiplantibacillus is significantly more abundant in black natural olives than in alkali treated
green olives (including specialties), but *Lacticaseibacillus* is more abundant in Green alkali
treated olives compared to Black natural.
The same approach was used for fungal communities. We first visualized community
relationships using non-constrained ordination (nMDS) (Figure 9). Most ripeness/trade
preparation groups exhibited overlapping, suggesting shared fungal community
characteristics. By examining preparations separately (Supplementary Figures 48-50), we
noted that composition of fungal is often highly variable within-variety. This was especially
true for Oliva di Gaeta (Supplementary Figure 48). Although spoiled samples frequently
appeared distinct within a given producer, our limited sample size for defective olives
prevented definitive statistical conclusions regarding spoilage markers.

566
Figure 8. Differential abundance of bacterial genera across olive ripeness and trade preparation groups. Heatmap showing Z-score transformed
variance-stabilized counts (VST) of all bacterial genera significantly differentially abundant (DESeq2, adjusted p-value < 0.01) in at least one
pairwise contrast between olive groups. Columns represent individual samples grouped by ripeness/trade preparation (Green natural, Green
specialties, Green alkali treated, Turning color natural, Black natural, Ripe trade) and ordered by olive variety within each group. Rows
represent bacterial genera and are ordered by Spearman correlation-based hierarchical clustering. Row annotation bars (right) show the
taxonomic order of each genus, the mean log10 relative abundance across all samples, and the log2 fold change for each pairwise contrast;
grey cells in the log2 fold change bars indicate that the genus was not significantly differentially abundant in that contrast. Heatmap colors
represent Z-scores capped at ±2 for visualization purposes; values exceeding this range are shown in the most extreme color.

As for bacteria, PERMANOVA confirmed that the combination of ripeness and trade
preparation significantly shaped composition of fungal community (p = 0.001), as did the
olive variety within the main groups (Supplementary Table 25).
However, in the test of effect of ripeness and trade preparation inhomogeneity of dispersion
is a contributing factor, largely due to the high variance in Green and Black natural olives
(see also Figure 9).

581
582 **Figure 9**. Configuration plot for the first two dimensions of a non-monotonic
583 Multidimensional Scaling of the Bray-Curtis distance matrix of the composition of fungal
584 communities of table olives. See Supplementary Table 2 for abbreviations of olive varieties.
585
The influence of the producer was particularly prominent. For Kalamata, Itrana nera, and
Halkidiki, the producing firm was a significant factor (Supplementary Table 26). This indicates
that house microbiota and local practices are major drivers in determining which fungi
dominate the fermentation.
As for bacteria significant differences between the fungal communities of black Itrana
varieties (non-PDO Itrana nera and PDO Oliva di Gaeta) were found both when all firms were
included and when only firm F23 was used (p = 0.002) thus showing that even within the
same firm, systematic differences in microbiota are due to starting material and production
practices.
We then carried out differential abundance analysis using DeSeq2 with alpha = 0.01. The
results are shown in Supplementary Figures 50 to 57 and in Supplementary Table 27 and
summarized in the heatmap shown in Figure 10.
Even with all the variability within each group, a pattern is evident. A first group of genera
(*Saturnispora, Gibellulopsis, Foliophoma, Acremonium, Chytridiomycota incertae sedis, Fungi*
gen incertae sedis, Alternaria, Vishniacozyma, Cladosporium, Aureobasidium and *Fusarium*),
whose origin is likely the carposphere or the environment and whose average abundance is
characteristically low, were enriched in Green specialties, in which fermentation is slowed or
prevented by low temperature.

605
606
Figure 10. Differential abundance of bacterial genera across olive ripeness and trade preparation groups. Heatmap showing Z-score
transformed variance-stabilized counts (VST) of all bacterial genera significantly differentially abundant (DESeq2, adjusted p-value < 0.01) in at
least one pairwise contrast between olive groups. Columns represent individual samples grouped by ripeness/trade preparation (Green
natural, Green specialties, Green alkali treated, Turning color natural, Black natural, Ripe trade) and ordered by olive variety within each group.
Rows represent bacterial genera and are ordered by Spearman correlation-based hierarchical clustering. Row annotation bars (right) show the
taxonomic order of each genus, the mean log10 relative abundance across all samples, and the log2 fold change for each pairwise contrast;
grey cells in the log2 fold change bars indicate that the genus was not significantly differentially abundant in that contrast. Heatmap colors
represent Z-scores capped at ±2 for visualization purposes; values exceeding this range are shown in the most extreme color.

615
A second group, which included both abundant and prevalent genera (*Trichosporon,*
Kodamea, Meyerozyma, Kluyveromyces, Geotrichum, Debaryomyces, Wickerhamomyces,
Schwanniomyces) was significantly more abundant in one or more alkali treated varieties
compared to natural olives. The genera *Schwanniomyces* (which is abundant in Halkidiki
olives) and *Starmerella* (which is abundant in Halkidiki, Konservolea and Kalamata) stand out
becaue of their association with specific varieties, and this may explain its pattern in some
contrasts. The last group includes genera whose pattern is more variable. *Pichia,* the most
abundant and prevalent genus, is clearly less abundant in specialties but its abundance is
variable in other groups. Significant differences may be obscured by the agglomeration at
the genus level, since the pattern of abundance of the species within this genus varies in
different groups (see Supplementary Figure 25). *Dekkera* was found to be more abundant in
black and turning color natural compared to specialties. *Citeromyces* was more abundant in
all types of naturally fermented olives compared to green alkali treated olives. The
abundance of the genus *Candida* was lower on average in black natural compared to both
green alkali treated and green specialties: however, this may not be true for all varieties and
all species, since *C. boidinii* was very abundant in some samples of black natural olives (see
Supplementaty Figure 25).
633
634 4. **Discussion**.
635
This work presents the largest survey to date of the bacterial and fungal communities of
table olives, with a particular focus on Italian varieties. Among these, Oliva di Gaeta —
produced from naturally fermented black Itrana olives and one of the most commercially
important Italian table olive varieties — is described here for the first time using culture-
independent approaches, thus extending significantly previously available findings based on
culture dependent approaches (Sacchi et al., 2022; Servili et al., 2006; Tofalo et al., 2012).
For several other Italian PDO and non-PDO varieties, including Bella di Daunia, Ascolana del
Piceno, Nocellara Etnea, Nocellara del Belìce and Tonda di Cagliari, prior metataxonomic
data were limited to small, single-producer studies (Campus et al., 2025; Cocolin et al., 2013;
De Angelis et al., 2015; Maoloni et al., 2022; Randazzo et al., 2017; Zinno et al., 2017), while
for many other minor varieties (see Supplementary Table 2) this is the first study reporting
data con bacterial and fungal communities. The dataset also includes substantial
representation of Greek (Halkidiki, Kalamata, Conservolea) and Spanish (Manzanilla, Aloreña
and others) varieties, providing a broader comparative framework and enabling, for the first
time, a direct cross-variety analysis at this scale. A key finding emerging from this dataset is
the striking variability in microbiota composition observed not only across varieties, but
within the same variety and even within the same producer. This variability is consistent
with a strong role of stochastic colonization events, which can obscure the selective
pressures — salt concentration, ripeness, trade preparation — that might otherwise
generate variety-specific microbial signatures. This has direct implications for the
development of control strategies: harnessing the diversity of these communities through
microbiome-based starter cultures offers a promising avenue, provided that the rich
diversity of flavours and aromas that characterizes artisanal Italian table olives is preserved
rather than homogenized.
We confirmed that the composition of microbial communities in table olives is shaped by
physicochemical factors associated with olive ripeness, and trade preparation

(Anagnostopoulos and Tsaltas, 2022; López-García et al., 2025; Perpetuini et al., 2020),
which exert consistent selective pressures across producers and varieties, as confirmed by
our inferential analysis.
The strongest structuring factor in our dataset was the contrast between alkali-treated and
naturally fermented olives. PERMANOVA and differential abundance analysis confirmed that
the combination of ripeness and trade preparation significantly affected the composition of
both bacterial and fungal communities. Naturally fermented black olives were characterised
by a diverse array of *Lactobacillaceae* — particularly heterofermentative genera including
Levilactobacillus, Leuconostoc and *Weissella* — alongside a higher prevalence of
Pseudomonadota, including *Enterobacteriaceae*. The diversity of lactic and non-lactic
microbiota of natural olives and the widespread occurrence of *Enterobacteriaceae* and other
Pseudomonatota has been confirmed by several authors (see Portilha-Cunha et al., 2020;
Ricciardi et al., 2025; Tsoungos et al., 2023, for recent reviews). Among Pseudomonadota
Celerinatantimonas is especially notable because of its spoilage potential (de Castro et al.,
2022) and its very strong association with black natural olives (Kazou et al. 2020; Penland et
al., 2020, 2021), a pattern which is confirmed by our data (Supplementary Figure 14, Figure
8). Alkali-treated olives, by contrast, were distinguished by enrichment of Halophilic and
Alkalophilic Lactic Acid Bacteria (HALAB), including *Enterococcus, Alkalibacterium,*
Marinilactobacillus, and Halolactibacillus. Within alkali-treated olives, a further important
distinction emerged between Spanish-style olives, which undergo fermentation after lye
treatment, and Castelvetrano-style olives, in which low-temperature storage suppresses
fermentation (Alfonzo et al., 2024). The signature presence of these genera in alkali treated
olives has been confirmed by several authors (Arroyo-López et al., 2021; López-García et al.,
2022; Lucena-Padrós and Ruiz-Barba, 2016) and emerges as a very robust signal in a recent
study using machine-learning approaches (López-García et al., 2025), especially when these
are compared with directly brined counterparts of the same variety (Ruiz-Barba et al.,
2023bb). The latter were characterised by the most consistent dominance of HALAB and,
strikingly, by the highest fungal diversity in the dataset, with enrichment of fungal genera
which have been associated with the carposphere and the environment (Ferrocino et al.,
2025; Penland et al., 2021)— a pattern consistent with the less selective conditions
prevailing in the absence of fermentation-driven community succession, which in naturally
fermented olives progressively reduces fungal diversity as fermentation time increases.
These broad contrasts provide the interpretive framework within which finer-grained effects
— olive variety, producer identity, and stochastic colonization — should be understood.
However, even within the same variety and production process, significant differences
between producers were observed, again supported by inferential methods. This producer-
level variability is consistent with the existence of a house microbiota — a resident microbial
community associated with each producer's facility and equipment — whose composition
introduces a further layer of differentiation, a well-known phenomenon in spontaneous
fermentation processes (Bokulich et al., 2015; Comasio et al., 2020; Li et al., 2022), which
has also been confirmed in several studies on table olives (Kazou et al., 2020, Lucena-Padrós
and Ruiz Barba, 2019; Penland et al., 2021). The widespread use of small fermentation
vessels (240–280 L plastic fermenters at uncontrolled temperature) likely amplifies this
effect: in small, semi-closed systems, stochastic colonization events have an outsized
influence on the trajectory of each individual fermentation, producing divergent
communities even from nominally identical starting conditions.

Culture-based enumeration and culture-independent analysis provided complementary but
not always concordant pictures of the microbial communities of table olives. Among the
media tested, only mMRS+NA and GYEAC showed sufficient selectivity for LAB and yeasts,
respectively, while PCA+NaCl supported the growth of both groups and is therefore of
limited diagnostic value when their counts are high. LAB counts were highly variable and
frequently below the detection limit, consistent with the known decline of LAB over the
course of fermentation and their scarcity in high-salt brines (Benítez-Cabello et al., 2020;
Sánchez et al., 2018; Traina et al., 2024). Yeast counts were more consistently high across
varieties and trade preparations, including in varieties which do not undergo fermentation,
confirming the ubiquity of yeasts as a major microbial group in table olives
(Anagnostopoulos and Tsaltas, 2022; Arroyo-López et al., 2012; Giavalisco et al., 2024).
A noteworthy discrepancy emerged between culture-based and amplicon-based results for
LAB. *Lactiplantibacillus* was frequently recovered even in samples where it was not the
dominant taxon according to amplicon sequencing: only 29% of samples had a
Lactiplantibacillus fraction among *Lactobacillaceae* sequences exceeding 0.20, and only 7%
exceeded 0.80, yet *Lactiplantibacillus* was the sole genus isolated in 87% of samples. Colony
picking was performed randomly, so this discrepancy most likely reflects either a systematic
cultivability advantage of *Lactiplantibacillus* on mMRS+NA over other LAB, and relatively low
recovery of colonies from other highly abundant genera (*Leuconostoc, Pediococcus,*
Lentilactobacillus, among others) may reflect the fact that these genera were not viable or
able to grow at the time of enumeration and isolation; a visibility bias in colony picking due
to differences in colony morphology on this medium is less likely.
Enterobacteriaceae counts were below the detection limit in most samples, but values
exceeding 100 cfu/g were occasionally found in naturally fermented and alkali-treated olives.
Metataxonomic analysis revealed the presence of DNA assigned to several
Enterobacteriaceae genera at high relative abundance in some varieties even at the end of
fermentation; however, whether this represents viable cells, viable but non-culturable
forms, or residual DNA from dead cells cannot be determined from these data alone.
The same argument applies to the detection of genomic DNA of pathogenic or potentially
pathogenic bacteria. We confirmed our previous finding of the occurrence of sequences
assigned to bacterial genera containing pathogenic microorganisms (Ricciardi et al., 2025).
The occurrence of *Salmonella* at relative abundance close to 0.1% in at least two samples,
together with previous reports on outbreaks caused by *Salm. enterica* from olives (Donachie
et al., 2018) raises some concerns on the occurrence of this pathogen in olive producing
environments although we cannot infer the presence of viable pathogenic microorganisms
from this data. It is well known that the low abundance of pathogenic microorganisms in
foods and the low taxonomic resolution of amplicon targeted metagenomics pose
challenges to the use of metagenomic approaches in the detection of microorganisms
relevant for food safety (Ferrocino et al., 2022; Wang et al., 2025).
Bacterial and fungal alpha diversity, as measured by Chao1, were both highly variable within
and between groups, consistent with the high variability in community composition
documented throughout this study. Bacterial diversity showed a decreasing trend from
alkali-treated olives to naturally fermented olives to specialties, while the opposite trend
was observed for fungi — specialties harboured the highest fungal diversity, reflecting the
less selective conditions prevailing in the absence of prolonged fermentation. Fermentation-
driven acidification and the progressive dominance of competitive fermentative taxa reduce
fungal diversity over time, as supported by the significant negative correlation between

fungal Chao1 and fermentation duration observed across most groups. However, previous
findings on the effect of fermentation time on diversity of fungal communities are
contradictory (López-García et al., 2024; Ruiz-Barba et al., 2023a, 2023b).
Producer identity was a significant contributor to within-variety variability in diversity for
both bacteria and fungi, consistent with the house microbiota effect discussed above. Chao1
values for both bacteria and fungi were broadly comparable to those reported for similar
varieties and trade preparations in previous studies (Cocolin et al., 2013; De Angelis et al.,
2015; de Castro et al., 2018; Kamilari et al., 2023; Maoloni et al., 2022; Medina et al., 2016;
Penland et al., 2020, 2021), although differences in bioinformatic pipelines and diversity
indices limit direct comparisons.
Among LAB, *Lactiplantibacillus* was the most prevalent and abundant genus in naturally
fermented olives, consistent with its well-documented adaptation to the brine environment
and its widespread use as a starter and functional culture in table olives (Giavalisco et al.,
2024; Portilha-Cunha et al., 2020; Zotta et al., 2022). However, it was absent or
outnumbered by other genera in a substantial proportion of samples, underscoring that the
LAB ecology of naturally fermented olives is more complex than the dominant role assigned
to this genus in much of the starter literature. *Lacticaseibacillus*, which also has documented
starter and functional potential in table olives (Pino et al., 2019), was among the dominant
genera particularly in alkali-treated varieties — a pattern confirmed by our differential
abundance analysis — suggesting that starter development for alkali-treated olives should
more systematically consider this genus alongside *Lactiplantibacillus*. Several
heterofermentative LAB — particularly *Lentilactobacillus*, and occasionally
Limosilactobacillus, *Levilactobacillus*, *Paucilactobacillus* and *Leuconostoc* — were frequently
subdominant or dominant in naturally fermented olives. Their contribution to flavour
development and fermentation dynamics remains poorly understood, although
Levilactobacillus and *Leuconostoc* have been evaluated as starters in Greek olives (Chytiri et
al., 2020; Vougiouklaki et al., 2024). The frequent dominance of *Lentilactobacillus* is of
particular concern given its association with spoilage and biogenic amine production in
protein-rich environments (O'Sullivan et al., 2015; Penland et al., 2021; Suzzi and Gardini,
2003), and warrants systematic investigation of its role across olive varieties and trade
preparations. Finally, the occasional occurrence of HALAB in naturally fermented olives —
where alkaline and high-salt conditions are less sustained — is an unexpected finding whose
ecological and technological significance deserves further study.
The core mycobiota of table olives — dominated by *Pichia, Candida, Wickerhamomyces*, and
Saccharomyces — was largely consistent with previous surveys (Ricciardi et al., 2025;
Tsoungos et al., 2023), although their abundance was highly variable within and across
varieties, and the pattern of individual *Pichia* species differed between trade preparations
and varieties. Several findings extend or add to the existing literature.
Dekkera/Brettanomyces, which is frequently reported in table olives (Medina et al., 2018;
Michailidou et al., 2021; Tsoungos et al., 2023), was subdominant in several naturally
fermented varieties in our dataset, a pattern that warrants attention given its spoilage
potential in other fermented products (Tubia et al., 2018). *Zygotorulaspora*, whose role in
the aromatic profile of table olives has been documented (Arroyo-López et al., 2016;
Mougiou et al., 2023; Montaño et al., 2021) and which has been used as a starter in Nyons
olives (Penland et al., 2022), was occasionally dominant in our dataset, reinforcing its
relevance as a candidate for microbiome-based starter development. *Schwanniomyces*
showed a strong association with Halkidiki olives, extending a previous report (Argyri et al.,

2020), while *Starmerella* was consistently associated with Greek varieties (Conservolea,
Kalamata and, to a lesser extent, Halkidiki) — a finding not previously reported and
suggesting a variety- or origin-specific nature for this genus. *Debaryomyces,* which has been
evaluated as starter cultures (Chytiri et al., 2020, Tarantini et al., 2024), was also among the
genera identified in our dataset, further supporting the potential for diversifying yeast
starter selection beyond the currently dominant *Wickerhamomyces* and *Saccharomyces*
(Giavalisco et al., 2024). Together, these findings highlight the richness of the fungal
communities of table olives and their largely untapped potential for variety-specific starter
development.
The beta-diversity analysis confirmed, at a global level, the patterns already evident from
the differential abundance analysis: ripeness and trade preparation were the dominant
structuring factors, while olive variety within trade preparation groups and producer identity
within varieties introduced additional, statistically significant layers of differentiation. The
striking within-group variability visible in the ordination plots — even for samples of the
same variety from the same producer — reflects the combined influence of stochastic
colonization, small fermentation vessel effects, and house microbiota, as discussed above.
Technical replication confirmed that this variability is biological rather than methodological,
with Bray-Curtis distances between technical replicates being very low (data not shown).
Our dataset included spoiled samples across several varieties, and fungal communities of
spoiled samples were often visually distinct from those of unspoiled samples in the
ordination plots. However, the limited number of spoiled samples and the heterogeneity of
spoilage types prevented inferential analysis, and no firm conclusions on microbial markers
of spoilage can be drawn from these data, although other authors have identified consistent
spoilage markers for selected varieties (Arroyo-López et al., 2021; de Castro et al., 2018,
2022; Penland et al., 2021). This remains an important open question, particularly for fungal
communities where dissimilatiries between spoiled and non-spoiled samples suggests that
diagnostic signals may exist.
Several authors have proposed that metataxonomic analysis of microbial communities may
assist in identifying the geographical origin of table olives, invoking the concept of terroir —
originally developed for wine — to describe the influence of environment and raw material
on microbial community composition and, ultimately, organoleptic properties (Argyri et al.,
2020; Belda et al., 2017; Ferrocino et al., 2025; Kamilari et al., 2023; Kazou et al., 2020). Our
data are partially consistent with this view: producers of the same variety share common
microbial patterns, indicating that ecological factors associated with raw material and
production environment do shape community structure. However, significant and
measurable differences among producers within the same variety — confirmed by
PERMANOVA — indicate that house microbiota and producer-specific practices introduce
variation that cannot be attributed to terroir alone. This confounding between terroir and
house microbiota effects has important implications for authentication: producers within the
same PDO area can harbour more divergent microbial communities than PDO and non-PDO
producers of the same variety, making microbiota-based discrimination inherently
challenging. Indeed, while statistically significant differences in both bacterial and fungal
communities between PDO Oliva di Gaeta and its non-PDO counterpart Itrana nera were
detectable — both within the largest single producer and across all producers — the signal
was weak and heavily influenced by between-firm variability. Similar results were obtained
for Bella di Daunia and Nocellara del Belìce. These findings suggest that, with current
amplicon-based approaches and unbalanced sampling designs, Italian PDO table olive

varieties cannot be reliably discriminated from similar non-PDO varieties on the basis of
microbiota composition alone. Realising the potential of microbiota-based authentication
will require larger, balanced sampling designs, integration with metabolomic data
(Beteinakis et al., 2024), and the application of supervised machine learning approaches
(López-García et al., 2025) capable of extracting weak but consistent signals from highly
variable community data. The availability of large, well annotated databases, like OliveFMBN
(Ricciardi et al, 2025), in which the data of this survey have been integrated, may support
efforts in this field.
The high variability in microbial community composition documented in this study,
combined with the limited control that artisanal and semi-industrial producers currently
exert over fermentation conditions, provides a strong rationale for the development of
starter cultures tailored to specific olive varieties and trade preparations. The current starter
market for table olives is dominated by a small number of commercial producers, virtually all
of whom rely on *Lactiplantibacillus* as the primary — and often sole — inoculant. Examples
are the Lal'Olive Crispy *Lactipl. pentosus* OM13 (Alfonzo et al., 2023a; Vege-Start 10 (Corsetti
et al., 2012) and the well documented functional strain *Lactipl. pentosus* LPG1 (Benítez-
Cabello et al., 2020). While the fermentative, biopreservative, and functional potential of
this species in table olives is well established (Giavalisco et al., 2024; Portilha-Cunha et al.,
2020; Zotta et al., 2022), our data suggest that this approach captures only a fraction of the
ecologically and technologically relevant microbial diversity present in table olives. The
continued reliance on a narrow range of *Lactiplantibacillus*-based starters risks progressive
homogenization of the organoleptic diversity that currently characterises artisanal varieties
— a diversity that is itself a reflection of the complex, variety-specific microbial communities
documented here.
Microbiome-based starters offer a promising alternative (Nikoloudaki et al., 2024): rather
than relying on one or a few well-characterised strains, this approach uses microbiome data
to design complex starter communities that capture the diversity and ecological interactions
of the native microbiota of a specific variety and production context. Our findings provide
ecological foundations for this approach. For naturally fermented olives, candidate bacterial
starters should look beyond *Lactiplantibacillus* to include *Lacticaseibacillus*,
heterofermentative LAB such as *Levilactobacillus* and *Leuconostoc*. The role of HALAB is
more controversial since some of them have been associated with spoilage (Arroyo-López et
al., 2021). Fungal communities are an equally important: genera such as *Wickerhamomyces,*
Zygotorulaspora, and *Debaryomyces* have documented starter or flavour potential (Benítez-
Cabello et al., 2020; Giavalisco et al, 2024) and are consistent members of the mycobiota of
specific varieties. The development of mixed bacterial-fungal starters grounded in variety-
specific microbiome data represents a natural extension of this work, although a full
characterisation of microbial interactions and their functional consequences lies beyond the
scope of the present study.
Realising this potential will require not only larger and more targeted microbiome surveys,
but also functional studies linking community composition to fermentation dynamics and
organoleptic outcomes, and the development of competitive, variety-adapted strains
capable of driving fermentation consistently without suppressing the ecological diversity
that underpins the quality and identity of artisanal table olives.
893
5. Conclusions

This study presents the largest metataxonomic survey of table olive microbial communities
to date, providing the first culture-independent characterisation of several Italian PDO and
non-PDO varieties — most notably Oliva di Gaeta — and substantially extending existing
knowledge for others, including Bella di Daunia, Ascolana del Piceno, and Nocellara del
Belìce. The dominant structuring factors shaping both bacterial and fungal communities
were olive ripeness and trade preparation, with the contrast between alkali-treated and
naturally fermented olives representing the strongest and most consistent signal; within this
framework, variety- and producer-level effects introduced additional, statistically significant
differentiation consistent with the combined influence of ecological selection and house
microbiota. However, the striking variability observed within the same variety and even
within the same producer — driven by stochastic colonization events amplified by small-
scale fermentation conditions — obscures variety-specific microbial signatures and means
that, with current amplicon-based approaches, Italian PDO varieties cannot be reliably
discriminated from similar non-PDO counterparts on the basis of microbiota composition
alone, although integration with metabolomic data and supervised machine learning
approaches may improve authentication in the future. The ecological and taxonomic
complexity documented here — encompassing a diverse array of LAB, HALAB, halophilic
Gram-negatives, and fungal genera with documented starter and flavour potential —
provides the foundation for a new generation of variety- and preparation-specific
microbiome-based starter cultures that move beyond the current reliance on
Lactiplantibacillus-only commercial products; realising this potential will require functional
studies linking community composition to fermentation dynamics and organoleptic
outcomes, and the development of mixed bacterial-fungal starters grounded in variety-
specific microbiome data.
919

920 **Data availability**.
921 Sequence data have been deposited in NCBI SRA with bioproject accession PRJNA1292811.
922 The accession list of the runs used in this study is available in Supplementary data and
923 figures. The metataxonomic data have been integrated in the OliveFMBN database
924 (https://doi.org/10.5281/zenodo.13729346)
925

Acknowledgements.
The authors gratefully acknowledge the skilled technical assistance of Ms. Anita Aliano.
This work was carried out within the PRIN 2022 Project METAOlive 2022NN28ZZ and
received funding from the European Union Next-GenerationEU, CUP C53D23005460006
(PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2,
INVESTIMENTO 1.4 – D.D. 1048 14/07/2023).

6. References

Abarenkov, K., Nilsson, R.H., Larsson, K.-H., Taylor, A.F.S., May, T.W., Frøslev, T.G.,
Pawlowska, J., Lindahl, B., Põldmaa, K., Truong, C., Vu, D., Hosoya, T., Niskanen, T.,
Piirmann, T., Ivanov, F., Zirk, A., Peterson, M., Cheeke, T.E., Ishigami, Y., Jansson, A.T.,
Jeppesen, T.S., Kristiansson, E., Mikryukov, V., Miller, J.T., Oono, R., Ossandon, F.J.,
Paupério, J., Saar, I., Schigel, D., Suija, A., Tedersoo, L., Kõljalg, U., 2023. The UNITE
database for molecular identification and taxonomic communication of fungi and other

eukaryotes: sequences, taxa and classifications reconsidered. Nucleic Acids Res. 52,
D791–D797. https://doi.org/10.1093/nar/gkad1039.

Alfonzo, A., Alongi, D., Prestianni, R., Pirrone, A., Naselli, V., Viola, E., Pasquale, C.D., Croce,
F.L., Gaglio, R., Settanni, L., Francesca, N., Moschetti, G., 2024. Enhancing the quality and
safety of Nocellara del Belice green table olives produced using the Castelvetrano
method. Food Microbiol. 120, 104477. https://doi.org/10.1016/j.fm.2024.104477.

Alfonzo, A., Craparo, V., Prestianni, R., Naselli, V., Seminerio, V., Pirrone, A., Croce, F.L.,
Settanni, L., Moschetti, G., Francesca, N., 2023a. Application of technological protocols
on an industrial scale to improve Seville-style table olive production in Italy and Spain.
Appl. Food Res. 100349. https://doi.org/10.1016/j.afres.2023.100349.

Alfonzo, A., Naselli, V., Gaglio, R., Settanni, L., Corona, O., Croce, F.L., Vagnoli, P., Krieger-
Weber, S., Francesca, N., Moschetti, G., 2023b. Use of different nutrients to improve the
fermentation performances of *Lactiplantibacillus pentosus* OM13 during the production
of Sevillian Style Green table olives. Microorganisms 11, 825.
https://doi.org/10.3390/microorganisms11040825.

Anagnostopoulos, D.A., Tsaltas, D., 2022. Current status, recent advances, and main
challenges on table olive fermentation: the present meets the future. Front. Microbiol.
12, 797295. https://doi.org/10.3389/fmicb.2021.797295.

Argyri, K., Doulgeraki, A.I., Manthou, E., Grounta, A., Argyri, A.A., Nychas, G.-J.E., Tassou,
C.C., 2020. Microbial diversity of fermented greek table olives of Halkidiki and
Konservolia varieties from different regions as revealed by metagenomic analysis.
Microorganisms 8, 1241. https://doi.org/10.3390/microorganisms8081241.

Arroyo-López, F.N., Benítez-Cabello, A., Romero-Gil, V., Rodríguez-Gómez, F., Garrido-
Fernández, A., 2021. Delving into the bacterial diversity of spoiled green Manzanilla
Spanish-style table olive fermentations. Int. J. Food. Microbiol. 359, 109415.
https://doi.org/10.1016/j.ijfoodmicro.2021.109415.

Arroyo-López, F.N., Medina, E., Ruiz-Bellido, M.Á., Romero-Gil, V., Montes-Borrego, M.,
Landa, B.B., 2016. Enhancement of the knowledge on fungal communities in directly
brined Aloreña de Málaga green olive fermentations by metabarcoding analysis. PLoS
One 11, e0163135. https://doi.org/10.1371/journal.pone.0163135.

Arroyo-López, F.N., Romero-Gil, V., Bautista-Gallego, J., Rodriguez-Gómez, F., Jiménez-Díaz,
R., Garcia, P.G., Querol, A., Fernandez, A.G., 2012. Yeasts in table olive processing:
desirable or spoilage microorganisms? Int. J. Food Microbiol. 160, 42-49.
https://doi.org/10.1016/j.ijfoodmicro.2012.08.003.

Ballester, E., Ribes, S., Barat, J.M., Fuentes, A., 2021. Spoilage yeasts in fermented
vegetables: conventional and novel control strategies. Eur. Food Res. Technol. 1-14.
https://doi.org/10.1007/s00217-021-03888-7.

Belda, I., Zarraonaindia, I., Perisin, M., Palacios, A., Acedo, A., 2017. From vineyard soil to
wine fermentation: microbiome approximations to explain the "terroir" concept. Front.
Microbiol. 8, 821. https://doi.org/10.3389/fmicb.2017.00821.

Benítez-Cabello, A., Calero-Delgado, B., Rodríguez-Gómez, F., Bautista-Gallego, J., Garrido-
Fernández, A., Jiménez-Díaz, R., Arroyo-López, F.N., 2020. The use of multifunctional
yeast-lactobacilli starter cultures improves fermentation performance of Spanish-style
green table olives. Food Microbiol. 91, 103497.
https://doi.org/10.1016/j.fm.2020.103497.

Beteinakis, S., Papachristodoulou, A., Stathopoulos, P., Mikros, E., Halabalaki, M., 2024. A
multilevel LC-HRMS and NMR correlation workflow towards foodomics advancement:
application in table olives. Talanta 280, 126641.
https://doi.org/10.1016/j.talanta.2024.126641.

Blighe, K., Lun, A ., 2025. PCAtools: PCAtools: Everything Principal Components Analysis.
doi:10.18129/B9.bioc.PCAtools. https://doi.org/10.18129/B9.bioc.PCAtools.
Bokulich, N.A., Amiranashvili, L., Amiranashvili, L., Chitchyan, K., Chitchyan, K., Ghazanchyan,
N., Ghazanchyan, N., Darbinyan, K., Darbinyan, K., Gagelidze, N., Gagelidze, N.,
Sadunishvili, T., Sadunishvili, T., Goginyan, V., Goginyan, V., Kvesitadze, G., Kvesitadze, G.,
Torok, T., Torok, T., Mills, D.A., 2015. Microbial biogeography of the transnational
fermented milk matsoni. Food Microbiol. 50, 12–19.
https://doi.org/10.1016/j.fm.2015.01.018.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016.
DADA2: High-resolution sample inference from Illumina amplicon data. Nat. Methods 13,
581-583. https://doi.org/10.1038/nmeth.3869.

Campus, M., Cauli, E., Scano, E., Piras, F., Comunian, R., Paba, A., Daga, E., Salvo, R.D.,
Sedda, P., Angioni, A., Zurru, R., 2017. Towards controlled fermentation of table olives:
LAB starter driven process in an automatic pilot processing plant. Food Bioprocess
Technol. 10, 1063-1073. https://doi.org/10.1007/s11947-017-1882-7.

Campus, M., Corrias, F., Angioni, A., Arru, N., Sedda, P., Addis, M., Fiori, M., Paba, A., Chessa,
L., Comunian, R., 2025. Efficacy of a native microbial starter in promoting table olive
fermentation: an industrial-scale trial at controlled and ambient temperature. Foods 14,
2159. https://doi.org/10.3390/foods14132159.

Campus, M., Değirmencioğlu, N., Comunian, R., 2018. Technologies and trends to improve
table olive quality and safety. Front. Microbiol. 9, 617.
https://doi.org/10.3389/fmicb.2018.00617.Chytiri, A., Tasioula-Margari, M., Bleve, G.,
Kontogianni, V.G., Kallimanis, A., Kontominas, M.G., 2020. Effect of different inoculation
strategies of selected yeast and LAB cultures on Conservolea and Kalamàta table olives
considering phenol content, texture, and sensory attributes. J. Sci. Food Agric. 100, 926-
935. https://doi.org/10.1002/jsfa.10019.

Ciafardini, G., Zullo, B.A., 2019. Use of selected yeast starter cultures in industrial-scale
processing of brined Taggiasca black table olives. Food Microbiol. 84, 103250.
https://doi.org/10.1016/j.fm.2019.103250.

Cocolin, L., Alessandria, V., Botta, C., Gorra, R., De Filippis, F., Ercolini, D., Rantsiou, K., 2013.
NaOH-debittering induces changes in bacterial ecology during table olives fermentation.
PLoS One 8, e69074. https://doi.org/10.1371/journal.pone.0069074.s004.

Comasio, A., Verce, M., Kerrebroeck, S.V., Vuyst, L.D., 2020. Diverse Microbial Composition
of Sourdoughs From Different Origins. Front Microbiol 11, 1212.
https://doi.org/10.3389/fmicb.2020.01212.

Comunian, R., Ferrocino, I., Paba, A., Daga, E., Campus, M., Salvo, R.D., Cauli, E., Piras, F.,
Zurru, R., Cocolin, L., 2017. Evolution of microbiota during spontaneous and inoculated
Tonda di Cagliari table olives fermentation and impact on sensory characteristics. LWT -
Food Sci. Technol. 84, 64-72. https://doi.org/10.1016/j.lwt.2017.05.039.

Corsetti, A., Perpetuini, G., Schirone, M., Tofalo, R., Suzzi, G., 2012. Application of starter
cultures to table olive fermentation: an overview on the experimental studies. Front.
Microbiol. 3, 248. https://doi.org/10.3389/fmicb.2012.00248.

Davis, N.M., Proctor, D.M., Holmes, S.P., Relman, D.A., Callahan, B.J., 2018. Simple statistical
identification and removal of contaminant sequences in marker-gene and metagenomics
data. Microbiome 6, 226. https://doi.org/10.1186/s40168-018-0605-2.

De Angelis, M., Campanella, D., Cosmai, L., Summo, C., Rizzello, C.G., Caponio, F., 2015.
Microbiota and metabolome of un-started and started Greek-type fermentation of Bella
di Cerignola table olives. Food Microbiol. 52, 18-30.
https://doi.org/10.1016/j.fm.2015.06.002.

de Castro, A., Sánchez, A.H., López-López, A., Cortés-Delgado, A., Medina, E., Montaño, A.,
2018. Microbiota and metabolite profiling of spoiled spanish-style Green table Olives.
Metabolites 8, 73. https://doi.org/10.3390/metabo8040073

de Castro, A., Ruiz-Barba, J.L., Romero, C., Sánchez, A.H., García, P., Brenes, M., 2022.
Formation of gas pocket defect in Spanish-style green olives by the halophile
Celerinatantimonas sp. Food Control 136, 108868.
https://doi.org/10.1016/j.foodcont.2022.108868.

Donachie, A., Melillo, T., Bubba, L., Hartman, H., Borg, M.-L., 2018. National outbreak of
Salmonella give linked to a local food manufacturer in Malta, October 2016. Epidemiol.
Infect. 146, 1425-1432. https://doi.org/10.1017/s0950268818001656.

Ferrocino, I., Rantsiou, K., Cocolin, L., 2022. Microbiome and -omics application in food
industry. Int J. Food Microbiol. 377, 109781.
https://doi.org/10.1016/j.ijfoodmicro.2022.109781.

Ferrocino, I., Buzzanca, D., Pagiati, L., Kazou, M., Georgalaki, M., Hatzopoulos, I., Tsakalidou,
E., 2025. The microbial terroir of the Greek olive varieties. Int. J. Food Microbiol. 441,
111332. https://doi.org/10.1016/j.ijfoodmicro.2025.111332.

Gänzle, M., 2022. The periodic table of fermented foods: limitations and opportunities.
Appl. Microbiol. Biotechnol. 106, 2815-2826. https://doi.org/10.1007/s00253-022-
11909-y.

Garrido-Fernández, A., Benítez-Cabello, A., Rodríguez-Gómez, F., Jiménez-Díaz, R., Arroyo-
López, F.N., Morales, M.L., 2021. Relating starter cultures to volatile profile and potential
markers in green Spanish-style table olives by compositional data analysis. Food
Microbiol. 94, 103659. https://doi.org/10.1016/j.fm.2020.103659.

Giavalisco, M., Lavanga, E., Ricciardi, A., Zotta, T., 2024. Starter cultures for the production
of fermented table olives: current status and future perspectives. Fermentation 10, 351.
https://doi.org/10.3390/fermentation10070351.

Gu, Z., 2022. Complex heatmap visualization. iMeta 1, e43. https://doi.org/10.1002/imt2.43.

Hernández-Velázquez, R., Flörl, L., Lavrinienko, A., Sebechlebská, Z., Merk, L., Greppi, A.,
Bokulich, N.A., 2024. The future is fermented: microbial biodiversity of fermented foods
is a critical resource for food innovation and human health. Trends Food Sci. Technol.
150, 104569. https://doi.org/10.1016/j.tifs.2024.104569.

Kamilari, E., Anagnostopoulos, D.A., Tsaltas, D., 2023. Fermented table olives from Cyprus:
microbiota profile of three varieties from different regions through metabarcoding
sequencing. Front. Microbiol. 13, 1101515.
https://doi.org/10.3389/fmicb.2022.1101515.

Kassambara, A., 2023. rstatix: Pipe-Friendly Framework for Basic Statistical Tests.
https://doi.org/10.32614/CRAN.package.rstatix

Kassambara, A., 2025. _ggpubr: 'ggplot2' Based Publication Ready Plots.
https://doi.org/10.32614/CRAN.package.ggpubr.

Kauserud, H., 2023. ITS alchemy: On the use of ITS as a DNA marker in fungal ecology.
Fungal Ecol. 65, 101274. https://doi.org/10.1016/j.funeco.2023.101274.

Kazou, M., Tzamourani, A., Panagou, E.Z., Tsakalidou, E., 2020. Unraveling the Microbiota of
Natural Black cv. Kalamata Fermented Olives through 16S and ITS Metataxonomic
Analysis. Microorganisms 8, 672. https://doi.org/10.3390/microorganisms8050672.

International Olive Oil Council, 2004. Trade standard applying to table olives. COI/OT/NC no.
1. December 2004.

Langgut, D., Garfinkel, Y., 2022. 7000-year-old evidence of fruit tree cultivation in the Jordan
Valley, Israel. Sci. Rep. 12, 7463. https://doi.org/10.1038/s41598-022-10743-6.

Lanza, B., Zago, M., Marco, S.D., Loreto, G.D., Cellini, M., Tidona, F., Bonvini, B., Bacceli, M.,
Simone, N., 2020. Single and multiple inoculum of *Lactiplantibacillus plantarum* strains in
table olive lab-scale fermentations. Fermentation 6, 126.
https://doi.org/10.3390/fermentation6040126.

Lanza, B., Marco, S.D., Bacceli, M., Serio, M.G.D., Loreto, G.D., Cellini, M., Simone, N., 2021.
Lactiplantibacillus plantarum used as single, multiple, and mixed starter combined with
Candida boidinii for table olive fermentations: chemical, textural, and sensorial
characterization of final products. Fermentation 7, 239.
https://doi.org/10.3390/fermentation7040239.

Lee, H.M., Lee, Y., 2008. A differential medium for lactic acid-producing bacteria in a mixed
culture. Lett. Appl. Microbiol. 46, 676-681. https://doi.org/10.1111/j.1472-
765x.2008.02371.x.

Li, R., Yang, S., Lin, M., Guo, S., Han, X., Ren, M., Du, L., Song, Y., You, Y., Zhan, J., Huang, W.,
2022. The biogeography of fungal communities across different chinese wine-producing
regions associated with environmental factors and spontaneous fermentation
performance. Front. Microbiol. 12, 636639. https://doi.org/10.3389/fmicb.2021.636639.

López-García, E., Benítez-Cabello, A., Arroyo-López, F.N., 2025. Applying machine learning to
classify table olives using bacterial metataxonomic data. npj Sci. Food 9, 121.
https://doi.org/10.1038/s41538-025-00496-7.

López-García, E., Benítez-Cabello, A., Rodríguez-Gómez, F., Romero-Gil, V., Garrido-
Fernández, A., Jiménez-Díaz, R., Arroyo-López, F.N., 2022. Bacterial metataxonomic
analysis of industrial Spanish-style green table olive fermentations. Food Control 137,
108969. https://doi.org/10.1016/j.foodcont.2022.108969.

López-García, E., Romero-Gil, V., Arroyo-López, F.N., Benítez-Cabello, A., 2024. Impact of
lactic acid bacteria inoculation on fungal diversity during Spanish-style green table olive
fermentations. Int. J. Food Microbiol. 417, 110689.
https://doi.org/10.1016/j.ijfoodmicro.2024.110689.

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. 15, 550.
https://doi.org/10.1186/s13059-014-0550-8.

Lucena-Padrós, H., Ruiz-Barba, J.L., 2016. Diversity and enumeration of halophilic and
alkaliphilic bacteria in Spanish-style green table-olive fermentations. Food Microbiol. 53,
53-62. https://doi.org/10.1016/j.fm.2015.09.006.

Lucena-Padrós, H., Ruiz-Barba, J.L., 2019. Microbial biogeography of Spanish-style green
olive fermentations in the province of Seville, Spain. Food Microbiol. 82, 259-268.
https://doi.org/10.1016/j.fm.2019.02.004.

Maoloni, A., Cardinali, F., Milanović, V., Osimani, A., Garofalo, C., Ferrocino, I., Corvaglia,
M.R., Cocolin, L., Aquilanti, L., 2022. Microbial dynamics and key sensory traits of

laboratory-scale co-fermented green olives (*Olea europaea* L. cv. *Ascolana tenera*) and sea fennel (*Crithmum maritimum* L.). Food Biosci. 50, 102077. https://doi.org/10.1016/j.fbio.2022.102077.

Martorana, A., Alfonzo, A., Settanni, L., Corona, O., Croce, F.L., Caruso, T., Moschetti, G., Francesca, N., 2015. An innovative method to produce green table olives based on "pied de cuve" technology. Food Microbiol. 50, 126-140. https://doi.org/10.1016/j.fm.2015.03.008.

McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PLoS One 8, e61217. https://doi.org/10.1371/journal.pone.0061217.

Medina, E., Ruiz-Bellido, M.A., Romero-Gil, V., Rodríguez-Gómez, F., Montes-Borrego, M., Landa, B.B., Arroyo-López, F.N., 2016. Assessment of the bacterial community in directly brined *Aloreña de Málaga* table olive fermentations by metagenetic analysis. Int. J. Food Microbiol. 236, 47-55. https://doi.org/10.1016/j.ijfoodmicro.2016.07.014.

Medina, E., Brenes, M., García-García, P., Romero, C., de Castro, A., 2018. Microbial ecology along the processing of Spanish olives darkened by oxidation. Food Control 86, 35-41. https://doi.org/10.1016/j.foodcont.2017.10.035.

Michailidou, S., Trikka, F., Pasentsis, K., Petrovits, G.E., Kyritsi, M., Argiriou, A., 2021. Insights into the evolution of Greek style table olives microbiome stored under modified atmosphere: Biochemical implications on the product quality. Food Control 130, 108286. https://doi.org/10.1016/j.foodcont.2021.108286.

Montaño, A., Cortés-Delgado, A., Sánchez, A.H., Ruiz-Barba, J.L., 2021. Production of volatile compounds by wild-type yeasts in a natural olive-derived culture medium. Food Microbiol. 98, 103788. https://doi.org/10.1016/j.fm.2021.103788.

Nikoloudaki, O., Aheto, F., Di Cagno, R., Gobbetti, M., 2024. Synthetic microbial communities: A gateway to understanding resistance, resilience, and functionality in spontaneously fermented food microbiomes. Food Res. Int. 192, 114780. https://doi.org/10.1016/j.foodres.2024.114780.

Oksanen, J., Simpson, G., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O'Hara, R., Solymos, P., Stevens, M., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Borman, T., Carvalho, G., Chirico, M., De Caceres, M., Durand, S., Evangelista, H., FitzJohn, R., Friendly, M., Furneaux, B., Hannigan, G., Hill, M., Lahti, L., Martino, C., McGlinn, D., Ouellette, M., Ribeiro Cunha, E., Smith, T., Stier, A., Ter Braak, C., Weedon, J., 2025. _vegan: Community Ecology Package. https://doi.org/10.32614/CRAN.package.vegan.

O'Sullivan, D.J., Fallico, V., O'sullivan, O., McSweeney, P.L.H., Sheehan, J.J., Cotter, P.D., Giblin, L., 2015. High-throughput DNA sequencing to survey bacterial histidine and

tyrosine decarboxylases in raw milk cheeses. BMC Microbiol. 15, 266.
https://doi.org/10.1186/s12866-015-0596-0.

Paba, A., Chessa, L., Daga, E., Campus, M., Bulla, M., Angioni, A., Sedda, P., Comunian, R.,
2020. Do best-selected strains perform table olive fermentation better than undefined
biodiverse starters? A comparative study. Foods 9, 135.
https://doi.org/10.3390/foods9020135.

Parente, E., Zotta, T., Giavalisco, M., Ricciardi, A., 2023. Metataxonomic insights in the
distribution of *Lactobacillaceae* in foods and food environments. Int. J. Food Microbiol.
391, 110124. https://doi.org/10.1016/j.ijfoodmicro.2023.110124.

Penland, M., Deutsch, S.-M., Falentin, H., Pawtowski, A., Poirier, E., Visenti, G., Meur, C.L.,
Maillard, M.-B., Thierry, A., Mounier, J., Coton, M., 2020. Deciphering microbial
community dynamics and biochemical changes during nyons black olive natural
fermentations. Front. Microbiol. 11, 586614.
https://doi.org/10.3389/fmicb.2020.586614.

Penland, M., Mounier, J., Pawtowski, A., Tréguer, S., Deutsch, S.-M., Coton, M., 2021. Use of
metabarcoding and source tracking to identify desirable or spoilage autochthonous
microorganism sources during black olive fermentations. Food Res. Int. 144, 110344.
https://doi.org/10.1016/j.foodres.2021.110344.

Penland, M., Pawtowski, A., Pioli, A., Maillard, M.-B., Debaets, S., Deutsch, S.-M., Falentin,
H., Mounier, J., Coton, M., 2022. Brine salt concentration reduction and inoculation with
autochthonous consortia: impact on protected designation of origin nyons black table
olive fermentations. Food Res. Int. 155, 111069.
https://doi.org/10.1016/j.foodres.2022.111069.

Perpetuini, G., Prete, R., Garcia-Gonzalez, N., Alam, M.K., Corsetti, A., 2020. Table olives
more than a fermented food. Foods 9, 178. https://doi.org/10.3390/foods9020178.

Pino, A., De Angelis, M., Todaro, A., Hoorde, K.V., Randazzo, C.L., Caggia, C., 2018.
Fermentation of Nocellara Etnea table olives by functional starter cultures at different
low salt concentrations. Front. Microbiol. 9, 1125.
https://doi.org/10.3389/fmicb.2018.01125.

Pino, A., Vaccalluzzo, A., Solieri, L., Romeo, F.V., Todaro, A., Caggia, C., Arroyo-López, F.N.,
Bautista-Gallego, J., Randazzo, C.L., 2019. Effect of sequential inoculum of beta-
glucosidase positive and probiotic strains on brine fermentation to obtain low salt Sicilian
table olives. Front. Microbiol. 10, 174. https://doi.org/10.3389/fmicb.2019.00174.

Portilha-Cunha, M.F., Macedo, A.C., Malcata, F.X., 2020. A review on adventitious lactic acid
bacteria from table olives. Foods 9, 948. https://doi.org/10.3390/foods9070948.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O.,
2013. The SILVA ribosomal RNA gene database project: improved data processing and
web-based tools. Nucleic Acids Res.41, D590-6. https://doi.org/10.1093/nar/gks1219.

R Core Team, 2025. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Randazzo, C.L., Todaro, A., Pino, A., Pitino, I., Corona, O., Caggia, C., 2017. Microbiota and
metabolome during controlled and spontaneous fermentation of Nocellara Etnea table
olives. Food Microbiol. 65, 136-148. https://doi.org/10.1016/j.fm.2017.01.022.

Ricciardi, A., Parente, E., Tramutola, A., Guidone, A., Ianniello, R.G., Pavlidis, D., Tsakalidou,
E., Zotta, T., 2014. Evaluation of a differential medium for the preliminary identification
of members of the *Lactobacillus plantarum* and *Lactobacillus casei* groups. Ann.
Microbiol. 1-10. https://doi.org/10.1007/s13213-014-1004-y.

Ricciardi, A., López, F.N.A., Giavalisco, M., Pietrafesa, R., Parente, E., 2025. Determining the
core bacterial and fungal genera in table olive fermentations. Int. J. Food Microbiol. 442,
111344. https://doi.org/10.1016/j.ijfoodmicro.2025.111344.

Rodríguez-Gómez, F., Valero, A., Lara, E.V., Marín, A., Ramírez, E.M., 2023. LP309 a new
strain of *Lactiplantibacillus pentosus* that improves the lactic fermentation of Spanish-
style table olives. J. Food Sci. 88, 5191-5202. https://doi.org/10.1111/1750-3841.16802.

Rué, O., Coton, M., Dugat-Bony, E., Howell, K., Irlinger, F., Legras, J.-L., Loux, V., Michel, E.,
Mounier, J., Neuvéglise, C., Sicard, D., 2023. Comparison of metabarcoding taxonomic
markers to describe fungal communities in fermented foods. Peer Community J. 3, e97.
https://doi.org/10.24072/pcjournal.321.

Ruiz-Barba, J.L., Cortés-Delgado, A., Sánchez, A.H., López-López, A., Montaño, A., 2024.
Impact of selected wild yeasts starters on the volatilome and phenolic contents of
Gordal, Manzanilla and Hojiblanca naturally fermented green olives. LWT 195, 115811.
https://doi.org/10.1016/j.lwt.2024.115811.

Ruiz-Barba, J.L., Jimenéz-Diaz, R., 2012. A novel *Lactobacillus pentosus*-paired starter culture
for Spanish-style green olive fermentation. Food Microbiol. 30, 253-259.
https://doi.org/10.1016/j.fm.2011.11.004.

Ruiz-Barba, J.L., Sánchez, A.H., López-López, A., Cortés-Delgado, A., Montaño, A., 2023a.
Microbial and chemical characterization of natural-style green table olives from the
Gordal, Hojiblanca and Manzanilla cultivars. Foods 12, 2386.
https://doi.org/10.3390/foods12122386.

Ruiz-Barba, J.L., Sánchez, A.H., López-López, A., Cortés-Delgado, A., Montaño, A., 2023b.
Microbial community and volatilome changes in brines along the spontaneous
fermentation of Spanish-style and natural-style green table olives (Manzanilla cultivar).
Food Microbiol. 113, 104286. https://doi.org/10.1016/j.fm.2023.104286.

Sacchi, R., Corrado, G., Basile, B., Mandarello, D., Ambrosino, M.L., Paduano, A., Savarese,
M., Caporaso, N., Aponte, M., Genovese, A., 2022. Effect of salt addition and
fermentation time on phenolics, microbial dynamics, volatile organic compounds, and

sensory properties of the PDO table olives of Gaeta (Italy). Molecules 27, 8100.
https://doi.org/10.3390/molecules27228100.

Sánchez, A.H., López-López, A., Cortés-Delgado, A., Beato, V.M., Medina, E., de Castro, A.,
Montaño, A., 2018. Effect of post-fermentation and packing stages on the volatile
composition of Spanish-style green table olives. Food Chem. 239, 343-353.
https://doi.org/10.1016/j.foodchem.2017.06.125.

Servili, M., Settanni, L., Veneziani, G., Esposto, S., Massitti, O., Taticchi, A., Urbani, S.,
Montedoro, G.F., Corsetti, A., 2006. The Use of *Lactobacillus pentosus* 1MO to shorten
the debittering process time of black table olives (Cv. Itrana and Leccino): a pilot-scale
application. J. Agr. Food Chem. 54, 3869-3875. https://doi.org/10.1021/jf053206y.

Suzzi, G., Gardini, F., 2003. Biogenic amines in dry fermented sausages: a review. Int. J. Food
Microbiol. 88, 41-54. https://doi.org/ 10.1016/s0168-1605(03)00080-1.

Tamang, J.P., Cotter, P.D., Endo, A., Han, N.S., Kort, R., Liu, S.Q., Mayo, B., Westerik, N.,
Hutkins, R., 2020. Fermented foods in a global age: East meets West. Compr. Rev. Food
Sci. Food Saf. 19, 184-217. https://doi.org/10.1111/1541-4337.12520

Tarantini, A., Crupi, P., Ramires, F.A., D'Amico, L., Romano, G., Blando, F., Branco, P.,
Clodoveo, M.L., Corbo, F., Cardinali, A., Bleve, G., 2024. Study of the effects of
pasteurization and selected microbial starters on functional traits of fermented table
olives. Food Microbiol. 122, 104537. https://doi.org/10.1016/j.fm.2024.104537

Tedersoo, L., Bahram, M., Zinger, L., Nilsson, R.H., Kennedy, P.G., Yang, T., Anslan, S.,
Mikryukov, V., 2022. Best practices in metabarcoding of fungi: From experimental design
to results. Mol. Ecol. 31, 2769-2795. https://doi.org/10.1111/mec.16460.

Tofalo, R., Schirone, M., Perpetuini, G., Angelozzi, G., Suzzi, G., Corsetti, A., 2012.
Microbiological and chemical profiles of naturally fermented table olives and brines from
different Italian cultivars. Antonie van Leeuwenhoek 102, 121-131.
https://doi.org/10.1007/s10482-012-9719-x.

Traina, C., Ferrocino, I., Bonciolini, A., Cardenia, V., Lin, X., Rantsiou, K., Cocolin, L., 2024.
Monitoring the yeasts ecology and volatiles profile throughout the spontaneous
fermentation of Taggiasca cv. table olives through culture-dependent and independent
methods. Int. J. Food Microbiol. 417, 110688.
https://doi.org/10.1016/j.ijfoodmicro.2024.110688..

Tsoungos, A., Pemaj, V., Slavko, A., Kapolos, J., Papadelli, M., Papadimitriou, K., 2023. The
rising role of omics and meta-omics in table olive research. Foods 12, 3783.
https://doi.org/10.3390/foods12203783

Tubia, I., Prasad, K., Pérez-Lorenzo, E., Abadín, C., Zumárraga, M., Oyanguren, I., Barbero, F.,
Paredes, J., Arana, S., 2018. Beverage spoilage yeast detection methods and control
technologies: A review of *Brettanomyces*. Int. J. Food Microbiol. 283, 65-76.
https://doi.org/10.1016/j.ijfoodmicro.2018.06.020..

Tufariello, M., Anglana, C., Crupi, P., Virtuosi, I., Fiume, P., Terlizzi, B.D., Moselhy, N., Attay,
H.A., Pati, S., Logrieco, A.F., Mita, G., Bleve, G., 2019. Efficacy of yeast starters to drive
and improve Picual, Manzanilla and Kalamàta table olive fermentation. J. Sci. Food Agric.
99, 2504-2512. https://doi.org/10.1002/jsfa.9460.

Tzamourani, A., Kalogri, G., Kavvatha, M., Kochila, A., Manthou, E., Liu, Y., Panagou, E.Z.,
2024. Inoculated fermentation of cv. Conservolea natural black olives with
multifunctional starter cultures in reduced-sodium brines. Int. J. Food Sci. Technol. 59,
4093-4108. https://doi.org/10.1111/ijfs.17165.

Vaccalluzzo, A., Pino, A., Russo, N., De Angelis, M., Caggia, C., Randazzo, C.L., 2020.
FoodOmics as a new frontier to reveal microbial community and metabolic processes
occurring on table olives fermentation. Food Microbiol. 92, 103606.
https://doi.org/10.1016/j.fm.2020.103606.

Vougiouklaki, D., Letsiou, S., Mavrokefalidou, I., Tsakali, E., Akkermans, S., Impe, J.F.M.V.,
Houhoula, D., 2024. Fermentation of Kalamata natural black olives using selected lactic
acid bacteria as starters. Fermentation 10, 53.
https://doi.org/10.3390/fermentation10010053.

Wang, J., Bakker, H.C. den, Denes, T.G., 2025. A critical review of metagenomic approaches
for foodborne pathogen surveillance. Crit. Rev. Food Sci. Nutr. 65, 1-14.
https://doi.org/10.1080/10408398.2025.2503453.

Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naive Bayesian classifier for rapid
assignment of rRNA sequences into the new bacterial taxonomy. Applied and
Environmental Microbiology 73, 5261–5267. https://doi.org/10.1128/aem.00062-07.

White, T. J., Bruns, T., Lee, S., Taylor, J., 1990. Amplification and direct sequencing of fungal
ribosomal RNA genes for phylogenetics. In Innis, M. A., Gelfald, D. H., Sninsky, J. J. (Eds.)
PCR protocols: a guide to methods and applications. Academic Press, New York, pp. 315–
322.

Wickham, H., 2016 ggplot2: Elegant graphics for data analysis. Springer-Verlag New York.

Zago, M., Lanza, B., Rossetti, L., Muzzalupo, I., Carminati, D., Giraffa, G., 2013. Selection of
Lactobacillus plantarum strains to use as starters in fermented table olives:
oleuropeinase activity and phage sensitivity. Food Microbiol. 34, 81-87.
https://doi.org/10.1016/j.fm.2012.11.005.

Zinno, P., Guantario, B., Perozzi, G., Pastore, G., Devirgiliis, C., 2017. Impact of NaCl
reduction on lactic acid bacteria during fermentation of Nocellara del Belice table olives.
Food Microbiol. 63, 239–247. https://doi.org/10.1016/j.fm.2016.12.001.

Zotta, T., Giavalisco, M., Parente, E., Picariello, G., Siano, F., Ricciardi, A., 2022. Selection of
Lactiplantibacillus strains for the production of fermented table olives. Microorganisms
10, 625. https://doi.org/10.3390/microorganisms10030625.