A survey of bacterial and fungal communities of table olives.

Eugenio Parente[1], Rocchina Pietrafesa[1], Francesca De Filippis[2], Alessandra De Vivo[2], Maria

Grazia Labella[1], Marina Hidalgo[3], Emanuela Lavanga[1], Annamaria Ricciardi[1]

Affiliations

1 Dipartimento di Scienze Agrarie, Forestali, Alimentari ed Ambientali, Università degli Studi

della Basilicata, Potenza, Italy

2 Dipartimento di Agraria, Università degli Studi di Napoli "Federico II", Portici (NA), Italy

3 Departamento de Ciencias de la Salud, Universidad de Jaén, Jaén, Spain

Corresponding author

Prof. Eugenio PARENTE

Dipartimento di Scienze Agrarie, Forestali, Alimentari ed Ambientali

Università degli Studi della Basilicata

Via dell'Ateneo Lucano, 10

85100 Potenza, Italy

Abstract

Table olives are among the most ancient fermented foods. Their production extends beyond

the Mediterranean basin and the transformation of different olive varieties according to

various trade preparations results in high product diversity. Several varieties have Protected

Designation of Origin, and, in Italy the black naturally fermented Oliva di Gaeta is prominent.

In order to understand the diversity of bacterial and fungal communities and how it is

affected by olive variety, olive ripeness and trade preparation, we carried out a survey using

364 samples from 40 producers and 6 countries. We measured pH, salt concentration, viable

counts and carried out amplicon targeted metagenomics of bacterial and fungal

communities. Olive variety, olive ripeness and trade preparation all significantly affected the

diversity and composition of bacterial and fungal communities but dispersal plays a

confounding role: even within the same variety we found significant differences between

producing firms and a large variability even within the same firm. This may be due to limited

control of contamination (with house microbiota playing a role in initial contamination

events) and of technological factors and also to the widespread use of relatively small

fermenters (240-280 L). This variability obscured the differences that existed between PDO

(Oliva di Gaeta) and non PDO (Itrana nera) black naturally fermented olives of the Itrana

variety. Nonetheless, we were able to identify subset of bacterial and fungal genera which

were differentially abundant in selected trade preparations and confirmed that the diversity

of lactic acid bacteria communities of naturally fermented olives.

Keywords

Table olives; amplicon targeted metagenomics; Oliva di Gaeta; olive fermentation;

microbiota dynamics.

1. Introduction

Fermented foods have been part of the human diet for millennia and their diversity is huge

(Gänzle, 2022; Hernández-Velázquez et al., 2024; Tamang et al., 2020). Fermented

vegetables are very diverse, and their production is often based on artisanal practices,
without heat treatment of the raw materials and addition of starter culture, although back-
slopping is frequently used (Tamang et al., 2020). Table olives are among the most ancient
fermented vegetables from the Mediterranean basin (Langgut and Garfinkel, 2022) and this
has resulted in a great variety of products (Anagnostopoulos and Tsaltas, 2022; Perpetuini et
al., 2020). Fermentation of table olives has multiple functions: removing the bitter glucoside
oleuropein, enhancing the shelf life through acid production in the brine, and improving
sensory properties. Five main categories are recognized by the International Olive Oil
Council (https://www.internationaloliveoil.org/wp-content/uploads/2019/11/COI-OT-NC1-
2004-Eng.pdf). Alkali treated olives, also known as Spanish style olives, are treated with
sodium hydroxide to remove oleuropein before being washed and fermented in brine. Green
specialties, like Picholine style olives, on the other hand, are stored at low temperature in
brine after the alkali treatment, thus limiting fermentation. Naturally fermented olives
(Greek style) include both green and ripe olives directly fermented in brines.
Dried/shrivelled olives and olives darkened by oxidation, are preserved by methods other
than fermentation. Several varieties have received Protected Denomination of Origin (PDO).
In Italy four varieties (Oliva di Gaeta, Nocellara del Belice, Bella di Daunia, Oliva Ascolana del
Piceno) are included in the PDO food list and the Oliva Taggiasca Ligure has recently received
Protected Geographical Indication status
(https://ec.europa.eu/agriculture/eambrosia/geographical-indications-register/). Many
more are included in the Italian List of Traditional Agricultural Products
(https://www.gazzettaufficiale.it/eli/gu/2024/03/13/61/sg/pdf). Spain, Greece, France,
Portugal and Turkey also produce a wide variety of table olives, including PDO products.
Sensory quality, safety and stability of table olives are affected by several factors, including
olive ripeness, technological factors, and by the growth of microorganisms, either present as
natural contaminants on the fruits, salt, brines, and contact surfaces or added as starter
cultures (Anagnostopoulos and Tsaltas, 2022; Campus et al., 2018). Current trends in table
olive production aim at controlling the microbiota involved in fermentation, by sanitizing the
raw materials, by adding starter and functional cultures, and/or by reducing salt content
(Anagnostopoulos and Tsaltas, 2022; Campus et al., 2018). Addition of starters has been
tested repeatedly and, typically, includes lactic acid bacteria (*Lactiplantibacillus pentosus* or
Lactipl. plantarum, *Lacticaseibacillus paracasei* (Alfonzo et al., 2023; Comunian et al., 2017;
Lanza et al., 2021, 2020; Paba et al., 2020; Pino et al., 2019, 2018; Randazzo et al., 2017;
Rodríguez-Gómez et al., 2023; Ruiz-Barba and Jimenéz-Diaz, 2012; Servili et al., 2006;
Tzamourani et al., 2024; Vougiouklaki et al., 2024; Zago et al., 2013)), yeasts (several species
belonging to the genera *Candida, Wyckerhamomyces, Pichia, Cyteromyces* etc. have been
tested: (Ciafardini and Zullo, 2019; Montaño et al., 2021; Ruiz-Barba et al., 2024; Tufariello et
al., 2019)) or their combinations (Benítez-Cabello et al., 2020; Garrido-Fernández et al.,
2021). Back-slopping practices (Martorana et al., 2015) have also been described.
The bacterial and fungal microbiota of table olives is complex, usually due to the lack of
control of unit operations affecting dispersal and selection and includes microbial groups
which are considered beneficial, like lactic acid bacteria, yeasts, and other halophilic
microorganisms (Halophilic and Alkalophilic Lactic Acid Bacteria, HALAB, and halophilic
Pseudomonadota), but also microorganisms associated with spoilage, like some halophiles
(*Alkalibacterium, Marinilactobacillus, Celerinatantimonas*, some *Vibrio*, proprionibacteria,
clostridia and yeasts (Arroyo-López et al., 2021, 2012; Ballester et al., 2021; de Castro et al.,
2018; Penland et al., 2021) . The composition and dynamics of microbial communities in

table olives have been extensively reviewed (Anagnostopoulos and Tsaltas, 2022; Campus et
al., 2018; Portilha-Cunha et al., 2020; Tsoungos et al., 2023; Vaccalluzzo et al., 2020). We
have recently reviewed metataxonomic studies and found that a core microbiota of
approximately 30 genera is characteristic of table olive fermentations, with wide variations
among trade preparations and varieties (Ricciardi et al., 2025a). Geographic origin is also
believed to be a contributing factor (Argyri et al., 2020; Kamilari et al., 2023).
With a few exceptions (Penland et al., 2022, 2021, 2020) studies on table olives microbiota
are relatively small (<50 samples) and often focus on a limited number of varieties (see
reviews by (Ricciardi et al., 2025a; Tsoungos et al., 2023) for details). This, and the variety of
sampling plans and experimental approaches used, complicates the interpretation and the
integration of these data to obtain a comprehensive view of the factors affecting table olive
microbial community assembly and dynamics.
In this study, we carried out a large survey of table olives belonging to a wide array of
varieties and trade preparations from 6 countries (Italy, Greece, Spain, Cyprus, Peru, Egypt)
and including both sound and spoiled samples with the following objectives:
a. obtaining a comprehensive view of the bacterial and fungal microbial communities of
table olives, understanding their variability due to ripeness and trade preparation,
variety, between and within producing company.
b. using naturally fermented black Itrana olives as a case study to evaluate if detectable
differences between PDO and non-PDO products exist
115
116 **2. Materials and methods**
117
2.1 Sampling.
Table olive samples (n=340) were obtained from 40 producers across 6 countries (Italy,
Greece, Spain, Cyprus, Peru, Egypt). Producers were required to deliver olives ready for
packaging in their fermentation brine, or immediately after packaging and to fill a
questionnaire reporting essential metadata (location of the company, olive ripeness and
variety, the techniques used for transformation, addition of starter cultures, if any; spoilage;
duration of fermentation and of storage post fermentation, if applicable). Twenty sample,
for which the fermentation brine was not available, were dispatched in the final packages,
with the storage brine. Furthermore, 24 metagenomic DNA from table olives were obtained
from Instituto de la Grasa, CSIC (Seville, Spain). Summary data are shown in Table 1, while
detailed data are shown in Supplementary Table 1. All samples were dispatched in insulated
containers with ice packs and reached our laboratory within 48 (samples from Italy) to 72 h.
They were stored at 4°C for no more than 24 h before being used for microbiological
analyses and extraction of DNA. Supernatants (12,000 x g) of the brine were stored at -20°C
for 7-30 gg before the measurement of pH, titratable acidity and chlorides.
2.2 Chemical and physicochemical analyses.
pH and chlorides were measured in brines except for samples of dehydrated olives. The
latter were homogenized (1:4) with deionized water in a Stomacher Lab Blender for 1 min
using a filter bag, and pH and chlorides were measured in the filtered liquid phase.
pH was measured using a digital pH meter (CyberScan pH 110, Eutech Instrument Pte Ltd.,
Thermo Fisher Scientific, Monza, Italy) with a combination electrode (Double Pore Slim
electrode, Hamilton Company, Reno, Nevada, USA). NaCl concentration was estimated from
the measurement of chlorides on diluted (D = 10-50) brines using a selective electrode
(Crison ISE Chlorid 51340400) with a reference electrode (Crison 5241 with KNO_3 1 mol/L as

a reference electrolyte), connected to an Orion 420 A plus ion-analyzer. 5 mol/L NaNO$_3$ was
used as ionic strength adjuster (1 ml / 50 mL diluted brine) and a calibration curve (0.001-0.1
mol/L NaCl)
Titratable acidity was only measured in brines using phenophthalein (Sigma) as an indicator
using NaOH 0.1 mol/L and a digital burette (Brand Titrette, Brand, Wertheim, Germany).
Two technical replicates were used for each measurement.
148
149

Table 1. The samples used in this project, by olive ripeness and trade preparation.

olive ripeness	olive trade preparation	n
Green olives	Alkali treated olives	68
Green olives	Natural olives	65
Green olives	Olives darkened by oxidation	5
Green olives	Specialties (Picholine, etc.)	30
Olives turning color	Natural olives	41
Olives turning color	Olives darkened by oxidation	9
Black olives	Dehydrated olives	4
Black olives	Natural olives	152

150
2.3 Enumeration of microbial groups, isolation and identification.
Counts were performed on a 1:1 mixture of olives (bones were aseptically removed) and
brine after homogenization (2 min) in a filtered bag (Nasco Whirlpak, Pleasant Prairie, WI)
using a Stomacher Lab Blender 400 (Cole-Parmer, now Antylia Scientific, Vernon Hills, IL,
USA) in peptone water (Bacteriological peptone 0.1% w/v, NaCl 0.85% w/v).
Serial ten-fold dilutions were performed in peptone water. Lactic acid bacteria were
enumerated in modified MRS agar (Lee and Lee, 2008; Ricciardi et al., 2014) with 0.02%
sodium azide (mMRS+NA, 25°C, 72 h), yeasts and moulds on Glucose Yeast Extract agar with
100 mg/L chloramphenicol (GYEAC, 25°C, 5 days), halophilic microorganisms on Plate Count
Agar Standard with 5% NaCl (w/v) (PCA + NaCl, 30°C, 48 h), contaminants on *Pseudomonas*
agar base (PAB, 25°C, 48 h), and *Enterobacteriaceae* in Violet Red Bile Glucose Agar (VRBGA,

30°C, 24 h). All microbiological media were inoculated using a spiral plater (WASP Spiral
Plater, bioMérieux Italia SpA, Bagno a Ripoli, Italy) with the exception of VRBGA, for which
conventional pour plating was used. Colony counts were performed using a digital colony
counter (EasyCount 2, bioMérieux Italia).
Colonies were isolated and streaked for purification on the same medium and, after control
of cell morphology by light microscopy, identification was carried out with MALDI-ToF
Biotyper Sirius (Bruker Co, Mallerica, MA, USA) using the "Extended Direct Transfer" SOP
provided by the producer.
170
171 *2.4 Microbiological media, ingredients and reagents.*
172 All microbiological media and ingredients were obtained from Oxoid (Basingstoke, United
173 Kingdom), and reagents from Sigma-Aldrich (Merck KGaA, Darmstadt, Germany).
174
2.5 Amplicon targeted metagenomics.
Olives (40 g) were mixed with an equal volume of brine (for dehydrated olives, sterile
distilled water) to which Tween 80 (final concentration 0.1% vol/vol) was aseptically added.
After shaking for 10 min on a rotary shaker (100 rpm) in ice, the liquid was centrifuged
(8,000xg for 10 min) and the pellet was washed twice in sterile NaCl 0.85% and immediately
frozen at -80°C. DNA extraction and purification were performed using DNeasy PowerFood
Microbial Kit (QIAGEN Italia, Milan, Italy) followed by quantitation using NanoDrop One
spectrophotometer (Thermo-Fisher Scientific, Milan, Italia). Library preparation and
sequencing was performed by Novogene (Cambridge, UK). The V3-V4 region of the 16S rRNA
gene was amplified using primers Bakt_341F_Nov (5'-CCTAYGGGRBGCASCAG-3') and
Bakt_805R_Nov (5'-GGACTACNNGGGTATCTAAT-3'). The ITS2 region was amplified using
primers ITS3F (5'-GCATCGATGAAGAACGCAGC-3') and ITS4R (5'-TCCTCCGCTTATTGATATGC-3')
for Fungi. Paired end sequencing (2x250 bp) was performed on Illumina NovaSeq X.
Raw sequences were processed using a pipeline based on DADA2 (Callahan et al.,
2016) using R (R Core Team, 2025). Model scripts are available at
https://github.com/ep142/FoodMicrobionet/tree/master/dada2_pipeline. Taxonomic
assignment was carried out using SILVA 138.2 (Quast et al., 2013) for bacteria and UNITE
v10 general release 2024-04-04 for fungi ((Abarenkov et al., 2023). The results were
assembled in a phyloseq object (McMurdie and Holmes, 2013) which was then used for
further analysis. Extraction blanks and a mock community (ZymoBIOMICS Microbial
Community DNA Standard, Zymo Research, Invine, CA, USA) were also included.
Contaminating ASVs were removed using package "decontam" (Davis et al., 2018).
197
2.6 Statistical analysis
Non-parametric tests (Wilcoxon-Signed Rank Sum test) were used for comparing
distributions of physico-chemical and chemical variables and counts (as log(cfu/g brine and
olives)). Holm correction for multiple testing was used. Statistical analyses and graphs were
produced with scripts combining the functions of the packages stats (R Core Team, 2025),
PCAtools (Blighe and Lun, 2025), phyloseq (McMurdie and Holmes, 2013), ggplot2
(Wickham, 2016), ggpubr (Kassambara, 2025) and rstatix (Kassambara, 2023). Alpha
diversity analyses, ordinations and inferential analyses on microbial community data were
performed using functions of the packages phyloseq (McMurdie and Holmes, 2013), DESeq2
(Love et al., 2014) and vegan (Oksanen et al., 2025).
208

209 3. **Results**.
210
3.1 Sampling.
A summary of the samples grouped according to ripeness and trade preparation is shown in
Supplementary Table 1. Most of the samples (natural olives, 44%, Spanish-style, alkali
treated Spanish-style olives, 19%) had undergone a fermentation step, while 27% had
undergone an alkali treatment. The latter included both Spanish-style olives and specialties
like Nocellara del Belice produced with the "Castelvetrano" method (Perpetuini et al., 2020).
The remaining samples were produced with other methods, including drying. Spoiled
samples were also made available by some producers (Supplementary Table 3).
219
3.2 pH, titratable acidity and salt content.
The pH of brines and olives in our study ranged from 3.7 to 8.1 (see Supplementary Table 4,
Supplementary Table 5 for summary statistics). The high variability of pH of main varieties
(those with at least 9 samples) is highlighted in Supplementary Figure 1.
The null hypothesis of lack of differences between the distribution of pH values (Wilcoxon
Rank-Sum Test, WSRST, with correction for multiple testing) was discarded with $p < 0.001$ for
the comparisons of black dehydrated olives and specialties (higher median pH) and the
other groups. Even within a single variety, the pH was highly variable (see Supplementary
Figure 1 and Supplementary Table 5). Results for titratable acidity (TA) of brines reflect those
of pH with the exception of olives darkened by oxidation, to which gluconate was added. A
box plot for the main varieties is shown in Supplementary Figure 2, while summary statistics
are reported in Supplementary Table 6. After excluding olives darkened by oxidation and
packaged olives, the titratable acidity ranged from 0.3 to 24 g lactic acid/L, with significantly
higher median values (WSRT) for naturally fermented olives and lower values for alkali
treated olives and specialties in particular. Again, high variability was observed even within
the same variety. There was a weak (-0.58) but highly significant correlation between pH and
the $\log_{10}(TA)$.
Summary statistics for salt content for brines and olives are shown in Supplementary Tables
7 and 8. A box plot for the main varieties is shown in Supplementary Figure 3. NaCl content
ranged from 2.2 g/L to near saturation (246 g/L) but median values ranged from 23 to 82 g/L
for different combinations of ripeness and trade preparation. High variability was observed.
For Oliva di Gaeta samples, which were the most numerous, NaCl content of brines varied
between 30 and 116 g/L. Dehydrated olives and olives darkened by oxidation had
significantly lower median values (WSRT).
244
3.3 Microbial counts.
Viable microbial counts were obtained using selective or elective media for undesirable
microorganisms (*Enterobacteriaceae*, contaminants on *Pseudomonas* agar base) and
potentially beneficial microorganisms. Based on MALDI-ToF or molecular identification of
isolates (Supplementary Table 9) we found that both mMRS+NA and GYEAC were selective
(all isolates were LAB or yeasts, respectively), while PAB (Supplementary Figure 5) supported
the growth of yeasts (which formed pinpoint colonies) and PCA+NaCl supported both the
growth of yeast and lactic acid bacteria together with staphylococci and other minor
halophiles. The latter is therefore of limited use when counts of yeasts and LAB are high.

Counts of *Enterobacteriaceae* are shown in Supplementary Figure 4. For most samples
counts were below the detection limit (<10 cfu/ml), while values > 100 cfu/g (brine + olives)
were found in several samples of naturally fermented or alkali treated olives.
Counts of LAB are shown in figure 1 while summary statistics are shown in Supplementary
Tables 10 and 11. Several samples (30%) had viable counts below the detection limit of our
method (200 cfu/g), with 10^9 cfu/g detected as maximum counts. Spoiled samples of
naturally fermented olives had almost always high counts of LAB, while this pattern was not
observed in alkali treated olives. As for other variables, variability within variety was very
high but some of them showed significantly higher (WSRT) counts compared to others (for
example, Bella di Daunia vs Halkidiki for Green alkali treated olives; Kalamata and
Konservolea vs Oliva di Gaeta for black naturally fermented olives). High counts of LAB were
also found in alkali treated olives which are stored at low temperature after the alkali
treatment, like Nocellara del Belice produced with the Castelvetrano method (Alfonzo et al.,
2024).

268
269 **Figure 1**. Distribution of viable counts of lactic acid bacteria in table olives used in this study.
270

271 Summary results on the counts of yeasts (moulds were only occasionally found) are shown
272 in Supplementary Tables 12 and 13 and their distribution is shown in Figure 2. Only 6% of
273 the samples had counts below the detection limit of our method and, for all olive groups,
274 median values were higher than 10^5 cfu/g.
275

276
277 **Figure 2.** Distribution of viable counts of yeast and moulds in table olives used in this study.
278
Beyond the obvious negative correlation (Sperman rho -0.541) between pH and titratable
acidity, absolute values of the Spearman rho correlation coefficient were often <0.3
(although many were significantly different from 0). As a result of the lack of correlation
structure, the first two components of a Principal Component Analysis carried out on the
correlation matrix explained only 43% of the variance, with no clear separation between
different olive groups (see score plot in Supplementary Figure 6). This again, might be the
result of the very high variability within groups.
286
287
288

3.4 The composition of microbial communities of table olives.

3.4.1 Alpha-diversity of microbial communities.
Extraction of DNA proved to be challenging for some low biomass samples, but we were able to carry out sequencing of the V3-V4 region of the 16S rRNA gene (336 samples) and of the ITS2 region (337 samples) of most of the samples. Mock communities (2) and blanks (3) were included for quality control. The number of sequences after QC filtering, ASV inference and bimera removal was 17.2×10^6 for bacteria and 17.0×10^6 for fungi. 90% of the samples had 29.3×10^3 sequences or more for bacteria, while the 90° percentile for fungi was 34.9×10^3 sequences.
Our pipeline for 16S rRNA sequences theoretically allowed taxonomic assignment down to the species level. However, this is not always possible for *Lactobacillaceae*, even when using a custom database (Parente et al., 2023). Therefore, alpha and beta-diversity analysis for bacterial communities was performed after agglomeration at the genus level. Chao1 values calculated on the bacterial communities are shown in Figure 3, while summary statistics are reported in Supplementary Tables 14 and 15.

Figure 3. Alpha-diversity (Chao1) of bacterial communities in table olives analysed in this study.

Chao1 ranged from 3 to 134 (with median values for different groups ranging from 16.5 to
45). A large variability existed within each group and even within the same variety but some
trends were evident. If olive ripeness is ignored, there is a trend in the decrease of diversity
from Alkali treated olives, to natural olives, to olives darkened by oxidation, to specialties.
We decided to investigate if between-firm and within-firm variability was a contributing
factor. Distribution of Chao1 is shown for Itrana, Bella di Daunia and Halkidiki in
Supplementary Figures 7, 8 and 9. Within the same variety the diversity of bacterial
communities varied between different producers. We also evaluated the correlation
(Spearman ρ) between diversity and physical-chemical parameters and counts for selected
combinations of ripeness and trade preparation (Supplementary Table 16): with the
exception of a strong negative correlation between pH and diversity for Specialties (i.e.
diversity decreases as the pH increases possibly due to the selective effect of fermentation
occurring over time), all the other correlation coefficients were low ($<|0.40|$) although
highly significant and were not necessarily of the same sign for different trade preparations.
Again, differences exist among varieties, possibly because of systematic combinations of
specific factors affecting dispersal and selection: in fact, in Oliva di Gaeta, there was a strong
positive correlation (0.57) between pH and Chao1 compared to other black naturally
fermented olives (data not shown).
Given the larger sequence variability of ITS2 compared to 16S RNA V3-V4 sequences, we
decided to estimate diversity indices for fungi before agglomeration at the genus level. The
distribution of Chao1 values is shown in Figure 4, while summary statistics are in
Supplementary Tables 17 and 18.
Chao1 ranged from 1 to 209 (with median values for different groups ranging from 9 to 41,
lower than those for bacteria for most groups). As for other variables, a large variability
existed within each group and even within the same variety but some trends were evident: if
olive ripeness is ignored, the median value for Chao1 decreased from Specialties to alkali
treated olives, to naturally fermented olives. Significant differences among the distributions
were found using WSRT for the comparison between alkali treated olives (including
specialties) and natural olives. We investigated if between-firm and within-firm variability
was a contributing factor. As for bacterial communities, distribution of Chao1 values within
the same variety (Itrana olives, Bella di Daunia and Halkidiki olives: Supplementary Figures
10, 11 and 12) varied among different producers.
We also evaluated the correlation (Spearman ρ) between selected combinations of ripeness
and trade preparation (Supplementary Table 19): a rather strong negative correlation was
found with yeasts and molds counts and duration of fermentation. Again, differences exist
among varieties, possibly because of systematic combinations of specific factors affecting
dispersal and selection: in fact, in Oliva di Gaeta, there was a positive correlation (0.35)
between pH and Chao1; as for bacteria, higher pH is associated with higher diversity. On the
other hand, there was no significant correlation between Chao1 for fungi and bacteria.
349

Figure 4. Alpha-diversity (Chao1) of fungal communities in table olives used in this study.

After prevalence and abundance filtering (> 1% maximum abundance and >1% prevalence), rare taxa were removed but >99.8% of the sequences were preserved for bacteria (Supplementary Figure 13). The top 24 bacterial taxa selected on the basis of abundance and prevalence and are shown in Supplementary Table 20. The large number of genera retained highlights the high microbial diversity found in table olives. The distribution of the abundance of LAB and HALAB is shown in Figure 5. Although abundances of the main genera are highly variable within each olive group, *Alkalibacterium* and *Marinilactibacillus* are abundant and prevalent in alkali treated olives, while a diverse set of *Lactobacillaceae*, among which *Lactiplantbacillus* is clearly both prevalent and abundant, dominating naturally fermented olives communities. Several heterofermentative LAB are often (*Lentilactobacillus*) or occasionally *(Limosilactobacillus, Levilactobacillus, Paucilactobacillus* and *Leuconostoc*) dominant or subdominant. It is interesting to note that, even in samples where *Lactiplantibacillus* was not dominant, *Lactiplantibacillus pentosus* was the most frequently isolated species anyway (Supplementary Table 9).
The distribution of halophilic genera is shown in Supplementary Figure 14. These included *Celerinatantimonas*, which was the most prevalent and abundant, but other Gram-negatives, such as *Halomonas, Vibrio, Salinicola* and *Idiomarina* were prevalent and occasionally abundant.
In a recent review we have found that DNA of genera including pathogenic bacteria can be found, albeit at very low abundance, in table olives (Ricciardi et al., 2025b). We repeated the

373 analysis for this survey and found that only in very few samples sequences assigned to
374 genera *Salmonella, Yersinia* and *Listeria* were detectable, usually at very low abundance,
375 with one notable exception (Supplementary Figure 15).
376

377
378
379 **Figure 5**. Distribution of the relative abundance of LAB and HALAB in table olives. Genera are
380 ordered by alphabetical order within family.
381
Bar plots of the relative abundances of the 24 most abundant and prevalent genera are
shown in Supplementary Figures 16 (pooled data for all varieties, by ripeness and trade
preparation group) and 17 to 21 (main varieties within ripeness and trade preparation
groups). Some general patterns are evident: more frequent and abundant occurrence of
HALAB in alkali treated olives, while LAB, mainly *Lentilactobacillus, Lactiplantibacillus* and
Pediococcus were dominating members in naturally fermented olives. However, variety-
specific patterns appear. Two examples, for Bella di Cerignola (used for different variations of
Bella di Daunia PDO) and Itrana (used for the production of two non-PDO types and of Oliva
di Gaeta PDO) are shown in Supplementary Figures 22 and 23. In both cases, while there is
some similarity in the composition of the bacterial community within a given variety,
differences between different firms are visually evident.
Abundance and prevalence filtering was performed on fungal communities using
approaches similar to those described above. Our pipeline allowed the assignment at the
species level for the vast majority of ASVs, but given the debate on the reliability of species
assignment using regions of the ribosomal RNA operon in fungi (Kauserud, 2023; Rué et al.,

2023; Tedersoo et al., 2022) we decided to perform the analysis both at the species and
genus levels. Abundance and prevalence filtering at the species level retained 195 taxa out
of 795, with 99.6% of sequences, mostly belonging to phylum Ascomycota (Supplementary
Figure 24). Twenty-nine taxa had a prevalence > 0.1 and a maximum abundance > 0.01 and
may be considered as the core mycobiota of table olives (Supplementary Table 10). Six
species (*Pichia membranifaciens, P. manshurica, Candida boidinii, Dekkera custersiana,*
Wickerhamomyces anomalus, Saccharomyces bayanus) have a prevalence >0.5 and a
maximum relative abundance >0.90, but with one exception, their median abundance is
<0.01: in fact, they may dominate some samples but may have a low abundance in others
(see Supplementary Figure 25 for a boxplot of the distribution of core species).
The most abundant species were all cultivable, and were in fact isolated from counting
plates (Supplementary table 9).
To simplify the analysis, we performed aggregation at the genus level. Core fungal genera
are shown in Supplementary Figure 26 and, remarkably, they largely overlap with those
found in a recent review (Ricciardi et al., 2025b). To avoid losing genera which are important
only in one group of olives we used the same approach used for bacterial communities (the
union of the sets of taxa which passed the filter in each group was retained): this resulted in
107 genera. Figure 6 shows a box plot of the most important families (*Pichiaceae,*
Saccharomycetales_fam_Incertae_sedis and *Phaffomycetaceae*), while other genera are
shown in Supplementary Figures 27 and 28.
As for bacterial genera there is a very wide variability within each ripeness and trade
preparation group and sometimes even within the same variety. However, some patterns
appear to emerge: *Pichia* is by far the most abundant and prevalent genus in all types of
olives which have undergone a fermentation (including green alkali treated olives) while the
distribution of other important genera is more variable and often clearly multimodal, with
very high abundance in some samples and very low in others. Specialties clearly have the
largest variety of fungal genera, possibly due to the lower duration of fermentation and the
less selective conditions.
A summary bar plot with the relative abundance of the main fungal genera in ripeness/trade
preparation groups is shown in Supplementary Figure 29. Overall, *Pichia* dominated most of
the groups, and the sum of relative abundance of *Pichia, Candida, Wickerahmomyces,*
Saccharomyces, Zygotorulaspora and *Dekkera* usually exceeded 75%. On the other hand,
there are notable differences among varieties within each group (Supplementary Figures 30-
34). The high fungal diversity in specialties, even within the same firm (most samples of
Nocellara del Belìce were provided by one firm over two years) is evident from
Supplementary Figure 33. While the composition of fungal communities of samples of the
same combination of olive variety and trade preparation is remarkably similar, samples from
some firms have a substantially different composition (Supplementary Figures 35-36).
435
436 **3.4.2 Beta-diversity of bacterial and fungal communities.**
437
438 In order to compare the microbiota of individual samples we used different ordination
439 approaches on the Bray-Curtis distance matrix. A non-monotonic Multidimensional Scaling
440 for bacterial communities in three dimensions resulted in a low stress (0.13) and the
441 configuration for the first two is shown in Figure 7.
442

443
444 **Figure 6**. Distribution of fungal genera belonging to families *Pichiaceae, Saccharomycetales*
445 *fam Incertae sedis* and *Phaffomycetaceae* in table olives used in this study. The main
446 varieties are shown (see Supplementary table 2 for abbreviations).
447
The variability within each group is evident, although groups tend to occupy a specific area
of the graph. No clear pattern for the clustering of spoiled samples (which were often but
not always far from non-spoiled samples) is evident, but this may be due to the variety of
spoilage types in our dataset. The between-firms and within-firms variability was
substantial: selected examples are shown in Supplementary Figures 37-39.
Simple ordination plots are insufficient to clearly visualize differences among varieties and
this may due to the inability to separate variance associated with predictors (chemical and
microbiological analyses, ripeness/trade preparation groups) from that due to random
variation. When constrained ordination approaches, like Canonical Correspondence Analysis
ordination using quantitative variables (microbial counts, chemical analyses) were used, only
a very low amount of the variance was explained by the first components.
However, inferential methods (PERMANOVA with adonis2) showed that a general pattern
existed and that the combination of ripeness and trade preparation significantly (0.001)
affected the composition of the bacterial communities. The effect of olive varieties within a
gives ripeness and trade preparation was also significant for all main combinations except
for turning color naturally fermented olives (Supplementary Table 22). For selected varieties
more than one producer with at least 5 samples were available: we therefore tested the
hypothesis that, within a given variety, producing firm was not a contributing factor to the

 composition of bacterial communities: the null hypothesis was discarded for all the varieties
467 tested (Itrana nera, Kalamata, Bella di Daunia and Halkidiki) (Supplementary Table 23).
468

469
470 **Figure 7.** Configuration plot for the first two dimensions of a non-monotonic
471 Multidimensional Scaling of the Bray-Curtis distance matrix of the composition of bacterial
472 communities of table olives. See Supplementary table 2 for abbreviations of olive varieties.
473
We finally explored differentially abundant taxa in olive ripeness / trade preparation groups
using DeSeq2 with alpha = 0.01, and then created contrasts for comparing groups of olives.
The results are shown in Supplementary Figures 40 to 47 and in Supplementary Table 24.
The differentially abundant genera were not necessarily those with high relative abundance
(see Figure 5 and Supplementary Figures 16-23) but the patterns found in alpha-diversity
and inferential analysis clearly emerge. HALAB and, in general alkalophilic microorganisms,
are significantly more abundant in alkali treated olives; several Pseudomonadota are
typically more abundant in black and turning color naturally fermented olives.
Heterofermentative genera, like *Levilactobacillus*, *Leuconostoc* and *Weissella* are more
abundant in black natural olives compared to all types of green olives. The pattern for
Lactobacillaceae is more variable. *Lactiplantibacillus* is significantly more abundant in black
natural olives than in alkali treated green olives (including specialties), but *Lacticaseibacillus*
is more abundant in Green natural compared to Black natural olives. Overall, a variety of
homofermentative and heterofermentative *Lactobacillaceae* are significantly more abundant
in naturally fermented olives compared to alkali treated olives.
Finally, we used PERMANOVA to test the null hypothesis of lack of differences in the
structure of bacterial communities for Itrana nera and Oliva di Gaeta (its PDO counterpart).
The null hypothesis was rejected with p = 0.025. However, this result might be due
exclusively to samples provided from one firm (F23 provided 90% of the samples for these
varieties) and might be difficult to confirm if more samples were added from the same or
other firms.

Non-constrained ordination for fungal communities was carried out using the same
approach used for bacterial communities and the configuration for the first two dimensions
is shown in Figure 8. The only group which is clearly separated is that of specialties. To
visualize more clearly the effect of spoilage and the within-group variability the same
configuration is shown separately for Black natural, Green alkali treated and Green natural
olives in Supplementary Figures 48-50. There is sometimes a large variability for the same
variety produced by different firms (this is evident for both Oliva di Gaeta PDO
(Supplementary Figure 48), and other varieties, like Halkidiki and Bella di Daunia
(Supplementary Figure 49). Samples produced by the same firm tend to be close among
them, but this not always true. For some varieties spoiled samples are clearly different from
unspoiled ones in terms of composition of fungal communities. Unfortunately, as for
bacterial communities, we do not have enough spoiled samples nor a balanced distribution
for defects to draw any specific conclusion.
Constrained ordination explained a negligible amount of variance (results not shown).
We also used PERMANOVA with adonis2 to test the null hypothesis that combination of
ripeness and trade preparation did not affect the composition of fungal communities (which
was rejected with p=0.001) and that olive varieties (within the main combinations of
ripeness and trade preparation) did not affect composition of fungal communities, which
was rejected for all main groups (Supplementary Table 25). For the three varieties (Oliva di
Gaeta, Kalamata and Bella di Daunia) for which we had more firms, we also tested the
hypothesis that there were no differences among the firms: the null hypothesis was rejected
for Kalamata and Itrana Nera, Halkidiki, but not for Bella di Daunia (Supplementary Table
26). Both findings support the hypotheses that while selective factors related to variety and
technology are important, house microbiota and specific practices within firms also play an
important role in the assembly of fungal communities.
We also performed PERMANOVA to test the null hypothesis of lack of differences in the
structure of fungal communities for Itrana nera and Oliva di Gaeta (its PDO counterpart). The
null hypothesis was rejected with p = 0.002. Again, this result might be due exclusively to
samples provide from firm F23.
We then carried out differential abundance analysis using DeSeq2 with alpha = 0.01, and
then created contrasts for comparing groups of olives. The results are shown in
Supplementary Figures 50 to 57 and in Supplementary Table 27. The differentially abundant
genera were not necessarily those with high relative abundance (see Figure 7 and
Supplementary Figures 33-34) but the patterns found in alpha-diversity and PERMANOVA
analysis emerge.
For example, many filamentous fungal genera appear among those which are differentially
abundant in one or more contrasts but their relative abundance is often very low. In other
cases (the yeast *Trichosporon* is an example) the significance in the differential abundance
test simply depends on the absence of a genus in some groups. Given the low abundance it
is likely that these species originate from the carposphere environment and do not grow
well during fermentation. If we limit the analysis to the most abundant and prevalent core
genera, we confirm the patterns found with alpha and beta-diversity analyses.
537

538
539 **Figure 8**. Configuration plot for the first two dimensions of a non-monotonic
540 Multidimensional Scaling of the Bray-Curtis distance matrix of the composition of fungal
541 communities of table olives. See Supplementary table 2 for abbreviations of olive varieties.
542
Schwanniomyces and *Geotrichum* were more abundant in Green alkali treated olives that in
any type of naturally fermented olives. *Wickerhamomyces* was significantly less abundant in
black natural compared to Green and Turning color natural and Green alkali treated olives.
Dekkera was found to be more abundant in black and turning color natural compared to
specialties. *Pichia* was more abundant in black and turning color natural compared to
specialties, but also in green alkali treated compared to green natural: this may in fact reflect
differential abundance of species within this genus (see Supplementaty Figure 25).
Citeromyces was more abundant in all types of naturally fermented olives compared to
green alkali treated olives. *Kluyveromyces* is more abundant in alkali treated compared to
naturally fermented olives. *Saccharomyces* abundance was significantly higher in green
natural olives but this result may be affected by the very high relative abundance of this
genus in all Tonda di Cagliari samples. The abundance of the genus *Candida* was lower on
average in black natural compared to both green alkali treated and green specialties:
however, this may not be true for all varieties and all species, since *C. boidinii* was very
abundant in some samples of black natural olives (see Supplementaty Figure 25).
558
559 4. **Discussion**.
560
In this work, we have presented what is, to date, the largest survey on the composition of
bacterial and fungal communities of table olives.

Our main emphasis was on olives produced in Italy, including PDO varieties such as Ascolana
del Piceno, Bella di Daunia, Nocellara del Belìce, and, especially, Oliva di Gaeta
(Supplementary Table 1). The latter is produced using black Itrana olives, with a long natural
fermentation process and, together with its non-PDO counterpart (Itrana nera), is one of the
most important table olive varieties in Italy, but studies on its microbiology are
comparatively rare and have used traditional cultivation-based approaches. Although some
authors have reported high numbers of LAB and yeasts in pulp and or/brine (Sacchi et al.,
2022; Tofalo et al., 2012) others have reported low LAB abundance in uninoculated
fermentations (Servili et al., 2006) . Addition of starters (*Lpl. plantarum/pentosus* with or
without yeasts) has been reported to improve the control of fermentation (Lanza et al.,
2021, 2020; Servili et al., 2006) . However, to our knowledge, no metataxonomic or
metagenomic studies on the microbial communities of naturally fermented Itrana olives are
available.
Given the heterogeneity of olive variety, production processes and the limited control
artisanal or semi-industrial producers have on key factors affecting contamination and
microbial community dynamics (Anagnostopoulos and Tsaltas, 2022; Perpetuini et al.,
2020) , variability among products belonging to the same trade preparations or even within
the same variety is expected, but the extent of this variability has never been explored in
great detail and attempts at relating factors like origin or trade preparation with microbiota
are rare (López-García et al., 2025) .
Indeed, almost all the olives fermented under brine in our dataset were obtained from
plastic fermenters (240-280 L) left at uncontrolled temperature. This inevitably leads to a
lack of control of the fermentation and high variability in the microbial communities.
Addition of starter cultures has often been shown to result in more uniform quality and
more rapid acidification (Benítez-Cabello et al., 2020; Campus et al., 2017; Giavalisco et al.,
2024; Leal-Sánchez et al., 2003; Vaccalluzzo et al., 2022) and should be encouraged,
provided suitably competitive cultures are available.
pH and salt content, together with nutrient availability and polyphenol contents are among
the most important ecological factors affecting the survival and growth of microorganisms in
table olives (Campus et al., 2018; Tsoungos et al., 2023) with NaCl acting as the main factor
controlling the early growth of undesirable microorganisms (Perpetuini et al., 2020).
It is interesting to note that many of our samples have combinations of pH, titratable acidity
and NaCl which are less inhibitory for microorganisms than those indicated in the trade
standards for packaged olives (IOC, 2024): this is not surprising, since one of the criteria of
inclusion in our study was the delivery of samples in their fermentation brine at or past the
time at which the fermentation was deemed to be completed. Replacement of the
fermentation brine with a preservation brine containing organic acids is common at
packaging (IOC trade standards, IOC, 2024). However, we did not find any indication that
spoilage was related to pH, TA or salt content for individual varieties.
We used a combination of selective and differential culture media for counting LAB, yeast
and molds, halophilic and potentially deleterious microorganisms (*Enterobacteriaceae* and
other contaminants). Only mMRS+NA and GYEAC showed sufficient selectivity for the
enumeration of LAB and yeasts, respectively. MRS with sodium azide is frequently used for
the enumeration of LAB in table olives (Benítez-Cabello et al., 2020; Castro et al., 2018;
Sánchez et al., 2018) although other inhibitors, like actidione (Argyri et al., 2020; Kazou et
al., 2020; Maoloni et al., 2022; Tzamourani et al., 2021) and natamycin (Anagnostopoulos et
al., 2020) have been used to make MRS selective. PCA+5% NaCl may be of limited use for the

counts of halophilic microorganisms when the counts of LAB and yeasts are high because both groups may grow on this medium. With the exception of PAB these media are commonly used for enumeration of microorganisms in table olives (see (Benítez-Cabello et al., 2020; Castro et al., 2022; Penland et al., 2020) for examples) but we are not aware of scientific papers questioning their selectivity. LAB counts were highly variable in our samples and, often, they were below the detection limits of our method. LAB are among the most important members of table olives microbiota (Portilha-Cunha et al., 2020). High counts in brine are commonly found at the beginning of fermentation, although they can decline over time (Benítez-Cabello et al., 2020; Sánchez et al., 2018) and especially in high salt brines, LAB have not been found with either culture based or culture independent approaches (Portilha-Cunha et al., 2020; Traina et al., 2024).

High yeast counts were more consistently found. This is common in olives and brines and the role of different species of yeasts can be either beneficial or detrimental, although they are frequently included in starter cultures (Anagnostopoulos and Tsaltas, 2022; Arroyo-López et al., 2012; Giavalisco et al., 2024) where is has been shown that they can stimulate the growth of LAB (Benítez-Cabello et al., 2020).

Enterobacteriaceae are commonly found at the beginning of fermentation in table olives but disappear as pH becomes lower. However, genus *Enterobacter* has been found to be both prevalent and abundant in table olives (Ricciardi et al., 2025b) and high counts of *Enterobacteriaceae* have been occasionally found in naturally fermented olives (Anagnostopoulos et al., 2020; Maoloni et al., 2022; Penland et al., 2020). In our study metataxonomic analysis revealed that DNA of several genera of *Enterobacteriaceae* was present in high proportions in several varieties even at the end of fermentation, but it is unclear if it represents viable but not cultivable portions of the microbiota.

Diversity of bacterial communities, as measured by Chao1 was within the range of those found by others for olive varieties used in this study (De Angelis et al., 2015; Cocolin et al., 2013) or for other table olives in general (de Castro et al., 2018; Maoloni et al., 2022; Medina et al., 2016; Penland et al., 2020).

Chao1 values were highly variable between and within different groups and the occurrence of significant correlations with physico-chemical variable, counts or the duration of fermentation are hard to interpret. The decrease of fungal diversity with the duration of fermentation is consistent with the hypothesis that fungal diversity decreases over time because of the dominance of a limited number of species over a much more diverse initial community. The number of studies on fungal communities of table olives is more limited than those on bacteria, and sampling criteria and methods differ. Findings on the effect of fermentation time on diversity of fungal communities are contradictory (López-García et al., 2024; Ruiz-Barba et al., 2023a, 2023b).Kamilari et al. (2023) and Penland et al. (2021) reported Chao1 for Cypriot varieties and Nyons olives in the same order of magnitude of those found in our study. Other studies are hard to compare with ours because of the use of different bioinformatic approaches or indices.

The union of the core genera for all types of olives resulted in a set of 149 genera, including several genera of LAB, HALAB (which together make up 58% of the core) and Gram-positive and -negative halophiles, but also *Pseudomonadota*. Most of these genera have been found to be abundant in one or more metataxonomic studies in table olives (see (Tsoungos et al., 2023) for recent reviews).

Lactiplantibacillus, a genus which includes the species *Lpl. pentosus*, which is considered to be well suited to the brine/table olive ecosystem because of its tolerance to salt and

polyphenols (Portilha-Cunha et al., 2020) was among the most important bacterial genera
but, in several samples was either missing or outnumbered by others. It is frequently found
in Italian table olives (Tofalo et al., 2014) and used as a starter because of its fermentation
ability, oleuropein degradation and biopreservation and probiotic potentials (Giavalisco et
al., 2024; Zotta et al., 2022) However, *Lpl. pentosus* was by far the most frequently isolated
LAB in our study (Supplementary Table 9). Given the fact that only 1-2 colonies were picked
for each sample, the disproportionate presence of this species among isolates might indicate
either that it is more frequently viable at the end of fermentation or a bias due to the
culture medium used. HALAB, like *Alkalibacterium* and *Marinilactibacillus* are abundant and
prevalent in alkali treated olives, as confirmed by other researchers (López-García et al.,
2022; Lucena-Padrós and Ruiz-Barba, 2016; Portilha-Cunha et al., 2020), but we also found
them in naturally fermented olives. *Lacticaseibacillus*, which has also potential for use as a
starter and functional culture in table olives (Pino et al., 2019) was also among the dominant
genera. Several heterofermentative LAB were often (*Lentilactobacillus*) or occasionally
(*Limosilactobacillus, Levilactobacillus, Paucilactobacillus* and *Leuconostoc*) dominant or
subdominant, but their importance in spoilage or in promoting the quality of table olives is
unclear, although *Levilactobacillus* and *Leuconostoc* have been tested as starters in Greek
olives (Chytiri et al., 2020; Vougiouklaki et al., 2024). A high proportion of *Lentilactobacillus*
may be of concern, because this genus has been associated with spoilage and, in protein rich
environments, is associated with the production of biogenic amines (O'Sullivan et al., 2015;
Penland et al., 2021; Suzzi and Gardini, 2003).
Celerinatantimonas, which is abundant in different green and black varieties of olives (e.g.,
black Nyons and Kalamata, green naturally fermented Aloreña de Malaga) (Kazou et al.,
2020; Medina et al., 2016; Penland et al., 2020), and may contribute to spoilage (de Castro
et al., 2022) was abundant and prevalent in Picholine style olives, turning color and black
naturally fermented olives, but to a lower extent in green naturally fermented and alkali
treated olive, in which *Halomonas, Vibrio, Salinicola* and *Idiomarina* were prevalent and
occasionally abundant. This confirms previous findings (Arroyo-López et al., 2021; Lucena-
Padrós et al., 2015; Posada-Izquierdo et al., 2021; Ricciardi et al., 2025b; Ruiz-Barba et al.,
2023b; Tsoungos et al., 2023). Some of these genera have been associated with spoilage,
and more systematic work is needed to clarify their role in table olive fermentation.
Because of their phenolic content, salt concentration, and, for most varieties, of their acidic
pH, table olives are a relatively safe product and have been rarely associated with food-
borne outbreaks, for example botulism (Cawthorne et al., 2005) and salmonellosis
(Donachie et al., 2018). The usual low abundance of pathogenic microorganisms in foods
and the low taxonomic resolution of amplicon targeted metagenomics pose challenges to
the use of metagenomic approaches in the detection of microorganisms relevant for food
safety (Ferrocino et al., 2022; Wang et al., 2025). We confirmed our previous finding of the
occurrence of sequences assigned to bacterial genera containing pathogenic microorganisms
(Ricciardi et al., 2025b). The occurrence of *Salmonella* at relative abundance close to 0.1% in
at least two samples, together with previous reports on outbreaks caused by *Salm. enterica*
from olives (Donachie et al., 2018) raises some concerns on the occurrence of this pathogen
in olive producing environments.
Some fungal genera (*Pichia, Candida, Saccharomyces, Wickerhamomyces, Dekkera*) were
highly prevalent (>0.50) in table olives, in agreement with what has been found by many
studies (see (Ricciardi et al., 2025b; Tsoungos et al., 2023) for recent reviews), although their
abundance was very variable. The occurrence of the first 4 genera is very common and they

are among the dominant yeasts in olives (Ricciardi et al., 2025b; Tsoungos et al., 2023) and
some species are included in starter cultures. On the other hand, *Zygototorulaspora* has
been found as a subdominant or dominant species only occasionally (Arroyo-López et al.,
2016; Mougiou et al., 2023). *Dekkera/Brettanomyces*, a genus which includes species which
cause spoilage in alcoholic beverages (Tubia et al., 2018) is usually found at low abundances
in olives and brines (de Castro et al., 2018; Medina et al., 2018) but can sometimes become
subdominant (Michailidou et al., 2021), as in several varieties of naturally fermented olives
in our study.
We revealed a striking variability of the composition of both bacterial and fungal microbiota,
even within the same olive variety, suggesting the presence of producer-specific, or even
barrel-specific, microbial fingerprint. Bar plots of abundances of dominating genera can only
provide a general indication of the composition of the microbiota of a specific olive group or
variety. Due to differences in methods, it is hard to compare our results with those of other
authors, although combining all data in well annotated databases (Ricciardi et al., 2025b)
could be of assistance in this task. Overall, the patterns of genera and species found in our
study match well with those found by other authors (see (Ricciardi et al., 2025b; Tsoungos et
al., 2023) for recent compilations of data).
Only few metataxonomic studies have used either several samples for the same variety from
the same firm or samples for the same variety from several firms (see (Argyri et al., 2020;
Arroyo-López et al., 2021; Kazou et al., 2020; Lucena-Padrós and Ruiz-Barba, 2019; Penland
et al., 2021, 2020) for examples). Although common patterns may emerge in terms of shared
composition at the species or genus level, differences emerge within the same firm or
variety and have been attributed, among other things, to biogeographical patterns. In our
case, we exclude that this variability is a merely technical issue, because Bray-Curtis distance
between technical replicates was very low. The diversity, even for the same variety within
the same firm, is well reflected in the ordination plots shown in Figures 7 and 8 and in
Supplementary Figures 37-39 and 48-50.
Our dataset included spoiled samples, although sensory evaluation was not carried out. It is
well known that spoilage affects the microbiota and, in some cases, differential
increases/decreases of selected taxa have been associated with specific types of spoilage
(Arroyo-López et al., 2021; Castro et al., 2022, 2018; Penland et al., 2021). In our data set,
although, at least for fungi, the microbial communities of spoiled samples were often
visually different from those of unspoiled samples (see Figure 8), the proportion of spoiled
samples was too low and the spoilage categories too varied to draw solid conclusions or to
perform inferential analysis.
Despite the variability, the use of inferential methods allowed us to clarify that combinations
of olive ripeness and trade preparation across all varieties and firms, varieties within a given
combination of ripeness and trade preparation, and firms within a same variety were all
factors which significantly affected the composition of bacterial and fungal communities.
Differential abundance analysis was harder to interpret but some patterns clearly emerged,
like the significantly higher abundance of HALAB in alkali treated olives compared to all
other groups, and of a diverse group of *Lactobacillaceae* and of several *Proteobacteria* in
naturally fermented olives. The use of inferential methods in the analysis of microbial
communities of table olives is comparatively rare. Most studies are small and have an
observational approach (Ricciardi et al., 2025b). Comparisons between started and non-
started batches, different processes and ripening stages within the same variety (López-
García et al., 2022; Penland et al., 2020; Ruiz-Barba et al., 2023b), different locations (López-

García et al., 2022; Lucena-Padrós and Ruiz-Barba, 2019) or between spoiled or unspoiled
batches for the same variety (Penland et al., 2021) are more common as are studies on the
geographical origin, while, to our knowledge, nobody has ever undertaken a large survey
encompassing as many different types and varieties of olives as in our study. (Kamilari et al.,
2023) identified differentially abundant genera in three Cypriot varieties; in their case,
however, olive variety was confounded with ripeness and trade preparation. Recently
(López-García et al., 2025) applied a machine learning approach on metataxonomic data
from several studies to evaluate the effect of several variables on bacterial communities of
table olives, and the findings on olive processing identify some HALAB as the most important
in discriminating alkali treated from naturally fermented olives. Finally, several authors have
proposed that metataxonomic analysis of microbial communities may assist in identifying
the geographical origin of table olives (Argyri et al., 2020; Ferrocino et al., 2025; Kamilari et
al., 2023; Kazou et al., 2020) thus reinforcing an idea of *terroir,* a concept which has
originally developed for wine (Belda et al., 2017) and the existence of measurable
biogeographical patterns for table olives. This would be consistent with the idea that both
dispersal and selection play a role in shaping the microbial communities and thus the
organoleptic properties of table olives, which is reinforced by our data. However, while there
is sufficient proof that chemometric approaches can assist in the authentication of table
olives (Kazou et al., 2025), according to our findings, when the technology is not particularly
selective Italian PDO varieties (Bella di Daunia, Nocellara del Belíce and especially, Oliva di
Gaeta) can hardly be discriminated from other similar varieties based on microbiota
composition. Better integration of metataxonomic data in large metastudies (López-García et
al., 2025; Ricciardi et al., 2025b) and integration with metabolomic data may improve this
discrimination (Beteinakis et al., 2024).
775
5. Conclusions
We have presented the largest survey, to date, on the composition of microbial communities
of table olives. It is also the first paper providing information on the structure of the
microbial communities of Itrana olives, one of the most prominent table olive varieties in
Italy. It fully reveals the diversity of bacterial and fungal taxa which may occur in a variety of
products over a wide range in geographical origin and technology. We feel that our data
confirm that, although selective factors (olive ripeness variety and technological factors, like
alkali treatment, salting etc.) play a role in shaping the microbiota, dispersal limitation and
insufficient control of key ecological factors (temperature, duration of fermentation, salt)
result in a striking variability. This is especially evident for lactic acid bacteria, whose
diversity goes well beyond the widespread occurrence of species, like *Lpl. pentosus,* which
are traditionally used as starters. Apparently, a wider variety of LAB and HALAB are able to
colonize and thrive in this ecological niche, although some seem specifically adapted to
some trade preparations, compared to yeasts, whose diversity was significantly lower, with a
limited set of species dominating most varieties. Differentially abundant genera and species
should be used in designed experiments to clarify to which extent selection, in the absence
of dispersal limitation and drift, due to the potential stochastic nature of contamination
patterns of barrels used for fermentation within the same firm, governs the establishment of
dominant taxa and which is the impact on the safety and sensory quality of table olives. Our
data also prove that small surveys (low number of firms and samples) may provide a skewed
and limited view even when a study focuses on a single variety. Even the selection of
biomarkers for the evaluation of authenticity of PDO varieties may prove tricky: we were

able to detect significant differences between non-PDO and PDO naturally fermented black
Itrana olives but the sampling was too unbalanced to warrant the use of machine learning
approaches. Better control of contamination and of production technologies may allow a
better differentiation among olive varieties.
802

803 **Data availability**.
804 Sequence data have been deposited in NCBI SRA with bioproject accession PRJNA1292811.
805 The accession list of the runs used in this study is available in Supplementary data and
806 figures. The metataxonomic data have been integrated in the OliveFMBN database
807 (https://github.com/ep142/Metaolive/tree/main/olive_FMBN)
808

Acknowledgements.
The authors gratefully acknowledge the skilled technical assistance of Ms. Anita Aliano.
This work was carried out within the PRIN 2022 Project METAOlive 2022NN28ZZ and
received funding from the European Union Next-GenerationEU, CUP C53D23005460006
(PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2,
INVESTIMENTO 1.4 – D.D. 1048 14/07/2023).

815
816

6. References

Abarenkov, K., Nilsson, R.H., Larsson, K.-H., Taylor, A.F.S., May, T.W., Frøslev, T.G.,
Pawlowska, J., Lindahl, B., Põldmaa, K., Truong, C., Vu, D., Hosoya, T., Niskanen, T.,
Piirmann, T., Ivanov, F., Zirk, A., Peterson, M., Cheeke, T.E., Ishigami, Y., Jansson, A.T.,
Jeppesen, T.S., Kristiansson, E., Mikryukov, V., Miller, J.T., Oono, R., Ossandon, F.J.,
Paupério, J., Saar, I., Schigel, D., Suija, A., Tedersoo, L., Kõljalg, U., 2023. The UNITE
database for molecular identification and taxonomic communication of fungi and other
eukaryotes: sequences, taxa and classifications reconsidered. Nucleic Acids Res. 52,
D791–D797. https://doi.org/10.1093/nar/gkad1039.

Alfonzo, A., Alongi, D., Prestianni, R., Pirrone, A., Naselli, V., Viola, E., Pasquale, C.D., Croce,
F.L., Gaglio, R., Settanni, L., Francesca, N., Moschetti, G., 2024. Enhancing the quality and
safety of Nocellara del Belice green table olives produced using the Castelvetrano
method. Food Microbiol. 120, 104477. https://doi.org/10.1016/j.fm.2024.104477.

Alfonzo, A., Craparo, V., Prestianni, R., Naselli, V., Seminerio, V., Pirrone, A., Croce, F.L.,
Settanni, L., Moschetti, G., Francesca, N., 2023. Application of technological protocols on
an industrial scale to improve Seville-style table olive production in Italy and Spain. Appl.
Food Res. 100349. https://doi.org/10.1016/j.afres.2023.100349.

Anagnostopoulos, D.A., Kamilari, E., Tsaltas, D., 2020. Evolution of bacterial communities,
physicochemical changes and sensorial attributes of natural whole and cracked Picual
table olives during spontaneous and inoculated fermentation. Front. Microbiol. 11, 1128.
https://doi.org/10.3389/fmicb.2020.01128.

Anagnostopoulos, D.A., Tsaltas, D., 2022. Current status, recent advances, and main
challenges on table olive fermentation: the present meets the future. Front. Microbiol.
12, 797295. https://doi.org/10.3389/fmicb.2021.797295.

Argyri, K., Doulgeraki, A.I., Manthou, E., Grounta, A., Argyri, A.A., Nychas, G.-J.E., Tassou,
C.C., 2020. Microbial diversity of fermented greek table olives of Halkidiki and
Konservolia varieties from different regions as revealed by metagenomic analysis.
Microorganisms 8, 1241. https://doi.org/10.3390/microorganisms8081241.

Arroyo-López, F.N., Benítez-Cabello, A., Romero-Gil, V., Rodríguez-Gómez, F., Garrido-
Fernández, A., 2021. Delving into the bacterial diversity of spoiled green Manzanilla
Spanish-style table olive fermentations. Int. J. Food. Microbiol. 359, 109415.
https://doi.org/10.1016/j.ijfoodmicro.2021.109415.

Arroyo-López, F.N., Medina, E., Ruiz-Bellido, M.Á., Romero-Gil, V., Montes-Borrego, M.,
Landa, B.B., 2016. Enhancement of the knowledge on fungal communities in directly
brined Aloreña de Málaga green olive fermentations by metabarcoding analysis. PLoS
One 11, e0163135. https://doi.org/10.1371/journal.pone.0163135.

Arroyo-López, F.N., Romero-Gil, V., Bautista-Gallego, J., Rodriguez-Gómez, F., Jiménez-Díaz,
R., Garcia, P.G., Querol, A., Fernandez, A.G., 2012. Yeasts in table olive processing:
desirable or spoilage microorganisms? Int. J. Food Microbiol. 160, 42-49.
https://doi.org/10.1016/j.ijfoodmicro.2012.08.003.

Ballester, E., Ribes, S., Barat, J.M., Fuentes, A., 2021. Spoilage yeasts in fermented
vegetables: conventional and novel control strategies. Eur. Food Res. Technol. 1-14.
https://doi.org/10.1007/s00217-021-03888-7.

Belda, I., Zarraonaindia, I., Perisin, M., Palacios, A., Acedo, A., 2017. From vineyard soil to
wine fermentation: microbiome approximations to explain the "terroir" concept. Front.
Microbiol. 8, 821. https://doi.org/10.3389/fmicb.2017.00821.

Benítez-Cabello, A., Calero-Delgado, B., Rodríguez-Gómez, F., Bautista-Gallego, J., Garrido-
Fernández, A., Jiménez-Díaz, R., Arroyo-López, F.N., 2020. The use of multifunctional
yeast-lactobacilli starter cultures improves fermentation performance of Spanish-style
green table olives. Food Microbiol. 91, 103497.
https://doi.org/10.1016/j.fm.2020.103497.

Beteinakis, S., Papachristodoulou, A., Stathopoulos, P., Mikros, E., Halabalaki, M., 2024. A
multilevel LC-HRMS and NMR correlation workflow towards foodomics advancement:
application in table olives. Talanta 280, 126641.
https://doi.org/10.1016/j.talanta.2024.126641.

Blighe, K., Lun, A ., 2025. PCAtools: PCAtools: Everything Principal Components Analysis.
doi:10.18129/B9.bioc.PCAtools. https://doi.org/10.18129/B9.bioc.PCAtools.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016.
DADA2: High-resolution sample inference from Illumina amplicon data. Nat. Methods 13,
581-583. https://doi.org/10.1038/nmeth.3869.

Campus, M., Cauli, E., Scano, E., Piras, F., Comunian, R., Paba, A., Daga, E., Salvo, R.D.,
Sedda, P., Angioni, A., Zurru, R., 2017. Towards controlled fermentation of table olives:
LAB starter driven process in an automatic pilot processing plant. Food Bioprocess
Technol. 10, 1063-1073. https://doi.org/10.1007/s11947-017-1882-7.

Campus, M., Değirmencioğlu, N., Comunian, R., 2018. Technologies and trends to improve
table olive quality and safety. Front. Microbiol. 9, 617.
https://doi.org/10.3389/fmicb.2018.00617.

Cawthorne, A., Celentano, L.P., D'Ancona, F., Bella, A., Massari, M., Anniballi, F., Fenicia, L.,
Aureli, P., Salmaso, S., 2005. Botulism and preserved green olives. Emerg. Infect. Dis. 11,
781-782. https://doi.org/10.3201/eid1105.041088.

Chytiri, A., Tasioula-Margari, M., Bleve, G., Kontogianni, V.G., Kallimanis, A., Kontominas,
M.G., 2020. Effect of different inoculation strategies of selected yeast and LAB cultures
on Conservolea and Kalamàta table olives considering phenol content, texture, and
sensory attributes. J. Sci. Food Agric. 100, 926-935. https://doi.org/10.1002/jsfa.10019.

Ciafardini, G., Zullo, B.A., 2019. Use of selected yeast starter cultures in industrial-scale
processing of brined Taggiasca black table olives. Food Microbiol. 84, 103250.
https://doi.org/10.1016/j.fm.2019.103250.

Cocolin, L., Alessandria, V., Botta, C., Gorra, R., De Filippis, F., Ercolini, D., Rantsiou, K., 2013.
NaOH-debittering induces changes in bacterial ecology during table olives fermentation.
PLoS One 8, e69074. https://doi.org/10.1371/journal.pone.0069074.s004.

Comunian, R., Ferrocino, I., Paba, A., Daga, E., Campus, M., Salvo, R.D., Cauli, E., Piras, F.,
Zurru, R., Cocolin, L., 2017. Evolution of microbiota during spontaneous and inoculated
Tonda di Cagliari table olives fermentation and impact on sensory characteristics. Lwt -
Food Sci. Technol. 84, 64-72. https://doi.org/10.1016/j.lwt.2017.05.039.

Davis, N.M., Proctor, D.M., Holmes, S.P., Relman, D.A., Callahan, B.J., 2018. Simple statistical
identification and removal of contaminant sequences in marker-gene and metagenomics
data. Microbiome 6, 226. https://doi.org/10.1186/s40168-018-0605-2.

De Angelis, M., Campanella, D., Cosmai, L., Summo, C., Rizzello, C.G., Caponio, F., 2015.
Microbiota and metabolome of un-started and started Greek-type fermentation of Bella
di Cerignola table olives. Food Microbiol. 52, 18-30.
https://doi.org/10.1016/j.fm.2015.06.002.

de Castro, A., Ruiz-Barba, J.L., Romero, C., Sánchez, A.H., García, P., Brenes, M., 2022.
Formation of gas pocket defect in Spanish-style green olives by the halophile
Celerinatantimonas sp. Food Control 136, 108868.
https://doi.org/10.1016/j.foodcont.2022.108868.

de Castro, A., Sánchez, A.H., López-López, A., Cortés-Delgado, A., Medina, E., Montaño, A.,
2018. Microbiota and metabolite profiling of spoiled spanish-style Green table Olives.
Metabolites 8, 73. https://doi.org/10.3390/metabo8040073.

Donachie, A., Melillo, T., Bubba, L., Hartman, H., Borg, M.-L., 2018. National outbreak of
Salmonella Give linked to a local food manufacturer in Malta, October 2016. Epidemiol.
Infect. 146, 1425-1432. https://doi.org/10.1017/s0950268818001656.

Ferrocino, I., Buzzanca, D., Pagiati, L., Kazou, M., Georgalaki, M., Hatzopoulos, I., Tsakalidou,
E., 2025. The microbial terroir of the Greek olive varieties. Int. J. Food Microbiol. 441,
111332. https://doi.org/10.1016/j.ijfoodmicro.2025.111332.

Ferrocino, I., Rantsiou, K., Cocolin, L., 2022. Microbiome and -omics application in food
industry. Int J. Food Microbiol. 377, 109781.
https://doi.org/10.1016/j.ijfoodmicro.2022.109781.

Gänzle, M., 2022. The periodic table of fermented foods: limitations and opportunities.
Appl. Microbiol. Biotechnol. 1-12. https://doi.org/10.1007/s00253-022-11909-y.

Garrido-Fernández, A., Benítez-Cabello, A., Rodríguez-Gómez, F., Jiménez-Díaz, R., Arroyo-
López, F.N., Morales, M.L., 2021. Relating starter cultures to volatile profile and potential
markers in green Spanish-style table olives by compositional data analysis. Food
Microbiol. 94, 103659. https://doi.org/10.1016/j.fm.2020.103659.

Giavalisco, M., Lavanga, E., Ricciardi, A., Zotta, T., 2024. Starter cultures for the production
of fermented table olives: current status and future perspectives. Fermentation 10, 351.
https://doi.org/10.3390/fermentation10070351.

Hernández-Velázquez, R., Flörl, L., Lavrinienko, A., Sebechlebská, Z., Merk, L., Greppi, A.,
Bokulich, N.A., 2024. The future is fermented: microbial biodiversity of fermented foods
is a critical resource for food innovation and human health. Trends Food Sci. Technol.
150, 104569. https://doi.org/10.1016/j.tifs.2024.104569.

Kamilari, E., Anagnostopoulos, D.A., Tsaltas, D., 2023. Fermented table olives from Cyprus:
microbiota profile of three varieties from different regions through metabarcoding
sequencing. Front. Microbiol. 13, 1101515.
https://doi.org/10.3389/fmicb.2022.1101515.

Kassambara, A., 2023. rstatix: Pipe-Friendly Framework for Basic Statistical Tests.
https://doi.org/10.32614/CRAN.package.rstatix

Kassambara, A., 2025. _ggpubr: 'ggplot2' Based Publication Ready Plots.
https://doi.org/10.32614/CRAN.package.ggpubr.

Kauserud, H., 2023. ITS alchemy: On the use of ITS as a DNA marker in fungal ecology.
Fungal Ecol. 65, 101274. https://doi.org/10.1016/j.funeco.2023.101274.

Kazou, M., Psarafti, K., Panagou, E.Z., 2025. The future of the table olive sector: unlocking
the potential of process analytical technology (PAT) in the era of digital transformation.
Eur. Food Res. Technol. 1-16. https://doi.org/10.1007/s00217-025-04892-x.

Kazou, M., Tzamourani, A., Panagou, E.Z., Tsakalidou, E., 2020. Unraveling the Microbiota of
Natural Black cv. Kalamata Fermented Olives through 16S and ITS Metataxonomic
Analysis. Microorganisms 8, 672. https://doi.org/10.3390/microorganisms8050672.

Langgut, D., Garfinkel, Y., 2022. 7000-year-old evidence of fruit tree cultivation in the Jordan
Valley, Israel. Sci. Rep. 12, 7463. https://doi.org/10.1038/s41598-022-10743-6.

Lanza, B., Marco, S.D., Bacceli, M., Serio, M.G.D., Loreto, G.D., Cellini, M., Simone, N., 2021.
Lactiplantibacillus plantarum used as single, multiple, and mixed starter combined with
Candida boidinii for table olive fermentations: chemical, textural, and sensorial
characterization of final products. Fermentation 7, 239.
https://doi.org/10.3390/fermentation7040239.

Lanza, B., Zago, M., Marco, S.D., Loreto, G.D., Cellini, M., Tidona, F., Bonvini, B., Bacceli, M.,
Simone, N., 2020. Single and multiple inoculum of *Lactiplantibacillus plantarum* strains in
table olive lab-scale fermentations. Fermentation 6, 126.
https://doi.org/10.3390/fermentation6040126.

Leal-Sánchez, M.V., Ruiz-Barba, J.L., Sánchez, A.H., Rejano, L., Jimenéz-Diaz, R., Garrido, A.,
2003. Fermentation profile and optimization of green olive fermentation using
Lactobacillus plantarum LPCO10 as a starter culture. Food Microbiol. 20, 421-430.
https://doi.org/10.1016/S0740-0020(02)00147-8.

Lee, H.M., Lee, Y., 2008. A differential medium for lactic acid-producing bacteria in a mixed
culture. Lett. Appl. Microbiol. 46, 676-681. https://doi.org/10.1111/j.1472-
765x.2008.02371.x.

López-García, E., Benítez-Cabello, A., Arroyo-López, F.N., 2025. Applying machine learning to
classify table olives using bacterial metataxonomic data. npj Sci. Food 9, 121.
https://doi.org/10.1038/s41538-025-00496-7.

López-García, E., Benítez-Cabello, A., Rodríguez-Gómez, F., Romero-Gil, V., Garrido-
Fernández, A., Jiménez-Díaz, R., Arroyo-López, F.N., 2022. Bacterial metataxonomic
analysis of industrial Spanish-style green table olive fermentations. Food Control 137,
108969. https://doi.org/10.1016/j.foodcont.2022.108969.

López-García, E., Romero-Gil, V., Arroyo-López, F.N., Benítez-Cabello, A., 2024. Impact of
lactic acid bacteria inoculation on fungal diversity during Spanish-style green table olive
fermentations. Int. J. Food Microbiol. 417, 110689.
https://doi.org/10.1016/j.ijfoodmicro.2024.110689.

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. 15, 550.
https://doi.org/10.1186/s13059-014-0550-8.

Lucena-Padrós, H., González, J.M., Caballero-Guerrero, B., Ruiz-Barba, J.L., Maldonado-
Barragán, A., 2015. *Vibrio olivae* sp. nov., isolated from Spanish-style green-olive
fermentations. Int. J. Syst. Evol. Microbiol. 65, 1895-1901.
https://doi.org/10.1099/ijs.0.000196.

Lucena-Padrós, H., Ruiz-Barba, J.L., 2016. Diversity and enumeration of halophilic and
alkaliphilic bacteria in Spanish-style green table-olive fermentations. Food Microbiol. 53,
53-62. https://doi.org/10.1016/j.fm.2015.09.006.

Lucena-Padrós, H., Ruiz-Barba, J.L., 2019. Microbial biogeography of Spanish-style green
olive fermentations in the province of Seville, Spain. Food Microbiol. 82, 259-268.
https://doi.org/10.1016/j.fm.2019.02.004.

Maoloni, A., Cardinali, F., Milanović, V., Osimani, A., Garofalo, C., Ferrocino, I., Corvaglia,
M.R., Cocolin, L., Aquilanti, L., 2022. Microbial dynamics and key sensory traits of
laboratory-scale co-fermented green olives (*Olea europaea* L. cv. *Ascolana tenera*) and
sea fennel (*Crithmum maritimum* L.). Food Biosci. 50, 102077.
https://doi.org/10.1016/j.fbio.2022.102077.

Martorana, A., Alfonzo, A., Settanni, L., Corona, O., Croce, F.L., Caruso, T., Moschetti, G.,
Francesca, N., 2015. An innovative method to produce green table olives based on "pied
de cuve" technology. Food Microbiol. 50, 126-140.
https://doi.org/10.1016/j.fm.2015.03.008.

McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. PLoS One 8, e61217.
https://doi.org/10.1371/journal.pone.0061217.

Medina, E., Ruiz-Bellido, M.A., Romero-Gil, V., Rodríguez-Gómez, F., Montes-Borrego, M.,
Landa, B.B., Arroyo-López, F.N., 2016. Assessment of the bacterial community in directly
brined *Aloreña de Málaga* table olive fermentations by metagenetic analysis. Int. J. Food
Microbiol. 236, 47-55. https://doi.org/10.1016/j.ijfoodmicro.2016.07.014.

Medina, E., Brenes, M., García-García, P., Romero, C., de Castro, A., 2018. Microbial ecology
along the processing of Spanish olives darkened by oxidation. Food Control 86, 35-41.
https://doi.org/10.1016/j.foodcont.2017.10.035.

Michailidou, S., Trikka, F., Pasentsis, K., Petrovits, G.E., Kyritsi, M., Argiriou, A., 2021. Insights
into the evolution of Greek style table olives microbiome stored under modified
atmosphere: Biochemical implications on the product quality. Food Control 130, 108286.
https://doi.org/10.1016/j.foodcont.2021.108286.

Montaño, A., Cortés-Delgado, A., Sánchez, A.H., Ruiz-Barba, J.L., 2021. Production of volatile
compounds by wild-type yeasts in a natural olive-derived culture medium. Food
Microbiol. 98, 103788. https://doi.org/10.1016/j.fm.2021.103788.

Mougiou, N., Tsoureki, A., Didos, S., Bouzouka, I., Michailidou, S., Argiriou, A., 2023.
Microbial and biochemical profile of different types of greek table olives. Foods 12, 1527.
https://doi.org/10.3390/foods12071527.

Oksanen, J., Simpson, G., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O'Hara, R.,
Solymos, P., Stevens, M., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B.,
Borcard, D., Borman, T., Carvalho, G., Chirico, M., De Caceres, M., Durand, S., Evangelista,
H., FitzJohn, R., Friendly, M., Furneaux, B., Hannigan, G., Hill, M., Lahti, L., Martino, C.,
McGlinn, D., Ouellette, M., Ribeiro Cunha, E., Smith, T., Stier, A., Ter Braak, C., Weedon, J.,
2025. _vegan: Community Ecology Package.
https://doi.org/10.32614/CRAN.package.vegan.

O'Sullivan, D.J., Fallico, V., O'sullivan, O., McSweeney, P.L.H., Sheehan, J.J., Cotter, P.D.,
Giblin, L., 2015. High-throughput DNA sequencing to survey bacterial histidine and
tyrosine decarboxylases in raw milk cheeses. BMC Microbiol. 15, 266.
https://doi.org/10.1186/s12866-015-0596-0.

Paba, A., Chessa, L., Daga, E., Campus, M., Bulla, M., Angioni, A., Sedda, P., Comunian, R.,
2020. Do best-selected strains perform table olive fermentation better than undefined
biodiverse starters? A comparative study. Foods 9, 135.
https://doi.org/10.3390/foods9020135.

Parente, E., Zotta, T., Giavalisco, M., Ricciardi, A., 2023. Metataxonomic insights in the
distribution of *Lactobacillaceae* in foods and food environments. Int. J. Food Microbiol.
391, 110124. https://doi.org/10.1016/j.ijfoodmicro.2023.110124.

Penland, M., Deutsch, S.-M., Falentin, H., Pawtowski, A., Poirier, E., Visenti, G., Meur, C.L.,
Maillard, M.-B., Thierry, A., Mounier, J., Coton, M., 2020. Deciphering microbial
community dynamics and biochemical changes during nyons black olive natural
fermentations. Front. Microbiol. 11, 586614.
https://doi.org/10.3389/fmicb.2020.586614.

Penland, M., Mounier, J., Pawtowski, A., Tréguer, S., Deutsch, S.-M., Coton, M., 2021. Use of
metabarcoding and source tracking to identify desirable or spoilage autochthonous
microorganism sources during black olive fermentations. Food Res. Int. 144, 110344.
https://doi.org/10.1016/j.foodres.2021.110344.

Penland, M., Pawtowski, A., Pioli, A., Maillard, M.-B., Debaets, S., Deutsch, S.-M., Falentin,
H., Mounier, J., Coton, M., 2022. Brine salt concentration reduction and inoculation with
autochthonous consortia: impact on protected designation of origin nyons black table
olive fermentations. Food Res. Int. 155, 111069.
https://doi.org/10.1016/j.foodres.2022.111069.

Perpetuini, G., Prete, R., Garcia-Gonzalez, N., Alam, M.K., Corsetti, A., 2020. Table olives
more than a fermented food. Foods 9, 178. https://doi.org/10.3390/foods9020178.

Pino, A., De Angelis, M., Todaro, A., Hoorde, K.V., Randazzo, C.L., Caggia, C., 2018.
Fermentation of Nocellara Etnea table olives by functional starter cultures at different
low salt concentrations. Front. Microbiol. 9, 1125.
https://doi.org/10.3389/fmicb.2018.01125.

Pino, A., Vaccalluzzo, A., Solieri, L., Romeo, F.V., Todaro, A., Caggia, C., Arroyo-López, F.N.,
Bautista-Gallego, J., Randazzo, C.L., 2019. Effect of sequential inoculum of beta-
glucosidase positive and probiotic strains on brine fermentation to obtain low salt Sicilian
table olives. Front. Microbiol. 10, 174. https://doi.org/10.3389/fmicb.2019.00174.

Portilha-Cunha, M.F., Macedo, A.C., Malcata, F.X., 2020. A review on adventitious lactic acid
bacteria from table olives. Foods 9, 948. https://doi.org/10.3390/foods9070948.

Posada-Izquierdo, G.D., Valero, A., Arroyo-López, F.N., González-Serrano, M., Ramos-
Benítez, A.M., Benítez-Cabello, A., Rodríguez-Gómez, F., Jimenez-Diaz, R., García-Gimeno,
R.M., 2021. Behavior of *Vibrio* spp. in table olives. Front. Microbiol. 12, 650754.
https://doi.org/10.3389/fmicb.2021.650754.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O.,
2013. The SILVA ribosomal RNA gene database project: improved data processing and
web-based tools. Nucleic Acids Res.41, D590-6. https://doi.org/10.1093/nar/gks1219.

R Core Team, 2025. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Randazzo, C.L., Todaro, A., Pino, A., Pitino, I., Corona, O., Caggia, C., 2017. Microbiota and
metabolome during controlled and spontaneous fermentation of Nocellara Etnea table
olives. Food Microbiol. 65, 136-148. https://doi.org/10.1016/j.fm.2017.01.022.

Ricciardi, A., Parente, E., Tramutola, A., Guidone, A., Ianniello, R.G., Pavlidis, D., Tsakalidou,
E., Zotta, T., 2014. Evaluation of a differential medium for the preliminary identification
of members of the *Lactobacillus plantarum* and *Lactobacillus casei* groups. Ann.
Microbiol. 1-10. https://doi.org/10.1007/s13213-014-1004-y.

Ricciardi, A., López, F.N.A., Giavalisco, M., Pietrafesa, R., Parente, E., 2025a. Determining the
core bacterial and fungal genera in table olive fermentations. bioRxiv 2025.03.16.643505.
https://doi.org/10.1101/2025.03.16.643505.

Ricciardi, A., López, F.N.A., Giavalisco, M., Pietrafesa, R., Parente, E., 2025b. Determining the
core bacterial and fungal genera in table olive fermentations. Int. J. Food Microbiol. 442,
111344. https://doi.org/10.1016/j.ijfoodmicro.2025.111344.

Rodríguez-Gómez, F., Valero, A., Lara, E.V., Marín, A., Ramírez, E.M., 2023. LP309 a new
strain of *Lactiplantibacillus pentosus* that improves the lactic fermentation of Spanish-
style table olives. J. Food Sci. 88, 5191-5202. https://doi.org/10.1111/1750-3841.16802.

Rué, O., Coton, M., Dugat-Bony, E., Howell, K., Irlinger, F., Legras, J.-L., Loux, V., Michel, E.,
Mounier, J., Neuvéglise, C., Sicard, D., 2023. Comparison of metabarcoding taxonomic

markers to describe fungal communities in fermented foods. Peer Community J. 3, e97.
https://doi.org/10.24072/pcjournal.321.

Ruiz-Barba, J.L., Jimenéz-Diaz, R., 2012. A novel *Lactobacillus pentosus*-paired starter culture
for Spanish-style green olive fermentation. Food Microbiol. 30, 253-259.
https://doi.org/10.1016/j.fm.2011.11.004.

Ruiz-Barba, J.L., Sánchez, A.H., López-López, A., Cortés-Delgado, A., Montaño, A., 2023a.
Microbial and chemical characterization of natural-style green table olives from the
Gordal, Hojiblanca and Manzanilla cultivars. Foods 12, 2386.
https://doi.org/10.3390/foods12122386.

Ruiz-Barba, J.L., Sánchez, A.H., López-López, A., Cortés-Delgado, A., Montaño, A., 2023b.
Microbial community and volatilome changes in brines along the spontaneous
fermentation of Spanish-style and natural-style green table olives (Manzanilla cultivar).
Food Microbiol. 113, 104286. https://doi.org/10.1016/j.fm.2023.104286.

Ruiz-Barba, J.L., Cortés-Delgado, A., Sánchez, A.H., López-López, A., Montaño, A., 2024.
Impact of selected wild yeasts starters on the volatilome and phenolic contents of
Gordal, Manzanilla and Hojiblanca naturally fermented green olives. LWT 195, 115811.
https://doi.org/10.1016/j.lwt.2024.115811.

Sacchi, R., Corrado, G., Basile, B., Mandarello, D., Ambrosino, M.L., Paduano, A., Savarese,
M., Caporaso, N., Aponte, M., Genovese, A., 2022. Effect of salt addition and
fermentation time on phenolics, microbial dynamics, volatile organic compounds, and
sensory properties of the PDO table olives of Gaeta (Italy). Molecules 27, 8100.
https://doi.org/10.3390/molecules27228100.

Sánchez, A.H., López-López, A., Cortés-Delgado, A., Beato, V.M., Medina, E., de Castro, A.,
Montaño, A., 2018. Effect of post-fermentation and packing stages on the volatile
composition of Spanish-style green table olives. Food Chem. 239, 343-353.
https://doi.org/10.1016/j.foodchem.2017.06.125.

Servili, M., Settanni, L., Veneziani, G., Esposto, S., Massitti, O., Taticchi, A., Urbani, S.,
Montedoro, G.F., Corsetti, A., 2006. The Use of *Lactobacillus pentosus* 1MO to shorten
the debittering process time of black table olives (Cv. Itrana and Leccino): a pilot-scale
application. J. Agr. Food Chem. 54, 3869-3875. https://doi.org/10.1021/jf053206y.

Suzzi, G., Gardini, F., 2003. Biogenic amines in dry fermented sausages: a review. Int. J. Food
Microbiol. 88, 41-54. https://doi.org/ 10.1016/s0168-1605(03)00080-1.

Tamang, J.P., Cotter, P.D., Endo, A., Han, N.S., Kort, R., Liu, S.Q., Mayo, B., Westerik, N.,
Hutkins, R., 2020. Fermented foods in a global age: East meets West. Compr. Rev. Food
Sci. Food Saf. 19, 184-217. https://doi.org/10.1111/1541-4337.12520

Tedersoo, L., Bahram, M., Zinger, L., Nilsson, R.H., Kennedy, P.G., Yang, T., Anslan, S.,
Mikryukov, V., 2022. Best practices in metabarcoding of fungi: From experimental design
to results. Mol. Ecol. 31, 2769-2795. https://doi.org/10.1111/mec.16460.

Tofalo, R., Schirone, M., Perpetuini, G., Angelozzi, G., Suzzi, G., Corsetti, A., 2012.
Microbiological and chemical profiles of naturally fermented table olives and brines from
different Italian cultivars. Antonie van Leeuwenhoek 102, 121-131.
https://doi.org/10.1007/s10482-012-9719-x.

Tofalo, R., Perpetuini, G., Schirone, M., Ciarrocchi, A., Fasoli, G., Suzzi, G., Corsetti, A., 2014.
Lactobacillus pentosus dominates spontaneous fermentation of Italian table olives. LWT -
Food Sci. Technol. 57, 710-717. https://doi.org/10.1016/j.lwt.2014.01.035.

Traina, C., Ferrocino, I., Bonciolini, A., Cardenia, V., Lin, X., Rantsiou, K., Cocolin, L., 2024.
Monitoring the yeasts ecology and volatiles profile throughout the spontaneous
fermentation of Taggiasca cv. table olives through culture-dependent and independent
methods. Int. J. Food Microbiol. 417, 110688.
https://doi.org/10.1016/j.ijfoodmicro.2024.110688..

Tsoungos, A., Pemaj, V., Slavko, A., Kapolos, J., Papadelli, M., Papadimitriou, K., 2023. The
rising role of omics and meta-omics in table olive research. Foods 12, 3783.
https://doi.org/10.3390/foods12203783

Tubia, I., Prasad, K., Pérez-Lorenzo, E., Abadín, C., Zumárraga, M., Oyanguren, I., Barbero, F.,
Paredes, J., Arana, S., 2018. Beverage spoilage yeast detection methods and control
technologies: A review of *Brettanomyces*. Int. J. Food Microbiol. 283, 65-76.
https://doi.org/10.1016/j.ijfoodmicro.2018.06.020..

Tufariello, M., Anglana, C., Crupi, P., Virtuosi, I., Fiume, P., Terlizzi, B.D., Moselhy, N., Attay,
H.A., Pati, S., Logrieco, A.F., Mita, G., Bleve, G., 2019. Efficacy of yeast starters to drive
and improve Picual, Manzanilla and Kalamàta table olive fermentation. J. Sci. Food Agric.
99, 2504-2512. https://doi.org/10.1002/jsfa.9460.

Tzamourani, A.P., Napoli, E.D., Paramithiotis, S., Economou-Petrovits, G., Panagiotidis, S.,
Panagou, E.Z., 2021. Microbiological and physicochemical characterisation of green table
olives of Halkidiki and Conservolea varieties processed by the Spanish method on
industrial scale. Int. J. Food Sci. Technol. 56, 3845-3857.
https://doi.org/10.1111/ijfs.15000.

Tzamourani, A., Kalogri, G., Kavvatha, M., Kochila, A., Manthou, E., Liu, Y., Panagou, E.Z.,
2024. Inoculated fermentation of cv. Conservolea natural black olives with
multifunctional starter cultures in reduced-sodium brines. Int. J. Food Sci. Technol. 59,
4093-4108. https://doi.org/10.1111/ijfs.17165.

Vaccalluzzo, A., Pino, A., Russo, N., De Angelis, M., Caggia, C., Randazzo, C.L., 2020.
FoodOmics as a new frontier to reveal microbial community and metabolic processes
occurring on table olives fermentation. Food Microbiol. 92, 103606.
https://doi.org/10.1016/j.fm.2020.103606.

Vaccalluzzo, A., Celano, G., Pino, A., Calabrese, F.M., Foti, P., Caggia, C., Randazzo, C., 2022.
Metagenetic and volatilomic approaches to elucidate the effect of *Lactiplantibacillus*

plantarum starter cultures on Sicilian table olives. Front. Microbiol. 12, 771636.
https://doi.org/10.3389/fmicb.2021.771636.

Vougiouklaki, D., Letsiou, S., Mavrokefalidou, I., Tsakali, E., Akkermans, S., Impe, J.F.M.V.,
Houhoula, D., 2024. Fermentation of Kalamata natural black olives using selected lactic
acid bacteria as starters. Fermentation 10, 53.
https://doi.org/10.3390/fermentation10010053..

Wang, J., Bakker, H.C. den, Denes, T.G., 2025. A critical review of metagenomic approaches
for foodborne pathogen surveillance. Crit. Rev. Food Sci. Nutr. 65, 1-14.
https://doi.org/10.1080/10408398.2025.2503453.

Wickham, H., 2016 ggplot2: Elegant graphics for data analysis. Springer-Verlag New York.

Zago, M., Lanza, B., Rossetti, L., Muzzalupo, I., Carminati, D., Giraffa, G., 2013. Selection of
Lactobacillus plantarum strains to use as starters in fermented table olives:
Oleuropeinase activity and phage sensitivity. Food Microbiol. 34, 81-87.
https://doi.org/10.1016/j.fm.2012.11.005.

Zotta, T., Giavalisco, M., Parente, E., Picariello, G., Siano, F., Ricciardi, A., 2022. Selection of
Lactiplantibacillus strains for the production of fermented table olives. Microorganisms
10, 625. https://doi.org/10.3390/microorganisms10030625.

1188
1189