1

# Nuclear export governs TDP-43 phase transitions and cytoplasmic aggregation

Natalie Chin [1#], Qi Zhang [2#], Jizhong Zou [3], Ken Chih-Chien Cheng [4], Wei Zheng [2], Yihong Ye [1]

[1] Laboratory of Molecular Biology, National Institute of Diabetes, Digestive, and Kidney Diseases,

National Institutes of Health, Bethesda, MD, 20892, USA

[2] Therapeutic Development Branch, National Center for Advancing Translational Sciences, National

Institutes of Health, Bethesda, MD, 20892, USA

[3] iPSC Core, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD,

20892, USA

[4] Functional Genomics Laboratory, National Center for Advancing Translational Sciences, National

Institutes of Health, Bethesda, MD, 20892, USA

Correspondence: yihongy@mail.nih.gov (Y.Y.)

# Equal contribution

**Keywords**: TDP-43, anisosome, protein aggregation, nuclear transport, XPO1, liquid-liquid phase

separation/LLPS.

**RNA-binding protein TAR DNA-binding protein 43 (TDP-43) can form liquid-like, nuclear**

**assemblies whose phase transitions are linked to protein aggregation and neurodegeneration. The**

**mechanisms that determine whether these assemblies remain dynamic or convert into solid states**

**are poorly defined. Here we combined chemical and genome-wide genetic screenings to identify**

**cellular processes influencing the phase behavior of an RNA-binding defective TDP-43 mutant**

**associated with amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD). While the**

**screens identified multiple cellular processes—including RNA splicing, translation, and**

**proteostasis—as modulators of TDP-43 phase states, our analyses underscore nuclear transport as**

**a critical determinant of liquid-to-solid transitions. Inhibition of nuclear export prevents**

**cytoplasmic accumulation of immobile TDP-43 assemblies, whereas enhanced export promotes**

**irreversible, solid TDP-43 structures. We validated this mechanism in a brain organoid model of**

**ALS, showing nuclear retention limits pathogenic aggregation and its associated defects. These**

**findings establish nuclear export as a key regulator of TDP-43 phase transitions and define a**

**mechanistic link between altered phase dynamics and neurodegeneration.**

The RNA-binding protein TAR DNA-binding protein 43 (TDP-43), encoded by the *TARDBP* gene, plays a central role in RNA transport, splicing and metabolism (Guo and Shorter, 2017; Suk and Rousseaux, 2020), and is the major pathological component of cytoplasmic inclusions in amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD) patients (Kellett et al., 2025; Neumann et al., 2006; Rummens and Da Cruz, 2025). TDP-43 contains two RNA-recognition motifs (RRMs), a bipartite nuclear-localization sequence, and a glycine-rich C-terminal low-complexity domain (LCD) that is intrinsically prone to aggregation (Tziortzouda et al., 2021). Under physiological conditions, TDP-43 binds RNAs in the nucleus, participating in multiple RNA-metabolic processes. In contrast, disease-causing mutations trigger its mis-localization to the cytoplasm, forming detergent-resistant inclusions. TDP-43-positive protein aggregates were present in ~97% of ALS and ~45% of FTD cases, as well as in a subset of Alzheimer's disease patients. These diseases are now collectively termed TDP-43 proteinopathies (Brettschneider et al., 2014; Neumann et al., 2006). Both loss of nuclear function and gain of cytoplasmic toxicity have been implicated in neurodegeneration (Suk and Rousseaux, 2020; Tziortzouda et al., 2021), underscoring the importance of nucleocytoplasmic transport in regulating TDP-43 patho-physiological functions.

Like many RNA-binding proteins (RBPs), TDP-43 undergoes liquid-liquid phase separation to form membraneless biomolecular condensates (Conicella et al., 2016; Shorter, 2019; Tziortzouda et al., 2021). Under conditions of cellular stress or in the presence of RNA-binding defective mutations, these condensates can transition from dynamic liquid droplets to more rigid gel-like or solid assemblies (Conicella et al., 2016; Gasset-Rosa et al., 2019; Lang et al., 2024). To date, more than 70 disease-linked mutations in *TARDBP* have been identified in ALS or FTD, many within the LCD, while others clustering near or in the RRMs disrupt RNA binding (Tziortzouda et al., 2021). Defects in RNA binding appears to be a major governor of TDP-43 phase behavior (Babinchak et al., 2019; Mann et al., 2019). Additionally, disease-associated mutations alter TDP-43's regulatory functions in RNA metabolism, causing widespread cryptic exon inclusion (Ling et al., 2015; Ma et al., 2022; Mehta et al., 2023; Seddighi et al., 2024; Zhang et al., 2025b), alternative polyadenylation and other RNA maturation defects that collectively contribute to neurotoxicity (Zeng et al., 2025).

A major advance in the field of TDP-43 phase regulation is the recent report of a unique form of TDP-43 demixing, forming "anisosomes" that contain an anisotropic spherical shell of an RNA-binding-defective TDP-43 mutant and a central liquid core formed by the HSP70 chaperone (Yu et al., 2021). These structures, observed mostly in the nucleus, were postulated to be an intermediate en route to pathological aggregation when cellular proteostasis or nucleocytoplasmic transport is compromised. Additionally, in vitro studies with purified TDP-43 and selected small molecules have begun to elucidate

the molecular basis of TDP-43 self-association (Maurel et al., 2025; Rubien et al., 2022), offering insights

into its phase behavior and aggregation mechanisms.

Despite these advances, the cellular pathways modulating TDP-43 phase behavior (droplet

formation, anisosomal shell–core architecture, conversion to gel/solid structures) and its link to disease-

associated aggregation remain poorly defined. Here, we address this knowledge gap by combining a

chemical-genetic screen approach with a genome-wide siRNA screen to identify molecular determinants

that modulate the phase behavior of an RNA-binding defective TDP-43 mutant. Our study reveals critical

contributions of RNA splicing, protein translation, and the HSP90/ubiquitin–proteasome proteostasis

network to TDP-43 phase transitions. Importantly, our work demonstrates how nuclear export can

influence the transition of TDP-43 from a nuclear liquid-demixed form to a cytoplasmic immobile

structure. We validated the impact of nuclear export inhibition on phosphorylated-TDP-43 cytoplasmic

accumulation using an iPSC-derived 3-D organoid model bearing an ALS-associated mutation (Zhang et

al., 2025b). Collectively, these findings establish a mechanistic framework linking altered phase dynamics

and TDP-43 aggregation to nuclear transport defects, a process known to modulate neurodegeneration in

ALS and FTD.

**Results**

**A chemical genetic screen identified modulators of TDP-43 phase behavior**

To identify cellular pathways modulating TDP-43 phase behavior, we conducted a chemical genetic

screen using a small molecule library targeting diverse known cellular pathways and processes. We

employed a previously established DLD1 cell model stably expressing an RNA-binding defective TDP-

43 mutant (2KQ) tagged with the green fluorescence protein Clover because TDP-43 2KQ undergoes

rapid demixing in the nucleus upon expression induction (Yu et al., 2021). We seeded cells in 384-well

plates and induced the expression of TDP-43 2KQ with doxycycline for 24 hours. We then treated these

cells with a LOPAC library containing 1280 drugs for 24 hours (Figure 1A). This approach could avoid

false positive drugs that affect the induction of TDP-43 2KQ.

High content imaging combined with algorithm-based image analyses (see methods) detected ~20

spherical anisosomes per cell in controls and most treated cells. However, in cells treated with a subset of

the drugs, the number of anisosomes was noticeably reduced (Supplemental Table 1). Importantly, cells

treated with Bortezomib, a potent proteasome inhibitor, had fewer but enlarged TDP-43 positive punctae

with irregular shapes (Figure 1B), similar to cells treated with another proteasome inhibitor reported

previously (van Eersel et al., 2011; Yu et al., 2021). This result validated our screening design.

To further confirm the identified hits, we performed a titration experiment, which showed that

these inhibitors dose-dependently reduced the number of anisosomes with $IC_{50}$ ranging from sub-

micromolar to micromolar (Figure 1B, Supplemental Table 1). Interestingly, the screen identified

cytoplasmic signaling kinases including PAK4, BTK, and GSK-3β as well as the receptor tyrosine kinase

PDGFRβ as modulators of TDP-43 phase behavior (Figure 1B, Supplemental Table 1). PAK4 and GSK-

3β are known regulators of the YAP signaling pathway (Jin et al., 2025; Mo et al., 2024), and YAP was

recently identified as a TDP-43-interacting protein that influences its phase dynamics (Zhang et al.,

2025a). These connections provide additional physiological support for the relevance of the modulators

uncovered in our screen. Our screen also linked TDP-43 2KQ demixing to protein translation because

cycloheximide, a ribosome elongation inhibitor could modulate TDP-43 phase dynamics (see below).

**Live cell imaging revealed two distinct types of TDP-43 phase modulators**

We next used confocal microscopy to further characterize the impact of a subset of the identified

inhibitors on anisosome dynamics. We chose Tripterin (a drug targeting the heat shock protein HSP90),

CP-673451 (a PDGFRβ inhibitor), two deubiquitinase inhibitors VLX-1570 and Spautin-1, and the

Exportin-1 inhibitor KPT-276 because previous reports have linked the targets of these compounds to

TDP-43 proteinopathies (Archbold et al., 2018; Farrawell et al., 2020; Furukawa et al., 2015; Garcia-

Toscano et al., 2024). We first induced anisosome formation for 24 hours and then treated cells with the

inhibitors. Compared to DMSO-treated controls, cells treated with Spautin-1 had fewer and smaller

anisosomes. KPT-276 treatment also reduced anisosome number, albeit to a lesser extent (Figure 2A, top

panels). In contrast, cells treated with CP-673451, Tripterin, or Bortezomib had fewer but larger TDP-43

punctae. Since Tripterin was also reported as a 20S proteasome inhibitor, we confirmed the role of HSP90

in this process by treating cells with the well-established HSP90 inhibitor Geldanamycin, which caused

similar phenotypes as Tripterin (Supplemental Figure 1A). These results suggest that TDP-43 phase

dynamics are modulated by the cellular folding capacity, the ubiquitin proteasome system, and

nucleocytoplasmic transport.

128            To further dissect the effects of these inhibitors on TDP-43 phase separation, we combined drug

treatment with live cell fluorescence imaging, using Fluorescence Recovery After Photobleaching (FRAP)

or reversed FRAP to define the dynamics of TDP-43 2KQ. In FRAP, when the green fluorescence of

anisosomes was bleached by a laser in untreated cells, it rapidly recovered due to the recruitment of TDP-

43 2KQ from the nucleoplasm (Figure 2B, top panels, Figure 2C, D). Conversely, in reversed FRAP,

bleaching TDP-43 2KQ surrounding an anisosome and then imaging the anisosome over time revealed a

gradual decline in fluorescence intensity within the unbleached anisosome, while the surrounding

bleached anisosomes regained some fluorescence (Figure 2E, top panel). This result suggests that TDP-43

2KQ in anisosomes could also be released into the nucleoplasm. Interestingly, in cells treated with

Spautin-1 or KPT-276, anisosome-associated TDP-43 2KQ could freely exchange with the nucleoplasmic

pool (Figure 2B, D). However, in cells treated with Bortezomib, Tripterin or Geldanamycin, the enlarged

TDP-43 puncta appeared static; TDP-43 2KQ fluorescence within anisosomes failed to recover after

photobleaching (Figure 2B, C, Supplemental Figure 1B), nor did it decrease over time when surrounding

TDP-43 2KQ was bleached (Figure 2E-G). Collectively, these results suggest that these inhibitors affect

anisosomes via two distinct mechanisms: one maintains TDP-43 in a demixed liquid state, while the other

converts it to a gel-like, solid state.

**A genome-wide siRNA screen identified genetic modulators of TDP-43 phase separation**

To further decipher the molecular determinants of TDP-43 phase separation, we conducted an unbiased

genome-wide siRNA knockdown (KD) screen (Figure 3A). To this end, we transfected a siRNA library

targeting 21,404 human genes each with three siRNAs into DLD1 cells stably expressing Clover-tagged

TDP-43 2KQ. After gene KD, anisosomes were induced for 24 hours. High content imaging and

automated analysis identified 1,533 candidates whose KD reduced the anisosome number per cell (Z-

score > 2) (Supplementary Table S2). Based on a STRING gene network analysis, we selected 211

networked genes with top Z-scores and re-screened each of them with three additional siRNAs. This

confirmed the involvement of 110 genes in TDP-43 anisosome regulation (Figure 3A, Supplementary

Table S3). GO pathway analyses categorized identified genes into several major pathways including RNA

splicing, protein translation, proteasomal degradation, and nuclear transport (Figure 3B). The

identification of large number of ribosomal proteins and proteasome subunits is consistent with our

chemical genetic screen, which showed that translation and proteasome inhibitors modulate anisosomes

(Supplemental Table 1). Interestingly, GO analyses using 'molecular function' further linked many

identified genes to neurodegenerative diseases, particularly ALS (Figure 3C, D).

**Anisosome dynamics is modulated by RNA splicing and protein translation**

Given the well-established function of TDP-43 in RNA binding and processing, we investigated the role

of RNA splicing in anisosome regulation. To this end, we first incubated Clover-tagged TDP-43 2KQ

cells with doxycycline to induce anisosomes and then treated cells with the potent RNA splicing inhibitor

Pladienolide-B (PlaB). Confocal microscopy confirmed that RNA splicing inhibition resulted in fewer but

larger TDP-43-positive puncta in a dose dependent manner (Figure 4A, B, Supplemental Figure 2A).

Despite the size difference, anisosomes in drug-treated cells were morphologically indistinguishable from

controls. FRAP experiments further showed that TDP-43 within PlaB-treated condensates remained

highly mobile, although the fluorescence recovery rate was slightly reduced compared to controls (Figure

4C, D). Time-lapse live cell imaging frequently detected fusion of TDP-43 puncta after PlaB treatment

(Figure 4E, Movie S1). Together, these results suggest that disrupting RNA splicing stabilizes TDP-43 in

demixed liquid state, resulting in larger condensates.

Next, we explored the effect of translation inhibition on anisosome dynamics using

Cycloheximide (CHX) and Anisomycin (ANS). Confocal imaging showed that after anisosome induction,

translational shutdown by CHX or ANS did not significantly reduce TDP-43 expression, suggesting a

long half-life. Similar to PlaB-treated cells, CHX- and ANS-treated cells contained fewer, enlarged

anisosomes (Figure 4F-H, Supplemental Figure S2B). FRAP analysis revealed that TDP-43 dynamics

within anisosomes were similar between ANS-treated and control cells (Figure 4I, J). These results

suggest that translation inhibition also stabilizes TDP-43 in liquid condensates.

**Nuclear export regulates TDP-43 liquid-to-solid phase transition**

Our screen also identified several nuclear transport regulators (e.g., Exportin-1/XPO1, Exportin-2/CSE1L)

and nuclear pore components as anisosome regulators (Figure 3B). We focused on XPO1 because our

chemical genetic screen identified two XPO1 inhibitors, KPT-276 and Verdinexor as potent anisosome

modulators (Figure 1B).

To explore the role of XPO1 in TDP-43 anisosome regulation, we induced anisosome formation

in Clover-TDP-43 2KQ cells and treated them with Leptomycin B (LMB), a potent XPO1 inhibitor. LMB

treatment resulted in a time- and dose-dependent reduction in anisosome number, while increasing their

sizes (Figure 5A-C). The enlarged TDP-43 puncta showed the typical hollowed anisosome ring structure

(Figure 5D), indicating that TDP-43 remained in the liquid state. Indeed, FRAP experiments

demonstrated that LMB treatment did not significantly affect the fluorescence recovery rate of TDP-43

within bleached anisosomes (Figure 5D, E). Time-lapse confocal imaging showed that shortly after LMB

treatment, preformed anisosomes began to fuse with each other (Figure 5F, Movie S2). Thus, XPO1

inhibition stabilizes TDP-43 in a liquid phase.

To corroborate the inhibitor study, we overexpressed mCherry-tagged XPO1. Surprisingly, we

observed fewer and enlarged TDP-43 positive puncta in mCherry-XPO1-expressing cells (Figure 5G,

panels 1, 2). At first glance, this phenotype resembled that of XPO1-inhibited cells. However, careful

examination revealed several distinct features. First, despite the size increase, we did not observe the

typical hollowed anisosome ring structure. Secondly, while anisosomes in mCherry-XPO1 negative cells

were almost exclusively nuclear, ~30% of mCherry-XPO1-expressing cells contained cytoplasm-

localized TDP-43 puncta (Figure 5G, panels 1, 2, 3, Figure 5H). Thirdly, FRAP experiments

demonstrated that Clover-TDP-43 fluorescence did not recover after photobleaching, suggesting that

TDP-43 was in a gel-like state transitioning to aggregates (Figure 5I). Notably, mCherry-XPO1 could be

detected within TDP-43 puncta in some cells (Figure 5G, panels 3-6). These findings are consistent with a

previous report (Archbold et al., 2018), demonstrating that XPO1 promotes TDP-43 nuclear egression

and cytoplasmic aggregate formation. Immunostaining of endogenous XPO1 in anisosome-induced cells

further confirmed this conclusion. Although antibodies could not penetrate anisosomes (Yu et al., 2021),

anisosome induction depleted XPO1 from the nucleoplasm, presumably due to its sequestration into

anisosomes (Figure 5J, K). Since XPO1 does not bind TDP-43 directly (Pinarbasi et al., 2018), additional

factors likely facilitate XPO1-mediated TDP-43 nuclear egression under this condition. Collectively,

these results suggest that the stability and dynamics of anisosomes are modulated by XPO1: reduced

XPO1 activity stabilizes the liquid state and favors large TDP-43 condensates, whereas increasing XPO1

activity promotes its transition to a gel-like structure and cytoplasmic redistribution (Figure 5L). This

conclusion underscores a critical regulatory link between XPO1-dependent nuclear egression and the

TDP-43 liquid-to-solid phase transition.

**XPO1 inhibition mitigates TDP-43 hyperphosphorylation and modest rescues neuronal defects in**

**an anisosome-bearing organoid model**

Since increased XPO1 activity is associated with more cytoplasmic, gel-like TDP-43, we tested whether

reducing XPO1 activity could mitigate TDP-43 cytosolic aggregation using a recently established

anisosome-bearing organoid model of TDP-43 proteinopathies. This model is derived from an induced

pluripotent stem cell (iPSC) line carrying the ALS-associated mutation (K181E) (Zhang et al., 2025b),

which like 2KQ disrupts the RNA binding activity of TDP-43 (Chen et al., 2019). Additionally, the

K181E mutant accumulates in a liquid-condensed phase in association with HSP70, and like anisosomes,

these condensates can convert to solid aggregates during cellular stress or HSP70 depletion (Gu et al.,

2021).

We treated homozygous K181E organoids or wild-type (WT) controls with KPT-276 at a low

dose (20 nM) for 35 days, starting on day 87 based on pilot data indicating good tolerance with prolonged

treatment. Since hyperphosphorylated TDP-43 (p-TDP-43) is a major aggregate component, we stained

organoids with antibodies against p-TDP-43 and total TDP-43. Confocal imaging revealed that WT and

K181E organoids had similar amounts of total TDP-43, but p-TDP-43 was only significantly present in

mutant organoids (Figure 6A), as reported previously (Zhang et al., 2025b).  KPT-276 did not alter the

total TDP-43 levels in either genotype (Figure 6A) but significantly reduced the cytosolic p-TDP-43

positive puncta in K181E organoids (Figure 6A, B).

To assess whether KPT-276 treatment also rescued the neuronal defects in TDP-43 K181E

organoids, we performed bulk RNAseq using KPT-276-treated K181E and WT organoids. A comparison

of treated mutant organoids to WT or untreated K181E organoids showed that KPT-276 reversed

expression defects in 58 downregulated (2.90%) and 45 (1.99 %) upregulated genes (Supplemental Figure

3, Supplemental Table S4). GO pathway analysis of genes rescued from downregulation by KPT-276

highlighted neuronal pathways including neuron differentiation and brain development; for genes

upregulated in K181E organoids, rescued genes are linked to G-protein coupled neurotransmitter receptor

activities at the plasma membranes (Figure 6C, Supplementary Table S4). Together, these results

demonstrate that XPO1 inhibition by KPT-276 modestly restores gene transcription defects in a subset of

neuronal pathways in K181E mutant organoids, while robustly mitigating FTD-associated p-TDP-43

accumulation. Thus, reducing TDP-43 cytoplasmic aggregation alone may offers only limited benefit in

disease treatment.

**Discussion**

ALS- and FTD-associated TDP-43 mutants defective in RNA binding are known to undergo self-

association through their intrinsically disordered regions and RNA-binding domains, generating demixed

liquid, gel, and aggregated states that remain in dynamic exchange. This protein phase behavior is shared

by many RNA binding proteins and is increasingly recognized as a contributor to protein aggregation in

neurodegenerative diseases (Shorter, 2019). Intriguingly, TDP-43 phase separation forms unique layered

nuclear structures, with an HSP70-filled central core surrounded by a protein shell made of TDP-43.

Although it is unclear how these structures are nucleated in the cell, our observations suggest that, once

formed, they can grow by recruiting additional TDP-43 from the surrounding environment. Small

anisosomes may also fuse with one another, although such fusion events become infrequent and difficult

to detect after anisosomes reach a steady state,

259           Our study supports the notion that TDP-43, at high concentrations, can transition from a

260   demixing-prone, liquid state into a gel-like state en route to forming amorphous aggregates (Babinchak et

261   al., 2019; Carey and Guo, 2022). We found that inhibition of RNA splicing, protein translation, or nuclear

262   export favors the liquid state of TDP-43, while proteasome and HSP90 inhibition or enhancing XPO-1-

263   mediated nuclear egression promotes the conversion of anisosomes to a gel-like state.

264           Because the TDP-43 2KQ mutant lacks RNA binding activity, the effects of splicing inhibition

are likely mediated by other RNA-binding factors involved in splicing regulation. TDP-43 is known to

interact with many splicing factors including members of the hnRNP family like hnRNP A1/A2 and PSF

(Freibaum et al., 2010); Inhibition of RNA splicing may alter TDP-43's interactions with these factors,

trapping it in certain complexes that favor its retention in the liquid state. Alternatively, splicing inhibition

may change the overall nuclear environment, allowing large anisosomes to fuse with each other as

observed in our study. This could explain why PlaB treatment reduces the number of TDP-43 anisosomes

while maintaining their dynamic liquid property. Likewise, inhibition of nuclear export may influence

anisosome dynamics by reshaping the nuclear environment, lowing the barrier to anisosome fusion.

Future studies are required to elucidate the precise mechanisms by which these pathways regulate
anisosome dynamics.

Our validation of proteasome inhibition as a driver of TDP-43 droplet enlargement and hardening
reinforces the idea that impaired protein clearance promotes the transition from liquid condensates to gel-
like or solid aggregates. This is consistent with previous studies in ALS models, which showed that
proteasome stress accelerates TDP-43 aggregation and toxicity (van Eersel et al., 2011). In contrast,
inhibition of deubiquitinases reduced the number of TDP-43 anisosomes, probably by promoting the
proteasome-mediated clearance of TDP-43.  The role of HSP90 in this process may be linked to TDP-43
folding stability, as HSP90 can bind TDP-43 to prevent promiscuous interactions (Lin et al., 2021).
Collectively, these observations suggest that TDP-43 condensates are maintained by a delicate balance
between formation and clearance, modulated by proteolytic and chaperone activities.

While out chemical and genetic screens have identified multiple pathways influencing TDP-43
phase states, a particularly compelling aspect of our study is the discovery that the nuclear export receptor
XPO1 governs TDP-43 liquid-to-solid transitions and subcellular localization. Pharmacological inhibition
of XPO1 mitigates cytoplasmic p-TDP-43 accumulation, whereas XPO1 overexpression promotes the
cytoplasmic accumulation of TDP-43-containing puncta. However, XPO1 may modulate TDP-43 nuclear
egression indirectly, as suggested previously (Duan et al., 2022; Pinarbasi et al., 2018). While previous
studies have not detected strong interactions between XPO-1 and TDP-43 (Pinarbasi et al., 2018), a weak
or indirect interaction between XPO-1 and TDP-43 may still exist, especially when TDP-43 undergoes
demixing. This model would explain why anisosome induction depletes endogenous XPO-1 from the
nucleoplasm. Our findings also hint at a potential feedback mechanism in which TDP-43 demixing
perturbs nuclear export, analogous to how cytoplasmic TDP-43 aggregates disrupt nuclear import. Given
that ALS is characterized by cytoplasmic TDP-43 mis-localization and hyperphosphorylation, the XPO-1-
dependent nuclear export pathway may constitute a key molecular switch linking physiological phase
behavior to pathological aggregation.

Notably, in 3-D brain organoids bearing the K181E mutation, inhibition of nuclear export by
KPT-276 only modestly rescues the transcriptional defects in homozygous TDP-43 K181E organoids
despite an almost complete ablation of cytoplasmic phospho-TDP-43. This observation suggests that
cytoplasmic TDP-43 aggregation is only a modest contributor to TDP-43 proteinopathies that lead to
neurodegeneration. Major defects may still be attributed to deregulated RNA processing due to altered
RNA binding (Hergesheimer et al., 2019; Joseph et al., 2025; Mehta et al., 2023; Zhang et al., 2025b).
Thus, effective therapy likely requires both inhibition of cytoplasmic TDP-43 aggregation and restoration
of normal RNA processing and nuclear transport. Together, our results support a model in which multiple
cellular pathways including proteostasis, RNA metabolism, translation, and nuclear export—

cooperatively determine the phase behavior of TDP-43. These findings have deepened our understanding

of how physiological phase separation can evolve into pathogenic aggregation through cumulative

failures of cellular quality control and nuclear export systems.

**Methods**

**RNAi-based genetic screen for anisosome regulators**

Genome-wide RNA interference (RNAi) screen was performed using DLD1 cells stably expressing

Clover-tagged TDP-43 2KQ. Cells were transfected with siRNAs targeting 21,404 genes for 72 hours,

followed by induction of anisosome formation with doxycycline (0.5 μg/mL) for 24 h. Specifically, the

screen was carried out in 384-well format with 3 individual siRNAs for each gene. To prepare siRNA

transfection, control siRNA (40 nmol) was mixed with 2 mL water to create a 50X (20 μM) stock and

siRNA-death (20 nmol) was mixed with 1 mL of water for the same concentration. Dilute siRNA stocks

to a 400 nM working concentration in water. For each well, mix 0.1 μL transfection reagent in 20 μL of

cell culture medium without FBS or P/S. Add 2 μL of siRNA solution to each well and incubate with

RNAiMAX for 30 minutes at room temperature. While incubating RNAiMAX, prepare trypsinized cells

by ensuring thorough separation and accurate cell counting. Add 20 μL of cell suspension at a density of

$0.43 \times 10^5$ cells/mL (850 cells/well) in medium containing 20% FBS and 2 x P/S to achieve a final

volume of 40 μL per well. Incubate cells for 3 days before addition of 10 μL 10 x doxycycline solution to

each well (final volume 50 μL) for anisosome induction. 24 h later, high-throughput (HT) imaging and

automated phenotypic screening were conducted to identify genes that modulated anisosome formation.

Fluorescence imaging was conducted using an Opera Phenix High-Content Screening System

(PerkinElmer). Whole-well images were acquired in confocal mode using a 20X water objective lens after

anisosome induction for 24 h. Images were reconstructed and analyzed using the PerkinElmer Columbus

server (v2.9.1). Nuclei were segmented based on Hoechst33342 staining, and anisosomes were detected

via YFP signals within the nuclear region. The mean anisosome count per nucleus was calculated to

assess changes in anisosome levels, while toxicity was evaluated by counting the total number of nuclei.

Data were normalized to the siRNA-Neg control for each plate, and Z-scores were calculated to identify

significant hits. Identified 1,533 genes whose knockdown significantly reduced the number of anisosomes

per cell (Z-score > 2) were subjected to protein network analysis by STRING to identify candidate

anisosome regulatory genes that were subject to a second-round screen.

**A chemical genetic screen for anisosome regulators**

We also performed a chemical genetic screen using LOPAC[R1280] (Sigma) compound library on DLD1

cells expressing Clover-tagged TDP-43-2KQ. Briefly, cells were induced with doxycycline for 24 h to

promote anisosome formation, followed by treatment with individual compounds for an additional 24

hours. High-throughput (HT) imaging and automated phenotypic screening were conducted to identify

compounds that modulated anisosome formation.

**Visualization and analysis of anisosome dynamics**

To examine the impact of drug treatment on anisosome size, cells were induced to form anisosome for 24

h and then treated by different inhibitors as follows: cycloheximide (CHX, 20 µg/mL), anisomycin (ANS,

200 nM) for 16 hours. For splicing inhibition, cells were exposed to Pladienolide-B (PlaB) at

concentrations of 5 or 20 nM, or DMSO (control) for 16 hours before imaging. Cells were fixed with 4%

paraformaldehyde in Phosphate Buffer Saline (PBS). Randomly selected fields were used to quantify the

number of anisosomes per cell with at least two biological repeats.

To image anisosome fusion, live cell imaging was performed by treating cells with Leptomycin B

(200 nM) or PlaB (20 nM) for 5 hours and then imaged by confocal 3-D sectioning for 30 minutes to

capture anisosome fusion events. For Fluorescence Recovery After Photobleaching (FRAP), cells were

treated with DMSO, Leptomycin B, Geldanamycin or PlaB as indicated in figure legends. Either part of

an anisosome or entire anisosome was photobleached and imaged. TDP-43 fluorescence recovery after

photobleaching was quantified by ImageJ and calculated to assess anisosome dynamics. At least 10

anisosomes for each condition were bleached and analyzed.

To evaluate the time course of Leptomycin B (LMB)-induced anisosome changes, DLD1 TDP-43

2KQ-Clover cells were seeded and induced with 0.5 µg/mL doxycycline for 24 hours. Cells were treated

with 10 µM LMB and fixed at 0-, 8-, and 16-hours post-treatment. Fixed cells were stained with Hoechst

to visualize nuclei. 10-20 confocal 3-D sections were obtained for each randomly selected field. Images

were converted to maximum projected view by ImageJ before anisosome counting and intensity

measurement. The density plot of anisosome counts per cell was generated using R, illustrating dose-

dependent effects of leptomycin on anisosome formation.

To test the effect of XPO1 overexpression on TDP-43 phase regulation, DLD1 TDP-43 2KQ-

Clover cells were co-transfected with mCherry-tagged XPO1 (mCh-XPO1) for 24 h before doxycycline

was added at 0.5 µg/mL to induce anisosome formation for 24 h before confocal imaging.

To study the relationship between anisosome induction and endogenous XPO1 localization, cells

were seeded, induced with 0.5 µM doxycycline for 48 hours, and fixed with paraformaldehyde in PBS.

Immunostaining was performed using an XPO1 antibody (Cell Signaling, 46249S, 1:125) and Hoechst

nuclear stain. Fluorescence intensity of XPO1 in both induced and uninduced cells was measured by

ImageJ.

**Experiments with organoids**

Stem cell gene editing of the patient K181E mutation in iPSC cells was reported previously (Zhang et al., 2025b). Differentiation of iPSCs into forebrain neurons was performed following a previously published protocol (Fernandopulle et al., 2018). Briefly, the procedure used an induction medium (IM-N2) and a neuronal culture medium (CM). IM-N2 was prepared with Knockout DMEM/F12, N2 supplement, NEAA, Gluta-MAX, Chroman I, and doxycycline. On Day 0, iPSCs were observed for confluency, dissociated with Accutase, centrifuged, and resuspended in IM-N2 with Chroman I, and then seeded into Matrigel-coated plates. Over the next four days, cells were monitored microscopically for neurite extensions while media containing doxycycline is refreshed daily. By Day 4, cells exhibit neurite growth and are ready for replating onto poly-L-ornithine (PLO)-coated dishes, prepared in advance by coating with PLO solution, incubating, washing, and drying. Replated cells were cultured in CM comprising BrainPhys medium, B27+ supplement, neurotrophic factors (GDNF, BDNF, NT-3), laminin, and doxycycline, to promote neuronal maturation.

3-D culture and organoid growth were performed as previously described (Nguyen, 2022). In brief, iPSCs grown on Matrigel were dissociated into single cell suspension by Versene solution (ThermoFisher) and seeded into a 12-well Aggrewell plate (Stemcell Technologies) at 4,000 cells/ well. Next day, spheroids were transferred to an ultralow attachment plate (Corning) containing phase I medium: DMEM/F12, 20% Gibco KnockOut Serum Replacement, 1X Glutamax, 1X MEM Nonessential Amino Acid, 55 μM β-Mercaptoethanol, 1X Pen/Strep, 2 μM Dorsomorphine, 2 μM A83-01. After 5 days, medium was switched to phase II medium: DMEM/F12, 1X N-2 Supplement, 1X Glutamax, 1X MEM Nonessential Amino Acid solution, 1X Pen/Strep, 4 ng/mL WNT3a, 1 μM CHIR-99021, 1 μM SB-431542. On day 7, spheroids were embedded into Matrigel and allowed to continue growing for 7 more days. On day 14, individual spheroids were manually freed from the Matrigel and transferred to a SpinOmega bioreactor spinning at 120 RMP with phase III media: DMEM/F12, 1X N-2 Supplement, 1X B-27 Supplement, 1X Glutamax, 1X MEM Nonessential Amino Acid, 55 μM 2-Mercaptoethanol, 1X Pen/Strep, 2.5 μg/mL insulin. 50 days later, the medium was switched to final differentiation medium: Neurobasal medium, 1X B-27 Supplement, 1X Glutamax, 55 μM 2-Mercaptoethanol, 1X Pen/Strep, 0.2 mM Ascorbic acid, 0.5 mM cAMP, 20 ng/mL brain-derived neurotrophic factor (BDNF), 20 ng/mL glial-derived neurotrophic factor (GDNF). 107-day-old organoids were dissociated into single cells using a 50:50 mixed of Accutase and 0.25% Trypsin with DNase I (1 mg/mL) and plated onto chambered glass slides pretreated with 1% Matrigel (Corning, Inc). Alternatively, organoids were fixed and sectioned for immunostaining (see below). For drug treatment, forebrain organoids (87 Day) were transferred to a 12-well plate on an orbital shaker (120 RPM) and treated with 20 nM KPT276 in final differentiation

medium. Medium was changed every other day (Dexoregen, Inc). The organoids were harvested and

frozen after 35 days of treatment for sectioning or RNAseq analysis.

**Tissue preparation and immunostaining**

Organoids were fixed in 4% paraformaldehyde in PBS for 1 hour at room temperature, washed with PBS,

and immersed in 15% sucrose/PBS solution overnight. Subsequently, organoids were embedded in O.C.T.

compound (Sakura) in a plastic mold and frozen down in an ultralow freezer. Embedded organoids were

sliced onto glass slides using a Cryostat (Leica). Slides were rinsed with PBS, permeabilized with 0.5%

Triton-X/PBS solution for 1 hour at room temperature and blocked using 1% donkey serum in 0.1%

Tween-20/PBS solution for 1 hour. Primary antibodies, chicken anti-TUJ1 (Abcam, Ab41489,1:1,000

dilution), rabbit anti-TDP-43 (10782-2-AP, 1/1000 dilution), mouse anti-phospho-TDP-43 (Cosmobio,

CAC-TIP-PTD-M01A, 1/500 dilution), were added to the slides and the slides were incubated in a

humidified chamber at 4 °C overnight. After several washes in 0.1% Tween-20/PBS solution, DAPI

(Sigma) or secondary antibodies, donkey anti-rabbit, anti-mouse, and anti-chicken (Jackson

ImmunoResearch), diluted 1:1,000 in blocking solution, were added to the slides. After 1 hour of

incubation at room temperature, the slides were washed and mounted in antifade mounting solution

(Fisher scientific).

**Image acquisition and processing**

Fluorescence confocal images were acquired using a Nikon CSU-W1 SoRa microscope equipped with

temperature and $CO_2$ control enclosures and a 60x TIRF lens. Image reconstructions and analyses,

including 3-D and time lapse visualizations, were performed using Imaris software or Image J.

Fluorescence intensity was quantified using the open-source Fiji/Image J software. Images of randomly

selected fields were split into individual channels. A consistent threshold method was applied to

individual channel. The particle analysis function was used to automatically identify all anisosomes or

other fluorescence structures for size and intensity measurement. Statistical analyses were conducted

using Excel (for Student's t-test) or GraphPad Prism versions 8.0 and 9.0. P-values were calculated using

Student's t-test in Excel or one-way ANOVA in GraphPad Prism. Curve fitting, including linear and

nonlinear models, as well as $IC_{50}$ calculations, were also performed using GraphPad Prism versions 8.0

and 9.0.

**Bulk RNA-seq library preparation and data analysis**

800 ng of total RNA samples to generate the library with the NEBNext rRNA Depletion Kit v2 (Human/

mouse/rat) (NEB #E7405), NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (NEB

#E7760) and NEBNext® Multiplex Oligos for Illumina® (96 Unique Dual Index Primer Pairs, NEB,

E6440) as described in the manufacturer's manual. Sequencing carried out in the Illumina NextSeq2000

instrument with 101x101 pair end configuration by NCI genomic core.

Adapters were trimmed using Cutadapt (v4.7), and reads were aligned to the GRCh38 genome

build with STAR (v2.7.11b) using gene annotations from GENCODE v42. Gene expression was

quantified with HTSeq (v2.0.7) (Putri et al., 2022), utilizing the same gene annotations. For differential

gene expression analysis, all samples were processed uniformly following the standard DESeq2 (Love et

al., 2014) workflow. Differential expression was defined based on an adjusted P-value threshold of <0.05

and |log2 fold change ($log_2FC$) > 1. To identify gene expression rescued by exportin-1 inhibitor treatment,

the fold change of expression level should be less than without inhibitor treatment, with P-value < 0.05.

**Statistic and reproducibility**

All statistical analyses were conducted with GraphPad Prism v10 or Excel. Statistical methods and the

number of cells or brain organoid samples (N) are indicated in figure legends or shown in figures as

individual data point. Biological repeats (n) are specified in figure legends. No biological repeat was

excluded from the analyses. All experiments were repeated at least twice with individual data point

labeled in figures unless specified. For imaging analyses, cells in randomly selected fields were analyzed.

The researchers were not blinded. The iPSC cell line was derived from a ADRDs genetic risk-free clone,

which was initially obtained from a male. Figures were prepared using ImageJ 1.54f, Imaris 9.9.0, Adobe

Photoshop v25.12.1, and Adobe Illustrator 28.7.4.

**Data and code availability**

Raw and processed bulk RNA-seq data can be accessed from Sequence Read Archive (SRA), NCBI with

Bioproject ID: PRJNA1195353, http://www.ncbi.nlm.nih.gov/bioproject/1195353.

**Author contributions**

YY conceived and designed the study. YY, KC and WZ supervised the study. QZ and KC conducted the

chemical and siRNA screens. NC performed post-screen validation and characterization of anisosomes

under different conditions. J Z constructed the iPSC knock-in lines and QZ conducted the organoid study.

QZ, NC and YY analyzed the data. YY wrote the manuscript. All authors read, edited, and approved the

manuscript.

**Corresponding authors**

Correspondence to Yihong Ye.

**Funding**

This work is supported by the Intramural Research Programs of NCATS (WZ & KC) and NIDDK (YY)

at NIH.

**Declaration of Interests**

The authors declare no competing interests.

**Acknowledgements**

We thank the NIDDK Advanced Imaging Core with data acquisition. We thank D.W. Cleveland (UCSD)

and H. Yu (Genentech) for providing the doxycycline-inducible DLD1 TDP-43 2KQ-Clover cell line and

protocol. We thank K. Bharti and W. Li (NEI), Dr. T. Zhang (NIA) for critical reading of the manuscript.

We also thank Drs. L. El Touny, D. Blivis, and T. Voss (NCATS) for their valuable support, as well as

the NCATS Compound Management Group and the NCATS Automation Group for their contributions to

the chemical and genetic screens.

**References**

Archbold, H.C., K.L. Jackson, A. Arora, K. Weskamp, E.M. Tank, X. Li, R. Miguez, R.D. Dayton, S. Tamir, R.L. Klein, and S.J. Barmada. 2018. TDP43 nuclear export and neurodegeneration in models of amyotrophic lateral sclerosis and frontotemporal dementia. *Sci Rep*. 8:4606.

Babinchak, W.M., R. Haider, B.K. Dumm, P. Sarkar, K. Surewicz, J.K. Choi, and W.K. Surewicz. 2019. The role of liquid-liquid phase separation in aggregation of the TDP-43 low-complexity domain. *J Biol Chem*. 294:6306-6317.

Brettschneider, J., K. Del Tredici, D.J. Irwin, M. Grossman, J.L. Robinson, J.B. Toledo, L. Fang, V.M. Van Deerlin, A.C. Ludolph, V.M. Lee, H. Braak, and J.Q. Trojanowski. 2014. Sequential distribution of pTDP-43 pathology in behavioral variant frontotemporal dementia (bvFTD). *Acta Neuropathol*. 127:423-439.

Carey, J.L., and L. Guo. 2022. Liquid-Liquid Phase Separation of TDP-43 and FUS in Physiology and Pathology of Neurodegenerative Diseases. *Front Mol Biosci*. 9:826719.

Chen, H.J., S.D. Topp, H.S. Hui, E. Zacco, M. Katarya, C. McLoughlin, A. King, B.N. Smith, C. Troakes, A. Pastore, and C.E. Shaw. 2019. RRM adjacent TARDBP mutations disrupt RNA binding and enhance TDP-43 proteinopathy. *Brain*. 142:3753-3770.

Conicella, A.E., G.H. Zerze, J. Mittal, and N.L. Fawzi. 2016. ALS Mutations Disrupt Phase Separation Mediated by alpha-Helical Structure in the TDP-43 Low-Complexity C-Terminal Domain. *Structure*. 24:1537-1549.

Duan, L., B.L. Zaepfel, V. Aksenova, M. Dasso, J.D. Rothstein, P. Kalab, and L.R. Hayes. 2022. Nuclear RNA binding regulates TDP-43 nuclear localization and passive nuclear export. *Cell Rep*. 40:111106.

Farrawell, N.E., L. McAlary, J.S. Lum, C.G. Chisholm, S.T. Warraich, I.P. Blair, K.L. Vine, D.N. Saunders, and J.J. Yerbury. 2020. Ubiquitin Homeostasis Is Disrupted in TDP-43 and FUS Cell Models of ALS. *iScience*. 23:101700.

Fernandopulle, M.S., R. Prestil, C. Grunseich, C. Wang, L. Gan, and M.E. Ward. 2018.
Transcription Factor-Mediated Differentiation of Human iPSCs into Neurons. *Curr Protoc*
*Cell Biol*. 79:e51.

Freibaum, B.D., R.K. Chitta, A.A. High, and J.P. Taylor. 2010. Global analysis of TDP-43
interacting proteins reveals strong association with RNA splicing and translation
machinery. *J Proteome Res*. 9:1104-1120.

Furukawa, T., N. Matsui, K. Fujita, H. Nodera, F. Shimizu, K. Miyamoto, Y. Takahashi, T. Kanda,
S. Kusunoki, Y. Izumi, and R. Kaji. 2015. CSF cytokine profile distinguishes multifocal
motor neuropathy from progressive muscular atrophy. *Neurol Neuroimmunol*
*Neuroinflamm*. 2:e138.

Garcia-Toscano, L., H.N. Currey, J.C. Hincks, J.G. Stair, N.J. Lehrbach, and N.F. Liachko. 2024.
Decreased Hsp90 activity protects against TDP-43 neurotoxicity in a C. elegans model
of amyotrophic lateral sclerosis. *PLoS Genet*. 20:e1011518.

Gasset-Rosa, F., S. Lu, H. Yu, C. Chen, Z. Melamed, L. Guo, J. Shorter, S. Da Cruz, and D.W.
Cleveland. 2019. Cytoplasmic TDP-43 De-mixing Independent of Stress Granules Drives
Inhibition of Nuclear Import, Loss of Nuclear TDP-43, and Cell Death. *Neuron*. 102:339-
357 e337.

Gu, J., C. Wang, R. Hu, Y. Li, S. Zhang, Y. Sun, Q. Wang, D. Li, Y. Fang, and C. Liu. 2021.
Hsp70 chaperones TDP-43 in dynamic, liquid-like phase and prevents it from amyloid
aggregation. *Cell Res*. 31:1024-1027.

Guo, L., and J. Shorter. 2017. Biology and Pathobiology of TDP-43 and Emergent Therapeutic
Strategies. *Cold Spring Harb Perspect Med*. 7.

Hergesheimer, R.C., A.A. Chami, D.R. de Assis, P. Vourc'h, C.R. Andres, P. Corcia, D.
Lanznaster, and H. Blasco. 2019. The debated toxic role of aggregated TDP-43 in
amyotrophic lateral sclerosis: a resolution in sight? *Brain*. 142:1176-1194.

Jin, Y., Y. Xue, J. Yao, C. Xu, and R. Yu. 2025. FBXO9 mediated the ubiquitination and
degradation of YAP in a GSK-3beta-dependent manner. *J Biol Chem*. 301:110652.

Joseph, B.J., K.A. Marshall, P. Harley, J.R. Mann, F. Alessandrini, C.G. Vanoye, W. Chi, M.
Prudencio, D. Simkin, T.T. Kao, R.R. Desai, M.J. Keuss, S. Barattucci, M. Zanovello, P.R.
Mehta, J.M. DeKeyser, F. Limone, J. Lee, A.L. Brown, M.F. Leyton-Jaimes, L.A. Nash,
I.G.S. Juan, E. Aronica, B.J. Wainger, M. Shah, A. Goswami, N.A. Shneider, D.W.
Dickson, J. Burrone, C. Zhang, H. Wichterle, L. Petrucelli, J.K. Watts, A.L. George, Jr., P.
Fratta, K. Eggan, and E. Kiskinis. 2025. TDP-43-dependent mis-splicing of KCNQ2
triggers intrinsic neuronal hyperexcitability in ALS/FTD. *Nat Neurosci*.

Kellett, E.A., A.T. Bademosi, and A.K. Walker. 2025. Molecular mechanisms and consequences
of TDP-43 phosphorylation in neurodegeneration. *Mol Neurodegener*. 20:53.

Lang, R., R.E. Hodgson, and T.A. Shelkovnikova. 2024. TDP-43 in nuclear condensates: where,
how, and why. *Biochem Soc Trans*. 52:1809-1825.

Lin, L.T., A. Razzaq, S.E. Di Gregorio, S. Hong, B. Charles, M.H. Lopes, F. Beraldo, V.F. Prado,
M.A.M. Prado, and M.L. Duennwald. 2021. Hsp90 and its co-chaperone Sti1 control
TDP-43 misfolding and toxicity. *FASEB J*. 35:e21594.

Ling, J.P., O. Pletnikova, J.C. Troncoso, and P.C. Wong. 2015. TDP-43 repression of
nonconserved cryptic exons is compromised in ALS-FTD. *Science*. 349:650-655.

Love, M.I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol*. 15:550.

Ma, X.R., M. Prudencio, Y. Koike, S.C. Vatsavayai, G. Kim, F. Harbinski, A. Briner, C.M.
Rodriguez, C. Guo, T. Akiyama, H.B. Schmidt, B.B. Cummings, D.W. Wyatt, K. Kurylo,
G. Miller, S. Mekhoubad, N. Sallee, G. Mekonnen, L. Ganser, J.D. Rubien, K. Jansen-
West, C.N. Cook, S. Pickles, B. Oskarsson, N.R. Graff-Radford, B.F. Boeve, D.S.
Knopman, R.C. Petersen, D.W. Dickson, J. Shorter, S. Myong, E.M. Green, W.W.

568           Seeley, L. Petrucelli, and A.D. Gitler. 2022. TDP-43 represses cryptic exon inclusion in
569           the FTD-ALS gene UNC13A. *Nature*. 603:124-130.
570  Mann, J.R., A.M. Gleixner, J.C. Mauna, E. Gomes, M.R. DeChellis-Marks, P.G. Needham, K.E.
571           Copley, B. Hurtle, B. Portz, N.J. Pyles, L. Guo, C.B. Calder, Z.P. Wills, U.B. Pandey, J.K.
572           Kofler, J.L. Brodsky, A. Thathiah, J. Shorter, and C.J. Donnelly. 2019. RNA Binding
573           Antagonizes Neurotoxic Phase Transitions of TDP-43. *Neuron*. 102:321-338 e328.
574  Maurel, C., N.M. Scherer, L. Luu, R.A. Radford, A. Hogan, J. Merjane, A. Vidal-Itriago, E.K. Don,
575           T. Chapman, S. Cull, P. Vourc'h, L. Lisowski, A. Lee, R.S. Chung, and M. Morsch. 2025.
576           Critical impact of lysine 136 in TDP-43 phase separation, compartmentalization, and
577           aggregation in living vertebrates. *iScience*. 28:112761.
578  Mehta, P.R., A.L. Brown, M.E. Ward, and P. Fratta. 2023. The era of cryptic exons: implications
579           for ALS-FTD. *Mol Neurodegener*. 18:16.
580  Mo, M.H., J.L. Xie, Z.W. Chen, Q.Y. Yang, X.W. Hu, X.H. Zhou, J.L. Pathak, L.J. Wang, C. Liu,
581           and L.J. Chen. 2024. Niobium Nanoparticles Activates Endothelial Tip Cells and
582           Promotes Angiogenesis by FOXA1-PAK4-YAP Signaling Pathway. *Acs Appl Nano Mater*.
583           7:20123-20137.
584  Neumann, M., D.M. Sampathu, L.K. Kwong, A.C. Truax, M.C. Micsenyi, T.T. Chou, J. Bruce, T.
585           Schuck, M. Grossman, C.M. Clark, L.F. McCluskey, B.L. Miller, E. Masliah, I.R.
586           Mackenzie, H. Feldman, W. Feiden, H.A. Kretzschmar, J.Q. Trojanowski, and V.M. Lee.
587           2006. Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic
588           lateral sclerosis. *Science*. 314:130-133.
589  Nguyen, H.N. 2022. Generation of iPSC-Derived Brain Organoids for Drug Testing and
590           Toxicological Evaluation. *Methods Mol Biol*. 2474:93-105.
591  Pinarbasi, E.S., T. Cagatay, H.Y.J. Fung, Y.C. Li, Y.M. Chook, and P.J. Thomas. 2018. Active
592           nuclear import and passive nuclear export are the primary determinants of TDP-43
593           localization. *Sci Rep*. 8:7083.
594  Putri, G.H., S. Anders, P.T. Pyl, J.E. Pimanda, and F. Zanini. 2022. Analysing high-throughput
595           sequencing data in Python with HTSeq 2.0. *Bioinformatics*. 38:2943-2945.
596  Rubien, J., E. Lavorando, G. Padilla, and J. Shorter. 2022. Identifying Therapeutic Inhibitors of
597           TDP43 Phase-Separation. *Faseb Journal*. 36.
598  Rummens, J., and S. Da Cruz. 2025. RNA-binding proteins in ALS and FTD: from pathogenic
599           mechanisms to therapeutic insights. *Mol Neurodegener*. 20:64.
600  Seddighi, S., Y.A. Qi, A.L. Brown, O.G. Wilkins, C. Bereda, C. Belair, Y.J. Zhang, M. Prudencio,
601           M.J. Keuss, A. Khandeshi, S. Pickles, S.E. Kargbo-Hill, J. Hawrot, D.M. Ramos, H.
602           Yuan, J. Roberts, E.K. Sacramento, S.I. Shah, M.A. Nalls, J.M. Colon-Mercado, J.F.
603           Reyes, V.H. Ryan, M.P. Nelson, C.N. Cook, Z. Li, L. Screven, J.Y. Kwan, P.R. Mehta, M.
604           Zanovello, M. Hallegger, A. Shantaraman, L. Ping, Y. Koike, B. Oskarsson, N.P. Staff,
605           D.M. Duong, A. Ahmed, M. Secrier, J. Ule, S. Jacobson, D.S. Reich, J.D. Rohrer, A.
606           Malaspina, D.W. Dickson, J.D. Glass, A. Ori, N.T. Seyfried, M. Maragkakis, L. Petrucelli,
607           P. Fratta, and M.E. Ward. 2024. Mis-spliced transcripts generate de novo proteins in
608           TDP-43-related ALS/FTD. *Sci Transl Med*. 16:eadg7162.
609  Shorter, J. 2019. Phase separation of RNA-binding proteins in physiology and disease: An
610           introduction to the JBC Reviews thematic series. *J Biol Chem*. 294:7113-7114.
611  Suk, T.R., and M.W.C. Rousseaux. 2020. The role of TDP-43 mislocalization in amyotrophic
612           lateral sclerosis. *Mol Neurodegener*. 15:45.
613  Tziortzouda, P., L. Van Den Bosch, and F. Hirth. 2021. Triad of TDP43 control in
614           neurodegeneration: autoregulation, localization and aggregation. *Nat Rev Neurosci*.
615           22:197-208.
616  van Eersel, J., Y.D. Ke, A. Gladbach, M. Bi, J. Gotz, J.J. Kril, and L.M. Ittner. 2011. Cytoplasmic
617           accumulation and aggregation of TDP-43 upon proteasome inhibition in cultured
618           neurons. *PLoS One*. 6:e22850.

Yu, H., S. Lu, K. Gasior, D. Singh, S. Vazquez-Sanchez, O. Tapia, D. Toprani, M.S. Beccari, J.R.
Yates, 3rd, S. Da Cruz, J.M. Newby, M. Lafarga, A.S. Gladfelter, E. Villa, and D.W.
Cleveland. 2021. HSP70 chaperones RNA-free TDP-43 into anisotropic intranuclear
liquid spherical shells. *Science*. 371.
Zeng, Y., A. Lovchykova, T. Akiyama, S.L. Rayner, V. Maheswari Jawahar, C. Liu, O. Sianto, C.
Guo, A. Calliari, M. Prudencio, D.W. Dickson, L. Petrucelli, and A.D. Gitler. 2025. TDP-43
nuclear loss in FTD/ALS causes widespread alternative polyadenylation changes. *Nat*
*Neurosci*. 28:2180-2189.
Zhang, J., J. Hu, R. Liu, T. Zhou, X. Luo, P. Liang, Z. Xie, Q. Zhao, Y. Chen, D. Du, C. Liu, Y.
Zheng, D. Li, and B. Wang. 2025a. YAP maintains the dynamics of TDP-43 condensates
and antagonizes TDP-43 pathological aggregates. *Nat Cell Biol*. 27:1148-1160.
Zhang, Q., M. Liu, X. Fan, N. Chin, Y. Xu, J. Suh, S. Amniouel, K. Linask, J. Zou, M. Hafner, L.
Ma, W. Zheng, and Y. Ye. 2025b. A human forebrain organoid model phenocopies
dysregulated RNA and protein homeostasis in ALS/FTD-associated TDP-43
proteinopathies. *BioRxiv*.

**Figure legends**

**Figure 1 A chemical genetic screen identifies TDP-43 phase modulators**

**(A)** The workflow of the chemical genetic screen.

**(B)** Dose dependent reduction of anisosome number by identified chemicals. DLD1 TDP-43 2KQ-Clover

cells were treated with doxycycline to induce anisosome for 24 h and then treated with the drugs as

indicated for 15 h. Cells were imaged for anisosome count.

**Figure 2 Two distinct types of TDP-43 phase modulators**

**(A)** DLD1 TDP-43 2KQ-Clover cells treated with doxycycline for 24 h were further treated with the

indicated compounds for 6 h and imaged (SPA, Spautin-1, 5 μM; KPT, KPT-276, 15 μM; CP, CP-673451,

30 μM; TRP, Tripterin, 1 μM; BTZ, Bortezomib, 10 nM). Scale bar, 5 μm.

**(B)** FRAP analyses of anisosomes after treatment with the indicated drugs for 3-5 h. Circles indicate

bleached areas. Scale bars, 1 μm.

**(C, D)** Quantification of the experiments represented in B. FL, Fluorescence.

**(E)** Reverse FRAP analyses of anisosome dynamics. The areas indicated by the dashed lines were

bleached. Scale bar, 5 μm.

**(F)** Quantification of the fluorescence loss over time in unbleached anisosomes as shown in E. N=4-6.

**(G)** Quantification of the initial rate of fluorescence loss in unbleached anisosomes. ****, $p<0.0001$; ***,

$p<0.001$ by unpaired Student's t-test.

**Figure 3 A genome-wide siRNA screen identifies modifiers of TDP43 phase behavior**

**(A)** The workflow of the siRNA genetic screen. GW, genome-wide; AS, anisosome; HT, high-throughput.

**(B)** Pathway analysis of genes whose knockdown reduces anisosome number in cells. The relative

anisosome count are indicated by both color and size of the nodes.

**(C)** GO molecular function analysis of TDP-43 phase modifiers.

**(D)** A list of identified genes linked to ALS in C.

**Figure 4 Anisosome phase behavior is modulated by RNA splicing and protein translation**

**(A)** The splicing inhibitor Pladienolide-B (PlaB) reduces anisosome number in a dose dependent manner.

Representative images of anisosome-induced cells treated with DMSO (control), 5 nM, or 20 nM PlaB for

16 h. Scale bar: 10 μm.

**(B)** Quantification showing the number of anisosome (AS) per cell in TDP-43 2KQ-Clover cells treated

with PlaB as indicated. * $p <0.05$, ** $p <0.01$, **** $p < 0.0001$ by One-way ANOVA. n=3 biological

repeats.

**(C)** Representative FRAP images of anisosomes in cells treated for 16 h with DMSO or PlaB (20 nM).

BB, before photobleaching, right after photobleaching (0 s), or 4 and 30 seconds after photobleaching (4 s

and 30 s). Scale bar, 1 μm.

**(D)** The graph shows the quantification of the remaining TDP-43 fluorescence (FL) in C. Error bars

indicate mean ± SD, N = 28 for control and 23 for PlaB-treated cells.

**(E)** Live cell imaging of anisosome fusion in TDP-43 2KQ-Clover cells treated with 20 nM PlaB for 5 h

before tracking the fusion. Representative images from Movie S1 showing fusion events indicated by

arrows. Scale bar, 1 μm.

**(F)** Representative images of anisosome-induced (24 h) cells treated with DMSO (control) or ANS (200

nM) for 16 h. Scale bar, 1 μm.

**(G)** Quantification of the number of anisosomes per cell in randomly selected images of DMSO-,

Cycloheximide (CHX)-, or ANS-treated cells. **** p < 0.0001 by one-way ANOVA; each dot represents

an image with at least 20 cells counted. n=3 biological repeats.

**(H)** Quantification of anisosome size in control or cells treated with CHX or ANS. ****, p < 0.0001 by

One-way ANOVA. n=3 biological repeats. AU, arbitrary unit.

**(I)** As in **C** except that anisosome-induced cells were treated with CHX or ANS before photobleaching.

Scale bar 1 μm.

**(J)** Quantification of fluorescence recovery in CHX- or ANS-treated cells vs the DMSO control.

**Figure 5 XPO1 regulates anisosome liquid-to-solid transition**

**(A)** Pharmacological inhibition of XPO-1 with Leptomycin B (LMB) reduces the number of anisosome.

The graph on top indicates the experimental design. Arrows show cells with enlarged anisosomes. Scale

bar 10 μm

**(B)** Quantification of anisosome numbers per cell in experiments represented by A. Error bars indicate

mean ± SD; **** p <0.0001 by one-way ANOVA. n=3 biological repeats.

**(C)** LMB dose dependently reduces anisosome number. Anisosome were induced in TDP-43 2KQ-Clover

cells followed by treatment with LMB at the indicated concentrations for 16 h. The histogram shows the

distribution of cells (50-90 cells/condition) by anisosome count.

**(D)** FRAP experiments demonstrate that anisosomes remain in a liquid phase following LMB treatment

(200 nM, 16 h). Anisosome-induced (24 h) cells were treated with DMSO or LMB for 16 h and then

photobleached at the indicated areas. Scale bar 1 μm.

**(E)** Quantification of the FRAP experiment in D.

**(F)** Time-lapse confocal microscopy detects anisosome fusion after LMB (200 nM, 5 h) treatment in

TDP-43 2KQ-Clover cells. Arrows indicate fusion events. Scale bar, 1 μm

**(G)** Overexpression of mCherry-tagged XPO-1 in TDP-43 2KQ-Clover cells induces cytoplasmic TDP-43 puncta. TDP-43 2KQ-Clover cells (green) transfected with mCherry-XPO1 (red) were stained with DAPI (blue) to label nuclei (dashed lines). Cells were imaged 48 h post-transfection. Panels 1, 2 show a representative confocal section, while panels 3-6 show reconstructed 3-D views. The position and volume of anisosomes were also presented in magenta in a surface-rendered view (SRV) in panel 5. The arrow in panel 3 indicates an example of cytoplasmic TDP-43 aggregate. Scale bar, 10 μm.

**(H)** Quantification of the percentage of cells showing cytosolic TDP-43 puncta in XPO-1 positive (pos) and negative (neg) cells in randomly selected fields from 3 independent experiments. **** $p < 0.0001$ by unpaired Student's t-test.

**(I)** FRAP-based confocal imaging reveals the transition of anisosomes into a gel-like state upon XPO-1 overexpression. Scale bar, 1 μm.

**(J)** Anisosome formation changes endogenous XPO-1 localization. TDP-43 2KQ-Clover before or after anisosome induction were stained with anti-XPO-1 antibodies (red) and DAPI (blue). The bleached areas were indicated by dashed lines. Scale bar, 10 μm.

**(K)** Quantification of nuclear endogenous XPO-1 levels in individual cells (indicated by dots) before or after anisosome induction. **** $p < 0.001$ by unpaired Student's t-test. n=2 independent repeats.

**(L)** A schematic model depicting how the nuclear XPO-1 activity influences TDP-43 liquid-to-phase transition. AS, anisosome.

**Figure 6 Inhibition of XPO-1 reduces TDP-43 hyperphosphorylation but only partially restore neuronal gene expression in K181E organoids**

**(A)** Confocal fluorescence imaging reveals significant reduction in phosphorylated TDP-43 (p-TDP43) in K181E/K181E organoids following KPT-276 treatment. Organoids of the indicated genotypes at day 87 were treated with DMSO as a control, or with KPT-276 (20 nM) for 35 days. Organoids were fixed, sectioned, and stained with antibodies against TUJ1, a neuronal marker (Magenta), TDP-43 (green) and p-TDP-43 (red). n= 3-5 organoids from two individual batches.

**(B)** Quantification of p-TDP-43 Mean Spot Volume (top) and percentage of cells lacking nuclear TDP-43 (bottom) in experiments represented by **A**. Error bars indicate mean ± s.e.m., * $p < 0.05$, **** $p < 0.0001$ by one-way ANOVA, n=3 organoids.

**(C)** Gene Ontology (GO) analysis of genes whose expression was disrupted in K181E/K181E organoids but restored by KPT-267 as shown in Supplemental Fig. S3. BP, Biological Pathway; CC, Cellular Component; MF, Molecular Function.

**A**

DMSO  SPA  KPT  CP

TRP  BTZ

**B**

DMSO  0"  4"  20"

BTZ  BB  0"  4"  20"

TRP  BB  0"  4"  20"

SPA

KPT

**C**

Control
BTZ
TRP

FL remaining (%)

Time (sec)

**D**

Control

KPT
SPA

FL remaining (%)

Time (sec)

**E**

DMSO  BB  0"  56"  104"

TRP

BZM

SPA

**F**

BTZ

TRP

DMSO

SPA

FL remaining (%)

Time (sec)

**G**

***
****

FL loss (%/Sec)

DMSO  BZM  TPR  SPA

748 **Supplemental Materials**

749 Movie S1 PlaB treatment induces anisosome fusion.

750 Movie S2 Blocking XPO-1-mediated nuclear export induces anisosome fusion.

752 **Supplemental Tables**

753 Table S1 A chemical genetic screen identified compounds that reduce anisosome numbers.

754 Table S2 A genome-wide siRNA screen identified potential anisosome regulators.

755 Table S3 Top 110 genes confirmed from a secondary screen as modulators of anisosome dynamics.

756 Table S4 Differentially expressed genes in K181E organoids rescued by KPT-276 (KPT).

 **Supplemental Figures**

Figure S1

**Figure S1. HSP90 inhibitor Geldanamycin converts TDP-43 anisosome to a gel-like state**

**(A)** DLD1 cells were treated with doxycycline to induce anisosome formation and then treated with

Geldanamycin (GA) at 10 μM for 4 h. Cells were stained with Hoechst (Blue) and imaged by a confocal

microscope. Scale bar, 5 μm.

**(B)** A TDP-43-bearing anisosome (dashed circle) in a GA-treated cell was photobleached and then

imaged. Scale bar, 1 μm. The graph shows the quantification. Error bars, s.e.m. (N>5 anisosomes). FL,

fluorescence.

**Figure S2. The impact of splicing inhibitor and translation inhibitor on anisosomes**

**(A)** A violin plot showing the relative size distribution of anisosomes (AS) in DLD1 cells treated with

Pladienolide-B at the indicated concentration for 16 h. ****, $p<0.0001$ by one-way ANOVA. n=3

biological repeats.

**(B)** DLD1 cells were treated with doxycycline to induce anisosome formation and then treated with

Cycloheximide (CHX) at 20 µg/mL or DMSO as a control for 16 h. Cells were stained with Hoechst

(Blue) and imaged by a confocal microscope. Scale bar, 5 µm.

Figure S3

779
780
**Figure S3. KPT-276 modestly reverses gene expression defects in K181E mutant organoids**

Heatmap showing genes whose expression was altered in K181E/K181E organoids and restored after

KPT-276 treatment.