Tracking active heterotrophic microbial communities in the Guaymas Basin deep biosphere

with BONCAT-FACS

Andrew Montgomery[1,2,3,*,#], Sylvia Nupp[1,2,3,*], Chase R. Gray[1,2,3], Zackary J. Jay[1,2,3], Virginia

Edgcomb[4], Roland Hatzenpichler[1,2,3,5,#]

1.   Department of Chemistry and Biochemistry, Montana State University, Bozeman, MT,

USA

2.   Thermal Biology Institute, Montana State University, Bozeman, MT, USA

3.   Center for Biofilm Engineering, Montana State University, Bozeman, MT, USA

4.   Woods Hole Oceanographic Institution, Woods Hole, MA, USA

5.   Department of Microbiology and Cell Biology, Montana State University, Bozeman, MT,

USA

*These authors contributed to this work equally.

#correspondence to: R.H., rolandhatzenpichler@gmail.com; A.M.,

andymontgomery2009@gmail.com

ORCIDs: A.M., 0000-0003-2798-1390; S.N., 0009-0008-1814-1211; R.H., 0000-0002-5489-

3444; Z.J.J., 0000-0003-3062-4933; C.R.G., 0009-0005-0296-1491; V.E., https://orcid.org/0000-

0001-6805-381X.

Abstract

The marine deep biosphere harbors microbial communities that drive organic matter transformations and biogeochemical cycles. Previous work on these communities has focused either on genomic characterization or metabolic activity measurements. However, to understand microbial ecophysiology in the deep biosphere taxonomic identity and metabolic function must be connected on both single-cell and ecosystem scales. In this work, we optimized a bioorthogonal non-canonical amino acid tagging fluorescence-activated cell sorting (BONCAT-FACS) workflow for low-biomass deep-biosphere sediments obtained during International Ocean Discovery Program Expedition 385 (IODP 385). BONCAT-FACS with 16S rRNA gene amplicon sequencing as well as metagenomics of sediment communities was applied to characterize translationally active communities in hydrothermally altered subsurface sediments of the Guaymas Basin. Our results revealed a heterotrophic microbial population throughout all sediments examined, with taxa translationally active down to our deepest sampling point, 154 meters below the seafloor. Based on 16S rRNA gene identities, the translationally active microbial community was dominated by heterotrophic members of the *Gammaproteobacteria*, *Bacilli*, *Deinococci*, and *Alphaproteobacteria*. These taxa are likely key contributors to cycling the large quantities of hydrothermally altered organic matter in Guaymas Basin sediments. To further elucidate the metabolic capacity of active taxa, we mapped 16S rRNA gene amplicons to metagenome assembled genomes (MAGs) previously obtained from IODP 385. These MAGs contained genes associated with C1 metabolism, carbohydrate degradation, and fermentation, indicating that active taxa leverage these metabolisms for energy conservation. Our results demonstrate that BONCAT-FACS provides high-throughput and single-cell insights into the metabolic activity of microbes in the low-biomass marine subsurface.

Importance

The marine deep biosphere spans the entire global ocean and offers a wide range of environmental

conditions for microbes to flourish. However, the identities and metabolic functions of these

microbial populations are not well understood. In this study, we investigated translationally active

microbial communities and elucidated their metabolic activity in hydrothermally impacted low-

biomass subsurface marine sediments. By combining activity measurements, amplicon

sequencing, and metagenomics data, we improve our understanding of carbon transformations and

microbial ecology of the deep biosphere. Our findings emphasize the ecological significance of

heterotrophic microbes and the carbon-degrading metabolisms they use to survive extreme

conditions. This work advances the applications of next-generation physiology techniques to

document single-cell metabolic activity in a high-throughput manner in the marine deep biosphere.

Introduction

The marine deep biosphere is one of the largest ecosystems on Earth, supporting an estimated 10^{29} cells (1, 2). Defined as sediments >1 m below the seafloor, the deep biosphere encompasses a wide spectrum of redox conditions that extend to the energetic and thermal limits of life. Biological and geological transformations of organic matter drive a dynamic, albeit slow, carbon cycle in these sub-seafloor environments (3–6). While it is clear that microorganisms are metabolically active in the deep biosphere (7, 8), low cell abundances mean the significance of their activity and contributions to biogeochemical cycles remains largely unconstrained. Direct connections between the taxonomic identity of active microbial populations and metabolic functions are necessary to understand microbial ecology and biogeochemical cycling in the marine deep biosphere.

Uncovering the metabolic capabilities of constituents of microbial communities *in situ* requires a coordinated effort that simultaneously identifies taxa and links them to ecosystem functions. Historically, cultured isolates were required in order to draw conclusions about the metabolic capabilities of specific microbial taxa. However, less than 1% of microbes can be reliably maintained in a laboratory setting, and enrichment and isolation conditions can differ substantially from *in situ* conditions, making it difficult to draw conclusions about how microbes live in their natural habitat. Further, cultivation efforts are impeded by low biomass inocula and energy limitations in the deep biosphere that select for microorganisms that grow at extremely slow rates (9, 10). More recently, culture-independent techniques have been used to elucidate connections between individual microorganisms and their metabolic functions under (near) *in situ* conditions. With the rise of high-throughput sequencing technologies and bioinformatics tools, culture-independent techniques have facilitated studies of the deep biosphere with metagenomics (11–13)

to identify microbial taxa, or metatranscriptomics (14–18) to reveal which genes are expressed *in*

situ. However, while these approaches offer powerful insights into metabolic potential and

community level activity, they do not link specific taxa with metabolic activity *in situ*.

Bioorthogonal non-canonical amino acid tagging (BONCAT) paired with fluorescence-activated

cell sorting (BONCAT-FACS) is a Next-Generation Physiology approach (19) that can be used to

identify which taxa were translationally active in complex environmental samples during a

relatively short incubation without culturing individual organisms (20–24). BONCAT leverages

the incorporation of a synthetic amino acid, such as *L*-homopropargylglycine (HPG), an analog of

L-methionine, into newly synthesized proteins and its later detection using azide-alkyne click

chemistry (20, 25). Using fluorescence as the mode of detection, FACS can be used to sort

translationally active cells for downstream genomic or amplicon sequencing and identification of

BONCAT-active taxa (21) and even for cultivation efforts (20, 26). BONCAT-FACS, coupled

with 16S rRNA gene amplicon sequencing, offers a powerful approach to identify anabolically

active taxa on a community level. This approach has been used to identify active taxa in marine

sediments (27), soils (24, 28), the human microbiome (29–31), and other complex environments

(32–35). Here, we used BONCAT-FACS to identify the translationally active community and,

coupled with community shotgun metagenomics, assess the metabolic capacity of translationally

active taxa in the Guaymas Basin deep biosphere.

Guaymas Basin, located in the Gulf of California, is a hydrothermally active rift basin

characterized by rapid seafloor spreading and high sedimentation rates. Terrestrial inputs and

highly productive waters in the Gulf of California deposit organic matter on the seafloor at rates

>1 mm yr^{-1}, generating sediment packages that can be hundreds of meters thick (36).

Thermocatalytic alteration of the organic matter, influenced by basaltic sill intrusions and

hydrothermal circulation (37, 38), produces a complex mixture of potentially bioavailable organic

compounds in the sediment (39). This includes methane, low-molecular-weight organics,

hydrocarbons, and crude oil (40–45). International Ocean Discovery Program Expedition 385

(IODP 385) drilled the subsurface of the Guaymas Basin to investigate how microbial communities

drive organic carbon cycling in the hydrothermally altered deep biosphere. In total, holes were

drilled at eight sites (U1545-U1552) to capture unaltered background sediment and a spectrum of

hydrothermal sediments including extremely hot sediments above a more recently intruded sill-

sediment system (46). To explore the microbial ecology of the Guaymas Basin subsurface, we

tracked translationally active taxa from six depths from five sites drilled during IODP 385. We

optimized a previously established BONCAT-FACS workflow with 16S rRNA gene amplicon

sequencing to identify translationally active taxa (27, 32) and to assess diversity across

biogeochemical gradients. We coupled this with metagenomic analyses from IODP 385 (12) to

provide direct insight into the metabolic capacity of active taxa within the Guaymas Basin deep

biosphere.

Materials and Methods

Sample collection and incubation

Sediment cores were collected from Guaymas Basin during IODP expedition 385 onboard drilling

vessel *JOIDES Resolution.* Samples were taken at six depths, ranging from 0.76 to 154 mbsf, from

five drill holes (Table 1). Intact, round samples were aseptically transferred to the microbiology

lab after retrieval on deck and stored at 4°C in heat-sealed gas-tight bags to maintain anoxic

conditions until processing. Working in an anoxic glove chamber, sediments were homogenized

with N_2-gassed sterile bottom water diluted three times to form a slurry. For each sample, four

replicate aliquots of 40 or 120 mL, depending on depth (shallow vs. deep), were distributed into

Pyrex bottles and sealed with a butyl stopper and aluminum crimp. We increased the volume of

deeper samples during incubation to increase the likelihood of detecting enough active cells for

sorting. The amino acid analog *L*-homopropargylglycine (HPG; Vector Laboratories, previously

Click Chemistry Tools), a synthetic analog of *L*-methionine, was added to three replicates to a

final concentration of 50 µM. The fourth replicate, which served as a control, was not amended

with HPG. After HPG addition, samples were incubated at *in situ* temperature (Table 1) for seven

days.

After seven days, samples were washed with 1X PBS to terminate incubations and remove unused

HPG. A 5 mL subsample was frozen in a cryovial at -80°C for whole sediment DNA extraction.

The remaining sediment was transferred to 50 mL centrifuge tubes, amended with 10% glycerol-

TE (9:1 sediment to glycerol-TE), and stored at -80°C for subsequent steps of the BONCAT-FACS

workflow back at Montana State University.

Cell Extractions

Cells were extracted following a modified protocol (27, 47) that we optimized to extract cells from

low-biomass subsurface marine sediments. Samples preserved in glycerol-TE were thawed

overnight at 4°C and centrifuged at 3,200 × *g* for 30 min to pellet sediment. For each sample,

sediment was diluted 1:3 in a filter-sterilized 3.5% NaCl solution (0.2 µm pore size) and briefly

homogenized by vortex. From this slurry, 200 µL aliquots were further diluted in 100 µL of pure

methanol and 700 µL of filter-sterilized 3.5% NaCl solution. For each experimental replicate, eight

extraction replicates were prepared in 2 mL microcentrifuge tubes to ensure enough biomass was

extracted for downstream processing. Extraction blanks with phosphate-buffered saline (1X PBS)

were processed in parallel to assess contaminant introduction during extraction. Extraction

replicates were homogenized by vortex for 30 min at room temperature on setting 7 (approximately

2,000 rpm, VortexGenie2, Scientific Industries, Inc., Bohemia, NY, United States). Cells were

separated by density centrifugation by layering 1 mL of 50% Nycodenz solution below the

sediment slurry with a syringe and needle and centrifuging at $3,000 \times g$ for 10 min. After

centrifugation, the upper and inner phases of the supernatant were collected by pipetting and

pooling extraction replicates in a 15 mL centrifuge tube. The remaining supernatant was discarded,

leaving only the sediment pellet. To detach cells from particles, the pellet was resuspended in 800

µL of 3.5% NaCl solution, 100 µL of pure methanol, and 100 µL of detergent mix (100 mM

ethylenediamine tetraacetic acid (EDTA), 100 mM sodium pyrophosphate, 1% Tween80 in 3.5%

NaCl solution (47), briefly vortexed to homogenize, and sonicated. The sonication step was done

on ice at 125 W and 200 µm amplitude for 5 sec, followed by a 20 sec pause and repeated for 5

cycles to minimize cell rupture. Subsequently, samples were subjected to the same Nycodenz

density gradient centrifugation described above. The upper and inner phases of the supernatant

from each extraction replicate were combined with the pooled supernatant from the first density

gradient centrifugation step. The pooled fractions were centrifuged at $16,000 \times g$ for 10 min to

pellet extracted cells, and the supernatant was gently decanted, leaving ~1 mL of supernatant

remaining to resuspend pelleted cells. This cell suspension was transferred to a 1.5 mL

microcentrifuge tube, and cells were centrifuged again at $18,000 \times g$ for 10 min and concentrated

in 100 µL of PBS:ethanol (1:1 ratio of 1X PBS and pure ethanol). Concentrated cells were stored

overnight at -20°C to permeabilize cells for click chemistry.

Click Chemistry and Fluorescence-Activated Cell Sorting

The following day, cell extracts were click-stained to fluorescently tag cells that incorporated HPG

into newly synthesized proteins (48). A bulk click solution was prepared, and 200 µL was added

to each cell extract. The final click reaction comprised of 500 µM tris-

hydroxypropyltriazolylmethylamine (THPTA; Vector Laboratories), 100 µM copper sulfate

pentahydrate, 5 mM sodium *L*-ascorbate, 5 mM aminoguanidine HCl, and 5 µM of Cy5 picolyl-

azide dye (Vector Laboratories) in 1X PBS. The click solution and cell extract were homogenized

and then incubated at room temperature, in the dark, for 30 min. The click solution was washed

away with three washes in 1 mL of 1X PBS with centrifugation at $18{,}000 \times g$ for 5 min between

each wash. A 50 µL aliquot of the final suspension was kept aside to be used as a "presort" total

cell-extractable fraction for PCR. This subsample was stored in a fresh 1.5 mL microcentrifuge

tube at -80°C until freeze-thaw DNA extraction.

The remaining cell extract was immediately subjected to fluorescence-activated cell sorting

(FACS). Prior to loading, each sample was passed through a 35 µm mesh filter to remove sediment

particles and diluted to a final volume of ~1.5 mL with 1X PBS. FACS was performed with a Sony

SH800S (Sony Biotechnology, San Jose, CA) equipped with a 70 µm sorting chip using 1X PBS

as sheath fluid. The sheath fluid was 0.2 µm filtered and UV sterilized overnight while stirring to

remove salt crystals in the PBS and minimize contamination of the FACS fluidics system (32). A

series of gates was manually drawn to exclude large sediment particles. The final sorting gates

were drawn based on the Cy5 relative fluorescence and forward scatter (Fig. S1) of the No-HPG

controls. A "Cy5-positive" gate was drawn so that only BONCAT-positive cells that incorporated

HPG into newly synthesized proteins were sorted. Each sample was sorted until the sample had

been exhausted, regardless of the number of events sorted (Table S1). Sorted fractions were

centrifuged at $18,000 \times g$ for 5 min to pellet cells. After removing the supernatant, cells were

resuspended in 50 µL of nuclease-free water and stored at -80°C for freeze-thaw DNA extraction.

DNA extraction

DNA from whole sediments was extracted for each replicate sample after HPG incubation using

the FastDNA Spin Kit for Soil (MP Biomedicals, Irvine, CA), following the manufacturer's

protocol. After elution, DNA was quantified via the Qubit High Sensitivity assay (Invitrogen,

Carlsbad, CA) and stored at -80°C until PCR. DNA was extracted from presort and sorted

(BONCAT active) fractions by triplicate repeated freeze-thaw cycles to lyse cells (32). The freeze

cycles were done at -80°C for 20 min, and thawing was done at 99°C for 10 min. These DNA

extracts were stored at -80°C for further processing.

16S rRNA Gene Amplicon Sequencing

Amplification of bacterial and archaeal 16S rRNA genes was performed following the Earth

Microbiome protocol (49) using updated primers 515F (forward primer, 5′-

GTGYCAGCMGCCGCGGTAA-3' (50) and 806R (reverse primer, 5′-

GGACTACNVGGGTWTCTAAT-3' (51). Polymerase chain reaction (PCR) was performed in a

final volume of 50 µL consisting of 20 µL of template DNA, 20 µL Invitrogen Platinum Taq II 2x

Master Mix, 1 µL of each primer at a concentration of 10 µM, and 8 µL nuclease-free water for

freeze/thaw extracted cells, and 25 µL consisting of 5 µL of template DNA, 10 µL Invitrogen

Platinum Taq II 2x Master Mix, 0.5 µL of each primer at a concentration of 10 µM, and 9 µL

nuclease-free water for DNA extracted with the FastDNA Spin Kit for Soil. The thermocycler

conditions were: 94°C for 3 min, then 28 cycles of 94°C for 45 sec, 50°C for 60 sec, and 72°C for

90°C, followed by an elongation step of 72°C for 10 min. A selection of PCR products was checked

for the expected length on a 1% agarose gel. All products were then purified using AMPure XP

beads following the manufacturer's protocol with a final elution volume of 35 µL. For Illumina

sequencing, Illumina dual index barcodes and sequencing adapters were attached in a second PCR

reaction. The final volume for this reaction was 25 µL, consisting of 5 µL of purified amplicon,

10 µL Invitrogen Platinum Taq II 2x Master Mix, 0.5 µL of each primer at a concentration of 10

µM (i5 and i7 primers), and 9 µL nuclease-free water. The thermocycler conditions were: 95°C

for 3 min, then 8 cycles of 95°C for 30 sec, 55°C for 30 sec, 72°C for 30 sec, followed by an

elongation step of 72°C for 5 min. The PCR products were again purified using AMPure XP beads,

and then were quantified in triplicate reactions using Quant-iT Picogreen dsDNA assay

(Invitrogen) on a Biotek Synergy H1 Hybrid microplate reader. Purified amplicons were then

pooled with 7.5 ng of DNA each, or up to 30 µL if a sample had less than 7.5 ng of DNA. The

pooled library was quantified with the Qubit high-sensitivity assay. Purification, quality

assessment, and sequencing of pooled libraries were performed at the Molecular Research Core

Facility at Idaho State University using Illumina's MiSeq chemistry for 2 x 250 bp paired-end

reads.

Bioinformatics Analysis

Amplicon sequence data were processed with QIIME2 (version 2025.7) (52). Barcoding and

primer sequences were removed using *cutadapt* before truncating the forward (137 bp) and reverse

(139 bp) reads based on sequencing quality. DADA2 was subsequently used to filter, denoise, and

merge the reads. Taxonomic assignments for each ASV (amplicon sequence variant) were

determined with the Silva 138.2 database using the *sklearn* method (53). Contaminant ASVs were

identified using the prevalence model (threshold = 0.5) in the R package *decontam* (54) and manual

inspection of the dataset to compare samples to the no HPG controls and process blanks. Observed

diversity and Shannon's diversity metrics were calculated using the *phyloseq* package in R (55),

with replicate samples pooled for analysis. Linear fold enrichment was calculated in Equation 1 as

the ratio of the fractional relative abundance of ASVs affiliated with a given class between the

presort and sorted fractions:

$Fold\ Enrichment\ = (\frac{max(RAsort,RApresort)\ +0.1}{min(RAsort,RApresort)\ +0.1} - 1)\ * sgn(RA_{sort} - RApresort)$ (1)

Where RA$_{sort}$ and RA$_{presort}$ are the fractional relative abundances of a given taxonomic class in the

sorted and presort fractions, respectively. Fold enrichment is calculated in the first part of the

function, and a small value (0.1) is added to each relative abundance to avoid dividing by zero.

Subtracting that value by one shifts the ratio so that equal relative abundances give zero. The *sgn*

function returns the sign of the differences to force the fold enrichment value positive.

Analysis of Metagenome Assembled Genomes

Metagenome assembled genomes (MAGs) were previously generated from IODP 385 (12). In

short, trimmed reads from all drill holes were pooled and co-assembled using MEGAHIT 1.2.9,

and MAGs that were at least 50% complete and contained less than 10% contamination were then

used for downstream analyses. Taxa in control samples were identified as potential contaminants

and were removed from downstream analysis. Taxonomic identities of these MAGs were assigned

using GTDB-Tk 2 (56). MAG contamination and completeness were checked using CheckM (57).

Barrnap (v0.9) was used to identify 16S rRNA gene sequences from these MAGs. The 16S rRNA

gene sequences of these MAGs, if present, were then compared to the 16S rRNA gene amplicon

sequences from sorted samples generated in this study for sequence similarity. From a BLAST

nucleotide similarity comparison, we used a cutoff of 90% for %ID and 99% for %hit to identify

sequences that matched.

The organism identities of these matches were then compared between the MAGs and amplicon

sequences to determine if they were identified as the same organism. Importantly, the MAG and

16S rRNA gene amplicon sequences were independently sequenced and taxonomy was assigned

using different databases (Silva 138.2 for 16S rRNA gene amplicon sequences, and GTDB-Tk

2.1.0 for MAGs), and taxonomic identification varied between the two. Organism identity was

matched based on nomenclature changes in the literature to the best of our ability. For consistency

across our datasets, the Silva 138.2 taxonomic identification is used throughout the manuscript.

Six matches were identified that met our cutoff criteria of 90% for %ID and 99% for %hit, five

bacteria and one archaea.

MAGs were annotated with Prokka separately for archaea and bacteria. Individual genes of interest

for the environment were selected based on Dombrowski *et al.* (58). Carbon metabolism pathways

were identified using KEGG BlastKOALA. Genes for the Wood-Ljungdahl pathway were

identified using an internal protein database generated from cultured methanogens because of the

presence of MAGs in the same class as other methanogens.

Results & Discussion

Sampling Sites

Samples for BONCAT incubations were collected from drill hole sites, U1545 to U1549, along a

transect across Guaymas Basin during IODP Expedition 385 (Table 1). Sites U1545 and U1546

are located ~52 km off-axis of the central spreading center, and include a relatively unaltered

reference (U1545) site and a more thermally altered (U1546) site with deep sediments (~350-430

mbsf) influenced by a now thermally-equilibrated ancient sill emplacement. Sites U1547 and

U1548 were drilled within and adjacent to the much hotter Ringvent site. Ringvent is shaped by a

shallow, more recently emplaced sill, exhibits a sill-driven hydrothermal mound, and allows us to

investigate links between sediment alteration, geochemical gradients, and microbial responses to

elevated temperatures. Site U1549 targeted an off-axis cold seep near a subsurface sill, where

methane seepage and hydrates indicate persistent subsurface fluid migration. Sample depths

ranged from 0.8 m at site U1546 to 155 m at U1545, and temperatures were between 2.8°C and

46.6°C (Table 1). Sediment and porefluid geochemical data relevant to microbial activity are

outlined in Table 1, and comprehensive datasets are available in the IODP 385 expedition reports

(59–62).

Workflow development

We expanded the applications of BONCAT by refining a workflow for detecting translational

activity of microbial communities in extremely low-biomass marine deep-biosphere sediments that

maximized cell recovery and minimized interference from sediment autofluorescence. To optimize

this workflow and test the efficacy of cell extraction protocols, we spiked IODP 385 sediment with

Escherichia coli (*E. coli*) cells that had been grown with HPG. These sediments had been preserved

analogously to the IODP 385 BONCAT samples and incubated without HPG. The presence of

BONCAT-positive *E. coli* in cell extracts was used to assess the efficacy of different cell extraction

protocols to maximize cell recovery. Additionally, we sought to minimize particulate matter in the

final cell suspension. Autofluorescent sediment particles can interfere with FACS analyses when

the signal overlaps with the target fluorophore and distorts forward and back scatter, complicating

the accurate detection of cells.

The final cell extraction procedure was adapted from an established protocol (27, 47) and consisted

of two primary cell separations. The first wash with methanol only was used to extract cells that

are more susceptible to lysis from detergents. The second step combined sonication with a

detergent mixture (sodium pyrophosphate, Tween80, and methanol) to detach cells from sediment

particles. A 50% Nycodenz gradient was used between detergent wash steps to phase-separate

detached cells from sediment. Based on the number of cells and amount of autofluorescent

sediment in the extracted suspension, we determined that pooling eight technical replicate extracts

was sufficient to extract cells from low-biomass samples for downstream processing with FACS.

We visualized labeling of BONCAT-positive cells from click chemistry in the presort and sorted

fractions with fluorescence microscopy (Fig. S2). During method testing and sample analysis, the

presort fraction contained cells with and without HPG incorporation, while the sorted fraction only

contained HPG-positive cells, attesting to the sorting of BONCAT-positive cells only from low-

biomass marine deep biosphere sediments.

Initially, we aimed to assess the translational activity of fungi with BONCAT-FACS in this study,

so we developed this cell extraction protocol to include fungal cells. We used two fungal isolates

from Guaymas Basin that are representative of the fungal community in IODP 385 sediment (63),

and tested extraction methods similar to our use of *E.coli* above. Based on our testing, this cell

extraction method can be used to separate yeast and filamentous fungal cells from sediments.

Furthermore, these fungal strains incorporate HPG in liquid media and can be detected with click

chemistry and fluorescence microscopy (Fig. S3), demonstrating that BONCAT workflows can be

applied to marine fungi. However, due to issues with downstream processing, we could not reliably

detect fungi in the presort and sorted fractions with internal transcribed spacer (ITS) amplicon

sequencing. While the exact reason for this issue is unknown, poor extraction efficiency, low

fungal biomass (or absence of fungi), the inability to sort certain fungal cells with FACS, or poorly

matched PCR primers for the ITS region in natural fungal populations (64, 65) may have all

impeded the successful detection of fungi in our samples. Therefore, we chose to exclude any

further discussion of fungi from this manuscript and focus on bacteria and archaea inhabiting the

Guaymas Basin deep biosphere.

Diversity of the total, cell extractable, and active prokaryotic community

We combined BONCAT-FACS with 16S rRNA gene amplicon sequencing of sorted cells to

identify translationally active bacteria and archaea in the Guaymas Basin deep biosphere. To assess

community compositions, the observed diversity (the quantity of unique ASVs) and the Shannon's

diversity index were determined for each sample (Fig. 1A-B). For all samples, observed and

Shannon's diversity indices were higher in the whole sediment DNA extract fraction than in the

cell-extracted presort and BONCAT-active sorted fractions (Fig. S4). Generally, the diversity of

the total community decreased with depth, reflecting the decrease in cell abundance with depth.

The two deepest samples, 1547B-8H2 and 1545B-19F3 (66 and 155 mbsf, respectively), exhibited

the smallest difference in diversity metrics across all three fractions, especially with respect to the

DNA extract. Compared to shallower samples, a larger portion of the total microbial community

was detected in the BONCAT active fraction in these deeper sediments, where there are relatively

few cells, carbon is more recalcitrant, energetic conditions are less favorable, and geochemical

conditions remain stable over extended timescales (66, 67). This suggests that the microbial

population is highly specialized to fill limited available niches in the deep biosphere.

To evaluate the changes in diversity metrics between the presort and sorted fractions, we calculated

fold enrichment (Fig. 1C-F) between these fractions for the most abundant ASVs, which were

affiliated with *Gammaproteobacteria*, *Alphaproteobacteria*, *Bacilli*, and *Deinococci* (Fig. 2 and

Table S2). Generally, these abundant taxa were not overly represented in either fraction

(enrichment values near zero) or were enriched in the presort fraction (*Deinococci* and *Bacilli*),

suggesting that only a portion of those taxa detected in the presort fraction were actually

translationally active. In several cases, corresponding ASVs were not detected in either the presort

or sorted fraction, which skewed enrichment toward the respective other fraction. For instance, in

sample 1548B-2H3, no sequences affiliated with *Alphaproteobacteria* (Fig. 1D) were detected in

the presort fraction, artificially inflating the fold enrichment toward the sorted fraction. Similarly,

no *Deinococci* affiliated ASVs were observed in the sorted fraction for one replicate from 1547B-

1H2 (Fig. 1F). The inability to detect a given ASV in a specific fraction could skew diversity

indices, making it difficult to interpret diversity data in those samples.

Observed diversity of unique ASVs increased from the presort to the sorted fractions in all samples

(Fig. 1A), and Shannon's diversity increased in 1549B-3H2 and 1547B-8H2 between those

fractions (Fig. 1B). It should be expected that diversity would decrease between the presort and

the BONCAT active fraction to reflect that only a portion of the community is translationally active

and detectable with BONCAT-FACS. One explanation for this outcome is that there were

substantially fewer reads in many of the presort samples than in the sorted samples (Fig. S5), likely

due to the relatively small subsample (50 µL) taken from the presort samples for DNA extraction

and low amount of biomass in the sample. More ASVs were detected when the number of reads

was higher (Fig. S5), which would influence alpha diversity metrics. In the two samples in which

the Shannon's diversity increased in the sorted fraction (1547B-8H2 and 1549B-3H2), across all

replicates, we detected between three and eleven unique class-level ASVs in the presort samples

and between 12 and 30 in the sorted samples. This indicates that sequencing depth directly impacts

the detection of additional and/or rare taxa that may have been, in some cases, missed in the presort

fraction.

We observed major differences in Shannon's diversity indices across the BONCAT workflow

(DNA extract to presort to sorted fractions). Shannon's diversity was highest in the DNA extract

relative to the presort or sorted fractions for nearly every sample (the only exception was the sorted

fraction for 1547B-8H2) (Fig. S4). This is expected for the marine subsurface as extracellular DNA

in the sediments may have been amplified with PCR, altering community composition and

artificially inflating diversity metrics in the initial DNA extract fraction (24). Diversity indices

were generally lower in the presort and sorted fraction, as cell extraction for BONCAT-FACS is

known to introduce biases (27, 32) that could cause large discrepancies in diversity metrics

between the DNA extract and presort fractions. To address this, we optimized the cell extraction

procedure for these low-biomass deep biosphere samples, but biased cell recovery is a challenge

for any complex sample (19, 68), especially from Guaymas Basin sediments that are enriched with

hydrothermally altered organic carbon. Ideally, cell extraction procedures should be tuned to

specific samples based on experimental goals and biogeochemical conditions.

Abundant Taxa and Active Community

To assess the active community within the Guaymas Basin deep biosphere, we analyzed the

relative abundances of ASVs in each fraction and compared those frequencies between samples

(Fig. 2 and Fig. S6). In the presort cell extract and BONCAT active sorted fractions, we identified

a total of 40 bacterial classes and six archaeal classes, which were frequently dominated by a few

classes. Across all three fractions (DNA extract, presort, and sorted), ASVs affiliated with

Gammaproteobacteria were consistently the most abundant taxa detected. *Gammaproteobacteria*

had greater than 50% relative abundance within the BONCAT active community in 12 of the sorted

fraction samples (Fig. 2). In three of the six samples (1546B-1H2, 1547B-1H2, and 1545B-19F3),

all *Gammaproteobacterial* genera were detected in higher relative abundance in the sorted fraction

than in the presort fraction (Fig. 2), which indicates members of this group are active in the

Guaymas Basin deep biosphere. High abundances of *Gammaproteobacteria* in the BONCAT

active fraction likely reflect the heterotrophic nature of this class, which is capable of degrading

diverse organic substrates typically found in Guaymas Basin sediments (69–71). While the relative

frequency of *Gammaproteobacteria* was elevated in most samples, differences in relative

abundance between fractions could be explained by enrichment of *Gammaproteobacteria* during

BONCAT incubation, or methodological biases against other taxonomic groups.

Across all three fractions, we identified 61 genera affiliated with the *Gammaproteobacteria* (Fig.

S7 and Table S3) that are associated with diverse heterotrophic metabolisms. As observed for the

whole community analysis, the highest number of gammaproteobacterial genera was observed in

the DNA extract, and ASVs affiliated with the genus *Thiomicrohabdus* were amongst the most

abundant. In contrast, the presort and sorted fractions were primarily dominated by members of

Halomonas, and to a lesser extent, *Burkholderia-Caaballerionia-Paraburkholderia* and

Acinetobacter, which had the highest relative frequency in the deepest sample (1545B-19F3). In

several samples, ASVs belonging to *Halomonas* (1546B-1H2, 1547B-1H2, and 1549B-3H2) and

Acinetobacter (1545B-19F3) exhibited a relative frequency >75% in the BONCAT active fraction

in at least two replicates. This was a sharp increase compared to the other fractions which, across

all of those samples, were typically less than 20% for the DNA extract and all less than 3% for the

presort fractions. The relative abundance of *Halomonas* demonstrated the largest observed

increase between the presort and sorted fractions (Fig. S7). In particular, BONCAT active

Halomonas ASVs were detected at elevated frequencies in samples with temperatures <15°C

(Table 1), compared to warmer samples (~38°C). Sediment temperatures can fluctuate by >50°C

on relatively small temporal and spatial scales due to hydrothermal activity (72), which may create

a selective pressure on the microbial community. This indicates that some members of the

Halomonas genus remain metabolically active at lower temperatures and withstand large

temperature changes characteristic of hydrothermal sediments. Genomic analyses of these

communities revealed genes associated with central carbon metabolism, CAZymes, hydrocarbon

degradation, and nitrogen, sulfur, and arsenic cycling, suggesting that *Gammaproteobacteria* are

key drivers of organic matter remineralization and redox transformations in the subsurface (71,

73).

In addition to *Gammaproteobacteria*, abundant ASVs in the sorted fraction were affiliated with

Alphaproteobacteria throughout our samples. The relative frequency of translationally active

Alphaproteobacteria was between 28-52% for three replicates in the relatively warm (~14°C)

sediment sample from 1548B-2H3 and up to 19% in the deepest sample from 155 mbsf at the more

thermally moderate site 1545B (~39°C). After BONCAT incubation, *Alphaproteobacteria*

exhibited higher relative frequencies in the sorted fraction in 4 samples compared to the presort

fraction (Fig. 1D). Some of the supposed increase in frequency between fractions was due to low

detection of *Alphaproteobacteria* ASVs in the presort fraction. Regardless, the presence of

Alphaproteobacteria in the sorted fractions demonstrates that members of this class are active in

the Guaymas Basin deep biosphere. Similar to our observations for the *Gammaproteobacteria*,

this is consistent with previous studies that have demonstrated high abundances of

Alphaproteobacteria and associated heterotrophic activity in the marine deep biosphere (69, 74–

76).

Members of the class *Deinococci* were detected in all fractions, and consistently exhibited the

highest relative abundance in the presort fraction compared to the DNA extract and active sorted

fractions (Fig. 2 and Fig. S6). At Ringvent, in sample 1547B-1H2 (2.2 mbsf), *Deinococci* ASVs

constituted up to 54% of the presort fraction, compared to less than 1% in the DNA extract. This

higher frequency in the presort fraction suggests our cell extraction method is biased in favor of

Deinococci cells, which may influence relative abundances in the BONCAT active fraction. In

sample 1547B-8H2 (66 mbsf), members of this class accounted for up to 15% relative abundance

in the BONCAT active fraction and were less than 6% in the corresponding presort fraction. In

hydrothermally impacted habitats, *Deinococci*-affiliated sequences have been detected via 16S

rRNA gene amplicon survey (77). Additionally, analysis of 52 MAGs assigned to *Deinococci* from

IODP 360 to Atlantis Bank revealed genes associated with broad anaerobic and heterotrophic

metabolisms, including fatty acid and aromatic compound degradation (*e.g.*, acyl-CoA

dehydrogenase and benzoyl-CoA), acetogenesis, nitrogen and sulfur cycling, and metal reduction

(78). This versatility can promote *Deinococci* metabolism across the dynamic geochemical

gradients observed in the Guaymas Basin deep biosphere (Table 1).

A total of 7 classes affiliated with phylum *Firmicutes* were detected throughout our samples, and

the most abundant ASVs from this phylum were assigned to the *Bacilli* class in the presort and

active sorted fractions. Although ASVs affiliated with *Bacilli* were detected throughout the

sediment, the highest relative abundances in the BONCAT sorted fraction, up to ~20% (one

replicate was above 95%), were observed in the shallower samples (1546B-1H2 and 1547B-1H2)

(Fig. 2). In those shallower depths, above the sulfate methane transition zone, where sulfate is the

dominant terminal electron acceptor (Table 1) for anaerobic metabolisms, BONCAT-active ASVs

affiliated with *Firmicutes* may couple sulfate reduction to organic matter degradation (79).

Firmicutes isolates cultured from several marine deep biosphere sites degrade a broad suite of

organic substrates, including acetate, polysaccharides, amino acids, alcohols, nucleotides, and

microbial necromass, indicating a key role in heterotrophic carbon cycling (27, 69, 80, 81). This

metabolic capacity was substantiated by metagenomic analysis that also demonstrated high

thermal tolerance, exceeding 80°C (82). Additionally, members of the *Firmicutes* phylum within

the *Clostridia* class were associated with degradation of crude oil and polycyclic aromatic

hydrocarbon degradation in methanogenic enrichments (83). Hydrothermally induced pyrolysis of

organic matter in Guaymas Basin sediments generates petroleum hydrocarbons that are dispersed

throughout the sediment column and can fuel diverse microbial metabolisms (41, 42, 84). In the

Guaymas Basin specifically, *Firmicutes* have been linked to both sulfate and nitrate reduction (79)

and have dominated enrichment cultures amended with methanol, monomethylamine, and

ammonium (85). Given the wide metabolic capabilities of the *Firmicutes* and the high frequency

in the BONCAT active fraction, members of this phylum likely contribute significantly to organic

matter degradation and elemental cycling in Guaymas Basin sediments.

Implications for deep biosphere carbon cycling

The BONCAT active fractions were dominated by a heterotrophic microbial population capable

of metabolizing a wide range of organic carbon in the Guaymas Basin deep biosphere. To further

assess this metabolic capacity and explore potential implications for biogeochemical cycling in

hydrothermal sediments, we mapped 16S rRNA gene ASVs from the BONCAT active fraction to

annotated metagenome assembled genomes (MAGs) from IODP 385 (12). Six unique MAGs

associated with the BONCAT active fraction were identified as members of 5 bacterial phyla

(*Planctomycetota*, *Chloroflexota*, *Acidobacteriota*, *Aerophobota*, and *TA06*) and 1 archaeal

phylum (*Thermoproteota*). These MAGs were 52.7-95.9% complete and all had contamination

below 10% (Table S4).

The translationally active MAGs harbored genes associated with diverse carbon cycling pathways,

including carbohydrate degradation, C1 metabolism, and fermentation, plus nitrogen and sulfur

cycling (Fig. 3). MAG_005 (95.91% complete, 9.60% contamination) was assigned to the phylum

Planctomycetota and encodes complete pathways for glycolysis and galactose degradation, and a

nitrite reductase gene. Recently, the first anaerobic culture of *Planctomycetota* isolated from

marine sediments experimentally demonstrated functional capacity for these metabolisms (86),

validating the genomic potential of that isolate and MAG_005. MAG_054 (52.66% complete,

3.54% contamination), assigned to the class *Anaerolineae*, contained genes associated with

carbohydrate degradation and fermentation, both of which have been demonstrated by cultured

Anaerolineae (87). Similarly, NanoSIMS experiments implicated diverse *Chloroflexota* (phylum

of *Anaerolineae*) in fermentation and broad heterotrophic metabolisms in hypersaline

photoheterotrophic mats (88). In their experiments, Lee *et al*. (2014) ascertained ^{13}C-substrate

assimilation by *Chloroflexota* members using phylum-specific FISH probes, but could not resolve

taxonomic affiliations on class level or below, or implicate other phyla in carbon cycling (88). In

contrast, BONCAT-FACS workflows provide a high-throughput community-level analysis of

active taxa, and when coupled with metagenomics, support the investigation of the metabolic

potential of sorted cells (89). Taken together, BONCAT-FACS, paired with metagenomics, can

corroborate the results from culture-dependent and -independent approaches, while also providing

metabolic insights at the community level.

Most importantly, metagenomic analysis, corroborated by 16S rRNA gene identities, revealed a

diverse, translationally active microbial community in the Guaymas Basin deep biosphere with the

metabolic capacity for a range of carbon transformations and broad anaerobic metabolisms. Our

results confirm that these taxa are active, and that under the right biogeochemical conditions (*e.g.,*

substrate availability, temperature, etc.), the organisms represented by these MAGs likely utilize

these metabolisms for energy conservation in the deep biosphere. However, experimental evidence

is necessary to confirm which of these metabolisms are used *in situ* under different conditions.

Additionally, the specific ASVs affiliated with these MAGs were not necessarily amongst the most

abundant or widespread taxa in the BONCAT sorted fraction. Thus, the extent to which these

microbes drive carbon transformations is yet unknown. Future experiments that combine

BONCAT-FACS with substrate amendment (27, 32) and paired metagenomics (89) and/or

metatranscriptomics could provide deeper, community-level insights into the metabolic

capabilities of translationally active taxa in the marine deep biosphere.

Conclusion

In this study, we assessed the total, cell-extracted (presort), and translationally active (sorted)

microbial community in contrasting anoxic sediments of the Guaymas Basin deep biosphere. We

identified a diverse extant microbial community down to 154 meters below the sediment surface,

and the DNA extract contained 139 unique classes. We optimized our cell extraction and FACS

procedures for low-biomass deep biosphere samples and successfully identified 46 classes in the

presort and sorted fractions. Alpha diversity metrics decreased sharply from the DNA extract

fraction to the presort fraction, likely due to challenges associated with separating certain taxa

from the sediments, but were comparable between the presort and sorted fractions..

Gammaproteobacteria overwhelmingly dominated the BONCAT-active fraction, followed by

Alphaproteobacteria, *Firmicutes*, and *Deinococci*. These phyla have the capacity for a wide range

of heterotrophic metabolisms and drive microbial-mediated recycling of hydrothermally altered

organic matter in the Guaymas Basin deep biosphere.

We demonstrated that BONCAT-FACS can be used to assess microbial activity in low-biomass

marine sediments on a community level. Additionally, we paired this with metagenomic analysis

from IODP 385 to identify six translationally active MAGs and assess their metabolic capacity.

BONCAT-FACS coupled metagenomics links single-cell activity and genetic potential to

understand community-level dynamics in a high-throughput manner (19, 89). This approach could

be further optimized in the future to generate metagenomes directly from the sorted fraction and/or

by combining experimental amendments with whole sediment metagenomics to more tightly link

taxa with metabolic function. Importantly, our workflow, while optimized for low-biomass

samples, was likely biased against rare taxa, given the decrease in diversity between fractions.

These biases could be exacerbated if combining with metagenomics, which would require greater

quantities of biomass for FACS and sequencing compared to gene amplicon sequencing. With

careful benchmarking and optimization, BONCAT-FACS and metagenomics, when combined,

offer a powerful approach for connecting microbial taxa and metabolic function in the marine deep

biosphere and other complex environmental samples.

Acknowledgments

We would like to thank the scientific party and D/V JOIDES Resolution crew from IODP 385 for

their assistance in sample collection. Amplicon sequencing was performed by the Molecular

Research Core Facility at Idaho State University (RRID:SCR_012598). This work was supported

by NSF funding to A.M. (PRFB 2010880), R.H. (OCE-2049445), R.H. and V.E. (OCE-2046056),

V.E. and R.H. (OCE-2046799), and V.E. (OCE-1829903). Large language models (LLMs) were

used to assist in developing data visualization scripts.

Author Contributions

R.H. and V.E. conceptualized and supervised this study. V.E. collected the samples and performed

shipboard sample processing for BONCAT incubations. A.M., S.N., and C.G. developed the

methodology, performed experiments, and curated data. A.M., S.N., and Z.J. performed

bioinformatics analysis. A.M. and S.N. analyzed the data and wrote the manuscript. All authors

edited and gave critical feedback on the final manuscript. A.M., V.E., and R.H. raised funding.

A.M. and S.N. contributed to this work equally.

Data Availability

Amplicon sequence data generated for this manuscript are publicly available under NCBI

Sequence Read Archive (SRA) Bio Project accession number PRJNA1356905

(https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA1356905) and reads are available under

accession numbers SRR35956508-SRR35956585. The metagenome assembled genomes are

publicly available under NCBI SRA Bio Project accession number PRJNA909197

(https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA909197), and reads are under accession

numbers SRR22580794-SRR22580807 and SRR23614663-SRR23614677. The full

biogeochemical datasets from IODP 385, including data presented in this manuscript, are available

in the IODP expedition report (http://publications.iodp.org/proceedings/385/385title.html

References

1.   Whitman WB, Coleman DC, Wiebe WJ. 1998. Prokaryotes: The unseen majority.
Proceedings of the National Academy of Sciences 95:6578–6583.

2.   Kallmeyer J, Pockalny R, Adhikari RR, Smith DC, D'Hondt S. 2012. Global distribution
of microbial abundance and biomass in subseafloor sediment. Proceedings of the National
Academy of Sciences of the United States of America 109:16213–16216.

3.   Oremland RS, Culbertson C, Simoneit BRT. 1982. Methanogenic activity in sediment from
Leg 64, Gulf of California. Initial reports DSDP, Leg 64, Mazatlan to Long Beach, 1978-79 Part
2 759–762.

4.   D'Hondt S, Rutherford S, Spivack AJ, Hondt SD, Rutherford S, Spivack AJ. 2002.
Metabolic Activity of Subsurface Life in Deep-Sea Sediments. Science 295:2067–2070.

5.   Lever MA. 2012. Acetogenesis in the energy-starved deep biosphere-a paradox? Frontiers
in Microbiology 2:1–18.

6.   Teske A, Callaghan AV, LaRowe DE. 2014. Biosphere frontiers of subsurface life in the
sedimented hydrothermal system of Guaymas Basin. Frontiers in Microbiology 5:1–11.

7.   Santelli CM, Orcutt BN, Banning E, Bach W, Moyer CL, Sogin ML, Staudigel H, Edwards
KJ. 2008. Abundance and diversity of microbial life in ocean crust. Nature 453:653–656.

8.   Jørgensen BB. 2011. Deep subseafloor microbial cells on physiological standby.
Proceedings of the National Academy of Sciences 108:18193–18194.

9.   Lever MA, Alperin M, Engelen B, Inagaki F, Nakagawa S, Steinsbu BO, Teske A. 2006.

Trends in basalt and sediment core contamination during IODP expedition 301. Geomicrobiology

Journal 23:517–530.

10.   Orcutt BN, LaRowe DE, Biddle JF, Colwell FS, Glazer BT, Reese BK, Kirkpatrick JB,

Lapham L, Mills HJ, Sylvan JB, Wankel SD, Wheat CG. 2013. Microbial activity in the marine

deep biosphere: Progress and prospects. Frontiers in Microbiology 4:1–15.

11.   Gittins DA, Desiage P-A, Morrison N, Rattray JE, Bhatnagar S, Chakraborty A, Zorz J, Li

C, Horanszky O, Cramm MA, Bisiach F, Bennett R, Webb J, MacDonald A, Fowler M, Campbell

DC, Hubert CRJ. 2022. Geological processes mediate a microbial dispersal loop in the deep

biosphere. Science Advances 8:eabn3485.

12.   Mara P, Geller-McGrath D, Edgcomb V, Beaudoin D, Morono Y, Teske A. 2023.

Metagenomic profiles of archaea and bacteria within thermal and geochemical gradients of the

Guaymas Basin deep subsurface. Nat Commun 14:7768.

13.   Sun J, Hirai M, Takaki Y, Evans PN, Nunoura T, Rinke C. 2025. Metagenomic insights

into taxonomic and functional patterns in shallow coastal and deep subseafloor sediments in the

Western Pacific. Microbial Genomics 11:001351.

14.   Orsi WD, Edgcomb VP, Christman GD, Biddle JF. 2013. Gene expression in the deep

biosphere. Nature 499:205–208.

15.   Coskun ÖK, Pichler M, Vargas S, Gilder S, Orsi WD. 2018. Linking Uncultivated

Microbial Populations and Benthic Carbon Turnover by Using Quantitative Stable Isotope

Probing. Applied and Environmental Microbiology 84.

16.   Coskun ÖK, Özen V, Wankel SD, Orsi WD. 2019. Quantifying population-specific growth

in benthic bacterial communities under low oxygen using H218O. The ISME Journal 13.

17.   Mara P, Zhou Y-L, Teske A, Morono Y, Beaudoin D, Edgcomb V. 2023. Microbial gene

expression in Guaymas Basin subsurface sediments responds to hydrothermal stress and energy

limitation. 11. ISME J 17:1907–1919.

18.   Mara P, Beaudoin D, Aiello I, Morono Y, Geller-McGrath D, Edgcomb VP, Teske A.

2024. Deep subseafloor sediments in Guaymas Basin harbor cosmopolitan microbiota and traces

of hydrothermal populations. Commun Earth Environ 5:505.

19.   Hatzenpichler R, Krukenberg V, Spietz RL, Jay ZJ. 2020. Next-generation physiology

approaches to study microbiome function at single cell level. Nature Reviews Microbiology

18:241–256.

20.   Hatzenpichler R, Scheller S, Tavormina PL, Babin BM, Tirrell DA, Orphan VJ. 2014. In

situ visualization of newly synthesized proteins in environmental microbes using amino acid

tagging and click chemistry. Environmental Microbiology 16:2568–2590.

21.   Hatzenpichler R, Connon SA, Goudeau D, Malmstrom RR, Woyke T, Orphan VJ. 2016.

Visualizing in situ translational activity for identifying and sorting slow-growing archaeal -

bacterial consortia. Proceedings of the National Academy of Sciences 113:E4069–E4078.

22.   Samo TJ, Smriga S, Malfatti F, Sherwood BP, Azam F. 2014. Broad distribution and high

proportion of protein synthesis active marine bacteria revealed by click chemistry at the single cell

level. Frontiers in Marine Science 1:1–18.

23.   Leizeaga A, Estrany M, Forn I, Sebastián M. 2017. Using click-chemistry for visualizing

in situ changes of translational activity in planktonic marine bacteria. Frontiers in Microbiology

8:1–11.

24.   Couradeau E, Sasse J, Goudeau D, Nath N, Hazen TC, Bowen BP, Chakraborty R,

Malmstrom RR, Northen TR. 2019. Probing the active fraction of soil microbiomes using

BONCAT-FACS. Nature Communications 10.

25.   Kiick KL, Saxon E, Tirrell DA, Bertozzi CR. 2002. Incorporation of azides into

recombinant proteins for chemoselective modification by the Staudinger ligation. Proceedings of

the National Academy of Sciences 99:19–24.

26.   Mulay SA, Vishnivetskaya TA, Hochanadel LH, Klingeman DM, Lloyd KG, Pelletier DA,

Podar M. 2025. BONCAT-Live for isolation and cultivation of active environmental bacteria.

mBio.

27.   Krukenberg V, Reichart NJ, Spietz RL, Hatzenpichler R. 2021. Microbial Community

Response to Polysaccharide Amendment in Anoxic Hydrothermal Sediments of the Guaymas

Basin. Frontiers in Microbiology 12.

28.   Harris JE, Bledsoe RB, Guha S, Omari H, Crandall SG, Burghardt LT, Couradeau E. 2025.

The activity of soil microbial taxa in the rhizosphere predicts the success of root colonization.

mSystems 0:e00458-25.

29.   Riva A, Kolimár D, Spittler A, Wisgrill L, Herbold CW, Abrankó L, Berry D. 2020.

Conversion of Rutin, a Prevalent Dietary Flavonol, by the Human Gut Microbiota. Front Microbiol

11.

30.   Valentini TD, Lucas SK, Binder KA, Cameron LC, Motl JA, Dunitz JM, Hunter RC. 2020.

Bioorthogonal non-canonical amino acid tagging reveals translationally active subpopulations of

the cystic fibrosis lung microbiota. Nat Commun 11:2287.

31.   Taguer M, Shapiro BJ, Maurice CF. 2021. Translational activity is uncoupled from nucleic

acid content in bacterial cells of the human gut microbiota. Gut Microbes 13:1903289.

32.   Reichart NJ, Jay ZJ, Krukenberg V, Parker AE, Spietz RL, Hatzenpichler R. 2020.

Activity-based cell sorting reveals responses of uncultured archaea and bacteria to substrate

amendment. The ISME Journal 14:2851–2861.

33.   Chen L, Zhao B, Li X, Cheng Z, Wu R, Xia Y. 2021. Isolating and characterizing

translationally active fraction of anammox microbiota using bioorthogonal non-canonical amino

acid tagging. Chemical Engineering Journal 418:129411.

34.   Du Z, Behrens SF. 2021. Tracking *de novo* protein synthesis in the activated sludge

microbiome using BONCAT-FACS. Water Research 205:117696.

35.   Trexler RV, Van Goethem MW, Goudeau D, Nath N, Malmstrom RR, Northen TR,

Couradeau E. 2023. BONCAT-FACS-Seq reveals the active fraction of a biocrust community

undergoing a wet-up event. Front Microbiol 14.

36.   Calvert SE. 1966. Origin of Diatom-Rich, Varved Sediments from the Gulf of California.

The Journal of Geology 74:546–565.

37.   Einsele G, Gieskes JM, Curray J, Moore DM, Aguayo E, Aubry M-P, Fornari D, Guerrero

J, Kastner M, Kelts K, Lyle M, Matoba Y, Molina-Cruz A, Niemitz J, Rueda J, Saunders A,

Schrader H, Simoneit B, Vacquier V. 1980. Intrusion of basaltic sills into highly porous sediments,

and resulting hydrothermal activity. Nature 283:441–445.

38.   Lonsdale P, Becker K. 1985. Hydrothermal plumes, hot springs, and conductive heat flow

in the Southern Trough of Guaymas Basin. Earth and Planetary Science Letters 73:211–225.

39.   Lin YS, Koch BP, Feseker T, Ziervogel K, Goldhammer T, Schmidt F, Witt M, Kellermann

MY, Zabel M, Teske A, Hinrichs KU. 2017. Near-surface heating of young rift sediment causes

mass production and discharge of reactive dissolved organic matter. Scientific Reports 7:1–10.

40.   Simoneit BRT, Mazurek MA, Brenner S, Crisp PT, Kaplan IR. 1979. Organic

geochemistry of recent sediments from Guaymas Basin, Gulf of California. Deep Sea Research

Part A, Oceanographic Research Papers 26:879–891.

41.   Kawka OE, Simoneit BRT. 1987. Survey of hydrothermally-generated petroleums from

the Guaymas Basin spreading center. Organic Geochemistry 11:311–328.

42.   Bazylinski DA, Farrington JW, Jannasch HW. 1988. Hydrocarbons in surface sediments

from a Guaymas Basin hydrothermal vent site. Organic Geochemistry 12:547–558.

43.   Whelan JK, Simoneit BRT, Tarafa ME. 1988. C1-C8 hydrocarbons in sediments from

Guaymas Basin, Gulf of California-Comparison to Peru Margin, Japan Trench and California

Borderlands. Organic Geochemistry 12:171–194.

44.   Martens CS. 1990. Generation of short chain acid anions in hydrothermally altered

sediments of the Guaymas Basin, Gulf of California. Applied Geochemistry 5:71–76.

45.   Zhuang G, Montgomery A, Samarkin VA, Song M, Liu J, Schubotz F, Teske A, Hinrichs

K-U, Joye SB. 2019. Generation and Utilization of Volatile Fatty Acids and Alcohols in

Hydrothermally Altered Sediments in the Guaymas Basin, Gulf of California. Geophysical

Research Letters 46:2637–2646.

46.   Teske A, Lizarralde D, Höfig TW, Aiello IW, Ash JL, Bojanova DP, Buatier MD,

Edgcomb VP, Galerne CY. 2021. "Expedition 385 Summary" in: Guaymas Basin Tectonics and

Biosphere. International Ocean Discovery Program.

47.   Kallmeyer J, Smith DC, Spivack AJ, D'Hondt S. 2008. New cell extraction procedure

applied to deep subsurface sediments. Limnology and Oceanography: Methods 6:236–245.

48.   Hatzenpichler R, Orphan VJ. 2015. Detection of Protein-Synthesizing Microorganisms in

the Environment via Bioorthogonal Noncanonical Amino Acid Tagging (BONCAT).

Hydrocarbon and Lipid Microbiology Protocols Vol 7: Single-Cell and Single-Molecule Methods

171–195.

49.   Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, Prill RJ, Tripathi

A, Gibbons SM, Ackermann G, Navas-Molina JA, Janssen S, Kopylova E, Vázquez-Baeza Y,

González A, Morton JT, Mirarab S, Zech Xu Z, Jiang L, Haroon MF, Kanbar J, Zhu Q, Jin Song

S, Kosciolek T, Bokulich NA, Lefler J, Brislawn CJ, Humphrey G, Owens SM, Hampton-Marcell

J, Berg-Lyons D, McKenzie V, Fierer N, Fuhrman JA, Clauset A, Stevens RL, Shade A, Pollard

KS, Goodwin KD, Jansson JK, Gilbert JA, Knight R. 2017. A communal catalogue reveals Earth's

multiscale microbial diversity. Nature 551:457–463.

50.   Parada AE, Needham DM, Fuhrman JA. 2016. Every base matters: Assessing small subunit

rRNA primers for marine microbiomes with mock communities, time series and global field

samples. Environmental Microbiology 18:1403–1414.

51.   Apprill A, McNally S, Parsons R, Weber L. 2015. Minor revision to V4 region SSU rRNA

806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquatic Microbial

Ecology 75:129–137.

52.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H,

Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown

CT, Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da Silva R, Diener C, Dorrestein

PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz

JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H,

Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR,

Keim P, Kelley ST, Knights D, Koester I, Kosciolek T, Kreps J, Langille MGI, Lee J, Ley R, Liu

Y-X, Loftfield E, Lozupone C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik

AV, Metcalf JL, Morgan SC, Morton JT, Naimey AT, Navas-Molina JA, Nothias LF, Orchanian

SB, Pearson T, Peoples SL, Petras D, Preuss ML, Pruesse E, Rasmussen LB, Rivers A, Robeson

MS, Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR, Swafford AD,

Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, van der Hooft JJJ,

Vargas F, Vázquez-Baeza Y, Vogtmann E, von Hippel M, Walters W, Wan Y, Wang M, Warren

J, Weber KC, Williamson CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R,

Caporaso JG. 2019. Reproducible, interactive, scalable and extensible microbiome data science

using QIIME 2. Nat Biotechnol 37:852–857.

53.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.

2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based

tools. Nucleic Acids Res 41:D590-596.

54.   Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. 2018. Simple statistical

identification and removal of contaminant sequences in marker-gene and metagenomics data.

Microbiome 6:226.

55.   McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. PLOS ONE 8:e61217.

56.   Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. 2020. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. Bioinformatics 36:1925–1927.

57.   Chklovski A, Parks DH, Woodcroft BJ, Tyson GW. 2023. CheckM2: a rapid, scalable and

accurate tool for assessing microbial genome quality using machine learning. Nat Methods 1–10.

58.   Dombrowski N, Seitz KW, Teske A, Baker BJ. 2017. Genomic insights into potential

interdependencies in microbial hydrocarbon and nutrient cycling in hydrothermal sediments.

Microbiome 5:106.

59.   Teske AP, Lizarralde D, Höfig TW, Aiello IW, Ash JL, Bojanova DP, Buatier MD,

Edgcomb VP, Galerne CY, Gontharet S, Heuer VB, Jiang S, Kars MAC, Singh SK, Kim J-H,

Koornneef LMT, Marsaglia KM, Meyer N, Morono Y, Negrete-Aranda R, Neumann F, Pastor LC,

Peña-Salinas ME, Pérez-Cruz LL, Ran L, Riboulleau A, Sarao JA, Schubert F, Stock JM, Toffin

LMAA, Xie W, Yamanaka T, Zhuang G. 2021. Site U1545. In Teske, A., Lizarralde, D., Höfig,

T. W., and the Expedition 385 Scientists, Guaymas Basin Tectonics and Biosphere. Proceedings

of the International Ocean Discovery Program 385.

60.   Teske AP, Lizarralde D, Höfig TW, Aiello IW, Ash JL, Bojanova DP, Buatier MD,

Edgcomb VP, Galerne CY, Gontharet S, Heuer VB, Jiang S, Kars MAC, Singh SK, Kim J-H,

Koornneef LMT, Marsaglia KM, Meyer N, Morono Y, Negrete-Aranda R, Neumann F, Pastor LC,

Peña-Salinas ME, Pérez-Cruz LL, Ran L, Riboulleau A, Sarao JA, Schubert F, Stock JM, Toffin

LMAA, Xie W, Yamanaka T, Zhuang G. 2021. Site U1546. In Teske, A., Lizarralde, D., Höfig,

T. W., and the Expedition 385 Scientists, Guaymas Basin Tectonics and Biosphere. Proceedings

of the International Ocean Discovery Program 385.

61.   Teske AP, Lizarralde D, Höfig TW, Aiello IW, Ash JL, Bojanova DP, Buatier MD,

Edgcomb VP, Galerne CY, Gontharet S, Heuer VB, Jiang S, Kars MAC, Singh SK, Kim J-H,

Koornneef LMT, Marsaglia KM, Meyer N, Morono Y, Negrete-Aranda R, Neumann F, Pastor LC,

Peña-Salinas ME, Pérez-Cruz LL, Ran L, Riboulleau A, Sarao JA, Schubert F, Stock JM, Toffin

LMAA, Xie W, Yamanaka T, Zhuang G. 2021. Sites U1547 and U1548. In Teske, A., Lizarralde,

D., Höfig, T. W., and the Expedition 385 Scientists, Guaymas Basin Tectonics and Biosphere.

Proceedings of the International Ocean Discovery Program 385.

62.   Teske AP, Lizarralde D, Höfig TW, Aiello IW, Ash JL, Bojanova DP, Buatier MD,

Edgcomb VP, Galerne CY, Gontharet S, Heuer VB, Jiang S, Kars MAC, Singh SK, Kim J-H,

Koornneef LMT, Marsaglia KM, Meyer N, Morono Y, Negrete-Aranda R, Neumann F, Pastor LC,

Peña-Salinas ME, Pérez-Cruz LL, Ran L, Riboulleau A, Sarao JA, Schubert F, Stock JM, Toffin

LMAA, Xie W, Yamanaka T, Zhuang G. 2021. Site U1549. In Teske, A., Lizarralde, D., Höfig,

T. W., and the Expedition 385 Scientists, Guaymas Basin Tectonics and Biosphere. Proceedings

of the International Ocean Discovery Program 385.

63.   Keeler E, Burgaud G, Teske A, Beaudoin D, Mehiri M, Dayras M, Cassand J, Edgcomb

V. 2021. Deep-sea hydrothermal vent sediments reveal diverse fungi with antibacterial activities.

FEMS Microbiology Ecology 97:fiab103.

64.   Taylor DL, Walters WA, Lennon NJ, Bochicchio J, Krohn A, Caporaso JG, Pennanen T.

2016. Accurate Estimation of Fungal Diversity and Abundance through Improved Lineage-

Specific Primers Optimized for Illumina Amplicon Sequencing. Applied and Environmental

Microbiology 82:7217–7226.

65.   Tedersoo L, Anslan S, Bahram M, Põlme S, Riit T, Liiv I, Kõljalg U, Kisand V, Nilsson

H, Hildebrand F, Bork P, Abarenkov K. 2015. Shotgun metagenomes and multiple primer pair-

barcode combinations of amplicons reveal biases in metabarcoding analyses of fungi. MycoKeys

10:1–43.

66.   Inagaki F, Hinrichs KU, Kubo Y, Bowles MW, Heuer VB, Hong WL,

Hoshino T, Ijiri A, Imachi H, Ito M, Kaneko M, Lever MA, Lin YS, Methé BA,

Morita S, Morono Y, Tanikawa W, Bihan M, Bowden SA, Elvert M, Glombitza

C, Gross D, Harrington GJ, Hori T, Li K, Limmer D, Liu CH, Murayama M,

Ohkouchi N, Ono S, Park YS, Phillips SC, Prieto-Mollar X, Purkey M,

Riedinger N, Sanada Y, Sauvage J, Snyder G, Susilawati R, Takano Y,

Tasumi E, Terada T, Tomaru H, Trembath-Reichert E, Wang DT, Yamada Y.

2015. Exploring deep microbial life in coal−bearing sediment down to

~2.5 km below the ocean floor. Science 349:420−424.

67.   D'Hondt S, Pockalny R, Spivack AJ, Fulfer VM. 2019. Subseafloor life and its

biogeochemical impacts. Nature Communications 10:1–13.

68.   Morono Y, Terada T, Kallmeyer J, Inagaki F. 2013. An improved cell separation technique

for marine subsurface sediments: applications for high-throughput analysis using flow cytometry

and cell sorting. Environmental Microbiology 15:2841–2849.

69.   Batzke A, Engelen B, Sass H, Cypionka H. 2007. Phylogenetic and physiological diversity

of cultured deep-biosphere bacteria from Equatorial Pacific Ocean and Peru Margin sediments.

Geomicrobiology Journal 24:261–273.

70.   Biddle JF, Sylvan JB, Brazelton WJ, Tully BJ, Edwards KJ, Moyer CL, Heidelberg JF,

Nelson WC. 2012. Prospects for the study of evolution in the deep biosphere. Frontiers in

Microbiology 2:1–7.

71.   McKay L, Klokman VW, Mendlovitz HP, Larowe DE, Hoer DR, Albert D, Amend JP,

Teske A. 2016. Thermal and geochemical influences on microbial biogeography in the

hydrothermal sediments of Guaymas Basin, Gulf of California. Environmental Microbiology

Reports 8:150–161.

72.   Teske A, De Beer D, McKay L, Tivey MK, Biddle JF, Hoer D, Lloyd KG, Lever MA, Røy

H, Albert DB, Mendlovitz HP, MacGregor BJ. 2016. The Guaymas Basin hiking guide to

hydrothermal mounds, chimneys, and microbial mats: Complex seafloor expressions of subsurface

hydrothermal circulation. Frontiers in Microbiology 7.

73.   D'Angelo T, Goordial J, Poulton NJ, Seyler L, Huber JA, Stepanauskas R, Orcutt BN.

2022. Oceanic Crustal Fluid Single Cell Genomics Complements Metagenomic and

Metatranscriptomic Surveys With Orders of Magnitude Less Sample Volume. Frontiers in

Microbiology 12:1–13.

74.   Orcutt BN, Sylvan JB, Knab NJ, Edwards KJ. 2011. Microbial Ecology of the Dark Ocean

above, at, and below the Seafloor. Microbiology and Molecular Biology Reviews 75:361–422.

75.   Meier DV, Bach W, Girguis PR, Gruber-Vodicka HR, Reeves EP, Richter M, Vidoudez

C, Amann R, Meyerdierks A. 2016. Heterotrophic Proteobacteria in the vicinity of diffuse

hydrothermal venting. Environmental Microbiology 18:4348–4368.

76.   Dong X, Rattray JE, Campbell DC, Webb J, Chakraborty A, Adebayo O, Matthews S, Li

C, Fowler M, Morrison NM, MacDonald A, Groves RA, Lewis IA, Wang SH, Mayumi D,

Greening C, Hubert CRJ. 2020. Thermogenic hydrocarbon biodegradation by diverse depth-

stratified microbial populations at a Scotian Basin cold seep. Nat Commun 11:5825.

77.   Wee SY, Edgcomb VP, Beaudoin D, Yvon-Lewis S, Sylvan JB. 2021. Microbial

Abundance and Diversity in Subsurface Lower Oceanic Crust at Atlantis Bank, Southwest Indian

Ridge. Applied and Environmental Microbiology 87.

78.   Zhou Z, St. John E, Anantharaman K, Reysenbach A-L. 2022. Global patterns of diversity

and metabolism of microbial communities in deep-sea hydrothermal vent deposits. Microbiome

10:241.

79.   Callac N, Rommevaux-Jestin C, Rouxel O, Lesongeur F, Liorzou C, Bollinger C, Ferrant

A, Godfroy A. 2013. Microbial colonization of basaltic glasses in hydrothermal organic-rich

sediments at Guaymas Basin. Frontiers in Microbiology 4.

80.   L'Haridon S, Miroshnichenko ML, Kostrikina NA, Tindall BJ, Spring S, Schumann P,

Stackebrandt E, Bonch-Osmolovskaya EA, Jeanthon C. 2006. Vulcanibacillus modesticaldus gen.

nov., sp. nov., a strictly anaerobic, nitrate-reducing bacterium from deep-sea hydrothermal vents.

International Journal of Systematic and Evolutionary Microbiology 56:1047–1053.

81.   Guo L-Y, Ling S-K, Li C-M, Chen G-J, Du Z-J. 2018. Bacillus marinisedimentorum sp.

nov., isolated from marine sediment. Int J Syst Evol Microbiol 68:198–203.

82.   Bell E, Rattray JE, Sloan K, Sherry A, Pilloni G, Hubert CRJ. 2022. Hyperthermophilic

endospores germinate and metabolize organic carbon in sediments heated to 80°C. Environmental

Microbiology n/a.

83.   Berdugo-Clavijo C, Dong X, Soh J, Sensen CW, Gieg LM. 2012. Methanogenic

biodegradation of two-ringed polycyclic aromatic hydrocarbons. FEMS Microbiology Ecology

81:124–133.

84.   Edgcomb VP, Teske AP, Mara P. 2022. Microbial Hydrocarbon Degradation in Guaymas

Basin—Exploring the Roles and Potential Interactions of Fungi and Sulfate-Reducing Bacteria.

Front Microbiol 13.

85.   Trembath-Reichert E, Morono Y, Ijiri A, Hoshino T, Dawson KS, Inagaki F, Orphan VJ.

2017. Methyl-compound use and slow growth characterize microbial life in 2-km-deep subseafloor

coal and shale beds. Proceedings of the National Academy of Sciences 114:E9206–E9215.

86.   Zheng R, Wang C, Liu R, Cai R, Sun C. 2024. Physiological and metabolic insights into

the first cultured anaerobic representative of deep-sea Planctomycetes bacteria. eLife 12:RP89874.

87.   Sekiguchi Y, Yamada T, Hanada S, Ohashi A, Harada H, Kamagata Y. 2003. Anaerolinea

thermophila gen. nov., sp. nov. and Caldilinea aerophila gen. nov., sp. nov., novel filamentous

thermophiles that represent a previously uncultured lineage of the domain Bacteria at the

subphylum level. International Journal of Systematic and Evolutionary Microbiology 53:1843–

1851.

88.   Lee JZ, Burow LC, Woebken D, Everroad RC, Kubo MD, Spormann AM, Weber PK, Pett-

Ridge J, Bebout BM, Hoehler TM. 2014. Fermentation couples Chloroflexi and sulfate-reducing

bacteria to Cyanobacteria in hypersaline microbial mats. Front Microbiol 5.

89.   Schweitzer HD, Smith HJ, Barnhart EP, McKay LJ, Gerlach R, Cunningham AB,

Malmstrom RR, Goudeau D, Fields MW. 2022. Subsurface hydrocarbon degradation strategies in

low- and high-sulfate coal seam communities identified with activity-based metagenomics. npj

Biofilms Microbiomes 8:7.

Table 1. Depth and temperature data for BONCAT incubations. BD = below detection. Geochemistry data are compiled from IODP 385 site chapters as published by Mara *et al.* (2023). Sediment temperatures are based on linear interpolation from published temperature gradients

Hole	Sample ID	Depth (mbsf)	Temp (°C)	Alkalinity (mM)	SO_4^{2-} (mM)	PO_4^{3-} (µM)	H_2S (mM)	NH_4^+ (mM)	CH_4 (mM)	CO (nM)	DOC (mg/L)	DIC (mM)	TOC (wt%)	TN (wt%)	TOC/TN
U1545B	1545B-19F3	155.0	39.8	35.3	0.3	46.9	3.2	15.9	1.9	116	48.2	10.7	2.17	0.34	7.5
U1546B	1546B-1H2	0.8	2.8	5.1	27.7	12.4	428	0.7	BD	665	21.8	2.2	4.36	0.52	9.7
U1547B	1547B-1H2	2.2	14.2	3	27.9	16.5	68	0.1	BD	350	13.9	1.2	3.76	0.45	9.7
U1547B	1547B-8H2	65.8	46.6	13	20.2	25.4	6606	2.9	BD	91	23.3	1.5	1.25	0.15	9.5
U1548B	1548B-2H3	8.9	13.7	4.8	26.8	24.5	732.5	0.3	BD	554	13.1	1.4	2.4	0.22	12.9
U1549B	1549B-3H2	16.5	6.4	18.7	17.5	98.1	2567	3.3	BD	188	23.6	5.7	3.12	0.59	6.1

Figure 1. Alpha diversity indices of each sample and fold enrichment changes of the 4 most abundant Classes. Plots of Observed Diversity (A) and Shannon's diversity index (B) for all sites. Fold enrichment of relative abundance between the pre-sort and sorted fractions for Gammaproteobacteria (C), Alphaproteobacteria (D), Bacilli (E), and Deinococci (F). Fold enrichment was calculated as the ratio of relative abundances of ASVs affiliated with a given class between the pre-sort and sorted fraction, normalized to zero so that equal frequencies equate to zero. In fold-enrichment plots (C-F), yellow bars indicate that a class was more represented in the sorted fraction, and blue bars show that the class was more abundant in the pre-sort fraction. Bars represent the average of triplicate samples, and each point shows the value of each replicate. A * indicated that no ASVs affiliated with that class were detected in either the presort or sorted fraction.

Figure 2. Community composition of cell extract (pre-sort) and BONCAT-positive sorted samples depicted as the relative abundance of ASVs of the 16S rRNA gene collapsed to class level. Fractions are grouped separately on the y-axis. Replicate numbers are indicated by 1-3; C refers to the no-HPG control

Figure 3. Heat map of metabolic potential of the six translationally active MAGs. A subset of genes was assessed. Squares indicate the presence of a functional gene (blue), complete metabolic pathway (red), or partial metabolic pathway (pink). White squares indicate that no gene(s) affiliated with that pathway were detected. MAGs are numbered following identities in (Mara et al., 2023). Abbreviations: PPP (pentose phosphate pathway), ED pathway (Entner-Doudoroff pathway). MAG_012 (84.65% complete, 9.60% contamination), MAG_005 (95.91% complete, 7.52% contamination), MAG_022 (87.01% complete, 4.23% contamination), MAG_054 (52.66% complete, 3.54% contamination), MAG_013 (86.40% complete, 0.78% contamination), MAG_014 (88.80% complete, 8.25% contamination).

Figure S1. Representative FACS plots for 1545B-19F3 No HPG control (A) and 1545B-19F3 Replicate A (B). No-HPG control samples were used to draw the positive (Cy5; purple polygon) and negative (Neg; black polygon) gates for each sample set. The Cy5 positive gate represents BONCAT positive events and is sorted into collection tubes for downstream analysis. The percent values represent the fraction of the total events in each gate.

DAPI (DNA) **BONCAT (new proteins)**

Figure S2. Fluorescence microscopy images of IODP 385 BONCAT samples. Images from 1545B-19F3 Replicate A pre-sort (A) and sorted (B), 1548B-3H2 Replicate B pre-sort (C) and sorted (D), and 1548B-3H2 no HPG control pre-sort (E). All cells are stained with the general nucleic acid stain 4′,6-diamidino-2-phenylindole (DAPI) in blue. Cells that incorporated HPG were stained with Cy5 using click chemistry and are depicted as red. Note that the pre-sort samples (A and C) contain cells with and without HPG incorporation, while sorted samples only contain cells that had incorporated HPG (red). Some sediment leftover from cell extraction appears blue, either from autofluorescence or non-specific DAPI staining. Scale bars depicted on each image are 10 μm.

Figure S3. Fluorescence microscopy image of fungal isolates showing successful BONCAT labeling. *Rhodotorula sp.* (A) and *Cadophora sp.* (B) were grown with HPG in liquid media, preserved with GlyTE, and subjected to click chemistry to identify HPG incorporation. The red stain indicates BONCAT-positive cells labelled with Cy5, and the blue stain represents the general DNA stain DAPI. The scale bar is 10 μm.

Figure S4. Alpha diversity indices of each sample. Observed diversity (A) and Shannon's diversity (B) for DNA extract, cell extracted pre-sort, and BONCAT-sorted fractions from all samples.

Figure S5. Sequence processing for 16S rRNA gene amplicons. Samples are grouped by fraction (DNA extract, cell extracted fraction/presort, sorted, and controls) and depth. The size and color of points reflect the number of reads at each processing step. Cleaned = number of reads after processing and removal of contaminant sequences. Control replicates for presort and sort samples are the no-HPG controls. Extraction blanks (EB) were extracted alongside samples and then treated as presort samples. Sorted control replicates, extraction blanks, and PCR blanks were used to assess contaminants and were removed from the dataset prior to community analysis.

Figure S6. Relative abundance of ASVs from 16S rRNA gene amplicon sequencing from whole sediment DNA extracts. Replicates are denoted Rep1-3.

Figure S7. Relative abundance of genus-level ASVs, affiliated with the Gammaproteobacteria, from 16S rRNA gene amplicon sequencing for each fraction. Fractions are faceted on the y-axis for the DNA extract (DNA), cell extract (presort), and BONCAT positive fractions (sort). Replicate numbers are 1-3; C refers to the no-HPG control.

Table S1. Sorting metrics from FACS for each sample. The number of events sorted for each sample and replicate is listed. The percent of total events is the number of positive, sorted events out of the total number of events detected by FACS.

Sample	Replicate	Sorted BONCAT+ events	Percent of total events
1546B-1H2	1	97991	0.69%
1546B-1H2	2	208025	1.31%
1546B-1H2	3	119443	0.58%
1545B-19F3	1	136468	0.63%
1545B-19F3	2	135643	0.59%
1545B-19F3	3	143970	0.57%
1547B-1H2	1	132251	0.55%
1547B-1H2	2	266329	0.77%
1547B-1H2	3	398515	0.91%
1549B-3H2	1	723234	6.72%
1549B-3H2	2	8567	0.06%
1549B-3H2	3	7786	0.06%
1548B-3H2	1	54497	0.28%
1548B-3H2	2	33976	0.18%
1548B-3H2	3	38075	0.15%
1547B-8H2	1	12239	0.06%
1547B-8H2	2	6959	0.06%
1547B-8H2	3	8718	0.08%