Previous legume identity influence wheat rhizosphere microbial

communities and grain protein content

Emmy L'Espérance[1], Vincent Poirier[2], Stéphanie Lavergne[2], Étienne Yergeau[1]*

[1]Centre Armand-Frappier Santé Biotechnologie, Institut national de la recherche

scientifique, 531 boulevard des Prairies, Laval, QC, H7V 1B7, Canada.

[2]Institut de recherche en agriculture et agroalimentaire, Université du Québec en Abitibi-

Témiscamingue, 79 rue Côté, Notre-Dame-du-Nord, QC, J0Z 3B0, Canada.

*Correspondance to: Etienne Yergeau, Centre Armand-Frappier Santé Biotechnologie,

Institut national de la recherche scientifique, 531 boulevard des Prairies, Laval, QC, H7V

1B7, Canada. Tel.: +1 450-687-5010 ext. 8881, etienne.yergeau@inrs.ca

Abstract

In crop rotation systems, plant-soil feedback (PSF) effects can modulate nutrient cycling

in soil, like organic carbon (C) and nitrogen (N) cycling. Organic nitrogen is present in plant

and microbial necromass, and it must be depolymerized by microbes to become available

for crops. Since plant identity can modulate microbial diversity and community

composition, we thought that previous crop identity and its residue management would

also change the microbial functional capacity, and thereby impact soil N availability, and

the quality and yield of the following crop. To test this, two legumes (*Vicia faba* L., i.e.

faba bean, and *Pisum sativum* L., i.e., yellow pea) were grown in two fields (Cloutier, and

Palmarolle) in Abitibi-Témiscamingue, Québec, Canada (n=3 for each field). At the end of

the growing season, we harvested peas and faba beans in both fields. During the following

growing season, spring wheat (*Triticum aestivum* L.) was sown and received, or not

granulated chicken manure as an organic fertilizer. We determined the diversity and

composition of the microbial communities and their enzymatic depolymerization capacity

in the soil and the rhizosphere each growing season. During wheat growth, previous

legumes shaped bacterial (p-value = 0.006) and fungal (p-value = 0.001) communities

without modulating the enzymatic activity of wheat-associated rhizosphere microbes.

However, faba bean as a previous crop increased soil ammonium and wheat grain protein

content at harvest as compared to peas at Cloutier. Altogether, our results show that faba

beans can enhance wheat N nutrition, without a concomitant increase in potential protein

or cellulose depolymerization, suggesting more mineralization due to increases in fungal:

bacterial ratio or in the availability of substrates. Understanding plant-soil feedback in

crop rotation systems is crucial to improve our practices and sustainably meet crops'

nutritional needs.

Key words

Plant-Soil feedback, Microbial communities, Rhizosphere, Depolymerization, crop

rotation, Grain quality.

Highlights

- Previous crop identity distinctively shaped wheat rhizosphere microbial

communities

- Previous legume did not altered the enzymatic activity of wheat-associated

rhizosphere microbes, but increased soil ammonium

- Wheat grain protein content was higher following faba bean than peas

1. Introduction

Agricultural practices have a significant impact on belowground microbial communities, altering their functionality, community structure, and diversity (Calderon et al., 2001; Li et al., 2020; Zhao et al., 2020). Monocropping often leads to lower soil microbial diversity and decreased plant growth and yield compared to crop rotation systems (Bak et al., 2022; Adebayo et al., 2025). Crop rotation systems (i.e., growing different crop species sequentially on the same land) enhance soil nutrient content and nutrient cycling (Wang et al., 2023c), limit the establishment of plant pathogens (Zhou et al., 2023) and enhance soil health and microbial diversity (Yang et al., 2024). Crop rotation systems are therefore fundamental in organic farming to maximize yield without the use of synthetic chemical inputs (Gamage et al., 2023).

The benefits of crop rotation systems can be explained by positive plant-soil feedback (PSF). The previous crop modulates soil microbial communities and physicochemical properties, impacting the next crop (Mariotte et al., 2018; Wang et al., 2025). Indeed, plants release a myriad of compounds, shaping spatially and temporally distinct microbial communities (Zhalnina et al., 2018). Positive plant-soil feedback includes pathogen reduction and beneficial microbes' enrichment (Ma et al., 2017; Hannula et al., 2021). *Arabidopsis thaliana* changes its exudation pattern in response to the pathogenic bacterium *P. syringae* (Yuan et al., 2018). After six passages in the same soil, these changes created soil microbial communities antagonist to the pathogen, resulting in healthy plants (Yuan et al., 2018). Due to highly diversified plant traits, functions and

exudation patterns, PSF effects are species-specific, meaning that we cannot associate a

PSF effect with a plant family (Hannula et al., 2020). Plant-soil feedbacks are also not

limited to the original plant species and can affect other plants grown in the same soil,

creating multiple possibilities for crop rotation systems (Murrell et al., 2020). For instance,

winter wheat (*Triticum aestivum* L.) grown continuously had a lower yield compared to

winter wheat growing after oilseed rape, because oilseed rape changed the microbial

communities and the abundances of functional genes related to inorganic N-cycling

(Kaloterakis et al., 2025). This plant-specific rhizospheric effect can persist for many

growing seasons (Bay et al., 2021).

Besides changing the microbial community structure and abundance of specific genes,

crops also affect more generally the carbon (C) and nitrogen (N) cycles (Knops et al.,

2002). Plant-soil feedback effects on nutrient cycling can be attributed to both living and

dead crops. Plant litter origin (above vs belowground) and chemistry mediate the

decomposition rate (Freschet et al., 2013), and microbial decomposer abundance and

diversity over time (Bhatnagar et al., 2018). A classic use of PSF in agricultural settings is

legume-cereal rotations. Legumes are well known to enhance soil N through biological

fixation, and using them in crop rotation systems replenishes soil N without using

chemical fertilizers (Gan et al., 2015). But not all legumes are identical. For instance, faba

bean (*Vicia faba* L.) fixed 185 kg N ha^{-1} and forage peas (*Pisum sativum* L.) 165 kg ha^{-1},

resulting in different amounts of C and N in their residues (Lupwayi and Soon, 2015).

Living plants also influence microbial decomposer communities and their functionality

through their C-rich root exudates. Since litter and root exudates composition differ

among plant species, different crops in rotation systems will not have the same effect on

soil nutrient cycling and on the next crop grown (Huo et al., 2017). Therefore, the choice

of rotation systems design, especially the choice of crop species, is crucial to optimize PSF

effects associated with nutrient cycling. The mechanistic link between shifts in microbial

communities caused by the crop's identity, soil nutrients availability, and the following

crop's growth and nutrition is, however, not clear, especially in organic agroecosystems.

One possible explanation could be changes in organic N depolymerization.

Organic N depolymerization is a limiting step of the enzymatic cascade leading to

mineralization and nitrification, producing inorganic N available for crops (Jan et al.,

2009). The input of C-rich compounds, from plant litter and/or root exudates, steers up

the C: N ratio, forcing microbes to mine for nitrogen (Jones et al., 2009; Kuzyakov and Xu,

2013) by depolymerizing soil organic matter (SOM) containing organic N, such as protein,

chitin and peptidoglycan (Zhu et al., 2014). When their elemental demand is met,

microbes will mineralize the excess N into inorganic form (ammonium and nitrate), usable

by plants. Not to mention that small organic N forms (e.g. amino acids) are also usable by

plants, highlighting the importance of N depolymerization for crop nutrition (Näsholm et

al., 1998). Extracellular enzymes produced by microbial decomposers mediate soil N

depolymerization. Microbes will release extracellular enzymes in soil, and their enzymatic

activities will be induced by the presence of their substrate (Geisseler and Horwath,

2008). Nitrogen release from depolymerization depends on the microbial decomposer

communities, soil physicochemical properties, and the organic substrate availability
(Séneca et al., 2021), all of which are influenced by the identity of the plant and the crop
rotation system. Increased N depolymerization may be one of the mechanisms
responsible for the benefits of crop rotation systems. Yet, to our knowledge, few studies
have tested that hypothesis under organic crop production conditions.

Here, we hypothesize that peas and faba beans, two legumes, have different lasting
influences on the microbial community diversity and composition, changing the microbial
depolymerization capacity, and thereby impacting soil inorganic N availability and quality
and yield of wheat grown the following season. To test this hypothesis, we grew wheat
with and without organic fertilizer in plots where peas or faba beans had been previously
grown, under production conditions at two different fields in the Abitibi-Témiscamingue
region (Québec Canada).

2. Materials and methods

2.1 Experimental design

We conducted a two-year field experiment in Abitibi-Témiscamingue (Québec Canada)

on two farms under actual farm operating conditions, one in Palmarolle (48.665°N,

79.201°W) and in Cloutier (48.00297°N, 79.15°W). The goal of the experiment was to

compare the effect of two legume crops (faba bean and peas) on the microbial

communities and activities of wheat planted the following year. We also wanted to

compare four fertilization strategies for wheat: granulated chicken manure (organic

fertilizer), clover (*Trifolium repens*, cv. Huia) cover crop, fertilizer+cover crop, control

without fertilization nor cover crop. We set up a split- plot experiment, with three blocks

per field. In the first year (legume growing season), we defined randomly two plots (faba

bean and peas) within each block. The following year (wheat growing season), we

randomly defined two subplots (organic fertilizer and no fertilizer), and then two sub-

subplots (clover cover crop and no cover crop). This resulted in 24 plots per field (48 total)

(Fig. 1).

[Insert Figure 1 here]

Both sites were grassland previously cropped to perennial forages for at least three years

before our experiment. Palmarolle soil texture is clay with 80, 355 and 565 g kg^{-1} of sand

(2000–50 µm), silt (50-20 µm) and clay (<2 µm) particles at 0-15 cm depth, respectively.

It belongs to the Gleysolic order according to the World Reference Base for Soil Resource

system (WRB, 2007) and the Canadian System of Soil Classification (Group et al., 1998).

Cloutier soil texture is a silty clay with 38, 481 and 520 g kg^{-1} of sand, silt and clay particles

at 0-15 cm depth, respectively, and also belongs to the Gleysolic order (Group et al., 1998)

(WRB, 2007).

In fall 2022, both sites were moldboard plowed to a depth of 17 cm. In spring 2023,

successive passes with a disk harrow and a field cultivator were carried out to prepare the

seedbed. Faba beans and peas were sown in 2023, using the producer's equipment (Table

1). Plots at Palmarolle were 11 m wide by 100 m long, while those at Cloutier were 4 m

wide by 120 m long. At Palmarolle, faba bean was buried at the end of the growing season

since it did not reach full maturity, while pea was harvested. At Cloutier, both crops were

harvested at the end of the season (Table 1). But, since faba bean grains did not reach full

maturity at Cloutier, residues were spread and buried a few days after harvest. Therefore,

faba beans were used as green manure at both fields. Thus, residue management differed

between both site (Cloutier: fragmented residues were buried a few days after and

Palmarolle: residues were buried at the time of harvest).

The following year (2024), the subplots were fertilized or not with the equivalent of 0.5

Mg N ha^{-1} and 1 Mg N ha^{-1} of granulated chicken manure (N:5-P:3-K:2) for Palmarolle and

Cloutier, respectively. Wheat was sown on the same day as the fertilization occurred (i.e.,

May 30th at Palmarolle and June 04 at Cloutier). The sub-subplots were assigned or not to

receive clover seeds as a cover crop (sown on the same day and not incorporated into the

soil). However, at the time of sampling, the clover had not established and we disregarded

this treatment, treating the two sub-subplots from these treatments as replicates. At the

end of the growing season, wheat was sampled and harvested (Table 1).

Table 1: Sowing, Harvesting and sampling dates for Cloutier and Palmarolle in 2023 and 2024

Field	Crops	Sowing rate (kg ha^{-1})	Sowing date	Harvesting date	Incorporation	Beginning of the season soil sampling date	Mid-growing season soil sampling date
Clou	Faba beans	265	2023-06-07	2023-10-17 Buried afterward	Yes	2023-06-06	2023-08-01
	Peas	265		2023-09-27	No		
	Wheat	170	2024-06-04	2024-09-12	No	2024-05-23	2024-07-31
Pal	Faba beans	265	2023-05-21	NA Buried	Yes	2023-05-21	2023-07-31
	Peas	265		2023-09-18	No		
	Wheat	170	2024-05-30	2024-09-04	No	2024-05-21	2024-07-30

2.2 Soil sampling

Every year, we sampled twice during the season, in the beginning and mid-growing season

(Table 1). Wheat was approximately in the tillering phase when we sampled in 2024. We

sampled five points per sub-subplot (one in the center and four in the corners, at

approximately 0.5 m from the sub-subplot borders), and we combined all samples into a

composite, creating one composite sample for each of the 48 sub-subplots. From the five

points, we sampled the bulk soil by collecting soil (0-15cm) between rows. When present

(mid-growing season), we sampled the rhizosphere of the main crops (faba bean, peas or

wheat) by gently removing plants from the same five points, shaking the roots to remove

the excess soil and considering the remaining soil attached to the rhizosphere (i.e., the

soil influenced by plant roots).

Soil samples were kept in a cooler until they were brought back to the laboratory.

Samples were kept at 4 °C before sieving and aliquoting. We separated the rhizospheric

soil from the roots before sieving. We sieved (4mm) rhizospheric and bulk soil, and we

prepared two aliquots for each sample: one stored at 4 °C for enzymatic assays and

ammonium/nitrate quantification, and another stored at -20 °C for DNA extraction.

2.3 Wheat sampling

At the end of the growing season, we sampled wheat to determine yield and protein

content. Each sub-subplot was subsampled at four randomly distributed locations within

the plot. At each location, we sampled a quadrat of wheat by placing two 76 cm long half-

frames on the soil at about 45 cm width from each other, including 5 wheat rows

(approximately 0.342 m^2). At each subsample point, we measured the internal width

between half-frames to calculate precisely the sampling area, and we harvested wheat.

We stored wheat in paper bags and brought it back to the laboratory. Wheat was then

dried at 35 °C for several days and kept for further analysis.

2.4 DNA extraction and sequencing

We extracted soil DNA from 0.25 g of bulk and rhizosphere soil (previously sieved at 4

mm) samples using the DNeasy PowerLyzer power soil kit (Qiagen) following the

manufacturer's protocol. DNA was eluted in 50 μL of elution buffer and was stored at

−20°C until further analyses. We prepared libraries for 16S and ITS amplicon sequencing

according to Illumina's "16S Metagenomic Sequencing Library Preparation" guide

(Part#15044223Rev.B). We prepared libraries for the bacterial 16 S rRNA gene using

primers 16S: 515Y and 926R, and for the fungal ITS1 region, we used the primers ITS1F

and 58A2R. NextSeq (Illumina) sequencing (PE300 10M reads) was performed at the

Centre d'expertise et de service Génome Québec (CESGQ, Montréal, Canada).

2.5 Enzymatic potential activities quantification

We quantified the potential activity of three exoenzymes involved in SOM

depolymerization (deaminases, β-glucosidases and proteases) using colorimetric

methods. Every protocol follows the same principle: soil samples were incubated with an

excess of substrate under optimal conditions. Therefore, we measured the maximum

enzymatic capacity under optimal lab conditions, meaning that these results do not

reflect real-time activity in the soil.

2.5.1 Protease activity

To quantify the potential activity of soil protease, we used a protocol adapted from Jesmin

et al. (2022). Briefly, we incubated 1 g of dried sieved (4 mm) soil with 2.5 mL sodium

caseinate (2%) and 2.5 mL Tris buffer for two hours at 50°C and 100 rpm. After incubation,

we added 5 mL Trichloroacetic acid (1 g L^{-1}) to stop the enzymatic reaction, and we used

0.75 mL Na_2CO_3 and 0.25 mL Folin & Ciocalteu's phenol (1:3) to quantify tyrosine, the

product that produces a blue colour. We used 500 μL of every sample to measure the

absorbance at 650-700 nm using a TECAN Infinite M plate reader. The potential protease

activity is expressed in μmol tyrosine release g^{-1} dry soil h^{-1} incubation and was quantified

using a tyrosine standard curve (2-fold serial dilution from 1.0 to 0.005 mM).

2.5.2 β-glucosidases activity

To quantify the potential activity of soil β-glucosidase, we used a protocol adapted from

Eivazi and Tabatabai (1988). Briefly, we incubated 1 g of dried sieved (4 mm) soil with a

buffer solution containing p-nitrophenyl glycoside (25 mM) at 37°C for 1 h. After

incubation, we added calcium chloride (0.5 M) and Tris buffer (0.1 M, pH 12) to extract p-

nitrophenol, resulting from the degradation of the substrate. Extraction of the product

produces a yellow colour, and we measured absorbance at 415 nm using a TECAN Infinite

M plate reader. The potential activity is expressed in μmol p-nitrophenol released g^{-1} dry

soil h^{-1} incubation and was quantified using a p-nitrophenol standard curve (2-fold serial

dilution from 0.5 to 0.0078 g L^{-1}).

2.5.3　Deaminase activity

To quantify the potential activity of soil deaminases, we used a protocol adapted from

Killham and Rashid (1986). Briefly, we incubated 1 g of dried sieved (4 mm) soil with 1.2-

diamino-4-nitrobenzene (0.06 g L^{-1}) for 20 h at 25 °C, 100 rpm. The substrate is red, and

the colour intensity decreases with the deamination of the benzene ring. After incubation,

we extracted twice the remaining substrate using methanol 100 % and filtered (Whatman

no1) the solution to remove any impurities. We measured the absorbance at 405 nm

using a TECAN Infinite M plate reader. The potential activity is expressed in and was

quantified using a 1.2-diamino-4-itrobenzene standard curve

(0/0.008/0.011/0.014/0.019/0.025/0.034/0.045/0.06 g L^{-1}).

2.6 Available inorganic nitrogen quantification

To quantify ammonium and nitrate, we first needed to extract nitrogen from the soil,

using a protocol adapted from Maynard et al. (1993).

To do so, we incubated 1 g of dried sieved (4 mm) soil with 10 mL of KCl (2 M) for 1 h at

25 °C at 100 rpm. After incubation, we filtered (Whatman no. 42) the soil slurry to remove

any soil and debris. Soil extracts were kept at 4 °C until further analyses.

2.6.1 Ammonium quantification

To quantify ammonium in soil extracts, we adapted a protocol from Rhine et al. (1998).

Briefly, we added 100 µL of the sample (standard and/or soil extract) to a 96-well plate.

Using a multi-channel pipette, we added 50 µL of citrate reagent, 50 µL of phenylphenol

nitroprusside, 25 µL of buffered hypochlorite reagent and 50 µL of deionized water to

each well and incubated 45 min at room temperature. Based on the Berthelot method,

ammonium in soil extracts reacts with the buffered hypochlorite solution and the

phenylphenol nitroprusside, producing a blue product. After incubation, we measured

the absorbance at 660 nm using a TECAN Infinite M plate reader and the concentration

of ammonium is expressed in mg g^{-1} of air-dried soil and quantified using a $(NH_4)_2SO_4$

standard curve (2-fold serial dilution from 20 to 0.0195 mg L^{-1}).

2.6.2 Nitrate quantification

To quantify nitrate in soil extracts, we adapted a protocol from Hood-Nowotny et al.

(2010). Briefly, we added 100 μL of the sample (standard and/or soil extract) to a 96-well

plate. Using a multi-channel pipette, we added 100 μL of VCl_3 solution and 100 μL of the

Griess reagent mix (a solution containing 1:1 Griess solution 1 and Griess solution 2) to

each well, and we incubated the plate for 1 h at 37 °C. Based on the VCL_3/Griess method,

nitrate in soil extracts is converted to nitrite with VCL_3, and nitrite reacts with both Griess

solutions to produce a pink color. After incubation, we measured absorbance at 540 nm

using a TECAN Infinite M plate reader and the concentration of nitrate is expressed in mg

g^{-1} of soil and quantified with a standard curve of KNO_3 (2-fold serial dilution from 10 to

0.0195 mg L^{-1}).

2.7 Wheat grain yield and protein content

At the end of the 2024 growing season, wheat was hand harvested (see section 2.3) to be

threshed using a laboratory thresher (DL-350, Wintersteiger, Ried im Innkreis, Austria).

Grains were then cleaned with a sample cleaner (MNL Pfeuffer, Kitzingen, Germany).

Grain yield (Y_{FH}, kg ha^{-1}) was determined in both fields at the subsample level by weighing

clean grain harvested for the given surface area sampled at each subsampling point (see

section 2.3). Grain protein content (%) was determined only at Cloutier and was analyzed

using a Near-infrared spectrometer (Infratec 1241, FOSS, Hillerød, Denmark). Wheat yield

was reported on a 13.5 % relative humidity content ($Y_{13.5}$, kg ha^{-1}) as follows: $Y_{13.5}$ = (Y_{FH} x

(100 – W_{FH})) / (100 – 15). We used $Y_{13.5}$ for our statistical analyses. We determined yield

and protein content (when applicable) for each subsample harvested in the field (four

subsamples), and we used the mean of the subsamples to perform our statistical analyses.

2.8 Bioinformatic analyses

We analyzed amplicon sequencing data (16S rRNA gene and ITS region) using the DADA2

pipeline in R to process raw reads. For 16S, we used truncLen (280,250) and maxEE (2,2)

to filter and trim forward and reverse reads. We used the pooled method for removing

chimeras, and we assigned taxonomy with the SILVA reference database. After the

assignation, we removed chloroplasts and mitochondria from our dataset before further

analysis. For ITS, we used truncLen (270,250) and maxEE (2,2) to filter and trim forward

and reverse reads. We used the pooled method for removing chimeras, and we assigned

taxonomy with the UNITE reference database.

2.9 Statistical analysis and data visualization

We performed Statistical analyses in R (version 4.5.1, *The R Foundation for Statistical*

Computing), and we created figures using the package ggplot2.   We divided and

performed statistical analyses for each field (i.e., Cloutier and Palmarolle) and for each

year (i.e., 2023 and 2024) separately. First, we performed Shapiro-Wilk tests to check the

normality of the residuals and Bartlett tests to check the homogeneity of variances. When

the assumptions were met, we performed a full factorial analysis of variance (ANOVA).

For 2024, we performed ANOVAs for the effect of the previous legume crop (i.e., pea and

faba bean), organic fertilization (i.e., with or without granulated chicken manure), soil

compartment (i.e., rhizosphere and bulk soil), while including the split-split-plot design in the model. For 2023, we performed ANOVAs for the effect of the previous legume crop (i.e., pea and faba bean) and soil compartment (i.e., rhizosphere and bulk soil), while including the split-split-plot design in the model. We did this for potential enzymatic activities, soil ammonium and nitrate content. When the assumptions were not met, we first tried to transform our data (log or square root transformation), and if that did not work, we performed a Kruskal-Wallis non-parametric test, testing each factor independently. Specifically, in the 2024 second sampling, Cloutier nitrate did not meet the normality of residuals assumption, Cloutier ammonium and protease activity did not meet the homogeneity of variances assumption, and deaminase data did not meet both assumptions. Palmarolle deaminase data did not meet the normality of residuals assumption. In the 2024 first sampling, Cloutier nitrate did not meet the homogeneity of variances assumption. Palmarolle nitrate and ammonium did not meet both assumptions and deaminase data did not meet the homogeneity of variances assumption. Finally, in 2023 second sampling, Cloutier nitrate did not meet the homogeneity of variances assumption. Palmarolle nitrate did not meet both assumptions, Palmarolle ammonium did not meet the normality of residuals assumption and deaminase data did not meet the homogeneity of variances assumption. We performed Kruskal-Wallis nonparametric tests to assess the effects of previous crop, organic fertilization, the interaction between previous crop and fertilization and block on wheat yield ($Y_{13.5}$, kg ha^{-1}) and protein content. We performed Kruskal-Wallis nonparametric tests to assess the effects of previous crop, organic fertilization, soil compartment, and block on alpha-diversity and

relative abundance data, since this data did not meet the assumption of parametric

ANOVA. To analyze the bacterial and fungal community structure, we first transformed

our dataset using the Hellinger transformation. We then used Euclidean distance matrices

based on our transformed data to visualize the community structure using a principal

coordinate analysis (PCoA). We first used betadisper to test the homogeneity of

multivariate dispersion. We tested the effect of the previous crop, organic fertilization

and soil compartment on the community using PERMANOVA (999 permutations). To

consider the experimental design, we used the argument strata = block. To assess ASVs

differently abundant in wheat plots after faba bean and peas, we performed three

differential analysis (DEseq2, ANCOM-BC and LEfsE) and kept only ASVs found by all

methods.

2.10 Data availability

The R code used to analyze the data and generate the figures is available on GitHub

(https://github.com/le-labo-yergeau/). Raw sequencing reads were deposited in the NCBI

SRA repository under BioProject accession PRJNA1390138.

3. Results

We focused our microbial analysis on the mid-growing season samples (2024) to assess

previous legume crops' PSF effects, except when it is mentioned.

3.1 Previous crop PSF on microbial communities' composition

3.1.1 Cloutier 2024

Before performing PERMANOVA, we tested the homogeneity of multivariate dispersion

for both bacterial and fungal communities, and it was not significant, meaning that the

results of the tests were not influenced by differential dispersion across treatments. The

bacterial and fungal community compositions of wheat plots differed significantly

depending on the previous legume sown in 2023 and the addition of organic fertilization

(i.e. granulated chicken manure) in 2024 (Fig. 2.A and 2.C; Table S.1 and S.2). For the

bacterial community (Fig. 2.A), there was also a significant interaction between the

previous crop and the fertilization, meaning that depending on the previous crop sown,

fertilization's effect varied. In Figure 2.A, we can see that the addition of granulated

chicken manure increases the variability of the bacterial community composition,

especially when the previous crop was peas. Therefore, fertilization reduced the PSF

effect of the previous crop. For the fungal community (Fig. 2.B), there was no interaction

between the effect of the previous crop and the organic fertilization. The soil

compartment (rhizosphere and bulk soil) modulated the fungal community composition

(Fig. 2.B), but not the bacterial community.

After assessing that the previous legume crop identity shapes microbial communities in

wheat plots differently, we tried to find which microbes' relative abundance was different

between plots that received faba beans and peas. After performing DESeq2, ANCOM-BC

and LEfSe analysis, we did not find any bacterial ASV differentially abundant in wheat

plots that received faba bean and peas (Fig. 3.B). On the other hand, we found 6 fungal

ASVs differentially abundant in wheat plot depending on previous legume crops (Fig. 3.D;

Table 2). Specifically, fungal ASVs # 16, 87, 387 and 686 were significantly more abundant

in wheats plots that received peas as the previous legume crop and ASVs # 169 and 271

were significantly more abundant in wheat plots that received faba bean as the previous

legume crop.

3.1.2 Palmarolle 2024

Before performing PERMANOVA, we tested the homogeneity of multivariate dispersion,

and it was not significant for fungi. However, we found significant differences for bacterial

communities (p-value = 0.0249), which suggests that the PERMANOVA results could also

be due to differential dispersion of the bacterial communities among treatments. The

bacterial and fungal community composition of wheat plots is significantly different

depending on the previous crop sown in 2023 and the soil compartment (Fig. 2.C and 2.D;

Table S.1 and S.2). There is also a significant effect (p = 0.043) of the interaction between

the previous crop and the fertilization on the fungal communities' composition in

Palmarolle (Fig. 2.D). Like the bacterial community composition in Cloutier, fertilization

makes the fungal community composition more variable (Fig. 2.C).

After performing differential analysis (DESeq2, ANCOM-BC and LEfSe analysis), we found

that the bacterial ASV # 520 was significantly more abundant in wheat plots that received

faba bean as the previous legume crop (Fig. 4.B). We also found 8 fungal ASVs

differentially abundant in wheat plots depending on previous legume crops (Fig. 4.D;

Table 2). Specifically, fungal ASVs # 16, 162, 288, 323, 390, 580, 1155, 1159 were

significantly more abundant in wheat plots that received peas as the previous legume

crop.

[Insert Figure 2 here]

[Insert Figure 3 here]

[Insert Figure 4 here]

Table 2: Bacterial and fungal ASVs differentially abundant in wheat plot at Cloutier and Palmarolle found with DESeq2, ANCOM-BC and LEfSe.

	#ASV	Kingdom	Phylum	Class	Genus	Previous crop	log2FC D	logFC A	lda lefse
Cloutier	16	Fungi	Ascomycota	Dothideomycetes	*Didymella*	peas	2.84	3.04	4.12
	87	Fungi	Ascomycota	Sordariomycetes	*Phomatospora*	peas	4.26	2.48	3.95
	169	Fungi	Ascomycota	Leotiomycetes	*Cadophora*	faba	-3.02	-1.66	3.18
	271	Fungi	Basidiomycota			faba	-3.76	-1.63	2.91
	387	Fungi	Chytridiomycota	Rhizophydiomycetes		peas	4.10	1.54	2.33
	686	Fungi	Ascomycota	Sordariomycetes	*Phomatospora*	peas	4.04	1.89	2.65
Palmarolle	16	Fungi	Ascomycota	Dothideomycetes	*Didymella*	peas	3.99	3.05	4.27
	162	Fungi	Ascomycota	Leotiomycetes		peas	4.56	2.29	3.59
	288	Fungi	Ascomycota	Eurotiomycetes	*Cladophialophora*	peas	3.62	1.95	3.44
	323	Fungi	Mortierellomycota	Mortierellomycetes	*Mortierella*	peas	5.56	2.57	3.03
	390	Fungi	Chytridiomycota	C_GS13		peas	3.12	1.73	3.22
	580	Fungi	Ascomycota	Eurotiomycetes	*Cladophialophora*	peas	4.10	2.12	2.95
	1155	Fungi	Ascomycota	Dothideomycetes	*Stemphylium*	peas	3.86	1.53	2.41
	1159	Fungi	Ascomycota	Sordariomycetes	*Metapochonia*	peas	2.73	1.25	2.31
	520	Bacteria	Actinobacteriota		*Cryobacterium*	faba	-2.51	-1.48	2.62

3.2 Potential Enzymatic activities associated with depolymerization

3.2.1 Cloutier 2024 and 2023

During wheat growth (2024 second sampling), there was significantly more β-glucosidase and deaminase potential activities in the wheat rhizosphere than in the bulk soil (Fig. 5.B and 5.C; Table 3). The potential β-glucosidase activity in wheat plots was higher, but not significant, after faba bean than after peas. The addition of organic fertilizer at the beginning of the season did not affect enzymatic activities during wheat growth. Before wheat growth (2024 first sampling), the potential protease activity was higher in wheat plots after peas than faba bean (Table 3). Fertilization did not affect enzymatic activities

before wheat growth. In 2023, when faba bean and peas were growing, there was

significantly more protease and deaminase potential activities in both legumes'

rhizosphere than in the bulk soil (Table 3).

3.2.2 Palmarolle 2024 and 2023

During wheat growth (2024 second sampling), there was significantly more protease and

β-glucosidase potential activities in the wheat rhizosphere than in the bulk soil (Fig. 5.D

and 5.E; Table 3). After faba bean, wheat plots had more protease and β-glucosidase

activities than after peas. The addition of organic fertilizer at the beginning of the season

did not affect enzymatic activities during wheat growth. Before wheat growth (2024 first

sampling), fertilization and the previous crop did not influence the three exoenzymes

potential activities tested. In 2023, when faba bean and peas were growing, there was

significantly more protease, β-glucosidase, and deaminase potential activities in both

legumes' rhizosphere than in the bulk soil (Table 3).

[Insert Figure 5 here]

Table 3: Effects of previous crop, organic fertilization and soil compartments (rhizospheric vs. bulk soil) on proteases, β -glucosidases and deaminases potential activities at 0-15 cm depth at Cloutier and Palmarolle during the wheat mid-season (2024 2 sampling), wheat beginning of the season (2024 1 sampling) and legumes mid-season (2023 2 sampling).

Clou 2024 2 sampling

		Proteases (Kruskal)				B-glucosidases (ANOVA square transformed)				Deaminases (Kruskal)			
Stratum						B:PC	B:PC:F	B:PC:F	Within				
Factor/term		PC	C	F	B	PC	F	PC:F	C	PC	C	F	B
Statistic		0.41	2.93	1.58	2.09	2.40	0.04	0.42	105.61	0.82	4.42	0.79	4.35
df		1	1	1	2	1	1	1	1	1	1	1	2
p-value		0.523	**0.087** .	0.208	0.352	0.262	0.856	0.552	**1.16E-11** ***	0.364	**0.035** *	0.375	0.113

Clou 2024 1 sampling

		Proteases (ANOVA square transformed)			B-glucosidases (ANOVA log transformed)			Deaminases (ANOVA square transformed)		
Stratum		B:PC	B:PC:F	B:PC:F	B:PC	B:PC:F	B:PC:F	B:PC	B:PC:F	B:PC:F
Factor/term		PC	F	PC:F	PC	F	PC:F	PC	F	PC:F
Statistic		32.03	0.73	0.72	0.005	0.45	0.003	0.003	0.06	5.66
df		1	1	1	1	1	1	1	1	1
p-value		**0.030** *	0.440	0.445	0.951	0.540	0.957	0.962	0.820	**0.076** .

Clou 2023 2 sampling

		Proteases (ANOVA square transformed)			B-glucosidases (ANOVA square transformed)			Deaminases (ANOVA square transfromed)		
Stratum		B:PC	Within	Within	B:PC	Within	Within	B:PC	Within	Within
Factor/term		PC	C	PC:C	PC	C	PC:C	PC	C	PC:C
Statistic		0.53	49.67	0.30	0.37	0.04	2.19	48.71	35.89	0.07
df		1	1	1	1	1	1	1	1	1
p-value		0.542	**1.61E-08** ***	0.586	0.605	0.846	0.147	**0.020** *	**4.87E-07** ***	0.798

Pal 2024 2 sampling

		Proteases (ANOVA)				B-glucosidases (ANOVA log transformed)				Deaminases (Kruskal)			
Stratum		B:PC	B:PC:F	B:PC:F	Within	B:PC	B:PC:F	B:PC:F	Within				
Factor/term		PC	F	PC:F	C	PC	F	PC:F	C	PC	C	F	B
Statistic		1.65	0.70	0.82	39.61	0.79	0.14	0.22	11.10	0.27	0.08	3.33	7.65
df		1	1	1	1	1	1	1	1	1	1	1	2
p-value		0.328	0.451	0.417	**4.65E-07** ***	0.467	0.731	0.667	**0.002** **	0.606	0.775	**0.068** .	**0.022** **

Pal 2024 1 sampling

		Proteases (ANOVA square transformed)			B-glucosidases (ANOVA square transformed)			Deaminases (Kruskal)		
Stratum		B:PC	B:PC:F	B:PC:F	B:PC	B:PC:F	B:PC:F			
Factor/term		PC	F	PC:F	PC	F	PC:F	PC	F	B
Statistic		1.89	0.01	1.39	0.27	0.18	0.90	0.03	0.03	1.81
df		1	1	1	1	1	1	1	1	2
p-value		0.303	0.916	0.304	0.654	0.694	0.395	0.862	0.862	0.406

	Stratum	Kruskal			ANOVA (log transfromed)			ANOVA (no transformed)		
		PC	C	B	B:PC PC	Within C	Within PC:C	B:PC PC	Within C	Within PC:C
Pal 2023 2	Factor/term									
sampling	Statistic	0.02	14.35	1.82	5.924	9.024	2.58E-05	2.194	3.205	2.401
	df	1	1	2	1	1	1	1	1	1
	p-value	0.882	**0.0002** ***	0.402	0.248	**0.005** **	0.996	0.378	**0.081** .	0.129

.: p<0.1; *: p<0.05; **p<0.01; ***:p<0.001

B: Block; PC: Previous crop, F: Fertilization

Factor/Term, Statistic, df and p-value in bold are the significant and/or almost significant values

3.3 Soil inorganic N content

3.3.1 Cloutier 2024 and 2023

Figure 6 shows the amount of inorganic N in the bulk soil and the rhizosphere of wheat,

at mid-growing season in 2024. In the wheat rhizospheric soil at Cloutier, there is

significantly less inorganic N in ammonium and nitrate forms than in the bulk soil (Fig. 6.A

and 6.B; Table 4). There is also a trend toward an increased concentration of ammonium

(P=0.08) in the faba bean plots as compared to the pea plots (Fig. 6.A, Table 4). This could

be related to the nearly significant (P= 0.084) compartment x previous legume interaction

at Cloutier in 2023 (legume growing season) that stemmed from a higher concentration

of ammonium in the rhizosphere vs. bulk soil, but only for faba bean (Table 4). In the first

sampling of 2024, Cloutier nitrate concentration significantly differed between blocks.

When legumes were the main crop (2023 second sampling), there was less nitrate in both

legumes' rhizosphere than in the bulk soil.

3.3.2 Palmarolle 2024 and 2023

In the wheat rhizospheric soil at Palmarolle, there is significantly less nitrate, but more

ammonium than in the bulk soil (Fig. 6.C and 6.D; Table 6). Also, at Palmarolle, there was

less nitrate in wheat plots after faba beans than after peas, regardless of the

compartment (Fig. 6.D).   In the first sampling of 2024, Palmarolle ammonium

concentration significantly differed between blocks. When legumes were the main crop

(2023 second sampling), there was less nitrate in both legumes' rhizosphere than in the

bulk soil and ammonium concentration differed significantly between blocks.

[Insert Figure 6 here]

Table 4: Effects of previous crop, organic fertilization and soil compartments (rhizospheric vs. bulk soil) on

nitrate and ammonium content at 0-15 cm depth at Cloutier and Palmarolle during the wheat mid-season

(2024 2 sampling), wheat beginning of the season (2024 1 sampling) and legumes mid-season (2023 2

sampling).

		Nitrate				Ammonium			
Clou 2024 2 sampling		Kruskal				Kruskal			
	Stratum								
	Factor/term	PC	C	F	B	PC	C	F	B
	Statistic	2.10	20.58	1.35	1.84	3.07	6.22	0.00	2.14
	df	1	1	1	2	1	1	1	2
	p-value	0.147	**5.73E-06** ***	0.246	0.399	**0.08**	**0.013** *	0.984	0.343
Clou 2024 1 sampling		Kruskal				ANOVA (square transformed)			
	Stratum					B:PC	B:PC:F	B:PC:F	
	Factor/term	PC	F	B		PC	F	PC:F	
	Statistic	0.56	2.43	6.50		0.96	3.17	0.68	
	df	1	1	2		1	1	1	
	p-value	0.453	0.119	**0.039** *		0.431	0.149	0.455	
Clou 2023 2 sampling		Kruskal				ANOVA			
	Stratum					B:PC	Within	Within	
	Factor/term	PC	C	B		PC	C	PC:C	
	Statistic	0.38	3.92	2.34		0.04	1.11	3.13	
	df	1	1	2		1	1	1	
	p-value	0.536	**0.048** *	0.311		0.855	0.299	**0.084** .	
Pal 2024 2 sampling		ANOVA (square transformed)				ANOVA (square transformed)			
	Stratum	B:PC	B:PC:F	B:PC:F	Within	B:PC	B:PC:F	B:PC:F	Within
	Factor/term	PC	F	PC:F	C	PC	F	PC:F	C
	Statistic	6.23	0.05	0.99	32.68	1.17	3.39	0.04	14.95
	df	1	1	1	1	1	1	1	1
	p-value	0.130	0.834	0.375	**2.47E-06** ***	0.393	0.139	0.845	**5.08E-04** ***
Pal 2024 1 sampling		Kruskal				Kruskal			
	Stratum								
	Factor/term	PC	F	B		PC	F	B	
	Statistic	0.013	1.14	0.88		0.10	0.08	9.91	
	df	1	1	2		1	1	2	
	p-value	0.908	0.285	0.646		0.751	0.773	**0.007** **	
Pal 2023 2 sampling		Kruskal				Kruskal			
	Stratum								
	Factor/term	PC	C	B		PC	C	Block	
	Statistic	0.07	5.990	0.43		0.41	0.02	13.72	
	df	1	1	2		1	1	2	
	p-value	0.798	**0.014** *	0.807		0.523	0.898	**0.001** **	

.: p<0.1; *: p<0.05; **p<0.01; ***:p<0.001

B: Block; PC: Previous crop, F: Fertilization

Factor/Term, Statistic, df and p-value in bold are the significant and/or almost significant values

3.4 Wheat yield and grain quality

3.4.1 Cloutier 2024

The addition of organic fertilizer significantly enhanced wheat grain protein content (Fig.

7: Table 5). Also, faba bean, when used as the previous crop, significantly increased wheat

grain protein content as compared to peas (Fig. 7).

[Insert Figure 7 here]

Table 5: Kruskal-Wallis results for the effect of organic fertilization and previous crops on grain protein

content and yield ($Y_{13.5}$) at Cloutier

Response variable	Factor	Statistic	p-value	p.adj	
Yield(kg ha^{-1})	Fertilization	0.093	0.7609	0.7609	
	Previous crop	1.245	0.2645	0.3174	
	Fertilization : Previous crop	7.265	0.0639	0.1024	
	Block	**10.619**	**0.0050**	**0.0148**	*
Protein content (%)	**Fertilization**	**14.087**	**0.0002**	**0.0007**	***
	Previous crop	**5.927**	**0.0149**	**0.0298**	*
	Fertilization : Previous crop	**20.106**	**0.0002**	**0.0006**	***
	Block	**10.048**	**0.0066**	**0.0158**	*

*: p<0.05; **p<0.01; ***:p<0.001

p.adj refers to the p-value with Benjamini-Hochberg correction

Factor/Term, Statistic, df, p-value and p.adj in bold are the significant values

3.4.2 Palmarolle 2024

For Palmarolle, we were only able to determine the grain yield at the end of the growing

season. We found no significant effect of fertilization and the previous crop on grain yield

(Table 5).

Table 5: Kruskal-Wallis results for the effect of organic fertilization and previous crops on grain yield ($Y_{13.5}$)

at Palmarolle

Response variable	Factor	Statistic	p-value	p.adj
	Fertilization	0.257	0.6120	0.9180
	Previous crop	0.833	0.3613	0.7225
Yield (kg ha-1)	Fertilization : Previous crop	1.461	0.6913	0.9217
	Block	0.247	0.8838	1.0000

*: p<0.05; **p<0.01; ***:p<0.001

p.adj refers to the p-value with Benjamini-Hochberg correction

Factor/Term, Statistic, df, p-value and p.adj in bold are the significant values

4. Discussion

This study aimed to evaluate the impact of the identity of the previous legume crop on

microbial community composition and functional capacity in the following growing

season. We hypothesized that a part of the positive legume-cereal rotation effect on

wheat yield and grain quality could be mediated by changes in the C and N

depolymerization capacity of the microbial community. First, we confirmed that legume

identity modifies soil microbial communities in the following growing season, and that

PSF effects can be transferred to other plant species (in our case, from legumes to wheat).

We also found, at Cloutier, that faba bean as previous crop increased soil ammonium

content and wheat grain protein content, a key parameter of grain baking quality (Nuttall

et al., 2017). We did not, however, find significant changes in potential protein and

cellulose depolymerization activities, infirming our main hypothesis. Instead, the

mechanism behind the effect of PSF on wheat grain quality could be an increased

mineralization linked to shifts in the relative dominance of bacteria and fungi or a higher

substrate availability in faba bean plots.

Under real farming conditions in fields located 100 km apart, we confirmed that the

previous crops' PSF influenced the composition of microbial communities. In fact, among

all factors tested, the previous crop was the only one that consistently shaped microbial

communities in both fields. This highlights the strength of the previous crop effect on

microbial communities. Indeed, due to logistics (e.g., the size of plots needed to match

the seeder and the fertilizer applicator) and field size, we could not set up more than

three complete blocks. Additionally, there was quite some variation between the blocks,

but we still found significant evidence of a PSF effect. Previous crop PSF effects on

microbial community composition has been observed before (Wang et al., 2023a; Dang

et al., 2024). We also found ASVs significantly differently abundant depending on the

previous crop identity, proving the legacy effect of previous legume crops on microbial

communities. In both fields, fungal communities were more altered by the previous crop

identity and residue management than bacterial communities. Fungi are known to be key

players in decomposition and can form close relationships with plants, as symbionts and

pathogens (Zeilinger et al., 2016). Therefore, changes in fungal community composition

and relative abundance due to the previous crop identity are expected (Li et al., 2023).

Fungi are also known to be key players in plant compounds degradation, especially lignin

and cellulose (Schneider et al., 2012; Arcand et al., 2016). For instance, in Cloutier's wheat

plots after faba bean, Leotiomycetes, more specifically *Cadophora luteo-olivacea,* were

more abundant. *Cadophora spp.* possess a wide variety of exoenzymes associated with C

and N depolymerization, such as protease and carbohydrate active enzymes (Bruyant et

al., 2024). They could have potentially played a role in faba bean residue

depolymerization and wheat nutrition. In Palmarolle's wheat plots after faba bean,

Cryobacterium spp. was more abundant. *Cryobacterium spp.* is often correlated to higher

lignin and C content (Shamshitov et al., 2025). The incorporation of faba bean residue

could have enhanced *Cryobacterium spp.* abundance during the wheat growing season.

Fungi have a higher C:N ratio than bacteria and increases in fungal dominance in

depolymerization processes would result in higher mineralization rate of the

depolymerized organic N (Strickland and Rousk, 2010), without necessarily changing

potential enzymatic activities.

Faba beans and peas are two legumes often used in crop rotation systems to replenish

soil N pools. Both crops fix atmospheric N through a symbiosis with rhizobia (Jensen et

al., 2010). Their PSF effects on microbial community composition were, however, distinct,

supporting other findings (Yixuan et al., 2025). Faba bean is particularly efficient in

increasing soil N for the following crop (Herridge et al., 2008; Peoples et al., 2009). Faba

beans have a coarser root system (i.e., higher mean root diameter and lower specific root

length) and allocate a higher proportion of total plant biomass belowground than pea

(Guerrero-Ramírez et al., 2021), therefore adding more organic matter to the soil

(Nuruzzaman et al., 2006). Differences in root system sizes can also alter soil physical

properties, therefore changing microbial communities. With a root system four times

bigger than peas, faba beans promote aggregate formation, creating different

microenvironments for microbes in the next growing season (Oliveira et al., 2019). At

Cloutier, we also found higher ammonium concentration in the rhizosphere of faba beans

as compared to adjacent bulk soil, while there was no difference for peas. This difference

between faba beans and peas was maintained during the wheat growing season, which

could explain the higher grain protein content of wheat that followed faba beans. Indeed,

even with similar potential depolymerization activities, the soil with more substrate (i.e.

faba bean plots) will mineralize more N, as soil depolymerization activities are substrate-

limited (Greenfield et al., 2020).

Fertilization alters soil properties and microbial communities (Luo et al., 2018), and it was

expected that organic fertilization would affect microbial communities in our study.

Interestingly, microbial communities from wheat growing season did not respond the

same way to organic fertilization in both fields. It is important to note that the organic

fertilizer amount was not the same in both fields, which could in part explain the

difference between fields (Zhou et al., 2021). Nevertheless, organic fertilization increased

the variability of microbial communities coming from peas as compared to faba beans. It

suggests that faba beans have a stronger PSF effects due to a larger addition of root-

derived organic matter and, incorporation of aboveground biomass. It creates a resource-

rich environment, priming the microbial community to mineralize organic fertilizer and

making it less susceptible to changes caused by fertilizer addition (Aschi et al., 2017).

The only significant difference in potential enzymatic activity we found between the two

legumes was for protease activity that was higher in plots that received peas at Cloutier

early in the season, before wheat was sown. In contrast to our main hypothesis, this

increased protease activity did not seem to be beneficial for the wheat, as the wheat

grains harvested from the pea plots had in fact a lower protein content than the ones

harvested in the faba bean plots. This increase in protease potential activity, early in the

season and in the absence of the crops, probably led to microbial immobilization of the

depolymerized N. In that case, the immobilized N would have needed to be mineralized

or re-mobilized from the microbial biomass to reach the wheat. Other enzymes that

depolymerize chitin or peptidoglycan, two abundant soil organic N sources that make

fungal and bacterial cell walls, respectively, could have been good targets to test this idea.

However, since we observed less protein content in wheat grains from the pea plots, the

N produced from protease activity probably remained immobilized. This suggest that the

timing of depolymerization activities is paramount for it to be beneficial to crops. In

wheat, the N assimilated by the plant is mostly taken up before and during the anthesis

stage (Cox et al., 1985; Sharma et al., 2023). Our mid-season sampling might have missed

the potential increase in depolymerization activities that could have increased wheat N

uptake at key growth stages, resulting in higher protein content at the end of the season

at Cloutier. Alternatively, as mentioned above, enzymatic activities are substrate-limited,

and changes in potential enzymatic activities measured in the laboratory with excess

substrate might not represent activities in the field.

Almost every exoenzyme's potential activities were higher in the wheat rhizosphere than

in the bulk soil, suggesting that wheat, through its root exudates, selects for a community

proficient in depolymerization. Our results align with the rhizosphere priming theory, a

plant competitive strategy for N (Kuzyakov and Xu, 2013; L'Espérance et al., 2024). Plants

change the rhizosphere C: N ratio through their C-rich exudates to force microbes to

depolymerize soil organic matter to scavenge for N. When the C-rich exudates are all

consumed, microbes will mineralize the N excess, providing inorganic N to plants (Zhu et

al., 2014). To confirm this hypothetical rhizosphere priming effect, future studies should

measure rhizosphere C: N ratios and root exudates.

Unlike previous studies (Li et al., 2015; Lupwayi et al., 2019), the addition of organic

fertilizer did not have a significant effect on the enzymatic activities, except for the

deaminase activity at Palmarolle. This is surprising, since we added a substrate that can

stimulate microbes to produce and release exoenzymes. In contrast, Breza et al., (2023)

found that the addition of granular urea in more complex rotation systems disturbs the

organic N cycle, reducing the N depolymerization rate to a similar rate as monocropping.

In both studies, fertilization affects exoenzymes activities mediating N depolymerization

(Breza et al., 2023). In our case, the lack of fertilization effects can be due to the fertilizer

type, the application rate and/ or the sampling date. We chose to use a fertilizer used in

this region for organic farms, and the equivalent of only 25-50 kg N ha^{-1} was applied. The

amount applied – which is less than the recommended amount for spring wheat under

organic management (~100 kg N ha^{-1}) – and type of fertilizer was chosen to accommodate

farmers who must reconcile crop needs and fertilizer prices. The lack of fertilization

effects can also be due to the sampling date. Our mid-growing season sampling point in

2024 was approximately two months after the application of the organic fertilizer.

Organic fertilizer availability depends on the soil mineralization rate and usually takes

longer to be available for plants and microbes (Wang et al., 2023b). However, Lazicki et

al. (2020) showed that, under laboratory conditions, poultry manure-based granulated

fertilizer was already mineralized after a few weeks of incubation. Therefore, nutrients

derived from organic fertilizer were probably already immobilized in microbial or wheat

biomass at the time of sampling, which would explain the lack of a fertilizer effect on

enzymatic activities and soil ammonium and nitrate content.

Conducting a two-year field experiment under actual farm operating conditions is a

daunting task, and differences between fields are unavoidable. Pedoclimatic conditions,

residue management, cultivation history, plot sizes, and fertilization rates differed

between our sites. All these differences can alter PSF effects on microbial community

composition and depolymerization capacity. In our case, PSF effects were not found

across both fields for every parameter tested. This may be due to differences in

pedoclimatic conditions, which alter soil physicochemical properties across both fields

(A'Bear et al., 2014). Limitations created by different agricultural practices in our

experiment also highlighted the need to investigate PSF effects more deeply. As

mentioned above, different faba bean incorporation at Cloutier and Palmarolle certainly

influenced microbial communities and their functions. But, even with different residue

management, we found that the previous crops' PSF influenced microbial community

composition and soil ammonium content. Alternatively, any of the above-mentioned

differences between the two fields could have played a role. It seems, therefore, possible

to enhance PSF when combining crop rotations with other agricultural practices.

5. Conclusion

In conclusion, even with limitations encountered by doing an experiment under actual

farm operating conditions (field size, block number, residue management), we showed

that faba bean increased soil ammonium and wheat grain protein content as compared

to peas, at Cloutier. We had to reject our hypothesis that this effect on wheat grain

protein content was due to changes in potential soil enzymatic activities, as we found

little statistical support for it. We have, however, come up with two alternative

explanations. First, most significant shifts in microbial communities due to previous

legume identity were in fungal ASVs. This could result in a shift in the relative dominance

of bacteria and fungi, which, because of differences in C:N ratio between the two groups,

could result in differences in mineralization rates. Second, as soil enzymatic activities are

substrate-limited, the larger root system of faba bean could have brought more substrate

to these plots, increasing mineralization. Depolymerization of cellulose and proteins are

widely distributed functions among soil microbial communities, and their *in-situ* activities

are probably restricted by substrate availability, whereas the ratio between

immobilization/mineralization probably varies with the soil microbial community

composition. More field experiments with *in-situ* measurements would be needed to fully

understand the role of plant-soil feedback in crop rotation systems to improve our

practices and sustainably meet crops' nutritional needs.

6. Funding Sources

This work was supported by the FRQNT Programme de recherche en partenariat (grant

number 322604), the Fondation de l'université du Québec en Abitibi-Témiscamingue and

the Entente sectorielle biolalimentaire de l'Abitibi-Témiscamingue. ELE was supported by

a Fondation Armand-Frappier doctoral scholarship.

703
7. Acknowledgements

We extend our gratitude to our interns: Raphaël Boisvert, Sopheaktra Honn Kuhn , Marc-

Antoine Duchesne, Hubert Côté and Lisa Colleoni whose hard work during field sampling

greatly contributed to this project. We are thankful to Johanne Massy, Martin Jacques,

Werbson Lima Barroso, Mahefa Ravoavison, Sendy Augustin Salomon and Thierry

Nousseassi for their technical assistance, especially with harvest and assessment of wheat

yield and protein content. We are also thankful to the farmers who generously provided

access to their fields and supported this project. Their collaboration and commitment

were essential to the success of this study.

8.References

A'Bear, A.D., Jones, T.H., Kandeler, E., Boddy, L., 2014. Interactive effects of temperature

and soil moisture on fungal-mediated wood decomposition and extracellular enzyme

activity. Soil Biology and Biochemistry 70, 151-158.

Adebayo, A.A., Enagbonma, B.J., Babalola, O.O., 2025. Comparative metagenomics on

community structure and diversity of rhizomicrobiome associated with monoculture and

soybean precedent carrot. Scientific Reports 15, 28161.

Arcand, M.M., Helgason, B.L., Lemke, R.L., 2016. Microbial crop residue decomposition

dynamics in organic and conventionally managed soils. Applied Soil Ecology 107, 347-359.

Aschi, A., Aubert, M., Riah-Anglet, W., Nélieu, S., Dubois, C., Akpa-Vinceslas, M.,

Trinsoutrot-Gattin, I., 2017. Introduction of Faba bean in crop rotation: Impacts on soil

chemical and biological characteristics. Applied Soil Ecology 120, 219-228.

Bak, G.-r., Lee, G.-j., Lee, J.-t., Jee, S.-n., 2022. Crop rotation affects biological properties

of rhizosphere soil and productivity of Kimchi cabbage (Brassica rapa ssp. pekinensis)

compared to monoculture. Horticulture, Environment, and Biotechnology 63, 613-625.

Bay, G., Lee, C., Chen, C., Mahal, N.K., Castellano, M.J., Hofmockel, K.S., Halverson, L.J.,

2021. Agricultural management affects the active rhizosphere bacterial community

composition and nitrification. Msystems 6, 10.1128/msystems. 00651-00621.

Bhatnagar, J.M., Peay, K.G., Treseder, K.K., 2018. Litter chemistry influences

decomposition through activity of specific microbial functional guilds. Ecological

Monographs 88, 429-444.

Breza, L.C., Mooshammer, M., Bowles, T.M., Jin, V.L., Schmer, M.R., Thompson, B.,

Grandy, A.S., 2023. Complex crop rotations improve organic nitrogen cycling. Soil Biology

and Biochemistry 177, 108911.

Bruyant, P., Moënne-Loccoz, Y., Almario, J., 2024. Root-associated helotiales fungi:

overlooked players in plant nutrition. Soil Biology and Biochemistry 191, 109363.

Calderon, F.J., Jackson, L.E., Scow, K.M., Rolston, D.E., 2001. Short-term dynamics of

nitrogen, microbial activity, and phospholipid fatty acids after tillage. Soil Science Society

of America Journal 65, 118-126.

Cox, M.C., Qualset, C.O., Rains, D.W., 1985. Genetic variation for nitrogen assimilation

and translocation in wheat. II. Nitrogen assimilation in relation to grain yield and protein

1. Crop Science 25, 435-440.

Dang, K., Ma, Y., Liang, H., Fan, Z., Guo, S., Li, Z., Li, H., Zhang, S., 2024. Distinct planting

patterns exert legacy effects on the networks and assembly of root-associated

microbiomes in subsequent crops. Science of the Total Environment 946, 174276.

Eivazi, F., Tabatabai, M., 1988. Glucosidases and galactosidases in soils. Soil Biology and

Biochemistry 20, 601-606.

Freschet, G.T., Cornwell, W.K., Wardle, D.A., Elumeeva, T.G., Liu, W., Jackson, B.G.,

Onipchenko, V.G., Soudzilovskaia, N.A., Tao, J., Cornelissen, J.H., 2013. Linking litter

decomposition of above-and below-ground organs to plant–soil feedbacks worldwide.

Journal of Ecology 101, 943-952.

Gamage, A., Gangahagedara, R., Gamage, J., Jayasinghe, N., Kodikara, N., Suraweera, P.,

Merah, O., 2023. Role of organic farming for achieving sustainability in agriculture.

Farming System 1, 100005.

Gan, Y., Hamel, C., O'Donovan, J.T., Cutforth, H., Zentner, R.P., Campbell, C.A., Niu, Y.,

Poppy, L., 2015. Diversifying crop rotations with pulses enhances system productivity.

Scientific Reports 5, 14625.

Geisseler, D., Horwath, W.R., 2008. Regulation of extracellular protease activity in soil in

response to different sources and concentrations of nitrogen and carbon. Soil Biology and

Biochemistry 40, 3040-3048.

Greenfield, L.M., Hill, P.W., Seaton, F.M., Paterson, E., Baggs, E.M., Jones, D.L., 2020. Is

soluble protein mineralisation and protease activity in soil regulated by supply or

demand? Soil Biology and Biochemistry 150, 108007.

Group, C.A.S.C.C.S.C.W., Canada, N.R.C., Agriculture, C., Branch, A.-F.C.R., 1998. The

Canadian system of soil classification. NRC Research Press.

Guerrero-Ramírez, N.R., Mommer, L., Freschet, G.T., Iversen, C.M., McCormack, M.L.,

Kattge, J., Poorter, H., van Der Plas, F., Bergmann, J., Kuyper, T.W., 2021. Global root traits

(GRooT) database. Global Ecology and Biogeography 30, 25-37.

Hannula, S.E., Heinen, R., Huberty, M., Steinauer, K., De Long, J.R., Jongen, R., Bezemer,

T.M., 2021. Persistence of plant-mediated microbial soil legacy effects in soil and inside

roots. Nature Communications 12, 5686.

Hannula, S.E., Ma, H.k., Pérez-Jaramillo, J.E., Pineda, A., Bezemer, T.M., 2020. Structure

and ecological function of the soil microbiome affecting plant–soil feedbacks in the

presence of a soil-borne pathogen. Environmental microbiology 22, 660-676.

Herridge, D.F., Peoples, M.B., Boddey, R.M., 2008. Global inputs of biological nitrogen

fixation in agricultural systems. Plant and Soil 311, 1-18.

Hood-Nowotny, R., Umana, N.H.-N., Inselbacher, E., Oswald-Lachouani, P., Wanek, W.,

2010. Alternative methods for measuring inorganic, organic, and total dissolved nitrogen

in soil. Soil Science Society of America Journal 74, 1018-1027.

Huo, C., Luo, Y., Cheng, W., 2017. Rhizosphere priming effect: a meta-analysis. Soil Biology

and Biochemistry 111, 78-84.

Jan, M.T., Roberts, P., Tonheim, S.K., Jones, D.L., 2009. Protein breakdown represents a

major bottleneck in nitrogen cycling in grassland soils. Soil Biology and Biochemistry 41,

2272-2282.

Jensen, E.S., Peoples, M.B., Hauggaard-Nielsen, H., 2010. Faba bean in cropping systems.

Field crops research 115, 203-216.

Jesmin, T., Margenot, A.J., Mulvaney, R.L., 2022. A comprehensive method for casein-

based assay of soil protease activity. Communications in Soil Science and Plant Analysis

53, 507-520.

Jones, D.L., Nguyen, C., Finlay, R.D., 2009. Carbon flow in the rhizosphere: carbon trading

at the soil–root interface. Plant Soil 321, 5–33.

Kaloterakis, N., Giongo, A., Braun-Kiewnick, A., Rashtbari, M., Zamberlan, P., Razavi, B.S.,

Smalla, K., Reichel, R., Brüggemann, N., 2025. Rotational diversity shapes the bacterial

and archaeal communities and confers positive plant-soil feedback in winter wheat

rotations. Soil Biology and Biochemistry 203, 109729.

Killham, K., Rashid, M., 1986. Assay of activity of a soil deaminase. Plant and Soil 92, 15-

21.

Knops, J., Bradley, K., Wedin, D., 2002. Mechanisms of plant species impacts on ecosystem

nitrogen cycling. Ecology letters 5, 454-466.

Kuzyakov, Y., Xu, X., 2013. Competition between roots and microorganisms for nitrogen:

mechanisms and ecological relevance. New Phytologist 198, 656-669.

L'Espérance, E., Bouyoucef, L.S., Dozois, J.A., Yergeau, E., 2024. Tipping the plant-microbe

competition for nitrogen in agricultural soils. Iscience 27.

Lazicki, P., Geisseler, D., Lloyd, M., 2020. Nitrogen mineralization from organic

amendments is variable but predictable. Wiley Online Library.

Li, B., Liu, X., Zhu, D., Su, H., Guo, K., Sun, G., Li, X., Sun, L., 2023. Crop diversity promotes

the recovery of fungal communities in saline-alkali areas of the Western Songnen Plain.

Frontiers in microbiology 14, 1091117.

Li, J., Cooper, J.M., Lin, Z.a., Li, Y., Yang, X., Zhao, B., 2015. Soil microbial community

structure and function are significantly affected by long-term organic and mineral

fertilization regimes in the North China Plain. Applied Soil Ecology 96, 75-87.

Li, X., Panke-Buisse, K., Yao, X., Coleman-Derr, D., Ding, C., Wang, X., Ruan, H., 2020.

Peanut plant growth was altered by monocropping-associated microbial enrichment of

rhizosphere microbiome. Plant and Soil 446, 655-669.

Luo, G., Rensing, C., Chen, H., Liu, M., Wang, M., Guo, S., Ling, N., Shen, Q., 2018.

Deciphering the associations between soil microbial diversity and ecosystem

multifunctionality driven by long-term fertilization management. Functional Ecology 32,

1103-1116.

Lupwayi, N.Z., Soon, Y.K., 2015. Carbon and nitrogen release from legume crop residues

for three subsequent crops. Soil Science Society of America Journal 79, 1650-1659.

Lupwayi, N.Z., Zhang, Y., Hao, X., Thomas, B.W., Eastman, A.H., Schwinghamer, T.D., 2019.

Linking soil microbial biomass and enzyme activities to long-term manure applications

and their nonlinear legacy. Pedobiologia 74, 34-42.

Ma, H.-K., Pineda, A., Van der Wurff, A.W., Raaijmakers, C., Bezemer, T., 2017. Plant–soil

feedback effects on growth, defense and susceptibility to a soil-borne disease in a cut

flower crop: species and functional group effects. Frontiers in plant science 8, 2127.

Mariotte, P., Mehrabi, Z., Bezemer, T.M., De Deyn, G.B., Kulmatiski, A., Drigo, B., Veen,

G.C., Van der Heijden, M.G., Kardol, P., 2018. Plant–soil feedback: bridging natural and

agricultural sciences. Trends in ecology & evolution 33, 129-142.

Maynard, D., Kalra, Y., Crumbaugh, J., 1993. Nitrate and exchangeable ammonium

nitrogen. Soil sampling and methods of analysis 1, 25-38.

Murrell, E.G., Ray, S., Lemmon, M.E., Luthe, D.S., Kaye, J.P., 2020. Cover crop species

affect mycorrhizae-mediated nutrient uptake and pest resistance in maize. Renewable

Agriculture and Food Systems 35, 467-474.

Näsholm, T., Ekblad, A., Nordin, A., Giesler, R., Högberg, M., Högberg, P., 1998. Boreal

forest plants take up organic nitrogen. Nature 392, 914-916.

Nuruzzaman, M., Lambers, H., Bolland, M.D., Veneklaas, E.J., 2006. Distribution of

carboxylates and acid phosphatase and depletion of different phosphorus fractions in the

rhizosphere of a cereal and three grain legumes. Plant and Soil 281, 109-120.

Nuttall, J., O'leary, G., Panozzo, J., Walker, C., Barlow, K., Fitzgerald, G., 2017. Models of

grain quality in wheat—A review. Field crops research 202, 136-145.

Oliveira, M., Barre, P., Trindade, H., Virto, I., 2019. Different efficiencies of grain legumes

in crop rotations to improve soil aggregation and organic carbon in the short-term in a

sandy Cambisol. Soil and Tillage Research 186, 23-35.

Peoples, M.á., Brockwell, J., Herridge, D., Rochester, I., Alves, B., Urquiaga, S., Boddey, R.,

Dakora, F., Bhattarai, S., Maskey, S., 2009. The contributions of nitrogen-fixing crop

legumes to the productivity of agricultural systems. Symbiosis 48, 1-17.

Rhine, E., Mulvaney, R., Pratt, E., Sims, G.K., 1998. Improving the Berthelot reaction for

determining ammonium in soil extracts and water. Soil Science Society of America Journal

62, 473-480.

Schneider, T., Keiblinger, K.M., Schmid, E., Sterflinger-Gleixner, K., Ellersdorfer, G.,

Roschitzki, B., Richter, A., Eberl, L., Zechmeister-Boltenstern, S., Riedel, K., 2012. Who is

who in litter decomposition? Metaproteomics reveals major microbial players and their

biogeochemical functions. The ISME journal 6, 1749-1762.

Séneca, J., Söllinger, A., Herbold, C.W., Pjevac, P., Prommer, J., Verbruggen, E., Sigurdsson,

B.D., Peñuelas, J., Janssens, I.A., Urich, T., 2021. Increased microbial expression of organic

nitrogen cycling genes in long-term warmed grassland soils. ISME communications 1, 69.

Shamshitov, A., Kadžienė, G., Pini, F., Supronienė, S., 2025. The role of tillage practices in

wheat straw decomposition and shaping the associated microbial communities in

Endocalcaric–Epigleyic Cambisol soil. Biology and Fertility of Soils 61, 211-231.

Sharma, S., Kumar, T., Foulkes, M.J., Orford, S., Singh, A.M., Wingen, L.U., Karnam, V.,

Nair, L.S., Mandal, P.K., Griffiths, S., 2023. Nitrogen uptake and remobilization from pre-

and post-anthesis stages contribute towards grain yield and grain protein concentration

in wheat grown in limited nitrogen conditions. CABI Agriculture and Bioscience 4, 12.

Strickland, M.S., Rousk, J., 2010. Considering fungal:bacterial dominance in soils –

Methods, controls, and ecosystem implications. Soil Biology and Biochemistry 42, 1385-

1395.

Wang, G., van der Putten, W.H., Klironomos, J., Zhang, F., Zhang, J., 2025. Steering plant-

soil feedback for sustainable agriculture. Science 389, eads2506.

Wang, P., Nie, J., Yang, L., Zhao, J., Wang, X., Zhang, Y., Zang, H., Yang, Y., Zeng, Z., 2023a.

Plant growth stages covered the legacy effect of rotation systems on microbial

community structure and function in wheat rhizosphere. Environmental Science and

Pollution Research 30, 59632-59644.

Wang, Y., Li, Q., Li, C., 2023b. Organic fertilizer has a greater effect on soil microbial

community structure and carbon and nitrogen mineralization than planting pattern in

rainfed farmland of the Loess Plateau. Frontiers in Environmental Science 11, 1232527.

Wang, Y., Wu, P., Qiao, Y., Li, Y., Liu, S., Gao, C., Liu, C., Shao, J., Yu, H., Zhao, Z., 2023c.

The potential for soil C sequestration and N fixation under different planting patterns

depends on the carbon and nitrogen content and stability of soil aggregates. Science of

the Total Environment 897, 165430.

WRB, I.W.G., 2007. World Reference Base for Soil Ressources 2006, First Update, World

Soil Resources Report, Rome.

Yang, X., Xiong, J., Du, T., Ju, X., Gan, Y., Li, S., Xia, L., Shen, Y., Pacenka, S., Steenhuis, T.S.,

2024. Diversifying crop rotation increases food production, reduces net greenhouse gas

emissions and improves soil health. Nature Communications 15, 198.

Yixuan, C., Zhijie, D., Yu, W., Qiong, L., Ruohan, Y., Tida, G., Zhenke, Z., 2025. Faba bean

enhances soil multifunctionality through shaping rhizosphere microbial communities in

legume-cereal crop rotations. Frontiers of Agricultural Science & Engineering 12.

Yuan, J., Zhao, J., Wen, T., Zhao, M., Li, R., Goossens, P., Huang, Q., Bai, Y., Vivanco, J.M.,

Kowalchuk, G.A., 2018. Root exudates drive the soil-borne legacy of aboveground

pathogen infection. Microbiome 6, 156.

Zeilinger, S., Gupta, V.K., Dahms, T.E., Silva, R.N., Singh, H.B., Upadhyay, R.S., Gomes, E.V.,

Tsui, C.K.-M., Nayak S, C., 2016. Friends or foes? Emerging insights from fungal

interactions with plants. FEMS Microbiology Reviews 40, 182-207.

Zhalnina, K., Louie, K.B., Hao, Z., Mansoori, N., Da Rocha, U.N., Shi, S., Cho, H., Karaoz, U.,

Loqué, D., Bowen, B.P., 2018. Dynamic root exudate chemistry and microbial substrate

preferences drive patterns in rhizosphere microbial community assembly. Nature

Microbiology 3, 470-480.

Zhao, Z.-B., He, J.-Z., Quan, Z., Wu, C.-F., Sheng, R., Zhang, L.-M., Geisen, S., 2020.

Fertilization changes soil microbiome functioning, especially phagotrophic protists. Soil

Biology and Biochemistry 148, 107863.

Zhou, J., Kong, Y., Zhao, W., Wei, G., Wang, Q., Ma, L., Wang, T., Shu, F., Sha, W., 2021.

Comparison of bacterial and archaeal communities in two fertilizer doses and soil

compartments under continuous cultivation system of garlic. Plos one 16, e0250571.

Zhou, Y., Yang, Z., Liu, J., Li, X., Wang, X., Dai, C., Zhang, T., Carrión, V.J., Wei, Z., Cao, F.,

2023. Crop rotation and native microbiome inoculation restore soil capacity to suppress

a root disease. Nature communications 14, 8126.

Zhu, B., Gutknecht, J.L.M., Herman, D.J., Keck, D.C., Firestone, M.K., Cheng, W., 2014.

Rhizosphere priming effects on soil carbon and nitrogen mineralization. Soil Biology and

Biochemistry 76, 183-192.

Figure captions

Figure 1: Experimental design of Cloutier and Palmarolle field in 2024. Colors (yellow and

green) represent the previous crop sown in 2023. Brown dots line represents fertilized

plots in 2024, and clovers represent plots that received clover as a cover crop. The cover

crop was not used in our analysis, since it was not grown when we sampled mid-growing

season (2024). Our experimental design allowed us to test the effects of previous crop,

fertilization and wheat compartment on exoenzymes potential activities (Protease,

deaminase and B-glucosidase), on inorganic N content and on the microbial community

diversity and composition.

Figure 2: Beta diversity of the bacterial (16S) and fungal communities (ITS) from

rhizosphere and bulk soil under wheat cropping at Cloutier (A, B) and Palmarolle (C, D) in

2024 using a Euclidean distance matrix and principal coordinate analysis. The effect of the

previous crop (color) and organic fertilization (form and ellipse) is presented. For each

field, all blocks are represented (n=3). The R^2 and p-value correspond to the PERMANOVA

results for these variables and the compartment (i.e., rhizospheric and bulk soil) variable.

Figure 3: Compositions of the soil bacterial (A) and fungal (C) communities and Venn plot

illustrating the number of differentially abundant bacterial (B) and fungal (D) ASV from

rhizosphere and bulk soil under wheat cropping at Cloutier in 2024. Stacked bar charts

contain the relative abundance of phyla for bacteria (A) and Classes for fungi. Venn plots

illustrate the number of significantly differentially abundant ASVs found by DESeq2,

ANCOM-BC and LEfSe analysis (p<0.05).

Figure 4: Compositions of the soil bacterial (A) and fungal (C) communities and Venn plot

illustrating the number of differentially abundant bacterial (B) and fungal (D) ASV from

rhizosphere and bulk soil under wheat cropping at Palmarolle in 2024. Stacked bar charts

contain the relative abundance of phyla for bacteria (A) and Classes for fungi. Venn plots

illustrate the number of significantly differentially abundant ASVs found by DESeq2,

ANCOM-BC and LEfSe analysis (p<0.05).

Figure 5: Protease, β-glucosidase and deaminases potential activities in the rhizosphere

and bulk soil at 0-15 cm depth under wheat cropping at Cloutier (A, B, C) and Palmarolle

(D, E, F) mid-growing season in 2024. The effects of the soil compartment, the previous

crop, and fertilization are represented. The organic fertilization is represented by points

(mean) and error bars (95% confidence interval) of both compartments and the previous

crop. For each field and potential enzymatic activities, all blocks are represented (n=3).

Significant results from Kruskal-Wallis or ANOVA tests for each enzyme's potential

activities (see Table 2) are identified with a *(p<0.05).

Figure 6: Wheat rhizosphere and bulk soil ammonium and nitrate content mid-growing

season at Cloutier (A, B) and Palmarolle (C, D) at 0-15 cm depth. The effects of the

compartment, the previous crop, and fertilization are represented. The organic

fertilization is represented by points (mean) and error bars (95% confidence interval) of

both compartments and the previous crop. For each field and N content, all blocks are

represented (n=3). Significant results from Kruskal-Wallis or ANOVA tests for ammonium

and nitrate content (see Table 3) are identified with a *($p<0.05$).

Figure 7: Wheat grain protein content of each organic fertilization treatment and previous

crops after harvest at Cloutier. The effects of the previous crop and fertilization are

represented. Significant results from Kruskal-Wallis tests (see Table 3) are identified with

a **($p<0.05$).

974 **Figure 1.**

978 **Figure 2.**

Figure 3.

986   **Figure 4.**

Figure 5.

Figure 6.

997   **Figure 7.**

1000