Gene Editing of Caffeic-O-methyltransferase (COMT1) in the model grass Setaria viridis
Improves Biomass Saccharification Without Compromising Plant Growth or Abiotic Stress
Tolerance

Fernanda de Oliveira Menezes [a].; Karoline Estefani Duarte [a].; Gabriel Garon Carvalho [b].;
Leonardo Dario Gomez [c].; Igor Cesarino [b].; Wagner Rodrigo de Souza [a]*

[a] Federal University of ABC (UFABC), Center for Natural and Human Sciences. Alameda da
Universidade, s/n – Anchieta 09606-045, São Bernardo do Campo - SP, Brazil

[b] Department of Botany, Institute of Biosciences, University of São Paulo, Rua do Matão,
27 – Cidade Universitária 73091-7507. São Paulo, Brazil.

[c] Centre for Novel Agricultural Products (CNAP), Department of Biology, University of York,
Heslington YO10 5DD, York, UK

Gene Editing of Caffeic-O-methyltransferase (COMT1) in the model grass Setaria viridis
Improves Biomass Saccharification Without Compromising Plant Growth or Abiotic Stress
Tolerance

Highlights

CRISPR knockout of *COMT1* in *Setaria viridis* markedly enhances biomass digestibility by
reducing lignin monomers, without compromising plant growth or salt-stress tolerance.

Genome editing of *COMT1* in *Setaria viridis* reveals unexpected lignin plasticity, reducing S, G, H
and tricin units and improving saccharification efficiency without developmental penalties.

Combined thiacidolysis/GC-MS and metabolomics suggests a compensatory mechanism in the
lignin portion of the *S. viridis* cell wall.

Abstract

Second-generation bioethanol production is limited by the recalcitrance of lignocellulosic
biomass, largely driven by lignin content and composition. Genetic strategies targeting lignin
biosynthesis offer a promising alternative to improve biomass digestibility without relying on
energy-intensive pretreatments. Here, we used CRISPR/Cas9 editing to disrupt the caffeic-O-
methyltransferase gene (*COMT1*) in *Setaria viridis*, a C_4 monocot model related to major
bioenergy crops. *COMT1* was selected based on its strong expression in lignifying tissues and its
central role in the biosynthesis of syringyl (S) units, guaiacyl (G) units, and the flavone tricin, a
non-canonical lignin monomer in grasses. Edited plants showed no visible growth defects,
maintaining normal development and salt-stress tolerance. Chemical analyses revealed a drastic
reduction in S units and tricin, accompanied by moderate but significant decreases in G and *p*-
hydroxyphenyl (H) monomers. Despite these compositional changes, total soluble lignin content
was only slightly reduced, suggesting compensatory incorporation of non-canonical monomers.
Metabolome analysis of edited plants suggested a redirection of carbon flux towards
phenylpropanoid and flavonoid pathways. These modifications resulted in a substantial increase
in saccharification efficiency, demonstrating that *COMT1* disruption can enhance biomass
digestibility without compromising plant viability. Our findings highlight *COMT1* as a key target for
engineering improved feedstocks for bioenergy production.

Keywords

Seteria viridis, *COMT1*, CRISPR/Cas9, lignin biosynthesis, tricin, biomass recalcitrance,
saccharification efficiency, cell wall engineering, bioenergy crops

1. Background

As highlighted in the *World Energy Outlook* (U.S. Energy Information Administration, 2021), the energy sector accounts for nearly three-quarters of all anthropogenic greenhouse gas (GHG) emissions worldwide. This alarming statistic underscores the urgent need for transformative changes in how energy is produced and consumed to mitigate climate change and promote long-term ecological resilience effectively. Within this context, the development of sustainable energy systems has gained prominence as a strategic pathway to curb emissions. Among the proposed alternatives, second-generation (E2G) bioethanol derived from lignocellulosic biomass such as agricultural residues has attracted significant international attention as a renewable, low-emission substitute for fossil fuels (Reshmy *et al.*, 2023; Yılbaşı, 2025).

E2G is produced from lignocellulosic biomass derived mainly from agricultural residues such as sugarcane bagasse. The production process begins with a physical and/or chemical pretreatment aimed at reducing biomass recalcitrance by disrupting the lignocellulosic matrix. This step is essential to enhance biomass digestibility, as the rigid cell wall structure limits enzymatic accessibility. Following pretreatment, enzymatic hydrolysis converts structural carbohydrates into fermentable sugars, which are subsequently fermented to ethanol. Improving biomass digestibility remains a major technological bottleneck in E2G production, with a direct impact on process efficiency and economic viability (Liu *et al.*, 2019, Preprint; Kordala *et al.*, 2024, Preprint).

Lignocellulosic biomass primarily comprises three biopolymers that form plant cell walls: cellulose, hemicelluloses, and lignin (Ma *et al.*, 2019, Preprint; Shi *et al.*, 2019). The cellulose polymer forms a scaffold of microfibrils embedded in a matrix of hemicelluloses, which are interlinked with lignin. Lignin, a complex phenolic polymer mainly composed of *p*-hydroxyphenyl (H), guaiacyl (G), and syringyl (S) units, confers mechanical strength to the cell wall, facilitates water transport, and plays key roles in plant defense mechanisms against biotic and abiotic stressors. However, from a bioindustrial perspective, lignin represents a major barrier to biomass deconstruction. Its hydrophobic nature and cross-linking capacity restrict enzymatic access to polysaccharides, thereby reducing sugar release during hydrolysis and compromising fermentation efficiency (Broda *et al.*, 2022, Preprint; Abolore *et al.*, 2024, Preprint).

Although physicochemical pretreatments can partially mitigate biomass recalcitrance, they often require high energy input, increase processing costs, and may produce environmentally harmful byproducts (Basak *et al.*, 2023; Abolore *et al.*, 2024, Preprint). As an alternative, targeted genetic modifications of cell wall components have emerged as a promising strategy to enhance biomass digestibility (Damm *et al.*, 2016, Preprint; Rao and Dixon, 2018, Preprint). Reducing lignin content or altering its composition enhances the release of fermentable sugars from structural carbohydrates, representing a sustainable approach to improving biomass conversion efficiency and reducing second-generation bioethanol production costs (Mosier *et al.*, 2005; Tye *et al.*, 2016, Preprint). However, these genetic modifications often lead to growth penalties, such as reduced biomass yield and increased susceptibility to abiotic stresses, which limit their scalability.

Therefore, optimizing the balance between lignin modification, plant productivity, and stress
resilience is essential for advancing the next generation of bioenergy crops (De Vries *et al.*, 2018).

Currently, twelve enzyme families have been identified as key players in the biosynthetic pathway
towards canonical lignin monomers (Figure 1). The primary precursor phenylalanine is
deaminated into cinnamic acid, which is converted into distinct hydroxycinnamyl alcohols
(generally known as monolignols) via hydroxylations/methylations of the aromatic ring and
reduction of the propane tail. Multiple routes have been described to contribute to the biosynthesis
of monolignols in different plant species, including a tyrosine by-pass via tyrosine ammonia-lyase
activity (Barros *et al.*, 2016), the "acid pathway" via bifunctional ascorbate peroxidase activity
(Barros *et al.*, 2019), and the highly evolutionarily conserved "shikimate shunt" (Dixon and Barros,
2019, Preprint). The complexity of lignin biosynthesis drastically increases when considering
lignin monomers beyond the canonical monolignols, as 49 metabolites have been detected to be
incorporated into lignin polymers in different cellular contexts and plant species (Pesquet *et al.*,
2025, Preprint) . Among key lignin biosynthetic enzymes, Caffeic Acid O-Methyltransferase
(COMT, E.C. 2.1.1.68) is a *S*-adenosyl-L-methionine-dependent *O*-methyltransferase involved in
multiple steps of the pathway. Originally, COMT was thought to catalyze the methylation of caffeic
and 5-hydroxyferulic acid, but latter studies demonstrated that this enzyme preferentially
methylate 5-hydroxyconiferaldehyde and/or 5-hydroxyconiferyl alcohol to produce sinapaldehyde
and/or sinapyl alcohol, respectively (Humphreys J.M and Chapple C, 2002) . Thus, together with
Ferulic Acid 5´-Hydroxylase (F5H), COMT constitutes the specific pathway towards the
biosynthesis of syringyl lignin units. More recently, COMT-mediated methylation of caffeic acid
into ferulic acid was demonstrated to be of great importance for lignification in grasses, being
proposed as the most favored route towards G and S units in this group of plants (Barros *et al.*,
2019). Additionally, COMT was demonstrated to play dual roles in the lignin pathway in grasses,
as it catalyzes not only the methylation steps within the S-lignin branch but also the methylation
steps towards the biosynthesis of the flavone tricin (Eudes *et al.*, 2017) , an authentic lignin
monomer in grasses (Lan *et al.*, 2015) . This enzyme also plays an important role in responses
to abiotic stresses, such as drought and salinity, through the synthesis of melatonin. COMT
catalyzes the conversion of N-acetyl-5-hydroxytryptamine into melatonin, an important regulator
of oxidative stress and redox balance in plants, helping to protect cells against damage caused
by abiotic stress (Li *et al.*, 2022). Due to its key position in the lignin biosynthetic pathway, the
COMT gene is considered a suitable target for regulating lignin monomers aimed at reducing
biomass recalcitrance.

Studies involving the negative regulation of the *COMT* gene using RNA interference (*RNAi*)
techniques in plants of *Brachypodium distachyon* (Ho-Yue-Kuang *et al.*, 2016), *Panicum virgatum*
(Wu et al., 2018), and *Hordeum vulgare* (Daly *et al.*, 2019) observed significant changes in lignin
composition and biomass digestibility. However, the negative regulation of *COMT* in *Zea mays*
and *Sorghum bicolor* plants resulted in negative effects on biomass yield (Miller *et al.*, 1983;
Sattler *et al.*, 2010, Preprint). In sugarcane (*Saccharum* sp.), Jung et al. (2012) silenced the
COMT gene, addressing biomass digestibility. Although these plants showed increased biomass

saccharification, there was also a decrease in productivity. In a more recent study, Kannan et al. (2018) used TALEN-mediated mutagenesis for the knockout of 100 copies/alleles of the *COMT* gene in sugarcane increasing saccharification without compromising biomass.

Setaria viridis is an emerging monocot model plant for molecular and genetic research. This short, fast-growing, C$_4$ species has a fully sequenced genome (Bennetzen *et al.*, 2012). It belongs to the *Panicoideae* subfamily, which includes the bioenergy crops *Sorghum bicolor, Zea mays, and Saccharum sp.* In a study conducted by Ferreira et al. (2019), six *COMT* genes were identified in the *Setaria viridis* genome. According to this study, the *COMT1 and COMT6* genes showed phylogenetic proximity to the *COMT* genes involved in lignification in *Brachypodium distachyon, Zea mays, Oryza sativa,* and *Sorghum bicolor* (Piquemal *et al.*, 2002*a*; Saballos *et al.*, 2008; Koshiba *et al.*, 2013; Trabucco *et al.*, 2013*a*; Ho-Yue-Kuang *et al.*, 2016). Furthermore, it was observed that most of the *COMT* genes showed almost null expression in the elongated internode of *Setaria viridis,* except for *COMT1,* which demonstrated strong and preferential expression in the transition and maturation zones Ferreira et al. (2019) This work shows that *COMT1* is the main candidate gene to play a crucial role in lignification during development in *Setaria viridis.*

Here, we demonstrate that CRISPR/Cas9 knockout of the *COMT1* gene in *Setaria viridis* represents a promising strategy to enhance biomass digestibility. The targeted genome editing significantly improved saccharification efficiency without compromising overall plant viability. Notably, despite substantial alterations in lignin composition, the mutant plants maintained normal growth and did not exhibit increased sensitivity to salt stress. These findings underscore the feasibility of engineering lignin pathways without detrimental effects on plant development.

2.   Materials and methods

2.1.   Strategies for *COMT1* gene editing in *Setaria viridis* using CRISPR/Cas9

The sequence of the target gene, *COMT1* from *Setaria viridis*, was retrieved from the public database Phytozome v13 (https://phytozome-next.jgi.doe.gov/). The genome version used was *Setaria viridis* v4.1, with the corresponding *COMT1* gene ID Sevir. 6G053300. CRISPRPLANT v.2 software (Minkenberg *et al.*, 2019) was utilized to identify the sgRNAs corresponding to the selected gene-editing target, focusing on the region immediately upstream of the PAM sequence (5′-NGG), specific to each gene region to be deleted. The software also enabled the analysis of off-target regions for the selected sgRNAs, and target sites with low probabilities of mis-targeting were chosen for gene editing. The sequences of the sgRNAs for *COMT1* are available in Supplementary Table 1.

After identifying the sgRNAs, we proceeded with the construction of vectors for the genetic transformation of *S. viridis*. The selected sequences were cloned into a CRISPR vector (E856; p6i2xoR-UcasW-U6Os4), with codon optimization for monocot species. Cloning was performed by DNA Cloning Service (Hamburg, Germany). The schematic representation of the optimized

CRISPR/Cas9 vector is shown in Supplementary Figure 1A, and a schematic representation of the sgRNA target sites in the *COMT1* gene is shown in Supplementary Figure 1B.

2.2. Genetic transformation of *Setaria viridis* and CRISPR/Cas screening

The CRISPR vector was introduced into *Agrobacterium tumefaciens* (strain EAH105α) via heat shock, and genetic transformation of *S. viridis* A10.1 was carried out by co-inoculating embryogenic callus with *A. tumefaciens*, following the protocol adapted from (Martins *et al.*, 2015). Transformed plants were confirmed by PCR using primers specific to the amplification of the *hptII* (244 bp) and *Cas9* (500 bp) genes. Primers used to confirm the transformed events are listed in Supplementary Table 2. To detect insertions or deletions (indels) in the target sites, PCR amplification was performed to generate gene fragments spanning DSB sites (gRNA regions), followed by PCR product purification and Sanger sequencing to validate potential mutations. Chromatograms from sequencing were visualized using BioEdit® software (version 7.2.5). For mutation analysis, we utilized SYNTHEGO software, specifically the ICE (Inference of CRISPR Edits) Analysis tool, as described by (Hsiau *et al.*, 2018, Preprint) (available at https://ice.synthego.com/#/). Samples with R^2 values below 0.8 were excluded due to the lower reliability of the results. In addition to R^2, we also considered the percentages of indels formed, the nature of these indels, and the modifications induced by mutations in the chromatogram. Thus, our analysis included both the percentage of indels and the knockout score.

2.3. Plant material and growth conditions

Independent lines with frameshift mutations were propagated up to the T_2 generation. They were cultivated in a substrate mixed with vermiculite at a 3:1 ratio, in a controlled environment at 28 ± 2 °C, with a photoperiod of 16 hours of light and 8 hours of darkness. Irrigation was performed six times a week, maintaining field capacity, while a nutrient solution containing 1 mL of Plant Magic Plus Bio Silicon® per liter of water was applied weekly. The fertilizer composition included 11.7% potassium oxide, 14.3% water-soluble silicon, and 0.05% water-soluble EDTA-chelated iron. Cell wall characterization was conducted with sampling during the senescence phase, when the plants were naturally aging. Plant material (leaves and stems) was processed into powder using a ball mill. Each mutant was treated as an individual biological replicate, and four technical replicates were performed for subsequent analyses

2.4. Cell wall characterization

Independent lines with frameshift mutations were propagated up to the T_2 generation. They were cultivated in a substrate mixed with vermiculite at a 3:1 ratio, in a controlled environment at 28 ± 2 °C, with a photoperiod of 16 hours of light and 8 hours of darkness. Irrigation was performed six times a week, maintaining field capacity, while a nutrient solution containing 1 mL of Plant Magic Plus Bio Silicon® per liter of water was applied weekly. The fertilizer composition included 11.7% potassium oxide, 14.3% water-soluble silicon, and 0.05% water-soluble EDTA-chelated iron. Cell wall characterization was conducted with sampling during the senescence phase, when the plants were naturally aging. Plant material (leaves and stems) was processed into powder using a ball

199 mill. Each mutant was treated as an individual biological replicate, and four technical replicates
200 were performed for subsequent analyses.

2.5. Preparation of alcohol insoluble residue (AIR)

Alcohol-insoluble residue (AIR) was prepared as described by Möller et al. (2022), by adding 2
mL of water to 100 mg of powdered plant material in a 2 mL Eppendorf tube. The powder was
resuspended in the water and shaken by inverting the tube for 30 seconds. The plant material
was then pelleted by centrifugation at 4000 g for 10 minutes. The supernatant was removed and
replaced with 2 mL of 96% ethanol, followed by two additional ethanol washes. The resulting
residue was allowed to dry under ambient conditions.

2.6. Lignin quantification by acetyl bromide method

The determination of soluble lignin was carried out by the acetyl bromide method, as described
by Liu et al. (2016). Precisely 5 mg of AIR was added to a centrifuge tube with a screw cap, along
with 250 µL of acetyl bromide solution (25% acetyl bromide and 75% glacial acetic acid, v/v). The
mixture was heated at 50 °C for three hours. After cooling the samples to room temperature, the
resulting liquid was transferred to a 5 mL volumetric flask. Next, 1 mL of 2 M NaOH and 175 µL
of 0.5 M hydroxylamine hydrochloride (HCl) were added to the reaction. The final volume of the
reaction was adjusted precisely to 5 mL using glacial acetic acid. The absorbance of the
supernatant was measured on a spectrophotometer at 280 nm. The calculation of the total lignin
content in the dry sample was performed using the following formula, where the coefficient used
in the calculation was 17.75, specific for grasses (Foster et $al.$, 2010; Dwivedi et $al.$, 2024).
219

220 % Lignin = $\left(\dfrac{abs}{coeff \; X \; pathlength}\right) X \left(\dfrac{total \; volume \; X \; 100\%}{biomass \; weight}\right)$

2.7. Analysis of lignin composition by thioacidolysis method

Lignin composition analysis was also performed using the method described by Chen et al.
(2021), with some modifications. Briefly, 1 mL of thioacidolysis reagent (composed of 100 mL of
1,4-dioxane, 6.25 mL of boron trifluoride diethyl ether, and 25 mL of ethanethiol, in a v/v ratio) was
added to 2.5 mg of AIR. 4,4-Ethylidenebisphenol was added as an internal standard at a final
concentration of 12 µg/mL. The reaction was incubated for 4 hours at 100 °C. After cooling, 400
µL of the solution was transferred to glass vials, and 190 µL of 0.4 M NaHCO$_3$ were added. The
reaction mixture was incubated overnight at 55 °C. The resulting pellet was then homogenized in
100 µL of derivatization reagent (1:1 pyridine and BSTFA with 1% TCM v/v) at 55 °C for 30 minutes
prior to GC-MS analysis. For tricin analysis, 400 µL of the thioacidolysis reaction was transferred
to 1.5 mL tubes, followed by the addition of 190 µL of 0.4 M NaHCO3. This mixture was incubated
overnight at 55 °C. The resulting pellet was homogenized in a 1:1 (v/v) methanol:DMSO solution.
Samples were sonicated for 10 seconds, centrifuged at 22,000 g for 10 minutes, and 100 µL of

the supernatant was transferred to vials for HPLC analysis. An Agilent Zorbax Eclipse Plus C18 column (2.1 mm × 50 mm, 1.8 μm) was used to separate the compounds. The mobile phases used were: (A) 0.1% formic acid in water (v/v) and (B) 0.1% formic acid in acetonitrile (v/v). The column thermostat was maintained at 30 °C with a solvent flow rate of 0.45 mL/min. The elution gradients were as follows: initially 6% (B) for three minutes, followed by an increase from 6% to 95% (B) over 5 minutes. The mobile phase was maintained at 95% (B) for two minutes, followed by a return from 95% to 6% (B) in one minute, and a final phase of two minutes at 6% (B). The total run time was 13 minutes.

2.8. Determination of total *p*-CA and FA by alkaline hydrolysis

The extraction of *p*-coumaric acid (*p*-CA) and ferulic acid (FA) followed the protocol established by Möller et al. (2022). Approximately 10 mg of AIR were incubated with 1M NaOH at 30 °C for 12 hours. The reaction was then neutralized with 99% trifluoroacetic acid (TFA) and subjected to a liquid-liquid extraction using 700 μL of butanol. After the evaporation of butanol, the resulting residue was resuspended in a 50% methanol solution and analyzed by high-performance liquid chromatography (HPLC) with UV detection. The injection volume was 10 μL. Component separation was performed using an XBridge BEH Shield RP18 chromatographic column (Waters, 186003149) with a porosity of 130 Å, a particle size of 3.5 μm, and dimensions of 1 mm × 100 mm. The flow rate was set to 1.23 mL/min, and a binary gradient of 0.1% acetic acid and methanol was applied. The gradient program increased the methanol concentration from 5% to 70% over 0–30 minutes, then from 70% to 90% between 30 and 35 minutes and finally returned it to 5% between 35 and 40 minutes. Standard curves were generated using commercial *p*-CA and FA standards (Sigma) over a concentration range of 0.02–3.28 μg/mL for *p*-CA and 0.02–3.88 μg/mL for FA.

2.9. Determination of *p*-CA-arabinose and FA-arabinose by mild acidolysis

The quantification of *p*-coumaric acid and ferulic acid bound to arabinose, *p*-CA-ara and Fa-ara, respectively, was carried out following the protocol described by Lapierre et al. (2018). Briefly, 1 mL of the acidolysis reagent (composed of dioxane, methanol, and 2M aqueous hydrochloric acid in a 60:30:10 (v/v/v) ratio) was added to 15 mg of AIR in a screw-cap vial. The incubation was carried out for three hours at 80 °C with shaking at 163 *rpm*. After cooling, the reaction was stopped by the addition of 3 mL of water. The resulting diluted mixture was subjected to two extractions with 1 mL of ethyl acetate, followed by the evaporation of the ethyl acetate and dissolution of the pellet in 100 μL of methanol. The analysis was performed by HPLC, using a binary gradient of 0.1% acetic acid and methanol, with an injection volume of 10 μL. Separation of the sample components was achieved using an XBridge BEH Shield RP18 chromatographic column (Waters - 186003149). The gradient was programmed as follows: for the first 0-60 minutes, the methanol concentration was increased from 5% to 65%; from 60-65 minutes, the methanol concentration was increased from 65% to 90%; and from 65-70 minutes, the methanol concentration was returned to 5%.

2.10. Monosaccharide analysis

The analysis of the monosaccharide composition of the AIR fraction was performed through the hydrolysis of polysaccharides into monosaccharides. For this, 4 mg of AIR were hydrolyzed in 0.5 mL of 2 M trifluoroacetic acid (TFA) for 4 hours at 100 °C. After cooling to room temperature, the solution was completely evaporated using a centrifugal evaporator. The resulting pellet was washed twice with 2-propanol and evaporated. Subsequently, the material was resuspended in 200 µL of distilled water, and the suspension was filtered through a 0.45 µm PTFE filter. The analysis was carried out by HPAEC using a Dionex Carbopac PA-10 column with pulsed amperometric detection, as described by Möller et al. (2022) and Wannitikul et al. (2024). The quantification of the isolated monosaccharides was performed using an external calibration curve containing nine monosaccharide standards at concentrations of 250, 500, and 700 µM (glucose, fucose, arabinose, galactose, mannose, rhamnose, xylose, galacturonic acid, glucuronic acid).

2.11. Determination of cellulose content

The quantification of cellulose was carried out according to the methodology outlined by Acevedo et al. (2019). The TFA pellets, obtained as detailed in the previous section, were subjected to a washing procedure with 1.5 mL of water, followed by three additional washes with 1.5 mL of acetone. Subsequently, the samples were air-dried at room temperature overnight. For complete hydrolysis, 90 µL of 72% (w/v) sulfuric acid were added to the dry pellets. Then, 1.89 mL of water was added, and the mixture was heated at 120 °C for 4 hours. The glucose content in the resulting supernatant was determined using the Anthrone colorimetric assay (Morse, 1947) , with a glucose standard curve for quantification. The analysis was performed at a final volume of 1.2 mL, using 40 µL of sample and 800 µL of Anthrone (2 mg/mL). The blue color developed after incubation at 80 °C for 30 minutes. Standard reactions with glucose concentrations ranging from 0.5 µg/mL to 10 µg/mL were included for calibration purposes. The absorbance of the reaction was measured at 620 nm.

2.12. Biomass enzymatic hydrolysis (saccharification)

Enzymatic saccharification was carried out according to the procedures outlined by (Gomez *et al.*, 2010). Two types of biomasses were used in the saccharification process: untreated biomass and pretreated biomass. Biomass consisted of whole aerial parts of the plants (leaves + stem). For the pretreatment, 0.5 mL of 500 mM NaOH was added to 100 mg of biomass, and the reaction was heated to 90 °C for 30 minutes with occasional stirring. The material was then decanted to the bottom of the tube. Subsequently, it was washed three times with 1 mL of 25 mM sodium acetate buffer at pH 4.5. For enzymatic saccharification, an enzyme cocktail with a 4:1 ratio of Celluclast to Novozyme 188 was used, where Novozyme 188 acted as a cellulase and was derived from *Aspergillus niger* (both enzymes were obtained from Novozymes, Bagsvaerd, Denmark). Before use, the enzymes were filtered using a Hi-Trap desalting column (GE Healthcare, Little Chalfont, Buckinghamshire, UK). The hydrolysis process was carried out in a

25 mM sodium acetate buffer (pH 4.5) at 50 °C for specified time intervals. The determination of
sugars released after hydrolysis was performed using a modified version of the method introduced
by (Anthon and Barrett, 2002), incorporating 3-methyl-2-benzothiazolinone hydrazone (MBTH).
The analysis was conducted in a final volume of 1 mL, with 300 µL of the sample used for the
assay. The detection reaction consisted of 0.25 N NaOH, 3 mg/mL MBTH, and 1 mg/mL
dithiothreitol (DTT). Color development occurred after incubation at 70 °C for 20 minutes,
achieved by adding the oxidizing reagent (0.2% FeNH4(SO4)2, 0.2% sulfamic acid, and 0.1%
HCl). Standard reactions with glucose concentrations of 10, 20, 50, and 100 nmol were included
for calibration purposes. The absorbance of the reaction was measured at 620 nm. This method
was validated to detect a variety of sugars released from the cell wall and demonstrated sensitive
detection of several monosaccharides (Gomez *et al.*, 2010).

2.13. Untargeted metabolome profiling

For metabolome analysis, frozen aerial parts were milled with a Grindomix GM200 using
precooled stainless-steel jars (Retsch). Metabolites were extracted using 5% acetonitrile, adding
a necessary volume to obtain a concentration of 1 mg/mL. The samples were homogenized in
vortex for 10 min. at 6720 x *g*. The supernatant was collected and filtered in microfilter PTFE 0.22
µm. The filtered samples were injected (5 µL) on a UHPLC system (Waters ACQUITY UPLC)
equipped with a HSS T3 column (100 mm by 2.1 mm, 1.8 µm; Waters) and coupled to a Waters
Synapt XS qToF (quadrupole time-of-flight) hybrid (Waters) mass spectrometer. A gradient of two
buffers (A and B) was utilized: buffer A (0.1 % :formic acid, v/v) and buffer B (99:1:0.1
acetonitrile:water:formic acid, pH 3, v/v), as follows: 95% A, 5% B for 3 min, decreased to 50% A
and 50% B in 15 min, decreased to 100% B in 18 min and 95% A, 5% B in 20 min , flow rate was
0.280 mL per min. LockSpray ion source was used in negative electrospray ionization mode under
the following specific conditions: capillary voltage, 2.50 kV; source temperature, 120°C;
desolvation gas temperature, 500°C; desolvation gas flow, 1000 L per h; and cone gas flow, 100
L/h. The mass range was set from 50 to 1200 Da. For DDA-MSMS, the low mass ramp was
ramped between 15 and 50 eV. The peak selection was performed when a peak was > 1% of
baseline peak spectra. Data processing was performed with Progenesis QI software version 2.4
(Waters). All peaks (mass-to-charge ratio [m/z] features) were integrated in the chromatograms
of WT and *SvCOMT* Ev1, Ev2 and Ev3 plants. The following filters were applied to analyze the
data: average ion intensity within at least one group >100, P value Student's t-test < 0.001, and
fold change >2 or <0.5 compared to the respective control (WT). Peaks that appeared in the blank
(30 blank samples) were also excluded from analysis.

2.14. Salt stress assay in hydroponic system

345 Initially, the seeds were germinated on Petri dishes containing Murashige and Skoog medium with
346 the pH adjusted to 5.8. After germination, the plants were transferred to a hydroponic system and
347 maintained in Hoagland medium (Phytotech Labs, Product ID H353) for 20 days, until the stage
348 just before flowering onset. After the initial growth period, the salinity stress experiment was
349 initiated with three treatments: 0 mM (control), 100 mM, and 250 mM NaCl. The pH of the medium

was maintained at 5.8, and the treatment solutions were adjusted according to the specified NaCl concentrations. The growth environment was controlled to maintain a photoperiod of 16 hours of light and 8 hours of darkness, with a constant temperature of 26 °C during both day and night. The nutrient solutions were renewed every 4 days to ensure the maintenance of the desired nutrient and salt concentrations. Plants were collected in quadruplicate after 5 days and 20 days of treatment exposure.

2.15.   Determination of relative water content and dry biomass

The relative water content (RWC) of control and stressed plants was determined with 0.2 g of fresh leaves (fresh weight; FW) that were immediately rehydrated in distilled water for 24 h in the dark, when the turgid weight (TW) was recorded. Next, leaves were transferred to an oven set at 60 °C and kept until reaching a constant dry weight (DW). Leaf relative water content was calculated using the following equation (Menezes *et al.*, 2020).

$\%RWC = 100 \times (FW - DW)/(TW - DW)$

In addition to RWC, the dry biomass of the plants was evaluated after harvesting. Initially, 0.2 g of dry biomass from the aerial part was weighed and then dried in an oven at 60 °C for 24 hours. After this period, the samples were reweighed to determine the final weight.

3.   Results
3.1.   CRISPR/Cas9 of *COMT1* editing resulted in plants with different frameshift mutations

The strategy for *COMT1* gene knockout was based on the non-homologous end joining (NHEJ) DNA repair pathway, which repairs double-stranded DNA breaks (DSBs) generated by nuclease activity through error-prone ligation. Four sgRNAs were designed to direct the CRISPR/Cas9 system to the *COMT1* gene in *Setaria viridis*. The first three (sgRNA1, sgRNA2, and sgRNA3) targeted nucleotides 653–752 within exon 1, a critical coding region essential for gene function, whereas sgRNA4 was designed to target position 1771 in exon 2. This multi-site targeting strategy was intended to maximize knockout efficiency by inducing DSBs at key functional regions of the gene (Supplementary Figure 1B).

Agrobacterium-mediated transformation generated multiple independent events, confirmed by PCR and Sanger sequencing. Three independent T_2 lines (Ev.1, Ev.2, and Ev.3) carrying frameshift mutations induced by sgRNAs 1, 2, and 3 were selected for further analysis (Supplementary Figure 2 to 4 and Supplementary Table 3). The selected lines exhibited knockout-scores above 70%, as determined by Synthego ICE analysis. This score represents the proportion of indels predicted to cause functional knockouts—specifically frameshift mutations, indels ≥21 nucleotides, or edits likely to disrupt protein function. Unlike total indel frequency, which includes all insertions and deletions differing from the wild-type sequence, the knockout-score reflects only those mutations expected to abolish gene function. Scores above 50% are generally associated with biallelic mutations (Synthego, 2018; Conant et al., 2022). Based on these criteria, we

selected three homozygous lines with distinct knockout-scores (Ev.1, 91%; Ev.2, 75%; and Ev.3, 88%) for subsequent biochemical and phenotypic characterization.

3.2. *COMT1* mutants lack the brown midrib phenotype and demonstrate normal growth and development

Reduced lignin content and/or altered lignin composition has been frequently associated with the so-called "brown midrib" phenotype in grasses, a characteristic reddish-brown coloration of leaf blade midribs (Sattler *et al.*, 2010, Preprint) . This phenotype has been classically applied to identify and isolate lignin mutants in grasses, given that all mutants arising by either spontaneous or chemical mutagenesis were shown to be associated with loss-of-function of lignin-related genes (Sattler *et al.*, 2010, Preprint; Tang *et al.*, 2014; Li *et al.*, 2015; Tetreault *et al.*, 2021). After generating knockout mutants for *COMT1* in *S. viridis*, we first assessed whether these plants presented the expected brown midrib phenotype. However, the *COMT1* mutants did not exhibit any obvious visible phenotypes. These plants did not show the presence of a brown central vein in the leaves (Figure 2A) nor significant changes in growth patterns (Figure 2B). After senescence, the mutant lines (Ev.1, Ev.2 and Ev.3) were harvested as described in Section 4. Subsequently, we analyzed the effects of the *COMT1* knockout on the cell wall and biomass saccharification of these plants. Since the individual plants exhibited different editing patterns (Supplementary Figures 2, 3 and 4), each mutant was treated as an independent biological replicate, with four technical replicates in the subsequent experiments.

3.3. Quantification of lignin content and analysis of its monomeric composition reveal differences in *COMT*-edited plants

To investigate the effects of the *COMT1* gene knockout on lignin content and composition, we first quantified the total soluble lignin content in the mutant lines and wild-type plants, following the method described by Liu et al. (2016) . A small but statistically significant reduction in total soluble lignin content was observed in the mutant plants, with a decrease ranging from 1.9% to 3.1% compared to the wild-type plants (Figure 3A). The lignin composition was also analyzed using the thioacidolysis method, as described by Chen et al. (2021). The results of this analysis reveal a significant reduction in the G, S, and H monomers in the edited plants compared to the wild-type (WT) plants. For the G monomer, the reduction ranged from 13.3% to 28.8% (Figure 3B). In the case of the S monomer, reductions were more pronounced, ranging from 81.9% to 76.6% (Figure 3C), while the H monomer presented a reduction between 44.1% and 52.5% (Figure 3D). Additionally, tricin, another important component of lignin in grasses, decreased by 44.1% to 51% (Figure 3E).

3.4. Total and p-CA-lignin contents are drastically reduced in edited plants

Subsequently, the total content of *p*-CA and FA in *Setaria viridis* was measured by HPLC after alkaline hydrolysis in both mutant and wild-type plants. The method used effectively releases nearly all HCAs that are ester-linked, both in arabinoxylan (AX) and lignin (Möller *et al.*, 2022). Our results revealed alterations in the *p*-CA levels in *COMT1* plants. Specifically, as shown in

Figure 4A, the total content of *p*-CA decreased by 45% to 51% compared to wild-type plants. Significant changes in FA were also observed in the Ev.2 and Ev.3 lines, with a modest increase of 6% and 4%, respectively. Due to the potential association of ester-linked *p*-CA with both AX and lignin, we employed the methodology described in Lapierre et al. (2018) to extract and quantify HCAs and their conjugates with arabinose (FA-Ara and *p*-CA-Ara) using mild acidolysis. This method allowed us to distinguish between HCAs associated with AX as their arabinose conjugates and those associated with lignin, by subtracting the values of *p*CA-Ara and FA-Ara obtained through mild acidolysis from the values of *p*-CA and FA obtained through alkaline hydrolysis. In this analysis, no significant differences were observed in the levels of *p*-CA and FA esterified to arabinose (*p*-CA-Ara and FA-Ara) in the *COMT1* knockout lines (Ev.1, Ev.2, and Ev.3) compared to the wild-type plants (Figure 4B). However, regarding the *p*-CA bound to lignin, we observed a significant reduction in the *COMT1* knockout plants, with values ranging from 45% to 51%. This decrease was also associated with a 6% and 4% increase in lignin-bound FA in lines #2 and #3, respectively (Figure 4C).

3.5. Cellulose content and structural monosaccharide composition did not change in mutants compared to wild type plants

The monosaccharide composition of cell wall polymers, determined by high-performance anion-exchange chromatography (HPAEC), showed no significant changes in response to *COMT1* gene knockout (Supplementary Figure 5A). Following this analysis, we evaluated cellulose content, which displayed a slight increase in *COMT1*-knockout plants compared to wild-type controls; however, this difference was not statistically significant (Supplementary Figure 5B). Collectively, these results suggest that the major cell wall modification in the edited plants is associated primarily with alterations in lignin composition.

3.6. The enzymatic hydrolysis of biomass revealed that loss of *COMT1* increases the saccharification

To assess the impact of *COMT1* knockout on biomass saccharification, enzymatic hydrolysis assays were performed. Biomass from mutant plants exhibited a significant increase in saccharification efficiency, ranging from 34% to 55% relative to wild-type plants, without pretreatment (Figure 5A). Following pretreatment with 500 mM NaOH, saccharification efficiency increased up to threefold in all lines, likely reaching a sugar release plateau. Although *COMT1* knockout plants showed a modest increase in saccharification compared to wild-type, statistical significance was observed only in line #3 (Figure 5B), with no significant differences detected in lines #1 and #2.

3.7. *COMT mutants present altered metabolic flow*

Lignin pathway perturbations frequently lead to transcriptional and metabolic reprogramming of the phenylpropanoid pathway, which has been observed for different target genes and plant species (Vanholme *et al.*, 2012; Saleme *et al.*, 2017; Ferreira *et al.*, 2022). Thus, we anticipated that loss of COMT activity results in a redirection of the carbon flux, especially through the

phenolic metabolism, given its involvement in different steps of this pathway. To gain insight into the consequences of *COMT1* mutation in the metabolic profile of *S. viridis*, we analyzed acetonitrile-soluble metabolites in aerial parts (leaves + stem) extracts using ultra-high-performance liquid chromatography-mass spectrometry (UHPLC-MS). We were able to detect 956 peaks [mass-to-charge ratio (*m/z*) features] with intensities ≥ 1% of the spectra baseline peak (Supplementary Table 4). Then we filtered these peaks based on blank samples, in which all peaks present in the blank samples were removed from analysis when their intensities were comparable to the plant samples, resulting in 160 peaks (Supplementary Table 5). Next, we focused on peaks present in all replicates of at least one sample type, with at least a twofold difference in intensity in *COMT1* mutants compared with controls, resulting in 22 peaks with higher and lower abundances in mutant plants (Table 1).

From the 22 peaks differentially abundant in *COMT1* mutants, identified 17 peaks (partially) structurally annotated based on tandem mass spectrometry (MS/MS) fragmentation spectra, using different database analysis. Most were phenolic or flavonoid compounds coupled to sugar moieties, such as hexoses and glucuronic acid, including caffeic acid-*O*-hexoside, kaempferol-3-*O*-caffeoyl glucoside, 2-glucosyloxy-4-methoxycinnamic acid, di-*O*-caffeoyl-hexoside, luteolin-3,7-diglucoside, quercetin-diglucoside, feruloyl-*O*-hexoside and caffeoyl-glucoside. Two compounds are putative G-G dimers coupled to a hexoside moiety, G(8–*O*–4)G-hexoside and G(8–*O*–4) ferulic acid di-(acetylhexoside), increased by two-fold in *COMT* mutants. We also observed one compound involved in phenylpropanoid biosynthesis directly linked to lignin, caffeic acid, increased by 2-fold in *COMT1* mutants. From these 22 peaks, only 4 were decreased in mutant plants, syringaresinol *O*-beta-D-glucoside (decreased by 2-fold), sinapoyl-feruloyl-glucoside (decreased by 6-fold), and 2 unidentified compounds, with *m/z* 577.1576 and 403.1044 and retention time 9.55 and 13.18, respectively. These results suggest that disruption of *COMT1* leads to rewiring of the phenolic metabolism in *S. viridis*.

3.8. *COMT1* knockout plants did not present altered sensitivity to moderate or severe salinity stress

Genetic modifications in the plant cell wall can result in pleiotropic effects, including altered responses to abiotic stresses such as salinity. To determine whether *COMT1* knockout influences plant tolerance to salt stress, a major environmental factor limiting growth and agricultural productivity, we evaluated *COMT1* knockout lines under different NaCl concentrations. Wild-type (WT) plants and knockout lines (Ev.1, Ev.2, and Ev.3) were exposed to 0, 100, and 250 mM NaCl for 5 and 21 days. These NaCL concentrations have been previously shown to cause stress responses in S. viridis (Duarte *et al.*, 2019). Representative phenotypes after 5 days across all treatments are shown in Figures 6A and 6B, while phenotypes after 21 days are presented in Supplementary Figure 6. At 250 mM NaCl, plants of all genotypes failed to survive beyond 9 days (Figure 6C). Conversely, plants submitted to 100 mM NaCl, despite a slight decrease in biomass compared to control (Figure 6B), did not present differences in relative water content (Figure 6A), suggesting a moderate tolerance to salt stress. As shown in Figure 6, this tolerance was not

502 affected by mutations in *COMT1*, as no significant differences in relative water content (RWC) or
503 dry mass were detected between WT and knockout lines after 5 days of 100 mM NaCl (Figures
504 6D and 6E) or 21 days under 100 mM NaCl (Supplementary Figure 6). These results indicate that
505 *COMT1* loss does not impair growth or salt tolerance, as knockout plants maintained RWC levels
506 comparable to WT under saline conditions.

4. Discussion

The conversion efficiency of lignocellulosic biomass into biofuels is inversely correlated with the
lignin content in feedstock plants, as demonstrated by Chen and Dixon (2007) and Studer et al.
(2011), among many other studies. This correlation arises from the recalcitrance imposed by
lignin in the cell wall, which limits enzyme access to structural polysaccharides. Consequently,
strategies aimed at reducing lignin content or modifying its structure have been extensively
explored to improve biofuel yields from lignocellulosic biomass (Dixon *et al.*, 2001; Fu *et al.*, 2011;
Baxter *et al.*, 2014). Different genetic engineering approaches have been employed to decrease
lignin content and alter its composition in different species. However, disrupting lignin
biosynthesis can lead to undesirable pleiotropic effects, such as reduced mechanical strength
and decreased stem rigidity. For instance, in *Arabidopsis thaliana*, mutants with a 50% reduction
in lignin content exhibit weaker stems (Jones *et al.*, 2001). Likewise, in tobacco, downregulation
of an *O-methyltransferase* reduces lignin levels but also negatively impacts plant growth (Pichon
et al., 2006).

It is currently known that COMT enzymes play a crucial role in lignin biosynthesis in angiosperms
by catalyzing the methylation of the hydroxyl group at the 5-position (5-OH) of intermediates in
the phenylpropanoid pathway (Bonawitz and Chapple, 2010, Preprint; Vanholme *et al.*, 2010).
This process is essential for the formation of sinapyl alcohol, the main precursor of the S lignin
units (Figure 1). Studies indicate that the primary *in vivo* substrates for COMTs are 5-
hydroxyconiferaldehyde and 5-hydroxyconiferyl alcohol, intermediates required to produce
sinapyl alcohol (Zubieta *et al.*, 2002; Louie *et al.*, 2010; Green *et al.*, 2014). In addition to the
methylation of S unit precursors, it is currently accepted that COMT is a bifunctional enzyme in
lignin biosynthesis, at least in grasses. In this context, COMT is also responsible for methylation
steps that lead to the production of tricin, a member of the flavone subclass of flavonoid
compounds, which is incorporated into the lignin polymer via 4-O-β-coupling and is considered
the first lignin non-canonical monomer outside the monolignol biosynthetic pathway (Del Río *et*
al., 2012; Lan *et al.,* 2016*a,b*).

In *Setaria viridis*, a study conducted by Ferreira et al. (2019) six *COMT* genes were identified in
its genome, all belonging to class II of the *O-methyltransferase* superfamily. These *COMTs* form
a distinct phylogenetic group within the monocots, with *COMT1* and *COMT6* showing significant
similarity to *COMT* genes involved in lignification in grasses such as *Brachypodium distachyon*,
maize, rice, and sorghum. The strong expression of *COMT1* in the transition and maturation
zones of the elongated internode in *S. viridis* suggests a central role in lignification, making it an

ideal target for genetic manipulation to reduce biomass recalcitrance and improve saccharification
efficiency. Therefore, we have chosen *SvCOMT1* as a target for the generation of CRISPR/Cas9
mutants in *S. viridis* to disrupt this gene.

The disruption of *COMT1* gene in *Setaria viridis* did not show any visible phenotype

In some C4 grasses, such as *Zea mays* and *Sorghum bicolor*, mutations in genes involved in
lignin metabolism, including *COMT*, result in the brown midrib phenotype (*bm* or *bmr*), which is
characterized by brown-colored central veins and, often, other parts of the plants. This phenotype
is associated with reduced lignin content and increased biomass digestibility (Vignols *et al.*, 1995;
Bout and Vermerris, 2003), and it is widely used to identify mutants affected in lignin deposition
(Sattler *et al.*, 2010, Preprint). Whereas the majority of the isolated mutants with this phenotype
have been characterized as affected in lignin biosynthesis, loss-of-function of genes encoding
enzymes that synthesize the methyl donor *S*-adenosyl-L-methionine, used by lignin *O*-
methyltransferases, was also shown to result in brown midribs in maize (Tang *et al.*, 2014; Li *et*
al., 2015). Additionally, the sorghum *bmr30* mutant has been characterized as disrupted in the
tricin biosynthetic gene encoding chalcone isomerase (Tetreault *et al.*, 2021). Until recently, brown
midribs have been described only in a few lignin mutants of panicoid C4 grasses such as maize
and sorghum, but disruption of some lignin biosynthetic genes in the C3 grass *Brachypodium*
distachyon, including *COMT*, also resulted in the accumulation of brown pigment in leaf midribs
and other lignified tissues (Bouvier D'Yvoire *et al.*, 2013; Trabucco *et al.*, 2013*b*). Conversely, this
phenotype has not been found in rice or barley, despite their extensive mutant stocks. Regarding
COMT, their corresponding mutants in maize and sorghum, as well as their silenced lines in
Brachypodium and switchgrass, showed the brown midrib phenotype ('Bout 2003; Piquemal *et*
al., 2002*b*; Trabucco *et al.*, 2013*b*; Wu *et al.*, 2019). In addition to the *bmr* phenotype, different
studies demonstrated that null *comt* mutant plants presented an abnormal growth (Yu *et al.*,
2025), while silenced *COMT* gene plants have reduced lignin content without growth penalties
(Vermerris *et al.*, 2007; Saballos *et al.*, 2008; Dien *et al.*, 2009; Fu *et al.*, 2011; Jung *et al.*, 2012,
2013; Lam *et al.*, 2019*a*). Here, the *COMT1* edited *S. viridis* plants did not display any visible
phenotype, neither the brown midrib trait, nor observable changes in plant growth (Figure 2). In
the context of generating optimized feedstock for bioeconomy, the lack of detrimental effects in
plant growth and development is a desirable trait. A similar phenotype was found when more than
100 *COMT* copies/alleles were gene-edited in sugarcane using TALEN technology (Kannan *et*
al., 2018), demonstrating that targeting *COMT* genes using gene editing might be an effective
strategy for biomass optimization without yield penalty.

COMT1 mutants display a drastic reduction of S monomers and a decrease of G, H and
tricin moieties in the lignin portion of _S. viridis_

Our results demonstrated a drastic reduction (from 40 to 80% reduction) in S units and tricin in
the AIR of edited *COMT1* plants, while G and H monomers demonstrated less pronounced effects
(Figure 4). It is expected that disruption of COMT activity leads to reduction of S units, as the
enzyme is mostly involved in the pathway steps that generates S monomers in grasses. The

novelty here is that tricin, G and H monomers were also significantly decreased, despite the total
soluble lignin content only slightly reduced (Figure 4A). These results may indicate that other
steps within the phenylpropanoid/flavonoid pathways might be compensating for lignin
biosynthesis through the incorporation of non-canonical monomers, to the extent that normal plant
development remains unaffected. The fact that total soluble lignin was only slightly decreased
may reinforce the speculation that more soluble, non-canonical monomers, could be incorporated
into the lignin.

Although the mechanisms underlying lignin homeostasis remain unclear, previous research
suggests that plants can preserve lignin deposition by incorporating non-canonical aromatic
monomers when canonical monolignols are limited. Reports of flavonoid-derived units,
hydroxycinnamate amides, and other atypical phenolics support the view that lignin is more
flexible than previously assumed, providing compensatory routes that may stabilize lignin levels
despite major pathway disruptions (Ralph et al., 2001b,a; Bonawitz and Chapple, 2010; Vanholme
et al., 2010). Also, it is worth mentioning that reduction in S, G and H monolignols may reflect
both a global down-regulation of phenylpropanoid flux through feedback inhibition (see Table 1)
or/and an altered lignin architecture with increased condensed linkages (C-C linkages instead of
β-O-4), which are poorly released by degradative methods.

Although there are studies that partially characterize lignin biosynthesis in plants such as
Arabidopsis, *Oryza sativa*, and *Zea mays*, there are still significant gaps in our understanding of
the dynamics of this process in grasses. For example, recent research indicates that many stages
of monolignol biosynthesis in grasses do not occur in the same way as in eudicots. In *Arabidopsis*,
loss-of-function mutations in genes such as *F5H* and *COMT* lead to a significant reduction in
sinapyl lignin units, resulting in a polymer composed solely of G units (Meyer *et al.*, 1998; Goujon
et al., 2003). However, in *Oryza sativa*, the loss of function of the *F5H* gene does not result in a
complete absence of S units, while in *COMT*-deficient mutants of maize and sorghum, S units are
significantly reduced but not completely absent. This suggests that grasses may have alternative
pathways to produce these units (Piquemal *et al.*, 2002*a*; Fornalé *et al.*, 2017; Takeda *et al.*,
2019). Our results on lignin monomers align with those previously reported in the literature for
Brachypodium distachyon, *Zea mays*, and *Hordeum vulgare* with *COMT* silencing (Piquemal *et*
al., 2002*a*; Barrière *et al.*, 2004; Daly *et al.*, 2019). However, these studies reported that the
reduction in S lignin content does not affect G monomers, which contrasts with the findings of the
present work. As mentioned above, in *S. viridis* the *COMT1* knockout resulted in significant
reductions in G lignin units (ranging from 13.3% to 28.8%), contrary to what was reported in those
studies. In alfalfa plants, the negative regulation of *COMT* also caused a decrease in G units and
an almost complete loss of S units (Guo *et al.*, 2001). While the formation of G lignin units
predominantly depends on CCoAOMT activity, *COMT* may play a secondary role by contributing
to the methylation of intermediates feeding into this pathway, such as coniferaldehyde. Thus, the
absence of *COMT* could reduce the efficiency of the G unit formation pathway, which explains the
decrease in G lignin units observed in our study. Moreover, the reduction in H units observed in
S. viridis (ranging from 44.1% to 52.5%) can be attributed to the overall modulation of the lignin

biosynthetic pathway, where disruption of methylation affects both the accumulation and utilization

of precursor monomers. These combined effects confirm that the knockout of the *COMT1* gene

not only drastically reduces S units but also impacts the synthesis of other lignin monomers.

COMT1 mutants demonstrate substantial differences in *p*CA pools

In grasses, lignin and xylan polymers can be substituted with hydroxycinnamic acids (HCAs),

such as ferulic acid (FA) and *p*-coumaric acid (*p*-CA). These HCA groups can be esterified or

etherified in lignin, playing a crucial role in modulating the structure and properties of the cell wall

(Hatfield *et al.*, 2017). In particular, the feruloylation of xylan facilitates the formation of covalent

cross-links between xylan chains or between xylan and lignin through the formation of a diferulate

(diFA) bridges. This cross-linking process contributes to the rigidity and mechanical strength of

the cell wall, as well as affecting biomass digestibility (Hatfield *et al.*, 2017; Terrett and Dupree,

2019, Preprint; Feijao *et al.*, 2022).

Recent studies suggest that, in grasses, the phenylpropanoid pathway is predominantly directed

toward the synthesis of hydroxycinnamic acids (HCAs) (Barros et al. 2022; Karlen et al. 2018;

Ralph 2010). This pattern can be attributed to the higher abundance of *p*-coumaric acid (*p*-CA)

and ferulic acid (FA) in the cell walls of grasses compared to dicotyledonous species. In this

context, *p*-CA plays a crucial role as a precursor in the biosynthesis of various phenylpropanoids

and flavonoids (Vogt, 2010) and is commonly conjugated to syringyl (S) and guaiacyl (G) lignin

monomers (Peracchi *et al.*, 2024, Preprint). Although the exact function of *p*-CA–lignin conjugates

is not fully understood (Hatfield et al. 2017), two main hypotheses have been proposed: (1) *p*-CA

may act as a radical transfer agent during the polymerization of S units (Ralph *et al.*, 2004), or (2)

p-CA-S monomers could terminate polymerization, resulting in a more linear lignin polymer and

consequently altering its structural properties (Kishimoto *et al.*, 2005; Hatfield *et al.*, 2017). In

addition to its conjugation with S lignin monomers, *p*-coumaric acid (*p*-CA) is also acylated to

arabinoxylans in the cell walls of grasses. This process is mediated by BAHD acyltransferases,

as demonstrated in species such as *Oryza sativa* (Bartley *et al.*, 2013) and *Setaria viridis* (de

Souza *et al.*, 2018; Mota *et al.*, 2021).

In the present study, using alkaline hydrolysis and mild acidolysis, we were able to quantify total

*p*CA and FA, in addition to *p*CA-Ara and FA-Ara, respectively. Further calculations allowed us to

estimate the amount of *p*CA and FA bound to lignin (Figure 4). We observed that there was a

large reduction of total *p*CA and *p*CA bound to lignin, while *p*CA-Ara did not show any significant

changes in *COMT1* mutants. These results indicate a substantial modulation in the *p*-CA pools,

demonstrating that the knockout of *COMT1* directly affects the incorporation of *p*-CA into cell

walls, thereby influencing lignin formation and structure. The reduction in total and lignin-bound

p-CA content observed in our study is consistent with previous reports investigating negative

regulation of *COMT* in maize (Pichon *et al.*, 2006) and *Brachypodium distachyon* (Ho-Yue-Kuang

et al., 2016), in which decreased levels of *p*-CA and S-type lignin monomers were also observed.

Moreover, our results reveal a slight increase in total FA content (ranging from 4% to 6%) in

COMT1 knockout plants. This increase may be associated with compensatory adjustments in the

phenylpropanoid biosynthesis caused by disruption of *COMT1*. Evidence of this mechanism has
already been reported in genetic modification studies involving the addition of HCAs in *Oryza*
sativa and *Setaria viridis* (de Souza *et al.*, 2018; Möller *et al.*, 2022). Therefore, we propose that
in *S. viridis* ferulic acid biosynthesis is not directly impacted by the knockout of *COMT1*, as its
production may proceed via an alternative pathway that bypasses caffeic acid methylation, as
suggested by previous studies (Barrière *et al.*, 2004; Chen *et al.*, 2006).

***COMT1* disruption increased biomass saccharification without altering cell wall**
polysaccharides

COMT1 knockout revealed no significant changes in the contents of structural polysaccharides
(Supplementary Figure 5A). This suggests that, while *COMT* plays a key role in lignin
biosynthesis, it does not directly affect cell wall polysaccharides. This result is supported by
studies in Arabidopsis and barley, in which loss-of-function of *COMT* did not lead to changes in
cellulose or structural carbohydrate content (Lee *et al.*, 2021). Conversely, the amounts of
arabinose and xylose released from hemicellulosic polymers were significantly increased in both
COMT downregulated and knockout rice plants (Lam *et al.*, 2019*b*), demonstrating different
results depending on the plant species. As expected for plants with diminished lignin contents,
the *COMT1* knockout plants showed a significant increase in biomass saccharification efficiency,
ranging from 34% to 55% compared to WT plants, without the necessity for pretreatments (Figure
5A). When subjected to pretreatment with 500 mM NaOH, saccharification increased up to
threefold, regardless of the genotype, potentially reaching a plateau in sugar release. However,
the *COMT1* knockout plants showed only a slight increase in saccharification efficiency compared
to the WT plants upon alkaline pretreatment, with no statistically significant differences (Figure
5B). The absence of significant differences after pretreatment may be explained by the
effectiveness of the pretreatment in breaking down the structural bonds of the cell wall, making
the biomass more accessible to enzymes and masking potential differences between the groups.
Our results are in line with previous studies that reported a negative correlation between lignin
content and the efficiency of hemicellulose enzymatic hydrolysis and fermentation in species such
as switchgrass, sugarcane, sorghum, and alfalfa (Chen and Dixon, 2007; Fu *et al.*, 2011; Xu *et*
al., 2011; Jung *et al.*, 2012).

Metabolome profiling of *COMT1* mutants suggests a redirection of carbon flux towards
phenylpropanoid and flavonoid pathways

Our metabolome analysis demonstrated that, as expected, *COMT1* disruption led to accumulation
of caffeic acid, a known (but not preferential) substrate of COMT enzymes (Table 1). However,
most tentatively annotated compounds accumulating in *COMT1* mutants were conjugated forms
of either flavonoids or phenylpropanoids, such as kaempferol-3-*O*-caffeoyl-glucoside (200-fold
increase), 2-glucosyloxy-4-methoxycinnamic acid (4.5-fold increase), caffeoyl-glucoside (44-fold
increase) and quercetin-diglucoside (2.6-fold increase). It is not surprising that most of the
accumulated metabolites found in edited plants were found as their sugar-conjugated forms,
given that several phenylpropanoid derivatives are either cytotoxic and/or bioactive and

glycosylation is known to play key roles in modulating phenylpropanoid properties (including
solubility, toxicity and bioactivity) and compartmentalization (Rates and Cesarino, 2023, Preprint)
. Our results are consistent with some recent studies showing that perturbations of lignin
metabolism in grasses lead to the accumulation of glycosylated phenylpropanoids (Barros *et al.*,
2022; Ferreira *et al.*, 2022; Oliveira *et al.*, 2025; Priya *et al.*, 2025, Preprint). Importantly, we
observed that selgin, the direct precursor of tricin biosynthesis and a known substrate for grass
COMTs, is highly accumulated in mutant plants (4.5-fold increase). In addition, luteolin, which is
also an intermediate of the tricin pathway and a COMT substrate, also accumulated, but in its
conjugated form as luteolin-3',7-diglucoside (2-fold increase). The fact that we did not find in our
metabolome many intermediates of the phenylpropanoid pathway suggests that *S. viridis* may
have other OMTs involved in lignin biosynthesis through this pathway, or the disruption of *COMT1*
is redirecting the carbon flux to flavonoid synthesis. Thus, our results further corroborate that
Setaria viridis COMT1 play an important role in the synthesis of tricin. The accumulated flavonoids
could also be responsible for the resilience of the mutants under salt stress observed in our work.

Salinity stress tolerance is not altered in *COMT1* mutants

Lignin plays a crucial role in plant defense against pathogens and in adaptation to stresses, such
as salinity. Given the central role of the *COMT* gene in lignin biosynthesis, we investigated how
its knockout affects plant responses to abiotic stresses, particularly salt stress. Soil salinization is
a major form of land degradation, characterized by the excessive accumulation of water-soluble
salts, especially sodium (Na^+) and chloride (Cl^-) ions (Tian *et al.*, 2021). Several studies have
demonstrated that the *COMT* gene regulates plant stress responses, including salinity (Sun *et al.*,
2020; Wang *et al.*, 2024), cold (Chang *et al.*, 2021), and drought (Yao *et al.*, 2022).

Interestingly, although previous studies have demonstrated that genetic modifications in the
COMT gene can influence plant responses to salinity, our results show that under moderate
salinity levels (100 mM), not only the WT but also genetically modified *Setaria viridis* plants are
able to maintain relative water content (RWC) in their tissues (Figure 6), suggesting a moderate
tolerance of *S. viridis* to salt stress. Under more severe salinity conditions (250 mM), a reduction
of 10.8% in RWC was observed only in line Ev.3, whereas the other edited lines did not show
significant changes. Moreover, plant survival was not affected compared to WT, as both WT and
COMT-knockout plants were able to survive for up to nine days under high salt concentrations.
Furthermore, dry biomass measurements (Figure 6) revealed no statistically significant
differences between WT and *COMT*-knockout plants at any of the tested NaCl concentrations,
both after 5 and 21 days of treatment. These findings suggest that *COMT* gene editing does not
significantly affect biomass accumulation under the salinity conditions evaluated.

In conclusion, this study highlights the potential of CRISPR/Cas9-mediated knockout of the
COMT1 gene as an effective strategy to alter lignin and improve biomass digestibility in grasses.
This has direct implications for enhancing saccharification efficiency and, consequently, biofuel
production, without increasing the plants' susceptibility to salt stress. Although the typical *bmr*
phenotype was not observed in the mutant lines, the changes in lignin composition and the

significant reduction in S, G, H, and tricin monomers reinforce the role of *COMT1* in modulating
lignin structural diversity in grasses. Moreover, the alterations in *p*-coumaric acid (*p*-CA) and
ferulic acid (FA) content, combined with metabolomics, suggest the activation of a compensatory
mechanism that maintains cell wall structural integrity under both normal and saline conditions.
In this context, the application of techniques such as nuclear magnetic resonance (NMR)
spectroscopy would be valuable for gaining deeper insights into how *COMT1* knockout-driven
modifications affect lignin structure, especially regarding incorporation of non-canonical
monomers. Such information could significantly contribute to the rational engineering of plant cell
walls without compromising growth. These findings support the notion that genetic manipulation
of lignin biosynthetic genes represents a promising approach for optimizing lignocellulosic
biomass conversion, with applications in bioenergy and animal nutrition, offering a meaningful
contribution toward the development of more efficient and sustainable renewable energy
technologies.

Author Contributions

Fernanda de Oliveira Menezes: methodology; investigation; formal analysis; data curation;
writing original draft; writing review & editing.
Karoline Estefani Duarte: supervision; writing review & editing.
Gabriel Garon Carvalho: formal analysis.
Leonardo Dario Gomez: methodology; supervision; writing review & editing.
Igor Cesarino: formal analysis; writing review & editing.
Wagner Rodrigo de Souza: conceptualization; supervision; project administration; funding
acquisition; writing review & editing.

Conflict of Interest

759 No conflict of interest declared.

Funding Statement

762 This research was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo
763 (Fapesp Proc. 2022/04419-5, 2019/26761-4 and 2019/04878-7).

5. References

Abolore RS, Jaiswal S, Jaiswal AK. 2024. Green and sustainable pretreatment methods for cellulose extraction from lignocellulosic biomass and its applications: A review. Elsevier Ltd.

Acevedo A, Simister R, McQueen-Mason SJ, Gómez LD. 2019. Sudangrass, an alternative lignocellulosic feedstock for bioenergy in Argentina. PLoS ONE 14.

Anthon GE, Barrett DM. 2002. Determination of reducing sugars with 3-methyl-2-benzothiazolinonehydrazone. Analytical Biochemistry 305, 287–289.

Barrière Y, Ralph J, Méchin V, Guillaumie S, Grabber JH, Argillier O, Chabbert B, Lapierre C. 2004. Genetic and molecular basis of grass cell wall biosynthesis and degradability. II. Lessons from brown-midrib mutants. Comptes Rendus - Biologies 327, 847–860.

Barros J, Escamilla-Trevino L, Song L, *et al.* 2019. 4-Coumarate 3-hydroxylase in the lignin biosynthesis pathway is a cytosolic ascorbate peroxidase. Nature Communications 10.

Barros J, Serrani-Yarce JC, Chen F, Baxter D, Venables BJ, Dixon RA. 2016. Role of bifunctional ammonia-lyase in grass cell wall biosynthesis. Nature Plants 2.

Barros J, Shrestha HK, Serrani-Yarce JC, Engle NL, Abraham PE, Tschaplinski TJ, Hettich RL, Dixon RA. 2022. Proteomic and metabolic disturbances in lignin-modified Brachypodium distachyon. Plant Cell 34, 3339–3363.

Bartley LE, Peck ML, Kim SR, *et al.* 2013. Overexpression of a BAHD acyltransferase, OsAt10, alters rice cell wall hydroxycinnamic acid content and saccharification. Plant Physiology 161, 1615–1633.

Basak B, Kumar R, Bharadwaj AVSLS, Kim TH, Kim JR, Jang M, Oh SE, Roh HS, Jeon BH. 2023. Advances in physicochemical pretreatment strategies for lignocellulose biomass and their effectiveness in bioconversion for biofuel production. Bioresource Technology 369.

Baxter HL, Mazarei M, Labbe N, *et al.* 2014. Two-year field analysis of reduced recalcitrance transgenic switchgrass. Plant Biotechnology Journal 12, 914–924.

Bennetzen JL, Schmutz J, Wang H, *et al.* 2012. Reference genome sequence of the model plant Setaria. Nature Biotechnology 30, 555–561.

Bonawitz ND, Chapple C. 2010. The genetics of lignin biosynthesis: Connecting genotype to phenotype.

Bout 2003. *in press*.

Bout S, Vermerris W. 2003. A candidate-gene approach to clone the sorghum Brown midrib gene encoding caffeic acid O-methyltransferase. Molecular Genetics and Genomics 269, 205–214.

Bouvier D'Yvoire M, Bouchabke-Coussa O, Voorend W, *et al.* 2013. Disrupting the cinnamyl alcohol dehydrogenase 1 gene (BdCAD1) leads to altered lignification and improved saccharification in Brachypodium distachyon. Plant Journal 73, 496–508.

Broda M, Yelle DJ, Serwańska K. 2022. Bioethanol Production from Lignocellulosic Biomass—Challenges and Solutions. MDPI.

Chang J, Guo Y, Yan J, *et al.* 2021. The role of watermelon caffeic acid O-methyltransferase (ClCOMT1) in melatonin biosynthesis and abiotic stress tolerance. Horticulture Research 8.

Chen F, Dixon RA. 2007. Lignin modification improves fermentable sugar yields for biofuel production. Nature Biotechnology 25, 759–761.

Chen F, Srinivasa Reddy MS, Temple S, Jackson L, Shadle G, Dixon RA. 2006. Multi-site genetic modulation of monolignol biosynthesis suggests new routes for formation of syringyl lignin and wallbound ferulic acid in alfalfa (Medicago sativa L.). Plant Journal 48, 113–124.

Chen F, Zhuo C, Xiao X, Pendergast TH, Devos KM. 2021. A rapid thioacidolysis method for biomass lignin composition and tricin analysis. Biotechnology for Biofuels 14.

Daly P, McClellan C, Maluk M, *et al.* 2019. RNAi-suppression of barley caffeic acid O-methyltransferase modifies lignin despite redundancy in the gene family. Plant Biotechnology Journal 17, 594–607.

Damm T, Commandeur U, Fischer R, Usadel B, Klose H. 2016. Improving the utilization of lignocellulosic biomass by polysaccharide modification. Elsevier Ltd.

Dien BS, Sarath G, Pedersen JF, Sattler SE, Chen H, Funnell-Harris DL, Nichols NN, Cotta MA. 2009. Improved sugar conversion and ethanol yield for forage sorghum (sorghum bicolor L. Moench) lines with reduced lignin contents. Bioenergy Research 2, 153–164.

Dixon RA, Barros J. 2019. Lignin biosynthesis: Old roads revisited and new roads explored. Royal Society Publishing.

Dixon RA, Chen F, Guo D, Parvathi K. 2001. *The biosynthesis of monolignols: a '"metabolic grid"', or independent pathways to guaiacyl and syringyl units?*

Duarte KE, de Souza WR, Santiago TR, Sampaio BL, Ribeiro AP, Cotta MG, da Cunha BADB, Marraccini PRR, Kobayashi AK, Molinari HBC. 2019. Identification and characterization of core abscisic acid (ABA) signaling components and their gene expression profile in response to abiotic stresses in Setaria viridis. Scientific Reports 9.

Dwivedi N, Yamamoto S, Zhao Y, Hou G, Bowling F, Tobimatsu Y, Liu CJ. 2024. Simultaneous suppression of lignin, tricin and wall-bound phenolic biosynthesis via the expression of monolignol 4-O-methyltransferases in rice. Plant Biotechnology Journal 22, 330–346.

Eudes A, Dutta T, Deng K, *et al.* 2017. SbCOMT (Bmr12) is involved in the biosynthesis of tricin-lignin in sorghum. PLoS ONE 12.

Feijao C, Morreel K, Anders N, Tryfona T, Busse-Wicher M, Kotake T, Boerjan W, Dupree P. 2022. Hydroxycinnamic acid-modified xylan side chains and their cross-linking products in rice cell walls are reduced in the Xylosyl arabinosyl substitution of xylan 1 mutant. Plant Journal 109, 1152–1167.

Ferreira SS, Goeminne G, Simoes MS, *et al.* 2022. Transcriptional and metabolic changes associated with internode development and reduced cinnamyl alcohol dehydrogenase activity in sorghum. Journal of Experimental Botany 73, 6307–6333.

Ferreira SS, Simões MS, Carvalho GG, de Lima LGA, Svartman RM de A, Cesarino I. 2019. The lignin toolbox of the model grass Setaria viridis. Plant Molecular Biology 101, 235–255.

Fornalé S, Rencoret J, García-Calvo L, Encina A, Rigau J, Gutiérrez A, Del Río JC, Caparros-Ruiz D. 2017. Changes in cell wall polymers and degradability in maize mutants lacking 30- and 50-o-methyltransferases involved in lignin biosynthesis. Plant and Cell Physiology 58, 240–255.

Foster CE, Martin TM, Pauly M. 2010. Comprehensive compositional analysis of plant cell walls (Lignocellulosic biomass) part I: Lignin. Journal of Visualized Experiments doi: 10.3791/1745.

Fu C, Mielenz JR, Xiao X, *et al.* 2011. Genetic manipulation of lignin reduces recalcitrance and improves ethanol production from switchgrass. Proceedings of the National Academy of Sciences of the United States of America 108, 3803–3808.

Gomez LD, Whitehead C, Barakate A, Halpin C, McQueen-Mason SJ. 2010. Automated saccharification assay for determination of digestibility in plant materials. Biotechnology for Biofuels 3.

Goujon T, Sibout R, Pollet B, *et al.* 2003. *A new Arabidopsis thaliana mutant deficient in the expression of O-methyltransferase impacts lignins and sinapoyl esters*.

Green AR, Lewis KM, Barr JT, Jones JP, Lu F, Ralph J, Vermerris W, Sattler SE, Kang C. 2014. Determination of the structure and catalytic mechanism of sorghum bicolor caffeic acid O-methyltransferase and the structural impact of three brown midrib12 mutations. Plant Physiology 165, 1440–1456.

Guo D, Chen F, Inoue K, Blount JW, Dixon RA. 2001. *Downregulation of Caffeic Acid 3-O-Methyltransferase and Caffeoyl CoA 3-O-Methyltransferase in Transgenic Alfalfa: Impacts on Lignin Structure and Implications for the Biosynthesis of G and S Lignin*.

Hatfield RD, Rancour DM, Marita JM. 2017. Grass cell walls: A story of cross-linking. Frontiers in Plant Science 7.

Ho-Yue-Kuang S, Alvarado C, Antelme S, *et al.* 2016. Mutation in Brachypodium caffeic acid O-methyltransferase 6 alters stem and grain lignins and improves straw saccharification without deteriorating grain quality. Journal of Experimental Botany 67, 227–237.

Hsiau T, Conant D, Rossi N, *et al.* 2018. Inference of CRISPR Edits from Sanger Trace Data. doi: 10.1101/251082. [Preprint].

Humphreys J.M and Chapple C. 2002. Rewriting the lignin roadmap. Current Opinion in Plant Biology 5, 224–229.

Jones L, Ennos AR, Turner SR. 2001. Cloning and characterization of irregular xylem4 (irx4): A severely lignin-deficient mutant of Arabidopsis. Plant Journal 26, 205–216.

Jung JH, Fouad WM, Vermerris W, Gallo M, Altpeter F. 2012. RNAi suppression of lignin biosynthesis in sugarcane reduces recalcitrance for biofuel production from lignocellulosic biomass. Plant Biotechnology Journal 10, 1067–1076.

Jung JH, Vermerris W, Gallo M, Fedenko JR, Erickson JE, Altpeter F. 2013. RNA interference suppression of lignin biosynthesis increases fermentable sugar yields for biofuel production from field-grown sugarcane. Plant Biotechnology Journal 11, 709–716.

Kannan B, Jung JH, Moxley GW, Lee SM, Altpeter F. 2018. TALEN-mediated targeted mutagenesis of more than 100 COMT copies/alleles in highly polyploid sugarcane improves saccharification efficiency without compromising biomass yield. Plant Biotechnology Journal 16, 856–866.

Kishimoto T, Uraki Y, Ubukata M. 2005. Easy synthesis of β-O-4 type lignin related polymers. Organic and Biomolecular Chemistry 3, 1067–1073.

Kordala N, Walter M, Brzozowski B, Lewandowska M. 2024. 2G-biofuel ethanol: an overview of crucial operations, advances and limitations. Springer Science and Business Media Deutschland GmbH.

Koshiba T, Hirose N, Mukai M, Yamamura M, Hattori T, Suzuki S, Sakamoto M, Umezawa T. 2013. Characterization of 5-hydroxyconiferaldehyde O-methyltransferase in Oryza sativa. Plant Biotechnology 30, 157–167.

Lam PY, Tobimatsu Y, Matsumoto N, *et al.* 2019*a*. OsCAldOMT1 is a bifunctional O-methyltransferase involved in the biosynthesis of tricin-lignins in rice cell walls. Scientific Reports 9.

Lam PY, Tobimatsu Y, Matsumoto N, *et al.* 2019*b*. OsCAldOMT1 is a bifunctional O-methyltransferase involved in the biosynthesis of tricin-lignins in rice cell walls. Scientific Reports 9.

Lan W, Lu F, Regner M, Zhu Y, Rencoret J, Ralph SA, Zakai UI, Morreel K, Boerjan W, Ralph J. 2015. Tricin, a flavonoid monomer in monocot lignification. Plant Physiology 167, 1284–1295.

Lan W, Morreel K, Lu F, Rencoret J, del Río JC, Voorend W, Vermerris W, Boerjan W, Ralph J. 2016*a*. Maize tricin-oligolignol metabolites and their implications for monocot lignification. Plant Physiology 171, 810–820.

Lan W, Rencoret J, Lu F, Karlen SD, Smith BG, Harris PJ, del Río JC, Ralph J. 2016*b*. Tricin-lignins: occurrence and quantitation of tricin in relation to phylogeny. Plant Journal 88, 1046–1057.

Lapierre C, Voxeur A, Karlen SD, Helm RF, Ralph J. 2018. Evaluation of Feruloylated and p-Coumaroylated Arabinosyl Units in Grass Arabinoxylans by Acidolysis in Dioxane/Methanol. Journal of Agricultural and Food Chemistry 66, 5418–5424.

Lee JH, Won HJ, Hoang Nguyen Tran P, Lee S mi, Kim HY, Jung JH. 2021. Improving lignocellulosic biofuel production by CRISPR/Cas9-mediated lignin modification in barley. GCB Bioenergy 13, 742–752.

Li L, Hill-Skinner S, Liu S, Beuchle D, Tang HM, Yeh CT, Nettleton D, Schnable PS. 2015. The maize brown midrib4 (bm4) gene encodes a functional folylpolyglutamate synthase. Plant Journal 81, 493–504.

Li R, Yang R, Zheng W, Wu L, Zhang C, Zhang H. 2022. Melatonin Promotes SGT1-Involved Signals to Ameliorate Drought Stress Adaption in Rice. International Journal of Molecular Sciences 23.

Liu B, Gómez LD, Hua C, *et al.* 2016. Linkage mapping of stem saccharification digestibility in rice. PLoS ONE 11.

Liu CG, Xiao Y, Xia XX, Zhao XQ, Peng L, Srinophakun P, Bai FW. 2019. Cellulosic ethanol production: Progress, challenges and strategies for solutions. Elsevier Inc.

Louie G V., Bowman ME, Tu Y, Mouradov A, Spangenberg G, Noel JP. 2010. Structure-function analyses of a caffeic acid O-methyltransferase from perennial ryegrass reveal the molecular basis for substrate preference. Plant Cell 22, 4114–4127.

Ma J, Shi S, Jia X, Xia F, Ma H, Gao J, Xu J. 2019. Advances in catalytic conversion of lignocellulose to chemicals and liquid fuels. Elsevier B.V.

Martins PK, Ribeiro AP, Cunha BADB Da, Kobayashi AK, Molinari HBC. 2015. A simple and highly efficient Agrobacterium-mediated transformation protocol for Setaria viridis. Biotechnology Reports 6, 41–44.

Menezes FO, Carvalho V, Moreira VA, Rigui AP, Gaspar M, Nievola CC. 2020. Juvenile plants of an epiphytic bromeliad exhibit rapid changes in carbohydrate content and antioxidant capacity to cope with water withholding. Theoretical and Experimental Plant Physiology 32, 89–98.

Meyer K, Shirley AM, Cusumano JC, Bell-Lelong DA, Chapple C. 1998. *Lignin monomer composition is determined by the expression of a cytochrome P450-dependent monooxygenase in Arabidopsis*.

Miller JE, Geadelmann JL, Marten GC. 1983. Effect of the Brown Midrib-Allele on Maize Silage Quality and Yield 1 . Crop Science 23, 493–496.

Minkenberg B, Zhang J, Xie K, Yang Y. 2019. CRISPR-PLANT v2: an online resource for highly specific guide RNA spacers based on improved off-target analysis. Blackwell Publishing Ltd.

Möller SR, Lancefield CS, Oates NC, Simister R, Dowle A, Gomez LD, McQueen-Mason SJ. 2022. CRISPR/Cas9 suppression of OsAT10, a rice BAHD acyltransferase, reduces p-coumaric acid incorporation into arabinoxylan without increasing saccharification. Frontiers in Plant Science 13.

Morse EE. 1947. Anthrone in Estimating Low Concentrations of Sucrose. Analytical Chemistry 19, 1012–1013.

Mosier N, Wyman C, Dale B, Elander R, Lee YY, Holtzapple M, Ladisch M. 2005. Features of promising technologies for pretreatment of lignocellulosic biomass. Bioresource Technology 96, 673–686.

Mota TR, de Souza WR, Oliveira DM, *et al.* 2021. Suppression of a BAHD acyltransferase decreases p-coumaroyl on arabinoxylan and improves biomass digestibility in the model grass Setaria viridis. Plant Journal 105, 136–150.

Oliveira DM, Saleme M de LS, Smith RA, *et al.* 2025. CRISPR/Cas9 editing of p-COUMAROYL-CoA:MONOLIGNOL TRANSFERASE 1 in maize alters phenolic metabolism, lignin structure, and lignin-first biomass processing. Trends in Biotechnology 43, 1166–1195.

Peracchi LM, Panahabadi R, Barros-Rios J, Bartley LE, Sanguinet KA. 2024. Grass lignin: biosynthesis, biological roles, and industrial applications. Frontiers Media SA.

Pesquet E, Cesarino I, Kajita S, Pawlowski K. 2025. Physiological roles of lignins – tuning cell wall hygroscopy and biomechanics. John Wiley and Sons Inc doi: 10.1111/nph.70505. [Preprint].

Pichon M, Deswartes C, Gerentes D, Guillaumie S, Lapierre C, Toppan A, Barrière Y, Goffner D. 2006. Variation in lignin and cell wall digestibility in caffeic acid O-methyltransferase down-regulated maize half-sib progenies in field experiments. Molecular Breeding 18, 253–261.

Piquemal J, Chamayou S, Nadaud I, *et al.* 2002*a*. Down-regulation of caffeic acid O-methyltransferase in maize revisited using a transgenic approach. Plant Physiology 130, 1675–1685.

Piquemal J, Chamayou S, Nadaud I, *et al.* 2002*b*. Down-regulation of caffeic acid O-methyltransferase in maize revisited using a transgenic approach. Plant Physiology 130, 1675–1685.

Priya S, Lin H-H, Menon S, *et al.* 2025. Engineered lignin composition in Brachypodium distachyon modulates the root-associated microbiome. doi: 10.1101/2025.07.30.667798. [Preprint].

Ralph J, Bunzel M, Marita JM, Hatfield RD, Lu F, Kim H, Schatz PF, Grabber JH, Steinhart H. 2004. *Peroxidase-dependent cross-linking reactions of p-hydroxycinnamates in plant cell walls*.

Ralph J, Lapierre C, Lu F, Marita JM, Pilate G, Doorsselaere J Van, Boerjan W, Jouanin L. 2001*a*. NMR Evidence for Benzodioxane Structures Resulting from Incorporation of 5-Hydroxyconiferyl Alcohol into Lignins of O-Methyltransferase-Deficient Poplars. doi: 10.1021/jf001042.

Ralph J, Lapierre C, Marita JM, *et al.* 2001*b*. *Elucidation of new structures in lignins of CAD-and COMT-deficient plants by NMR*.

Rao X, Dixon RA. 2018. Current models for transcriptional regulation of secondary cell wall biosynthesis in grasses. Frontiers Media S.A.

Rates A de B, Cesarino I. 2023. Pour some sugar on me: The diverse functions of phenylpropanoid glycosylation. Elsevier GmbH.

Reshmy R, Philip E, Unni R, Paul SA, Sindhu R, Madhavan A, Sirohi R, Pandey A, Binod P. 2023. Bioethanol—A promising alternative fuel for sustainable future. Valorization of Biomass to Bioproducts: Organic Acids and Biofuels. Elsevier, 179–196.

Del Río JC, Rencoret J, Prinsen P, Martínez ÁT, Ralph J, Gutiérrez A. 2012. Structural characterization of wheat straw lignin as revealed by analytical pyrolysis, 2D-NMR, and reductive cleavage methods. Journal of Agricultural and Food Chemistry 60, 5922–5935.

Saballos A, Vermerris W, Rivera L, Ejeta G. 2008. Allelic Association, Chemical Characterization and Saccharification Properties of brown midrib Mutants of Sorghum (Sorghum bicolor (L.) Moench). BioEnergy Research 1, 193–204.

Saleme M de LS, Cesarino I, Vargas L, *et al.* 2017. Silencing CAFFEOYL SHIKIMATE ESTERASE affects lignification and improves saccharification in poplar. Plant Physiology 175, 1040–1057.

Sattler SE, Funnell-Harris DL, Pedersen JF. 2010. Brown midrib mutations and their importance to the utilization of maize, sorghum, and pearl millet lignocellulosic tissues.

Shi N, Liu D, Huang Q, Guo Z, Jiang R, Wang F, Chen Q, Li M, Shen G, Wen F. 2019. Product-oriented decomposition of lignocellulose catalyzed by novel polyoxometalates-ionic liquid mixture. Bioresource Technology 283, 174–183.

de Souza WR, Martins PK, Freeman J, *et al.* 2018. Suppression of a single BAHD gene in Setaria viridis causes large, stable decreases in cell wall feruloylation and increases biomass digestibility. New Phytologist 218, 81–93.

Studer MH, DeMartini JD, Davis MF, Sykes RW, Davison B, Keller M, Tuskan GA, Wyman CE. 2011. Lignin content in natural populus variants affects sugar release. Proceedings of the National Academy of Sciences of the United States of America 108, 6300–6305.

Sun S, Wen D, Yang W, Meng Q, Shi Q, Gong B. 2020. Overexpression of Caffeic Acid O-Methyltransferase 1 (COMT1) Increases Melatonin Level and Salt Stress Tolerance in Tomato Plant. Journal of Plant Growth Regulation 39, 1221–1235.

Takeda Y, Suzuki S, Tobimatsu Y, Osakabe K, Osakabe Y, Ragamustari SK, Sakamoto M, Umezawa T. 2019. Lignin characterization of rice CONIFERALDEHYDE 5-HYDROXYLASE loss-of-function mutants generated with the CRISPR/Cas9 system. Plant Journal 97, 543–554.

Tang HM, Liu S, Hill-Skinner S, Wu W, Reed D, Yeh CT, Nettleton D, Schnable PS. 2014. The maize brown midrib2 (bm2) gene encodes a methylenetetrahydrofolate reductase that contributes to lignin accumulation. Plant Journal 77, 380–392.

Terrett OM, Dupree P. 2019. Covalent interactions between lignin and hemicelluloses in plant secondary cell walls. Elsevier Ltd.

Tetreault HM, Gries T, Liu S, Toy J, Xin Z, Vermerris W, Ralph J, Funnell-Harris DL, Sattler SE. 2021. The Sorghum (Sorghum bicolor) Brown Midrib 30 Gene Encodes a Chalcone Isomerase Required for Cell Wall Lignification. Frontiers in Plant Science 12.

Tian A, Zhao J, Tang B, Zhu D, Fu C, Xiong H. 2021. Study on the Pretreatment of Soil Hyperspectral and Na + Ion Data under Different Degrees of Human Activity Stress by Fractional-Order Derivatives. 13, 3974.

Trabucco GM, Matos DA, Lee SJ, Saathoff AJ, Priest HD, Mockler TC, Sarath G, Hazen SP. 2013*a*. Functional characterization of cinnamyl alcohol dehydrogenase and caffeic acid O-methyltransferase in Brachypodium distachyon. BMC Biotechnology 13.

Trabucco GM, Matos DA, Lee SJ, Saathoff AJ, Priest HD, Mockler TC, Sarath G, Hazen SP. 2013*b*. Functional characterization of cinnamyl alcohol dehydrogenase and caffeic acid O-methyltransferase in Brachypodium distachyon. BMC Biotechnology 13.

Tye YY, Lee KT, Wan Abdullah WN, Leh CP. 2016. The world availability of non-wood lignocellulosic biomass for the production of cellulosic ethanol and potential pretreatments for the enhancement of enzymatic saccharification. Elsevier Ltd.

U.S. Energy Information Administration E. 2021. *Annual Energy Outlook 2021 Narrative*.

Vanholme R, Demedts B, Morreel K, Ralph J, Boerjan W. 2010. Lignin biosynthesis and structure. Plant Physiology 153, 895–905.

Vanholme R, Storme V, Vanholme B, Sundin L, Christensen JH, Goeminne G, Halpin C, Rohde A, Morreel K, Boerjana W. 2012. A systems biology view of responses to lignin biosynthesis perturbations in Arabidopsis. Plant Cell 24, 3506–3529.

Vermerris W, Saballos A, Ejeta G, Mosier NS, Ladisch MR, Carpita NC. 2007. Molecular breeding to enhance ethanol production from corn and sorghum stover. Crop Science.

Vignols F, Rlgau J, Torres MA, Capellades M, Puigdomènech ' P. 1995. *The brown midrib3 (bm3) Mutation in Maize Occurs in the Gene Encoding Caffeic Acid O-Methyltransferase*. American Society of Plant Physiologists.

Vogt T. 2010. Phenylpropanoid biosynthesis. Oxford University Press.

De Vries L, Vanholme R, Van Acker R, De Meester B, Sundin L, Boerjan W. 2018. Stacking of a low-lignin trait with an increas ed guaiacyl and 5-hydroxyguaiacyl unit trait leads to additive and synergistic effects on saccharification efficiency in Arabidopsis thaliana. Biotechnology for Biofuels 11.

Wang H, Chen Z, Luo R, Lei C, Zhang M, Gao A, Pu J, Zhang H. 2024. Caffeic Acid O-Methyltransferase Gene Family in Mango (Mangifera indica L.) with Transcriptional Analysis under Biotic and Abiotic Stresses and the Role of MiCOMT1 in Salt Tolerance. International Journal of Molecular Sciences 25.

Wannitikul P, Dachphun I, Sakulkoo J, Suttangkakul A, Wonnapinij P, Simister R, Gomez LD, Vuttipongchaikij S. 2024. In Vivo Proximity Cross-Linking and Immunoprecipitation of Cell Wall Epitopes Identify Proteins Associated with the Biosynthesis of Matrix Polysaccharides. ACS Omega 9, 31438–31454.

Wu Z, Wang N, Hisano H, Cao Y, Wu F, Liu W, Bao Y, Wang ZY, Fu C. 2019. Simultaneous regulation of F5H in COMT-RNAi transgenic switchgrass alters effects of COMT suppression on syringyl lignin biosynthesis. Plant Biotechnology Journal 17, 836–845.

Xu B, Escamilla-Treviño LL, Sathitsuksanoh N, Shen Z, Shen H, Percival Zhang YH, Dixon RA, Zhao B. 2011. Silencing of 4-coumarate: Coenzyme A ligase in switchgrass leads to reduced lignin content and improved fermentable sugar yields for biofuel production. New Phytologist 192, 611–625.

Yao Z, Zhang X, Liang Y, Zhang J, Xu Y, Chen S, Zhao D. 2022. NtCOMT1 responsible for phytomelatonin biosynthesis confers drought tolerance in Nicotiana tabacum. Phytochemistry 202.

Yılbaşı Z. 2025. Biofuels, E-Fuels, and Waste-Derived Fuels: Advances, Challenges, and Future Directions. Sustainability 17, 6145.

Yu H, Zhang G, Liu J, *et al.* 2025. A functional cascading of lignin modification via repression of caffeic acid O-methyltransferase for bioproduction and anti-oxidation in rice. Journal of Advanced Research doi: 10.1016/j.jare.2025.01.048.

Zubieta C, Kota P, Ferrer JL, Dixon RA, Noel JP. 2002. Structural basis for the modulation of lignin monomer methylation by caffeic acid/5-hydroxyferulic acid 3/5-O-methyltransferase. Plant Cell 14, 1265–1277.

TABLES

Table 1. Differentially accumulated compounds in aerial parts of *SvCOMT1* mutants

Name	RT (min)	Exp. *m/z*	Peak Area±SD				Fold-Change
			WT	Ev.1	Ev.2	Ev.3	
Caffeic acid-O-hexoside	1.29	341.1090	324±117	3572±5689	573±174	426±206	**2.4**
tricetin 3'-O-methyl ether (Selgin)	5.07	315.0719	1261±233	2357±1063	2376±447	1819±322	**4.2**
kaempferol-3-O-caffeoyl-glucoside	6.24	563.1620	ND	232±213	236±89	212±37	**200**
2-Glucosyloxy-4-methoxycinnamic acid	8.23	355.1037	73±22	306±13	519±49	152±20	**4.5**
Di-O-caffeoyl-hexoside	8.86	677.1338	1186±55	2375±105	2769±143	2311±65	**2**
Unknown	8.86	745.1220	99±3	198±5	223±19	201±2	**2**
Luteolin-3',7-diglucoside	8.87	609.1465	8817±332	16666±830	19149±831	16135±663	**2**
Quercetin-diglucoside	8.87	672.1423	344±24	799±41	1102±53	768±65	**2.6**
Unknown	8.87	708.0647	73±2	143±2	172±14	147±14	**2.1**
Feruloyl-O-hexoside	8.98	525.1623	512±59	1323±40	1154±30	923±21	**2.2**
Unknown	9.55	577.1576	214±40	106±2	95±3	115±8	**-2**
G(8–O–4)G-hexoside	9.94	535.1825	1751±338	3879±181	4736±210	2526±279	**2.1**
Flavonoid-diglucuronide	10.12	693.2056	43±18	127±14	184±10	69±5	**3**
Caffeoyl-glucoside	10.17	499.1244	17±13	741±66	1036±13	500±71	**44**
Syringaresinol O-beta-D-glucoside	10.58	579.2041	925±170	424±24	551±9	401±52	**-2**
Caffeic acid	10.64	179.0564	81±34	210±122	179±13	101±13	**2**
Tri-caffeoylquinic acid	11.95	785.1943	762±265	1636±229	2263±530	1660±412	**2.4**
Unknown	11.95	853.1824	62±36	163±18	217±55	165±35	**2.4**
Sinapoyl-feruloyl-glucoside	12.48	583.2194	602±180	104±55	111±10	102±9	**-6**
G(8–O–4) ferulic acid di (acetylhexoside)	12.57	799.2105	1327±508	2235±268	3179±361	2970±450	**2.1**

Dicaffeoyl-hexaric acid	12.58	867.1984	16±8	60±14	138±27	152±37	**7**
Unknown	13.18	403.1044	461±142	174±20	182±52	301±38	**-2**

FIGURE LEGENDS

Figure 1. Lignin biosynthetic pathway in grasses
Current model of lignin biosynthesis encompassing the phenylpropanoid and flavonoid pathways. Preferential carbon flux is indicated by green arrows, while dashed arrows denote unconfirmed steps. The "shikimate shunt" of the phenylpropanoid pathway is highlighted in gray. Solid arrows indicate experimentally confirmed enzymatic steps, and dashed arrows represent suggested conversions. **Phenylpropanoid-related enzymes:** PAL (phenylalanine ammonia-lyase), PTAL (phenylalanine/tyrosine ammonia-lyase), C4H (cinnamate 4-hydroxylase), 4CL (4-hydroxycinnamate:CoA ligase), C3H (4-coumarate 3-hydroxylase/ascorbate peroxidase), CSE (caffeoyl shikimate esterase), HCT (hydroxycinnamoyl-CoA:shikimate/quinate hydroxycinnamoyl transferase), C3'H (4-coumaroyl shikimate 3'-hydroxylase), CCoAOMT (caffeoyl-CoA 3-O-methyltransferase), CCR (cinnamoyl-CoA reductase), F5H (ferulate/coniferaldehyde 5-hydroxylase), COMT (caffeic acid O-methyltransferase), CAD (cinnamyl alcohol dehydrogenase), PMT (p-coumaroyl-CoA:monolignol transferase), FMT (feruloyl-CoA:monolignol transferase). **Flavonoid- and flavone-related enzymes:** CHS (chalcone synthase), CHI (chalcone isomerase), FNSII (flavone synthase II), A3'H/C5'H (apigenin 3'-hydroxylase/chrysoeriol 5'-hydroxylase). The red star marks the intermediates that were altered in *COMT1*-edited plants.

Figure 2. Phenotypes of wild-type (WT) plants and *COMT1* knockout lines.

(A) Leaves of WT plants and *COMT1* knockout lines (Ev.1, Ev.2, and Ev.3). The typical symptoms of *bmr* (brown midrib) mutations, characterized by brown coloration in the central vein, were not observed in the knockout lines. (B) Representative image showing no visible growth differences between WT plants and *COMT1* knockout lines (Ev.1, Ev.2, and Ev.3).

Figure 3. Quantification of lignin content and monomeric composition in wild-type and *COMT1* knockout plants.

(A) Soluble lignin content determined by the acetyl bromide method, and levels of (B) guaiacyl (G), (C) syringyl (S), (D) p-hydroxyphenyl (H), and (E) tricin monomers in alcohol-insoluble residue (AIR) extracts. Lignin content and monomeric composition were analyzed in wild-type (WT) plants and in three independent *COMT1* knockout lines (Ev.1, Ev.2, and Ev.3). Values are expressed as µg/mg AIR and represent means ± SEM (n = 4). Statistical differences between WT and knockout lines were assessed using one-way ANOVA followed by Tukey's post hoc test (*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ***$P < 0.0001$; n.s., not significant).

Figure 4. Determination of hydroxycinnamic acid (HCA) content.

(A) Total HCA content, (B) HCA content linked to arabinose, and (C) HCA content bound to lignin. The analysis was performed on wild-type (WT) plants and *COMT1* knockout lines (Ev.1, Ev.2, and Ev.3). Values were obtained by alkaline hydrolysis per mg of the AIR (Alcohol-Insoluble Residue)

fraction and are expressed as means ± SEM (n = 4). Statistical significance between *COMT1* knockout lines and WT was assessed by one-way ANOVA followed by Tukey's post hoc test. Significance levels: $*P < 0.05$; $**P < 0.01$; $***P < 0.001$; $****P < 0.0001$; n.s., not significant

Figure *5*. Biomass saccharification analysis.

The analysis was conducted on wild-type (WT) plants and *COMT1* knockout lines (Ev.1, Ev.2, and Ev.3) using untreated biomass (A) and biomass treated with 500 mM NaOH (B). Values are expressed as nmol of glucose released per mg of AIR (alcohol-insoluble residue), and data are presented as means ± SEM (n = 4). Statistical significance between *COMT1* knockout lines and WT was assessed by one-way ANOVA followed by Tukey's post hoc test. Significance levels: $*P < 0.05$; $**P < 0.01$; $***P < 0.001$; $****P < 0.0001$; n.s., not significant.

Figure 6. Responses of Relative Water Content and Dry Mass in Wild-Type and *COMT1* Knockout Plants to 5-Day NaCl Treatments

(A–B) Representative phenotypes after 5 days of exposure to 0, 100, and 250 mM NaCl. (C) Survival of plants exposed to 250 mM NaCl, showing that all genotypes failed to survive beyond 9 days. (D–E) Relative water content (RWC) and dry mass after 5 days under the indicated NaCl concentrations. Values are presented as means ± SEM (n = 4). Statistical significance relative to WT was determined by one-way ANOVA followed by Tukey's post hoc test ($p \leq 0.05$); n.s., not significant.

Figure 1. Lignin biosynthetic pathway in grasses

Current model of lignin biosynthesis encompassing the phenylpropanoid and flavonoid pathways. Preferential carbon flux is indicated by green arrows, while dashed arrows denote unconfirmed steps. The "shikimate shunt" of the phenylpropanoid pathway is highlighted in gray. Solid arrows indicate experimentally confirmed enzymatic steps, and dashed arrows represent suggested conversions. **Phenylpropanoid-related enzymes:** PAL (phenylalanine ammonia-lyase), PTAL (phenylalanine/tyrosine ammonia-lyase), C4H (cinnamate 4-hydroxylase), 4CL (4-hydroxycinnamate:CoA ligase), C3H (4-coumarate 3-hydroxylase/ascorbate peroxidase), CSE (caffeoyl shikimate esterase), HCT (hydroxycinnamoyl-CoA:shikimate/quinate hydroxycinnamoyl transferase), C3'H (4-coumaroyl shikimate 3'-hydroxylase), CCoAOMT (caffeoyl-CoA 3-O-methyltransferase), CCR (cinnamoyl-CoA reductase), F5H (ferulate/coniferaldehyde 5-hydroxylase), COMT (caffeic acid O-methyltransferase), CAD (cinnamyl alcohol dehydrogenase), PMT (p-coumaroyl-CoA:monolignol transferase), FMT (feruloyl-CoA:monolignol transferase). **Flavonoid- and flavone-related enzymes:** CHS (chalcone synthase), CHI (chalcone isomerase), FNSII (flavone synthase II), A3'H/C5'H (apigenin 3'-hydroxylase/chrysoeriol 5'-hydroxylase). The red star marks the intermediates that were altered in *COMT1*-edited plants.

Figure 2. Phenotypes of wild-type (WT) plants and *COMT1* knockout lines.

(A) Leaves of WT plants and *COMT1* knockout lines (Ev.1, Ev.2, and Ev.3). The typical symptoms of *bmr* (brown midrib) mutations, characterized by brown coloration in the central vein, were not observed in the knockout lines. (B) Representative image showing no visible growth differences between WT plants and *COMT1* knockout lines (Ev.1, Ev.2, and Ev.3).

Figure 3. Quantification of lignin content and monomeric composition in wild-type and *COMT1* knockout plants.

(A) Soluble lignin content determined by the acetyl bromide method, and levels of (B) guaiacyl (G), (C) syringyl (S), (D) p-hydroxyphenyl (H), and (E) tricin monomers in alcohol-insoluble residue (AIR) extracts. Lignin content and monomeric composition were analyzed in wild-type (WT) plants and in three independent *COMT1* knockout lines (Ev.1, Ev.2, and Ev.3). Values are expressed as µg/mg AIR and represent means ± SEM (n = 4). Statistical differences between WT and knockout lines were assessed using one-way ANOVA followed by Tukey's post hoc test (*P < 0.05; **P < 0.01; ***P < 0.001; ***P < 0.0001; n.s., not significant).

Figure 4. Determination of hydroxycinnamic acid (HCA) content.

(A) Total HCA content, (B) HCA content linked to arabinose, and (C) HCA content bound to lignin. The analysis was performed on wild-type (WT) plants and *COMT1* knockout lines (Ev.1, Ev.2, and Ev.3). Values were obtained by alkaline hydrolysis per mg of the AIR (Alcohol-Insoluble Residue) fraction and are expressed as means ± SEM (n = 4). Statistical significance between *COMT1* knockout lines and WT was assessed by one-way ANOVA followed by Tukey's post hoc test. Significance levels: *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; n.s., not significant

Figure *5*. Biomass saccharification analysis.

The analysis was conducted on wild-type (WT) plants and *COMT1* knockout lines (Ev.1, Ev.2, and Ev.3) using untreated biomass (A) and biomass treated with 500 mM NaOH (B). Values are expressed as nmol of glucose released per mg of AIR (alcohol-insoluble residue), and data are presented as means ± SEM (n = 4). Statistical significance between *COMT1* knockout lines and WT was assessed by one-way ANOVA followed by Tukey's post hoc test. Significance levels: *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; n.s., not significant.

Figure 6. Responses of Relative Water Content and Dry Mass in Wild-Type and *COMT1* Knockout Plants to 5-Day NaCl Treatments

(A–B) Representative phenotypes after 5 days of exposure to 0, 100, and 250 mM NaCl. (C) Survival of plants exposed to 250 mM NaCl, showing that all genotypes failed to survive beyond 9 days. (D–E) Relative water content (RWC) and dry mass after 5 days under the indicated NaCl concentrations. Values are presented as means ± SEM (n = 4). Statistical significance relative to WT was determined by one-way ANOVA followed by Tukey's post hoc test (p ≤ 0.05); n.s., not significant.