IGHV3-53 antibody abundance drives divergent SARS-CoV-2 immune imprinting

Xiao Niu[1,2,3,#], Fanchong Jian[1,#], Yutong Li[1,2], Ke Li[1,2], Siyu Lei[1,2,4], Weiliang Song[1,2], Ruoxi Kong[1,2],

Xingan Cai[1,2], Ran An[2], Yao Wang[2], Yifei Huang[5,6,7], Lingling Yu[2], Wenjing Wang[2], Haiyan Sun[2],

Yuanling Yu[2], Jing Wang[2], Binmao Zhang[8], Tianen Zhu[8,9], Jianheng Huang[8,10], Fei Shao[2], Xiaotian

Tan[8], Sai Luo[5,6,7], Yunlong Cao[1,2,4*]

[1]Biomedical Pioneering Innovation Center (BIOPIC), School of Life Sciences, Peking University,

Beijing, P.R. China.

[2]Changping Laboratory, Beijing, P.R. China.

[3]College of Chemistry and Molecular Engineering, Peking University, Beijing, P.R. China.

[4]Peking–Tsinghua Center for Life Sciences, Peking University, Beijing, P. R. China.

[5]Center for Infectious Disease Research, Tsinghua University, Beijing, P.R. China.

[6]Tsinghua-Peking Joint Center for Life Sciences, Tsinghua University, Beijing, P.R. China.

[7]School of Basic Medical Sciences, Tsinghua University, Beijing, P.R. China.

[8]Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences.

[9]Shenzhen University General Hospital.

[10]Shenzhen University.

*Correspondence: Yunlong Cao (yunlongcao@pku.edu.cn).

#These authors contributed equally.

Abstract:

The mechanisms driving divergent SARS-CoV-2 immune imprinting in populations primed with different COVID-19 vaccines remain unclear. Recipients of inactivated vaccines readily develop Omicron-specific antibodies through repeated breakthrough infections, whereas mRNA-vaccinated individuals exhibit severe ancestral-strain imprinting that suppresses *de novo* Omicron-specific responses. These differences could result in distinct antibody landscapes, leading to regional epidemiological divergence and necessitating region-specific vaccine update strategies. Importantly, conventional wild-type mouse models fail to recapitulate strong human SARS-CoV-2 imprinting, which significantly hinders imprinting-related mechanistic investigation and vaccine update evaluation. Here, we surprisingly found that V(D)J-humanized mice could faithfully recapitulate human severe SARS-CoV-2 immune imprinting phenotypes. Comprehensive antibody repertoire and epitope mapping of 583 monoclonal antibodies from these models revealed that the abundance of pre-existing IGHV3-53/66-encoded SARS-CoV-2 antibody responses determines imprinting severity following Omicron exposure through antibody-mediated masking of Omicron-specific epitopes. Both passive transfer of IGHV3-53/66 antibodies and knock-in of the human IGHV3-53 gene were sufficient to induce severe SARS-CoV-2 imprinting in wild-type mice. Concordantly, head-to-head comparison also showed that mRNA vaccine recipients retained higher IGHV3-53/66 antibody abundance and thus stronger imprinting than inactivated vaccine recipients. Consequently, compared to NB.1.8.1, XFG exhibits greater immune evasion in mRNA-vaccinated individuals but not in inactivated vaccine recipients. This explains the regional predominance of XFG in mRNA-vaccinated populations, while NB.1.8.1 prevails in inactivated vaccine-dominated countries. Together, these findings demonstrate that the V(D)J germline repertoire—even a single germline-encoded antibody response—can profoundly shape humoral imprinting severity. Accordingly, we constructed a human IGHV3-53 knock-in mouse model that can accurately recapitulate human SARS-CoV-2 antibody landscape, providing a valuable tool for guiding future COVID-19 vaccine updates.

Main

After extensive global circulation, SARS-CoV-2 continues to co-evolve with human humoral immunity, repeatedly generating variants capable of evading neutralizing antibodies elicited by prior infections and vaccinations [1–11]. Recently, a divergence in SARS-CoV-2 viral circulation has emerged between East and West. In Eastern countries (primarily China), the NB.1.8.1 lineage predominates while XFG remains virtually absent [9]. Conversely, XFG has rapidly risen to dominance in Western regions, leaving NB.1.8.1 as a minor lineage (Extended Data Fig. 1a).

The global deployment of diverse COVID-19 vaccine platforms has consequently established distinct humoral immune landscapes across populations [12–17]. Recipients of mRNA vaccines, primarily in Western countries, exhibit pronounced immune imprinting, persistently recalling ancestral Wuhan-Hu-1 (Wuhan) spike-reactive antibodies even upon repeated Omicron exposures, significantly limiting the development of Omicron-specific neutralizing responses [18–25]. In contrast, individuals only receiving inactivated vaccines, predominantly in China, demonstrate greater adaptability in humoral responses, developing robust Omicron-specific antibodies upon repeated infections or boosters, thus overriding Wuhan immune imprinting [26–29]. These imprinting disparities may potentially contribute to the observed divergence in SARS-CoV-2 epidemiology, raising potential needs for region-specific vaccine update strategies.

The mechanism that drives the divergent SARS-CoV-2 immune imprinting in populations primed with different COVID-19 vaccines is unclear and needs immediate investigation. However, prior studies have shown that, unlike humans, wild-type mice only exhibit weak SARS-CoV-2 ancestral-strain imprinting whether receiving inactivated or mRNA-based vaccine priming [18], limiting the utility of this animal model for both mechanistic investigation of the divergent imprinting and translational vaccine update preclinical research.

In this paper, we present a head-to-head comparison of antibody repertoires, imprinting severity, and responses to emerging variants in individuals primed with mRNA versus inactivated vaccines. These analyses reveal that differences in immune imprinting are already influencing real-world epidemiology, contributing to the distinct regional dominance of XFG and NB.1.8.1. Using V(D)J-humanized mice, we demonstrate faithful recapitulation of strong human-like imprinting and

identify pre-existing IGHV3-53/66-encoded antibodies as the key driver of imprinting severity via epitope masking. Finally, we establish a human IGHV3-53 knock-in mice model that accurately recapitulate human SARS-CoV-2 antibody responses, offering a practical tool for evaluating future vaccine update strategies.

Distinct immune imprinting in humans

To directly compare the humoral immunity in individuals primed with mRNA versus inactivated vaccines, we recruited 20 participants stratified into two cohorts based on vaccination history (Supplementary Table 1 and Fig. 1a). All participants had resided in the same region since 2021 and had experienced multiple Omicron infections. The inactivated-only cohort (n = 12) received three doses of CoronaVac, with samples collected 46.5 ± 5.4 months after the first vaccine dose. The mRNA-vaccinated cohort (n = 8) received at least one mRNA vaccine dose (BNT162b2 or mRNA-1273), with samples collected 46.0 ± 2.3 months after initial vaccination. Although detailed infection histories were unavailable, widespread Omicron transmission following China's reopening strongly suggests that most participants were exposed to at least BA.5 and JN.1 sublineages.

To assess immune imprinting, we first used fluorescence-activated cell sorting (FACS) to quantify cross-reactive memory B cells, defined as the proportion of Wuhan$^+$ cells among JN.1$^+$ memory B cells (Fig. 1b and Supplementary Information Fig. 1a). The mRNA-vaccinated cohort indeed exhibited a significantly higher proportion of cross-reactive memory B cells than the inactivated-only cohort, indicating stronger imprinting (Fig. 1c). Additionally, the mRNA group showed a higher proportion of Wuhan$^+$ memory B cells, reflecting persistent responses to the ancestral strain. In contrast, JN.1$^+$ memory B cell frequencies were comparable between groups, suggesting comparable responses to recent variants despite different priming histories (Fig. 1c).

To examine antibody specificity, we depleted Wuhan RBD-binding antibodies from serum and measured the remaining IgG titres against the JN.1 RBD (Fig. 1d). This depletion nearly abolished JN.1-binding activity in the mRNA-vaccinated cohort (19-fold reduction), indicating that the response was almost entirely contributed by cross-reactive antibodies. In contrast, the inactivated-

Figure 1

Figure 1 | mRNA and inactivated vaccine priming induce distinct immune imprinting level.
a, Schematic of the SARS-CoV-2-related immune histories and sampling time of the inactivated-only and mRNA-vaccinated cohorts involved in this study. b, Representative flow cytometry dot plots of cross-reactive memory B cells from individuals from the inactivated-only (top) and mRNA-vaccinated (bottom) cohorts. APC, allophycocyanin; PE, phycoerythrin; BV421, Brilliant Violet 421. c, Scatter plots showing the proportion of cross-reactive memory B cells (left), JN.1-reactive memory B cells (middle), and Wuhan-reactive memory B cells (right). d, Serum IgG midpoint titer against Wuhan (left) or JN.1 (right) RBD before and after Wuhan RBD depletion. Statistical significance of the fold-reduction in titers was assessed between two cohorts. Dashed lines indicate the limit of detection (midpoint titre = 10). e, NAb response of the two cohorts against a panel of SARS-CoV-2 variant pseudoviruses. Geometric mean titres (GMTs) are shown on the top. Dashed lines indicate the limit of detection (NT_{50} = 10). f, Heavy chain V gene usage proportion of the two cohorts. IGHV3-53 and IGHV3-66 were combined for analysis. Two-tailed Wilcoxon rank-sum tests were used in c-e. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$; NS, not significant ($P > 0.05$).

only cohort showed a more modest decrease (4.4-fold), consistent with the presence of a subset of JN.1-specific antibodies (Fig. 1d).

Neutralization assays against a pseudovirus panel further supported this divergence (Fig. 1e). In the mRNA-vaccinated cohort, 50% neutralization titres (NT_{50}) were highest for D614G and declined rapidly with newer variants. In contrast, the inactivated-only cohort maintained relatively stable neutralization, declining only from JN.1 onward. Interestingly, the NT_{50} curves intersected at JN.1, beyond which the inactivated cohort showed higher neutralization. These data suggest that, owing to stronger immune imprinting, the mRNA cohort's neutralizing responses to emerging variants are strongly confined by the recall and expansion of antibodies originally elicited by ancestral exposure, whereas the more adaptable response in the inactivated cohort enables superior neutralization of later-emerging variants.

Notably, the mRNA-vaccinated cohort showed a marked reduction in titres against XFG relative to NB.1.8.1, whereas the inactivated-only cohort displayed comparable neutralizing activity against both variants. This disparity provides a serological explanation for the regional predominance of XFG in mRNA-vaccinated populations and NB.1.8.1 in inactivated vaccine-dominated countries [9]. Additionally, neutralization titres against BA.3.2 were significantly higher in the mRNA cohort (Fig. 1e). This variant carries unique mutations that have been shown to evade Class 1/4 antibodies prevalent in inactivated vaccinees but not the dominant Class 1 antibodies induced by mRNA vaccines [9]. Collectively, these head-to-head comparisons of serological and flow cytometry data confirm robust imprinting in the mRNA-vaccinated cohort and weaker imprinting in the inactivated cohort that can be partially overcome. Such imprinting differences have resulted in distinct antibody landscapes, contributing to regional epidemiological divergence and raising the potential necessity for region-specific vaccine update strategies.

While the lack of longitudinal samples makes it challenging to trace the evolutionary mechanism of this divergence in humans, distinct signatures within the BCR repertoire may offer critical insights. To explore these correlates, we performed plate-based single B cell V(D)J sequencing to compare heavy-chain V-gene usage of JN.1⁺ memory B cells between the two cohorts (Fig. 1f) [30]. The mRNA-vaccinated cohort showed enriched usage of IGHV3-53/66—germline genes encoding "public" antibodies targeting the Class 1 epitope, which overlaps with the ACE2-binding site [31–38].

These neutralizing antibodies are broadly elicited by Wuhan vaccination or infection and are well

known for their potent neutralizing capacity [31,37]. In contrast, the inactivated-only cohort showed

greater representation of IGHV5-51 and IGHV2-5, associated with Omicron-specific Class 1/4

antibodies [39]. The differences in light chain V-gene usage between the two cohorts were less

pronounced, with both groups showing a preference for IGKV1-39, IGKV3-20, and IGKV1-33

(Extended Data Fig. 2a).

These findings suggest that the IGHV3-53/66–encoded Class 1 antibodies—originally elicited by

ancestral Wuhan strain exposure—are preferentially recalled and expanded upon repeated Omicron

encounters, and the mRNA-vaccinated cohort exhibits much higher IGHV3-53/66 antibody

abundance compared to the inactivated-only cohort, due to its higher priming vaccine

immunogenicity. To further dissect the underlying mechanisms and confirm the relationship

between immunoglobulin germline genes and severe SARS-CoV-2 imprinting, animal models that

faithfully recapitulate these human phenotypes are therefore required.

V(D)J-humanized mice mimic strong imprinting

Previous studies have shown that wild-type mice do not recapitulate the strong SARS-CoV-2

immune imprinting observed in the human mRNA-vaccinated cohort [18]. Given that our human data

identified a critical role for the IGHV3-53/66 germline genes in severe SARS-CoV-2 imprinting,

we hypothesized that the absence of these specific human V-genes in the murine repertoire explains

this discrepancy. Therefore, we employed V(D)J humanized mice for comparative analysis. These

BALB/c-derived mice, commonly used in therapeutic antibody discovery [40], carry human

immunoglobulin V(D)J genes for all heavy chains and kappa light chains. Both V(D)J humanized

and wild-type mice were primed with three doses of ancestral mRNA vaccine (encoding the spike

protein of SARS-CoV-2 ancestral strain), followed by two booster doses of BA.5 mRNA vaccine to

mimic human exposure history to SARS-CoV-2 variants (Fig. 2a). Blood, spleens, and lymph nodes

were collected one week after the third ancestral dose and each BA.5 dose for serological and flow

cytometry analyses.

We first performed FACS analysis on mouse lymph nodes to assess cross-reactivity among memory

Figure 2

Figure 2 | V(D)J humanized mice recapitulates severe SARS-CoV-2 imprinting.
a, Schematic of the mouse immunization regimen and sampling timeline for wild-type and V(D)J-humanized mice. The number of mice is indicated at the time point for each endpoint experiment. b, Scatter plots showing the proportion of cross-reactive memory B cells in draining lymph nodes of the two mouse strains after one (left) or two (right) BA.5 boosts. c, Serum IgG midpoint titer of the two mouse strains after one (left) or two (right) BA.5 boosts against Wuhan or BA.5 RBD before and after Wuhan RBD depletion. Statistical significance of the fold-reduction in titers was assessed between humanized and wild-type mice. Dashed lines indicate the limit of detection (midpoint titre = 100). d, Serum neutralization titers (NT_{50}) of the two mouse models after Wuhan priming (left), one BA.5 boost (middle), or two BA.5 boosts (right) against a panel of SARS-CoV-2 variant pseudoviruses. Geometric mean titres (GMTs) are shown on the top. Dashed lines indicate the limit of detection ($NT_{50} = 60$). Two-tailed Wilcoxon rank-sum tests were used in b-d.

B cells and germinal-center (GC) B cells (Supplementary Information Fig. 1b). As expected, wild-type mice harbored abundant BA.5-specific memory and GC B cells after each BA.5 booster (Fig. 2b and Extended Data Fig. 3a). In striking contrast, the V(D)J-humanized mice exhibited a strong imprinting phenotype, developing a high proportion of cross-reactive memory B cells (89% and 82% after the first and second boosters, respectively) that mirrored the response observed in the human mRNA cohort. Despite similar overall percentages of GC B cells, humanized mice showed a significantly higher frequency of cross-reactive GC B cells after one BA.5 booster, indicating that their GC response was dominated by Wuhan-strain imprinted B cells (Extended Data Fig. 3a and 3b). Additionally, they presented a significantly higher frequency of class-switched memory B cells, which aligns with the preferential recall and expansion of a pre-existing memory population typical of immune imprinting (Extended Data Fig. 3c).

The imprinting was also faithfully reflected in the serological response. Wuhan RBD-depletion caused a substantially greater drop in BA.5-binding IgG titres in humanized mice than in wild-type mice (10-fold vs. 2.1-fold), although the magnitude of this effect varied among individual humanized mice (Fig. 2c). Specifically, titres in three humanized mice dropped to undetectable levels, while the remaining three exhibited modest reductions. The post-depletion titre ratio was inversely correlated with the proportion of cross-reactive memory B cells (Extended Data Fig. 3d), demonstrating consistency between cellular and serological assays and reflecting individual heterogeneity in imprinting strength.

We next compared serum neutralization profiles between the two mouse models (Fig. 2d and Extended Data Fig. 4a). Initially, following Wuhan priming, humanized mice mounted a significantly more potent and broad response than wild-type mice. Two BA.5 boosters successfully shifted the NAb preferences of wild-type mice toward Omicron variants. In contrast, the neutralization curves of humanized mice closely mirrored those of the mRNA cohort, with titres remaining maximal against D614G despite repeated Omicron exposure, indicating strong immune imprinting that could not be overcome by two doses of BA.5 boosting.   (Extended Data Fig. 4a). Together, these data collectively demonstrate that humanized mice can mirror SARS-CoV-2 imprinting. This divergence between the two mouse models strongly suggests that the presence of human immunoglobulin V(D)J genes is a critical determinant of the imprinting phenotype.

Distinct antibody landscapes of mouse models

To dissect the molecular basis of imprinting, we interrogated the evolution of antibody repertoire from the two mouse models at single-B-cell resolution. We sorted RBD-specific memory B cells from spleens of humanized and wild-type mice following Wuhan priming (sorted on Wuhan RBD) and after each BA.5 booster (sorted on BA.5 RBD; Supplementary Information Fig. 1c). Single-cell V(D)J sequencing of paired heavy- and light-chain variable regions generated 583 unique monoclonal antibodies from the six mice groups, which were recombinantly expressed as human IgG1 (Supplementary Table 2). Consistent with the established imprinting signature, antibodies from humanized mice showed a higher proportion of cross-reactivity than those from wild-type mice (Extended Data Fig. 5a and 5b).

To systematically dissect the divergence in antibody epitope distribution between humanized and wild-type mice, we employed high-throughput yeast-display-based deep mutational scanning (DMS) to map the RBD mutations that could escape the isolated mAbs and define the epitope targeted [28,41,42]. We built single-site saturation mutant libraries based on the Wuhan-Hu-1 and BA.5 RBDs and performed DMS for antibodies isolated following Wuhan priming or BA.5 boosting, respectively. The resulting escape profiles of 583 monoclonal antibodies define the critical residues that mediate immune evasion and facilitating their precise categorization into distinct epitope clusters.

IC50-weighted escape profiles show that neutralizing antibodies from wild-type and humanized mice exhibit distinct escape hotspot sites (Fig. 3a-3c and Extended Data Fig. 6a-6c). Following Wuhan priming, the NAbs elicited in humanized mice substantially focused on Class 1 epitopes, featured by hotspots at residues 417, 456, 460, and 473. Although wild-type mice displayed a comparable landscape after Wuhan priming, they lacked these Class 1 peaks, instead featuring residue 477 as a characteristic hotspot (Fig. 3a). BA.5 boosting leads to further divergence between the two mouse models. In humanized mice, the prominent Class 1 hotspots (e.g., residue 456) persisted obstinately throughout the boosting regimen. In contrast, the wild-type repertoire diverged significantly upon boosting, characterized by the emergence of distinct Class 1/4 escape

Figure 3

Figure 3 | Distinct antibody landscapes between mRNA vaccinated humanized and wild-type mice.
a–c, Normalized average DMS escape scores for mAbs isolated from humanized (top) and wild-type (bottom) mice following Wuhan priming (a), the first BA.5 booster (b), or the second BA.5 booster (c). Escape scores were aggregated and weighted by the IC_{50} of each individual mAb (To focus on the neutralizing mAbs, lower IC_{50} corresponds to greater weight) against D614G for Wuhan-primed groups or BA.5 for BA.5-boosted groups. Codon constraints were applied (see Methods). In each panel, the ten residues with the highest cumulative escape scores are annotated, with their specific mutational escape profiles visualized as logo plots above the scatter plots. To highlight divergent immune pressure, residues exhibiting pronounced differences between humanized and wild-type strains (as identified in Extended Data Fig. 6) are colored red in both the scatter plots and logo plots.

peaks, such as residue 504 (Fig. 3b-3c).

To translate these differences in escape sites into a more intuitive view of epitope-level repertoire

differences, we categorized the 583 mAbs into 9 epitope groups (Supplementary Information Fig.

2a) [28]. Names of the epitope groups were generally assigned in line with the epitope groups on

Wuhan RBD defined previously [43,44]. We then comprehensively analyzed the antibody repertoires

elicited in humanized and wild-type mice after each timepoint (Fig. 4a and 4b, Extended Data Fig.

7a-7d).   Following Wuhan priming, humanized mice uniquely elicited a dominant population of

antibodies targeting the A1 epitope (Fig. 4a). Notably, antibodies targeting this epitope are closely

related to the IGHV3-53 germline and target key residues such as 456, aligning with the escape

hotspots. The foundational divergence between the two mouse models was established immediately

after Wuhan priming. After two BA.5 boosting doses, while dynamic shifts occurred—including the

abrogation of Group B and the emergence of Group A2—the A1 response remained persistent in

humanized mice (Fig. 4c and Extended Data Fig. 7a and 7c). In contrast, Omicron-specific

antibodies in epitope groups B, D, and F3 contribute mostly to the neutralization against BA.5

exclusively in wild-type but not the V(D)J-humanized mice.

Overall, DMS analysis demonstrates that Class 1 (A1) antibodies play a pivotal role in the response

to Wuhan priming, serving as a defining characteristic of the V(D)J-humanized mouse model,

compared to the wild-type mice. This antibody subset substantially contributed to the polyclonal

neutralization landscape and could be robustly recalled upon BA.5 boosting. Notably, these A1

antibodies exhibited remarkable germline convergence, consistently dominated by IGHV3-53 after

both Wuhan priming and subsequent BA.5 boosting (Fig. 4a). This mirrors the IGHV3-53/3-66

dominance observed in mRNA-vaccinated human cohorts. Collectively, these findings led us to

hypothesize that the dominant, IGHV3-53/66-driven A1 antibody response induced by Wuhan

priming is the mechanistic driver of SARS-CoV-2 immune imprinting.

Molecular mechanism of imprinting

Given that immune imprinting relies on the recall of pre-existing memory B cells, we then confirmed

the cross-reactivity of these IGHV3-53/66-encoded A1 antibodies to determine their potential for

Figure 4

Figure 4 | VDJ germline difference shapes vaccination- induced SARS-CoV-2 antibody epitope distribution.
Epitope distribution of the antibody repertoire generated after Wuhan priming in humanized (a) and wild-type mice (b). The Wuhan RBD structure (PDB: 6m0j) is displayed as a surface map colored by normalized aggregate escape scores, with epitope groups outlined in distinct colors. Their neutralizing category (Strong, Weak, or Non-neutralizing), antibody count, and percentage are indicated. Key escape residues for each group are displayed as logos, and the sites with the highest escape scores per group are labeled on the structure. Paired heavy and light chain V-gene usage for each epitope is shown in Sankey plots. c, Pyramidal bar charts showing the proportional distribution of epitope groups in antibodies isolated from humanized and wild-type mice after Wuhan priming and after one or two BA.5 boosts. Bars are colored according to the log10 geometric mean IC_{50} of antibodies within each group.

reactivation by the Omicron booster (Fig. 5a). 38% of A1 mAbs from Wuhan-primed humanized

mice were cross-reactive, constituting a pre-existing memory pool that was efficiently recalled to

achieve 100% cross-reactivity upon BA.5 boosting. In contrast, the negligible A1 response in

primed wild-type mice was entirely Wuhan-specific, making subsequent boosters elicited only rare,

de novo BA.5-specific A1 antibodies. Competitive SPR mapping revealed that A1 antibodies

compete with ACE2 and the majority of antibodies targeting neutralizing epitopes (A2, B, C, D4,

F3). Conversely, they showed minimal to no competition with Group D1 or the weakly or non-

neutralizing E and F antibodies (Fig. 5b).

These results indicate that A1 antibodies can sterically mask most other major neutralizing sites on

Omicron RBD from access by other antibodies and their corresponding B cell receptors. Enabled

by the IGHV3-53/66 germline genes, Wuhan priming induces a high-frequency pool of these potent

"masking" antibodies, the abundance of which scales directly with the intensity of the priming.

Critically, a significant portion of this pool is not escaped by Omicron and remains cross-reactive,

making it available for recall upon subsequent Omicron exposure. Therefore, during Omicron

vaccination, this pre-existing and cross-reactive memory B cell population is preferentially

reactivated and expanded. This potent, recalled response actively outcompetes and suppresses the

de novo activation of B cells targeting novel, Omicron-specific epitopes by masking, providing a

direct mechanistic basis for the strong immune imprinting observed in humanized mice. (Fig. 5c).

To test whether these antibodies can indeed drive strong imprinting *in vivo*, we performed passive

antibody transfer experiments in wild-type mice (Fig. 5d). Wuhan-primed wild-type mice were

infused with BD55-1205 (a representative IGHV3-66-encoded A1 broad neutralizing antibody [45])

at various doses and formats (human or mouse IgG1) one day prior to BA.5 boosting (Fig. 5d-5e).

Controls received either PBS or a non-neutralizing antibody (BD57-2665, a representative F1

antibody targeting the cryptic sites of RBD, Fig. 5e). Flow cytometry analysis of lymph nodes

showed that mice receiving 400 μg or 200 μg BD55-1205 hIgG1 and 200 μg BD55-1205 mIgG1

developed pronounced imprinting, evidenced by significantly elevated frequencies of cross-reactive

memory B cells at both post-boost time points compared with PBS controls (Fig. 5f). After the first

BA.5 boost, GC B cells in BD55-1205–treated mice also showed markedly higher cross-reactivity,

demonstrating that early maturation and clonal expansion within the GC was dominated by recalled,

Figure 5

Figure 5 | IGHV3-53/66-encoded Class 1 antibody drives SARS-CoV-2 imprinting through epitope masking.
a, Donut plots showing the cross-reactivity of A1 antibodies from humanized or wild-type mice. The number of antibodies are indicated in the centre of the donut. b, Heatmap of competitive SPR for various antibody groups. The definition of the competition score is described in the Methods section. c, Schematic of the molecular mechanism by which pre-existing IGHV3-53/66-encoded A1 antibodies cause strong immune imprinting. d, Schematic of the antibody passive transfer experiment. Timing of mRNA vaccinations, antibody injection, blood collection, and FACS analysis are indicated. Mice were divided into experimental groups (receiving 400 µg BD55-1205 hIgG1, 200 µg BD55-1205 hIgG1, or 200 µg BD55-1205 mIgG1) and control groups (receiving 200 µg BD57-2665 mIgG1 or PBS). The number of mice per group is indicated at the endpoint. e, DMS escape map logoplots for BD55-1205 and BD57-2665 and their projection onto the SARS-CoV-2 Wuhan RBD (PDB: 6m0j). f, Scatter plots showing the proportion of cross-reactive memory B cells in draining lymph nodes after one (left) or two (right) BA.5 boosts. g, Scatter plots showing the proportion of cross-reactive germinal center B cells in draining lymph nodes after one BA.5 boost. h, Serum IgG midpoint titer against BA.5 RBD before and after Wuhan RBD depletion. Statistical significance of the fold-reduction in titers was assessed between groups. Dashed lines indicate the limit of detection (midpoint titre = 100). Two-tailed Wilcoxon rank-sum tests were used in f-h.

imprint-driven B cells (Fig. 5g). In contrast, BD57-2665 mIgG1–treated mice showed no significant change in cross-reactive memory B cell or GC B cell frequencies, indicating that non-neutralizing antibody blockade does not induce imprinting, ruling out the possibility that the transferred antibody simply depleted the vaccine antigen and prevented a successful BA.5 immunization (Fig. 5f-5g). Correspondingly, Wuhan-RBD depletion of serum from BD55-1205–treated mice produced a more severe drop in BA.5 IgG titres, whereas BD57-2665 mIgG1 and PBS groups showed minimal titre reduction (Fig. 5h). Notably, the mIgG1 format induced a significantly stronger imprinting phenotype than hIgG1, which is likely attributable to the species-matched Fc region (Fig. 5f-5h). Pharmacokinetic analysis confirmed that the passively transferred antibodies had decayed to near-undetectable levels by the time of analysis, thus eliminating their potential interference with ELISA measurements (Extended Data Fig. 8a-8b). Importantly, we demonstrated that the suppressive effect of BD55-1205 on the BA.5-specific response was dose-dependent, with doses as low as 12.5 μg being sufficient to achieve inhibition (Extended Data Fig. 8c-8d). Together, these results confirm that IGHV3-53/66-encoded A1 neutralizing antibodies not only compete with other neutralizing antibodies *in vitro* but also suppress the emergence of Omicron-specific B cell and antibody responses to induce severe imprinting *in vivo*.

This explains why immune imprinting is recapitulated in V(D)J-humanized mice but not in wild-type mice—only the former harbor this human germline gene required to produce such imprinting-prone antibodies. Similarly, in mRNA-vaccinated individuals, the potent Wuhan priming response by mRNA vaccination (elicited by monovalent Wuhan vaccines and, in certain populations, bivalent Wuhan/BA.1 or Wuhan/BA.5 vaccines) induces a larger pool of IGHV3-53/66-encoded A1 antibodies, thereby establishing pronounced imprinting. In contrast, recipients of inactivated vaccines typically experience weaker priming responses and, following stringent containment policies during 2021-2022, undergo substantial waning of Wuhan antibodies. This attenuated pre-existing immunity, leading to lower IGHV3-53/66 antibody abundance, fails to fully suppress subsequent Omicron-specific responses, resulting in more flexible and adaptive humoral evolution.

IGHV3-53 knock-in mouse model

Since wildtype mice could not recapitulate human SARS-CoV-2 immune imprinting observed in mRNA vaccine recipients, evaluating SARS-CoV-2 vaccine updates and broad-spectrum vaccine design in mice is highly problematic. Although V(D)J-humanized mice could capture human imprinting phenotypes, its significant cost creates a pressing need for a more accessible and cost-effective model for vaccine evaluation. Building on our finding that the human IGHV3-53/66 germline is the primary driver of immune imprinting, we engineered a knock-in mouse model by replacing the murine Ighv3-1 gene with human IGHV3-53. This generated both heterozygous (IGHV3-53$^{+/-}$) and homozygous (IGHV3-53$^{+/+}$) mice. High-throughput Genome-wide Translocation Sequencing (HTGTS) of the naïve B cell repertoire confirmed the successful knock-in, revealing IGHV3-53 usage at 0.5% in homozygous and 1.5% in heterozygous mice, with a corresponding ablation of IGHV3-1 usage (0.1% and 0.0%, respectively, Fig. 6a).

To test the model, we immunized wild-type, heterozygous, and homozygous mice using the same regimen of three Wuhan primes followed by two BA.5 boosts (Fig. 6b). Subsequent flow cytometry analysis and depletion assays confirmed that both IGHV3-53$^{+/-}$ and IGHV3-53$^{+/+}$ mice faithfully recapitulated the strong immune imprinting phenotype (Fig. 6c-6d). The serum neutralization profiles of IGHV3-53$^{+/-}$ and IGHV3-53$^{+/+}$ mice also showed a trend remarkably similar to that of the V(D)J-humanized mice (Fig. 6e and Extended Data Fig. 9a). These results demonstrate that both heterozygous and homozygous IGHV3-53 knock-in mice faithfully recapitulate the strong immune imprinting phenotype.

We subsequently utilized this model to simulate the real-world SARS-CoV-2 exposure history characteristic of mRNA-vaccinated populations (Fig. 6f). Specifically, mice received a two-dose Wuhan priming series, mimicking the standard primary course. This was followed by sequential boosters with bivalent Wuhan/BA.1 and Wuhan/BA.5 vaccines, a monovalent XBB.1.5 vaccine, and a monovalent JN.1 vaccine, corresponding to the recommended mRNA booster updates during the Omicron era. These regimens effectively reconstruct the antigenic trajectory of SARS-CoV-2 evolution encountered by humans. Compared to wild-type controls, IGHV3-53$^{+/+}$ mice recapitulated a more pronounced immune imprinting phenotype following the six-dose regimen. Specifically, while retaining potent neutralization against D614G and early Omicron variants (antecedent to JN.1, Fig. 6g), IGHV3-53$^{+/+}$ mice exhibited a compromised breadth of neutralization against the newly

Figure 6

Figure 6 | IGHV3-53 knock-in mouse faithfully reflects SARS-CoV-2 antibody map in human.
a, Usage percentage of IGHV3-53 and Ighv3-1 in naïve B cells from HTGTS sequencing of IGHV3-53[+/+] (balb/c), IGHV3-53[+/-] (balb/c* C57B6/J), and wild-type (C57B6/J) mice. b, Schematic of the immunization regimen and time points for blood collection and FACS analysis of the three mouse models involved in this study. The number of mice is indicated above the timeline. c, Scatter plots showing the proportion of cross-reactive memory B cells in draining lymph nodes of the three mouse models after two BA.5 boosts. d, Serum IgG midpoint titer of the three mouse strains after two BA.5 boosts against BA.5 RBD before and after Wuhan RBD depletion. Statistical significance of the fold-reduction in titers was assessed between groups. Dashed lines indicate the limit of detection (midpoint titre = 100). e, Serum neutralization titers (NT_{50}) of the three mouse strains after one (left) or two (right) BA.5 boosts against a panel of SARS-CoV-2 variant pseudoviruses. Geometric mean titres (GMTs) are shown on the top. Dashed lines indicate the limit of detection ($NT50 = 60$). f, Schematic of the immunization regimen simulating real-world SARS-CoV-2 exposure history. g, Serum neutralization titers (NT_{50}) of IGHV3-53[+/+] and wild-type mice against a panel of SARS-CoV-2 variant pseudoviruses. Geometric mean titres (GMTs) are shown on the top. Dashed lines indicate the limit of detection ($NT_{50} = 180$). h, Antigenic cartography was performed using mouse (top) and human (bottom) plasma neutralization data. Each square indicates a plasma sample, and each circle indicates a SARS-CoV-2 variant. Two-tailed Wilcoxon rank-sum tests were used in c-e and g.

emerged JN.1 sublineages. Specifically, titers against KP.2, KP.3, LP.8.1.1, NB.1.8.1, and XFG were significantly suppressed compared to those in wild-type mice, indicating an inhibition of de novo responses to the newly emerged Omicron-specific epitopes in these variants. (Fig. 6g). Notably, IGHV3-53$^{+/+}$ mice showed higher neutralization titres against BA.3.2 than wild-type mice (Fig. 6g), mirroring the specific serological feature observed in the human mRNA cohort (Fig. 1e). Similar trends were observed after five doses (Extended Data Fig. 10a). Furthermore, comparing neutralization profiles pre- and post-JN.1 boost revealed a strong back-boosting effect in IGHV3-53$^{+/+}$ mice, where JN.1 immunization significantly elevated titres against antecedent strains (pre-JN.1). In contrast, wild-type mice displayed no such effect and even exhibited a significant decline in D614G titres (Extended Data Fig. 10b).

Consequently, the two mouse models exhibited distinct susceptibility to viral escape. Following XBB.1.5 immunization, neutralization against EG.5 was significantly reduced compared to XBB.1.5 in IGHV3-53$^{+/+}$ mice, a reduction not seen in wild-type mice (Extended Data Fig. 10c). Similarly, after the JN.1 boost, IGHV3-53$^{+/+}$ mice showed significant susceptibility to escape by KP.2 and KP.3 relative to JN.1, whereas wild-type mice did not. Consistently, the fold-reduction in titres against XFG relative to NB.1.8.1 was greater in IGHV3-53$^{+/+}$ mice than in wild-type mice (Extended Data Fig. 10c). This result faithfully recapitulates the serological divergence observed between the human mRNA-vaccinated and inactivated-only cohorts (Fig. 1e), and matches the variant escaping profiles of V3-53/66 Class 1 NAbs. By leveraging the genetic precision of the IGHV3-53 knock-in model to eliminate confounding immune histories, we definitively confirm that the imprinted, V3-53/66-dominated response inherently renders mRNA recipients more susceptible to immune evasion by such emerging variants

To visualize how IGHV3-53$^{+/+}$ mice could help with SARS-CoV-2 vaccine update evaluation, we constructed antigenic cartography using serum neutralization data from both mouse models and human cohorts (Fig. 6h). The resulting maps revealed striking topological differences. Wild-type mice exhibited a 'condensed' antigenic landscape similar to the inactivated-only cohort, characterized by short antigenic distances between all JN.1 sublineages. In contrast, the antigenic map of IGHV3-53$^{+/+}$ mice closely mirrors that of mRNA-vaccinated cohort, with both displaying substantial antigenic distances between JN.1, KP.2, KP.3/LP.8.1.1, and XFG, supporting vaccine

antigen updates. Crucially, only the IGHV3-53$^{+/+}$ model faithfully reproduces the substantial antigenic distance (~2) between XFG and LP.8.1.1 characteristic of the mRNA cohort. These findings further implicate the IGHV3-53 knock-in mice could faithfully recapitulates human SARS-CoV-2 immune imprinting, and can serve as a superior model for preclinical SARS-CoV-2 vaccine evaluation.

Discussion

Overall, our study establishes that the abundance of pre-existing IGHV3-53/66-encoded Class 1 antibodies is the primary determinant of SARS-CoV-2 immune imprinting severity. Specifically, three converging factors—the inherent potent immunogenicity of mRNA platforms in human [46], the widespread administration of bivalent mRNA boosters containing Wuhan components in Western populations, and the substantial antibody waning in inactivated vaccine recipients due to strict containment policies (e.g., zero-COVID in China) [47–49]—sufficiently infer that mRNA vaccinees harbored a significantly higher abundance of IGHV3-53/66-encoded Class 1 antibodies prior to their Omicron exposure. Consequently, this mRNA-vaccinated population exhibits a more robust humoral imprinting phenotype. As these distinct humoral immune landscapes became established, a parallel divergence in viral evolution emerged, characterized by the regional predominance of XFG and NB.1.8.1. This co-evolution highlights the necessity of incorporating regional immunological differences into vaccine update strategies.

Importantly, we developed an accessible and robust IGHV3-53 knock-in mouse model that overcomes the limitations of wild-type mice, which fail to recapitulate the SARS-CoV-2 imprinting phenotype. By simulating real-world SARS-CoV-2 exposure histories in this model, we successfully mapped the humoral response landscape of mRNA human vaccinees in mice. Moving forward, this model should be leveraged to conduct parallel evaluations of boosting regimens derived from diverse, newly emerging variants. Such comparative studies will provide the essential data support required to guide the further updating and optimization of SARS-CoV-2 vaccines. Furthermore, as a platform that explicitly exposes the 'imprinted barrier' of mRNA recipients, it would enable the future research of vaccine candidates capable of overcoming such entrenched imprinting.

Data and code availability

Information of individuals providing blood samples and isolated mAbs involved in this study have been included in the Supplementary Tables. All data and materials presented in this manuscript are available from the lead contact upon a reasonable request under a completed Material Transfer Agreement. Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, yunlongcao@pku.edu.cn (Y.C.).

Acknowledgments

We thank all volunteers who provided blood samples. This project is financially supported by the Ministry of Science and Technology of China (2022ZD0115002, Y.C.) and Changping Laboratory (2025D-04-01 to Y.C.).

Author contributions

Y.C. designed and supervised the study. X.N., F.J. and Y.C. wrote the manuscript with input from all authors. X.N., W.S., R.A. and Y.W. performed B cell sorting, single-cell V(D)J sequencing experiments and data analysis. X.N., Y.L., K.L., S.L., R.K., X.C., R.A., and Y.W. performed FACS analysis. H.S. and F.J. obtained and analyzed the DMS data. J.W. and F.S. performed antibody expression. W.W. constructed mRNA vaccines and conducted mouse immunization. Y.Y. constructed the pseudotyped virus. L.Y. performed the pseudovirus neutralization assays, ELISAs and SPR experiments. Y.H., S.L., X.N. and Y.L. performed the HTGTS-Rep-seq and data analysis. B.Z., T.Z., J.H., and X.T. recruited the patients and collected the blood samples.

Declaration of interests

Provisional patents related to the antibodies mentioned in this paper have been filed. Y.C. is a co-founder of Singlomics Biopharmaceuticals. Other authors declare no competing interests.

References

1. Addetia, A. *et al.* Neutralization, effector function and immune imprinting of Omicron

variants. *Nature* **621**, 592–601 (2023).

2.   Raharinirina, N. A. *et al.* SARS-CoV-2 evolution on a dynamic immune landscape. *Nature*
639, 196–204 (2025).

3.   Mellis, I. A. *et al.* Antibody evasion and receptor binding of SARS-CoV-2 LP.8.1.1,
NB.1.8.1, XFG, and related subvariants. 2025.07.18.662329 Preprint at
https://doi.org/10.1101/2025.07.18.662329 (2025).

4.   Carabelli, A. M. *et al.* SARS-CoV-2 variant biology: immune escape, transmission and
fitness. *Nat. Rev. Microbiol.* **21**, 162–177 (2023).

5.   Yuan, M. & Wilson, I. A. Structural Immunology of SARS-CoV-2. *Immunol. Rev.* **329**,
e13431 (2025).

6.   Holmes, E. C. The Emergence and Evolution of SARS-CoV-2. *Annu. Rev. Virol.* **11**, 21–42
(2024).

7.   Telenti, A., Hodcroft, E. B. & Robertson, D. L. The Evolution and Biology of SARS-CoV-2
Variants. *Cold Spring Harb. Perspect. Med.* **12**, a041390 (2022).

8.   Markov, P. V. *et al.* The evolution of SARS-CoV-2. *Nat. Rev. Microbiol.* **21**, 361–379 (2023).

9.   Guo, C. *et al.* Antigenic and virological characteristics of SARS-CoV-2 variants BA.3.2,
XFG, and NB.1.8.1. *Lancet Infect. Dis.* **25**, e374–e377 (2025).

10.   Liu, J. *et al.* Enhanced immune evasion of SARS-CoV-2 variants KP.3.1.1 and XEC through
N-terminal domain mutations. *Lancet Infect. Dis.* **25**, e6–e7 (2025).

11.   Liu, J. *et al.* Virological and antigenic characteristics of SARS-CoV-2 variants LF.7.2.1, NP.1,
and LP.8.1. *Lancet Infect. Dis.* **25**, e128–e130 (2025).

12.   García-Montero, C. *et al.* An Updated Review of SARS-CoV-2 Vaccines and the
Importance of Effective Vaccination Programs in Pandemic Times. *Vaccines* **9**, 433 (2021).

13.   Yang, Z.-R. *et al.* Efficacy of SARS-CoV-2 vaccines and the dose–response relationship
with three major antibodies: a systematic review and meta-analysis of randomised controlled
trials. *Lancet Microbe* **4**, e236–e246 (2023).

14.   Creech, C. B., Walker, S. C. & Samuels, R. J. SARS-CoV-2 Vaccines. *JAMA* **325**, 1318–1320
(2021).

15.   Kyriakidis, N. C., López-Cortés, A., González, E. V., Grimaldos, A. B. & Prado, E. O. SARS-
CoV-2 vaccines strategies: a comprehensive review of phase 3 candidates. *Npj Vaccines* **6**, 28
(2021).

16.   Krammer, F. SARS-CoV-2 vaccines in development. *Nature* **586**, 516–527 (2020).

17.   Thanh Le, T. *et al.* The COVID-19 vaccine development landscape. *Nat. Rev. Drug Discov.*
19, 305–306 (2020).

18.   Liang, C.-Y. *et al.* Imprinting of serum neutralizing antibodies by Wuhan-1 mRNA
vaccines. *Nature* 1–3 (2024) doi:10.1038/s41586-024-07539-1.

19.   Tortorici, M. A. *et al.* Persistent immune imprinting occurs after vaccination with the
COVID-19 XBB.1.5 mRNA booster in humans. *Immunity* **57**, 904-911.e4 (2024).

20.   Wang, Q. *et al.* Deep immunological imprinting due to the ancestral spike in the current
bivalent COVID-19 vaccine. *Cell Rep. Med.* **4**, (2023).

21.   Reynolds, C. J. *et al.* Immune boosting by B.1.1.529 (Omicron) depends on previous
SARS-CoV-2 exposure. *Science* **377**, eabq1841 (2022).

22.   Chemaitelly, H. *et al.* Long-term COVID-19 booster effectiveness by infection history and
clinical vulnerability and immune imprinting: a retrospective population-based cohort study.

Lancet Infect. Dis. **23**, 816–827 (2023).

23. Wang, Z. *et al.* Memory B cell responses to Omicron subvariants after SARS-CoV-2

mRNA breakthrough infection in humans. *J. Exp. Med.* **219**, e20221006 (2022).

24. Wang, Q. *et al.* XBB.1.5 monovalent mRNA vaccine booster elicits robust neutralizing

antibodies against XBB subvariants and JN.1. *Cell Host Microbe* **32**, 315-321.e3 (2024).

25. Johnston, T. S. *et al.* Immunological imprinting shapes the specificity of human antibody

responses against SARS-CoV-2 variants. *Immunity* **57**, 912-925.e4 (2024).

26. Yisimayi, A. *et al.* Prolonged Omicron-specific B cell maturation alleviates immune

imprinting induced by SARS-CoV-2 inactivated vaccine. *Emerg. Microbes Infect.* **13**, 2412623

(2024).

27. Chen, Y. *et al.* SARS-CoV-2 Omicron infection augments the magnitude and durability

of systemic and mucosal immunity in triple-dose CoronaVac recipients. *mBio* **15**, e02407-23

(2024).

28. Yisimayi, A. *et al.* Repeated Omicron exposures override ancestral SARS-CoV-2 immune

imprinting. *Nature* **625**, 148–156 (2024).

29. Yang, X. *et al.* Immune imprinting toward SARS-CoV-2 XBB: implications for vaccine

strategy and variant risk assessment. *Signal Transduct. Target. Ther.* **10**, 372 (2025).

30. Tiller, T. *et al.* Efficient generation of monoclonal antibodies from single human B cells by

single cell RT-PCR and expression vector cloning. *J. Immunol. Methods* **329**, 112–124 (2008).

31. Yuan, M. *et al.* Structural basis of a shared antibody response to SARS-CoV-2. *Science*

https://doi.org/10.1126/science.abd2321 (2020) doi:10.1126/science.abd2321.

32. Wang, Y. *et al.* A large-scale systematic survey reveals recurring molecular features of

public antibody responses to SARS-CoV-2. *Immunity* **55**, 1105-1117.e4 (2022).

33. Li L. *et al.* Breakthrough infection elicits hypermutated IGHV3-53/3-66 public antibodies

with broad and potent neutralizing activity against SARS-CoV-2 variants including the

emerging EG.5 lineages. *PLOS Pathog.* **19**, e1011856 (2023).

34. Vanshylla, K. *et al.* Discovery of ultrapotent broadly neutralizing antibodies from SARS-

CoV-2 elite neutralizers. *Cell Host Microbe* **30**, 69-82.e10 (2022).

35. Yan, Q. *et al.* Germline IGHV3-53-encoded RBD-targeting neutralizing antibodies are

commonly present in the antibody repertoires of COVID-19 patients. *Emerg. Microbes Infect.*

10, 1097–1111 (2021).

36. Zhang, Q. *et al.* Potent and protective IGHV3-53/3-66 public antibodies and their shared

escape mutant on the spike of SARS-CoV-2. *Nat. Commun.* **12**, 4210 (2021).

37. Nutalai, R. *et al.* Potent cross-reactive antibodies following Omicron breakthrough in

vaccinees. *Cell* **185**, 2116-2131.e18 (2022).

38. Tan, T. J. C. *et al.* Sequence signatures of two public antibody clonotypes that bind SARS-

CoV-2 receptor binding domain. *Nat. Commun.* **12**, 3815 (2021).

39. Jian, F. *et al.* Evolving antibody response to SARS-CoV-2 antigenic shift from XBB to JN.1.

Nature 1–3 (2024) doi:10.1038/s41586-024-08315-x.

40. Lee, E.-C. *et al.* Complete humanization of the mouse immunoglobulin loci enables

efficient therapeutic antibody discovery. *Nat. Biotechnol.* **32**, 356–363 (2014).

41. Cao, Y. *et al.* Omicron escapes the majority of existing SARS-CoV-2 neutralizing

antibodies. *Nature* **602**, 657–663 (2022).

42. Jian, F. *et al.* Evolving antibody response to SARS-CoV-2 antigenic shift from XBB to JN.1.

Nature **637**, 921–929 (2025).

43. Cao, Y. *et al.* BA.2.12.1, BA.4 and BA.5 escape antibodies elicited by Omicron infection.

Nature **608**, 593–602 (2022).

44. Cao, Y. *et al.* Imprinted SARS-CoV-2 humoral immunity induces convergent Omicron

RBD evolution. *Nature* **614**, 521–529 (2023).

45. Jian, F. *et al.* Viral evolution prediction identifies broadly neutralizing antibodies to

existing and prospective SARS-CoV-2 variants. *Nat. Microbiol.* 1–15 (2025)

doi:10.1038/s41564-025-02030-7.

46. Lim, W. W., Mak, L., Leung, G. M., Cowling, B. J. & Peiris, M. Comparative immunogenicity

of mRNA and inactivated vaccines against COVID-19. *Lancet Microbe* **2**, e423 (2021).

47. Xu, W., Wu, J. & Cao, L. COVID-19 pandemic in China: Context, experience and lessons.

Health Policy Technol. **9**, 639–648 (2020).

48. Bo, W. *et al.* The current COVID-19 pandemic in China: An overview and corona data

analysis. *Alex. Eng. J.* **61**, 1369–1381 (2022).

49. Gauffin K., Östergren O. & Cederström A. The global response to the pandemic: An

empirical cluster analysis of policies targeting COVID-19. *PLOS ONE* **20**, e0322692 (2025).

50. Yanez Arteta, M. *et al.* Successful reprogramming of cellular protein production through

mRNA delivered by functionalized lipid nanoparticles. *Proc. Natl. Acad. Sci.* **115**, E3351–E3360

(2018).

51. Li, H. *et al.* Establishment of replication-competent vesicular stomatitis virus-based

recombinant viruses suitable for SARS-CoV-2 entry and neutralization assays. *Emerg.*

Microbes Infect. **9**, 2269–2277 (2020).

52. Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. IgBLAST: an immunoglobulin variable domain

sequence analysis tool. *Nucleic Acids Res.* **41**, W34–W40 (2013).

53. Gupta, N. T. *et al.* Change-O: a toolkit for analyzing large-scale B cell immunoglobulin

repertoire sequencing data. *Bioinformatics* **31**, 3356–3358 (2015).

54. Lin, S. G. *et al.* Highly sensitive and unbiased approach for elucidating antibody

repertoires. *Proc. Natl. Acad. Sci.* **113**, 7846–7851 (2016).

55. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-

connected communities. *Sci. Rep.* **9**, 5233 (2019).

Methods

Human serum and PBMC isolation

Blood samples were obtained from convalescent individuals who had received various vaccine

platforms (detailed in Supplementary Table 1). The research protocol and the collection of human

blood samples were approved by the Institutional Review Broad (IRB) of Shenzhen Institute of

Advanced Science, Chinese Academy of Science (Ethics committee archiving no. SIAT-IRB-

230715-H0667) and Beijing Ditan Hospital, Capital Medical University (Ethics committee

archiving no. DTEC-KY2024-112-01). All participants provided their agreement for the collection,

storage and use of their blood samples strictly for research purposes and the subsequent publication

of related data.

To isolate serum and PBMCs, whole blood was diluted 1:1 with PBS (Invitrogen, C10010500BT)

containing 2% (v/v) fetal bovine serum (FBS; Hyclone, SH30406.05) and separated by density

gradient centrifugation using Ficoll (Cytiva, 17-1440-03). The upper serum layer was collected,

aliquoted, and stored at or below -20°C. Before use in assays, serum samples were heat-inactivated

and assessed for neutralizing titres against SARS-CoV-2 variant spike-pseudotyped vesicular

stomatitis virus (VSV).

The PBMC layer was harvested from the interface and processed further. Following red blood cell

lysis (Invitrogen eBioscience 1X RBC Lysis Buffer, 00-4333-57) and washing, PBMCs were either

used immediately or cryopreserved in FBS with 10% (v/v) DMSO (Solarbio, D8371) for storage in

liquid nitrogen. All PBMC samples were transported on dry ice. For B cell isolation, cryopreserved

PBMCs were thawed in PBS supplemented with 5% (v/v) FBS. B cells were subsequently enriched

by immunomagnetic positive selection using the EasySep™ Human CD19 Positive Selection Kit II

(STEMCELL, 17854) following the manufacturer's protocol. The resulting purified B cells were

resuspended in PBS with 2% (v/v) FBS, and cell counts and viability were determined using 0.4%

(w/v) trypan blue staining (Invitrogen, T10282) on a Countess Automated Cell Counter (Invitrogen).

Mice

All animal experiments were conducted under protocols approved by the Animal Welfare Ethics

Committee of HFK Biologics (Approval No. HFK-AP-20250313). The humanized mouse model

(NeoMab-IgG), provided by NeoMab Biotechnology Co., Ltd., was created by in situ replacing the

mouse heavy chain variable region genes and kappa light chain variable region genes with human

genes in BALB/c mouse background. The mouse constant region genes were retained, making sure

the Fc of the immunoglobulins can interact with the Fc receptors expressed on other immune cells

normally, supporting standard immune system development and response. The IGHV3-53$^{+/+}$ or

IGHV3-53$^{+/-}$ mice were generated by Cyagen Biosciences (Suzhou, China). To generate this mice,

the targeting construct replaced the sequence from ATG start codon to exon 3 of mouse Ighv3-1 by

the sequence from ATG start codon to exon 2 of the human IGHV3-53, including the introns. The

Cas9 protein, construct and guide RNAs (gRNA-A1: 5'-TGAGAGTGCTGATTCTTTTGTGG-3',

gRNA-A2: 5'-TGCAGTGCTGCTCTGCAAGGAGG-3' and gRNA-B1: 5'-

TGGCTGTTCACAGCCTTTCCTGG-3', gRNA-B2: 5'-TGGTGCAGTGCTGCTCTGCAAGG-3')

were microinjected into zygotes from BALB/cAnCya wild-type mice. The embryos were

transferred to recipient female mice to obtain F0 mice. The genotype of IGHV3-53$^{+/+}$ or IGHV3-

53$^{+/-}$ mice was confirmed by PCR using two pairs of primers (F4: 5'-

CACCATCTCCAGAGACAATTCCAAG-3', R3: 5'-GTTCTGGATACCAATGTGCCTTCAG-3'

and F3: 5'-TGCAGTCAGTAGCCACCTCGCCAAT-3', R4: 5'-

CATCAGACAGGATACCTTCAGAGAAG-3') and sequencing. Mice of all strains (6–8 weeks old)

were housed under specific-pathogen-free (SPF) conditions, maintained at 22 ± 2°C with 50–60%

humidity on a 12-hour light/dark cycle. All animals had *ad libitum* access to standard chow and

sterile water.

mRNA-LNP Synthesis and Formulation

A PSP73 plasmid bearing the antigen insert followed by a 120-nt poly(T) tract was linearized with

the appropriate restriction enzyme. This DNA served as a template for *in vitro* transcription process

to generate RNA that encoded the SARS-CoV-2 Wuhan, BA.1, BA.5, XBB.1.5 and JN.1 S6P (F817P,

A892P, A899P, A942P, K986P, V987P, R683A and R685A) protein. Linearized DNA template (1

μg) was transcribed for 2 hours at 37°C using the EasyCap T7 Co-transcription Kit with CAG

Trimer (Vazyme, DD4203). Following transcription, the DNA template was digested by incubation

with 1 U of RNase-free DNase I for 15 minutes at 37°C. The resulting mRNA was purified with

VAHTS RNA Clean Beads (Vazyme, N412-03). The concentration and purity of the purified mRNA

were determined by UV spectrophotometry (absorbance at 260 nm and A260/A280), and its

integrity was verified by agarose gel electrophoresis.

The mRNA was encapsulated in a functionalized lipid nanoparticle (LNP) as described previously

[50]. The lipid mixture (oil phase) was prepared by dissolving SM-102, DSPC, cholesterol, and DMG-

PEG2000 in 100 % ethanol at a molar ratio of 50:10:38.5:1.5 to a total concentration of 9.04 mg/mL.

The filter-sterilized (0.22 μm) aqueous phase, consisting of 50 mM sodium citrate buffer (pH 4.0),

and the purified mRNA was diluted within it to a concentration of 133 μg/mL. Oil and aqueous

phases were then rapidly mixed at a 3:1 volume ratio using a staggered herringbone micromixer,

operating at a total flow rate of 12 mL/min to induce LNP self-assembly. The crude LNP suspension

underwent buffer exchange and purification via overnight dialysis at 4°C against a solution of 10

mM Tris-HCl (pH 7.4) with 8-10% (w/v) sucrose, using 10 kDa MWCO dialysis cassettes. The

dialyzed LNPs were subsequently concentrated by centrifugation at 1,500-2,500 × g for 10-15

minutes at 4°C using 10 kDa MWCO ultrafiltration units (ULRC0100150P). The final formulation

was adjusted to a sucrose concentration of 8.7% (w/v), aliquoted, flash-frozen in liquid nitrogen,

and stored at -80°C.

The final product was subjected to rigorous quality control. Key attributes including particle size

(by dynamic light scattering), RNA encapsulation efficiency (by RiboGreen assay), mRNA integrity

(by capillary electrophoresis), osmolality, and endotoxin levels (by LAL assay) were assessed. All

manufactured batches were required to meet the following release criteria: a particle diameter of 80-

100 nm, ≥90% RNA encapsulation, ≥80% mRNA integrity, and an endotoxin level below 1 EU/mL.

Mouse immunization and passive antibody transfer

For immunization, mice were administered mRNA-LNP vaccines. The specific immunization

regimens, including vaccine type, dosage, and timelines, are detailed in the schematics of Figure 2a,

5d, 6b, 6f and Extended Data figure 11a and 11c. On the day of administration, LNP-mRNA vials

were thawed on ice and diluted in sterile 1× PBS to the appropriate concentration. Each mouse was

injected intramuscularly (i.m.) into the quadriceps muscle with 100 μL of the vaccine solution using

a 29 G insulin syringe. To prevent leakage of the inoculum, the needle was held in place for 3–5

seconds post-injection.

For passive antibody transfer, purified monoclonal antibodies (BD55-1205-mIgG1, BD55-1205-

hIgG1, or BD57-2665-mIgG1) with low endotoxin levels (<0.02 EU/mg) were used. The antibodies

were diluted in sterile PBS to the desired concentration for injection. Female BALB/c mice (6–8

weeks old) received a 200 μL dose via intraperitoneal (i.p.) injection in the left lower quadrant using

a 26 G needle. Mice were monitored for at least 5 minutes following the injection, and any

subsequent doses were administered according to the specified experimental schedules.

Pseudovirus preparation and neutralization assay

We generated SARS-CoV-2 variant spike protein pseudovirus as described previously [9–11,28,39,51].

Plasmids encoding a codon-optimized SARS-CoV-2 Spike (S) protein were constructed by inserting

the S gene into the pcDNA3.1 vector. To produce pseudovirus, 293T cells (ATCC, CRL-3216) were

transfected with the S protein expressing plasmids with Lipofectamine 3000 (Invitrogen, L3000015)

and subsequently infected with G*ΔG-VSV (Kerafast, EH1020-PM). After 24 hours, the

supernatant containing the pseudovirus was harvested, filtered through a 0.45 μm filter (Millipore),

aliquoted, and stored at -80°C.

Neutralization assays were performed using the Huh-7 cell line (JCRB, 0403). Monoclonal

antibodies or serum samples were serially diluted in DMEM (Hyclone, SH30243.01) and incubated

with the pseudovirus in 96-well plates for 1 hour at 37°C with 5% CO_2. Following incubation, Huh-

7 cells were seeded into the wells (2×10^4 cells per well) and cultured for an additional 24 hours at

37°C with 5% CO_2. To assess infection levels, the culture supernatant was removed and left 100 μ

l in each well. The Bright-Lite Luciferase Assay Substrate was reconstituted with its corresponding

Assay Buffer (Vazyme, DD1209-03-AB), and this mixture was added to the wells. After incubating

in the dark for 2 minutes, luminescence was measured using a microplate spectrophotometer

(PerkinElmer, HH3400). The NT_{50} or IC_{50} values were determined using a three-parameter logistic

regression model.

Mouse tissue processing and B cell extraction

Following euthanasia, the spleen, inguinal lymph nodes, and popliteal lymph nodes were harvested

and placed in RPMI 1640 culture medium (Invitrogen, C11875500CP) containing 5% (v/v) FBS.

Single-cell suspensions were prepared by mechanical disruption using the plunger of a syringe. The

popliteal and inguinal lymph nodes from each mouse were pooled, ground, and filtered through a

40 μm cell strainer. For the spleen, tissue was processed by the same grinding method and passed

through a 70 μm cell strainer, followed by centrifugation and lysis of red blood cells using 1× RBC

Lysis Buffer (Invitrogen eBioscience, 00-4333-57). After washing steps and centrifugation, the

resulting cell pellets were resuspended in PBS containing 2% (v/v) FBS.

Splenic B cells were enriched from the splenic single-cell suspensions via immunomagnetic

negative selection with the EasySep™ Mouse Pan-B Cell Isolation Kit (STEMCELL, 19844).

Following the manufacturer's protocol, the untouched, purified B cells were collected and washed

in PBS with 2% (v/v) FBS. The cell numbers of total lymph node cells and purified splenic B cells

were determined using 0.4% (w/v) trypan blue stain (Invitrogen, T10282) and a Countess

Automated Cell Counter (Invitrogen).

Flow cytometry analysis and antigen-specific B cell sorting

For the isolation of antigen-specific human B cells, enriched B cell populations from PBMCs were

prepared for fluorescence-activated cell sorting (FACS). The staining panel included FITC anti-

human CD20 (BioLegend, 302304), Brilliant Violet 605™ anti-human CD27 (BioLegend, 302824),

PE/Cyanine7 anti-human IgM (BioLegend, 314532), and PE/Cyanine7 anti-human IgD (BioLegend,

348210). Antigen-specific cells were detected using biotinylated JN.1 RBD (Sino Biological,

40592-V49H16-B) conjugated with PE-streptavidin (BioLegend, 405204) and APC-streptavidin

(BioLegend, 405207), and Wuhan RBD (Sino Biological, 40592-V27H-B) conjugated with BV421-

streptavidin (BioLegend, 405225). The viability dye 7-AAD (Invitrogen, 00-6993-50) was used to

exclude dead cells. A gating strategy was applied to sort single, viable (7-AAD$^-$), class-switched

(IgM$^-$ and IgD$^-$), CD20$^+$CD27$^+$ memory B cells that were positive for JN.1 RBD.

For the characterization of B cell responses in immunized mice, single-cell suspensions from the

inguinal and popliteal lymph nodes were stained for flow cytometry analysis. The cells were stained

with a panel including PE/Cyanine7 anti-mouse CD38 (BioLegend, 102718), Brilliant Violet 605™

anti-mouse/human B220 (BioLegend, 103244), APC/Cyanine7 anti-mouse IgD (BioLegend,

405716), Brilliant Violet 711™ anti-mouse IgM (BioLegend, 406539), and FITC anti-mouse/human

GL7 (BioLegend, 144603). The antigen probe cocktail consisted of biotinylated BA.5 RBD (Sino

Biological, 40592-V49H9-B) conjugated with PE- and APC-streptavidin, and Wuhan RBD

conjugated with BV421-streptavidin. Data for lymph node analysis were acquired on a Symphony

A5SE cytometer (BD Biosciences).

For the isolation of antigen-specific mouse splenic B cells, purified splenic B cells were stained

using the identical panel of antibodies and RBD probes as described for the lymph node analysis.

MoFlo Astrios EQ Cell Sorter (Beckman Coulter) was used for all sorting experiments, targeting

live (7-AAD$^-$), B220$^+$, CD38$^+$, class-switched (IgM$^-$ and IgD$^-$), non-germinal center (GL7$^-$) B cells

that bound to the BA.5 RBD.

For all procedures, data were collected via Summit 6.0 software (Beckman Coulter). Data from all

experiments were uniformly analyzed using FlowJo v10.8 (BD Biosciences).

Single-cell V(D)J sequencing

For the 10X Genomics workflow, sorted antigen-specific B cells, suspended in PBS with 10% (v/v)

FBS, were processed with the Chromium Next GEM Single Cell V(D)J Reagent Kits v1.1 (10X

Genomics, CG000208). The cell suspension was loaded onto a 10X Chromium Controller to

generate Gel Beads-in-Emulsion (GEMs), which facilitate the barcoding of mRNA and subsequent

reverse transcription within individual droplets. Following cDNA synthesis, the product was

purified using a SPRIselect Reagent Kit (Beckman Coulter, B23318) and pre-amplified. Targeted

enrichment of paired V(D)J sequences was then achieved using 10X-specific BCR primers, and the

resulting products were used for sequencing library construction. These final libraries were

sequenced on an Illumina NovaSeq 6000 platform with a NovaSeq 6000 S4 Reagent Kit v1.5 (300

cycles; Illumina, 20028312). The 10X Genomics V(D)J Illumina sequencing data were assembled

as B cell receptor contigs and aligned to the B cell V(D)J reference using Cell Ranger (v.6.1.1)

pipeline. For human-source IGH and IGK contigs, we use GRCh38 as reference. For mouse-source

IGL contigs, we use GRCm38 as reference. Only the productive contigs and B cells with one heavy

chain and one light chain were kept to remove doublets. The germline V(D)J genes were identified

and annotated using IgBlast (v1.17.1) [52]. SHM nucleotides and residues in the antibody variable

domain were detected using Change-O toolkit (v.1.2.0) [53].

For the plate-based method, single antigen-specific B cells were sorted directly into the individual

wells of 96-well plates, each containing 4 µL of lysis buffer (0.095% Triton X-100, 2.5 µM oligo-

dT   primer: 5'-

AAGCAGTGGTATCAACGCAGAGTACTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTVN-3',

2.5 mM dNTPs, 1 U/µL RNase Inhibitor). The plates were immediately processed by vortexing,

centrifugation, and incubation at 72°C for 3 min to denature RNA secondary structures, followed

by rapid chilling on ice. Reverse transcription was performed in each well by adding 5.7 µL of an

RT master mix (final concentrations: 1× HiScript III first-strand buffer, 10 U/µL HiScript III

reverse transcriptase, 2 U/μL RNase inhibitor, 5 mM DTT, 1 M betaine, 6 mM MgCl₂, 1 μM TSO

primer: 5'-AAGCAGTGGTATCAACGCAGAGTACATrGrG+G-3'). The RT reaction was

incubated at 37°C for 60 minutes, followed by enzyme inactivation at 85°C for 5 seconds. The

resulting cDNA served as a template for amplifying full-length immunoglobulin heavy and light

chain V(D)J sequences via a two-round nested PCR strategy using Phanta Max Master Mix (Vazyme,

P515). The first PCR round employed a multiplex primer pool targeting all VH, Vκ, and Vλ gene

families, along with constant region-specific reverse primers. A second, nested PCR round was then

performed with internal primers to enhance specificity and yield. Final amplicons were purified and

subsequently analyzed by Sanger sequencing. The germline V(D)J genes were identified and

annotated using IgBlast (v1.22.0) [52].

HTGTS-Rep-seq

0.5-4 μg of genomic DNA from purified splenic B cells was used for generating HTGTS-Rep-seq

libraries as previously described [54]. Four bait primers that target mouse JH1

(TGACATGGGGAGATCTGAGA), JH2 (CCCCAACAAATGCAGTAAAATCT), JH3

(GAGAATCTTGGTCCTGAAGGC), and JH4 (CTGCAATGCTCAGAAAACTCC) were mixed to

capture all heavy chain (HC) repertoire in one library. All primers carried a 5′ biotin modification

(5BiosG). These HTGTS-Rep-seq libraries were sequenced on an Illumina NextSeq platform (2 ×

150 bp paired-end) DNBSEQ-T7. Data were analyzed with the HTGTS-Rep-seq pipeline [54].

Monoclonal antibody expression and purification

The sequences for the antibody heavy and light chains were initially codon-optimized for expression

in human cells (GenScript). The variable regions (VH and VL) were then separately inserted into

corresponding expression vectors (pCMV3-CH, pCMV3-CL or pCMV3-CK). Plasmids of the

heavy and light chain constructs were transformed into *Escherichia coli* DH5α competent cells

(Tsingke, TSC-C01-96). After overnight incubation at 37°C, single colonies were picked for colony

PCR identification. Plasmid DNA of expanded cultures was extracted (CWBIO,CW2105) after

verified by Sanger sequencing.

For protein production, heavy and light chain plasmids were co-transfected into Expi293F cells

(Thermo Fisher, A14527) using polyethylenimine (PEI; Yeasen, 40816ES03). The plasmid-PEI

complexes were prepared in sterile 0.9% NaCl solution before being added to the cell culture. The

transfected cells were cultured at 36.5°C with 5% CO_2 and 175 rpm shaking for 6–10 days. A

nutrient supplement (OPM Biosciences, F081918-001) was added to each culture at 24 hours post-

transfection and every 48 hours thereafter.

To purify the antibodies, the culture supernatant was first clarified by centrifugation (3,000 × g, 10

minutes). The supernatant was then incubated with Protein A magnetic beads (GenScript, L00695)

for 2 hours to allow antibody binding. The beads were subsequently washed, and the bound

antibodies were eluted using a KingFisher automated purification system (Thermo Fisher). The

concentration of the purified antibody was determined using a NanoDrop spectrophotometer

(Thermo Fisher, 840-317400), and its purity was assessed by SDS-PAGE (LabLead, P42015).

RBD depletion of serum

To deplete RBD-specific antibodies from serum, 50 μL of Dynabeads™ MyOne™ Streptavidin T1

(Invitrogen, 65601) were washed once with PBS. 10 μg of biotinylated SARS-CoV-2 Wuhan RBD

was incubated with the washed beads for 1 hours with gentle rotation to allow binding via

streptavidin-biotin interaction. The beads were then collected using a magnetic rack for 2-3 min, the

supernatant was discarded, and the beads were washed three times with PBS to remove unbound

proteins. Subsequently, 200-400 μL of serum was incubated with the RBD-conjugated beads for

1 hours with gentle rotation to allow specific antibody binding. Finally, the tubes were placed on the

magnetic rack, and the supernatant representing the RBD-depleted serum was carefully collected

for downstream analyses.

Enzyme-linked immunosorbent assays

High-binding 96-well plates (NEST, 504201) were coated overnight at 4°C with SARS-CoV-2

Wuhan (Sino Biological, 40592-V27H-B) or BA.5 RBD proteins (Sino Biological, 40592-V49H9-

B). The following day, plates were washed three times with 1×PBST (Solarbio, P1033) and blocked

with 250 μL 3–5% bovine serum albumin (BSA; Solarbio, A8020) in 1×PBST for 2 hours at 37°C

to prevent non-specific binding. After three additional washes, 100 μL of serially diluted antibodies

or serum samples were added to the wells and incubated for 30 minutes at room temperature.

Unbound antibodies were removed by five washes with 1×PBST. Subsequently, 100 μL of HRP-

conjugated Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody (Invitrogen, G21040)

was added and incubated for 30 minutes at room temperature. Following a final five washes, the

signal was developed by adding 100 μL of TMB substrate (Solarbio, PR1200) to each well and

incubating for 8 minutes in the dark. The reaction was terminated by adding 50 μL of stop solution

(Solarbio, C1058). The optical density (OD) was measured at 450 nm with a reference wavelength

of 630 nm using a Multiskan FC microplate reader (Thermo Scientific). Final absorbance values

were obtained by subtracting the OD630 reading from the OD450 reading for each well.

Surface plasmon resonance

Surface plasmon resonance (SPR) experiments were conducted using a Biacore 8K+ system (Cytiva)

at room temperature. For competitive binding assays, His-tagged SARS-CoV-2 BA.5 RBD protein

(5 μg/mL) was immobilized onto an anti-His-tagged CM5 sensor chip (Cytiva) for 1 minute.

Subsequently, a saturating concentration of the first antibody (Ab1, 20 μg/mL) was injected for 2

minutes, immediately followed by the injection of the second antibody (Ab2, 20 μg/mL) for another

2 minutes. The sensor surface was regenerated between cycles using a glycine solution (pH 1.5).

All binding data were recorded and processed using Biacore 8K Evaluation Software (v4.0.8.20368).

The degree of epitope competition was calculated using the following formula:

781
$$score_{\text{Ab2}-\text{Ab1}} = 1 - \frac{response_{Ab2\ after\ Ab1}}{\overline{response_{Ab2\ alone}}}$$

Where $response_{Ab2\ after\ Ab1}$ represents the response units when Ab2 serves as the second

antibody and Ab1 as the first antibody, whereas $\overline{response_{Ab2\ alone}}$ denotes the mean response

units when Ab2 acts as the first antibody.

DMS library construction

Replicate deep mutational scanning (DMS) libraries of the SARS-CoV-2 BA.5 RBD (residues

N331–T531; Wuhan-Hu-1 numbering) were generated based on established protocols [28,41] with a

modification. Rather than using a pooled primer mix, we performed 201 individual PCR reactions,

each using a specific NNS primer pair to introduce all possible amino acid substitutions at a single

target residue. The resulting products from each single-site mutagenesis reaction were then

combined to form the final comprehensive library. Each unique RBD variant was subsequently

tagged with a distinct 26-nucleotide (N26) barcode via PCR. The mutagenized and barcoded RBD

sequences were then cloned into pETcon 2649 vector and the resulted plasmid libraries were

amplified in electrocompetent *E. coli* DH10B cells. The association between every RBD variant and

its corresponding N26 barcode was established by preparing PacBio sequencing libraries and

performing long-read sequencing on the Sequel II platform. Amplified DMS plasmid libraries were

transformed into *Saccharomyces cerevisiae* strain EBY100. Transformed yeast cells were initially

selected on SD-CAA agar plates. Positive clones were then expanded by culturing in SD-CAA liquid

media. The resulting comprehensive DMS yeast libraries were preserved by flash-freezing in liquid

nitrogen and stored at -80 °C.

Magnetic beads-based antibody mutation escape profiling

High-throughput mutation escape profiling for mAbs was performed using magnetic beads based

on previously established protocols [28,41]. DMS yeast libraries (Wuhan and BA.5) first underwent

functional pre-screening. Non-functional or misfolded RBD variants were removed using ACE2-

biotin conjugate (Sino Biological, 10108-H08H-B) bound to streptavidin magnetic beads (Thermo

Fisher, 11533D). ACE2-bound yeast cells were washed with PBS containing 0.1% (v/v) BSA,

released, expanded in SD-CAA liquid medium, and cryopreserved at -80°C as functional libraries.

For Antibody Escape Selection, thawed functional libraries were cultured overnight in SD-CAA

with shaking, then back-diluted into SG-CAA medium to induce RBD surface expression. Escape

variants of each mAb were isolated using a sequential selection strategy, with 2 rounds of negative

selection to deplete antibody-binding variants and 1 round of positive selection to capture the

antibody-escaping population using anti-c-Myc magnetic beads (Thermo Fisher, 88843). The final

sorted yeasts were washed, regrown overnight and subjected to plasmid extraction using a 96-well

kit (Coolaber, PE053). The unique N26 barcodes appended to each RBD variant were amplified by

PCR using extracted plasmid as template. PCR products were purified with Ampure XP beads

(Beckman Coulter, A63882) and subjected to high-throughput single-end sequencing (NextSeq

500/550 platforms or MGI200 platforms).

Antibody DMS data analysis

The raw sequencing data from the DMS were processed as previously described [28]. Specifically, the

barcode sequences detected from both the antibody-screened and reference libraries were aligned

with a barcode-variant dictionary derived from PacBio sequencing data of the Wuhan and BA.5

DMS libraries using the alignparse (v.0.6.2) and dms_variants (v.1.4.3) tools. Ambiguous barcodes

were excluded during the merging of yeast libraries. Only barcodes detected more than five times

in the reference library were considered for further analysis. The escape score for a variant X,

present in both the screened and reference libraries, was calculated as $F \times (n_{X,ab}/N_{ab})/(n_{X,ref}/N_{ref})$,

where F is a scaling factor to normalize the scores to a 0–1 range, and n and N represent the numbers

of detected barcodes for variant X and the total barcodes in the antibody-screened (ab) or reference

(ref) samples, respectively. For antibodies subjected to DMS with multiple replicates using different

mutant libraries, the final escape score for each mutation was averaged for subsequent analyses.

We used graph-based unsupervised clustering and embedding to assign an epitope group to each

antibody and visualize them in a two-dimensional space. Initially, site escape scores (sum of

mutation escape scores per residue) for each antibody were normalized to a sum of one, representing

a distribution over RBD residues. The dissimilarity between two antibodies is defined based on the

Pearson's correlation coefficient of their escape score vectors using numpy (v1.25.2). A k-nearest-

neighbour graph was constructed using the python-igraph module (v.0.9.6), and Leiden clustering

was applied to assign a cluster to each antibody [55]. Cluster names were manually annotated on the

basis of the characteristic sites in the average escape profiles of each cluster, using the same

nomenclature as our previously published DMS dataset [28]. To visualize the dataset in two

dimensions, uniform manifold approximation and projection was performed based on the k-nearest-

neighbour graph using umap-learn module (v.0.5.2), and figures were generated using R package

ggplot2 (v.3.3.3).

To compute the average immune pressure or identify escape hotspots using a collection of mAb

DMS profiles, we incorporating two types of weight to account for the impact of each mutation on

neutralizing activity and codon constraints at each residue. For codon usage constraints, mutations

inaccessible through single nucleotide changes were assigned a weight of zero, whereas others

received a weight of 1.0. We used Wuhan/D614G (Wuhan-Hu-1 reference genome) and BA.4/5

(EPI_ISL_11207535) to define one-nucleotide-accessible amino acid mutations. Neutralizing

activity weights were calculated as $-\log 10$ (IC_{50}), with IC_{50} values below 0.0005 or above 1.0

adjusted to 0.0005 or 1.0, respectively. Raw escape scores for each antibody were normalized by

the maximum score across all mutants. The weighted score for each antibody and mutation was

obtained by multiplying the normalized scores by the corresponding two weights, and the final

mutation-specific weighted score was the sum of scores for all antibodies in the designated set,

subsequently normalized to a 0–1 range. To visualize the calculated escape maps, sequence logos

were generated using the Python module logomaker (v.0.8).

Figure legends

Figure 1 | mRNA and inactivated vaccine priming induce distinct immune imprinting level.

a, Schematic of the SARS-CoV-2-related immune histories and sampling time of the inactivated-only and mRNA-vaccinated cohorts involved in this study. **b,** Representative flow cytometry dot plots of cross-reactive memory B cells from individuals from the inactivated-only (top) and mRNA-vaccinated (bottom) cohorts. APC, allophycocyanin; PE, phycoerythrin; BV421, Brilliant Violet 421. **c,** Scatter plots showing the proportion of cross-reactive memory B cells (left), JN.1-reactive memory B cells (middle), and Wuhan-reactive memory B cells (right). **d,** Serum IgG midpoint titre against Wuhan (left) or JN.1 (right) RBD before and after Wuhan RBD depletion. Statistical significance of the fold-reduction in titres was assessed between two cohorts. Dashed lines indicate the limit of detection (midpoint titre = 10). **e,** NAb response of the two cohorts against a panel of SARS-CoV-2 variant pseudoviruses. Geometric mean titres (GMTs) are shown on the top. Dashed lines indicate the limit of detection ($NT_{50} = 10$). **f,** Heavy chain V gene usage proportion of the two cohorts. IGHV3-53 and IGHV3-66 were combined for analysis. Two-tailed Wilcoxon rank-sum tests were used in **c-e**. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$; NS, not significant ($P > 0.05$).

Figure 2 | V(D)J humanized mice recapitulates severe SARS-CoV-2 imprinting.

a, Schematic of the mouse immunization regimen and sampling timeline for wild-type and V(D)J-humanized mice. The number of mice is indicated at the time point for each endpoint experiment. **b,** Scatter plots showing the proportion of cross-reactive memory B cells in draining lymph nodes of the two mouse strains after one (left) or two (right) BA.5 boosts. **c,** Serum IgG midpoint titre of the two mouse strains after one (left) or two (right) BA.5 boosts against Wuhan or BA.5 RBD before and after Wuhan RBD depletion. Statistical significance of the fold-reduction in titres was assessed between humanized and wild-type mice. Dashed lines indicate the limit of detection (midpoint titre = 100). **d,** Serum neutralization titres (NT_{50}) of the two mouse models after Wuhan priming (left), one BA.5 boost (middle), or two BA.5 boosts (right) against a panel of SARS-CoV-2 variant pseudoviruses. Geometric mean titres (GMTs) are shown on the top. Dashed lines indicate the limit of detection ($NT_{50} = 60$). Two-tailed Wilcoxon rank-sum tests were used in **b-d**.

Figure 3 | Distinct antibody landscapes between mRNA vaccinated humanized and wild-type mice.

a–c, Normalized average DMS escape scores for mAbs isolated from humanized (top) and wild-type (bottom) mice following Wuhan priming (**a**), the first BA.5 booster (**b**), or the second BA.5 booster (**c**). Escape scores were aggregated and weighted by the IC_{50} of each individual mAb (To focus on the neutralizing mAbs, lower IC_{50} corresponds to greater weight) against D614G for Wuhan-primed groups or BA.5 for BA.5-boosted groups. Codon constraints were applied (see Methods). In each panel, the ten residues with the highest cumulative escape scores are annotated, with their specific mutational escape profiles visualized as logo plots above the scatter plots. To highlight divergent immune pressure, residues exhibiting pronounced differences between humanized and wild-type strains (as identified in Extended Data Fig. 6) are colored red in both the scatter plots and logo plots.

Figure 4 | VDJ germline difference shapes vaccination- induced SARS-CoV-2 antibody epitope distribution.

Epitope distribution of the antibody repertoire generated after Wuhan priming in humanized (**a**) and wild-type mice (**b**). The Wuhan RBD structure (PDB: 6m0j) is displayed as a surface map colored by normalized aggregate escape scores, with epitope groups outlined in distinct colors. Their neutralizing category (Strong, Weak, or Non-neutralizing), antibody count, and percentage are indicated. Key escape residues for each group are displayed as logos, and the sites with the highest escape scores per group are labeled on the structure. Paired heavy and light chain V-gene usage for each epitope is shown in Sankey plots. **c,** Pyramidal bar charts showing the proportional distribution of epitope groups in antibodies isolated from humanized and wild-type mice after Wuhan priming and after one or two BA.5 boosts. Bars are colored according to the log10 geometric mean IC_{50} of antibodies within each group.

Figure 5 | IGHV3-53/66-encoded Class 1 antibody drives SARS-CoV-2 imprinting through epitope masking.

a, Donut plots showing the cross-reactivity of A1 antibodies from humanized or wild-type mice.

The number of antibodies are indicated in the centre of the donut. **b,** Heatmap of competitive SPR

for various antibody groups. The definition of the competition score is described in the Methods

section. **c,** Schematic of the molecular mechanism by which pre-existing IGHV3-53/66-encoded A1

antibodies cause strong immune imprinting. **d,** Schematic of the antibody passive transfer

experiment. Timing of mRNA vaccinations, antibody injection, blood collection, and FACS analysis

are indicated. Mice were divided into experimental groups (receiving 400 μg BD55-1205-hIgG1,

200 μg BD55-1205-hIgG1, or 200 μg BD55-1205-mIgG1) and control groups (receiving 200 μg

BD57-2665-mIgG1 or PBS). The number of mice per group is indicated at the endpoint. **e,** DMS

escape map logoplots for BD55-1205 and BD57-2665 and their projection onto the SARS-CoV-2

Wuhan RBD (PDB: 6m0j). **f,** Scatter plots showing the proportion of cross-reactive memory B cells

in draining lymph nodes after one (left) or two (right) BA.5 boosts. **g,** Scatter plots showing the

proportion of cross-reactive germinal center B cells in draining lymph nodes after one BA.5 boost.

h, Serum IgG midpoint titre against BA.5 RBD before and after Wuhan RBD depletion. Statistical

significance of the fold-reduction in titres was assessed between groups. Dashed lines indicate the

limit of detection (midpoint titre = 100). Two-tailed Wilcoxon rank-sum tests were used in **f-h**.

Figure 6 | IGHV3-53 knock-in mouse faithfully reflects SARS-CoV-2 antibody map in human.

a, Usage percentage of IGHV3-53 and Ighv3-1 in naïve B cells from HTGTS sequencing of IGHV3-

$53^{+/+}$ (balb/c), IGHV3-$53^{+/-}$ (balb/c* C57B6/J), and wild-type (C57B6/J) mice. **b,** Schematic of the

immunization regimen and time points for blood collection and FACS analysis of the three mouse

models involved in this study. The number of mice is indicated above the timeline. **c,** Scatter plots

showing the proportion of cross-reactive memory B cells in draining lymph nodes of the three mouse

models after two BA.5 boosts. **d,** Serum IgG midpoint titre of the three mouse strains after two BA.5

boosts against BA.5 RBD before and after Wuhan RBD depletion. Statistical significance of the

fold-reduction in titres was assessed between groups. Dashed lines indicate the limit of detection

(midpoint titre = 100). **e,** Serum neutralization titres (NT_{50}) of the three mouse strains after one (left)

or two (right) BA.5 boosts against a panel of SARS-CoV-2 variant pseudoviruses. Geometric mean

titres (GMTs) are shown on the top. Dashed lines indicate the limit of detection ($NT_{50} = 60$). **f,**

Schematic of the immunization regimen simulating real-world SARS-CoV-2 exposure history. **g,**

Serum neutralization titres (NT_{50}) of IGHV3-$53^{+/+}$ and wild-type mice against a panel of SARS-

CoV-2 variant pseudoviruses. Geometric mean titres (GMTs) are shown on the top. Dashed lines indicate the limit of detection ($NT_{50} = 180$). **h,** Antigenic cartography was performed using mouse (top) and human (bottom) serum neutralization data. Each square indicates a serum sample, and each circle indicates a SARS-CoV-2 variant. Two-tailed Wilcoxon rank-sum tests were used in **c-e** and **g**.

Extended Data Figure 1 | Different epidemiological landscape of XFG and NB.1.8.1 in 2025.

a, Prevalence of XFG and NB.1.8.1 in China and the rest of the world as of October 5, 2025. The asterisk (*) denotes the specific lineage and its sublineages. Data were retrieved from CovSpectrum (https://cov-spectrum.org/).

Extended Data Figure 2 | Enrichment of IGHV3-53/66 and high SHM rate characterize the mRNA-imprinted antibody repertoire.

a, Clonal analysis of memory B cells from individuals in the mRNA-vaccinated cohort. Single clones are colored white. The numbers of B cells analyzed are labelled in the centre of the donut plots. **b,** Paired heavy and light chain V-gene usage for memory B cells from the mRNA-vaccinated and inactivated-only cohorts. **c, d,** Violin plots comparing somatic hypermutation (SHM) rates between the two cohorts, shown for heavy and light chains (**c**) and representative heavy chain V-genes (**d**). The number of sequences (n) is indicated above each plot. Two-tailed Wilcoxon rank-sum tests were used in **d**.

Extended Data Figure 3 | V(D)J-humanized mice exhibit a cellular imprinting phenotype dominated by cross-reactive GC and memory B cells.

a-c, Scatter plots showing the proportion of cross-reactive B cells (**a**), total GC B cells (**b**), and class-switched memory B cells (**c**). The indicated percentages reflect the frequency relative to their respective parental gates defined in Supplementary Information Fig. 1b. **d,** Linear regression analysis of the correlation between memory B cell cross-reactivity proportion and BA.5-specific IgG titre.

Extended Data Figure 4 | Repeated Omicron boosters fail to elicit robust Omicron-specific

neutralization in imprinted humanized mice.

a, Line plots showing longitudinal pseudovirus neutralization titres (NT_{50}) against a panel of

SARS-CoV-2 variants in paired humanized and wild-type mice across Wuhan priming and BA.5

booster timepoints; lines connect data from the same mouse. Dashed lines indicate the limit of

detection ($NT_{50} = 60$). Two-tailed Wilcoxon rank-sum tests were used.

Extended Data Figure 5 | Neutralization potency and cross-reactivity of mAbs from

humanized and wild-type mice.

a,b, Comparison of antibody IC_{50} values against D614G and BA.5(**a**) and cross-reactivity

proportion (**b**) from humanized and wild-type mice. Two-tailed Wilcoxon rank-sum tests were

used in **a**.

Extended Data Figure 6 | Differential escape hotspots in humanized versus wild-type mice.

a–c, Scatter plots comparing the normalized average DMS escape scores of neutralizing

antibodies isolated from humanized (y-axis) versus wild-type (x-axis) mice following Wuhan

priming (**a**), the first BA.5 booster (**b**), and the second BA.5 booster (**c**). Residues falling along the

diagonal (y=x) indicate shared immune pressure between the two models. To identify divergent

hotspots—including those with low-to-moderate scores in one strain that are absent in the other—

a shaded tolerance region was defined by the boundaries y = 1.4x + 0.3 and y = 0.6x - 0.18. Points

falling outside this shaded region represent distinct escape hotspots. Among these outliers, the five

residues with the highest escape scores in each group are highlighted in red. Related to Fig. 3.

Extended Data Figure 7 | Distinct SARS-CoV-2 antibody epitope distribution after

sequential BA.5 boosts.

a-d, Epitope distribution of the antibody repertoire generated after one BA.5 boost (**a, b**) or two

BA.5 boosts (**c, d**) in humanized mice (**a, c**) and wild-type mice (**b, d**). Related to Figure 4. **e,**

Pyramidal bar charts showing the proportional distribution of epitope groups in antibodies isolated

from humanized and wild-type mice after one or two BA.5 boosts. Bars are colored according to

the log10 geometric mean IC_{50} of antibodies within each group.

Extended Data Figure 8 | Pharmacokinetics and dose-dependent suppressive effects of

antibodies used in passive transfer.

a, Schematic of the *in vivo* monoclonal antibody pharmacokinetic study. **b,** Serum IgG titres over

time following antibody injection (mIgG1 forms were detected using an anti-mouse Fc secondary

antibody, and hIgG1 forms using an anti-human Fc secondary antibody). Dashed lines indicate the

limit of detection (midpoint titre = 100). **c,** Schematic of the BD55-1205 dose-ranging experiment.

d, Corresponding serum IgG titres before and after Wuhan RBD depletion. Statistical significance

of the fold-reduction in titres was assessed between groups. Dashed lines indicate the limit of

detection (midpoint titre = 100). Two-tailed Wilcoxon rank-sum tests were used in **d**.

Extended Data Figure 9 | IGHV3-53 knock-in mice exhibit diminished Omicron

neutralization boosting compared to wild-type mice.

a, Line plots showing longitudinal pseudovirus neutralization titres (NT_{50}) against a panel of

SARS-CoV-2 variants in paired IGHV3-53$^{+/+}$ (top), IGHV3-53$^{+/-}$ (middle), and wild-type (bottom)

mice across BA.5 booster timepoints; lines connect data from the same mouse. Dashed lines

indicate the limit of detection ($NT_{50} = 60$). Two-tailed Wilcoxon rank-sum tests were used.

Extended Data Figure 10 | Strong back-boosting of Wuhan immunity in IGHV3-53 KI mice

restricts the breadth of neutralization against emerging variants.

a, Serum neutralization titres (NT_{50}) of IGHV3-53$^{+/+}$ and wild-type mice against a panel of SARS-

CoV-2 variant pseudoviruses following the fifth dose (XBB.1.5). Geometric mean titres (GMTs) are

shown on the top. Dashed lines indicate the limit of detection ($NT_{50} = 180$). **b,** Radar plot illustrating

the back-boosting effect of the JN.1 booster on neutralization titres against pre-JN.1 variants. **c,**

Neutralization profiles of IGHV3-53$^{+/+}$ and wild-type mice following XBB.1.5 and JN.1 boosters.

Fold changes and statistical significance between highlighted variants are indicated. Lines connect

data from the same mouse. Two-tailed Wilcoxon rank-sum tests were used in **a-c**.

Extended Data figure 1

Extended Data Figure 1 | Different epidemiological landscape of XFG and NB.1.8.1 in 2025.
a, Prevalence of XFG and NB.1.8.1 in China and the rest of the world as of October 5, 2025. The asterisk (*) denotes the specific lineage and its sublineages. Data were retrieved from CovSpectrum (https://cov-spectrum.org/).

Extended Data figure 2

Extended Data Figure 2 | Paired heavy and light chain V-gene usage for human memory B cells
a, Paired heavy and light chain V-gene usage for memory B cells from the mRNA-vaccinated and inactivated-only cohorts.

Extended Data figure 3

Extended Data Figure 3 | V(D)J-humanized mice exhibit a cellular imprinting phenotype dominated by cross-reactive GC and memory B cells.

a-c, Scatter plots showing the proportion of cross-reactive B cells (a), total GC B cells (b), and class-switched memory B cells (c). The indicated percentages reflect the frequency relative to their respective parental gates defined in Supplementary Information Fig. 1b. d, Linear regression analysis of the correlation between memory B cell cross-reactivity proportion and BA.5-specific IgG titer.

Extended Data figure 4

Extended Data Figure 4 | Repeated Omicron boosters fail to elicit robust Omicron-specific neutralization in imprinted humanized mice.

a, Line plots showing longitudinal pseudovirus neutralization titers (NT50) against a panel of SARS-CoV-2 variants in paired humanized and wild-type mice across Wuhan priming and BA.5 booster timepoints; lines connect data from the same mouse. Dashed lines indicate the limit of detection (NT50 = 60). Two-tailed Wilcoxon rank-sum tests were used.

Extended Data figure 5

Extended Data Figure 5 | Neutralization potency and cross-reactivity of mAbs from humanized and wild-type mice.
a,b, Comparison of antibody IC50 values against D614G and BA.5(a) and cross-reactivity proportion (b) from humanized and wild-type mice. Two-tailed Wilcoxon rank-sum tests were used in a.

Extended Data figure 6

Extended Data Figure 6 | Differential escape hotspots in humanized versus wild-type mice.
a–c, Scatter plots comparing the normalized average DMS escape scores of neutralizing antibodies isolated from humanized (y-axis) versus wild-type (x-axis) mice following Wuhan priming (a), the first BA.5 booster (b), and the second BA.5 booster (c). Residues falling along the diagonal (y=x) indicate shared immune pressure between the two models. To identify divergent hotspots—including those with low-to-moderate scores in one strain that are absent in the other—a shaded tolerance region was defined by the boundaries y = 1.4x + 0.3 and y = 0.6x - 0.18. Points falling outside this shaded region represent distinct escape hotspots. Among these outliers, the five residues with the highest escape scores in each group are highlighted in red. Related to Fig. 3.

Extended Data figure 7

Extended Data Figure 7 | Distinct SARS-CoV-2 antibody epitope distribution after sequential BA.5 boosts.
a-d, Epitope distribution of the antibody repertoire generated after one BA.5 boost (a, b) or two BA.5 boosts (c, d) in humanized mice (a, c) and wild-type mice (b, d). Related to Figure 4. e, Pyramidal bar charts showing the proportional distribution of epitope groups in antibodies isolated from humanized and wild-type mice after one or two BA.5 boosts. Bars are colored according to the log10 geometric mean IC50 of antibodies within each group.

Extended Data figure 8

Extended Data Figure 8 | Pharmacokinetics and dose-dependent suppressive effects of antibodies used in passive transfer.
a, Schematic of the in vivo monoclonal antibody pharmacokinetic study. b, Serum IgG titers over time following antibody injection (mIgG1 forms were detected using an anti-mouse Fc secondary antibody, and hIgG1 forms using an anti-human Fc secondary antibody). Dashed lines indicate the limit of detection (midpoint titre = 100). c, Schematic of the BD55-1205 dose-ranging experiment. d, Corresponding serum IgG titers before and after Wuhan RBD depletion. Statistical significance of the fold-reduction in titers was assessed between groups. Dashed lines indicate the limit of detection (midpoint titre = 100). Two-tailed Wilcoxon rank-sum tests were used in d.

Extended Data figure 9

Extended Data Figure 9 | IGHV3-53 knock-in mice exhibit diminished Omicron neutralization boosting compared to wild-type mice.

a, Line plots showing longitudinal pseudovirus neutralization titers (NT50) against a panel of SARS-CoV-2 variants in paired IGHV3-53+/+ (top), IGHV3-53+/- (middle), and wild-type (bottom) mice across BA.5 booster timepoints; lines connect data from the same mouse. Dashed lines indicate the limit of detection (NT50 = 60). Two-tailed Wilcoxon rank-sum tests were used.

Extended Data figure 10

Extended Data Figure 10 | Strong back-boosting of Wuhan immunity in IGHV3-53 KI mice restricts the breadth of neutralization against emerging variants.
a, Serum neutralization titers (NT50) of IGHV3-53+/+ and wild-type mice against a panel of SARS-CoV-2 variant pseudoviruses following the fifth dose (XBB.1.5). Geometric mean titres (GMTs) are shown on the top. Dashed lines indicate the limit of detection (NT50 = 180). b, Radar plot illustrating the back-boosting effect of the JN.1 booster on neutralization titers against pre-JN.1 variants. c, Neutralization profiles of IGHV3-53+/+ and wild-type mice following XBB.1.5 and JN.1 boosters. Fold changes and statistical significance between highlighted variants are indicated. Lines connect data from the same mouse. Two-tailed Wilcoxon rank-sum tests were used in a-c.