Transcriptional profiling of *Pseudomonas aeruginosa* biofilm life cycle stages reveals dispersal-specific biomarkers

Xavier Bertran i Forga[1,2], Kathryn E. Fairfull-Smith[3,4,] Jilong Qin[1,5*], and Makrina Totsika[1,2*]

[1]Centre for Immunity and Infection Control, School of Biomedical Sciences, Queensland University of Technology, Brisbane, QLD, Australia

[2]Max Planck Queensland Centre, Queensland University of Technology, Brisbane, QLD, Australia

[3]School of Chemistry and Physics, Queensland University of Technology, Brisbane, QLD, Australia

[4]Centre for Materials Science, Queensland University of Technology, Brisbane, QLD, Australia

[5]Institute for Molecular Bioscience, The University of Queensland, Brisbane, QLD, Australia

*correspondence: Jilong Qin (j.qin@uq.edu.au), and Makrina Totsika (makrina.totsika@qut.edu.au)

Abstract

Bacteria exhibit two lifestyles: planktonic free-floating individual cells or sessile multicellular

aggregates known as biofilms. The biofilm lifecycle is characterised by three distinct stages:

attachment, maturation and dispersal. Specific signals govern each stage triggering responses that

spatially and temporally regulate bacterial attachment to a surface, synthesis of extracellular matrix

components and their subsequent degradation. Characterising these signals is therefore a valuable

approach to develop novel antibiofilm therapies. Here, we used the model biofilm-forming bacterium

Pseudomonas aeruginosa PAO1 to characterise the transcriptional profiles of each stage of the

biofilm life cycle: attachment, biofilm maturation and spontaneous dispersal. We report that surface

attachment was accompanied by the upregulation of genes comprising the mechanosensor Pil-Chp,

whereas biofilm maturation characterised with the upregulation of genes involved in Pel

polysaccharide synthesis, *siaD* and PA4396 diguanylate cyclases as well as *pipA, fimX* and PA5442.

In contrast, dispersing cells upregulated genes responsible for the biosynthesis of alginate,

rhamnolipid, and extracellular nucleases (*eddA*, *eddB*), as well as the transcriptional regulator of

dispersal *amrZ*. Additionally, genes encoding the quorum sensing dispersal molecule cis-2-decenoic

acid (*dspS* and *dspI*), canonical phosphodiesterases (*nbdA* and *rbdA*) and nine other c-di-GMP–related

enzymes were also upregulated during dispersal. Our comprehensive analysis of transcriptional

profiles associated with different biofilm stages allowed us to define a subset of fourteen genes as

biomarkers whose transcription was largely upregulated during biofilm dispersal. Our study therefore

provides benchmarking stage-specific transcriptional profiles for *P. aeruginosa* biofilms in closed

culture systems*,* which led to the identification of a dispersal fingerprint marking the onset of

dispersal.

Importance

50 Biofilm infections by *P. aeruginosa* are a major medical challenge due to the increased tolerance to

51 antimicrobials displayed by bacteria living in sessile communities, which are forfeited during

52 spontaneous biofilm dispersal. Attachment, biofilm maturation and dispersal conform the main stages

of a dynamic process known as the biofilm lifecycle. However, the global responses governing the transitions through each stage remain understudied. Here, we combine live microscopy and biomass quantification to track the progression of *P. aeruginosa* cultures through the three main stages of the biofilm lifecycle. We show that cells from each stage recapitulated canonical stage-specific transcriptional responses and identify a set of dispersal-specific biomarkers. These biomarkers may offer a practical tool for rapidly screening dispersal-inducing compounds, aiding in the discovery of the next generation of antibiofilm therapeutics.

Introduction

Pseudomonas aeruginosa is a highly adaptable Gram-negative opportunistic pathogen, which survives in hostile environments by forming cell aggregates embedded in a protective extracellular matrix primarily composed of exopolysaccharides, extracellular DNA (eDNA) and structural proteins [1]. These aggregates, known as biofilms, exhibit increased tolerance to desiccation and host defences compared to their planktonic counterparts [2]. *P. aeruginosa* biofilms are associated with chronic infections, which exhibit high tolerance to a broad range of clinically employed antimicrobials [2].

In laboratories, *P. aeruginosa* biofilms are frequently studied using a variety of culture systems such as closed culture systems (i.e. microtiter plates) or open-flow systems (i.e. flow cells), in which three stages are exhibited: attachment, biofilm maturation and dispersal and each stage is characterised by distinct molecular mechanisms [3]. During attachment, cells transition from the motile to sessile lifestyle. This process is mediated by the mechanosensors Pil-Chp and Wsp which lead to the upregulation of genes involved in the synthesis of matrix components and fimbrial adhesins such as CupA [4–6]. During biofilm maturation, the components of the matrix are progressively secreted into the extracellular space. As biofilms grow and mature, environmental cues and intracellular signals may accumulate, such as the fatty acid signal cis-2-decenoic acid, eventually triggering spontaneous dispersal responses [7]. This phenomenon is characterised by biofilm-residing single cells regaining motility and resuming the planktonic lifestyle, during which they escape the biofilm matrix by hydrolysing its components [8]. This is achieved by the upregulation of glycosyl hydrolases such as

PslG and PelA for exopolysaccharide Psl and Pel degradation [8–10], and extracellular endonucleases

EndA, EddA and EddB [8–10]. This process is influenced by the intracellular concentration of the master

regulator cyclic-di-GMP (c-di-GMP) [6], which is controlled by a family of proteins harbouring

diguanylate cyclase domains (DGCs; possessing a GGDEF motif) and phosphodiesterase domains

(PDE; possessing a HD-GYP or EAL motifs), which respectively synthesise and degrade it [11–14].

The transition between the stages of the biofilm life cycle is driven by distinct signalling pathways. To

characterise them, transcriptomics has been applied to biofilms treated with biofilm-dispersing agents

such as NO donors or carbon sources, which primarily captured immediate transcriptional changes

triggered by these compounds leading to an abrupt interruption of the biofilm maturation process [15–17].

However, the progression through the biofilm stages is a dynamic process [18], urging a comprehensive

characterisation of transcriptional profiles between each stage. Therefore, we here performed RNA

sequencing to compare the currently understudied transcriptional variations across cells undergoing

attachment, biofilm maturation, and dispersal [33]. Subsequently, by tracking the transcription patterns

of genes throughout the three life cycle stages, we identified a subset of biomarkers for biofilm

dispersal.

Results

Microtiter plate-grown *P. aeruginosa* displays the three defined biofilm life stages

We previously showed that the biofilm life cycle of the model organism *Pseudomonas aeruginosa*

PAO1 in microtiter plates consisted of the following phases: 1) the progressive attachment of biomass

to the surface, reaching maximum biomass between 2-8 h in a process known as biofilm maturation

and, 2) a marked reduction in attached biomass known as the biofilm dispersal stage [18]. To further

characterise how these stages occur in closed culture systems such as microtiter plates, we quantified

the biofilm biomass (Fig 1A) and performed microscopy analysis simultaneously on forming and

dispersing biofilms at different time points to capture morphological changes during biofilm

formation (Fig 1B, Fig S1). Using this approach, we observed that the attachment stage occurs within

2 h of inoculation (Fig 1A and B; orange), during which cells adhered to the surface as a monolayer,

with a culture density (OD_{600}) of ~0.1 and a low level of biomass (stained by crystal violet with OD_{550}

of ~2.0). At 4-8 h post-inoculation, cells transitioned to the biofilm maturation stage (Fig 1A and B;

blue), organised in voluminous scattered clusters that grew over time and produced multicellular

biofilm structures (Fig 1B, Fig S1), as observed by the peak biofilm biomass staining quantified at the

8 h time point (Fig 1A). This build-up in biomass occurred between the early (OD_{600} ~0.3) and the

late-logarithmic phase (OD_{600} ~1.5). Beyond 8 h post-inoculation and at a higher culture density

(OD_{600}>1.5), biofilms biomass rapidly decreased. This phenomenon was characterised by

disassembled structures leaving clearly distinguishable biofilm remnants attached to the well surface

and a loss of 77% of accumulated biomass, (Fig 1A and B, Fig S1; green). Collectively, our data

reflects that cells grown in closed culture systems synchronically transition through distinctive phases

culminating in spontaneous biofilm dispersal. Therefore, these systems can effectively recapitulate the

stages of the biofilm life cycle originally described in open-flow culture systems (flow cells) [3,19,20].

Figure 1. Biofilm formation and dispersal kinetics of *P. aeruginosa* PAO1. (A) *P. aeruginosa* PAO1 biofilm biomass and

culture turbidity over 12 h. PAO1 cultures were seeded on 24-well plates and biofilm biomass was quantified at different

time points by crystal violet (CV) staining (●). Culture density was simultaneously monitored by OD_{600} (○) with means ± SD

shown. Data are representatives for at least 3 biological replicates for crystal violet staining and 1 biological replicate for

OD_{600}. **(B)** Photographic and microscopic images of the crystal violet-stained biofilms at each stage of the biofilm life cycle.

Crystal-violet-stained microscopic images of *P. aeruginosa* biofilm cultures in 24-wells. Images are representative of at least

3 biological replicates. Orange: attachment stage. Blue: biofilm maturation stage. Green: dispersal stage. This colour coding

scheme is used throughout the article to differentiate the three biofilm stages.

Transcriptional profiles recapitulate canonical pathways of attachment, maturation and dispersal

Having observed that *P. aeruginosa* cells grown in microtiter plates synchronically undergo the three

defined stages of the biofilm life cycle, we performed RNA sequencing of *P. aeruginosa* during

attachment, biofilm maturation and dispersal to characterise stage-specific transcriptional profiles (Fig

2A). Differential transcription analysis using a negative binomial test (P-value $\leq$ 0.01; Log_2 fold-

change $\geq$ |1|) identified a total of 3361 genes with significant transcriptional changes across all three

comparisons: i) Attachment vs. Biofilm Maturation; ii) Biofilm Maturation vs. Dispersal; and iii)

Attachment vs. Dispersal) (Table S1). Principal component analysis (PCA) further confirmed that

cells at each stage formed distinct clusters, indicating discrete transcriptional profiles were associated

with attachment, maturation, and dispersal (Fig 2B). Specifically, in cells undergoing biofilm

maturation 540 genes were upregulated, and 405 genes were downregulated compared to attaching

cells, whereas 1,240 genes were upregulated and 1,207 genes downregulated in maturing biofilms

relative to dispersing bacterial cells (Fig 2C).

Figure 2. Transcriptional profiles of different stages of the biofilm life cycle. (A) Schematic depicting RNA sample

collection for sequencing. During attachment, the free-swimming phase was collected to represent cells rapidly transitioning

from motile to sessile. During biofilm maturation, surface-attached cells were collected, and cells from both phases were

collected during the dispersal stage. **(B)** Principal component analysis of samples collected at distinct stages of the biofilm

life cycle (attachment – orange; biofilm maturation – blue; the planktonic phase of biofilm dispersal – light green; and the

biofilm phase of biofilm dispersal – dark green). Each biological replicate of a group is represented as a dot of the same

colour. The normalised reads of all genes were then extracted for each sample and transformed into PC loadings with

ClustVis. Values for PC1 and PC2, accounting for the largest variance percentage are represented in the plot. **(C)** Venn

diagrams showing differentially transcribed genes in the planktonic phases of attachment and dispersal (relative to biofilm

maturation). Downregulated (top) and upregulated (bottom) genes at each stage relative to biofilm maturation are

represented separately. Overlapping areas indicate genes with consistent transcriptional changes relative to biofilm

maturation stage.

Building on these global transcriptional variations and to further validate our datasets, we mined our

RNA-seq data for previously reported stage-specific changes in the transcription of genes previously

associated with attachment, biofilm formation or dispersal. During attachment and extending into

biofilm maturation, we observed the upregulation of the Pil-Chp surface-sensing complex known to

mediate attachment [6], whereas no upregulation of the complementary Wsp system was noted. Notably,

transcription of *pilA, pilB, pilD* and *pilGHIJK-chpABCDE,* encoding type IV pili structural and signal

transduction proteins, and of *cyaB* and *vfr,* which encode an adenylate cyclase and its associated

cyclic-AMP-controlled transcriptional regulator Vfr, was upregulated between 2.00- and 7.37-fold

relative to biofilm dispersal.

Biofilm maturation involves synthesis pathways for extracellular polysaccharides and eDNA, which

are the main components of the biofilm matrix [1]. During biofilm maturation, we observed the

upregulation of the exopolysaccharide Pel synthesis operon *pelBCDEF* but not *pelG* relative to

attachment (2.22 – 2.26-fold) and dispersal stages (2.16 – 2.99-fold) (Fig 3A, Table S1) [21].

Additionally, the alternative respiration process of denitrification is known to be active during biofilm

maturation due to cell layering inside the biofilm restricting oxygen availability [22]. Accordingly, we

found biofilm cells to upregulate the transcription of the nitrite reductase complex (*nirSMCFDLG*) by

up to 10-fold relative to attachment.

Planktonic cells at the biofilm dispersal stage, relative to attachment and biofilm maturation stages,

showed an increased transcription of genes involved in eDNA hydrolysis (*eddA,* 3.28-fold; and *eddB,*

2.95-fold; but not *endA*) (Fig 3A, Table S1), suggesting the remodelling of the extracellular matrix to

allow the escape of planktonic cells. We also observed a pronounced upregulation of quorum-sensing–

regulated genes previously shown to drive biofilm dispersal. The *rhlA*, *rhlB*, and *rhlC* genes, which

encode enzymes for the biosynthesis of the rhamnolipid biosurfactant implicated in quorum-sensing–

mediated dispersal [39–42]were upregulated by 88-, 45- and 12-fold, respectively (Fig 3A, Table S1) [23–26].

Another quorum-sensing pathway activated during dispersal involved the fatty acid signal cis-2-

decenoic acid [25,27]. This fatty acid signal is produced and sensed by the proteins encoded by *dspI* and

dspS, which were upregulated by 2.04- and 5.59-fold, respectively, during dispersal relative to the

biofilm maturation stage (Fig 3A, Table S1). Interestingly, genes associated to biofilm maturation

were also found upregulated during the dispersal stage. Adhesion factors, such as lectins *lecA* and

lecB were highly upregulated during dispersal by 279-fold and 115-fold, respectively, relative to the

biofilm maturation stage. Similarly, *flp*, which encodes the main subunit of the type IVb pili adhesin

was upregulated by 120-fold, together with several alginate biosynthesis genes (*alg8*, 2.26-fold*; algX,*

2.16-fold*; algJ*, 2.39-fold; *algA,* 2.74-fold) and their associated regulators *algU* (2.49-fold)*, algR*

(5.76-fold) and *amrZ* (6.27-fold) (Fig 3A, Table S1)*.* These findings were consistent with previous

studies showing that dispersed cells had significantly increased functional adhesion relative to

planktonic cells [28]. Therefore, our RNA-seq data captured transcriptional changes associated with

previously documented attachment, biofilm maturation and dispersal responses, further underscoring

the consistency of our dataset with established biofilm regulatory pathways and the robustness of our

analytical methods.

To further validate the upregulation of these known dispersal pathways, we extracted RNA from

biofilms at 4 h (early maturation), 8 h (late maturation), and planktonic cells collected at 12 h

(dispersal) and measured the transcript levels of representative genes involved in rhamnolipid

synthesis (*rhlAB, rhlC*), eDNA degradation (*eddA, eddB*), and cis-2-decenoic acid signalling (*dspI, dspS*) by RT-qPCR (Fig 3B). Relative to maturing biofilms, all genes were upregulated in mature and dispersed biofilms. These findings further validate our RNA-seq data and indicate that mature biofilms may begin to transcriptionally activate dispersal pathways before the onset of active detachment.

Figure 3. Genes involved in dispersal mechanisms are upregulated during late biofilm maturation and dispersal

compared to early biofilm maturation. (**A**) Comparison of the transcriptomes of planktonic cells during attachment or

dispersal to that of sessile cells during biofilm maturation. Genes containing Log2-fold change transcription values >|1| were

highlighted in biofilm stage specific colours. (**B and C**) RT-qPCR analysis of canonical dispersal genes (*rhlA, rhlB, rhlC,*

eddA, eddB, dpsS and *dspI*) and the PDEs PA4108 and PA4781, which were identified as putatively involved in dispersal.

RNA samples were extracted at different time points with Means ± SEM shown.

Specific c-di-GMP enzymes characterise the biofilm maturation and dispersal transcriptomes

Cyclic di-GMP (c-di-GMP) is a key signalling molecule governing the transition from biofilm

maturation to dispersal. Considering that the roles of enzymes responsible for its synthesis and

breakdown are stage-specific, we also analysed our transcriptomic data to identify genes encoding

enzymes with predicted PDE (EAL or HD-GYP) or DGC (GGDEF) domains (Table 1). Expectedly,

genes encoding proteins essential for biofilm development (*siaD,* 3.73-fold and *pipA,* 2.08-fold) were

upregulated during biofilm maturation [29–32]. Genes implicated in the formation of large microcolonies

[30], including PA4396 (2.08-fold), *fimX* (2.77-fold), and PA5442 (2.36-fold) were also significantly

upregulated. In contrast, we observed an upregulation of a combination of PDEs and DGCs in cells at

the dispersal stage, including sensory transducers of dispersal cues *nbdA* (7.56-fold) and *nicD* (6.73-

fold) [33–35]. Additionally, upregulation of *ismP* (12.99-fold) was noted, which encodes an iron sensor

that under low iron conditions hijacks and inhibits a cognate DGC ImcA [36].

Table 1: Differentially regulated genes encoding c-di-GMP-modifying enzymes during biofilm

maturation and dispersal.

Locus Tag	Gene name	Domain(s)	Activity	Fold-change
Biofilm maturation				(vs Dispersal)
PA0169	*siaD*	GGDEF	DGC	3.73
PA0285	*pipA*	GGDEF-EAL	PDE	2.08
PA4396	-	GGDEF	-	2.08

PA4959	*fimX*	GGDEF-EAL	PDE	2.77
PA5442	-	GGDEF-EAL	PDE	2.36
Biofilm dispersal				(vs Maturation)
PA0290	-	GGDEF	-	2.73
PA0575	*rmcA*	GGDEF-EAL	PDE	3.51
PA0847	-	GGDEF	DGC	2.09
PA0861	*rbdA*	GGDEF-EAL	PDE	3.03
PA1878	-	HD-GYP	-	3.39
PA2072	*ismP*	GGDEF-EAL	-	12.99
PA2572	-	HD-GYP	-	8.22
PA2771	-	GGDEF	DGC	2.75
PA3311	*nbdA*	GGDEF-EAL	PDE	7.56
PA3825	-	EAL	PDE	1.25
PA4108	-	HD-GYP	PDE	5.60
PA4781	-	HD-GYP	PDE	10.70
PA4929	*nicD*	GGDEF	DGC	6.73

Interestingly, dispersed cells upregulated all four genes encoding proteins with HD-GYP

phosphodiesterase domains present in the *P. aeruginosa* PAO1 genome (PA1878, PA2572, PA4108

and PA4781) (Table 1). Two of these, PA4108 and PA4781 were previously reported to have PDE

activity and to be necessary for swarming and the production of virulence factors [37,38]. We further

validated the transcriptional changes of these two genes using RT-qPCR (Figure 3C), confirming an

increase in expression of 7.5-fold and 4-fold, respectively, during dispersal relative to maturing

biofilms at 4 h. Moreover, the transcription of both genes was also upregulated in mature biofilms

(6.2- and 4-fold; Figure 3C), indicating a similar pattern of transcriptional priming prior to dispersal.

Altogether, our results indicate that canonical stage-specific c-di-GMP modulation responses are

accurately reflected in our transcriptomic dataset and suggest a potential undescribed role of HD-

GYP-containing PDEs in mediating biofilm dispersal.

Transcriptional biomarkers identify the onset of biofilm dispersal

Since the obtained transcriptomic profiles recapitulated canonical pathways associated with

attachment, maturation, and dispersal, our datasets are uniquely well suited to identify genes

associated with the dispersal phase of *P. aeruginosa* to serve as biomarkers of biofilm dispersal. We

reasoned that genes directly involved in biofilm dispersal would exhibit inverse regulation between

the biofilm maturation and dispersal stages relative to the initial attachment stage. Thus, pro-dispersal

genes would be repressed during maturation but strongly upregulated during dispersal, or vice versa.

To identify candidate biomarkers, we applied these selection criteria and identified 14 upregulated and

8 downregulated genes that could potentially serve as dispersal biomarkers (Table 2).

Among the upregulated genes, *cheR2* (2.1-fold, Table 2) encodes a methyltransferase component of

the Che2 chemotaxis pathway, which interacts with the heme-containing oxygen sensor McbP/Aer2

and the coupling protein CheW2 to modulate virulence through a cryptic phosphorylation cascade

[39,40]. Another notable gene, *pqqA* (2.55-fold; Table 2), is part of the *pqqABCDE* operon responsible for

the biosynthesis of pyrroloquinoline quinone (PQQ), a redox cofactor for the quinoprotein ethanol

dehydrogenases ExaA and ExaBC [41]. In addition, several genes under the transcriptional control of the

two-component system PprAB were also upregulated. These include *tadA*, *rcpA*, *rcpC*, and *flp* (13,66-

, 12.47-, 13.83- and 34.54-fold; Table 2) which are part of a contiguous cluster spanning PA4297

(*tadG*) to PA4306 (*flp*). This locus encodes the machinery for the synthesis and secretion of Flp type

IVb pili [42]. Notably, two of the dispersal biomarkers *amrZ* and PA2588 (*cdpR*) have known roles in

modulating biofilm physiology. *amrZ* encodes a recognised transcriptional regulator of dispersal and

cdpR is a quorum sensing and virulence transcriptional regulator [43–45].

Several uncharacterised genes, including PA0111, PA0743 and PA1353 were also identified as

upregulated biomarkers of biofilm dispersal, many of which appear to be associated with redox

processes. While no experimental data are available for PA0111, bioinformatic predictions suggest

that the directly upstream gene PA0110 encodes a protein putatively related to a mitochondrial

cytochrome *c* oxidase. Additionally, PA0112 and PA0113 are predicted to encode a heme A synthase

and a protoheme IX farnesyltransferase, respectively, implicating these loci in cytochrome maturation

or electron transport. Similarly, PA0743, which encodes an L-serine dehydrogenase, was also

upregulated, potentially reflecting altered amino acid catabolism related to redox homeostasis during

biofilm dispersal. No functional predictions are currently available for PA1353, which was also part of

the identified dispersal biomarkers.

Table 2. Genes differentially regulated at the biofilm maturation and dispersal stage compared

to the attachment stage.

Gene	Fold change in Biofilm maturation [a]	Fold change in Biofilm dispersal [a]	Gene product
Downregulated at 4h and upregulated at 8h			
PA0111	-2.01	12.21	hypothetical protein
cheR2	-2.22	2.10	chemotaxis protein methyltransferase
PA0743	-2.58	3.12	NAD-dependent L-serine dehydrogenase
PA1353	-2.36	3.32	hypothetical protein
pqqA	-5.10	2.55	coenzyme PQQ synthesis protein A
PA2588	-2.03	3.01	transcriptional regulator
amrZ	-2.30	2.55	alginate and motility regulator Z
tadA	-2.17	13.64	ATPase TadA
rcpA	-2.13	12.47	type II/III secretion system protein
rcpC	-2.07	13.83	hypothetical protein
flp	-3.27	34.54	type IVb pilin Flp
PA4523	-2.31	3.41	hypothetical protein
cupE1	-2.11	9.58	fimbrial subunit CupE1
cupE2	-2.04	3.95	fimbrial subunit CupE2
Upregulated at 4h and downregulated at 8h			
PA0485	2.93	- 2.91	hypothetical protein
PA1170	2.43	-4.76	hypothetical protein
metH	2.45	-2.13	B12-dependent methionine synthase
ilvH	3.53	-2.35	acetolactate synthase small subunit
ilvI	4.69	-2.57	acetolactate synthase 3 catalytic subunit
cysN	2.1	-2.62	bifunctional sulfate adenylyltransferase subunit1/adenylylsulfate kinase
PA4889	2.06	-14.92	oxidoreductase

| PA4907 | 2.41 | -2.27 | short-chain dehydrogenase |

We restricted our biomarker screen to upregulated transcripts, as elevated expression levels are more consistently detected and validated across analytic platforms. Due to the similarities in transcriptomic profiles between planktonic and biofilm cells collected during dispersal, we hypothesised that biofilms become primed for dispersal prior to phenotypically manifesting the response. Therefore, we collected samples from mature biofilms at 8 h and from dispersed cells at 12 h and validated these biomarker genes by RT-qPCR (Fig 4). Our results confirmed the upregulation of all fourteen genes during dispersal, with four of these (PA0743, PA1353, *pqqA*, and PA2588) only being significantly upregulated at 12 h (Fig 4). Genes PA0111, *cheR2*, *tadA*, *rcpC*, *flp*, and PA4523 showed a pattern where transcript abundance increased in mature biofilms (2.5- to 23-fold) and peaked during dispersal (4.6- to 117-fold relative to early biofilms) (Fig 4). In contrast, the transcription of *cupE1* and *cupE2* fimbrial genes peaked in mature biofilms (28.6- and 10.1-fold) but sharply declined upon dispersal (6.1- and 5.9-fold) (Fig 4). This transient transcriptional profile suggests that CupE fimbriae genes are likely characteristic of mature biofilms. In contrast, transcription of PA0111, *cheR2, tadA, rcpC, flp* and PA4523 peaked at dispersal, indicating that matured biofilms become primed for dispersal prior to its onset (Fig 4). Altogether, these biofilm dispersal markers exhibit strong reproducibility, with 12 out of 14 becoming progressively upregulated during the transition from mature biofilms to dispersal.

Figure 4. Validation of the transcriptional biomarkers of biofilm dispersal. Differences in transcriptional regulation measured by RT-qPCR for the indicated genes from total RNA samples extracted at 4 h (early biofilm maturation), 8 h (late

298   biofilm maturation) and 12 h (dispersal). Fold-changes at 8 h and 12 h are relative to 4 h. Means ± SEM are shown. Two

299   technical replicates are represented.

Discussion

The study of transcriptional changes between each biofilm stage provides valuable insights into the

key cellular signalling pathways underlying their transition between sessile and motile lifestyles.

Here, we characterised the transcriptional profiles of cells undergoing attachment, biofilm maturation

and dispersal, and identified 23 biomarkers characterising the dispersal response, 14 of which

comprised a subset strongly upregulated upon the initiation of dispersal.

Cells undergoing attachment sense and respond to surface contacts through mechanosensory systems

such as the Pil-Chp machinery, which relies on type IV pili-mediated twitching motility [46,47]. In our

study, we report an upregulation of type IV pili biogenesis genes (*pilA, pilB, pilD*) and of the

mechanosensory system Pil-Chp (*pilGHIJK-chpABCDE*) during the attachment and biofilm

maturation stages compared to dispersing cells. These were accompanied by the upregulation of *fimX*,

which encodes a putative phosphodiesterase essential for the assembly of type IV pili, and of *cupA1*

encoding the main subunit of the CupA adhesin [48]. Our findings therefore align with previous studies

indicating that type IV pili are indispensable to surface-attachment in microplates [49], and are

consistent with prior reports that surface sensing promotes the transcriptional upregulation of genes

involved in type IV pilus biogenesis [50].

The intracellular messenger c-di-GMP acts as a master regulator of the bacterial lifestyle, with its

accumulation leading to biofilm formation [8]. Our dataset revealed the upregulation of several stage-

specific genes encoding proteins with validated or hypothetical c-di-GMP modulating activity.

Specifically, *siaD* was the most upregulated during biofilm maturation, which encodes a DGC

involved in early microcolony formation and adhesion [46,51]. In contrast, biofilm dispersal is associated

with increased PDE activity and decreasing intracellular c-di-GMP [52]. This was reflected in our

dataset, where 6 genes encoding functionally active PDEs were upregulated. Genes *rbdA* and *nbdA*

encode enzymes with well-described roles in biofilm dispersal. Both have been reported to modulate

flagellar motility, with RbdA being a key component of the BdlA-regulated core dispersal response

[53,54]. Moreover, four upregulated genes (PA1878, PA2572, PA4108 and PA4781) encode PDEs

carrying the HD-GYP motif [55]. While the role of these genes in the biofilm dispersal stage is unclear,

it has been reported that disruption of PA4108 and PA4781 resulted in higher intracellular c-di-GMP

and decreased swarming activity, whereas PA2572 modulated swarming motility [55,56]. We here

validated by RNA-seq and RT-qPCR that both PA4108 and PA4781 are strongly upregulated during

late biofilm maturation and dispersal, which suggests a potential role in initiating dispersal responses.

Additionally, recent reports showed that deletion of *cdpR* led to increased biofilm biomass and

decreased swarming motility [45]. Considering that the upregulation of *cdpR* is strongly controlled by

the quorum sensing regulator RsaL [44], the stage-specific transcriptional activation of *cdpR* would

therefore suggest that dispersal in closed systems is mediated by a global quorum sensing-mediated

swarming response.

The synthesis of the rhamnolipid biosurfactant is important to the quorum sensing-mediated swarming

motility and dispersal [24]. In closed culture systems, rhamnolipid was proposed to drive biofilm

dispersal [57]. Here we report that the rhamnolipid and swarming modulator PA2572 together with

several rhamnolipid-associated genes were highly upregulated during dispersal. Moreover, four of the

identified PDEs upregulated during dispersal, including PA4108, PA4781, *nbdA* and *rbdA,* were

previously shown to influence the synthesis of rhamnolipid leading to loss of biofilm biomass [19,34,56].

Altogether, these data suggest biofilm dispersal in closed systems may be driven by rhamnolipids.

Among the upregulated biomarkers, *amrZ* was reported as a global regulator controlling the

upregulation of genes related to alginate biosynthesis and Type IV pili twitching motility while

downregulating Pel synthesis, therefore driving biofilm dispersal [43,58]. In contrast, *tadA*, *rcpA*, *rcpC*,

flp, cupE1 and *cupE2* were previously reported to be responsible for cell adhesion which are regulated

under the PprAB two-component system [59,60]. Our data therefore support previous studies indicating

that dispersing cells exhibit a hyperadhesive potential that primes them to re-attach to surfaces [28].

Additionally, we identified PA0110, *cheR2*, *pqqA,* PA0743, PA1353 and PA4523 to be uniquely

associated to the dispersal stage, yet their specific roles remain to be defined. Their marked

upregulation during dispersal warrants further investigation.

Understanding biofilm dispersal could prove a critical step towards combating chronic, biofilm-

associated infections. As dispersing cells become resensitised to antimicrobial treatment once they are

released from the protective biofilm environment, the underlying transcriptional changes occurring

during biofilm dispersal hold promise to unveiling the next generation of antibiofilm treatments and

adjuvants. We here report the transcriptional profiles of the *P. aeruginosa* biofilm life cycle, including

attachment, biofilm maturation and dispersal stages in closed systems, which can be used as

benchmarks for future studies. Additionally, we propose a robust set of biofilm dispersal biomarkers

with the potential to be used in the high throughput screens supporting the discovery and development

of novel dispersal-inducing antibiofilm therapeutics.

Materials and Methods

Strains, media and culture conditions

364 *Pseudomonas aeruginosa* PAO1 wild-type (WT). Cultures were routinely grown overnight in LB

365 (lysogeny broth) media at 37°C, 200rpm before incubation in fresh M9 media (9 mM NaCl, 22 mM

366 KH_2PO_4, 48 mM Na_2HPO_4, 19 mM NH_4Cl and 2 mM $MgSO_4$, 100 µM $CaCl_2$, 0.4% glucose, pH 7.0)

367 for each experiment. Where appropriate, cultures were supplemented with 10µg/ml of gentamicin.

Biofilm kinetics and microscopy

Biofilms were grown on Nunc 24-well plates (ThermoFisher Scientific) or tissue culture flasks (Nunc,

ThermoFisher Scientific) by inoculating 10^7 colony forming units (CFU)/ml of overnight bacterial

cultures for 2 h, 4 h, 8 h and 12h in 1 ml of M9 media at 37 $^{\circ}$C and 180 rpm (Fig S2). At the indicated

time points, supernatants were removed, and wells were delicately washed with 1ml phosphate

buffered saline (PBS, Gibco) to remove planktonic cells. Microscopy (400x; PrimoVert, Zeiss) was

used to confirm the removal of planktonic cells and preservation of biofilm structures. Remaining

biofilm cells were stained with 0.1% (w/v) crystal violet dissolved in 6.25% (v/v) methanol. Stained

biofilms were washed twice with PBS and solubilised with ethanol. Biofilm biomass was quantified

by measuring the optical density at 550 nm (OD_{550}) using a SPECTROStar Nano microplate reader

(BMG LabTech). Micrographs of stained biofilms were taken via microscopy as described above.

RNA sequencing and analysis

Biofilms were cultured and dispersed as previously described with some adjustments [18]. Briefly, tissue

culture flasks (Nunc, ThermoFisher Scientific) were inoculated with 10^7 CFU/ml cells in 50ml of M9

medium from overnight cultures. Cultures were incubated at 37^oC shaking at 70rpm for 2, 4 or 8h. At

2 h and 8 h, planktonic cells (10^8) were collected from the liquid phase and immediately mixed with 2

volumes of RNA-protect solution (QIAGEN). Biofilm cells were gently rinsed with 10 ml of PBS and

resuspended in 6 ml of RNAprotect mixed with 3ml of PBS using cell scrapers. Cells (10^8) were

subsequently pelleted at 5000 g (10 min, 25 ^{o}C). RNA was extracted with the RNeasy mini kit

(QIAGEN, Cat# 74104) as per the manufacturer's protocol for Gram negative bacteria grown in

minimal media. Extracted RNA samples were treated with DNAse I and rRNA was removed using

streptavidin-coated magnetic beads (BGI Genomics, Shenzhen, China). Library was prepared by RNA

fragmentation and cDNA synthesis followed by the ligation of adaptors at the 3' end of cDNA

fragments. These were further amplified by PCR and single-stranded fragments were circularised for

downstream DNBSEQ PE100 sequencing (BGI Genomics, Shenzhen, China). Clean reads were

mapped to the *P. aeruginosa* PAO1 reference genome (GenBank Accession number AE004091.2)

with Bowtie2 and comparison of expression levels between sample groups (attachment, biofilm

maturation and dispersal) was done with DESeq2 in Geneious Prime v2024.0.7.

The PCA plot was generated by extracting the reads per kilobase of transcript per million mapped

reads (RPKM) of every gene at each one of the stages (Table S1). Then, genes with <|2-fold|

transcription or *P*-value >0.01 for all groups were discarded, and the remaining differentially

regulated genes were plotted using GraphPad Prism 10.4.1 (Graph Pad, USA).

qRT-PCR

To validate RNA-seq data, total RNA was extracted from cells harvested from different biofilm stages

as described above. cDNA was transcribed using 1 µg of total RNA using the SuperScript III first-

strand synthesis kit (Invitrogen) as per the manufacturer's protocol. cDNA samples were then used as

templates for RT-qPCR conducted with the Quantinova SYBR Green master mix (Qiagen) as per the

manufacturer's protocol. The housekeeping gene *recA* was used as a control. Each sample was

independently assayed twice with the gene-specific primers listed in Table S2. Relative gene

expression was calculated using the $2^{-\Delta\Delta CT}$ method [61]. Transcriptional fold changes were plotted using

GraphPad Prism.

Acknowledgements

This work was funded by an Australian Research Council project grant (DP210101317), the Max

Planck Queensland Centre on the Materials Science of Extracellular Matrices, and a QUT Amplify

Scholarship provided by the Queensland University of Technology (Australia) to XB. The Ian Potter

Foundation sponsored the CLARIOStar high-performance microplate reader (BMG, Australia). The

funders had no role in study design, data collection and analysis, decision to publish, or preparation of

the manuscript. The authors would like to thank Professor Robert EW Hancock (University of

Columbia) for providing the *P. aeruginosa* PAO1 strain used in this study.

Competing interests

418 MT is an employee of the GSK group of companies. All other authors declare no competing interests.

419 This research was conducted in the absence of any commercial or financial relationships that could be

420 constructed as a potential conflict of interest.

Author contributions

422 JQ and MT conceptualised the project. XB, JQ and MT contributed to experimental design. XB

423 conducted all experiments, and contributed to data collection, analysis and visualisation. XB, JQ and

424 MT contributed to data interpretation. JQ, MT and KFS supervised the project. KFS and MT obtained

425   the funding. XB wrote the manuscript draft, JQ substantially revised the manuscript, all authors edited

426   the manuscript.

Data availability

428   The RNA-Seq data have been deposited in the NCBI Short Read Archive (SRA) database with

429   accession code PRJNA1291829.

References

431   1.   Flemming, H. C. & Wingender, J. The biofilm matrix. *Nat Rev Microbiol* **8**, 623–633
432   (2010).

433   2.   Høiby, N., Johansen, H. K., Moser, C. & Ciofu, O. Clinical Relevance of Pseudomonas
434   aeruginosa: A Master of Adaptation and Survival Strategies. in *Pseudomonas: Model
435   Organism, Pathogen, Cell Factory* (ed. Rehm, B.) 25–44 (John Wiley & Sons, Ltd,
436   Weinheim, 2008). doi:10.1002/9783527622009.CH2.

437   3.   Sauer, K., Camper, A. K., Ehrlich, G. D., Costerton, J. W. & Davies, D. G.
438   Pseudomonas aeruginosa displays multiple phenotypes during development as a
439   biofilm. *J Bacteriol* **184**, 1140–1154 (2002).

440   4.   Güvener, Z. T. & Harwood, C. S. Subcellular location characteristics of the
441   Pseudomonas aeruginosa GGDEF protein, WspR, indicate that it produces cyclic-di-
442   GMP in response to growth on surfaces. *Mol Microbiol* **66**, 1459 (2007).

443   5.   Vallet, I., Olson, J. W., Lory, S., Lazdunski, A. E. & Filloux, A. The chaperone/usher
444   pathways of Pseudomonas aeruginosa: Identification of fimbrial gene clusters (cup)
445   and their involvement in biofilm formation. *PNAS* **98**, 6911–6916 (2001).

446   6.   Patino, R. *et al.* Antagonistic response regulators spatially regulate receptor
447   methylation in the Pseudomonas aeruginosa Pil-Chp surface sensing system. *Cell
448   Rep* **44**, 115536 (2025).

449   7.   McDougald, D., Rice, S. A., Barraud, N., Steinberg, P. D. & Kjelleberg, S. Should we
450   stay or should we go: Mechanisms and ecological consequences for biofilm dispersal.
451   *Nature Reviews Microbiology* vol. 10 Preprint at https://doi.org/10.1038/nrmicro2695
452   (2012).

453   8.   Rumbaugh, K. P. & Sauer, K. Biofilm dispersion. *Nat Rev Microbiol* **18**, 571–586
454   (2020).

455   9.   Cherny, K. E. & Sauer, K. Pseudomonas aeruginosa Requires the DNA-Specific
456   Endonuclease EndA To Degrade Extracellular Genomic DNA To Disperse from the
457   Biofilm. *J Bacteriol* **201**, e00059-19 (2019).

458   10.   Cherny, K. E. & Sauer, K. Untethering and degradation of the polysaccharide matrix
459   are essential steps in the dispersion response of Pseudomonas aeruginosa biofilms. *J
460   Bacteriol* **202**, (2020).

11.   Randall, T. E. *et al.* Sensory Perception in Bacterial Cyclic Diguanylate Signal
Transduction. *J Bacteriol* **204**, (2022).

12.   Hengge, R. Trigger phosphodiesterases as a novel class of c-di-GMP effector
proteins. *Philosophical Transactions of the Royal Society B: Biological Sciences* vol.
371 Preprint at https://doi.org/10.1098/rstb.2015.0498 (2016).

13.   Hengge, R. High-specificity local and global c-di-GMP signaling. *Trends in*
Microbiology vol. 29 993–1003 Preprint at https://doi.org/10.1016/j.tim.2021.02.003
(2021).

14.   Povolotsky, T. L. & Hengge, R. 'Life-style' control networks in Escherichia coli:
Signaling by the second messenger c-di-GMP. *J Biotechnol* **160**, 10–16 (2012).

15.   Kalia, M. & Sauer, K. Distinct transcriptome and traits of freshly dispersed
Pseudomonas aeruginosa cells. *mSphere* **9**, (2024).

16.   Chua, S. L. *et al.* Dispersed cells represent a distinct stage in the transition from
bacterial biofilm to planktonic lifestyles. *Nature Communications 2014 5:1* **5**, 1–12
(2014).

17.   Zhu, X., Rice, S. A. & Barraud, N. Nitric Oxide and Iron Signaling Cues Have
Opposing Effects on Biofilm Development in Pseudomonas aeruginosa. *Appl Environ*
Microbiol **85**, (2019).

18.   Bertran i Forga, X., Hong, Y., Fairfull-Smith, K. E., Qin, J. & Totsika, M. Nitric oxide
donor sodium nitroprusside serves as a source of iron supporting *Pseudomonas*
aeruginosa growth and biofilm formation. *Microbiol Spectr*
https://doi.org/10.1128/SPECTRUM.02234-25 (2025)
doi:10.1128/SPECTRUM.02234-25.

19.   An, S., Wu, J. & Zhang, L. H. Modulation of pseudomonas aeruginosa biofilm
dispersal by a cyclic-di-gmp phosphodiesterase with a putative hypoxia-sensing
domain. *Appl Environ Microbiol* **76**, 8160–8173 (2010).

20.   Hsu, J. L., Chen, H. C., Peng, H. L. & Chang, H. Y. Characterization of the Histidine-
containing Phosphotransfer Protein B-mediated Multistep Phosphorelay System in
Pseudomonas aeruginosa PAO1. *Journal of Biological Chemistry* **283**, 9933–9944
(2008).

21.   Jennings, L. K. *et al.* Pel is a cationic exopolysaccharide that cross-links extracellular
DNA in the Pseudomonas aeruginosa biofilm matrix. *Proc Natl Acad Sci U S A* **112**,
11353–11358 (2015).

22.   Barraud, N. *et al.* Involvement of Nitric Oxide in Biofilm Dispersal of Pseudomonas
aeruginosa. *J Bacteriol* **188**, 7344–7353 (2006).

23.   Bhattacharjee, A., Nusca, T. D. & Hochbaum, A. I. Rhamnolipids mediate an
interspecies biofilm dispersal signaling pathway. *ACS Chem Biol* **11**, 3068–3076
(2016).

24.   Schooling, S. R., Charaf, U. K., Allison, D. G. & Gilbert, P. A role for rhamnolipid in
biofilm dispersion. *Biofilms* **1**, 91–99 (2004).

25. Davey, M. E., Caiazza, N. C. & O'Toole, G. A. Rhamnolipid surfactant production
affects biofilm architecture in Pseudomonas aeruginosa PAO1. *J Bacteriol* **185**, 1027–
1036 (2003).

26. Glick, R. *et al.* Increase in rhamnolipid synthesis under iron-limiting conditions
influences surface motility and biofilm formation in Pseudomonas aeruginosa. *J*
Bacteriol **192**, 2973–2980 (2010).

27. Kalia, M., Amari, D., Davies, D. G. & Sauer, K. cis-DA-dependent dispersion by
Pseudomonas aeruginosa biofilm and identification of cis-DA-sensory protein DspS.
mBio **14**, (2023).

28. Chambers, J. R., Cherny, K. E. & Sauer, K. Susceptibility of Pseudomonas aeruginosa
Dispersed Cells to Antimicrobial Agents Is Dependent on the Dispersion Cue and
Class of the Antimicrobial Agent Used. *Antimicrob Agents Chemother* **61**, e00846-17
(2017).

29. Eilers, K. *et al.* The dual GGDEF/EAL domain enzyme PA0285 is a Pseudomonas
species housekeeping phosphodiesterase regulating early attachment and biofilm
architecture. *Journal of Biological Chemistry* **300**, (2024).

30. Eilers, K. *et al.* Phenotypic and integrated analysis of a comprehensive Pseudomonas
aeruginosa PAO1 library of mutants lacking cyclic-di-GMP-related genes. *Front*
Microbiol **13**, 949597 (2022).

31. Wei, Q. *et al.* Diguanylate Cyclases and Phosphodiesterases Required for Basal-
Level c-di-GMP in Pseudomonas aeruginosa as Revealed by Systematic
Phylogenetic and Transcriptomic Analyses. *Appl Environ Microbiol* **85**, (2019).

32. Cai, Y. M. *et al.* The c-di-GMP Phosphodiesterase PipA (PA0285) Regulates
Autoaggregation and Pf4 Bacteriophage Production in Pseudomonas aeruginosa
PAO1. *Appl Environ Microbiol* **88**, (2022).

33. Cai, Y. ming *et al.* Differential impact on motility and biofilm dispersal of closely related
phosphodiesterases in Pseudomonas aeruginosa. *Sci Rep* **10**, (2020).

34. Liu, S. *et al.* Priority changes between biofilm exopolysaccharides synthesis and
rhamnolipids production are mediated by a c-di-GMP-specific phosphodiesterase
NbdA in Pseudomonas aeruginosa. *iScience* **25**, 105531 (2022).

35. Li, Y., Heine, S., Entian, M., Sauer, K. & Frankenberg-Dinkel, N. NO-induced biofilm
dispersion in Pseudomonas aeruginosa is mediated by an MHYT domain-coupled
phosphodiesterase. *J Bacteriol* **195**, 3531–3542 (2013).

36. Zhan, X. *et al.* A c-di-GMP signaling module controls responses to iron in
Pseudomonas aeruginosa. *Nature Communications 2024 15:1* **15**, 1–15 (2024).

37. Stelitano, V. *et al.* C-di-GMP hydrolysis by Pseudomonas aeruginosa HD-GYP
phosphodiesterases: analysis of the reaction mechanism and novel roles for pGpG.
PLoS One **8**, (2013).

38. Ryan, R. P. *et al.* HD-GYP domain proteins regulate biofilm formation and virulence in
Pseudomonas aeruginosa. *Environ Microbiol* **11**, 1126–1136 (2009).

39. Matilla, M. A., Velando, F., Monteagudo-Cascales, E. & Krell, T. Flagella, Chemotaxis
and Surface Sensing. in *Advances in Experimental Medicine and Biology* (eds.

Filloux, A. & Ramos, J. L.) vol. 1386 185–221 (Advances in Experimental Medicine
and Biology, Springer, Cham, 2022).

40. Tchigvintsev, A. *et al.* Biochemical and structural studies of uncharacterized protein
PA0743 from Pseudomonas aeruginosa revealed NAD +-dependent L-Serine
dehydrogenase. *Journal of Biological Chemistry* **287**, 1874–1883 (2012).

41. Gliese, N. & Khodaverdi, V. AgmR controls transcription of a regulon with several
operons essential for ethanol oxidation in Pseudomonas aeruginosa ATCC 17933.
https://doi.org/10.1099/mic.0.26882-0 doi:10.1099/mic.0.26882-0.

42. Tassinari, M., Rudzite, M., Filloux, A. & Low, H. H. Assembly mechanism of a Tad
secretion system secretin-pilotin complex. *Nat Commun* **14**, 1–13 (2023).

43. Kalia, M., Resch, M. D., Cherny, K. E. & Sauer, K. The Alginate and Motility Regulator
AmrZ is Essential for the Regulation of the Dispersion Response by Pseudomonas
aeruginosa Biofilms. *mSphere* https://doi.org/10.1128/MSPHERE.00505-22 (2022)
doi:10.1128/MSPHERE.00505-22.

44. Zhao, J. *et al.* Molecular Mechanism of CdpR Involved in Quorum-Sensing and
Bacterial Virulence in Pseudomonas aeruginosa. *PLoS Biol* **14**, 1002449 (2016).

45. Zhao, J. *et al.* Structural and Molecular Mechanism of CdpR Involved in Quorum-
Sensing and Bacterial Virulence in Pseudomonas aeruginosa. *PLoS Biol* **14**, (2016).

46. Park, S. & Sauer, K. Controlling Biofilm Development Through Cyclic di-GMP
Signaling. in *Advances in experimental medicine and biology* (eds. Filloux, A. &
Ramos, J. L.) vol. 1386 69–94 (Advances in Experimental Medicine and Biology,
Springer, Cham, 2022).

47. Webster, S. S. *et al.* Force-Induced Changes of PilY1 Drive Surface Sensing by
Pseudomonas aeruginosa. *mBio* **13**, (2021).

48. Kazmierczak, B. I., Lebron, M. B. & Murray, T. S. Analysis of FimX, a
phosphodiesterase that governs twitching motility in Pseudomonas aeruginosa. *Mol
Microbiol* **60**, 1026–1043 (2006).

49. Klausen, M. *et al.* Biofilm formation by Pseudomonas aeruginosa wild type, flagella
and type IV pili mutants. *Mol Microbiol* **48**, 1511–1524 (2003).

50. Cowles, K. N. & Gitai, Z. Surface association and the MreB cytoskeleton regulate
pilus production, localization and function in Pseudomonas aeruginosa. *Mol Microbiol*
76, 1411–1426 (2010).

51. Irie, Y. *et al.* Self-produced exopolysaccharide is a signal that stimulates biofilm
formation in Pseudomonas aeruginosa. *Proceedings of the National Academy of
Sciences* **109**, 20632–20636 (2012).

52. Hengge, R. Principles of c-di-GMP signalling in bacteria. *Nature Reviews Microbiology*
vol. 7 Preprint at https://doi.org/10.1038/nrmicro2109 (2009).

53. Petrova, O. E. & Sauer, K. PAS domain residues and prosthetic group involved in
BdlA-dependent dispersion response by Pseudomonas aeruginosa biofilms. *J
Bacteriol* **194**, 5817–5828 (2012).

54.   Morgan, R., Kohn, S., Hwang, S. H., Hassett, D. J. & Sauer, K. BdlA, a Chemotaxis
584   Regulator Essential for Biofilm Dispersion in Pseudomonas aeruginosa. *J Bacteriol*
585   **188**, 7335 (2006).

586   55.   Stelitano, V. *et al.* C-di-GMP Hydrolysis by Pseudomonas aeruginosa HD-GYP
587   Phosphodiesterases: Analysis of the Reaction Mechanism and Novel Roles for pGpG.
588   *PLoS One* **8**, e74920 (2013).

589   56.   Ryan, R. P. *et al.* HD-GYP domain proteins regulate biofilm formation and virulence in
590   Pseudomonas aeruginosa. *Environ Microbiol* **11**, 1126–1136 (2009).

591   57.   Wood, T. L. *et al.* Rhamnolipids from Pseudomonas aeruginosa disperse the biofilms
592   of sulfate-reducing bacteria. *NPJ Biofilms Microbiomes* **4**, (2018).

593   58.   Hou, L., Debru, A., Chen, Q., Bao, Q. & Li, K. AmrZ Regulates Swarming Motility
594   Through Cyclic di-GMP-Dependent Motility Inhibition and Controlling Pel
595   Polysaccharide Production in Pseudomonas aeruginosa PA14. *Front Microbiol* **10**,
596   (2019).

597   59.   Giraud, C. *et al.* The PprA-PprB two-component system activates CupE, the first non-
598   archetypal Pseudomonas aeruginosa chaperone-usher pathway system assembling
599   fimbriae. *Environ Microbiol* **13**, 666–683 (2011).

600   60.   Bernard, C. S., Bordi, C., Termine, E., Filloux, A. & De Bentzmann, S. Organization
601   and PprB-dependent control of the pseudomonas aeruginosa tad locus, involved in
602   Flp Pilus Biology. *J Bacteriol* **191**, 1961–1973 (2009).

603   61.   Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using
604   Real-Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. *Methods* **25**, 402–408 (2001).

605