Transcriptional profiling of *Pseudomonas aeruginosa* biofilm life cycle stages reveals dispersal-specific biomarkers

Xavier Bertran i Forga[1,2], Kathryn E. Fairfull-Smith[3,4,] Jilong Qin[1,5*], and Makrina Totsika[1,2*]

[1]Centre for Immunity and Infection Control, School of Biomedical Sciences, Queensland University of Technology, Brisbane, QLD, Australia

[2]Max Planck Queensland Centre, Queensland University of Technology, Brisbane, QLD, Australia

[3]School of Chemistry and Physics, Queensland University of Technology, Brisbane, QLD, Australia

[4]Centre for Materials Science, Queensland University of Technology, Brisbane, QLD, Australia

[5]Institute for Molecular Bioscience, The University of Queensland, Brisbane, QLD, Australia

*correspondence: Jilong Qin (j.qin@uq.edu.au), and Makrina Totsika (makrina.totsika@qut.edu.au)

Abstract

Bacteria exhibit two lifestyles: planktonic free-floating individual cells or sessile multicellular aggregates known as biofilms. The biofilm lifecycle is characterised by three distinct stages: attachment, maturation and dispersal. Distinct adaptations occur in each stage, determining cellular behaviours such as surface attachment or synthesis and degradation of extracellular matrix components. Characterising stage-specific bacterial profiles therefore represents a valuable strategy for the development of novel antibiofilm therapies. Here, we used the model biofilm-forming bacterium *Pseudomonas aeruginosa* PAO1 to characterise the transcriptional profiles of each stage of the biofilm life cycle: attachment, biofilm maturation and spontaneous dispersal in closed cultures. We report that surface attachment was accompanied by the upregulation of genes comprising the Pil-Chp mechanosensory system, whereas biofilm maturation was characterised by the upregulation of genes involved in Pel polysaccharide synthesis, *siaD* and PA4396 diguanylate cyclases as well as *pipA, fimX* and PA5442. In contrast, dispersing cells upregulated genes responsible for the biosynthesis of alginate, rhamnolipid, and extracellular nucleases (*eddA*, *eddB*), as well as the transcriptional regulator of dispersal *amrZ*. Additionally, genes encoding the spontaneous dispersal molecule cis-2-decenoic acid (*dspS* and *dspI*), canonical phosphodiesterases (*nbdA* and *rbdA*), four non-canonical HD-GYP phosphodiesterases and seven other c-di-GMP–related enzymes were also upregulated during dispersal. Our comprehensive analysis of transcriptional changes across biofilm stages therefore provides benchmarking stage-specific transcriptional profiles for *P. aeruginosa* biofilms in closed culture systems. Furthermore, it allowed the identification of a subset of fourteen genes as transcriptional biomarkers of dispersal, which were used to build reporter plasmids as tools to determine the onset of dispersal.

Importance

Biofilm infections by *P. aeruginosa* are a major medical challenge due to the increased tolerance to antimicrobials displayed by bacteria living in sessile communities, which is reduced during spontaneous biofilm dispersal. Attachment, biofilm maturation and dispersal represent the main stages

of a dynamic process known as the biofilm lifecycle. However, the global regulatory responses governing transitions between these stages remain understudied. Here, we combine live microscopy and biomass quantification to track the progression of *P. aeruginosa* cultures through the three main stages of the biofilm lifecycle. We show that cells from each stage recapitulate canonical, stage-specific transcriptional responses and identify a set of biomarkers associated with the onset of dispersal. These biomarkers may offer a practical tool for rapidly screening dispersal-inducing compounds, aiding in the discovery of the next generation of antibiofilm therapeutics.

Introduction

Pseudomonas aeruginosa is a highly adaptable Gram-negative opportunistic pathogen, which survives in hostile environments by forming cell aggregates embedded in a protective extracellular matrix primarily composed of exopolysaccharides, extracellular DNA (eDNA) and structural proteins [1]. These aggregates, known as biofilms, exhibit increased tolerance to desiccation and host defences compared to their planktonic counterparts [2]. *P. aeruginosa* biofilms are associated with chronic infections, which exhibit high tolerance to a broad range of clinically employed antimicrobials [2].

In laboratories, *P. aeruginosa* biofilms are frequently studied using a variety of culture systems such as closed culture systems (i.e. microtiter plates) or open-flow systems (i.e. flow cells). The latter have been extensively used to identify and study the three main stages of the biofilm life cycle (attachment, biofilm maturation and dispersal), as well as distinct stage-specific molecular mechanisms [3,4]. During attachment, cells transition from the motile to sessile lifestyle. This process is mediated by the mechanosensors Pil-Chp and Wsp which lead to the upregulation of genes involved in the synthesis of matrix components and fimbrial adhesins such as CupA [5–7]. During biofilm maturation, the components of the matrix are progressively secreted into the extracellular space. As biofilms grow and mature, environmental cues and native intracellular signals may accumulate, such as the fatty acid dispersal signal cis-2-decenoic acid, eventually triggering spontaneous dispersal responses [8]. This phenomenon is characterised by biofilm-residing single cells regaining motility and resuming the planktonic lifestyle, during which they escape the biofilm matrix by hydrolysing its components [9].

This is achieved by the upregulation of glycosyl hydrolases such as PslG and PelA for

exopolysaccharide Psl and Pel degradation [9–11], and extracellular endonucleases EndA, EddA and

EddB [9–11]. This process is influenced by the intracellular concentration of the secondary messenger

cyclic-di-GMP (c-di-GMP) [6], controlled by a family of proteins harbouring diguanylate cyclase

domains (DGCs; possessing a GGDEF motif) and phosphodiesterase domains (PDE; possessing a

HD-GYP or EAL motifs), which respectively synthesise and degrade it [12–15].

The transition between the stages of the biofilm life cycle is driven by distinct signalling pathways. To

characterise them, transcriptomics has been applied to biofilms treated with biofilm-dispersing agents

such as NO donors or carbon sources, which primarily captured immediate transcriptional changes

triggered by these compounds leading to an abrupt interruption of the biofilm maturation process [16–18].

However, the progression through the biofilm stages is a dynamic process [19], urging the temporal

resolution of stage-specific gene expression. Therefore, we here performed RNA sequencing to

compare the currently understudied transcriptional variations across cells undergoing attachment,

biofilm maturation, and dispersal [33]. Subsequently, by tracking the transcription patterns of genes

throughout the three life cycle stages, we identified a subset of biomarkers for biofilm dispersal,

which were used to build a collection of reporter plasmids as screening tools for the identification of

the onset of dispersal.

Results

P. aeruginosa **biofilms cultured in closed systems recapitulate the three defined life**

stages of open-flow biofilms

We previously reported that the biofilm growth kinetics of the model organism *P. aeruginosa* PAO1 in

closed cultures displayed progressive accumulation of surface-attached biomass, reaching maximum

biomass between 2-8 h, followed by a marked reduction [19]. While the stages of the biofilm life cycle

(attachment, maturation and dispersal) have been primarily described in open-flow cultures, these

remain understudied in closed culture systems such as microtiter plates. To further characterise how

these stages manifest in closed culture systems, we quantified the biofilm biomass via crystal violet

staining on forming and dispersing biofilms at different time points (Fig 1A), which we combined

with microscopy to capture stage-specific morphological features using microtiter plates (Fig 1B, Fig

S1). Using this approach, we observed that planktonic cells initiate attachment within 2 h of

inoculation (Fig 1A and B; orange), during which cells adhered to the surface as a monolayer, with a

culture density (OD_{600}) of ~0.1 and a low total biomass (stained by crystal violet with OD_{550} of ~2.0).

At 4-8 h post-inoculation, cells transitioned to the biofilm maturation stage (Fig 1A and B; blue),

organised in voluminous scattered clusters that grew over time and produced multicellular biofilm

structures (Fig 1B, Fig S1), as observed by peak biofilm biomass staining at the 8 h time point (OD_{550}

of ~5; Fig 1A). This build-up in biomass occurred between the early (OD_{600} ~0.3) and the late-

logarithmic phase (OD_{600} ~1.5). Beyond 8 h post-inoculation and at a higher culture density

(OD_{600}>1.5), biofilm biomass rapidly decreased. This phenomenon was characterised by disassembled

structures leaving clearly distinguishable biofilm remnants attached to the well surface and a loss of

77% of accumulated biomass (Fig 1A and B, Fig S1; green). Collectively, our data indicate that cells

grown in closed culture systems synchronously transition through distinctive phases culminating in

spontaneous (endogenously induced) biofilm dispersal at defined time-points. Therefore, closed

biofilm systems can effectively recapitulate the stages of the biofilm life cycle originally described in

open-flow culture systems in comparatively shorter time courses [3,20,21].

Figure 1. Biofilm formation and dispersal kinetics of *P. aeruginosa* PAO1. (A) *P. aeruginosa* PAO1 biofilm biomass and
culture turbidity over 12 h. PAO1 cultures were seeded on 24-well plates and biofilm biomass was quantified at different
time points by crystal violet (CV) staining (●). Culture density was simultaneously monitored by OD_{600} (○) with means ± SD

shown. 3 biological replicates are represented. **(B)** Micrographs of the crystal violet-stained biofilms at each stage of the

biofilm life cycle. Crystal-violet-stained microscopic images of *P. aeruginosa* biofilm cultures in 24-wells. Images are

representative of at least 3 biological replicates. Orange: attachment stage. Blue: biofilm maturation stage. Green: dispersal

stage. This colour coding scheme is used throughout the article to differentiate the three biofilm stages.

Transcriptional profiling of *P. aeruginosa* biofilm cultures displays canonical pathways

of attachment, maturation and dispersal

While *P. aeruginosa* cells grown in microtiter plates undergo the three canonical stages of the biofilm

life cycle, RNA recovered from biofilm cells cultured in microtiter plates is typically insufficient in

quantity to support downstream RNA sequencing (RNA-seq). Therefore, we replicated the biofilm

kinetics in tissue culture flasks as a larger-volume closed culture system, monitoring biofilm cell

density in addition to total biomass via staining with crystal violet. Biofilm kinetics displayed by *P.*

aeruginosa PAO1 in tissue culture flasks mimicked the temporal pattern of microtiter plates in total

biomass and microscopically (Fig S1), showing attachment, biofilm maturation and dispersal within

the same time frames.

Using the flask culture system, we performed RNA sequencing of *P. aeruginosa* during attachment,

early biofilm maturation and dispersal to characterise stage-specific transcriptional profiles (Fig 2A).

This study design enabled temporal resolution of both global and individual gene transcription

variation across the biofilm life cycle stages. Accordingly, biofilm maturation was used as the central

reference point, as it represents the intermediate stage between attachment and dispersal, (Fig 2C).

Differential transcription analysis using a negative binomial test (P-value ≤ 0.01; Log_2 fold-change $\geq$

$|1|$) identified a total of 3361 genes with significant transcriptional changes across all three

comparisons: i) Attachment vs. Biofilm Maturation; ii) Biofilm Maturation vs. Dispersal (Planktonic);

and iii) Attachment vs. Dispersal (Table S1). Cells collected from the liquid phase and from biofilms

during at the dispersal stage showed few transcriptional differences, with only 83 differentially

regulated genes (76 upregulated and 7 downregulated) between the two states, and with low fold

changes (Table S1). This suggested that cells at the dispersal stage measured in this study

synchronously altered their transcriptomic profiles regardless of remaining surface-associated or

becoming free-swimming. Due to the planktonic phase of dispersal containing larger transcriptional

variations relative to other stage transitions (Fig 2B), we used it for all subsequent analyses. Principal

component analysis (PCA) further confirmed that cells at each stage formed distinct clusters,

indicating discrete transcriptional profiles were associated with attachment, maturation, and dispersal

(Fig 2B).

Figure 2. Transcriptional profiles of different stages of the biofilm life cycle. (A) Schematic depicting RNA sample

collection for sequencing. During attachment, the free-swimming phase was collected to represent cells rapidly transitioning

from motile to sessile. During biofilm maturation, surface-attached cells were collected, and cells from both phases were

collected during the dispersal stage. **(B)** Principal component analysis of samples collected at distinct stages of the biofilm

life cycle (attachment – orange; biofilm maturation – blue; the planktonic phase of biofilm dispersal – light green; and the

biofilm phase of biofilm dispersal – dark green). Each biological replicate of a group is represented as a dot of the same

colour. The normalised reads of all genes were then extracted for each sample and transformed into PC loadings with

ClustVis [22]. Values for PC1 and PC2, accounting for the largest variance percentage are represented in the plot. **(C)** Venn

diagrams showing differentially transcribed genes in cells collected from the liquid phase undergoing attachment and

dispersal (relative to biofilm maturation). Downregulated (top) and upregulated (bottom) genes at each stage relative to

biofilm maturation are represented separately. Overlapping areas indicate genes with consistent transcriptional changes

relative to biofilm maturation stage.

Building on these global transcriptional variations and to further validate our datasets, we mined our

RNA-seq data for previously reported stage-specific changes in the transcription of genes previously

associated with attachment, biofilm formation or dispersal. During attachment and extending into

biofilm maturation, we observed the upregulation of the Pil-Chp surface-sensing complex known to

mediate attachment [7], whereas no upregulation of the complementary Wsp system was noted. Notably,

transcription of *pilA, pilB, pilD* and *pilGHIJK-chpABCDE,* encoding type IV pili structural and signal

transduction proteins, and of *cyaB* and *vfr,* which encode an adenylate cyclase and its associated

cyclic-AMP-controlled transcriptional regulator Vfr, was upregulated between 2.00- and 7.37-fold

relative to biofilm dispersal.

Biofilm maturation involves synthesis pathways for extracellular polysaccharides and eDNA, which

are the main components of the biofilm matrix [1]. During biofilm maturation, we observed the

upregulation of the exopolysaccharide Pel synthesis operon *pelBCDEF* but not *pelG* relative to

attachment (2.22 – 2.26-fold) and dispersal stages (2.16 – 2.99-fold) (Fig 3A, Table S1) [23].

Additionally, the alternative respiration process of denitrification is known to be active during biofilm

maturation due to cell layering inside the biofilm restricting oxygen availability [24]. Accordingly, we

found biofilm cells to upregulate the transcription of the nitrite reductase complex (*nirSMCFDLG*) by

up to 10-fold relative to attachment.

Planktonic cells at the biofilm dispersal stage, relative to attachment and biofilm maturation stages,

showed an increased transcription of genes involved in eDNA hydrolysis (*eddA,* 3.28-fold; and *eddB,*

2.95-fold; but not *endA*) (Fig 3A, Table S1), suggesting the remodelling of the extracellular matrix to

allow the escape of planktonic cells. We also observed a pronounced upregulation of quorum-sensing–

regulated genes previously shown to drive biofilm dispersal. *rhlA*, *rhlB*, and *rhlC*, which encode

enzymes for the biosynthesis of the rhamnolipid biosurfactant implicated in quorum-sensing–

mediated dispersal [39–42] were upregulated by 88-, 45- and 12-fold, respectively (Fig 3A, Table S1) [25–28].

Another quorum-sensing pathway activated during dispersal involved the fatty acid signal cis-2-

decenoic acid [27,29]. This fatty acid signal is produced and sensed by the proteins encoded by *dspI* and

dspS, which were upregulated by 2.04- and 5.59-fold, respectively, during dispersal relative to the

biofilm maturation stage (Fig 3A, Table S1). Interestingly, genes associated with biofilm maturation

were also found upregulated during the dispersal stage. Adhesion factors, such as lectins *lecA* and

lecB were highly upregulated during dispersal by 279-fold and 115-fold, respectively, relative to the

biofilm maturation stage. Similarly, *flp*, which encodes the main subunit of the type IVb pili adhesin

was upregulated by 120-fold, together with several alginate biosynthesis genes (*alg8*, 2.26-fold*; algX,*

2.16-fold*; algJ*, 2.39-fold; *algA,* 2.74-fold) and their associated regulators *algU* (2.49-fold)*, algR*

(5.76-fold) and *amrZ* (6.27-fold) (Fig 3A, Table S1). These findings were consistent with previous

studies showing that dispersed cells had significantly increased functional adhesion relative to

planktonic cells [30]. Therefore, our RNA-seq data captured transcriptional changes associated with

previously documented attachment, biofilm maturation and dispersal responses, further underscoring

the consistency of our dataset with established biofilm regulatory pathways and the robustness of our

analytical methods.

To further validate the upregulation of these known dispersal pathways, we extracted RNA from

biofilms at 4 h (early maturation), 8 h (late maturation), and planktonic cells collected at 12 h

(dispersal) and measured the transcript levels of representative genes involved in rhamnolipid

synthesis (*rhlAB, rhlC*), eDNA degradation (*eddA, eddB*), and cis-2-decenoic acid signalling (*dspI,*

dspS) by RT-qPCR (Fig 3B). Relative to maturing biofilms, all genes were upregulated in mature and

dispersed biofilms. These findings further validate our RNA-seq data and indicate that mature

biofilms may begin to transcriptionally activate dispersal pathways before the onset of active

detachment.

Figure 3. Genes involved in dispersal mechanisms are upregulated during late biofilm maturation and dispersal compared to early biofilm maturation. (**A**) Comparison of the transcriptomes of planktonic cells during attachment or

226 dispersal to that of sessile cells during biofilm maturation. Genes containing Log2-fold change transcription values >|1| were

227 highlighted in biofilm stage specific colours. (**B and C**) RT-qPCR analysis of canonical dispersal genes (*rhlA, rhlB, rhlC,*

228 *eddA, eddB, dpsS* and *dspI*) and the PDEs PA4108 and PA4781, which were identified as putatively involved in dispersal.

229 RNA samples were extracted at different time points with Means ± SEM shown.

231 Specific c-di-GMP enzymes characterise the biofilm maturation and dispersal
232 transcriptomes

Cyclic di-GMP (c-di-GMP) is a key signalling molecule governing the transition from biofilm

maturation to dispersal. Considering that the roles of enzymes responsible for its synthesis and

breakdown are stage-specific, we also analysed our transcriptomic data to identify genes encoding

enzymes with predicted PDE (EAL or HD-GYP) or DGC (GGDEF) domains (Table 1). Expectedly,

genes encoding proteins essential for biofilm development (*siaD,* 3.73-fold and *pipA,* 2.08-fold) were

upregulated during biofilm maturation [31–34]. Genes implicated in the formation of large microcolonies

[32], including PA4396 (2.08-fold), *fimX* (2.77-fold), and PA5442 (2.36-fold) were also significantly

upregulated. In contrast, we observed an upregulation of a combination of PDEs and DGCs in cells at

the dispersal stage, including sensory transducers of dispersal cues *nbdA* (7.56-fold) and *nicD* (6.73-

fold) [35–37]. Additionally, upregulation of *ismP* (12.99-fold) was noted, which encodes an iron sensor

that under low iron conditions hijacks and inhibits a cognate DGC ImcA [38].

Table 1: Differentially regulated genes encoding c-di-GMP-modifying enzymes during biofilm
maturation and dispersal.

Locus Tag	Gene name	Domain(s)	Activity	Fold-change
Biofilm maturation				(vs Dispersal)
PA0169	*siaD*	GGDEF	DGC	3.73
PA0285	*pipA*	GGDEF-EAL	PDE	2.08
PA4396	-	GGDEF	-	2.08
PA4959	*fimX*	GGDEF-EAL	PDE	2.77
PA5442	-	GGDEF-EAL	PDE	2.36

Biofilm dispersal				(vs Maturation)
PA0290	-	GGDEF	-	2.73
PA0575	*rmcA*	GGDEF-EAL	PDE	3.51
PA0847	-	GGDEF	DGC	2.09
PA0861	*rbdA*	GGDEF-EAL	PDE	3.03
PA1878	-	HD-GYP	-	3.39
PA2072	*ismP*	GGDEF-EAL	-	12.99
PA2572	-	HD-GYP	-	8.22
PA2771	-	GGDEF	DGC	2.75
PA3311	*nbdA*	GGDEF-EAL	PDE	7.56
PA3825	-	EAL	PDE	1.25
PA4108	-	HD-GYP	PDE	5.60
PA4781	-	HD-GYP	PDE	10.70
PA4929	*nicD*	GGDEF	DGC	6.73

Interestingly, dispersed cells upregulated all four genes encoding proteins with HD-GYP

phosphodiesterase domains present in the *P. aeruginosa* PAO1 genome (PA1878, PA2572, PA4108

and PA4781) (Table 1). Two of these, PA4108 and PA4781 were previously reported to have PDE

activity and to be necessary for swarming and the production of virulence factors [39,40]. We further

validated the transcriptional changes of these two genes using RT-qPCR (Figure 3C), confirming an

increase in expression of 7.5-fold and 4-fold, respectively, during dispersal relative to maturing

biofilms at 4 h. Moreover, the transcription of both genes was also upregulated in mature biofilms

(6.2- and 4-fold; Figure 3C), indicating a similar pattern of transcriptional priming prior to dispersal.

Altogether, our results indicate that canonical stage-specific c-di-GMP modulation responses are

accurately reflected in our transcriptomic dataset and suggest a potential undescribed role of HD-

GYP-containing PDEs in mediating biofilm dispersal.

Transcriptional biomarkers identify the onset of biofilm dispersal

Since the obtained transcriptomic profiles recapitulated canonical pathways associated with

attachment, maturation, and dispersal, our datasets are uniquely well suited to identify gene changes

associated with the dispersal phase of *P. aeruginosa* and serve as transcriptional biomarkers of biofilm

dispersal. We reasoned that genes directly involved in biofilm dispersal would exhibit a distinct

temporal pattern of transcription such as an inverse regulation between the biofilm maturation and

dispersal stages relative to the initial attachment stage. Thus, pro-dispersal genes would be repressed

during maturation but strongly upregulated during dispersal, or vice versa. To identify candidate

biomarkers, we applied these selection criteria and identified 14 upregulated and 8 downregulated

genes that could potentially serve as dispersal biomarkers (Table 2).

Among the upregulated genes, *cheR2* (2.1-fold, Table 2) encodes a methyltransferase component of

the Che2 chemotaxis pathway, which interacts with the heme-containing oxygen sensor McbP/Aer2

and the coupling protein CheW2 to modulate virulence through a cryptic phosphorylation cascade

[41,42]. Another notable gene, *pqqA* (2.55-fold; Table 2), is part of the *pqqABCDE* operon responsible for

the biosynthesis of pyrroloquinoline quinone (PQQ), a redox cofactor for the quinoprotein ethanol

dehydrogenases ExaA and ExaBC [43]. In addition, several genes under the transcriptional control of the

two-component system PprAB were also upregulated. These include *tadA*, *rcpA*, *rcpC*, and *flp* (13,66-

, 12.47-, 13.83- and 34.54-fold; Table 2) which are part of a contiguous cluster spanning PA4297

(*tadG*) to PA4306 (*flp*). This locus encodes the machinery for the synthesis and secretion of Flp type

IVb pili [44]. Notably, two of the dispersal biomarkers *amrZ* and PA2588 (*cdpR*) have known roles in

modulating biofilm physiology. *amrZ* encodes a recognised transcriptional regulator of dispersal and

cdpR is a quorum sensing and virulence transcriptional regulator [45–47].

Several uncharacterised genes, including PA0111, PA0743 and PA1353 were also identified as

upregulated biomarkers of biofilm dispersal, many of which appear to be associated with redox

processes. While no experimental data are available for PA0111, bioinformatic predictions suggest

that the directly upstream gene PA0110 encodes a protein putatively related to a mitochondrial

cytochrome *c* oxidase. Additionally, PA0112 and PA0113 are predicted to encode a heme A synthase

and a protoheme IX farnesyltransferase, respectively, implicating these loci in cytochrome maturation

or electron transport. Similarly, PA0743, which encodes an L-serine dehydrogenase, was also

upregulated, potentially reflecting altered amino acid catabolism related to redox homeostasis during

biofilm dispersal. No functional predictions are currently available for PA1353, which was also part of

the identified dispersal biomarkers.

Table 2. Genes differentially regulated at the biofilm maturation and dispersal stage compared

to the attachment stage.

Gene	Fold change in Biofilm maturation [a]	Fold change in Biofilm dispersal [a]	Gene product
Downregulated at 4h and upregulated at 8h			
PA0111	-2.01	12.21	hypothetical protein
*cheR*2	-2.22	2.10	chemotaxis protein methyltransferase
PA0743	-2.58	3.12	NAD-dependent L-serine dehydrogenase
PA1353	-2.36	3.32	hypothetical protein
pqqA	-5.10	2.55	coenzyme PQQ synthesis protein A
PA2588	-2.03	3.01	transcriptional regulator
amrZ	-2.30	2.55	alginate and motility regulator Z
tadA	-2.17	13.64	ATPase TadA
rcpA	-2.13	12.47	type II/III secretion system protein
rcpC	-2.07	13.83	hypothetical protein
flp	-3.27	34.54	type IVb pilin Flp
PA4523	-2.31	3.41	hypothetical protein
cupE1	-2.11	9.58	fimbrial subunit CupE1
cupE2	-2.04	3.95	fimbrial subunit CupE2
Upregulated at 4h and downregulated at 8h			
PA0485	2.93	- 2.91	hypothetical protein
PA1170	2.43	-4.76	hypothetical protein
metH	2.45	-2.13	B12-dependent methionine synthase
ilvH	3.53	-2.35	acetolactate synthase small subunit
ilvI	4.69	-2.57	acetolactate synthase 3 catalytic subunit
cysN	2.1	-2.62	bifunctional sulfate adenylyltransferase subunit1/adenylylsulfate kinase
PA4889	2.06	-14.92	oxidoreductase
PA4907	2.41	-2.27	short-chain dehydrogenase

[a] Relative to samples collected during attachment.

To evaluate the wider use of these biomarkers beyond the reference PAO1 strain, we investigated gene

presence across 1301 *P. aeruginosa* genomes. Indeed, we identified that across a diverse range of

genomic backgrounds, all fourteen transcriptional biomarkers of dispersal are present in ~100% of

complete *P. aeruginosa* genomes available in GenBank, with only *flp* showing an average pairwise

identity <99% (Table S2). We restricted our biomarker screen to upregulated transcripts, as elevated

expression levels are more consistently detected and validated across analytical platforms. Due to the

similarities in transcriptomic profiles between planktonic and biofilm cells collected during dispersal,

we hypothesised that biofilm cells initiate the expression of dispersal biomarkers prior to

phenotypically manifesting the dispersal response in culture. Therefore, we collected samples from

cells undergoing late biofilm maturation and dispersal and validated these biomarker genes by RT-

qPCR (Fig 4). Our results confirmed the upregulation of all fourteen genes during dispersal.

Furthermore, all genes but four (PA0743, PA1353, *pqqA*, and PA2588) were also significantly

upregulated during late biofilm maturation (Fig 4). Genes PA0111, *cheR2*, *tadA*, *rcpC*, *flp*,

and PA4523 showed a pattern where transcript abundance increased in mature biofilms (2.5- to 23-

fold) and peaked during dispersal (4.6- to 117-fold relative to early biofilms) (Fig 4). In contrast, the

transcription of *cupE1* and *cupE2* fimbrial genes peaked in mature biofilms (54.41- and 10.1-fold) but

sharply declined upon dispersal (25.55- and 5.9-fold) (Fig 4). This transient transcriptional profile

suggests that CupE fimbriae genes are likely characteristic of mature biofilms. In contrast,

transcription of PA0111, *cheR2, tadA, rcpC, flp* and PA4523 peaked at dispersal (Fig 4). Altogether,

these biofilm dispersal biomarkers exhibited strong reproducibility, and their increased transcription

prior to the onset of dispersal marks biofilm cells undergoing late biofilm maturation that are ready for

an incoming stage transition.

Figure 4. Validation of the transcriptional biomarkers of biofilm dispersal. Differences in transcriptional regulation

measured by RT-qPCR for the indicated genes from total RNA samples extracted at 4 h (early biofilm maturation), 8 h (late

biofilm maturation) and 12 h (dispersal). Fold-changes at 8 h and 12 h are relative to 4 h. Means ± SEM are shown. Two

technical replicates are represented. Statistical differences between groups were calculated using Student *t*-tests (*, *P*-value

<0.05).

Reporter plasmids as a low-cost rapid tool for monitoring biofilm dispersal

While RT-qPCR provides a direct and sensitive method to quantify transcriptional differences, we

constructed a more scalable reporter system using plasmids carrying *lacZ* transcriptional fusions to the

predicted promoter regions of each biomarker, excluding co-operonic genes (*rcpC* also represents

rcpA and *tadA*, and *cupE1* also represents *cupE2*). To evaluate this approach, *P. aeruginosa* PAO1

strains harbouring each reporter plasmid were cultured to early biofilm maturation (4 h) and dispersal

(12 h). β-galactosidase activity was significantly increased for 9 of 11 reporters, with only *cheR2* and

flp showing elevated levels of enzymatic activity at both early biofilm maturation and dispersal (Fig

5). In contrast, reporters containing PA0110, PA0743, PA1353, *pqqA*, PA2588, *amrZ*, *rcpC/rcpA*,

PA4523, and *cupE1/cupE2* showed significantly increased enzymatic activity during dispersal relative

to early biofilm maturation (Fig 5). Negative enzymatic activity values for *amrZ* at 4 h indicate a very

low presence of LacZ, where background noise levels (OD$_{550}$) are similar to total recorded activity

(OD$_{420}$; see Methods). Collectively, these results demonstrate that *lacZ* transcriptional reporter

plasmids provide a low-cost and rapid detection method for the transcriptional activation of the

identified biomarkers of dispersal and further validate these genes as robust biomarkers of the onset of

dispersal.

Figure 5. LacZ reporter plasmids indicate the upregulation of the identified biomarkers during dispersal. Differences in transcriptional regulation of biomarkers of dispersal measured by enzymatic activity of β-galactosidase in *P. aeruginosa* PAO1 in biofilm samples extracted at 4 h (early biofilm maturation) and 12 h (dispersal). Means ± SEM are shown. Three technical replicates are represented. Statistical differences between groups were calculated using Student's *t*-tests (**, *P*-

345 value <0.01; ***, *P*-value <0.001; ****, *P*-value <0.0001). One representative experiment is shown from two independently
346 performed assays.

Discussion

The study of transcriptional changes between each biofilm stage provides valuable insights into the
key cellular signalling pathways underlying their transition between sessile and motile lifestyles.
Here, we characterised the transcriptional profiles of cells undergoing attachment, biofilm maturation
and dispersal, and identified 23 biomarkers characterising the dispersal response, 14 of which
comprising a subset upregulated upon the initiation of dispersal.

Cells undergoing attachment sense and respond to surface contacts through mechanosensory systems
such as the Pil-Chp machinery, which relies on type IV pili-mediated twitching motility [48,49]. In our
study, we report an upregulation of type IV pili biogenesis genes (*pilA, pilB* and *pilD*) and of the
mechanosensory system Pil-Chp (*pilGHIJK-chpABCDE*) during the attachment and biofilm
maturation stages compared to dispersing cells. These were accompanied by the upregulation of *fimX*,
which encodes a putative phosphodiesterase essential for the assembly of type IV pili, and of *cupA1*
encoding the main subunit of the CupA adhesin [50]. Our findings therefore align with previous studies
indicating that type IV pili are indispensable to surface-attachment in microplates [51], and are
consistent with prior reports that surface-sensing promotes the transcriptional upregulation of genes
involved in type IV pilus biogenesis [52].

The intracellular messenger c-di-GMP acts as a master regulator of the bacterial lifestyle, with its
accumulation leading to biofilm formation [9]. Our dataset revealed the upregulation of several stage-
specific genes encoding proteins with validated or hypothetical c-di-GMP modulating activity.
Specifically, *siaD* was the most upregulated during biofilm maturation, which encodes a DGC
involved in early microcolony formation and adhesion [48,53]. In contrast, biofilm dispersal is associated
with increased PDE activity and decreasing intracellular c-di-GMP [54]. This was reflected in our
dataset, where 6 genes encoding functionally active PDEs were upregulated. Genes *rbdA* and *nbdA*
encode enzymes with well-described roles in biofilm dispersal. Both have been reported to modulate

flagellar motility, with RbdA being a key component of the BdlA-regulated core dispersal response

[55,56]. Moreover, four upregulated genes (PA1878, PA2572, PA4108 and PA4781) encode PDEs

carrying the HD-GYP motif [57]. While the role of these genes in the biofilm dispersal stage is unclear,

it has been reported that disruption of PA4108 and PA4781 resulted in higher intracellular c-di-GMP

and decreased swarming activity, whereas PA2572 modulated swarming motility [57,58]. We here

validated by RNA-seq and RT-qPCR that both PA4108 and PA4781 are strongly upregulated during

late biofilm maturation and dispersal, which suggests a potential role in initiating dispersal responses.

Additionally, recent reports showed that deletion of *cdpR* led to increased biofilm biomass and

decreased swarming motility [47]. Considering that the upregulation of *cdpR* is strongly controlled by

the quorum sensing regulator RsaL [46], the stage-specific transcriptional activation of *cdpR* would

therefore suggest that dispersal in closed systems is mediated by a global quorum sensing-mediated

swarming response.

The synthesis of the rhamnolipid biosurfactant is important to the quorum sensing-mediated swarming

motility and dispersal [26]. In closed culture systems, rhamnolipid was proposed to drive biofilm

dispersal [59]. Here we report that the rhamnolipid and swarming modulator PA2572 together with

several rhamnolipid-associated genes were highly upregulated during dispersal. Moreover, four of the

identified PDEs upregulated during dispersal, including PA4108, PA4781, *nbdA* and *rbdA,* were

previously shown to influence the synthesis of rhamnolipid leading to loss of biofilm biomass [20,36,58].

Altogether, these data suggest biofilm dispersal in closed systems may be driven by rhamnolipids.

Among the upregulated biomarkers, *amrZ* was reported as a global regulator controlling the

upregulation of genes related to alginate biosynthesis and Type IV pili twitching motility while

downregulating Pel synthesis, therefore driving biofilm dispersal [45,60]. In contrast, *tadA*, *rcpA*, *rcpC*,

flp, cupE1 and *cupE2*, which are regulated under the PprAB two-component system [61,62], were

reported to mediate surface attachment. Our data therefore support previous studies indicating that

dispersing cells exhibit a hyperadhesive potential that primes them to re-attach to surfaces [30].

Additionally, we identified PA0110, *cheR2*, *pqqA,* PA0743, PA1353 and PA4523 to be uniquely

associated with the dispersal stage, yet their specific roles remain to be defined. Their marked

upregulation during dispersal warrants further investigation.

Understanding biofilm dispersal may represent a critical step toward combating chronic biofilm-

associated infections. As dispersing cells become resensitised to antimicrobial treatment once they are

released from the protective biofilm matrix, the underlying transcriptional changes occurring during

biofilm dispersal hold promise for unveiling the next generation of antibiofilm treatments and

adjuvants. We here report the transcriptional profiles of the *P. aeruginosa* biofilm life cycle, including

attachment, biofilm maturation and dispersal stages in closed systems, which can be used as

benchmarks for future studies. Additionally, we propose a set of biofilm dispersal biomarkers that are

robustly activated at the onset of biofilm dispersal. As the upregulation of these biomarkers was

consistently reproduced by transcriptomic and RT-qPCR analyses, we generated gene fusion reporters

that enable the rapid monitoring of promoter activities *in vivo,* as well as being adaptable to high-

throughput (i.e. microplate-based) platforms. Therefore, these reporter plasmids present with potential

application in high throughput screens supporting the discovery and development of novel dispersal-

inducing antibiofilm therapeutics.

Materials and Methods

Strains, media and culture conditions

Pseudomonas aeruginosa PAO1 wild-type (WT). Cultures were routinely grown overnight in LB

(lysogeny broth) media at 37^{o}C, 200rpm before incubation in fresh M9 media (9 mM NaCl, 22 mM

KH_2PO_4, 48 mM Na_2HPO_4, 19 mM NH_4Cl and 2 mM $MgSO_4$, 100 µM $CaCl_2$, 0.4% glucose, pH 7.0)

for each experiment. Where appropriate, cultures were supplemented with 10µg/ml of gentamicin.

Biofilm kinetics and microscopy

Biofilms were grown on 24-well plates or tissue culture flasks (Nunc, ThermoFisher Scientific) by

inoculating 10^{7} colony forming units (CFU)/ml of overnight bacterial cultures for 2 h, 4 h, 8 h and

12h diluted in M9 media. 1 ml of culture was inoculated in each well of microtiter plates and 10 ml

were used in tissue culture flasks. Cultures were incubated at 37 ^{o}C and 180 rpm (microtiter plates) or

70 rpm (tissue culture flasks). Shaking speed was adjusted to prevent splashing. At the indicated time

points, supernatants were removed, and wells were delicately washed with 1ml phosphate buffered

saline (PBS, Gibco) to remove planktonic cells. Microscopy (PrimoVert, Zeiss) was used to confirm

the removal of planktonic cells and preservation of biofilm structures. Remaining biofilm cells were

stained with 0.1% (w/v) crystal violet dissolved in 6.25% (v/v) methanol. Stained biofilms were

washed twice with PBS and solubilised with ethanol. Biofilm biomass was quantified by measuring

the optical density at 550 nm (OD$_{550}$) using a SPECTROStar Nano microplate reader (BMG

LabTech). Micrographs of stained biofilms were taken via microscopy as described above.

RNA sequencing and analysis

Biofilms were cultured and dispersed as previously described with some adjustments [19]. Briefly, tissue

culture flasks (Nunc, ThermoFisher Scientific) were inoculated with 10^7 CFU/ml cells in 50 ml of M9

medium from overnight cultures. Cultures were incubated at 37^{o}C shaking at 70 rpm for 2, 4 or 8 h.

At 2 h and 8 h, planktonic cells (10^8) were collected from the liquid phase and immediately mixed

with 2 volumes of RNA-protect solution (QIAGEN). Biofilm cells were gently rinsed with 10 ml of

PBS and resuspended in 6 ml of RNAprotect (QIAGEN, Cat# 76506) mixed with 3ml of PBS using

cell scrapers. Cells (10^8) were subsequently pelleted at 5000 g (10 min, 25 ^{o}C). RNA was extracted

with the RNeasy mini kit (QIAGEN, Cat# 74104) as per the manufacturer's protocol for Gram

negative bacteria grown in minimal media. Extracted RNA samples were treated with DNase I and

rRNA was removed using streptavidin-coated magnetic beads (BGI Genomics, Shenzhen, China).

Library was prepared by RNA fragmentation and cDNA synthesis followed by the ligation of adaptors

at the 3' end of cDNA fragments. These were further amplified by PCR and single-stranded fragments

were circularised for downstream DNBSEQ PE100 sequencing (BGI Genomics, Shenzhen, China).

Clean reads were mapped to the *P. aeruginosa* PAO1 reference genome (GenBank Accession number

AE004091.2) with Bowtie2 v2.4.5 [63]. Geneious Prime v2024.0.7 was used to generate the read count

matrix and to subsequently run DESeq2 v1.50.2 to compare the expression levels between sample

groups (attachment, biofilm maturation and dispersal).

The Principal Component Analysis (PCA) plot was conducted exclusively as an exploratory

visualisation to assess overall transcriptomic similarity within biological replicates and to examine

separation between treatment groups. It was generated by extracting the reads per kilobase of

transcript per million mapped reads (RPKM) of every gene at each one of the stages (Table S1). Then,

genes with $<|1|$ $\log_2$ fold-change transcription or P-value >0.01 for all groups were discarded, and the

remaining differentially regulated genes were plotted using GraphPad Prism 10.4.1 (Graph Pad,

USA).

Gene sequence conservation analysis

A local database was generated with 1301 complete *P. aeruginosa* genomes (taxid: 287) were

downloaded from GenBank, excluding atypical, metagenome-assembled genomes and genomes from

large multi-isolate projects. Gene sequences were extracted from the annotated *P. aeruginosa* PAO1

reference genome AE004091.2. BLAST analyses were performed locally using the NCBI BLAST

plugin for Geneious Prime v2024.0.7 against the local *P. aeruginosa* database. In-built Megablast

analyses were configured with minimum E value of 10^{-8} and query cover of 75%.

RT-qPCR

To validate RNA-seq data, total RNA was extracted from cells harvested from different biofilm stages

as described above. cDNA was transcribed using 1 µg of total RNA using the SuperScript III first-

strand synthesis kit (Invitrogen) as per the manufacturer's protocol. cDNA samples were then used as

templates for RT-qPCR conducted with the Quantinova SYBR Green master mix (Qiagen) as per the

manufacturer's protocol. The housekeeping gene *recA* was used as a control. Each sample was

independently assayed twice with the gene-specific primers listed in Table S2. Relative gene

expression was calculated using the $2^{-\Delta\Delta CT}$ method [64]. Transcriptional fold changes were plotted using

GraphPad Prism.

Construction of *lacZ* fusion reporter plasmids and β-galactosidase assays

The cloning vector carrying a promoterless *lacZ* was constructed via Gibson Assembly with a 3.2 kb

fragment containing *lacZ* from *E. coli* K-12 MG1655, a 150 bp fragment containing the MCS of

pBBR1-MCS5 and a 3.8 kb fragment containing a gentamicin resistance cassette and the pBBR1

origin of replication from pBBR1-MCS5, resulting in the reporter construct pBBR1-lacZ. Gene

promoter regions were predicted using Sapphire.cnn.pseudomonas [65]. The 5' untranslated region

(~500 bp) of each gene/operon containing putative promoters were cloned into the MCS by BamHI

and HindIII upstream of *lacZ*. Reporters were introduced into *P. aeruginosa* PAO1 via

electroporation.

β-galactosidase-based reporter assays were performed as previously described [66]. Briefly, 10^8 cells

were collected by centrifugation from biofilm phases at early biofilm maturation or biofilm dispersal

stages and OD_{600} was recorded. Cells were lysed with 1mg/ml lysozyme in TE buffer (10 mM Tris-

HCl, 1 mM EDTA). Cell lysates were resuspended in 1ml of Z-buffer (0.06 M Na_2HPO_4, 0.04 M

NaH_2PO_4, 0.01 M KCl, 0.001 M $MgSO_4$, 0.05 M β-mercaptoethanol) and 160 µl were transferred to a

96-well plate in triplicate (Nunc, ThermoFisher Scientific). Then, 40 µl of an ONPG solution

(4mg/ml) in phosphate buffer (0.06M Na_2HPO_4, 0.04 M NaH_2PO_4; pH 7.0) were added to each well

and reactions were incubated for 2 h at 28 °C in a plate reader. Absorbances were recorded at 420nm

and 550nm every 10 min and Miller units (MU) were calculated using the following equation:

$$MU = 1000 \text{ x} \frac{OD420 - 1.75 * OD550}{t * v * OD600}$$

Where *t* is the reaction time, *v* is the volume of culture used and OD600 is the density of the culture

prior to centrifuging.

Statistical differences in enzymatic activity between cells collected at early biofilm maturation or

biofilm dispersal were calculated using Student t-tests (**, P-value <0.01; ***, P-value <0.001; ****,

P-value <0.0001). One representative experiment is shown from two independently performed assays.

Table 3: Bacterial strains and plasmids used in this study

Strain or Plasmid	Relevant Characteristics	Source
Strains		
P. aeruginosa PAO1	Wild-type, HER-1018 P1	[67]

E. coli DH10β	F–*mcr*A Δ(*mrr-hsd*RMS-*mcr*BC) φ80*lacZ*ΔM15 Δ*lac*X74 *rec*A1 *end*A1 *ara*D139 Δ(*ara-leu*)7697 *gal*U *gal*K λ–*rps*L(StrR) *nup*G	Invitrogen, Cat# 18297010
E. coli K-12 MG1655	*E. coli* K-12, Source of *lacZ*	[68]

Plasmids

pBBR1-MCS5	Gm^r, lac promoter, source of MCS5, resistance cassette and ori.	[69]
pBBR1-lacZ	Gm^r, promoterless *lacZ*.	This study
pBBR1-lacZ-PA0110	Gm^r, *PA0110::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-PA0179	Gm^r, *PA0179::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-PA0743	Gm^r, *PA0743::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-PA1353	Gm^r, *PA1353::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-PA1985	Gm^r, *PA1985::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-PA2072	Gm^r, *PA2072::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-PA2588	Gm^r, *PA2588::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-PA4305	Gm^r, *PA4305::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-PA4648	Gm^r, *PA4648::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-*amrZ*	Gm^r, *amrZ::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-*nicD*	Gm^r, *nicD::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-*nbdA*	Gm^r, *nbdA::lacZ* transcriptional fusion.	This study
pBBR1-lacZ-*flp*	Gm^r, *flp::lacZ* transcriptional fusion.	This study

Acknowledgements

This work was funded by an Australian Research Council project grant (DP210101317), the Max
Planck Queensland Centre on the Materials Science of Extracellular Matrices, and a QUT Amplify
Scholarship provided by the Queensland University of Technology (Australia) to XB. The Ian Potter
Foundation sponsored the CLARIOStar high-performance microplate reader (BMG, Australia). The
funders had no role in study design, data collection and analysis, decision to publish, or preparation of
the manuscript. The authors would like to thank Professor Robert EW Hancock (University of
Columbia) for providing the *P. aeruginosa* PAO1 strain used in this study.

Competing interests

506 MT is an employee of the GSK group of companies. All other authors declare no competing interests.
507 This research was conducted in the absence of any commercial or financial relationships that could be
508 constructed as a potential conflict of interest.

Author contributions

510 JQ and MT conceptualised the project. XB, JQ and MT contributed to experimental design. XB
511 conducted all experiments, and contributed to data collection, analysis and visualisation. XB, JQ and
512 MT contributed to data interpretation. JQ, MT and KFS supervised the project. KFS and MT obtained
513 the funding. XB wrote the manuscript draft, JQ substantially revised the manuscript, all authors edited
514 the manuscript.

Data availability

516 The RNA-Seq data have been deposited in the NCBI Short Read Archive (SRA) database with
517 accession code PRJNA1291829.

References

519 1. Flemming, H. C. & Wingender, J. The biofilm matrix. *Nat. Rev. Microbiol.* **8**, 623–633 (2010).

520 2. Høiby, N., Johansen, H. K., Moser, C. & Ciofu, O. Clinical Relevance of Pseudomonas
521 aeruginosa: A Master of Adaptation and Survival Strategies. in *Pseudomonas: Model*

Organism, Pathogen, Cell Factory (ed. Rehm, B.) 25–44 (John Wiley & Sons, Ltd, Weinheim, 2008). doi:10.1002/9783527622009.CH2.

3.   Sauer, K., Camper, A. K., Ehrlich, G. D., Costerton, J. W. & Davies, D. G. Pseudomonas aeruginosa displays multiple phenotypes during development as a biofilm. *J. Bacteriol.* **184**, 1140–1154 (2002).

4.   Southey-Pillig, C. J., Davies, D. G. & Sauer, K. Characterization of temporal protein production in Pseudomonas aeruginosa biofilms. *J. Bacteriol.* **187**, 8114–8126 (2005).

5.   Güvener, Z. T. & Harwood, C. S. Subcellular location characteristics of the Pseudomonas aeruginosa GGDEF protein, WspR, indicate that it produces cyclic-di-GMP in response to growth on surfaces. *Mol. Microbiol.* **66**, 1459 (2007).

6.   Vallet, I., Olson, J. W., Lory, S., Lazdunski, A. E. & Filloux, A. The chaperone/usher pathways of Pseudomonas aeruginosa: Identification of fimbrial gene clusters (cup) and their involvement in biofilm formation. *PNAS* **98**, 6911–6916 (2001).

7.   Patino, R. *et al.* Antagonistic response regulators spatially regulate receptor methylation in the Pseudomonas aeruginosa Pil-Chp surface sensing system. *Cell Rep.* **44**, 115536 (2025).

8.   McDougald, D., Rice, S. A., Barraud, N., Steinberg, P. D. & Kjelleberg, S. Should we stay or should we go: Mechanisms and ecological consequences for biofilm dispersal. *Nature Reviews Microbiology* vol. 10 Preprint at https://doi.org/10.1038/nrmicro2695 (2012).

9.   Rumbaugh, K. P. & Sauer, K. Biofilm dispersion. *Nat. Rev. Microbiol.* **18**, 571–586 (2020).

10. Cherny, K. E. & Sauer, K. Pseudomonas aeruginosa Requires the DNA-Specific Endonuclease EndA To Degrade Extracellular Genomic DNA To Disperse from the Biofilm. *J. Bacteriol.* **201**, e00059-19 (2019).

11. Cherny, K. E. & Sauer, K. Untethering and degradation of the polysaccharide matrix are essential steps in the dispersion response of Pseudomonas aeruginosa biofilms. *J. Bacteriol.* **202**, (2020).

12. Randall, T. E. *et al.* Sensory Perception in Bacterial Cyclic Diguanylate Signal Transduction. *J. Bacteriol.* **204**, (2022).

13. Hengge, R. Trigger phosphodiesterases as a novel class of c-di-GMP effector proteins. *Philosophical Transactions of the Royal Society B: Biological Sciences* vol. 371 Preprint at https://doi.org/10.1098/rstb.2015.0498 (2016).

14. Hengge, R. High-specificity local and global c-di-GMP signaling. *Trends in Microbiology* vol. 29 993–1003 Preprint at https://doi.org/10.1016/j.tim.2021.02.003 (2021).

15. Povolotsky, T. L. & Hengge, R. 'Life-style' control networks in Escherichia coli: Signaling by the second messenger c-di-GMP. *J. Biotechnol.* **160**, 10–16 (2012).

16. Kalia, M. & Sauer, K. Distinct transcriptome and traits of freshly dispersed Pseudomonas aeruginosa cells. *mSphere* **9**, (2024).

17. Chua, S. L. *et al.* Dispersed cells represent a distinct stage in the transition from bacterial biofilm to planktonic lifestyles. *Nature Communications 2014 5:1* **5**, 1–12 (2014).

18. Zhu, X., Rice, S. A. & Barraud, N. Nitric Oxide and Iron Signaling Cues Have Opposing Effects on Biofilm Development in Pseudomonas aeruginosa. *Appl. Environ. Microbiol.* **85**, (2019).

19. Bertran i Forga, X., Hong, Y., Fairfull-Smith, K. E., Qin, J. & Totsika, M. Nitric oxide donor
sodium nitroprusside serves as a source of iron supporting *Pseudomonas aeruginosa* growth
and biofilm formation. *Microbiol. Spectr.* https://doi.org/10.1128/SPECTRUM.02234-25
(2025) doi:10.1128/SPECTRUM.02234-25.

20. An, S., Wu, J. & Zhang, L. H. Modulation of pseudomonas aeruginosa biofilm dispersal by a
cyclic-di-gmp phosphodiesterase with a putative hypoxia-sensing domain. *Appl. Environ.*
Microbiol. **76**, 8160–8173 (2010).

21. Hsu, J. L., Chen, H. C., Peng, H. L. & Chang, H. Y. Characterization of the Histidine-
containing Phosphotransfer Protein B-mediated Multistep Phosphorelay System in
Pseudomonas aeruginosa PAO1. *Journal of Biological Chemistry* **283**, 9933–9944 (2008).

22. Metsalu, T. & Vilo, J. ClustVis: a web tool for visualizing clustering of multivariate data using
Principal Component Analysis and heatmap. *Nucleic Acids Res.* **43**, W566–W570 (2015).

23. Jennings, L. K. *et al.* Pel is a cationic exopolysaccharide that cross-links extracellular DNA in
the Pseudomonas aeruginosa biofilm matrix. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 11353–11358
(2015).

24. Barraud, N. *et al.* Involvement of Nitric Oxide in Biofilm Dispersal of Pseudomonas
aeruginosa. *J. Bacteriol.* **188**, 7344–7353 (2006).

25. Bhattacharjee, A., Nusca, T. D. & Hochbaum, A. I. Rhamnolipids mediate an interspecies
biofilm dispersal signaling pathway. *ACS Chem. Biol.* **11**, 3068–3076 (2016).

26. Schooling, S. R., Charaf, U. K., Allison, D. G. & Gilbert, P. A role for rhamnolipid in biofilm
dispersion. *Biofilms* **1**, 91–99 (2004).

27. Davey, M. E., Caiazza, N. C. & O'Toole, G. A. Rhamnolipid surfactant production affects
biofilm architecture in Pseudomonas aeruginosa PAO1. *J. Bacteriol.* **185**, 1027–1036 (2003).

28. Glick, R. *et al.* Increase in rhamnolipid synthesis under iron-limiting conditions influences
surface motility and biofilm formation in Pseudomonas aeruginosa. *J. Bacteriol.* **192**, 2973–
2980 (2010).

29. Kalia, M., Amari, D., Davies, D. G. & Sauer, K. cis-DA-dependent dispersion by
Pseudomonas aeruginosa biofilm and identification of cis-DA-sensory protein DspS. *mBio* **14**,
(2023).

30. Chambers, J. R., Cherny, K. E. & Sauer, K. Susceptibility of Pseudomonas aeruginosa
Dispersed Cells to Antimicrobial Agents Is Dependent on the Dispersion Cue and Class of the
Antimicrobial Agent Used. *Antimicrob. Agents Chemother.* **61**, e00846-17 (2017).

31. Eilers, K. *et al.* The dual GGDEF/EAL domain enzyme PA0285 is a Pseudomonas species
housekeeping phosphodiesterase regulating early attachment and biofilm architecture. *Journal*
of Biological Chemistry **300**, (2024).

32. Eilers, K. *et al.* Phenotypic and integrated analysis of a comprehensive Pseudomonas
aeruginosa PAO1 library of mutants lacking cyclic-di-GMP-related genes. *Front. Microbiol.*
13, 949597 (2022).

33. Wei, Q. *et al.* Diguanylate Cyclases and Phosphodiesterases Required for Basal-Level c-di-
GMP in Pseudomonas aeruginosa as Revealed by Systematic Phylogenetic and Transcriptomic
Analyses. *Appl. Environ. Microbiol.* **85**, (2019).

34. Cai, Y. M. *et al.* The c-di-GMP Phosphodiesterase PipA (PA0285) Regulates Autoaggregation
and Pf4 Bacteriophage Production in Pseudomonas aeruginosa PAO1. *Appl. Environ.*
Microbiol. **88**, (2022).

35. Cai, Y. ming *et al.* Differential impact on motility and biofilm dispersal of closely related
phosphodiesterases in Pseudomonas aeruginosa. *Sci. Rep.* **10**, (2020).

36. Liu, S. *et al.* Priority changes between biofilm exopolysaccharides synthesis and rhamnolipids
production are mediated by a c-di-GMP-specific phosphodiesterase NbdA in Pseudomonas
aeruginosa. *iScience* **25**, 105531 (2022).

37. Li, Y., Heine, S., Entian, M., Sauer, K. & Frankenberg-Dinkel, N. NO-induced biofilm
dispersion in Pseudomonas aeruginosa is mediated by an MHYT domain-coupled
phosphodiesterase. *J. Bacteriol.* **195**, 3531–3542 (2013).

38. Zhan, X. *et al.* A c-di-GMP signaling module controls responses to iron in Pseudomonas
aeruginosa. *Nature Communications 2024 15:1* **15**, 1–15 (2024).

39. Stelitano, V. *et al.* C-di-GMP hydrolysis by Pseudomonas aeruginosa HD-GYP
phosphodiesterases: analysis of the reaction mechanism and novel roles for pGpG. *PLoS One*
8, (2013).

40. Ryan, R. P. *et al.* HD-GYP domain proteins regulate biofilm formation and virulence in
Pseudomonas aeruginosa. *Environ. Microbiol.* **11**, 1126–1136 (2009).

41. Matilla, M. A., Velando, F., Monteagudo-Cascales, E. & Krell, T. Flagella, Chemotaxis and
Surface Sensing. in *Advances in Experimental Medicine and Biology* (eds. Filloux, A. &
Ramos, J. L.) vol. 1386 185–221 (Advances in Experimental Medicine and Biology, Springer,
Cham, 2022).

42. Tchigvintsev, A. *et al.* Biochemical and structural studies of uncharacterized protein PA0743
from Pseudomonas aeruginosa revealed NAD +-dependent L-Serine dehydrogenase. *Journal*
of Biological Chemistry **287**, 1874–1883 (2012).

43. Gliese, N. & Khodaverdi, V. AgmR controls transcription of a regulon with several operons
essential for ethanol oxidation in Pseudomonas aeruginosa ATCC 17933.
https://doi.org/10.1099/mic.0.26882-0 doi:10.1099/mic.0.26882-0.

44. Tassinari, M., Rudzite, M., Filloux, A. & Low, H. H. Assembly mechanism of a Tad secretion
system secretin-pilotin complex. *Nat. Commun.* **14**, 1–13 (2023).

45. Kalia, M., Resch, M. D., Cherny, K. E. & Sauer, K. The Alginate and Motility Regulator AmrZ
is Essential for the Regulation of the Dispersion Response by Pseudomonas aeruginosa
Biofilms. *mSphere* https://doi.org/10.1128/MSPHERE.00505-22 (2022)
doi:10.1128/MSPHERE.00505-22.

46. Zhao, J. *et al.* Molecular Mechanism of CdpR Involved in Quorum-Sensing and Bacterial
Virulence in Pseudomonas aeruginosa. *PLoS Biol* **14**, 1002449 (2016).

47. Zhao, J. *et al.* Structural and Molecular Mechanism of CdpR Involved in Quorum-Sensing and
Bacterial Virulence in Pseudomonas aeruginosa. *PLoS Biol.* **14**, (2016).

48. Park, S. & Sauer, K. Controlling Biofilm Development Through Cyclic di-GMP Signaling. in
Advances in experimental medicine and biology (eds. Filloux, A. & Ramos, J. L.) vol. 1386
69–94 (Advances in Experimental Medicine and Biology, Springer, Cham, 2022).

49. Webster, S. S. *et al.* Force-Induced Changes of PilY1 Drive Surface Sensing by Pseudomonas
aeruginosa. *mBio* **13**, (2021).

50. Kazmierczak, B. I., Lebron, M. B. & Murray, T. S. Analysis of FimX, a phosphodiesterase that
governs twitching motility in Pseudomonas aeruginosa. *Mol. Microbiol.* **60**, 1026–1043
(2006).

51. Klausen, M. *et al.* Biofilm formation by Pseudomonas aeruginosa wild type, flagella and type
IV pili mutants. *Mol. Microbiol.* **48**, 1511–1524 (2003).

52. Cowles, K. N. & Gitai, Z. Surface association and the MreB cytoskeleton regulate pilus
production, localization and function in Pseudomonas aeruginosa. *Mol. Microbiol.* **76**, 1411–
1426 (2010).

53. Irie, Y. *et al.* Self-produced exopolysaccharide is a signal that stimulates biofilm formation in
Pseudomonas aeruginosa. *Proceedings of the National Academy of Sciences* **109**, 20632–
20636 (2012).

54. Hengge, R. Principles of c-di-GMP signalling in bacteria. *Nature Reviews Microbiology* vol. 7
Preprint at https://doi.org/10.1038/nrmicro2109 (2009).

55. Petrova, O. E. & Sauer, K. PAS domain residues and prosthetic group involved in BdlA-
dependent dispersion response by Pseudomonas aeruginosa biofilms. *J. Bacteriol.* **194**, 5817–
5828 (2012).

56. Morgan, R., Kohn, S., Hwang, S. H., Hassett, D. J. & Sauer, K. BdlA, a Chemotaxis Regulator
Essential for Biofilm Dispersion in Pseudomonas aeruginosa. *J. Bacteriol.* **188**, 7335 (2006).

57. Stelitano, V. *et al.* C-di-GMP Hydrolysis by Pseudomonas aeruginosa HD-GYP
Phosphodiesterases: Analysis of the Reaction Mechanism and Novel Roles for pGpG. *PLoS*
One **8**, e74920 (2013).

58. Ryan, R. P. *et al.* HD-GYP domain proteins regulate biofilm formation and virulence in
Pseudomonas aeruginosa. *Environ. Microbiol.* **11**, 1126–1136 (2009).

59. Wood, T. L. *et al.* Rhamnolipids from Pseudomonas aeruginosa disperse the biofilms of
sulfate-reducing bacteria. *NPJ Biofilms Microbiomes* **4**, (2018).

60. Hou, L., Debru, A., Chen, Q., Bao, Q. & Li, K. AmrZ Regulates Swarming Motility Through
Cyclic di-GMP-Dependent Motility Inhibition and Controlling Pel Polysaccharide Production
in Pseudomonas aeruginosa PA14. *Front. Microbiol.* **10**, (2019).

61. Giraud, C. *et al.* The PprA-PprB two-component system activates CupE, the first non-
archetypal Pseudomonas aeruginosa chaperone-usher pathway system assembling fimbriae.
Environ. Microbiol. **13**, 666–683 (2011).

62. Bernard, C. S., Bordi, C., Termine, E., Filloux, A. & De Bentzmann, S. Organization and
PprB-dependent control of the pseudomonas aeruginosa tad locus, involved in Flp Pilus
Biology. *J. Bacteriol.* **191**, 1961–1973 (2009).

63. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature Methods*
2012 9:4 **9**, 357–359 (2012).

64. Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using Real-Time
Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25**, 402–408 (2001).

65. Coppens, L. & Lavigne, R. SAPPHIRE: a neural network based classifier for σ70 promoter
prediction in Pseudomonas. *BMC Bioinformatics 2020 21:1* **21**, 415- (2020).

66. Miller, J. H. Assay of B-galactosidase. in *Experiments in molecular genetics* 352–355 (Cold
Spring Harbor Laboratory, Cold Spring, N.Y., 1972).

67. HOLLOWAY, B. W. Genetic recombination in Pseudomonas aeruginosa. *J. Gen. Microbiol.*
13, 572–581 (1955).

68. Qin, J., Hong, Y., Morona, R. & Totsika, M. O antigen biogenesis sensitises Escherichia coli
K-12 to bile salts, providing a plausible explanation for its evolutionary loss. *PLoS Genet.* **19**,
e1010996 (2023).

69. Elzer, P. H. *et al.* In Vivo and in Vitro Stability of the Broad-Host-Range Cloning Vector
pBBR1MCS in Six Brucella Species. *Plasmid* **33**, 51–57 (1995).

696