1   Reinke crystals are immunoreactive for purine-synthesizing metabolic enzymes

3   John Woulfe MD PhD: Department of Pathology and Laboratory Medicine, University of
4   Ottawa and Ottawa Hospital Research Institute, Ottawa, Ontario, Canada. jwoulfe@toh.ca

5   Trevor Flood MD: Department of Pathology and Laboratory Medicine, University of Ottawa,
6   Ottawa, Ontario, Canada. tflood@toh.ca

7   Sharlene Faulkes PhD: Department of Pathology and Laboratory Medicine, University of
8   Ottawa, Ottawa, Ontario, Canada. sfaulkes@uottawa.ca

9   David G. Munoz MD: Department of Laboratory Medicine, St. Michael's Hospital, Unity Health
10   & Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto,
11   Ontario, Canada. David.munoz@unityhealth.to

12
13   Pages: 18
14   Tables: 0
15   Figures: 2
16   Running page title: Metabolic enzymes in Reinke crystals
17

Corresponding author:
John Woulfe
Dept. of Pathology, The Ottawa Hospital, Civic Campus
1053 Carling Avenue, Ottawa, Ontario, Canada
K1Y 4E9
Phone: 613-798-555 ext. 13345
Fax: 613-761-4199
E-mail: jwoulfe@toh.ca
ORCID ID: 0-0002-0007382-6343

Abstract

Reinke crystals are a defining histological feature of human adult Leydig cells, the testosterone

producing cells of the testis. These structures are present in the cytoplasm and the nucleus and

display quantitative alterations in a variety of physiological and pathological contexts. The

functional significance and protein composition of Reinke crystals have remained elusive for

over a century. Here, we demonstrate that Reinke crystals are intensely immunoreactive for

inosine monophosphate dehydrogenase (IMPDH), and phosphoribosyl pyrophosphate synthetase

(PRPS), two key rate-limiting enzymes in the *de novo* synthesis of purine nucleotides. IMPDH

and PRPS are two of several metabolic enzymes that are capable of forming mesoscale

filamentous aggregates as a mechanism to regulate enzyme activity. IMPDH is also able to form

crystals *in cellulo*. Our observations link Reinke crystal formation to purine nucleotide

metabolism in Leydig cells. We discuss how this novel finding may relate to the unique

dependence of Leydig cells on guanyl-based purine nucleotides for testosterone synthesis. The

results of this study may have important implications for understanding metabolic contributions

to male reproductive disorders as well as offering a novel diagnostic and theranostic tool

applicable to Leydig cell neoplasms.

Introduction

Leydig cells, the testosterone-producing cells of the testis, are essential for male reproductive

health [1]. Reinke crystals, first described by Frederich B. Reinke in 1896 [2], are a defining

cytological feature of both native and neoplastic Leydig cells (for a review, see [3]). The presence

of Reinke crystals in neoplastic Leydig cells is of diagnostic utility. They are well-

circumscribed, hyaline eosinophilic structures using conventional hematoxylin and eosin staining

and are also demonstrable in Masson's trichrome, Giemsa, and Gram preparations [4-7]. Reinke

crystals are 2-3 μm in size on average but can measure up to 20 μm [8]. Ultrastructurally, their

morphology is consistent with true crystals, showing "long-range order" in which 5 nm parallel

filaments are arranged in a regular array throughout the entire structure. The protein composition

and functional significance of Reinke crystals have remained enigmatic. Their abundance is

dependent on Leydig cell developmental stage and is altered in a variety of testicular disorders

(see [3]). To date, candidate proteins that have been implicated as Reinke crystal constituents

include nestin [9] and a 3β hydroxysteroid/calretinin complex [7,10]. However, definitive

confirmation of these is lacking [3].

Inosine monophosphate dehydrogenase (IMPDH) catalyzes the first committed step in

the *de novo* synthesis of guanine-based (guanyl) purine nucleotides guanosine mono-, di-, and

triphosphate (GMP, GDP and GTP). These, and especially GTP, are essential building blocks of

RNA and DNA, are used as signalling molecules in several molecular pathways and are

incorporated into co-enzymes. There are two isoforms of IMPDH that share 83% amino acid

identity and have differential expression patterns [11]. IMPDH1 is expressed constitutively in

many tissues whereas IMPDH2 expression is also ubiquitous but is upregulated in proliferating

tissues [12-15]. IMPDH is one of a growing list of metabolic enzymes capable of regulated assembly

to form long filaments as a mechanism to adjust enzyme availability and activity [16-18]. Each

filament consists of linearly stacked homo-octamers. Filament formation serves to resist

feedback inhibition, thereby sustaining guanyl nucleotide synthesis under conditions of increased

demand [19]. Individual enzyme filaments coalesce to form microscopically visible cytoplasmic or

intranuclear filament bundles called "cytoophidia" (Gk. For "cellular snakes") [18]. Cytoophidia

formation correlates with increased enzyme activity and thus increased guanyl nucleotide

synthesis. Accordingly, microscopically-detectable IMPDH filament assemblies serve as a

marker for guanyl nucleotide-dependent cells [16,17,20]. Importantly, IMPDH can also form large

intracellular crystals [21]. This spectrum suggests a temporally sequential series of assembly states

in which IMPDH crystallization represents the ultimate stage in GTP-dependent cells.

Intriguingly, *in cellulo* IMPDH crystals share morphological features with Reinke crystals. We

have provided evidence that IMPDH crystallization occurs *in vivo*. Specifically, we described

IMPDH filaments as well as crystals/paracrystals within the nuclei of neurons in the human brain

[22,23]. Interestingly, another metabolic enzyme, phosphoribosyl pyrophosphate synthase (PRPS)

which catalyzes one of the first steps in *de novo* purine nucleotide synthesis [23-26] co-localizes

with IMPDH in these neuronal structures [23,27]. This is consistent with studies demonstrating

PRPS filamentation/cytoophidia formation in cells, including neurons [24].

Relative to other endocrine cells, Leydig cells are uniquely dependent on GTP for

testosterone synthesis [28]. The latter is initiated in response to luteinizing hormone (LH) via a G-

protein-coupled receptor (GPCR). Interestingly, GPCR signaling has been linked to purine

metabolism by regulating the assembly of a purine-synthesizing multienzyme complex

(metabolon) known as the "purinosome" [29,30]. LH-receptor binding on Leydig cells results in

recruitment of GTP to replace GDP on the G-protein Gsα. GTP-bound Gs activates its effector,

adenylate cyclase resulting in the generation of cAMP. The latter stimulates cholesterol

translocation to mitochondria where it is metabolized to pregnenolone which in turn is converted

to testosterone by enzymes in the endoplasmic reticulum [1]. Whereas all endocrine cells utilize

GTP signaling to some extent, adult Leydig cells lack redundancy in this respect and are

exclusively dependent on the LH-GTP-cAMP signaling axis [28]. In contrast, fetal Leydig cells as

well as testosterone-producing adrenal cells, (both of which notably lack Reinke crystals), can

exploit alternate, non-GTP-dependent mechanisms [28,31]. Moreover, LH receptor signaling in

Leydig cells is characterized by maximal testosterone production even when only a tiny fraction

(typically <1%) of available LH receptors are occupied by a ligand [32-34]. This is critical for male

reproductive health because it allows the testes to maintain stable testosterone levels even under

low of fluctuating circulating LH [35,36]. Mechanistically, this is attributable to massive signal

amplification. There is evidence that this, in turn, is at least partly related to a stoichiometric

imbalance in the ratio of G-proteins to LH receptors whereby a single occupied receptor can

activate a large number of G-proteins, resulting in a 50-fold magnification of the LH binding

reaction [37]. This contrasts with catecholamine receptor signalling in which a closer numerical

relationship between hormone receptors and G-proteins exists. It seems reasonable to surmise

that this relative excess of G-proteins in Leydig cells would require a correspondingly robust

supply of GTP. In this context, Leydig cells are functionally constrained by the availability of

GTP, rendering them uniquely vulnerable to disruptions in purine nucleotide homeostasis.

Recognizing 1) the dependence of Leydig cells on guanyl nucleotide metabolism for their

function, 2) evidence that intracellular IMPDH macromolecular assemblies characterize GTP-

dependent cells and 3) the shared morphology of Reinke crystals, *in cellulo* IMPDH crystals, and

the IMPDH/PRPS structures we have described in neurons, we were compelled to investigate

whether Reinke crystals are immunoreactive for IMPDH and/or PRPS.

Materials and Methods

The study was conducted in accordance with the ethical standards of The Ottawa

Hospital. Tissue blocks of formalin-fixed, paraffin-embedded tissue from four cases of testicular

Leydig cell tumours with adjacent non-neoplastic testis were obtained from the tissue archive in

the Division of Anatomical Pathology of the Ottawa Hospital. These were resected from four

male participants aged 28, 29, 70, and 71. Diagnoses in all cases were well-circumscribed

Leydig cell tumours confined to the testicular parenchyma. The tumours measured from 1.3 to

1.6 cm. A representative block from each case was sectioned at a thickness of 5 μm, and

sections were mounted onto coated slides. Immunohistochemical staining was performed on

tissue sections using the Leica Bond™ system. Sections were deparaffinized and pre-treated

using heat mediated antigen retrieval with an EDTA buffer (pH 9) for 20 minutes. The sections

were then incubated in one of the following primary antibodies: rabbit polyclonal anti-IMPDH1

(1:400; Proteintech #22092-1 -AP, Lot #00018215; Rosemont IL USA), rabbit polyclonal anti-

PRPS1 (1:200; ProteinTech #15549-1-AP), or mouse monoclonal anti-PRPS1/2/3 (1:200; Santa

Cruz clone A11, sc-376440) for 30 minutes at room temperature and detected using an HRP

conjugated compact polymer system. Slides were then stained using DAB as the chromogen,

counterstained with hematoxylin, mounted and cover-slipped. As an experimental control for

antibody specificity, the polyclonal primary antibodies were pre-absorbed using a 20-fold molar

excess of the immunizing peptides (ProteinTech #Ag17473 for IMPDH and #Ag7907 for PRPS),

and incubated overnight at 4 degrees with gentle agitation which reduced staining. For method

specificity, omission of the primary antibodies resulted in a complete absence of staining.

Double-immunofluorescence staining was performed by incubating sections overnight at 4

degrees in a cocktail containing the polyclonal IMPDH (1:100) and monoclonal PRPS (1:200)

antibodies. Sections were washed with 1XTBST and then incubated with goat anti-mouse 594

(#A11005, Invitrogen) and donkey anti-rabbit 488 (#A21206) secondary antibodies for 2 h in the

dark at room temperature. This was followed by incubation with a quencher (Vector TrueView

Autofluorescence Quenching Kit #SP-8400, Vector Labs) to decrease autofluorescence. Sections

were then washed, incubated with 5 ug/ml of DAPI (ThermoScientific #62248) and

coverslipped. Microscopic imaging of was performed using Nikon Eclipse 80i light microscope.

Photomicrographs were acquired using a Nikon Digital Sight Fi-1 camera and processed using

NIS Elements imaging software version 4.60. For immunofluorescence, image acquisition was

performed using a Zeiss 880 Confocal Microscope equipped with a Plan-Apo 63×1.4 numerical

aperture. Processing of Airyscan 32-channel raw data was conducted using Zen Black 2.3 SP1

software.

Results

In all 4 cases, cytoplasmic Reinke crystals were visualized in hematoxylin- and eosin (H

and E)-stained sections in both tumour cells and native Leydig cells (Figure 1). Their frequency

varied among cases. Reinke crystals in the cytoplasm appeared as well-circumscribed

eosinophilic structures measuring from approximately 2 to 30 μm (Figure 1A). They were often

rod-shaped or rectangular although various more irregular morphologies were common. Larger

Reinke crystals, particularly the rod-shaped structures, appeared to "stretch" the plasma

membrane of the host cell. In this limited series, we did not detect consistent differences in the

size or morphology of cytoplasmic Reinke crystals between neoplastic and non-neoplastic

Leydig cells in H and E-stained sections.

Reinke crystals were also visualized within the nuclei of neoplastic and non-neoplastic

Leydig cells in H and E-stained sections (Figures 1A and B; arrowheads). The majority of these

consisted of an amorphous eosinophilic area devoid of hematoxylin staining. In rare cases,

nuclear crystals with a more solid rod-shaped morphology could be identified. Some of these

were longer than the nuclear diameter and distorted the nuclear membrane (Figure 1B, arrow).

Across all four cases, both cytoplasmic and nuclear Reinke crystals displayed intense

immunoreactivity for IMPDH and PRPS, concordant with the crystalline profiles seen on H&E

(Figure 1B and C). All antibodies produced a similar intensity and distribution of

immunostaining of cytoplasmic and nuclear Reinke crystals.   In both Leydig cell tumours

(Figure 1A-E) and native Leydig cell nests in the testicular interstitium (Figure 1F), Reinke

crystals were far more evident and abundant in IMPDH and PRPS-immunostained sections

relative to those stained using H and E (compare Figure 1A with C and D). Whereas less than

half of the Leydig cell population appeared to have nuclear Reinke crystals in the latter, IMPDH

or PRPS immunostaining revealed nuclear Reinke crystals in approximately 90% of the

neoplastic Leydig cell population (Figure 1E). Moreover, immunostaining revealed features that

were not appreciated on H and E staining. For example, whereas cytoplasmic crystals were

intensely stained and showed well-defined edges, many nuclear crystals showed "blush" staining

and ill-defined margins (Figure 1E, arrowheads). As shown in Figure 2, double-staining

immunofluorescence confirmed that all cytoplasmic and nuclear Reinke crystals were

immunoreactive for both IMPDH and PRPS.

Discussion

Despite having been described over a century ago [2], the protein composition of Reinke

crystals has remained a mystery. Here, we demonstrate intense immunoreactivity of Reinke

crystals for IMPDH and PRPS. These antibodies were generated using synthetic peptide

immunogens corresponding to distinct but overlapping epitopes in each of the enzymes. Our

study cannot distinguish the various IMPDH and PRPS isoforms. The target sequence of the

polyclonal anti-IMPDH1 antibodies (amino acids 170-235) shares 77% sequence identity with

the homologous domain of IMPDH2. Similarly, there are 3 highly homologous isoforms of

PRPS (PRPS1, 2, and 3) in humans. The target epitope of the polyclonal antibody used in the

present study (amino acids 1-318 of PRPS1) shares 98.7% and 98.1% sequence identity with

PRPS2 and PRPS3, respectively whereas that of the monoclonal antibody (amino acids 143-270

of PRPS2) shares 98.7% and 96.5% sequence identity with PRPS1 and PRPS3, respectively.

Interestingly, Keppeke and co-workers demonstrated co-localization of both IMPDH isoforms in

cytoophidia, including intranuclear structures, in mammalian cells [16]. Thus, it is possible that

Reinke crystals contain multiple IMPDH and PRPS variants.

Among the novel findings of this study, we show that, like IMPDH, PRPS filaments are

capable of incorporating into a crystalline array along with IMPDH. The co-assembly of distinct

metabolic enzymes to form heterotypic assemblies is an emerging theme in the field of enzyme

filamentation [24,38]. Interestingly, co-assembling enzymes tend to be those located at critical

"branch-points" in the purine biosynthetic pathway, including PRPS and IMPDH, suggesting that

co-assembly may have an important regulatory role [24]. In support of the possibility that these

two enzymes represent the core component(s) of the filaments in the crystal lattice, in contrast to

other proteins that have been nominated as Reinke crystal constituents, the ability of both

IMPDH and PRPS to assemble into macrostructures, including, for IMPDH, crystals with a light

microscopic and ultrastructural morphology similar to that described for Reinke crystals has been

convincingly demonstrated. However, further studies, such as super-resolution microscopy,

immunostaining studies at the electron microscopic level, and even cryoelectron microscopy are

required to confirm this as well as to elucidate the spatial relationships between IMPDH and

PRPS filaments within the crystals. [21,39].

By linking Reinke crystals to purine nucleotide metabolism in Leydig cells, our

findings contextualize what was once merely a morphological curiosity within a coherent and

physiologically relevant framework. As discussed above, Leydig cells are uniquely dependent

on a sufficient supply of GTP. This is not only because they lack recourse to alternative non-

guanyl-dependent signalling pathways [28], but also to satisfy the requirements of the signal

amplification provided by the relative multitude of G-proteins which stoichiometrically

outnumber their cognate LH receptors [37]. Moreover, because Leydig cells respond to even small

fluctuations in LH, GTP must be made available with high temporal fidelity. To meet these

demands, we postulate that Reinke crystals provide readily accessible storage depots for IMPDH

and PRPS to be rapidly deployed in response to sudden changes in GTP demand. This is

analogous to the storage function ascribed to insulin crystals in the exocytotic vesicles of

pancreatic beta cells and major basic protein 1 crystals in eosinophils [40-42]. Consistent with this

hypothesis, in conditions of higher LH levels and Leydig cell hyperplasia, including 17-βHSD

deficiency and androgen insensitivity syndrome, Reinke crystals are reduced or not present [3,43,44].

We propose that this reflects translocation of these key enzymes from the crystal to satisfy the

requirements imposed by upregulated LH signaling in these scenarios. On the other hand, it is

conceivable that Reinke crystals themselves provide a scaffold for guanyl-nucleotide synthesis in

situ. This model would be predicated on the following: 1) that IMPDH and PRPS are

enzymatically active in the crystalline context and 2) that the requisite enzymes catalyzing the

upstream and downstream steps in the pathway are spatially proximal to the crystal. Although

individual filaments contain enzymatically active oligomers, the status of IMPDH and PRPS

functionality in the crystalline context is unknown. Future studies should explore the nucleotide-

synthesizing capacity of Reinke crystals as well as their spatial relationship(s) with other purine-

synthesizing metabolons, most notably, purinosomes [30]. These hypothetical models are based on

the premise that Reinke crystals are somehow involved in supplying sufficient GTP to meet the

demands of Leydig cell signalling. The possibility that they are, alternatively or in addition,

linked to some other unique, purine-related aspect of Leydig cell metabolism cannot be

excluded.

Reinke crystals in the nucleus were also IMPDH and PRPS immunoreactive. Relative to

H and E, immunohistochemical detection of these antigens revealed a much larger proportion of

both neoplastic and native Leydig cells bearing nuclear inclusions. Nuclear Reinke crystals

appeared to be less intensely stained and were more morphologically amorphous than the "rigid"-

appearing cytoplasmic crystals. This may relate to observations suggesting that nuclear Reinke

crystals may be at an earlier, precursor stage of formation, corresponding to paracrystalline

structures [4,45]. Whether these represent a nuclear "overflow" storage compartment of excessive

cytoplasmic IMPDH/PRPS, or are functional is uncertain. In neurons, IMPDH/PRPS crystals

show consistent localization adjacent to the nucleolus [22]. The nucleolus consumes vast quantities

of GTP for the synthesis of ribosomal RNA [46,47]. It is tempting to speculate that juxtanucleolar

crystals may provide a proximate "hyperlocal" reservoir of IMPDH/PRPS to be deployed for the

synthesis of GTP. Alternatively, or in addition, IMPDH has been described in the cell nucleus

where it has been ascribed a non-enzymatic "moonlighting" role [48]. Specifically, nuclear

IMPDH binds to DNA and functions as a transcription factor, regulating E2F-dependent gene

expression in a cell-cycle dependent manner. In this context, Leydig cells might exploit the

capacity of IMPDH to form crystals as a mechanism to dynamically regulate its nucleoplasmic

availabilities to perform this moonlighting function.

The observation that children with Lesch-Nyhan syndrome, the prototypical disorder of

purine metabolism, display impaired testicular development and testosterone deficiency [49]

reinforces the link between Leydig cell function and purine metabolism in a pathophysiological

context with potential implications for understanding the pathogenesis of male reproductive

disorders. There may also be implications for cancer diagnosis and treatment. Leydig cell

tumours are a rare form of testicular cancer. Most of these are benign although malignant

variants can occur with metastatic potential. The identification of Reinke crystals in cytological

preparations or tissue sections is useful in diagnosing these tumours [6]. However, many tumours

have very few crystals. In addition, Reinke crystals are sensitive to formalin fixation,

contributing to false negative interpretation in H and E-stained sections [7]. Immunohistochemistry

for the detection of IMPDH or PRPS provides a novel, more sensitive histological marker for

Reinke crystal identification in the diagnostic setting.   With respect to therapeutic implications,

several studies suggest that, under steady-state conditions, IMPDH filamentation is a hallmark of

cells in which guanine nucleotide pools are limiting. This may represent a metabolic

vulnerability that could be leveraged by IMPDH inhibitors to further deplete GTP, culminating

in DNA replication stress and cell death. An important corollary of this is that the

immunohistochemical detection of Reinke crystals in Leydig cell tumours may provide a

histological biomarker predictive of a favourable therapeutic response to IMPDH inhibitors.

Currently approved IMPDH inhibitors like mycophenolic acid as well as newer generation

agents are receiving renewed attention as anti-neoplastic agents [50]. Future preclinical studies and

clinical trials should assess the susceptibility of Reinke-crystal bearing Leydig cell malignancies

to IMPDH inhibition.

Conclusions

Reinke crystals are immunoreactive for IMPDH and PRPS, linking these structures to

purine nucleotide metabolism and possibly a role in regulating guanyl-nucleotide availability.

In addition to solving a long-standing mystery, this finding has important implications for

understanding the role of purine nucleotide metabolism in testicular function and disease,

including cancer. Moreover, our previous demonstration of similar structures in neurons

suggests that IMPDH/PRPS crystal formation may represent a paradigmatic organizational motif

for understanding the regulation of purine nucleotide metabolism across diverse tissue types.

Acknowledgements

The authors wish to acknowledge the technical support provided by the University of

Ottawa Cell Biology Imaging and Acquisition Core (CBIA; RRID:SCR_021845) and the Louise

Pelletier Histology Core Facility (RRID: SCR_021737), Department of Pathology and

Laboratory Medicine, University of Ottawa. The authors specifically wish to acknowledge the

excellent technical support provided by Liyuan Wang of the CBIA. No funding was received for

conducting this study.

Disclosures

The authors have no conflict of interest to disclose.

Authors' Contributions

J.W. designed the project, performed experiments, analyzed the data, and wrote the

manuscript; T.F. provided tissue for the project, provided guidance regarding significance with

respect to Leydig cell function and edited the manuscript; S.F. performed the

immunohistochemical experiments and edited the manuscript; D.G.M. edited and provided

critical input regarding organization of the manuscript. J.W. is the guarantor of this work and, as

such, had full access to all of the data in the study and takes responsibility for the integrity of the

data and the accuracy of the data analysis.

Data Accessibility

All data pertaining to this study will be made readily available upon reasonable request.

References
313
1.   Zirkin BR, Papadopoulos V. Leydig cells: formation, function, and regulation. *Biol*
Reprod. Jul 1 2018;99(1):101-111. doi:10.1093/biolre/ioy059
2.   F R. Beitrage zur histologie des menschen. *Archiv fur mikroskopische Anatomie*.
1896;47:34-44.
3.   Planinic A, Maric T, Bojanac AK, Jezek D. Reinke crystals: Hallmarks of adult Leydig
cells in humans. *Andrology*. Sep 2022;10(6):1107-1120. doi:10.1111/andr.13201
4.   Gupta SK, Francis IM, Sheikh ZA, al-Rubah NA, Das DK. Intranuclear Reinke's crystals
in a testicular Leydig cell tumor diagnosed by aspiration cytology. A case report. *Acta Cytol*.
Mar-Apr 1994;38(2):252-6.
5.   Jain M, Aiyer HM, Bajaj P, Dhar S. Intracytoplasmic and intranuclear Reinke's crystals
in a testicular Leydig-cell tumor diagnosed by fine-needle aspiration cytology: a case report with
review of the literature. *Diagnostic cytopathology*. Sep 2001;25(3):162-4. doi:10.1002/dc.2029
6.   Handa U, Sood T, Punia RS. Testicular Leydig cell tumor diagnosed on fine needle
aspiration. *Diagnostic cytopathology*. Sep 2010;38(9):682-4. doi:10.1002/dc.21306
7.   Mesa H, Gilles S, Smith S, Dachel S, Larson W, Manivel JC. The mystery of the
vanishing Reinke crystals. *Hum Pathol*. Apr 2015;46(4):600-6.
doi:10.1016/j.humpath.2015.01.004
8.   Kozina V, Geist D, Kubinova L, et al. Visualization of Reinke's crystals in normal and
cryptorchid testis. *Histochemistry and cell biology*. Feb 2011;135(2):215-28.
doi:10.1007/s00418-011-0782-6
9.   Lobo MV, Arenas MI, Alonso FJ, et al. Nestin, a neuroectodermal stem cell marker
molecule, is expressed in Leydig cells of the human testis and in some specific cell types from
human testicular tumours. *Cell and tissue research*. Jun 2004;316(3):369-76.
doi:10.1007/s00441-003-0848-4
10.   Mesa H, Gilles S, Datta MW, et al. Immunophenotypic differences between neoplastic
and non-neoplastic androgen-producing cells containing and lacking Reinke crystals. *Virchows*
Arch. Dec 2016;469(6):679-686. doi:10.1007/s00428-016-2028-4
11.   Natsumeda Y, Ohno S, Kawasaki H, Konno Y, Weber G, Suzuki K. Two distinct cDNAs
for human IMP dehydrogenase. *The Journal of biological chemistry*. Mar 25 1990;265(9):5292-
5.
12.   Carr SF, Papp E, Wu JC, Natsumeda Y. Characterization of human type I and type II
IMP dehydrogenases. *The Journal of biological chemistry*. Dec 25 1993;268(36):27286-90.
13.   Hager PW, Collart FR, Huberman E, Mitchell BS. Recombinant human inosine
monophosphate dehydrogenase type I and type II proteins. Purification and characterization of
inhibitor binding. *Biochemical pharmacology*. May 11 1995;49(9):1323-9. doi:10.1016/0006-
2952(95)00026-v
14.   Jackson RC, Weber G, Morris HP. IMP dehydrogenase, an enzyme linked with
proliferation and malignancy. *Nature*. Jul 24 1975;256(5515):331-3. doi:10.1038/256331a0
15.   Senda M, Natsumeda Y. Tissue-differential expression of two distinct genes for human
IMP dehydrogenase (E.C.1.1.1.205). *Life Sci*. 1994;54(24):1917-26. doi:10.1016/0024-
3205(94)90150-3
16.   Keppeke GD, Chang CC, Peng M, et al. IMP/GTP balance modulates cytoophidium
assembly and IMPDH activity. *Cell Div*. 2018;13:5. doi:10.1186/s13008-018-0038-0

17. Chang CC, Lin WC, Pai LM, et al. Cytoophidium assembly reflects upregulation of
IMPDH activity. *Journal of cell science*. Oct 1 2015;128(19):3550-5. doi:10.1242/jcs.175265
18. Calise SJ, Chan EKL. Anti-rods/rings autoantibody and IMPDH filaments: an update
after fifteen years of discovery. *Autoimmun Rev*. Oct 2020;19(10):102643.
doi:10.1016/j.autrev.2020.102643
19. Johnson MC, Kollman JM. Cryo-EM structures demonstrate human IMPDH2 filament
assembly tunes allosteric regulation. *eLife*. Jan 30 2020;9doi:10.7554/eLife.53243
20. Valvezan AJ, Turner M, Belaid A, et al. mTORC1 Couples Nucleotide Synthesis to
Nucleotide Demand Resulting in a Targetable Metabolic Vulnerability. *Cancer cell*. Nov 13
2017;32(5):624-638 e5. doi:10.1016/j.ccell.2017.09.013
21. Nass K, Redecke L, Perbandt M, et al. In cellulo crystallization of Trypanosoma brucei
IMP dehydrogenase enables the identification of genuine co-factors. *Nature communications*. Jan
30 2020;11(1):620. doi:10.1038/s41467-020-14484-w
22. Woulfe J, Munoz DG, Gray DA, Jinnah HA, Ivanova A. Inosine monophosphate
dehydrogenase intranuclear inclusions are markers of aging and neuronal stress in the human
substantia nigra. *Neurobiol Aging*. Feb 2024;134:43-56.
doi:10.1016/j.neurobiolaging.2023.11.005
23. Woulfe J, Munoz D. Roncoroni Re-Visited: The Neuronal Intranuclear Rodlet Comes of
Age. *The Journal of comparative neurology*. Aug 2024;532(8):e25662. doi:10.1002/cne.25662
24. Noree C, Begovich K, Samilo D, Broyer R, Monfort E, Wilhelm JE. A quantitative
screen for metabolic enzyme structures reveals patterns of assembly across the yeast metabolic
network. *Molecular biology of the cell*. Oct 1 2019;30(21):2721-2736. doi:10.1091/mbc.E19-04-
0224
25. Hvorecny KL, Hargett K, Quispe JD, Kollman JM. Human PRPS1 filaments stabilize
allosteric sites to regulate activity. *Nat Struct Mol Biol*. Mar 2023;30(3):391-402.
doi:10.1038/s41594-023-00921-z
26. Hove-Jensen B, Andersen KR, Kilstrup M, Martinussen J, Switzer RL, Willemoes M.
Phosphoribosyl Diphosphate (PRPP): Biosynthesis, Enzymology, Utilization, and Metabolic
Significance. *Microbiol Mol Biol Rev*. Mar 2017;81(1)doi:10.1128/MMBR.00040-16
27. Ivanova A, Munoz DG, Woulfe J. Depletion of nuclear cytoophidia in Alzheimer's
disease. *Free Neuropathol*. Jan 2025;6:8. doi:10.17879/freeneuropathology-2025-6282
28. Sararols P, Stevant I, Neirijnck Y, et al. Specific Transcriptomic Signatures and Dual
Regulation of Steroidogenesis Between Fetal and Adult Mouse Leydig Cells. *Front Cell Dev
Biol*. 2021;9:695546. doi:10.3389/fcell.2021.695546
29. Verrier F, An S, Ferrie AM, et al. GPCRs regulate the assembly of a multienzyme
complex for purine biosynthesis. *Nature chemical biology*. Oct 23 2011;7(12):909-15.
doi:10.1038/nchembio.690
30. Pedley AM, Pareek V, Benkovic SJ. The Purinosome: A Case Study for a Mammalian
Metabolon. *Annu Rev Biochem*. Jun 21 2022;91:89-106. doi:10.1146/annurev-biochem-032620-
105728
31. Mori Sequeiros Garcia MM, Paz C, Castillo AF, et al. New insights into signal
transduction pathways in adrenal steroidogenesis: role of mitochondrial fusion, lipid mediators,
and MAPK phosphatases. *Front Endocrinol (Lausanne)*. 2023;14:1175677.
doi:10.3389/fendo.2023.1175677
32. Dufau ML, Catt KJ. Gonadotropin receptors and regulation of steroidogenesis in the
testis and ovary. *Vitam Horm*. 1978;36:461-592. doi:10.1016/s0083-6729(08)60989-9

33.   Catt KJ, Dufau ML. Spare gonadotrophin receptors in rat testis. *Nat New Biol*. Aug 15 1973;244(137):219-21. doi:10.1038/newbio244219a0

34.   Ascoli M, Fanelli F, Segaloff DL. The lutropin/choriogonadotropin receptor, a 2002 perspective. *Endocr Rev*. Apr 2002;23(2):141-74. doi:10.1210/edrv.23.2.0462

35.   Catt KJ, Harwood JP, Clayton RN, et al. Regulation of peptide hormone receptors and gonadal steroidogenesis. *Recent Prog Horm Res*. 1980;36:557-662. doi:10.1016/b978-0-12-571136-4.50021-8

36.   Ramaswamy S, Plant TM, Marshall GR. Pulsatile stimulation with recombinant single chain human luteinizing hormone elicits precocious sertoli cell proliferation in the juvenile male rhesus monkey (Macaca mulatta). *Biol Reprod*. Jul 2000;63(1):82-8. doi:10.1095/biolreprod63.1.82

37.   Dufau ML, Baukal AJ, Catt KJ. Hormone-induced guanyl nucleotide binding and activation of adenylate cyclase in the Leydig cell. *Proceedings of the National Academy of Sciences of the United States of America*. Oct 1980;77(10):5837-41. doi:10.1073/pnas.77.10.5837

38.   Liu JL. The Cytoophidium and Its Kind: Filamentation and Compartmentation of Metabolic Enzymes. *Annu Rev Cell Dev Biol*. Oct 6 2016;32:349-372. doi:10.1146/annurev-cellbio-111315-124907

39.   Nagano T, Otsuki I. Reinvestigation of the fine structure of Reinke's crystal in the human testicular interstitial cell. *The Journal of cell biology*. Oct 1971;51(1):148-61. doi:10.1083/jcb.51.1.148

40.   Schonherr R, Rudolph JM, Redecke L. Protein crystallization in living cells. *Biol Chem*. Jun 27 2018;399(7):751-772. doi:10.1515/hsz-2018-0158

41.   Dodson G, Steiner D. The role of assembly in insulin's biosynthesis. *Curr Opin Struct Biol*. Apr 1998;8(2):189-94. doi:10.1016/s0959-440x(98)80037-7

42.   Giembycz MA, Lindsay MA. Pharmacology of the eosinophil. *Pharmacol Rev*. Jun 1999;51(2):213-340.

43.   Soerensen RR, Johannsen TH, Skakkebaek NE, Rajpert-De Meyts E. Leydig cell clustering and Reinke crystal distribution in relation to hormonal function in adult patients with testicular dysgenesis syndrome (TDS) including cryptorchidism. *Hormones (Athens)*. Oct 2016;15(4):518-526. doi:10.14310/horm.2002.1708

44.   Chemke J, Shani A, Lancet M, Czernobilsky B. Testicular feminisation syndrome: unusual gonadal histology in an elderly patient. *J Med Genet*. Oct 1977;14(5):375-8. doi:10.1136/jmg.14.5.375

45.   Kerr JB. Ultrastructure of the seminiferous epithelium and intertubular tissue of the human testis. *J Electron Microsc Tech*. Oct 1991;19(2):215-40. doi:10.1002/jemt.1060190208

46.   Kofuji S, Hirayama A, Eberhardt AO, et al. IMP dehydrogenase-2 drives aberrant nucleolar activity and promotes tumorigenesis in glioblastoma. *Nature cell biology*. Aug 2019;21(8):1003-1014. doi:10.1038/s41556-019-0363-9

47.   Kofuji S, Sasaki AT. GTP metabolic reprogramming by IMPDH2: unlocking cancer cells' fuelling mechanism. *J Biochem*. Oct 1 2020;168(4):319-328. doi:10.1093/jb/mvaa085

48.   Kozhevnikova EN, van der Knaap JA, Pindyurin AV, et al. Metabolic enzyme IMPDH is also a transcription factor regulated by cellular state. *Molecular cell*. Jul 13 2012;47(1):133-9. doi:10.1016/j.molcel.2012.04.030

49.   Watts RW, Harkness RA, Spellacy E, Taylor NF. Lesch-Nyhan syndrome: growth delay,
testicular atrophy and a partial failure of the 11 beta-hydroxylation of steroids. *J Inherit Metab*
Dis. 1987;10(3):210-23. doi:10.1007/BF01800063
50.   Ogawa-Iio A, Takeuchi K, Shigemi K, et al. IMPDH and GTP Metabolism in Cancer:
Mechanisms, Regulation, and Translational Scope. *Cancer science*. Oct 3
2025;doi:10.1111/cas.70200
453

Figure Legends:

Figure 1: Reinke crystals are immunoreactive for IMPDH and PRPS. A-D: Reinke crystals in a

Leydig cell tumour. A) Cytoplasmic (arrows) and nuclear (arrowheads) Reinke crystals are

visible in an H and E stained section. B) A rod-shaped nuclear crystal extends to a length greater

than the nuclear diameter, distorting and stretching the nuclear membrane (arrowheads). C, D)

Immunostaining for IMPDH (C) and PRPS (D) reveals more abundant Reinke crystals

morphologically identical to those visualized in H and E-stained sections. E) IMPDH

immunohistochemistry is more sensitive for the demonstration of nuclear Reinke crystals. Many

nuclear crystals (eg. arrowheads) stain with reduced intensity relative to cytoplasmic crystals. F)

Both cytoplasmic and nuclear crystals are visualized in non-neoplastic Leydig cells using

IMPDH immunohistochemistry. Scale bars=50 µm in A, C, and D; 25 µm in B, E, and F.

Figure 2: IMPDH and PRPS immunoreactivity are co-localized in Reinke crystals. Double

immunofluorescence staining of neoplastic Leydig cells reveals immunoreactivity of cytoplasmic

and nuclear (arrow) Reinke crystals for both IMPDH (A; green) and PRPS (B; red). Sections are

counterstained with DAPI (blue). Scale bars=5 µm.

A **IMPDH / DAPI**, B **PRPS / DAPI**, C **Merge / DAPI**. Scale bars: 5 µm.