From Stability to Complexity: A Systematic Review of Long-term Divergence Exponents in Nonlinear Gait Analysis

Jeremy Torrent[1][¶], Roxane Coquoz[1][¶], Philippe Terrier[1][*]

1 Haute-Ecole Arc Santé, HES-SO University of Applied Sciences and Arts Western Switzerland, Neuchâtel, Switzerland

¶ These authors contributed equally to this work.

* Corresponding Author:

Philippe Terrier
Haute Ecole Arc Santé
Espace de l'Europe 11, CH-2000 Neuchâtel, Switzerland
Philippe.terrier@he-arc.ch

Keywords: Human locomotion; Walking; Gait analysis; Gait variability; Gait automaticity; Gait complexity; Nonlinear dynamics; Lyapunov exponent

Abstract

Background

Divergence exponents (DE), or maximum Lyapunov exponents, computed from stride-to-stride fluctuations have traditionally been interpreted as measures of gait stability. However, evidence suggests this measure may also reflect gait complexity and automaticity. This systematic review evaluated empirical support for reinterpreting long-term DE as a complexity measure.

Methods

We systematically searched Web of Science databases through August 2024 for studies applying Rosenstein's algorithm to human gait. We systematically extracted experimental conditions, and participant characteristics from each study. Study quality was assessed using a framework evaluating analytical rigor, outcome reporting, and sample size adequacy. We conducted a meta-analysis examining correlations between long-term DEs and detrended fluctuation analysis (DFA) scaling exponents, and synthesized evidence from perturbation (environmental disturbances), cueing (external rhythmic stimuli), and between-subject (clinical vs control) studies.

Results

Fifty-six studies published between 2000 and 2024 met inclusion criteria, with 43% achieving high overall quality scores. Meta-analysis from six datasets (209 participants) revealed a positive correlation between long-term DE and DFA scaling exponents (r=0.64, 95% CI 0.34 to 0.82; I²=82%). Perturbation studies consistently showed and increased short-term DE (lower local stability) while simultaneously decreasing long-term DE by up to 64%. External auditory and visual cueing interventions induced long-term DE decreases (up to -86%) while minimally affecting short-term DEs. Between-subject comparisons revealed heterogeneous patterns, with clinical populations exhibiting both increases and decreases in long-term DE depending on pathology.

Conclusions

Converging meta-analytic and experimental evidence supports reinterpreting long-term DE as the Attractor Complexity Index—a measure of gait complexity and automaticity rather than stability. Reduced long-term DE during controlled gait conditions reflects suppression of low-frequency variations in gait variability, which constrains phase space exploration and accelerates divergence curve saturation. The measure's sensitivity to prefrontal cortex engagement and attentional demands reveals that long-term DE captures the degree of gait

automaticity, with lower values indicating a shift from automatic subcortical control to

executive function-mediated regulation. This attention-dependent control reorganization

explains patterns observed in aging, clinical populations, and experimental perturbations.

This paradigm shift establishes long-term DE as a complementary biomarker for motor-

cognitive aspects of gait control, with implications for fall risk assessment, disease

monitoring, and rehabilitation evaluation.

Introduction

Walking is a complex motor activity that plays an important role in daily functioning [1]. Gait—the pattern of limb movement—can serve as a significant biomarker, providing insights into an individual's physical functioning [2,3], cognitive abilities [4], and level of independence [5]. Slower speeds and deviations from normal gait patterns have been shown to correlate with adverse outcomes, including an increased risk of falls [6,7], cognitive decline [7,8], and mortality [9]. For this reason, gait analysis is an essential component of clinical evaluation across medical specialties [10,11].

Nonlinear gait analysis is a methodological approach that examines the seemingly unpredictable dynamics of human gait using mathematical and statistical tools from nonlinear systems theory [12]. Nonlinear dynamical systems are characterized by outputs that are not proportional to changes in inputs, where the behavior cannot be described by simple linear differential equations. In the context of physiological signals, nonlinearity reflects how small changes in one component can lead to disproportionate or emergent effects throughout the system. These complex interactions arise from the integration of multiple regulatory mechanisms—such as the interplay between the autonomic nervous system and neuroendocrine factors in heart rate regulation—forming a dynamic network whose behavior cannot be fully understood by analyzing individual components in isolation [13,14]. Nonlinear analysis techniques have become an integral part of gait analysis, providing advanced methods such as maximum Lyapunov (or divergence) exponents for assessing resilience to perturbations [15], fractal dynamics for investigating the complexity and statistical persistence of time series [16], and entropy measures for quantifying signal irregularity [17]. These methods facilitate the detection of gait disorders [18–20], and support the development of innovative rehabilitation strategies [21]. By going beyond traditional kinematic and kinetic parameters, nonlinear approaches allow for a more comprehensive assessment of gait variability [22].

Nonlinear gait analysis also provides a window into the underlying neural control mechanisms [23], including the involvement of executive function in gait [4]. Attention, as a component of executive function, refers to the ability to allocate cognitive resources among tasks, including gait, that are performed concurrently [4,24]. Gait automaticity reflects the extent to which executive function contributes to gait control, with highly automatic gait requiring little attentional resources and less automatic gait requiring increased executive involvement [25]. One particularly powerful approach for investigating the interplay between executive function and gait is the use of dual-task paradigms. In these experimental setups,

participants are asked to perform a cognitive task while walking, which allows researchers to

assess how the allocation of attentional resources affects gait parameters [4]. Nonlinear

measures applied to dual-task walking data can reveal how the complexity and stability of

gait are modulated by concurrent cognitive demands, providing valuable information about

the adaptive capacity of the motor control system and the degree of gait automaticity

[26,27]. This approach has proven useful in understanding age-related changes in gait

control [28] and in identifying early markers of cognitive decline [29] and fall risk [30] in

various clinical populations.

Beyond dual-task paradigms, various factors—including sensory inputs, environmental

challenges, and individual physiological and psychological states—can influence the extent of

executive function involvement during walking, and nonlinear gait analysis provides valuable

tools for exploring these dynamics. For example, the influence of auditory cues on gait

automaticity can be observed through rhythmic auditory stimulation, where external rhythms

entrain motor rhythms, potentially bypassing impaired internal gait control mechanisms in

conditions such as Parkinson's disease [31,32]. Exposing older adults to visual perturbations

significantly reduces the complexity of their gait, a response not observed in younger adults.

This may be indicative of an increased reliance on visual feedback and a decrease in gait

automaticity with age [33]. Environmental constraints, such as treadmill walking, can alter

motor control strategies. Research has shown that stride-speed fluctuations exhibit statistical

anti-persistence during treadmill walking, indicating tight control over this gait parameter.

This tight control is further enhanced when maneuverability is restricted, suggesting that the

nervous system actively minimizes speed deviations [34]. These findings are consistent with

the idea that the need for continuous speed adjustments on a treadmill may increase the

demands on executive function to maintain a steady gait [35]. Individual physiological and

psychological states, such as pain, attentional focus, of anxiety, may also influence gait

automaticity. For example, research suggests that individuals with chronic low back pain

exhibit altered gait complexity, possibly due to attentional strategies focused on managing

pain [36]. In addition, fear of falling is associated with reduced local dynamic stability of gait

[37], which may be explained by an increased reliance on executive functions to maintain

balance and adapt to perceived threats during walking.

Among nonlinear measures, the maximum Lyapunov exponent [divergence exponent

(DE)] method stands out as a prominent measure for assessing dynamic stability and fall risk

[20,38]. This prominence its due to its ability to quantify local dynamic stability—a system's

sensitivity to small perturbations [15,39,40]. Modeling studies using dynamic walker systems

have validated its predictive validity, demonstrating strong correlations between short-term

DEs and simulated fall probabilities [41,42]. These computational validations, combined with
clinical observations of elevated DE in fall-prone elderly, position it as an important tool for
assessing gait stability and fall risk in both research and clinical applications [20]. However,
recent systematic reviews have shown heterogeneity in the implementation of the DE
method, with different parameterizations and measurement systems across studies [20,43],
highlighting the need for further research to standardize methodologies and facilitate cross-
study comparisons.

A persistent methodological challenge in DE-based gait analysis arises from the
differential responsiveness of short- and long-term DE computed by Rosenstein's algorithm.
Mathematically, Rosenstein's method embeds the time series in a state space and identifies
for each point a nearest neighbor such that their initial separation $d(0)$ evolves with time t
as $d(t) = d(0)e^{\lambda t}$, where λ is the DE characterizing the intensity of the exponential
divergence, which is also known as the "butterfly effect" [44]; taking the natural logarithm
gives $ln(d(t)) = ln(d(0)) + \lambda t$, and hence $\lambda \approx \frac{ln\,(d(t))}{\Delta t}$ [15,39]. The characteristic logarithmic
divergence curve is computed to assess the average progressive separation. Short-term DE is
computed as the average rate of divergence by linear regression of the initial segment of the
curve (0-1 step or stride), capturing local dynamic stability through immediate corrective
mechanisms. Conversely, long-term DE is derived from later intervals (4-10 strides) and
reflects long-term system behavior. While elevated short-term DE consistently correlates with
fall risk and other variability metrics [45], long-term DE exhibits paradoxical responses to
experimental manipulations, decreasing during cognitively demanding walking [33,46] and
showing weak, or even opposite, associations with stability measures [45,47].

Therefore, while long-term DE was initially designed to measure gait stability, it has
been reinterpreted as capturing distinct aspects of gait control compared to short-term DE.
Several studies have demonstrated strong associations between long-term DE and the
correlation structure of stride intervals, as quantified by scaling exponents that characterize
stride-to-stride fluctuation complexity [39,48,49]. This relationship is particularly evident in
experimental conditions that modify gait automaticity. Metronome walking, which requires
conscious sensorimotor synchronization and increased attentional demands, substantially
decreases both long-term DE and gait complexity while leaving short-term DE unchanged
[46,50]. Furthermore, it has been shown that gait acceleration signals with identical shapes,
but modified stride interval complexity exhibit different long-term DE values [51]. These
findings have led to its reconceptualization as the "Attractor Complexity Index" (ACI),
reflecting its role in quantifying gait complexity and automaticity rather than stability [51].

The primary objective of this systematic review is therefore to examine the use of long-
term DE in nonlinear gait analysis, with a particular focus on its reinterpretation as a
measure of gait complexity rather than stability. By synthesizing existing evidence, this
review seeks to reconcile historical interpretations of long-term DE as a stability metric with
emerging evidence highlighting its association with fractal dynamics and stride-to-stride
fluctuation complexity. This synthesis may clarify the potential of long-term DE in the
assessment of gait automaticity. A secondary objective is to address the substantial
methodological heterogeneity observed across studies using long-term DE. This includes
variations in measurement systems, signal processing techniques, and parameterization of
Rosenstein's algorithm. By evaluating these methodological differences, we aim to identify
best practices and propose standardized protocols for future research. Ultimately, this review
aims not only to refine our understanding of long-term DE as a measure of gait complexity
and automaticity, but also to facilitate more reliable and comparable applications in both
research and clinical contexts.
180

Methods

Reporting and transparency

The systematic review followed the PRISMA guidelines to ensure transparency and
comprehensive reporting, detailing the rationale, methodology, and results of the review
[52]. However, it is important to note that the protocol for this review was not registered in
PROSPERO or any other repository. Methodological reviews focusing on tool development or
reporting practices without direct clinical applicability are typically excluded from PROSPERO
registration requirements [53]. Despite the lack of formal registration, efforts were made to
maintain transparency by publishing the protocol as a non-peer-reviewed preprint [54]. The
readers are invited to consult the published protocol for further details about the review
procedures.
192

Study eligibility criteria

The review included original peer-reviewed research articles focusing on human locomotion
using the maximum Lyapunov exponent method with Rosenstein's algorithm to compute
long-term DEs. Database searches applied no language restrictions to ensure comprehensive
retrieval of potentially relevant literature. Eligible studies investigated gait in healthy
participants or patients across all age groups and were published between 2000 and 2024.

Observational, experimental, and intervention studies were included if they reported
sufficient methodological details on DE computation. Studies were excluded if they: focused
on modes of locomotion other than walking (e.g., running or jumping), relied exclusively on
short-term DEs or alternative analyses (e.g., Wolf's algorithm), were pure modeling or
simulation studies lacking empirical human data, or did not adequately describe the DE
computation parameters. Reviews, letters, editorials, and conference abstracts were also
excluded. Full-text screening was restricted to articles published in English or French based
on the linguistic competencies of the review team. Articles identified in languages other than
English or French underwent title, abstract, and methods section screening using automated
translation tools to assess eligibility against inclusion criteria.

Information sources and search strategy

212 We performed a systematic search using the Web of Science platform, including MEDLINE
213 and the Web of Science Core Collection databases. The search strategy combines themes
214 related to gait analysis, nonlinear dynamics, and Lyapunov exponents. Boolean operators and
215 truncations were used to ensure comprehensive coverage. The search keywords are
216 described in the supporting information (S1 Text).

Study selection process

The study selection process was conducted in two phases, the results of which are
summarized in a PRISMA flowchart (Fig 1). Initially, two independent reviewers screened the
titles and abstracts of all retrieved articles using broad inclusion criteria, as specific
references to long-term divergence exponents were most often missing from abstracts [54].
Articles potentially involving nonlinear gait analysis were advanced to the second phase,
where their full texts were thoroughly reviewed to confirm the use of long-term divergence
exponents calculated with Rosenstein's algorithm. Disagreements between reviewers were
resolved through discussion, with arbitration by a third reviewer when necessary [54].

Data collection process and items

Main outcomes. Study outcomes were collected using standardized extraction tables as
specified in the protocol [54], and the extracted results are available in the supporting
information for detailed reference (S1 Text). For each included study, three main categories
of information were extracted: study characteristics, methodological specifications, and

outcomes related to short-term and long-term DEs. Study characteristics included publication
year, authorship, sample size, participant demographics (age, health status), and research
context (e.g., methodological studies or clinical evaluations). Methodological specifications
included details on measurement methods (e.g., primary measurement system, signal type,
sampling rate), walking conditions (overground or treadmill), signal processing procedures,
and specific computational parameters for DEs (e.g., time delay, embedding dimension).
Finally, the extracted findings focused on study objectives, conclusions, and interpretations of
DE results.

Divergence curve characterization. In addition to extracting numerical outcomes, we
systematically identified studies that published graphical representations of divergence
curves. Divergence curves provide a visual representation of how the DE are calculated.
For each study that published divergence curves, we documented the figure location and
extracted relevant information about the visual presentation of the method. For articles
available through open-access licenses, we reproduced the published figures along with their
complete legends. For studies behind paywalls where figure reproduction was not allowed,
we documented the figure legends and descriptions to catalog the types of divergence curve
presentations.

Quality assessment

Quality assessment of the included studies was performed using a structured evaluation
framework across three domains. Each study was scored on a 15-point scale comprising
Domain 1: analytical rigor, 8 points; Domain 2: outcome reporting, 4 points; and domain 3
sample size, 3 points. Domain 1 assessed methodological transparency through four criteria:
experimental protocol description, measurement system reporting, signal processing clarity,
and algorithm application details. Domain 2 evaluated the quality of statistical reporting,
specifically whether descriptive statistics were clearly reported, and whether appropriate
statistical analyses with effect sizes were provided. Domain 3 considered sample size
adequacy relative to each study's research question. This structured approach allowed for
systematic comparison of methodological quality across the corpus while recognizing the
exploratory nature of the field and the developmental stage of long-term DE methodology.
Item-level domain scores and total quality scores for all 56 studies are provided in S1 Table
(XLSX).

Synthesis method: meta-analysis

We found four studies that examined associations between long-term DEs and scaling

exponents [39,48,55,56]. In addition, we added our recent study that also reported

correlation results [50], yielding five publications contributing six independent study sets.

These studies reported multiple correlation coefficients, corresponding to different

measurements (accelerometer axes, body positions). Therefore, to synthesize their results,

we performed a meta-analysis using a multilevel random-effects approach in accordance with

current methodological recommendations for handling dependent effect sizes [57]. We

implemented a correlated and hierarchical effects model using the metafor package in R

[58]. This model accounts for both between-study heterogeneity and within-study correlation

structures. We assumed within-study correlation of 0.8 between effect sizes from the same

study, as the different correlation coefficients consisted of repeated measures in the same

subjects. Within each study, repeated correlation coefficients were aggregated to a single

study-level estimate using a generalized least-squares average that respects the assumed

within-study correlation. Robust variance estimation (RVE) with small-sample correction was

applied using the clubSandwich package [59] to account for potential misspecification of the

working model and to generate valid inferences despite the dependence structure of the

data. Fisher's Z-transformed correlations were used for analysis and back-transformed to

correlation coefficients for interpretation. We also estimated heterogeneity and summarized it

with I^2. Results were visualized using forest plots displaying individual and overall estimates

with their respective 95% confidence intervals.

Protocol deviations

We acknowledge four protocol deviations that occurred during the conduct of this review.

First, the eligibility criteria for study inclusion originally specified a publication date range

from 2001 to 2024. However, it became apparent that seminal articles relevant to the

analysis of DE in nonlinear gait analysis were published as early as 2000. Consequently, the

inclusion criteria were adjusted to include publications from 2000 to 2024. Second, although

not explicitly stated in the original protocol, we decided during the data extraction phase to

identify and report studies using the detrended fluctuation analysis (DFA) method in parallel

to long-term DE. Third, we systematically documented divergence curve presentations to

capture how these visualizations demonstrate the extraction of short-term and long-term DEs

and reveal characteristics of the underlying gait dynamics. Fourth, the protocol specified a

narrative quality assessment approach. However, upon initiating the assessment process, we

recognized that a more structured framework would facilitate systematic evaluation and

comparison across studies. We therefore developed a formal quality assessment grid with

explicit scoring criteria across three methodological domains, enabling quantitative
characterization of reporting quality while maintaining the focus on methodological
transparency rather than risk of bias.
305

Results

307
As shown in Fig 1, our systematic search retrieved 673 records from database queries and 3
additional records from references of other articles, yielding 676 unique records for title-
abstract screening. Of these, 242 were excluded: 102 for non-relevant sources (proceedings,
reviews), 51 for modeling studies, and 89 for irrelevant topics or non-walking gaits (running).
Among the excluded records, 4 articles published in languages other than English or French
(two Spanish, one German, one Chinese) underwent screening based on English abstract;
none met the inclusion criteria. This left 434 full-text articles for eligibility assessment, all of
which were in English. The full-text review resulted in the exclusion of 378 articles, mainly
due to reporting only short-term DE. Other secondary reasons for exclusion were the use of
alternative DE algorithms (Wolf) or other stability metrics. As a result, 56 studies met the
criteria for outcome synthesis, and 4 studies provided the necessary data for inclusion in our
meta-analysis.
As shown in Figure 1, our systematic search retrieved 673 records from database queries
and 3 additional records from references of other articles, yielding 676 unique records for
title-abstract screening. Of these, 242 were excluded: 102 for non-relevant sources
(proceedings, reviews), 51 for modeling studies, and 89 for irrelevant topics or non-walking
gaits (running. Among the excluded records, 4 articles published in languages other than
English or French (two Spanish, one German, one Chinese) underwent screening based on
English abstract; none met the inclusion criteria. This left 434 full-text articles for eligibility
assessment, all of which were in English. The full-text review resulted in the exclusion of 378
articles, mainly due to reporting only short-term DE. Other secondary reasons for exclusion
were the use of alternative DE algorithms (Wolf) or other stability metrics. As a result, 56
studies met the criteria for outcome synthesis, and 4 studies provided the necessary data for
inclusion in our meta-analysis.
332

Methodological outcomes

Study characteristics, methodological scope, and sample sizes. Fig 2 summarizes the
characteristics of the 56 included studies, based on the data provided in the supporting
information (Table B in S1 Text). Panel a illustrates the annual number of publications from
2000 to 2023, showing a consistent upward trend starting in 2008, with a peak of eight
articles published in 2009. Panel b categorizes the type of research by participant category.
Exact definitions for each research type are provided in supporting information (Table A in S1
Text). In terms of populations studied, studies involving only healthy adults predominate,
with 40 studies. Within this category, most studies compared gait patterns under different
experimental conditions and aimed to develop, validate, or assess the reliability of DE
methods. Research on age-related gait changes and fall risk comprises 10 studies, while the
remaining 6 studies investigated gait patterns associated with other pathologies. Only a few
studies have used long-term DE for clinical investigations (4). Only one study was
interventional and longitudinal, i.e. it analyzed the gait patterns of patients before and after
surgery. Aside from studies aimed at predicting fall risk, we identified no studies that
evaluated the use of long-term DE as a diagnostic tool for detecting gait abnormalities. Panel
c shows that across the study samples, the median sample size was 20 participants
(interquartile range 14-39), with most studies enrolling fewer than 50 individuals; only a
handful exceeded 100, and the full range extended from 9 to 187 participants, indicating a
generally modest sample size framework in the field. Note that one article was counted as
two different studies because it reported two separate experiments with two different groups
of participants.

Experimental settings and sensor modalities. Per-study methodological outcomes
for all 56 studies are provided in Table C in S1 Text. Measurement environments were
overwhelmingly laboratory-bound (Fig 3a). Forty of the 56 studies (71 %) acquired data on
an instrumented treadmill in an indoor laboratory setting, typically in conjunction with 3-D
motion-capture systems. Nine studies (16 %) relied solely on overground walking, but even
here the majority preserved a controlled indoor context (indoor circuits or short walkways).
Only two overground protocols departed from this paradigm by recording prolonged outdoor
or quasi-natural bouts. Six further articles (11 %) combined treadmill and overground
sessions within the same experiment, whereas one study (2 %) captured free-living gait
during a week-long monitoring campaign. Instrumentation choices mirrored these
environmental constraints (Fig 3b). Optical motion-capture was used in 29 of the 56 studies
(52 %). Wearable inertial measurement units (IMUs) ranked second, appearing in 16 studies
(29 %). Dedicated instrumented treadmills—equipped with embedded force or pressure
arrays—were the unique measurement device in 3 studies (5 %). Although single-sensor

protocols remained the norm, a minority of eight studies (14 %) employed hybrid

instrumentation—most commonly electrogoniometers coupled with a trunk accelerometer

(4/8, 50 %), followed by optical motion-capture combined with body-worn IMUs (2/8, 25 %)

and motion capture and surface electromyography (EMG, 2/8, 25 %). Most studies prioritized

measurement at the trunk level (Fig 3c): 31 of the 56 studies (55 %) analyzed pelvis, lumbar

or thoracic spine kinematic. Lower-limb–only set-ups (hip, knee, ankle kinematics or EMG)

appeared in 11 studies (20 %). Nine studies (16 %) combined trunk and lower-limb

measurements within the same experiment. Foot-pressure–only configurations were rare

(3/56, 5 %), corresponding to the instrumented-treadmill studies. Other designs include

head or upper-limb recordings studies (2/56, 4 %).

Measurement lengths. Because Rosenstein's algorithm requires a single continuous

signal, we extracted the duration corresponding to one analyzed walking bout per

experimental condition, rather than aggregating multiple bouts. Reporting practices across

the included studies varied substantially: approximately one-third specified bout length

exclusively in minutes, another third exclusively in strides, and the remainder provided both

units. Due to this methodological heterogeneity, direct quantitative synthesis of durations

was not feasible. We therefore adopted an ordinal categorization to harmonize duration data

into four empirically defined classes (Fig 4). The threshold for the "very short" category (≤ 2

min or ≤ 100 strides) was anchored to the median cadence observed in studies reporting

both units (~55 strides·min^{-1}), yielding approximately 100 strides over two minutes—a

practical lower limit below which long-term DE estimates may become unreliable [60]. The

upper limit for the "short" category (> 2–5 min or 101–250 strides) aligns with the most

commonly adopted experimental protocol across studies (3–5 minutes of treadmill or

overground walking). The "moderate" category (> 5–10 min or 251–500 strides)

encompasses studies employing somewhat longer walking bouts, while the "long" category

(> 10 min or > 500 strides) includes the few studies focused on endurance-oriented walking

conditions. Applying this ordinal scale to the extracted data resulted in the following

distribution: "short" (28 studies; 50%), "very short" (19 studies; 34%), "moderate" (5

studies; 9%), and "long" (4 studies; 7%). Finally, five studies computed DE from averaged

signals of multiple short walking bouts within the same experimental condition; for these, we

categorized duration based on a single constituent bout, although averaging multiple bouts

may enhance the reliability of DE estimates [60].

Signal processing. Signal sources driving the computation of long-term DEs were

clustered into distinct categories (Fig 5a). Over half of the studies (29 out of 56, 52 %)

utilized kinematic data exclusively, acquired via optical motion capture or electrogoniometers.

Within these kinematic studies, angular joint trajectories (joint angles) were predominant,
representing most cases. Additionally, linear positional data (e.g., cervical or thoracic spine
positions) and derived velocity data were also commonly used. Accelerometric data from
inertial sensors constituted the second most frequent signal type, with 15 studies (27 %)
employing linear accelerations exclusively. Only two studies (4 %) incorporated combined
inertial measurements of linear acceleration and angular velocity from gyroscopes. No
studies used gyroscopic (angular velocity) signals alone. Three studies (5%) derived DEs
from COP trajectories obtained from instrumented treadmills. Surface EMG voltage signals
alone were used in a single study (2 %). Mixed signal sources appeared infrequently: five
studies (9 %) combined joint-angle kinematics with trunk acceleration, and one study (2%)
combined whole-body kinematics with multi-channel EMG. Reporting of raw-signal filtering
was inconsistent across the corpus. Eighteen of the 56 studies (32 %) explicitly described
applying a digital filter before computing the long-term DE, almost always a zero-lag low-
pass Butterworth between 8 Hz and 20 Hz. Nineteen papers (34 %) stated unambiguously
that no filtering was performed. For the remaining nineteen studies (34 %) no filtering
procedure was mentioned, making it probable—though not certain—that the raw signals
were analyzed without additional pre-processing. The next steps in DE computation typically
involve time normalization of raw data. The rationale is to maintain a consistent number of
steps and data points across participants and conditions. This approach prevents potential
bias arising from variations in DE values associated with sample length [60]. Panel 2 of Fig 4
shows the handling of temporal resampling. Explicit time normalization—typically realized as
linear or spline interpolation to a constant number of samples—was reported in 22 of the 56
investigations (41 %). Two studies (4 %) used an alternative method consisting in
resampling separately each stride to a fixed number of samples. In contrast, 29 papers (52
%) gave no information on resampling, very likely using raw, unequal-length signals, and /
or signals with a variable number of steps. The remaining two articles (4 %) explicitly did not
perform time normalization.

Algorithms. The subsequent processing involves reconstructing the state-space on
which DE is computed. Among the 56 studies, 48 (86 %) used the classical Takens approach
(see Apendix A in [39]), generating a d_E-dimensional trajectory from a single signal and its T-
delayed copies [61]. By contrast, 8 studies (14 %) constructed ad-hoc, multi-channel
attractors by concatenating measured variables—typically linear plus angular trunk
kinematics, or acceleration combined with gyroscope signals—before any delay embedding.
Within the subset of studies using the classical approach (48 studies), the tuning procedure
for the delay and embedding dimension was reported heterogeneously: 28 of 48 studies (58

%) applied the standard pair of average mutual information (AMI) for T and global false-

nearest-neighbors (GFNN) for d_E, seven (15 %) substituted the autocorrelation criterion for T

while still using GFNN, and the remaining thirteen (27 %) gave no explicit or unclear

justification for either parameter. Finally, after reconstructing the state-space and computing

divergence curves, the long-term DE is determined by performing a linear fit on a specific

segment of the divergence curve [39]. The interval used for this linear fit was initially defined

in seminal studies by Dingwell et al. [40,62]. Most studies (52 out of 56, 93%) adopted this

canonical interval, 4 to 10 strides. Four studies (7%) selected alternative ranges: one used

0.5–4 strides, one 2–6 strides, one 4–7 strides, and another explored multiple intervals (1–4,

4–7, and 7–10 strides). Additionally, most studies computed short-term DE, an approach first

introduced by Dingwell et al. in 2001. Only four studies (7%), including the two early

investigations by Dingwell and colleagues published before 2001, exclusively focused on

long-term DE without reporting short-term DE.

Divergence curve visualization. Divergence curves—graphical representations

plotting the average logarithmic divergence of neighboring trajectories against time—

demonstrate how short-term and long-term DEs are extracted as linear slopes from distinct

temporal regions. In addition, their shape may allow visualizing when divergence ceases to

grow as trajectories reach bounded regions of phase space. Among the 56 included studies,

29 (52 %) published divergence curves in their figures. We systematically categorized the

purpose for which divergence curves were presented across these studies. Eighteen studies

(62 %) used divergence curves exclusively for methodological illustration, displaying

representative curves to demonstrate the calculation procedure without comparing

experimental conditions. Six studies (21 %) employed divergence curves exclusively as

comparative results, presenting curves to highlight differences between groups or

experimental conditions. Five studies (17 %) included both types of presentations,

incorporating separate figures for methodological illustration and comparative outcomes.

Among studies presenting comparative results, typical comparisons included different walking

speeds, perturbation versus control conditions, young versus older adults, and with versus

without external cueing. The compiled divergence curves and associated figure descriptions

are provided in S2 Figures.

Quality Assessment

Quality assessment was completed using the three-domain scoring framework

(maximum 15 points). Overall methodological quality was moderate to high, with a mean

score of 11.45 (SD = 1.70, range 8-15) out of 15 possible points. Twenty-four studies

(42.9%) achieved high quality (12-15 points), while 32 studies (57.1%) demonstrated

moderate quality (8-11 points). No studies scored in the low-quality range (0-7 points).

Complete quality assessment scores for all studies are provided in S1_Table.

Domain-specific analysis revealed distinct patterns across methodological areas. Domain

1 (analytical rigor) demonstrated the strongest performance, with studies averaging 7.62 ±

0.65 out of 8 points (95.3%), and 40 studies (71.4%) achieving perfect scores (8/8). This

indicates robust methodological rigor in experimental protocols, measurement systems,

signal processing, and algorithm implementation across the included literature. In contrast,

Domain 2 (outcome reporting) showed moderate performance (mean 2.20 ± 1.13 out of 4

points; 54.9%), with only 10 studies (17.9%) achieving full credit for comprehensive

statistical reporting. Domain 3 (sample size) similarly revealed limitations, averaging 1.62 ±

0.75 out of 3 points (54.2%), with most studies (41.1%) demonstrating moderate sample

size justification and only 7 studies (12.5%) providing adequate sample size justification.

Long-term DE results

Perturbation studies. Among the 56 studies, 12 studies experimentally destabilized gait

using a range of perturbation modalities, including mechanical disturbances (e.g., compliant

surfaces, segmental loads, unexpected slips, platform instability), sensory manipulations

(e.g., galvanic vestibular stimulation, blindfolding, optical-flow distortions, foot

desensitization), or cognitive disturbance (dual-tasking). These experimental paradigms are

particularly relevant for testing the hypothesis that long-term DE reflects the degree of gait

automaticity. When gait is perturbed or destabilized, it is reasonable to assume that

individuals allocate greater attentional resources to gait control in order to manage the

imposed challenges. However, none of the 12 perturbation studies explicitly tested this

hypothesis, as they interpreted long-term DE in the same manner as short-term DE—that is,

as a direct index of gait stability sensitive to the magnitude of external perturbation. We

summarize the results of the perturbation studies in Table 1. Because long□term DE is

sensitive to signal length (number of data points or strides), valid comparisons require similar

data lengths across conditions. In treadmill perturbation paradigms—where walking speed is

held constant between baseline and challenge—data length remains comparable even

without time normalization. Therefore, we excluded studies with very□short bouts or

overground designs in which walking speed (and thus signal length) differed between

conditions, retaining only treadmill experiments with sufficient bout duration and matched

speeds. Note that the study by Manor et al. analyzed a reduced number of consecutive

strides (45), but reported results averaged across three different walking speeds, which may

enhance the reliability of the DE assessment.

Table 1. Results of the perturbation studies

Study	Measurement method	Perturbation	Short-term DE change with perturbation	Long-term DE change with perturbation
Manor et al. ID 15 [63]	2D motion capture, sagittal plane hip, knee, and ankle angles.	Ice-induced plantar desensitization	**+8%**	**+44%**
McAndrew et al. ID 22 [64]	3D motion capture, 3D velocity of the cervical spine.	Mechanical and visual	Mechanical: **+10% to +72%** Visual: **+3% to +86%**	Visual: **-7% to -51%** Mechanical: **+5% to -38%**
Van Schooten et al. ID 26 [47]	3D inertial sensor, accelerations and angular velocities of the lower back.	Vestibular	**+8 to +10%**	**-4% to -20%**
Sinitksi et al. ID 30 [65]	3D motion capture, mediolateral velocity of the cervical spine.	Mechanical and visual	Mechanical: **+30% to +34%** Visual: **+19% to +34%**	Mechanical: **-27% to -30%** Visual: **-9% to -35%**
Rabago et al. ID 42 [66]	3D motion capture, mediolateral velocity of the cervical spine.	Cognitive, mechanical, and visual	Cognitive: **-3% to 0%** Mechanical: **+8% to +28%** Visual: **+10% to +30%**	Cognitive: **+6% to -3%** Mechanical: **-16% to -39%** Visual: **-17% to -41%**
Reynard and Terrier ID 43 [67]	3D accelerometer. Mediolateral acceleration of the upper trunk (sternum).	Vision removal	**-2% to +7%**	**0% to -20%**
Franz et al. ID 41 [33]	3D motion capture. Right and left heel positions relative to the sacrum position.	Visual	Young: **+6%** Older: **+22%**	Young: **-13%** Older: **-17%**
Qiao et al. ID 47 [68]	3D motion capture, 3D velocity of the cervical spine.	Visual	Healthy young controls: **+19% to +24%** Elderly non-fallers: **+20% to +23%** Elderly fallers: **+19% to + 28%**	Healthy young controls: **-7% to -32%** Elderly non-fallers: **+2% to -49%** Elderly fallers: **-7% to -44%**

Perturbation types included sensory manipulation (ice-induced plantar desensitization),

mechanical disturbances (platform translations), visual alterations (moving room, optic flow),

vestibular stimulation (galvanic stimulation), and cognitive tasks (dual-task paradigms).

Percentage changes represent relative differences between perturbation and baseline

conditions. Positive values indicate increased divergence exponent (DE) while negative values

indicate decreased DE. Ranges reflect different measurement axes, experimental groups, or

perturbation levels within individual studies. ID refers to study identification number in the

systematic review database (S1 text in supplementary material).

Study designs differed in the nature and number of perturbation conditions, which

explains why we report ranges rather than single effect sizes. For most studies, numeric

values were extracted from figures using WebPlotDigitizer (https://automeris.io/). Manor et

al. [63] desensitized the plantar arch using ice to diminish sensory feedback from the feet,

anticipating subsequent gait destabilization. Unlike other perturbation studies that analyzed

trunk motion, Manor et al. focused on lower-limb kinematics. Their findings indicated an

increase in both short-term (+8%) and long-term (+44%) DE values, with results

aggregated across different walking speeds and joints. In McAndrew et al. [64], participants

experienced both mediolateral and anteroposterior platform oscillations and visual-field

perturbations while DE was computed along the AP, ML, and vertical axes, yielding short-

term DE increases of 10–72 % (mechanical) and 3–86 % (visual) and long-term DE

decreases of 5–38 % and 7–51 %, respectively. Sinitski et al. [65] systematically varied the

amplitude of both mechanical and visual perturbations, observing short-term DE rises of 30–

34 % (mechanical) and 19–34 % (visual) alongside long-term DE drops of 27–30 % and 9–

35 %, respectively. Rábago et al. [66] compared cognitive, mechanical, and visual

perturbations: cognitive load had negligible effects on short-term DE (−3% to 0%) with

near-zero long-term changes (−3% to +6%), while both mechanical and visual perturbations

produced a similar pattern of increased short-term DE (+8% to +30%) accompanied by

decreased long-term DE (−16% to −41%). Van Schooten et al. [47] and Reynard & Terrier

[67] compared walking at multiple speeds under galvanic vestibular stimulation and

blindfolded conditions. They found short-term DE increases of 8–10 % (GVS) and −2 to +7

% (blindfold), and long-term DE decreases of 4–20 % and 0–20 %, respectively. Franz et al.

[33] analyzed younger and older subjects, reporting short-term DE increases of 6 % (young)

and 22 % (older), and long-term DE decreases of 13 % (young) and 17 % (older) under

visual perturbation; they also found a reduction in the scaling exponent reflecting gait

complexity (DFA method) by 6 % in young and 26 % in older participants. The study by Qiao

et al. assessed the effects of varying intensities of optical-flow distortions on young and older

adults. Here again, opposite responses were observed between short-term DE, which

increased (+19% to +28%), and long-term DE, which mostly decreased (+2% to -48%).

Across all studies reviewed, the majority of perturbation-induced changes were statistically

significant (p < 0.05), though certain exceptions occurred, notably for long-term DE

measures that exhibited greater inter-individual variability.

Between-subject studies. Thirteen between☐subject studies compared cohorts with

presumed differences in gait stability and potentially in gait automaticity, including clinical

groups (e.g. diabetic or peripheral neuropathy, knee osteoarthritis, chronic pain, fallers vs

non☐fallers) and demographic cohorts (healthy older vs younger adults, inexperienced vs

experienced military cadets). However, most of these studies based their DE analysis on too

short walking bouts or on non-time normalized signals. After careful evaluation of study

methodologies, seven studies met the inclusion criteria for a valid comparison (Table 2).

Although Terrier et al. analyzed very short walking bouts (35 strides, 1 min), they

aggregated numerous bouts collected over one week under free-living conditions (on

average 77 bouts in patients and 34 in healthy controls). This aggregation substantially

enhances the reliability of the average DE values, mitigating concerns related to short

individual signal durations.

Table 2. Results of the between-subject studies

Study	Measurement method	Population	Short-term DE difference with controls	Long-term DE difference with controls
Manor et al. ID 9 [69]	2D motion capture. Sagittal plane joint angles for hip, knee and ankle	Patients with all-cause peripheral neuropathy	**+4%**	**-4%**
Franz et al. ID 41 [33]	3D motion capture. Right and left heel positions relative to the sacrum position.	Healthy older adults	Normal walking: **+1%** Perturbation: **+15%**	Normal walking: **-10%** Perturbation: **-15%**
Terrier et al. ID 45 [70]	3D acceleration, right hip.	Patients with chronic musculoskeletal pain of the lower limbs.	**+4%**	**-19%**
Bizovska et al. ID 46 [71]	3D acceleration of the lumbar spine	Older healthy female subjects	Overground: **+4% to +18%** Treadmill: **+3% to +25%**	Overground: **+56% to -11%** Treadmill: **+22% to -8%**
Qiao et al. ID 47 [68]	3D motion capture, 3D velocity of the cervical spine.	Older non-faller adults and older faller adults	Controls vs. older non-fallers: **+6% to +13%** Controls vs older fallers: **+14% to +19%**	Controls vs. older non-fallers: **+37% to -45%** Controls vs older fallers: **+49% to -36%**
Bizovska et al. ID 49 [72]	3D acceleration of the trunk	Elderly fallers and frequent fallers	Controls vs. fallers: **+7% to +8%** Controls vs frequent fallers: **-14% to +11%**	Controls vs. fallers: **+8% to -3%** Controls vs frequent fallers:

				-7% to -18%
Matinazad et al. ID 50 [73]	3D motion capture, sagittal angle of the knee	Older female patients with knee osteoarthritis	+41% to +43%	-43% to -85%

Percentage differences in short-term and long-term divergence exponents (DE) between
clinical/aging populations and control groups. Populations included patients with all-cause
peripheral neuropathy, chronic musculoskeletal pain of lower limbs, knee osteoarthritis, and
older adults (non-fallers, single fallers, and frequent fallers). Positive values indicate higher
DE in the comparison group relative to controls, while negative values indicate lower DE.
Ranges reflect different measurement axes, walking conditions, or perturbation levels within
individual studies. ID refers to study identification number in the systematic review database
(S1 text in supplementary material).

Study methodologies differed in signal processing approaches and reported outcomes.
Manor et al. [69] averaged their analyses across joint angles and walking speeds, reporting a
single DE value per group (patients with peripheral neuropathy vs. healthy controls).
Similarly, Terrier et al. (2017) computed the vector magnitude (norm) from three axes of hip
acceleration, yielding a single representative DE per group (chronic musculoskeletal pain
patients vs. controls). Using 3D motion capture during treadmill walking, Franz et al. [33]
examined trunk motion responses to visual perturbations. Similarly, the visual perturbation
study of Qiao et al. [68] examined cervical spine velocity under three increasing perturbation
levels. Bizovska et al. [71] reported separate DE outcomes for each axis of a 3D
accelerometer under two distinct walking conditions (overground vs. treadmill), resulting in a
range of observed differences between older and younger healthy females. In their second
study, Bizovska et al. also analyzed DEs across accelerometer axes, comparing a control
group of healthy older adults with groups of older individuals who fell infrequently or
frequently. Finally, Matinazad et al. studied knee kinematic at two different speeds.
Short-term DE demonstrated a pattern of limited increases in most populations
compared to their respective controls, though with heterogeneity in magnitude. The most
pronounced differences were observed in patients with knee osteoarthritis (41-43% higher).
Clinical populations with neuropathy and chronic musculoskeletal pain showed consistent
increases compared to controls (+4%). Regarding age comparison studies, Franz et al.
reported minimal differences during normal walking (1% higher in older adults as compared
to younger controls) but substantially larger increases under visual perturbations (15%
higher in older adults). Qiao et al. [68] found higher short-term DE in older adults versus
younger adults, with this difference amplified by perturbation intensity, especially in older

fallers. Bizovska et al. [71] found short-term DE differences ranging from 4% to 18% higher

in older compared to younger females during overground walking, and from 3% to 25%

higher during treadmill walking. However, it is worth noting that older group had a mean age

of only 57.5 years (±4.8), representing middle-aged rather than elderly adults. In their

second study focused on fall risk, Bizovska et al. [72] observed short-term DE increases of 7-

8% when comparing elderly non-fallers to single fallers, while comparisons between non-

fallers and frequent fallers showed highly variable results ranging from 14% decreases to

11% increases depending on measurement axis.

In contrast to the relatively consistent short-term increases, long-term DE exhibited

heterogeneous patterns. The most substantial reductions were observed in clinical

populations: patients with knee osteoarthritis showed decreases ranging from 43% to 85%

compared to controls, while those with chronic musculoskeletal pain demonstrated 19%

reductions. Neuropathy patients showed minimal reductions of 4%. Franz et al. reported

consistent long-term DE reductions in older adults during both normal walking (-10%) and

under visual perturbations (-15%). Despite reporting heterogeneous results, Qiao et al.'s

figures [68] show a trend toward lower long-term DE (-36%) in older participants under

high visual perturbation. The two Bizovska studies revealed higher variability depending on

measurement axis and experimental conditions. In their age-comparison study [71], long-

term DE differences ranged from increases of 56% to decreases of 11% when comparing

older and younger females during overground walking, and from increases of 22% to

decreases of 8% during treadmill walking. Their fall-risk study [72] showed long-term DE

differences ranging from increases of 8% to decreases of 3% when comparing non-fallers to

single fallers, and from decreases of 7% to 18% when comparing non-fallers to frequent

fallers.

Table 3. Results of the externally paced walking studies

Study	Measurement method	Cueing	Short-term DE change with cueing	Long-term DE change with cueing
Dingwell & Cusumano ID 3 [62]	Electrogoniometers (hip, knee, ankle) + sternum tri-axial accelerometer.	Treadmill speed constraint (vs overground).	**Trunk acceleration: -11% to -13% Lower limb kinematics: -11% to -16%**	**Trunk acceleration: -11% to +12% Lower limb kinematics: -11% to -20%**
Terrier & Dériaz ID 25	3D accelerometer on lower back (lumbar), trunk accelerations.	Treadmill speed constraint (vs overground).	**-8% to -9%**	**-5% to -18%**

[55]				
Sejdić et al. ID 29 [46]	Force-sensitive insole (heel strikes) + lower-back 3D accelerometer, trunk accelerations.	Rhythmic sensory cues during overground walking, step duration constraint: auditory, visual, and tactile.	**Visual / step duration:** **-4% to +3%** **Auditory / step duration:** **-11% to -4%** **Tactile / step duration:** **-8% to 0%** **Combined / step duration:** **-12% to -8%**	**Visual / step duration:** **-10% to -18%** **Auditory / step duration:** **-26% to -40%** **Tactile / step duration:** **-15% to -23%** **Combined / step duration:** **-34% to -43%**
Terrier & Dériaz ID 35 [39]	Instrumented treadmill (foot-pressure array); center-of-pressure trajectory.	Rhythmic auditory cueing (metronome), step duration constraint, during treadmill walking.	**-6% to +7%**	**-61% to -86%**
Bizovska et al. ID 46 [71]	Lumbar 3D accelerometer, trunk accelerations.	Treadmill speed constraint vs overground.	**Young:** **-19% to -24%** **Older:** **-16% to -21%**	**Young:** **+24% to +156%** **Older:** **0% to + 100%**
Terrier ID 51 [56]	Instrumented treadmill with force platform; center-of-pressure trajectory.	Visual (step length constraint) and auditory (step duration constraint) synchronization cues, during treadmill walking.	**Visual / step length:** **-4% to -2%** **Auditory / step duration:** **-3% to + 14%**	**Visual / step length:** **-37% to -66%** **Auditory / step duration:** **-43% to -78%**

Percentage changes in short-term and long-term divergence exponents (DE) during
externally paced walking compared to self-paced conditions. Interventions included treadmill
speed constraint (imposing constant velocity versus self-paced overground walking) and
rhythmic sensory cueing (auditory metronome, visual pacing, tactile stimulation). Negative
values indicate reduced DE with external pacing, while positive values indicate increased DE.
Ranges reflect different cueing modalities, measurement signals, or experimental conditions
within studies. Combined cueing indicates simultaneous presentation of multiple sensory
modalities. ID refers to study identification number in the systematic review database (S1
text in supplementary materials).

External cueing studies. Among the 56 studies, six within-subject experiments
examined either externally imposed gait synchronization or a fixed-speed treadmill
constraint. As these interventions likely shift the balance between automatic and voluntary
control of walking [74], they enable evaluation of the sensitivity of long-term DE to variations

in gait automaticity. Three studies implemented synchronization cues: Sejdić et al. [46] applied visual, auditory, tactile, and combined step-duration cues during overground walking using heel-strike detection with a force-sensitive insole and lumbar accelerometry; Terrier & Dériaz [39] and Terrier [56] imposed metronome-based step-duration or visual step-length targets during treadmill walking while recording center-of-pressure trajectories on an instrumented treadmill. Three studies compared constant-speed treadmill with unconstrained overground walking: Dingwell & Cusumano [62] recorded trunk accelerations (sternum) and lower-limb joint angles via electrogoniometers, and Terrier & Dériaz [55] used lower-back accelerometry; Bizovská et al. [71] extended this comparison in young and older women with lumbar accelerometry. Trial durations typically ranged from 3 to 10 min per condition (yielding ~100–500 strides), with speed held constant in cueing paradigms; overground–treadmill contrasts occasionally differed in bout duration or speed (e.g., 5 min overground vs 3 min treadmill in [71]).

Across studies that imposed external synchronization cues (auditory, visual, tactile), short-term divergence exponents (DEs) changed little or inconsistently (approximately −11% to +14%), whereas long-term DEs showed large, systematic reductions ($\approx$ −10% to −86%). In the overground study using rhythmic sensory cues [46], short-term DEs were minimally affected (visual: −4% to +3%; tactile: −8% to 0%) or modestly decreased under auditory and combined cueing (auditory: −11% to −4%; combined: −12% to −8%). By contrast, long-term DEs fell substantially (visual: −10% to −18%; tactile: −15% to −23%; auditory: −26% to −40%; combined: −34% to −43%). Similarly, a substantial decrease in gait complexity measured from force-sensitive insoles (scaling exponent, DFA) was observed (-50% to -13%). Two treadmill experiments that enforced step timing/length converged on the same pattern. Using an instrumented treadmill and center-of-pressure (COP) trajectories, Terrier & Dériaz [39] reported very large long-term DE decreases (−61% to −86%) with only small, direction-dependent short-term changes (−6% to +7%). In a mixed visual–auditory cueing paradigm [56], short-term DEs remained near baseline (visual/step-length: −4% to −2%; auditory/step-duration: −3% to +14%), whereas long-term DEs dropped markedly for both constraints (visual/step-length: −37% to −66%; auditory/step-duration: −43% to −78%).

By contrast, comparing treadmill speed constraint with unconstrained overground walking yielded consistent short-term DE reductions but heterogeneous long-term responses across datasets and signal types. In three studies that contrasted fixed-speed treadmill with overground walking, short-term DEs were lower on the treadmill in trunk accelerations ([62]: −11% to −13%, [55]: −8% to −9%, and [71] −16% to −24%), and lower-limb

677 kinematics ([62]: -11% to -16%). Long-term DEs, however, varied: they mostly decreased
678 in two datasets ([62]: trunk -11% to $+12\%$, lower-limb -11% to -20%; [55]: -5% to
679 -18%), but substantially increased in [71] ($+24\%$ to $+156\%$).
680

Meta-analysis

We synthesized six independent datasets reporting the association between long-term DE
and the scaling exponent from DFA. Measurement modalities spanned instrumented
treadmills with center-of-pressure trajectories, trunk accelerometry, and segmental
kinematics of the head and lower limbs; several experiments also varied speed (slow,
preferred, fast) or imposed synchronization tasks. One publication contributed two distinct
datasets because the same participants walked in separate overground and treadmill
experiments [55]; these were treated as independent due to the large contextual difference
between conditions. Where articles reported axis- or segment-specific associations, we
retained those dependent effects (e.g., anteroposterior, mediolateral, vertical, or head vs
ankle). This yielded 14 dependent correlations across the six datasets (k=14), while speed
manipulations within a given protocol were not counted as separate effects when
correlations were reported on the pooled observations [48].
Using a correlated-effects random-effects model with cluster-robust variance estimation,
Fisher's Z-transformed correlations were pooled and then back-transformed for
interpretation. The overall association was positive, with a pooled correlation of 0.64 (95%
CI 0.34 to 0.82) based on 14 effects (N=582 condition-level observations from 209
participants). Heterogeneity was considerable ($I^2 \approx 82\%$), with estimated between-study
variance $\tau^2 \approx 0.22$ and within-study variance $\omega^2 \approx 0.011$. Leave-one-out influence diagnostics
showed that the pooled correlation remained stable in magnitude and direction. Re-
estimating the model after omitting each dataset in turn yielded pooled r values between
0.57 and 0.71. Across all leave-one-out fits, the 95% confidence intervals stayed positive,
and the standard error varied only modestly (0.17–0.24). The forest plot summarizes study-
level estimates and the model summary (Fig 6).
705

Discussion

707 This systematic review synthesized evidence from 56 studies to critically examine the use of
708 long-term divergence exponents (DE) computed by Rosenstein's algorithm in human gait
709 analysis, with particular focus on its reinterpretation as a measure of gait complexity and
710 automaticity rather than stability.

Summary of main findings

Our meta-analysis (Fig 6) provides empirical validation for the theoretical reconceptualization of long-term DE as a complexity measure. The substantial positive correlation (r = 0.64) between long-term DE and the DFA scaling exponent demonstrates that these metrics capture a shared underlying construct—the fractal-like correlation structure of stride-to-stride fluctuations. This association persisted robustly across heterogeneous measurement contexts. Two additional studies [33,46] showed parallel reductions in both long-term DE and scaling exponent during visual perturbation and external cueing, qualitatively supporting the meta-analytic correlation despite not reporting correlation statistics.

The differential responsiveness of short-term and long-term DE across experimental designs provides further support for their distinct physiological interpretations. Perturbation studies (Table 1) revealed a consistent dissociation: destabilizing interventions systematically increased short-term DE while decreasing long-term DE, confirming that these indices may reflect separable control processes underlying gait. This pattern directly contradicts the interpretation of both measures as stability metrics, since a truly destabilizing intervention should increase both if they assessed the same construct. Between-subject comparisons (Table 2) yielded more heterogeneous patterns, with clinical populations sometimes exhibiting paradoxical long-term DE responses (both increases and decreases relative to controls), further highlighting its dissociation from stability measures that should consistently elevate with pathology. Finally, external cueing studies (Table 3) demonstrated that interventions requiring conscious temporal or spatial coordination—which engage voluntary control without perturbing balance—produced strong long-term DE reductions (up to -86%) with minimal short-term DE changes, as anticipated when gait automaticity is reduced without compromising stability. This selective sensitivity to sensory cueing, combined with the parallel reductions observed in DFA scaling exponents [74], further support long-term DE as a marker of gait complexity rather than stability.

Long-term divergence saturation as a signature of reduced gait complexity

Divergence in strange attractors. Rosenstein's algorithm was originally developed to estimate the maximal Lyapunov exponent in deterministic chaotic systems. The method's computational validity was established using well-characterized strange attractors, including the three-dimensional Lorenz system—a nonlinear dynamical system defined by three coupled differential equations that exhibits sensitive dependence on initial conditions [75].

When applied to such chaotic systems, the logarithmic divergence curve exhibits
characteristic behavior: an initial linear region from which the maximal Lyapunov exponent is
computed, followed by a plateau where further divergence ceases (see Appendix A in [39]).
This saturation occurs because strange attractors, despite their complex internal dynamics,
occupy a bounded region of phase space, constraining the maximum separation between
initially neighboring trajectories.

Divergence in gait dynamics attractors. Dynamical systems models suggest that
human gait may exhibit periodic limit cycle attractor dynamics with stochastic perturbations,
rather than strange attractors characteristic of deterministic chaos [15,38,76], a
characterization supported by both discrete stride interval analyses [77] and continuous
signal phase space reconstruction [76]. This distinction justifies the preference for
"divergence exponent" over "Lyapunov exponent" when describing gait dynamics [15]. The
divergence curves obtained from gait attractors reflect departure from purely chaotic
systems: they typically display an initial exponential divergence phase followed by gradual
saturation over subsequent strides but often lack the abrupt plateau characteristic of purely
deterministic chaotic systems with bounded strange attractors (S2 Figures).

Fractal stride patterns and phase space exploration. A gait attractor constructed
using Takens theorem or an alternative approach represents the repertoire of available
coordination patterns—the different kinematic combinations and their temporal
organization—with phase-space exploration describing how trajectories navigate through this
repertoire across consecutive gait cycles [78]. It can be hypothesized that the temporal
correlation structure of consecutive strides modulates how trajectories explore the phase
space, thereby influencing divergence curve saturation and hence long-term DE values. Low-
frequency variations distributed across dozens of consecutive strides may create drift
through multiple regions of the attractor, delaying the saturation process and yielding higher
long-term DE values. This mechanism may explain the observed correlation between long-
term DE and DFA scaling exponents. In support of this hypothesis, Terrier and Reynard [51]
manipulated stride interval correlation structure while preserving acceleration signal shape,
demonstrating that long-term DE scaled systematically with temporal complexity while short-
term DE remained unchanged. Divergence curves from correlated stride patterns exhibited
delayed saturation compared to constant or random patterns (S2 Document), providing
evidence that fractal temporal structure prevents premature saturation of phase space
exploration, thereby elevating long-term DE values. Consistent with this mechanism, two
included studies [79,80] that applied stride-level time normalization by resampling each

stride to a fixed number of data points (as in [51]) reported markedly suppressed long-term

DE values ($\lambda L \approx 0.00$ in [79]; no significant condition effects in [80]).

Attention-dependent control mechanisms. The selective reduction of long-term DE

during external pacing (Table 3) points to modifications in motor control strategy.

Sensorimotor synchronization with rhythmic cues induces error correction processes that

adjust each stride's timing to match external targets [34]. This correction mode contrasts

with the loosely coupled, endogenous control of self-paced walking [81]. The corrections

around target pace values generate oscillations at short timescales while suppressing the

low-frequency variations that characterize normal gait [82]. In fractal analysis terms (DFA),

short-term oscillations manifest as anti-persistent temporal structure, whereas low-frequency

variations represent the persistent, long-range correlations that define high-complexity gait

dynamics. Experimental evidence shows that indeed metronome walking induces anti-

persistent stride interval dynamics [74,83,84]. This transformation from persistent to anti-

persistent temporal structure constrains phase space exploration—trajectories converge more

rapidly within the attractor—thereby accelerating divergence curve saturation and reducing

long-term DE values. This is evident in divergence curves published in [39,56] (see S2

Figures). The reductions in both long-term DE and DFA scaling exponents during cued

walking [46,50] thus reflect a common underlying mechanism: the alteration of fractal

correlation structure through attention-dependent error correction mechanisms.

Dual constraint effects. The minimal sensitivity of long-term DE to constant-speed

treadmill constraint (Table 3) reflects parameter-specific alterations in temporal correlation

structure. Evidence from DFA demonstrates that treadmill walking selectively induces anti-

persistent dynamics in stride-speed fluctuations while preserving persistent correlations in

both stride intervals and stride lengths [74]. This selective transformation arises because

maintaining constant speed requires coordinated adjustments between stride time and stride

length—participants modulate both parameters in tandem to prevent positional drift, thereby

tightening speed control without imposing rigid constraints on either time series individually

[34]. DEs computed from center-of-pressure trajectories or trunk acceleration signals likely

capture a global fluctuation pattern, rather than the parameter-specific correlation structures

revealed by DFA of individual time series. When only stride-speed correlations become anti-

persistent while stride interval and stride length correlations remain persistent, this partial

disruption produces moderate effects on the global attractor dynamics quantified by long-

term DE. This interpretation is supported by the graded effects of constraint combinations

observed in Table 3. Metronome synchronization during overground walking—which

selectively modifies stride interval correlations while stride length and speed patterns remain

persistent [83]—produces moderate long-term DE reductions (-26% to -40% in [46], -12%
to -29% in [50]). In contrast, superimposing metronome pacing on treadmill walking
generates larger reductions (−66% to −86%, Table 3), reflecting combined modification of
both stride interval and stride length correlation structures. This graded response
demonstrates that dual constraints reorganize attractor dynamics more profoundly than
either single constraint alone, consistent with the parameter-specific nature of correlation
structure alterations captured by DFA.

Long-term divergence as an index of attractor complexity. Taken together, this
evidence suggests that long-term DE reflects gait complexity rather than stability: it
correlates strongly with fractal scaling exponents, responds systematically to manipulation of
stride interval correlation structure while short-term DE remains unchanged, and exhibits
graded sensitivity to constraint combinations that selectively modify correlations in different
gait parameters. These findings support reinterpreting long-term DE as an "Attractor
Complexity Index" that captures how temporal correlation structure shapes phase space
exploration—a measure reflecting the adaptive flexibility of motor control rather than
resistance to perturbations.

Long-term divergence and gait automaticity

Among the studies identified in this review, only six met methodological criteria for valid
between-subject comparisons (Table 2), having employed sufficiently long walking bouts and
time-normalized signals to ensure reliable long-term divergence exponent estimation. The
remaining clinical comparison studies, while potentially informative, warrant cautious
interpretation. Within these methodologically rigorous studies, long-term DEs exhibited
heterogeneous patterns across clinical populations. While short-term divergence exponents
consistently increased in patients relative to controls, long-term divergence exponents
showed both reductions and increases, which requires specific analysis separated by
conditions and pathologies.

Pain. Two included studies examining pain-related conditions converged on a
consistent pattern: patients with chronic lower-limb pain [56] and those with knee
osteoarthritis [73], demonstrated reduced long-term divergence. An earlier longitudinal study
of knee osteoarthritis patients ([85], N=16) awaiting replacement surgery provides additional
support for pain-related complexity reductions. Using only 30 strides per analysis, the study
nevertheless used global time normalization and generalized estimating equations that
appropriately modeled speed-dependent effects. At a clinically relevant speed of 3 km/h, the
affected limb demonstrated a dissociation between stability and complexity measures: short-

term divergence exponents were unchanged between patients and controls, while long-term
divergence exponents showed 19% reduction.

Pain may induce a shift toward more controlled, less automatic gait characterized by
greater attentional involvement in motor planning [36]. It can be assumed that pain-related
attentional focus on bodily sensations and movement execution constrains the natural stride-
to-stride variability that characterizes automatic walking.

Peripheral Neuropathy. Only one study met methodological criteria for valid between-
subject comparison (Table 2). However, the foundational work on this topic comes from the
two earlier studies by Dingwell and colleagues [40,86] examining the same cohort of diabetic
neuropathy patients (N=26). While these studies used sufficiently long walking bouts (10
minutes continuous), they did not use time-normalized signals. Nevertheless, they observed
a consistent pattern: patients with diabetic neuropathy showed reduced long-term
divergence exponents compared to controls (-26%). The pattern observed by Manor et al.
[69] using 100 consecutive strides with time-normalized signals confirmed this finding with a
4% reduction in long-term DE. In contrast, the within-subject study that examined acute
plantar desensitization in healthy adults showed an opposite effect: both short-term and
long-term DE increased ([63], Table 1). Such a result may point either to differential acute
and habituation effects on gait dynamics or to heterogeneity in study design and
measurement protocols.

Meta-analyses show that neuropathy patients walk more cautiously, with slower walking
speed, shorter steps, and, and longer stance time [87,88]. Reduced plantar sensory
information may make gait control more dependent on cognitive resources in diabetic
patients compared to controls [88]. This higher cognitive demand is reflected in poorer dual-
task performance [89]. Notably, these cautious gait adaptations are further amplified by
cognitive impairment in diabetic patients, independent of neuropathy status [90]. The
observed reduction in long-term DEs may therefore reflect a compensatory shift toward a
greater conscious control of gait characterized by a reduced reliance on automatic motor
control.

Aging and fall risk. Three studies examining young versus older adults were excluded
from this review due to methodological limitations (Tables B and C in S1 Text). These studies
generally showed trends toward increased short-term and long-term DE in older populations
[91], though the long-term DE increases rarely reached statistical significance [92,93].
Among the three included studies (Table 2), heterogeneous results and high variability in
long-term DE make difficult a clear interpretation of age-related effects. Franz et al. [33] and
Qiao et al. [68] showed consistent reductions in long-term DE in older adults, particularly

under highly visually perturbed conditions. In contrast, Bizovska et al. [71] reported variable

results across measurement directions and conditions. Comparisons between older fallers

and non-fallers in [72] yielded inconclusive results for long-term DE, although a notable

contrast emerged when comparing frequent fallers to non-fallers (−7% to −18% reduction).

To contextualize these findings within recent literature, the 2024 study included in the meta-

analysis [50] identified a significant and substantial reduction in long-term DE when

comparing older adults to younger controls (−21%).

Age-related decline in gait automaticity is well-documented [25], with older adults

demonstrating increased prefrontal cortex activation during usual walking compared to

standing [94], though patterns of cortex recruitment during cognitively demanding walking

may differ from younger adults [95]. This shift from automatic to more controlled gait

regulation has been attributed to age-related changes in neural networks, where diminished

function in cerebellum and basal ganglia regions necessitates greater compensatory

engagement of attention-related cortical networks, particularly the prefrontal cortex [96–98].

The observed reductions in long-term DE in older adults, particularly under conditions that

increase attentional demands such as visual perturbations, may therefore reflect this

transition toward less automatic gait control.

Perturbation studies. Neuroimaging studies provide direct support for the hypothesis

that perturbations increase cortical engagement and decrease gait automaticity. Mechanical

perturbations such as unpredictable slips during treadmill walking induce significant increases

in prefrontal cortex activation measured by functional near-infrared spectroscopy [99]. Visual

perturbations using manipulated optic flow similarly show enhanced frontal and parietal

cortical activation, indicating increased attentional demands to maintain gait stability under

challenging sensory conditions [100]. A recent systematic review synthesizing 35

neuroimaging studies of reactive balance concluded that cortical activation during

perturbations reflects higher sensory processing, error detection, and planning of

compensatory motor responses [101]. These neuroimaging results directly support the

biomechanical findings from within-subject perturbation studies in this review (Table 1:

experimental manipulations that likely increase cortical activation also consistently increase

short-term divergence exponents—reflecting higher gait instability—while simultaneously

decreasing long-term divergence exponents, reflecting greater attentional involvement in gait

control. The convergence of evidence across multiple experimental approaches, clinical

populations, measurement modalities, and independent research groups strongly supports

the interpretation that long-term divergence exponents reflect the degree of gait

automaticity, with lower values indicating a shift toward more cognitively-controlled, less

automatic locomotor patterns.

Methodological considerations

Measurement methods. The included studies employed diverse measurement approaches

(Fig 3), with signals derived from whole-body motion capture systems, wearable

accelerometry, and center-of-pressure trajectories. Despite this methodological

heterogeneity, examination of response patterns across experimental manipulations reveals

consistent findings, suggesting that long-term DE capture aspects of gait control that

transcend specific measurement modalities. The critical prerequisite is likely that the

measured signal captures the quasi-periodic gait signature—the rhythmic, repeating pattern

of locomotor cycles with their inherent stride-to-stride variations. Whether these dynamics

are sampled through trunk accelerations, joint angle trajectories, or center-of-pressure

displacements, the resulting measures converge on the same underlying phenomenon: the

temporal organization of consecutive gait cycles.

Among the measurement approaches employed, trunk-level assessment offers particular

advantages for capturing global gait coordination patterns. A majority of included studies

(31/56, 55%) prioritized trunk measurements at the pelvis, lumbar, cervical, or thoracic spine

levels. Dingwell and Kang [76] emphasized that trunk kinematics provide a comprehensive

view of locomotor control because trunk movements reflect the integrated outcome of

bilateral limb coordination and postural adjustments required to maintain upright

progression. For assessments targeting the temporal organization of overall locomotor

patterns, trunk-level measurements therefore represent the preferred approach, offering an

integrated readout of stride-to-stride control dynamics that reflects whole-body coordination

rather than isolated joint-level adjustments.

Measurement duration. Assessment of long-term DE as complexity measures

imposes specific data duration requirements that differ from stability assessments. The basis

for these requirements arises from the multiscale nature of gait complexity. Low-frequency

variations—the signature of persistent temporal correlations—manifest only across extended

observation windows. Truncated recordings artificially constrain the observable correlation

timescales, effectively filtering out the very low-frequency components. This explains why

studies employing shorter recordings (fewer than 50 strides) may capture local stability

properties adequately but potentially mischaracterize attractor complexity.

Among the 56 included studies, recording durations varied considerably, ranging from as few
as 30 strides to over 500 strides (Fig 4). This heterogeneity in measurement duration
represents a potential source of methodological variance that may have influenced reported
findings, particularly in between-subject comparisons where the reliability of individual
estimates critically determines statistical power to detect group differences (Table 2).
Methodological investigations within the included studies provide empirical guidance for
minimum data requirements. Bruijn et al. [102] showed that divergence exponent precision
improves markedly with recording length, identifying approximately 150 consecutive strides
as necessary for adequate measurement precision. Reynard and Terrier [60] confirmed these
requirements through systematic reliability analysis, showing that 70-stride estimates yielded
substantially higher reliability than 35-stride estimates for long-term DE. Riva et al. [103]
extended this work through within-session reliability assessment across 11 gait measures,
establishing stride-specific thresholds for achieving target precision levels. Their findings
revealed that long-term divergence measures required among the highest stride counts
(110+ strides for 50% reliability threshold) compared to other stability metrics.
A 2020 systematic review on DFA methodology [104] established consensus guidelines
recommending at least 500-600 consecutive strides for reliable estimation of scaling
exponents, acknowledging that shorter recordings may inadequately capture the long-range
correlation structure that defines fractal gait dynamics. The more modest stride requirements
for long-term DE (150-200 strides) compared to DFA's 500-600 stride recommendation
suggest that the DE approach may offer practical advantages for clinical gait assessment,
requiring shorter recording durations while still capturing essential features of temporal
correlation structure through phase space reconstruction [49]. Future investigations
employing long-term divergence exponents as complexity measures should prioritize
recording durations of at least 150-200 consecutive strides to ensure adequate precision for
detecting meaningful differences while maintaining practical feasibility for clinical
populations.
Data length effects and time normalization. Beyond precision considerations,
measurement duration influences DE magnitudes. Bruijn et al. [102] demonstrated through
systematic bootstrap analysis that both short-term and long-term DEs increase with data
series length. As recording duration extends, the phase space trajectory explores increasingly
distant regions of the attractor, allowing the algorithm to identify nearest neighbors
separated by larger distances. These more distant neighbor pairs, representing trajectories
sampling different attractor regions, exhibit different divergence characteristics than the
tightly clustered neighbors identified in shorter recordings. Reynard and Terrier confirmed

this effect in a larger sample (N=95) [60]. Consequently, standardizing recording duration

across experimental conditions and participant groups constitutes a critical methodological

requirement, as divergence exponent differences attributable to unequal measurement

lengths may confound or obscure true physiological differences between populations or

conditions. Time normalization became increasingly standardized in recent years, with most

high-quality studies reporting explicit normalization protocols (Table C in S1 text]). Earlier

studies occasionally analyzed time-based signals without normalization, potentially

introducing systematic bias.

Study quality and risk of bias

Analytical rigor and methodological transparency. Quality assessment revealed good

methodological transparency across the included studies (S3 Table), with comprehensive

reporting of experimental protocols, measurement systems, signal processing procedures,

and algorithm parameters necessary for study replication. This result may indicate that the

field has achieved methodological maturity, with recent studies providing sufficient detail to

enable independent replication. However, it is important to acknowledge that these favorable

quality metrics partly reflect our inclusion criteria, which required studies to provide

adequate methodological information to confirm use of Rosenstein's algorithm for long-term

DE computation.

Outcome reporting: descriptive statistics. While analytical procedures were

generally well-documented, the reporting of study outcomes showed considerable variability.

A substantial proportion of studies, particularly those published before 2013, presented

results exclusively through figures without accompanying numerical tables containing means

and measures of variability. This reporting approach necessitated manual extraction of

approximate values using specialized digitization software to enable quantitative synthesis

and calculation of percentage changes between conditions or groups (as presented in Tables

2 and 3). The reliance on visual estimation from figures introduces measurement error and

limits the precision of secondary analyses. Moreover, some studies reported only qualitative

descriptions of differences or provided statistical test results without the underlying

descriptive statistics, making meta-analytical synthesis unfeasible. A clear temporal

improvement emerged after 2013, with recent studies increasingly providing complete

descriptive statistics for all experimental conditions.

Outcome reporting: effect sizes and clinical significance. Beyond descriptive

statistics, a more persistent limitation concerned the reporting and interpretation of effect

sizes. The majority of studies across all time periods relied primarily on p-values and significance testing to characterize findings, with only 18% achieving full scores through comprehensive effect size reporting. Many studies reported results dichotomously as "significant" or "non-significant" without quantifying the magnitude of observed differences or discussing their potential clinical relevance. This approach limits the ability to distinguish between statistically significant but trivially small effects and meaningfully large effects that may warrant clinical attention. The absence of standardized effect size measures (such as Cohen's d, confidence intervals, or percentage differences with precision estimates) also constrains meta-analytic synthesis and prevents assessment of whether effects are consistent or variable across studies and populations. While some recent studies have begun incorporating effect size reporting and discussing practical significance, this remains an underdeveloped aspect of reporting standards in the field.

Sample size considerations and recruitment practices. Domain 3 (sample size adequacy) achieved an average score of 1.9 out of 3 points, reflecting the modest sample sizes typical of exploratory biomechanical research (S3 Table). The median sample size across studies was 20 participants, with most studies enrolling fewer than 50 individuals. Given the exploratory and methodological nature of much of this research, these sample sizes are not inappropriate, particularly considering that most studies used within-subjects designs and repeated measures that enhance statistical power relative to between-subjects comparisons. However, only a small minority of studies (<5%) provided explicit sample size justifications or reported formal power calculations, making it difficult to assess whether samples were systematically adequate for detecting meaningful effects. A more substantive concern relates to recruitment practices and sample representativeness. Although rarely discussed explicitly in the reviewed articles, convenience sampling appeared widespread, with many studies likely recruiting from institutional populations such as university students or laboratory staff members. This practice may introduce systematic biases, as such samples tend to overrepresent younger, healthier, more educated, and potentially more physically active individuals compared to the general population. For studies comparing healthy controls to clinical populations, the use of convenience samples may exaggerate group differences if controls represent an unusually homogeneous and high-functioning subset of the healthy population. While these recruitment practices are understandable during the methodological development phase of a field, the transition toward clinical application requires more systematic attention to sample representativeness, demographic diversity, and explicit justification of sample sizes relative to the precision needed for clinical decision-making.

1055

Limitations of the systematic review

1056

Study identification and selection limitations. This systematic review has several

1057

limitations that should be acknowledged. First, the favorable quality assessment results

1058

partly reflect selection bias inherent to our inclusion criteria. Studies were required to provide

1059

sufficient methodological detail, resulting in exclusion of less transparently reported research

1060

during full-text screening. Second, while our database searches applied no language

1061

restrictions to ensure comprehensive literature retrieval, our protocol specified that full-text

1062

screening would be limited to English and French publications based on the linguistic

1063

competencies of our review team. The language restriction remained theoretical rather than

1064

operational, as no relevant non-English/French studies required exclusion. Nevertheless, we

1065

acknowledge that future updates to this review would benefit from broader linguistic

1066

inclusion with appropriate translation resources to ensure maximum comprehensiveness.

1067

Third, our focus on peer-reviewed journal articles resulted in exclusion of grey literature

1068

including doctoral dissertations, conference abstracts, and technical reports, which may

1069

contain relevant methodological developments or findings not yet published in traditional

1070

venues. Fourth, publication bias may affect our synthesis, as studies reporting null findings

1071

or unsuccessful applications of the method may be less likely to appear in peer-reviewed

1072

literature. This bias may have been particularly pronounced given that most studies

1073

interpreted long-term DE as a stability measure: researchers observing paradoxical

1074

patterns—such as long-term DE reductions during destabilizing perturbations (Table 1)—may

1075

have regarded such findings as methodological failures rather than meaningful results,

1076

leading to selective non-reporting of studies with theoretically inconsistent outcomes under

1077

the traditional stability interpretation.

1078

Synthesis limitations. The scope of quantitative synthesis was constrained by several

1079

factors. Despite identifying 56 relevant studies, comprehensive meta-analysis across all

1080

research questions was not feasible due to heterogeneous outcome reporting, particularly in

1081

earlier studies that presented results exclusively through figures without numerical tables.

1082

Only four studies provided the necessary correlation data for our meta-analysis examining

1083

associations between long-term DE and DFA scaling exponents, limiting the precision of this

1084

key finding. Similarly, only six studies met methodological criteria for valid inclusion in

1085

between-subject clinical comparisons (Table 2), as most clinical studies used insufficiently

1086

long recording durations or lacked time normalization procedures necessary for reliable

1087

complexity assessment. The diverse measurement contexts—including different walking

1088

conditions (treadmill versus overground), signal types (kinematics, accelerometry, center-of-
pressure), and body locations—while potentially strengthening the generalizability of
findings, precluded direct quantitative comparison of effect magnitudes across studies. These
synthesis limitations reflect the exploratory developmental stage of the field rather than
fundamental obstacles to future evidence accumulation, as recent improvements in reporting
standards suggest that subsequent systematic reviews will be better positioned for
comprehensive meta-analytic synthesis.

Future directions and clinical translation

Research priorities and methodological advances. Several methodological priorities
emerge from this review to advance the field toward clinical application. First,
standardization of measurement protocols should prioritize recording durations of at least
150-200 consecutive strides for complexity assessment. Second, theoretical development
remains incomplete. Establishment of clinically meaningful change thresholds for long-term
DE is essential—what magnitude of reduction indicates functionally significant loss of gait
automaticity? Development of age-stratified and population-specific normative reference
values would enable interpretation of individual measurements relative to expected ranges.
Third, validation of the complexity interpretation requires experimental studies that directly
manipulate automaticity (e.g., dual-task paradigms, attention-demanding environments)
while simultaneously measuring long-term DE and neuroimaging markers of cortical
engagement. Finally, longitudinal research tracking long-term DE changes across disease
progression, rehabilitation, or aging trajectories would clarify whether complexity measures
predict clinically relevant outcomes such as fall risk, functional decline, or treatment
response. Multi-site replication studies ensuring generalizability across diverse populations
and measurement contexts remain particularly valuable given the current concentration of
research within specialized laboratories.

Clinical translation and practical applications. Long-term DE holds promise as a
complementary clinical measure that captures aspects of motor control not reflected in
traditional gait metrics such as speed or stride length variability. Its potential clinical utility
spans several domains. For diagnostic applications, reduced long-term DE may identify early-
stage neurological changes characterized by diminished gait automaticity before gross
functional impairments become apparent. In rehabilitation contexts, long-term DE could
monitor recovery of automatic motor control following neurological injury or track treatment
effects where restoration of natural gait patterns represents a therapeutic goal. For fall risk
assessment, integration of complexity measures with conventional stability metrics may

improve risk stratification. The method offers practical advantages over alternative

complexity measures such as DFA: the more modest recording requirements (150-200 versus

500-600 strides) enhance feasibility for clinical populations with limited walking tolerance.

Long-term DE should complement rather than replace existing clinical gait assessment tools,

contributing one element within comprehensive evaluation protocols. With these foundations

established, wearable sensor technologies could enable free-living complexity monitoring,

extending assessment beyond laboratory settings to capture gait automaticity in natural

environments where attentional demands and environmental challenges reflect real-world

function.

1132

1133

Conclusion

This systematic review provides converging evidence that long-term DE measure gait
complexity and automaticity rather than stability. The substantial correlation with DFA scaling
exponents (r = 0.64), combined with systematic dissociation from short-term stability
measures during perturbations and cueing, supports reinterpreting this metric as the
Attractor Complexity Index—a marker of the temporal organization and attentional demands
of locomotor control.
This paradigm shift has practical implications. Rather than duplicating information from
short-term stability measures, long-term divergence may uniquely capture motor-cognitive
aspects of gait control relevant to fall risk, disease progression, and rehabilitation
effectiveness. Establishing this measure as a complementary biomarker for gait automaticity
would enhance multidimensional assessment of locomotor function in clinical populations.
1146

Acknowledgments

1148

1149 We thank Marco Pedrotti and Florence Waelchli for their assistance with administrative tasks.

1150

1151

REFERENCES

1.   Perry J. Gait Analysis: Normal and Pathological Function, Second Edition. 2nd ed.
Thorofare: SLACK, Incorporated; 2010.

2.   Singh SJ, Puhan MA, Andrianopoulos V, Hernandes NA, Mitchell KE, Hill CJ, et al. An
official systematic review of the European Respiratory Society/American Thoracic
Society: measurement properties of field walking tests in chronic respiratory disease.
Eur Respir J. 2014;44: 1447–1478. doi:10.1183/09031936.00150414

3.   Giannitsi S, Bougiakli M, Bechlioulis A, Kotsia A, Michalis LK, Naka KK. 6-minute walking
test: a useful tool in the management of heart failure patients. Ther Adv Cardiovasc Dis.
2019;13: 1753944719870084. doi:10.1177/1753944719870084

4.   Yogev G, Hausdorff JM, Giladi N. The Role of Executive Function and Attention in Gait.
Mov Disord. 2008;23: 329–472. doi:10.1002/mds.21720

5.   Perera S, Patel KV, Rosano C, Rubin SM, Satterfield S, Harris T, et al. Gait Speed Predicts
Incident Disability: A Pooled Analysis. The Journals of Gerontology: Series A. 2016;71:
63–71. doi:10.1093/gerona/glv126

6.   Montero-Odasso M, Verghese J, Beauchet O, Hausdorff JM. Gait and Cognition: A
Complementary Approach to Understanding Brain Function and the Risk of Falling.
Journal of the American Geriatrics Society. 2012;60: 2127–2136. doi:10.1111/j.1532-
5415.2012.04209.x

7.   Allali G, Launay CP, Blumen HM, Callisaya ML, De Cock A-M, Kressig RW, et al. Falls,
Cognitive Impairment, and Gait Performance: Results From the GOOD Initiative. J Am
Med Dir Assoc. 2017;18: 335–340. doi:10.1016/j.jamda.2016.10.008

8.   Verghese J, Wang C, Lipton RB, Holtzer R, Xue X. Quantitative gait dysfunction and risk
of cognitive decline and dementia. Journal of Neurology, Neurosurgery & Psychiatry.
2007;78: 929–935. doi:10.1136/jnnp.2006.106914

9.   Dommershuijsen LJ, Isik BM, Darweesh SKL, van der Geest JN, Ikram MK, Ikram MA.
Unraveling the Association Between Gait and Mortality-One Step at a Time. J Gerontol A
Biol Sci Med Sci. 2020;75: 1184–1190. doi:10.1093/gerona/glz282

10.   Tinetti ME. Performance-Oriented Assessment of Mobility Problems in Elderly Patients.
Journal of the American Geriatrics Society. 1986;34: 119–126. doi:10.1111/j.1532-
5415.1986.tb05480.x

11.   Hulleck AA, Menoth Mohan D, Abdallah N, El Rich M, Khalaf K. Present and future of
gait assessment in clinical practice: Towards the application of novel trends and
technologies. Front Med Technol. 2022;4. doi:10.3389/fmedt.2022.901331

12.   Sprott JC. Chaos and time-series analysis. Oxford□; New York: Oxford University Press;
2003.

13.   Stergiou N, Harbourne R, Cavanaugh J. Optimal movement variability: a new theoretical
perspective for neurologic physical therapy. J Neurol Phys Ther. 2006;30: 120–129.
doi:10.1097/01.npt.0000281949.48193.d9

14.   Moghtadaei M, Dorey TW, Rose RA. Evaluation of non-linear heart rate variability using
multi-scale multi-fractal detrended fluctuation analysis in mice: Roles of the autonomic
nervous system and sinoatrial node. Front Physiol. 2022;13.
doi:10.3389/fphys.2022.970393

15.   Dingwell JB. Lyapunov Exponents. Wiley Encyclopedia of Biomedical Engineering. John
Wiley & Sons, Ltd; 2006. doi:10.1002/9780471740360.ebs0702

16.   Hausdorff JM, Mitchell SL, Firtion R, Peng CK, Cudkowicz ME, Wei JY, et al. Altered
fractal dynamics of gait: reduced stride-interval correlations with aging and
Huntington's disease. J Appl Physiol. 1997;82: 262–269.
doi:10.1152/jappl.1997.82.1.262

17.   Yentes JM, Raffalt PC. Entropy analysis in gait research – methodological considerations
and recommendations. Ann Biomed Eng. 2021;49: 979–990. doi:10.1007/s10439-020-
02616-8

18.   Lukšys D, Griškevičius J. Application of nonlinear analysis for the assessment of gait in
patients with Parkinson's disease. Technol Health Care. 2022;30: 201–208.
doi:10.3233/THC-219003

19.   Lahmiri S. Gait Nonlinear Patterns Related to Parkinson's Disease and Age. IEEE
Transactions on Instrumentation and Measurement. 2019;68: 2545–2551.
doi:10.1109/TIM.2018.2866316

20.   Amirpourabasi A, Lamb SE, Chow JY, Williams GKR. Nonlinear Dynamic Measures of
Walking in Healthy Older Adults: A Systematic Scoping Review. Sensors. 2022;22: 4408.
doi:10.3390/s22124408

21.   Freitas M, Pinho F, Pinho L, Silva S, Figueira V, Vilas-Boas JP, et al. Biomechanical
Assessment Methods Used in Chronic Stroke: A Scoping Review of Non-Linear
Approaches. Sensors. 2024;24: 2338. doi:10.3390/s24072338

22.   Stergiou N, editor. Nonlinear analysis for human movement variability. Boca Raton
London New York: CRC Press; 2016.

23.   Zanin M, Olivares F, Pulido-Valdeolivas I, Rausell E, Gomez-Andres D. Gait analysis
under the lens of statistical physics. Computational and Structural Biotechnology
Journal. 2022;20: 3257–3267. doi:10.1016/j.csbj.2022.06.022

24.   Stuss DT, Levine B. Adult clinical neuropsychology: lessons from studies of the frontal
lobes. Annu Rev Psychol. 2002;53: 401–433.
doi:10.1146/annurev.psych.53.100901.135220

25.   Clark DJ. Automaticity of walking: functional significance, mechanisms, measurement
and rehabilitation strategies. Front Hum Neurosci. 2015;9: 246.
doi:10.3389/fnhum.2015.00246

26.   Crowley P, Vuillerme N, Samani A, Madeleine P. The effects of walking speed and mobile
phone use on the walking dynamics of young adults. Sci Rep. 2021;11: 1237.
doi:10.1038/s41598-020-79584-5

27.   Walsh GS. Dynamics of Modular Neuromotor Control of Walking and Running during
Single and Dual Task Conditions. Neuroscience. 2021;465: 1–10.
doi:10.1016/j.neuroscience.2021.04.004

28.   Ahmadi S, Wu C, Sepehri N, Kantikar A, Nankar M, Szturm T. The Effects of Aging and
Dual Tasking on Human Gait Complexity During Treadmill Walking: A Comparative Study
Using Quantized Dynamical Entropy and Sample Entropy. J Biomech Eng. 2018;140.
doi:10.1115/1.4037945

29.   Lamoth CJ, van Deudekom FJ, van Campen JP, Appels BA, de Vries OJ, Pijnappels M.
Gait stability and variability measures show effects of impaired cognition and dual
tasking in frail people. J Neuroeng Rehabil. 2011;8: 2. doi:10.1186/1743-0003-8-2

30.   Tajali S, Mehravar M, Negahban H, van Dieën JH, Shaterzadeh-Yazdi M-J, Mofateh R.
Impaired local dynamic stability during treadmill walking predicts future falls in patients
with multiple sclerosis: A prospective cohort study. Clin Biomech (Bristol). 2019;67:
197–201. doi:10.1016/j.clinbiomech.2019.05.013

31.   Hove MJ, Suzuki K, Uchitomi H, Orimo S, Miyake Y. Interactive rhythmic auditory
stimulation reinstates natural 1/f timing in gait of Parkinson's patients. PLoS One.
2012;7: e32600. doi:10.1371/journal.pone.0032600

32.   Schaefer RS. Auditory rhythmic cueing in movement rehabilitation: findings and possible
1248   mechanisms. Philos Trans R Soc Lond B Biol Sci. 2014;369: 20130402.
1249   doi:10.1098/rstb.2013.0402

1250   33.   Franz JR, Francis CA, Allen MS, O'Connor SM, Thelen DG. Advanced age brings a
1251   greater reliance on visual feedback to maintain balance during walking. Human
1252   Movement Science. 2015;40: 381–392. doi:10.1016/j.humov.2015.01.012

1253   34.   Roerdink M, de Jonge CP, Smid LM, Daffertshofer A. Tightening Up the Control of
1254   Treadmill Walking: Effects of Maneuverability Range and Acoustic Pacing on Stride-to-
1255   Stride Fluctuations. Front Physiol. 2019;10. doi:10.3389/fphys.2019.00257

1256   35.   Decker LM, Cignetti F, Stergiou N. Executive function orchestrates regulation of task-
1257   relevant gait fluctuations. Gait Posture. 2013;38: 537–540.
1258   doi:10.1016/j.gaitpost.2012.12.018

1259   36.   Homs AF, Dupeyron A, Torre K. Relationship between gait complexity and pain attention
1260   in chronic low back pain. Pain. 2022;163: e31–e39.
1261   doi:10.1097/j.pain.0000000000002303

1262   37.   Toebes MJP, Hoozemans MJM, Furrer R, Dekker J, van Dieën JH. Associations between
1263   measures of gait stability, leg strength and fear of falling. Gait Posture. 2015;41: 76–80.
1264   doi:10.1016/j.gaitpost.2014.08.015

1265   38.   Bruijn SM, Meijer OG, Beek PJ, van Dieën JH. Assessing the stability of human
1266   locomotion: a review of current measures. J R Soc Interface. 2013;10: 20120999.
1267   doi:10.1098/rsif.2012.0999

1268   39.   Terrier P, Dériaz O. Non-linear dynamics of human locomotion: effects of rhythmic
1269   auditory cueing on local dynamic stability. Front Physiol. 2013;4: 230.
1270   doi:10.3389/fphys.2013.00230

1271   40.   Dingwell JB, Cusumano JP, Sternad D, Cavanagh PR. Slower speeds in patients with
1272   diabetic neuropathy lead to improved local dynamic stability of continuous overground
1273   walking. Journal of Biomechanics. 2000;33: 1269–1277. doi:10.1016/S0021-
1274   9290(00)00092-0

1275   41.   Roos PE, Dingwell JB. Influence of Simulated Neuromuscular Noise on the Dynamic
1276   Stability and Fall Risk of a 3D Dynamic Walking Model. J Biomech. 2011;44: 1514–1520.
1277   doi:10.1016/j.jbiomech.2011.03.003

1278   42.   Bruijn SM, Bregman DJJ, Meijer OG, Beek PJ, van Dieën JH. Maximum Lyapunov
1279   exponents as predictors of global gait stability: a modelling approach. Med Eng Phys.
1280   2012;34: 428–436. doi:10.1016/j.medengphy.2011.07.024

1281   43.   Mehdizadeh S. The largest Lyapunov exponent of gait in young and elderly individuals:
1282   A systematic review. Gait Posture. 2018;60: 241–250.
1283   doi:10.1016/j.gaitpost.2017.12.016

44.   Lorenz EN. The Essence of Chaos. Nachdr. Seattle: Univ. of Washington Press; 2008.

45. Toebes MJP, Hoozemans MJM, Furrer R, Dekker J, Van Dieën JH. Local dynamic stability
and variability of gait are associated with fall history in elderly subjects. Gait & Posture.
2012;36: 527–531. doi:10.1016/j.gaitpost.2012.05.016

46. Sejdić E, Fu Y, Pak A, Fairley JA, Chau T. The Effects of Rhythmic Sensory Cues on the
Temporal Dynamics of Human Gait. Ernst MO, editor. PLoS ONE. 2012;7: e43104.
doi:10.1371/journal.pone.0043104

47. Van Schooten KS, Sloot LH, Bruijn SM, Kingma H, Meijer OG, Pijnappels M, et al.
Sensitivity of trunk variability and stability measures to balance impairments induced by
galvanic vestibular stimulation during gait. Gait & Posture. 2011;33: 656–660.
doi:10.1016/j.gaitpost.2011.02.017

48. Jordan K, Challis JH, Cusumano JP, Newell KM. Stability and the time-dependent
structure of gait variability in walking and running. Human Movement Science. 2009;28:
113–128. doi:10.1016/j.humov.2008.09.001

49. Piergiovanni, S, Terrier, P. Effects of metronome walking on long-term attractor
divergence and correlation structure of gait: a validation study in older people. Scientific
Reports. 2024;14: 15784. doi:10.1038/s41598-024-65662-5

50. Piergiovanni S, Terrier P. Validity of Linear and Nonlinear Measures of Gait Variability to
Characterize Aging Gait with a Single Lower Back Accelerometer. SENSORS. 2024;24.
doi:10.3390/s24237427

51. Terrier P, Reynard F. Maximum Lyapunov exponent revisited: Long-term attractor
divergence of gait dynamics is highly sensitive to the noise structure of stride intervals.
Gait & Posture. 2018;66: 236–241. doi:10.1016/j.gaitpost.2018.08.010

52. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The
PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ.
2021;372: n71. doi:10.1136/bmj.n71

53. Schiavo JH. PROSPERO: An International Register of Systematic Review Protocols. Med
Ref Serv Q. 2019;38: 171–180. doi:10.1080/02763869.2019.1588072

54. Terrier P. From stability to complexity: A systematic review protocol on long-term
divergence exponents in gait analysis. medRxiv. 2024; 2024–12.

55. Terrier P, Dériaz O. Kinematic variability, fractal dynamics and local dynamic stability of
treadmill walking. J NeuroEngineering Rehabil. 2011;8: 12. doi:10.1186/1743-0003-8-
12

56. Terrier P. Complexity of human walking: the attractor complexity index is sensitive to
gait synchronization with visual and auditory cues. PeerJ. 2019;7: e7417.
doi:10.7717/peerj.7417

57. Pustejovsky JE, Tipton E. Meta-analysis with Robust Variance Estimation: Expanding the
Range of Working Models. Prev Sci. 2022;23: 425–438. doi:10.1007/s11121-021-
01246-3

58. Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. Journal of
Statistical Software. 2010;36: 1–48. doi:10.18637/jss.v036.i03

59.   Pustejovsky JE. clubSandwich. 2023. doi:10.32614/CRAN.package.clubSandwich

60.   Reynard P, Terrier P. Local dynamic stability of treadmill walking: Intrasession and week-
1327   to-week repeatability. Journal of Biomechanics. 2014;47: 74–80.
1328   doi:10.1016/j.jbiomech.2013.10.011

1329   61.   Takens F. Detecting strange attractors in turbulence. In: Rand D, Young L-S, editors.
1330   Dynamical Systems and Turbulence, Warwick 1980. Springer Berlin Heidelberg; 1981.
1331   pp. 366–381.

1332   62.   Dingwell JB, Cusumano JP, Cavanagh PR, Sternad D. Local Dynamic Stability Versus
1333   Kinematic Variability of Continuous Overground and Treadmill Walking. Journal of
1334   Biomechanical Engineering. 2001;123: 27–32. doi:10.1115/1.1336798

1335   63.   Manor B, Wolenski P, Guevaro A, Li L. Differential effects of plantar desensitization on
1336   locomotion dynamics. Journal of electromyography and kinesiology□: official journal of
1337   the International Society of Electrophysiological Kinesiology. 2009;19: e320-8.
1338   doi:10.1016/j.jelekin.2008.06.006

1339   64.   McAndrew PM, Wilken JM, Dingwell JB. Dynamic stability of human walking in visually
1340   and mechanically destabilizing environments. J Biomech. 2011;44: 644–649.
1341   doi:10.1016/j.jbiomech.2010.11.007

1342   65.   Sinitksi EH, Terry K, Wilken JM, Dingwell JB. Effects of perturbation magnitude on
1343   dynamic stability when walking in destabilizing environments. Journal of Biomechanics.
1344   2012;45: 2084–2091. doi:10.1016/j.jbiomech.2012.05.039

1345   66.   Rábago CA, Dingwell JB, Wilken JM. Reliability and Minimum Detectable Change of
1346   Temporal-Spatial, Kinematic, and Dynamic Stability Measures during Perturbed Gait.
1347   PLoS ONE. 2015;10: e0142083. doi:10.1371/journal.pone.0142083

1348   67.   Reynard F, Terrier P. Role of visual input in the control of dynamic balance: variability
1349   and instability of gait in treadmill walking while blindfolded. Exp Brain Res. 2015;233:
1350   1031–1040. doi:10.1007/s00221-014-4177-5

1351   68.   Qiao M, Truong KN, Franz JR. Does local dynamic stability during unperturbed walking
1352   predict the response to balance perturbations? An examination across age and falls
1353   history. Gait & Posture. 2018;62: 80–85. doi:10.1016/j.gaitpost.2018.03.011

1354   69.   Manor B, Wolenski P, Li L. Faster walking speeds increase local instability among people
1355   with peripheral neuropathy. Journal of Biomechanics. 2008;41: 2787–2792.
1356   doi:10.1016/j.jbiomech.2008.07.006

1357   70.   Terrier P, Le Carre J, Connaissa M-L, Leger B, Luthi F. Monitoring of Gait Quality in
1358   Patients With Chronic Pain of Lower Limbs. IEEE Trans Neural Syst Rehabil Eng.
1359   2017;25: 1843–1852. doi:10.1109/TNSRE.2017.2688485

1360   71.   Bizovska L, Svoboda Z, Kubonova E, Vuillerme N, Hirjakova Z, Janura M. The differences
1361   between overground and treadmill walking in nonlinear, entropy-based and frequency
1362   variables derived from accelerometers in young and older women - preliminary report.
1363   Acta of Bioengineering and Biomechanics. 2018;20: 93–100. doi:10.5277/ABB-00987-
1364   2017-02

72.   Bizovska L, Svoboda Z, Janura M, Bisi MC, Vuillerme N. Local dynamic stability during
gait for predicting falls in elderly people: A one-year prospective study. Plos One.
2018;13: e0197091. doi:10.1371/journal.pone.0197091

73.   Matinazad N, Esteki A, Ghomashchi H. Comprehensive characterization of gait variability
in patients with knee osteoarthritis for altered velocities. J Mech Med Biol. 2018;18:
1850041. doi:10.1142/S0219519418500410

74.   Terrier P. Fractal Fluctuations in Human Walking: Comparison Between Auditory and
Visually Guided Stepping. Ann Biomed Eng. 2016;44: 2785–2793. doi:10.1007/s10439-
016-1573-y

75.   Rosenstein MT, Collins JJ, De Luca CJ. A practical method for calculating largest
Lyapunov exponents from small data sets. Physica D: Nonlinear Phenomena. 1993;65:
117–134. doi:10.1016/0167-2789(93)90009-P

76.   Dingwell JB, Kang HG. Differences Between Local and Orbital Dynamic Stability During
Human Walking. Journal of Biomechanical Engineering. 2007;129: 586–593.
doi:10.1115/1.2746383

77.   Ahn J, Hogan N. Long-range correlations in stride intervals may emerge from non-
chaotic walking dynamics. PLoS One. 2013;8: e73239.
doi:10.1371/journal.pone.0073239

78.   Raffalt PC, Kent JA, Wurdeman SR, Stergiou N. To walk or to run – a question of
movement attractor stability. J Exp Biol. 2020;223: jeb224113. doi:10.1242/jeb.224113

79.   Rantalainen T, Hart NH, Nimphius S, Wundersitz DW. Associations Between Step
Duration Variability and Inertial Measurement Unit Derived Gait Characteristics. Journal
of Applied Biomechanics. 2016;32: 401–406. doi:10.1123/jab.2015-0266

80.   Sedighi A, Rashedi E, Nussbaum MA. A head-worn display ("smart glasses") has
adverse impacts on the dynamics of lateral position control during gait. Gait & Posture.
2020;81: 126–130. doi:10.1016/j.gaitpost.2020.07.014

81.   Dingwell JB, Cusumano JP. Re-interpreting detrended fluctuation analyses of stride-to-
stride variability in human walking. Gait Posture. 2010;32: 348–353.
doi:10.1016/j.gaitpost.2010.06.004

82.   Terrier P. Step-to-step variability in treadmill walking: influence of rhythmic auditory
cueing. PLoS ONE. 2012;7: e47171. doi:10.1371/journal.pone.0047171

83.   Terrier P, Turner V, Schutz Y. GPS analysis of human locomotion: further evidence for
long-range correlations in stride-to-stride fluctuations of gait parameters. Human
movement science. 2005;24: 97–115.

84.   Terrier P, Dériaz O. Persistent and anti-persistent pattern in stride-to-stride variability of
treadmill walking: influence of rhythmic auditory cueing. Human movement science.
2012;31: 1585–1597.

85.   Yakhdani HR, Bafghi HA, Meijer OG, Bruijn SM, Dikkenberg NVD, Stibbe AB, et al.
Stability and variability of knee kinematics during gait in knee osteoarthritis before and

after replacement surgery. Clinical Biomechanics. 2010;25: 230–236.
doi:10.1016/j.clinbiomech.2009.12.003

86.　Dingwell JB, Cusumano JP. Nonlinear time series analysis of normal and pathological
human walking. Chaos: An Interdisciplinary Journal of Nonlinear Science. 2000;10:
848–863. doi:10.1063/1.1324008

87.　Wang Z, Peng S, Zhang H, Sun H, Hu J. Gait Parameters and Peripheral Neuropathy in
Patients With Diabetes: A Meta-Analysis. Front Endocrinol (Lausanne). 2022;13:
891356. doi:10.3389/fendo.2022.891356

88.　Mustapa A, Justine M, Mohd Mustafah N, Jamil N, Manaf H. Postural Control and Gait
Performance in the Diabetic Peripheral Neuropathy: A Systematic Review. Biomed Res
Int. 2016;2016: 9305025. doi:10.1155/2016/9305025

89.　Paul L, Ellis BM, Leese GP, McFadyen AK, McMurray B. The effect of a cognitive or motor
task on gait parameters of diabetic patients, with and without neuropathy. Diabet Med.
2009;26: 234–239. doi:10.1111/j.1464-5491.2008.02655.x

90.　Roman de Mettelinge T, Delbaere K, Calders P, Gysel T, Van Den Noortgate N, Cambier
D. The impact of peripheral neuropathy and cognitive decrements on gait in older
adults with type 2 diabetes mellitus. Arch Phys Med Rehabil. 2013;94: 1074–1079.
doi:10.1016/j.apmr.2013.01.018

91.　Cignetti F, Decker LM, Stergiou N. Sensitivity of the Wolf's and Rosenstein's Algorithms
to Evaluate Local Dynamic Stability from Small Gait Data Sets. Ann Biomed Eng.
2012;40: 1122–1130. doi:10.1007/s10439-011-0474-3

92.　Kang HG, Dingwell JB. Effects of walking speed, strength and range of motion on gait
stability in healthy older adults. Journal of Biomechanics. 2008;41: 2899–2905.
doi:10.1016/j.jbiomech.2008.08.002

93.　Kang HG, Dingwell JB. Dynamic stability of superior vs. inferior segments during
walking in young and older adults. Gait & Posture. 2009;30: 260–263.
doi:10.1016/j.gaitpost.2009.05.003

94.　Mirelman A, Maidan I, Bernad-Elazari H, Shustack S, Giladi N, Hausdorff JM. Effects of
aging on prefrontal brain activation during challenging walking conditions. Brain and
Cognition. 2017;115: 41–46. doi:10.1016/j.bandc.2017.04.002

95.　Holtzer R, Mahoney JR, Izzetoglu M, Izzetoglu K, Onaral B, Verghese J. fNIRS Study of
Walking and Walking While Talking in Young and Old Individuals. J Gerontol A Biol Sci
Med Sci. 2011;66A: 879–887. doi:10.1093/gerona/glr068

96.　Holtzer R, Epstein N, Mahoney JR, Izzetoglu M, Blumen HM. Neuroimaging of Mobility in
Aging: A Targeted Review. J Gerontol A Biol Sci Med Sci. 2014;69: 1375–1388.
doi:10.1093/gerona/glu052

97.　Rosano C, Aizenstein HJ, Studenski S, Newman AB. A Regions-of-Interest Volumetric
Analysis of Mobility Limitations in Community-Dwelling Older Adults. J Gerontol A Biol
Sci Med Sci. 2007;62: 1048–1055. doi:10.1093/gerona/62.9.1048

98. Zwergal A, Linn J, Xiong G, Brandt T, Strupp M, Jahn K. Aging of human supraspinal
locomotor and postural control in fMRI. Neurobiology of Aging. 2012;33: 1073–1084.
doi:10.1016/j.neurobiolaging.2010.09.022

99. Lee B-C, Choi J, Martin BJ. Roles of the prefrontal cortex in learning to time the onset of
pre-existing motor programs. PLOS ONE. 2020;15: e0241562.
doi:10.1371/journal.pone.0241562

100. Malcolm BR, Foxe JJ, Butler JS, Molholm S, De Sanctis P. Cognitive load reduces the
effects of optic flow on gait and electrocortical dynamics during treadmill walking.
Journal of Neurophysiology. 2018;120: 2246–2259. doi:10.1152/jn.00079.2018

101. Purohit R, Bhatt T. Mobile Brain Imaging to Examine Task-Related Cortical Correlates of
Reactive Balance: A Systematic Review. Brain Sciences. 2022;12: 1487.
doi:10.3390/brainsci12111487

102. Bruijn SM, Van Dieën JH, Meijer OG, Beek PJ. Statistical precision and sensitivity of
measures of dynamic gait stability. Journal of Neuroscience Methods. 2009;178: 327–
333. doi:10.1016/j.jneumeth.2008.12.015

103. Riva F, Bisi MC, Stagni R. Gait variability and stability measures: Minimum number of
strides and within-session reliability. Computers in Biology and Medicine. 2014;50: 9–
13. doi:10.1016/j.compbiomed.2014.04.001

104. Ravi DK, Marmelat V, Taylor WR, Newell KM, Stergiou N, Singh NB. Assessing the
Temporal Organization of Walking Variability: A Systematic Review and Consensus
Guidelines on Detrended Fluctuation Analysis. Front Physiol. 2020;11: 562.
doi:10.3389/fphys.2020.00562

1465

1466

Supporting Information

1468

S1 Text. Systematic review database. Complete data extraction tables with study

characteristics (Table A: research types; Table B: study characteristics; Table C:

methodological specifications), search strategy, and detailed information for all 56 included

studies. (DOCX)

S2 Figures. Divergence curve figures. Compiled divergence curves from 29 studies with

detailed figure descriptions showing methodological illustrations and comparative results.

(PDF)

S3 Table. Quality assessment results. Complete quality assessment scores for all 56

studies using the three-domain framework (maximum 15 points). (XLSX)

1478

Figure legends

Fig 1. PRISMA flowchart of systematic review study selection. Flow diagram showing identification, screening, eligibility assessment, and final inclusion of studies examining long-term divergence exponents in human gait analysis.

Fig 2. Study characteristics. a) Annual publication trends. **b)** Distribution by research type and participant category. **c)** Sample size distribution. DE, divergence exponent.

Fig 3. Methodological characteristics. a) Measurement environments. **b)** Sensor modalities. **c)** Anatomical measurement locations. IMU, inertial measurement unit.

Fig 4. Measurement duration and temporal resampling Distribution of walking bout durations across four duration categories.

Fig 5. Signal processing characteristics. a) Signal sources used for computing DEs. **b)** Resampling methods to normalize signal durations. DE, divergence exponent; COP, center of pressure; EMG, electromyography.

Fig 6. Meta-analysis of correlations between long-term DE and DFA scaling exponents. Forest plot showing study-level correlations with 95% confidence intervals and pooled estimate using correlated and hierarchical effects (CHE) model with robust variance estimation. N, total observations; k, number of correlation estimates per study; CI, confidence interval; DFA, detrended fluctuation analysis; DE, divergence exponent.

Duration of a single analysed walking bout

Category	Time criterion	Stride criterion
Very short	≤ 2 min	≤ 100 strides
Short	> 2 – 5 min	101 – 250 strides
Moderate	> 5 – 10 min	251 – 500 strides
Long	> 10 min	> 500 strides

a Walking Conditions

- Overground — 16%
- Treadmill — 71%
- Mixed (OG+TW) — 11%
- Free-living — 2%

b Measurement Systems

- 3D motion capture — 52%
- Inertial sensors — 29%
- Instrumented treadmill — 5%
- Mixed — 14%

c Anatomical Measurement Sites

- Trunk — 55%
- Lower limbs — 20%
- Trunk + lower limbs — 16%
- Foot pressure — 5%
- Others — 4%