Robust ciliary flows protect early Xenopus embryos from pathogens independent of multiciliated cell patterning

Athullya Baby[a,b,c], **Alice Briole**[b], **Ayush Yadav**[a], **Isabelle Cheylan**[c], **Virginie Thomé**[a], **Camille Boutin**[a], **Annie Viallat**[b], **Julien Favier**[c,1], **Etienne Loiseau**[b,1], **and Laurent Kodjabachian**[a,1]

[a]Aix Marseille Univ, CNRS, IBDM, Turing Centre for Living Systems, Marseille, France
[b]Aix Marseille Univ, CNRS, CINAM, Turing Centre for Living Systems, Marseille, France
[c]Aix Marseille Univ, CNRS, M2P2, Turing Centre for Living Systems, Marseille, France

During the early stages of development, the skin of the Xenopus embryo is covered by around two thousand evenly distributed multiciliated cells (MCCs). This striking spatial distribution is believed to maximise the generation of superficial flows at the scale of the embryo. However, the specific role of regular MCC distribution and the physiological function of vigorous ciliary activity remain elusive. To investigate the extent to which superficial flows provide protection against external pathogens before the immune system matures, we combined experimental and computational approaches. First, we cultivated epithelial explants in order to quantify the distribution of MCCs, beating frequencies, and three-dimensional fluid flows. We then used these data to validate a computational fluid dynamics model. Using this model, seeded with structural data from whole embryos, our simulations reveal that the collective ciliary beatings create a robust liquid shield along the embryonic flank, which is highly effective at clearing pathogens from the vicinity of the epithelial surface. Through parametric analyses, we further demonstrate that this protective function is remarkably resilient. Indeed, the effectiveness of pathogen clearance is primarily governed by the overall characteristic velocity of the cilia and is less affected by moderate variations in MCC density and spatial organisation. Our findings suggest that, rather than optimising energy consumption, the biological system prioritises functional robustness to ensure reliable protection.

ciliated epithelium | flow generation | Xenopus embryo

Correspondence:
Julien.favier@univ-amu.fr
Etienne.loiseau@univ-amu.fr
Laurent.kodjabachian@univ-amu.fr

Introduction

During the larval development of the amphibian Xenopus laevis, the skin is temporarily covered by approximately two thousand multiciliated cells (MCCs), which are evenly distributed amongst secretory cells that produce a protective mucus layer (1, 2) . MCCS represent about 10% of the total number of cells on the surface of the embryo (3). The coordinated beating of their numerous cilia generates robust fluid flows directed from anterior-dorsal to posterior-ventral across the body surface (4, 5). MCCs emerge during embryonic development, and persist into the free-swimming tadpole stage, long after hatching, eventually disappearing (NF stage 40-47) through programmed cell death (apoptosis), or transdifferentiation into mucus-producing cells(2, 6–8). The pre-

cise physiological function of Xenopus MCCs and the vigorous flow they induce remains poorly understood. Several hypotheses have been proposed through works on different aquatic species , which include enhancement of respiratory gas exchange and, critically, clearance of pathogens in the vicinity of the embryo (2, 9–14). The hypothesis that cilia-powered flows prevent the attachment of post-hatching debris and external pathogens is particularly compelling. This is supported by the observation that maximum ciliation occurs after hatching (the breaking of the vitelline membrane), and that peak ciliary activity coincides with a period when the immune system is still non-functional (2, 15, 16). In that respect, this clearance function would offer a dual safety mechanism alongside the physical barrier of the 6 μm-thick mucus layer covering the embryo's surface (17).

The emergence of coherent cilia-powered flows depends on multiple parameters deployed at both cell and tissue-scales (5, 18–20). Such parameters, including cilia and MCC density, alignment, spacing and beating strength, are interdependent, making it difficult to evaluate experimentally their individual influence on flow production. Moreover, measuring a three-dimensional flow on the millimeter scale, especially over the non-transparent and non-symmetrical surface of a 3D embryo, is inherently difficult. While interesting theoretical and numerical approaches have been proposed, they often lack experimental validation at the tissue scale or comprehensive discussion regarding protection against infection (21).

A remarkable feature of Xenopus MCCs is their regular distribution across the embryonic skin, often at the nodes of a highly ordered network, that is established through active migration and homotypic repulsion, mediated by the Scf/Kit signalling system (22). This raises fundamental questions about the criticality and optimization of MCC distribution for cleaning efficiency: Is this precise spatial arrangement essential, and is it optimized for effective washing?

Here, we address these open questions through a multi-step strategy including both experiments and numerical modelling. First, we establish a protocol to quantify the rate of infection of individual embryos by Aeromonas hydrophila, a natural pathogen of Xenopus laevis (14, 17). Here, the aim is to test the importance of individual parameters of the

system on infection rates, including MCC density and spacing. Second, we establish the culture of transparent Xenopus embryonic epithelial explants on a planar substrate. We then measure the 3D flows generated by MCCs on these explants using video-microscopy. Next, we construct a numerical model, capable of simulating these flows, and validate its predictions against experimental measurements made on the explants. Finally, we leverage this validated numerical model to evaluate the relative influence on flow production and pathogen clearance of individual parameters, including MCC beating force, density, spacing and alignment. A first major outcome of this work is to provide a validated model to compute the flow at different distances above ciliated epithelia. Experiments on Xenopus embryos reveal that ciliated cells collectively create a robust liquid shield that effectively protects the embryo from external pathogens. Remarkably, this protective barrier demonstrates high resilience to variations in both MCC density and spatial distribution. This suggests that the biological system does not necessarily optimize its function in terms of a precise, singular MCC density or spatial arrangement. Instead, the biological system appears to prioritize robustness in physiological function, ensuring effective protection even amidst challenging environmental conditions and inherent imperfections in the density or homogeneity of the epithelial ciliated system. We discuss these results in the context of the effectiveness of the protective flow against bacterial infection.

Results

1. Cilia-powered flows prevent infection of Xenopus embryos

Previous work suggested that long-term exposure (72h) of Xenopus laevis embryos to A. hydrophila caused enhanced death, when cilia-powered flow was reduced ([14]). To better quantify infection rates, we established a protocol to score the number of colony-forming units (cfu) from the lysate of individual embryos, exposed to A. hydrophila for 9 hours at stage 30, when the ciliated epidermis is fully functional (see Methods). Uninjected control embryos, as well as Cas9-injected control embryos, displayed normal ciliogenesis, powerful directional flow and showed very little infection rates (1/15 for either condition, average cfu=26 and 44, respectively)(see Fig.1). In contrast, when embryos were co-injected with Cas9 and guide RNA against foxJ1, a key regulator of motile ciliogenesis ([23]), cilia were absent, flow was abolished, and a majority of embryos were infected (13/25, average cfu=474), half of them dying during the 9h period of exposure to A. hydrophila (6/13)(Fig.1H). This result confirms that cilia-powered flows help to protect Xenopus embryos from bacterial infection. Next, we wanted to evaluate the importance of regular MCC spacing for the resistance to infection by A. hydrophila. For this, we used Axitinib, a chemical inhibitor of the Kit receptor, which was reported to alter MCC homotypic repulsion and spacing ([22]). Unexpectedly, we observed that exposure to axitinib upset MCC distribution, but also caused their premature disappearance (see Fig. S1). This compound

phenotype prevented us from testing specifically the importance of MCC geometrical order for resistance against A. hydrophila. To overcome this experimental limitation, we developed a combined approach utilizing in vitro explants and a numerical model to facilitate the evaluation of how individual parameters of the system impact flow generation and pathogen clearance.

Fig. 1. MCCs are required to limit infections of Xenopus embryos by Aeoromas hydrophila. Distributions of MCCs on the flank of an embryo and trajectories of fluorescent beads propelled by the ciliary activity in the control case (A) & (B) and for a Cas 9 injected embryo (C) & (D). For an embryo co-injected with Cas9 and guide RNA against FoxJ1, the main gene involved in ciliogenesis, cilia are absent (E) and beads are not transported (F). (G) Quantification of the velocity of beads above the flank of embryos. (H) Infectability of embryos by Aeromonas hydrophila. The absence of cilia due to foxJ1 knockout causes the lack of flow and enhanced infection rates.

2. Static and Dynamic Features of MCCs in Embryos and Explants: Spatial Organization and Flow Fields

Embryos and Explant preparation. To characterize MCC organization, we prepared and fixed eight Xenopus embryos at stage 29-30. Additionally, twelve embryos at similar stage were fixed after treatment with Axitinib to induce disruptions in MCC spatial distribution. The epithelia of their flanks (surface areas ranging from 3.18 to 5.10 mm²) were subsequently imaged using immunofluorescence microscopy to visualize MCCs and their spatial arrangement. For experimental flow measurements, we prepared and imaged nine living explants using live-cell brightfield microscopy. Three of these explants exhibited flat surface areas exceeding 3.7 mm², while the remaining six displayed only small, non-contiguous flat regions, which were imaged across one to three tiles of 0.46 mm² each. We also prepared and imaged nine axitinib-treated explants, with three demonstrating flat areas larger than 1.75 mm² and six exhibiting smaller flat areas analyzed on one or two 0.46 mm² tiles. MCC and cilia development, along with ciliary dynamics, were assessed after 5 and 6 days of in-vitro culture following explant collection and transfer onto glass slides. Explants, cultured for 6 days (corresponding to embryonic stage 30-32), consistently displayed significantly

more vigorous flow compared to those cultured for 5 days (Fig. S2). Consequently, our subsequent analyses focused exclusively on day 6 explants to ensure optimal ciliary activity (see movie S1).

Fig. 2. Static parameters of the ciliated epidermis.(A) Spatial distribution of the multiciliated cells (MCCs) on the flank of an embryo. Binarised image computed from a fluorescent image (see methods section). (B) Delaunay's triangulation, computed from the image on panel (A), is used to compute the geometrical order. (C) Characteristic epidermal explant (3400 µm*2720 µm) visualized through stitching of 15 imaging tiles (680 µm*680 µm). The region inside the dashed line corresponds to location where the tissue is not perfectly flat. (D) MCC distribution corresponding to the explant in (C). Binarised image computed from movie acquired in bright field (see methods).(E) Comparison of the surface area of MCCs measured on fixed embryos and on live explants for both control and axitinib-treated condition. For explants, a correction is applied as described in the method section to compare with measurements done on fixed samples. (F) Comparison of the MCC density on embryos and explants for both control and axitinib-treated conditions. (G) Comparison of the geometrical order on embryos and on explants for both control and Axitinib-treated conditions.

Comparative Analysis of Embryos and Explants. Figure 2 illustrates our typical imaging and processing workflow on fixed embryos (Fig. 2A & Fig. 2B) and on living explants (Fig. 2C & Fig. 2D). Figure 2E shows that the size of MCCs is comparable between embryos and explants, suggesting that the explant culture largely preserves individual MCC morphology. However, due to spreading on the substrate, MCC density is significantly lower in explants than in embryos (Figure 2F). Despite this reduction, the MCC density in explants remained sufficiently high to generate robust surface flow (see below), confirming their suitability for experimental studies (5). The spatial distribution of MCCs is quantified using Delaunay triangulation (Fig. 2B), yielding a geometric order parameter (Fig. 2G) that ranges from 0 (random distribution) to 1 (perfectly regular distribution). This analysis indicates similar overall MCC spatial distributions in untreated embryos and explants, whereas an increased heterogeneity is observed in axitinib-treated explants compared

to untreated explants. We note that unlike in whole embryos, axitinib treatment on isolated epidermis explants does not cause MCC disappearance, which is compatible with previous studies indicating that MCC death signals emanate from the mesoderm (7, 8). Thus, axitinib treatment on explants provides a valuable paradigm to disturb MCC spatial organization, and measure the possible impact on the dynamic parameters of the ciliated epidermis.

Dynamic Features of Ciliary Flow. Due to the opacity and non-planar geometry of Xenopus embryos, directly measuring ciliary beat directions, as well as the direction and velocity of fluid flow at various distances from the embryonic surface, represent challenges that remain unmet. Therefore, we measured these critical parameters on explants, which offer the advantage of transparency and planarity (Fig. 3). These experimentally derived parameters will serve as inputs for the numerical model presented in the subsequent section, enabling a direct comparison between experimental and numerical values for model validation. Fluid streamlines were determined by Particle Imaging Velocimetry of fluorescent micro-beads on the surface of explant epithelia (Fig. 3 A & B). We observe that streamlines become more regular at greater distances from the surface, suggesting a smoothing effect of the flow field. We noticed that fluorescent beads remain in the focal plane over large distances. This is expected as at low Reynolds number flows are laminar. Yet, we also observe some beads moving up or down from time to time (See Movie SI2). Ciliary beat frequencies of MCCs in explants are in the range of 8 to 16 Hz, with a mean of 13 Hz (Fig. 3F), showing no significant difference between axitinib-treated and untreated explants.

The directions of ciliary beats are illustrated in Figure 3 C & D, for both untreated and axitinib-treated explants. These images clearly show that ciliary beats are relatively well aligned over short distances. The degree of this alignment was quantified using the polarity order parameter P (where P=1 indicates perfectly aligned beat directions and P=0 denotes random orientation). Figure 3 G depicts how this polarity varies across different length scales for both untreated and axitinib-treated explants. We observe progressive, partial loss of correlation with increasing distance between MCCs, which varies among explants. Notably, a strong degree of orientation, reaching up to 0.9, can be maintained at distances of up to 500 µm. No statistically significant difference in the polarity order was measured between untreated and axitinib-treated explants (Fig. 3G).

The flow velocity, extracted from PIV analysis, is also illustrated on maps for two heights, (30 µm and 90 µm) in Figure 3H & I. We observe rapid velocities, reaching several hundred micrometers per second, which become more homogeneous at greater distances from the tissue surface. The mean velocity at a given height as a function of that height is represented in Figure 3E for various explants. The velocity immediately above the tips of the MCC cilia, denoted U_0, is depicted on the surface of one explant in Figure 3J and for other large explants in Figure S5. U_0 represents the characteristic velocity of the fluid being dragged by an individ-

ual MCC. Strikingly, we observe significant heterogeneity in U_0 across different multiciliated cells within the same explant. This variability likely arises from inherent differences in MCC surface area, as well as variations in the frequency and number of cilia within each individual cell.

The experimentally determined frequency, polarity order, and distribution of U_0 values were all used as input parameters for our numerical model. The model predictions for streamlines and flow velocities at different heights above the epithelium were subsequently compared against these experimental measurements to validate its performance.

3. Numerical model of a ciliated epithelium

The model. Our numerical model, comprehensively detailed in the Materials and Methods section employs a three-dimensional Lattice Boltzmann method to simulate the fluid dynamics above the embryonic tissue. The ciliated cells are incorporated using the Immersed Boundary Method (IBM), where each cell is represented as a point force acting on a Lagrangian point. These forces, which represent the net ciliary propulsion, are located at a height of 20 μm above the epithelial surface, corresponding to the approximate height of the cilia.

Figure 4A provides a schematic representation of MCCs and the Xenopus epithelium within our model. The input parameters for the model are the position, direction, and magnitude of these point forces representing the ciliated cells. The positions of the point forces directly correspond to the centroids of MCCs obtained from our experimental images, their direction, in the horizontal plane, is determined based on the measured ciliary beating directions obtained from experimental data (Fig. S4), and their magnitudes are directly obtained from the U_0 values measured on our explant experiments.

The computational domain includes a solid wall at the bottom, representing the epithelial surface, on which a no-slip boundary condition for the fluid velocity is imposed. The four lateral walls in the x and y directions are modeled as open boundaries to ensure the fluid flow is unrestricted by artificial computational barriers. The domain size was optimized using a sensitivity analysis to ensure that boundary proximity had negligible effects on the numerical results (Fig. S6). For consistency and to ensure grid-converged results, a uniform grid size of dx=dy=dz=20 μm was used across all simulations.

The domain size was optimized using a sensitivity analysis to ensure that boundary proximity had negligible effects on the numerical results (Fig. S6).

Validation of the model. We validated our model by comparing the numerical flow fields with experimental measurements obtained from tissue explants. Figure 4B presents a side-by-side comparison of simulated and experimental streamlines at heights 30 and 90 μm for a representative explant (n°9). The local agreement is quantified by color maps of the angular error, defined as the local difference in flow direction between the simulation and experiments. Furthermore, Figure 4C compares the average planar velocity magnitudes across multiple explants at various heights. Note that vertical velocities were excluded from comparison due to the current inability to measure them experimentally.

Overall, there is good agreement between the model and experimental data. Streamlines from both simulations and experiments visually demonstrate a consistent lateral flow across the entire explant domain. Close to the tissue, significant but highly localized angular errors are observed. These errors are likely attribuable to small, unresolved tissue folds that harbor motile cilia. At greater distances from the tissue surface, these angular errors become minor, confirming that the model accurately reproduces the large-scale flow directions. Simulated average planar velocities also show very good agreement for all explants, with a deviation of less than 10% from experimental measurements (Figure 4C).

These results validate our simulation approach as an effective tool for describing fluid flows at the surface of ciliated epithelia. In the following section, we leverage this validated model to analyze flow over embryonic flanks, operating under the assumption that the tissue's slight convexity does not significantly alter the established flow patterns.

4. Computing 3d flows on embryo flanks

Leveraging this validated numerical model at hand, we proceeded to predict flow patterns at the surface of Xenopus embryo flanks.

Initial Conditions for the Model. The static input parameters, such as MCC position and distribution, were directly extracted from images of intact embryos. As MCC beat directions cannot be measured directly on fixed embryos, we adopted an average level of alignment based on explant tissue data, corresponding to a polarity order of 0.6 (which falls within the observed range of 0.4 to 0.8 at a radius of 500 micrometers for explants). To establish this polarity, beat directions were initially assigned randomly to each MCC and then iteratively adjusted until two criteria were met: (i) achieving a polarity order of 0.6 at a 500 μm radius and, (ii) reproducing the in-vivo major streamlines observed on the embryonic surface (see Figures 5A & (B). As shown in the horizontal projection of the streamlines (Figure 5(B)), the model successfully reproduces the characteristic diagonal flow (from anterior-dorsal to posterior-ventral) and the distinct streamline patterns on the dorsal side of the embryo (4).

Finally, the characteristic velocity U_0 for each MCC was randomly assigned following a Gaussian distribution with a mean of 350 μm/s and a standard deviation of 250 μm/s. These parameters were chosen based on explant measurements, where velocities ranged from 20 μm/s to 800 μm/s in a Gaussian-like distribution.

Flow characterization and spatial analysis. The resulting flow was characterized by analyzing the individual streamlines.

On the embryo flank, the streamlines are predominantly oriented along the horizontal anterior–posterior axis (Fig. 5(C)).

Fig. 3. (A) & (B) Visualisation of the streamlines at two heights z=30 µm and z=90 µm above the surface of an explant. Streamlines are obtained by computing the standard deviation over 100 frames (≃ 1s) of flowing fluorescent beads. (C) & (D) Ciliary beat directions for a control explant (C) and an axitinib-treated explant (D). Right panels show zooms to better visualise the coordination of the beat directions. (E) Average flow velocity profile above the surface of explants for both control and axitinib-treated samples. (F) Distributions of ciliary beat frequencies for both control and axitinib treated-explants. (G) The coordination of beat directions is measured via the polarity order at all length scales, from neighbouring cells (≃ 50 µm) till 500 µm. There is no significant difference between control and axitinib-treated explants, as shown in the two panels on the right side. (H - J) Maps of the amplitude of the flow velocity at different distances from the surface of an explant. U_0 map in (J) corresponds to the velocity measured at the tip of the cilia on each ciliated cell.

Fig. 4. Validation of the numerical model for ciliated epithelia. (A) Schematic of the virtual epithelium. Ciliated cells are modeled by point forces. (B) Comparison between measured streamlines and streamlines computed by the model when the position of the MCCs, U_0, an polarity order from experiment are used to initialise the numerical model. The numerical model reproduce the obbserved streamlines with a good agreement. Comparison is shwon for two differents heights (z=30 μm and z=90μm). The maps of angular errors quantify the orientation difference (in degree) between the local flow velocity vectors measured via PIV and the flow velocity vectors computed. (C) Comparison of the average velocity flow profiles between experiments (5 explants) and simulation.

Fig. 5. Characterisation of simulated streamlines at the embryo's scale. (A) Plotting of MCC orientation vectors on a virtual embryo to achieve an alignment order of 0.6 at 500 μm range. (B) Numerical streamlines projected in 2D, reproducing classical features observed experimentally, with an anterior–posterior flux along the flank. (C) Example of three numerically predicted streamlines showing anterior–posterior trajectories with fluctuations along the z-axis. The (x, z) dotted line plane corresponds to the cross section depicted in panel D. (D) Cross-section of the embryonic flank in the $(x-z)$ plane with Z-velocity shown as a colormap, highlighting alternating regions of upward (red) and downward (blue) flow, especially for $z < 180$ μm. (E) Projected streamlines and planar velocity as colormap for $z = 45$ μm (+ or- 15 μm). (F) Projected streamlines and planar velocity as colormap for $z = 195$ μm (+ or- 15 μm). Only partial streamline segments are captured due to fluctuations along z. (G) Mean planar velocity along streamlines as a function of height. (H) Relative standard deviation (RSD) of planar velocity and linearity as a function of height. The insert shows a streamline, with R and L definition. Linearity is defined as the ratio R/L. The color changes along the streamline represents the planar velocity variations, which are directly related to the RSD value. Error bars represent the standard deviation across the streamline population.

Variations in vertical height (along the z-axis) are primarily due to recirculation patterns in the plane. This is illustrated in Figure 5 (D), where the z-velocity is mapped, showing upward (red) and downward (blue) flow. The strongest z-velocities are observed close to the epithelial surface, at heights below 180 μm. However, importantly, the mean z-velocity along the streamlines is close to zero (Fig. S7), supporting the conclusion that the bulk flow exhibits an average two-dimensional character.

Figures 5 (E) & (F) show the streamlines at two different heights, $z = 45$ μm (+ or- 15 μm) and $z = 195$ μm (+ or- 15 μm). The color scale indicates the velocity magnitude in the epithelial plane. Since streamlines fluctuate in z, only short

segments (typically spanning about 1 mm) are captured at a given height. Consistent with explant observations, streamlines become less tortuous and flow velocities decrease with distance from the epithelial surface. Concurrently, the variability in velocity along a streamline also diminishes with height. Quantitative results (Figures 5 (G) & (H)) display the variation in streamline linearity, defined as the ratio of end-to-end displacement (R) to total length (L) (see insert of Figure 5 (H)), mean planar velocity, and its relative standard deviation (RSD), an indicator of the velocity heterogeneity, as a function of height. The plateau observed in both the RSD of planar velocity and the streamline linearity at a distance of approximatively 180μm allows us to define two distinct flow

regions: a Near-Field region (below 180 µm), where recirculation induces tortuous streamlines and inhomogeneous planar velocity, and a Far-Field region (above 180 µm), where the flow becomes more planar and uniform.

Parametric analysis: effects of simulation input parameters on flow characteristics. To understand how biological variability influences the flow driven by MCCs on the embryonic flanks, we investigated the impact of four key conditions: the characteristic velocity (U_0) of MCCs, the MCC density, their spatial organization (order parameter) and their beat direction alignment (polarity order at a 270 µm radius). The baseline conditions were $U_0 = 350, \mu$m/s, a density of $3.2 \cdot 10^{-4}$ MCC/µm^2, order parameter of 0.6, and polarity order of 0.6.
Each condition was varied as follows:

- The characteristic velocity (U_0) was reduced to 200µm/s and 100µm/s.

- The MCC density was varied to include a low-density scenario ($1.6 \cdot 10^{-4}$ MCC/µm^2) and a high-density scenario ($6.5 \cdot 10^{-4}$ MCC/µm^2).

- The spatial organization was varied by introducing MCC clustering, lowering the order parameter to 0.45, or creating an MCC-free zone (size 1 mm^2) near the head region, which also resulted in a reduced order parameter of 0.2.

- The beat direction alignment was varied to include a low-polarity scenario (polarity order of 0.4) and a high-polarity scenario (polarity order of 0.8).

Fig. 6. The mean planar velocity was computed for two representative z-slices: nearfield region (z = 45 µm) and far-field region (z = 195 µm), under various conditions illustrated schematically under the graphs.(A) Characteristic velocity U_0 at 100, 200, and 350 µm/s. (B) MCC density at low, medium, and high levels. (C) Geometrical order for ordered, clustered, and hole configurations. (D) Polarity order for low, medium and high MCC alignment. Error bars represent the standard deviation across the streamline population.

Figure 6 presents the mean planar velocity across all tested conditions at $z = 45$ µm (dark band, representing the tortuous,

near-field region) and $z = 195$ µm (light band, representing the uniform, far-field flow). The variations in RSD and linearity are presented in Supplementary Figure (Fig. S8). Qualitatively, the observed behaviors are similar at both tested z-values. The linearity of the streamlines is primarily influenced by the polarity order, while being relatively unaffected by other conditions (see Fig. SI8). Increasing U_0 yields a significant increase in planar velocities (Fig.6A), without altering the RSD (Fig. SI8). Increasing MCC density leads to a modest rise in planar velocities (Fig. 6B) and contributes to greater velocity homogeneity (reduced RSD, Fig. SI8). Likewise, increasing the polarity order raises planar velocities (Fig. 6D) and improves velocity homogeneity. In contrast, variations in the geometrical order parameter have no significant effect on the average streamline metrics. These findings suggest that flow velocity is primarily governed by the total ciliary force, density and beat direction alignment. Strikingly, the system appears robust to moderate levels of spatial disorder or small tissue defects.

5. Bacterial motion and infectability

To address the questions of epithelial clearance and its robustness against infection, we determined the trajectories of virtual swimming bacteria, modeled as point particles. Although bacteria are capable of active horizontal swimming, we note that their active horizontal velocity is negligible compared to the fast fluid flow. Therefore, we assume that their planar velocity is determined solely by the fluid streamline at their position above the epithelium. However, active vertical motion is critical: if a bacterium swims toward the epithelium, for example via chemotaxis, this velocity must be accounted for, as the mean fluid z-velocity is zero. Consequently, we introduced a tunable active downward bacterial velocity, v_{bac}.
We focus our analysis on bacteria moving above the embryo's head, as this region maximizes the transit time over the surface. This represents a worst-case scenario, offering the greatest opportunity for bacteria to reach and infect the epithelial surface.
Figure 7 (A) schematically represents the bacterial trajectories, showing the initial height (z_0) and the two possible outcomes: contact with the embryo (red) or cleared by the flow (grey). Only 3 out of 35 bacteria starting at $z_0 = 90$ µm with velocity $v_{bac} = 5$ µm/s reached the epithelial surface (Fig. 7(A)). These data were obtained from flow simulations conducted with baseline MCC conditions, $U_0 = 350, \mu$m/s, a density of $3.2 \cdot 10^{-4}$ MCC/µm^2, an order parameter of 0.6, and a polarity order of 0.6. The contact probability — defined as the fraction of bacteria reaching the epithelial surface— is shown in Figure 7(B) as a function of their initial z_0, for three distinct bacterial velocities ($v_{bac} = 0, 2, 5$ µm/s). This latter value is estimated to be among the highest for the vertical component of bacterial motility under shear flow [24–26]. The probability of reaching the epithelium is high for $z_0 < 40$ µm, which is the region where ciliary motion induces significant fluid mixing between flow layers. Above 40 µm, the contact probability decreases slowly below 10% with z_0, be-

Fig. 7. Parametric study of bacteria clearance by cilia-powered flow.(A) Schematized bacterial trajectories starting at an initial height z_0 above the embryo surface (left). A 2D projection of 35 bacterial trajectories starting at height $z_0 = 90$ μm with a bacterial speed of v_{bac}=5μm/s (right). Red trajectories indicate contact with the embryo; grey trajectories indicate successful clearance. (B&C) Contact probability as a function of the initial starting height z_0 for three different bacterial speeds (v_{bac}). Flow simulations were conducted with baseline MCC conditions, $U_0 = 350$, μm/s, a density of $3.2 \cdot 10^{-4}$ MCC/μm², an order parameter of 0.6, and a polarity order of 0.6 (B) Contact probability as a function of z_0 value. (C) Mean contact probability in $50 - 100$ μm z_0 range (light band) and $200 - 300$ μm z_0 range (dark band). Error bars correspond to the standard deviation of contact probabilities computed within each z_0 range. (D – G) Contact probability for various tested conditions and a bacterial velocity of $v_{bac} = 5$ μm/s in the two z_0 ranges. (D) Characteristic velocity U_0 at 100, 200, and 350 μm/s. (E) MCC density at low, medium, and high levels. (F) Geometrical order for ordered, clustered, and hole configurations. (G) Polarity order for low, medium and high MCC alignment. For low polarity order conditions, the contact probabilities for both z ranges overlap. Error bars correspond to the standard deviation of contact probabilities computed within each z_0 range.

coming zero at $z_0 = 150$ μm, for non-motile bacteria $v_{bac} = 0$ (Figure 7 (B) & (C)). In contrast, higher bacterial downward velocities consistently result in increased contact probabilities across all starting heights, maintaining a non-zero probability even at large z_0. This highlights that motile bacteria can reach the epithelium, whereas non-motile bacteria are efficiently cleared by the fluid flow.

The influence of ciliary velocity, U_0, MCC density, their spatial distribution and their beat direction alignment on the contact probability is illustrated in Figure 7 (D-G) for motile bacteria with velocity $v_{bac} = 5$ μm/s. This analysis clearly shows that U_0 is a strong protective factor, significantly reducing contact probability. Conversely, the spatial distribution (order parameter) has a limited effect; only the introduction of a large defect (hole) near the head region, which disrupts the local flow, has a noticeable impact on clearance (Fig.7 F). A low density of MCC results in a significant increase of contact probability, while higher densities provide a better protection (Fig.7 E). The polarity order also influences contact probability, with lower polarity (i.e. low beat direction alignment) associated with a higher contact probability (Fig.7 G).

Discussion

Three key points emerge from our results and warrant detailed discussion.

First, we address the seemingly contradictory observation of the high survival rate of healthy embryos despite the arrival

of some environmental bacteria onto the epithelial surface. Indeed, our physical model shows that even for bacteria lacking active movement towards the epithelium, the mixing of fluid layers near the surface ($< 150 \mu m$) allows a proportion of bacteria to contact the epithelial surface. We can estimate the rate of bacteria landing on the epithelium per unit of time for a typical embryo (MCC density = $3.2 \cdot 10^{-4}$ MCC/μm², $U_0 = 350$, μm/s, polarity = 0.6, and order = 0.6). The fluid volume passing over the embryo at its head (at a height z of $300 \mu m$) per unit of time is obtained by integrating the flow rate per 30-μm layer, based on numerical data. Assuming a typical bacterial concentration in pond water of 10^7 bacteria/ml (27), we deduce the bacterial arrival rate from our simulations. For non-vertically motile bacteria, this number is on the order of 50 bacteria/s. Considering that only 10% of these might be pathogenic, approximately 5 pathogenic bacteria per second land on each flank of the embryo. Given a flank surface area of about $5\,\text{mm}^2$, this equates to 1 bacteria $\cdot$ mm$^{-2} \cdot$ s^{-1}. These arriving bacteria first contact the 6-μm-thick mucus layer covering the embryo's skin, which possesses known antibacterial properties (17). Their subsequent elimination upon arrival provides a first, albeit coarse, estimate of the bacterial load that the mucus is capable of clearing per unit of time. The strong ciliary flow is coupled with the action of the mucus layer to successfully clean the surface and eliminates the arriving microbial load.

The second addressed point concerns the initial question of

whether the physical system is optimized for maximum clearance. Our results show that substantial perturbations to the spatial distribution of MCCs do not significantly affect bacterial clearance. Therefore, this specific cellular arrangement is not optimized to maximize the clearance rate. More generally, the system's characteristics (U_0, MCC density, polarity, and order) must be perturbed quite significantly to substantially impact the probability of bacterial contact. The biological system does not seek to optimize these parameters for maximum clearance but rather to ensure high robustness. This robustness allows embryos to develop without infection even if their physiological characteristics are somewhat degraded or if the environment is more hostile (e.g., higher bacterial concentration). This observation suggests that optimizing 'performance' metrics—such as speed or energy efficiency—may not be the primary guiding principle for the biological systems investigated here. Instead, these systems prioritize the ability to maintain function and survive even under compromised conditions, such as those resulting from injury, environmental stress, or developmental issues.

Finally, we reflect on the functional implications of the highly homogeneous spatial distribution of multiciliated cells, which is conserved feature in the embryonic skin of amphibian species (2). Our data suggest that the global flow powered by MCC collectives, and the associated clearance capacity, are not dependent on MCC geometrical order. It remains possible, however, that regular MCC distribution matters at a more local scale, and closer to the skin. Indeed, our data reveal fluctuating non-zero vertical flow velocities near ciliary bundles, that may optimize oxygen uptake. For such function, regular MCC spacing would appear as an optimal solution. Moreover, MCC distribution in the superficial cell layer is expected to indirectly impact the distribution of secretory cells, which may also be an important factor for functional mucociliary protection. A final potential implication that we would like to mention is linked to the death of MCCs at pre-metamorphic stages. Death and removal of cells in epithelia represent a challenge, as junctions with surrounding cells are temporarily lost and must be rebuilt to preserve tissue integrity (28). The concomittant death of clustered MCCs would represent a major risk of epithelium disintegration, which would be avoided through regular paving of MCCs.

In conclusion, by combining experimental measurements, on both embryos and in-vitro explants, with a computational fluid dynamics model, we contributed to a better understanding of the critical biophysical mechanisms involved in the protection of Xenopus embryo at early stages. Our parametric analysis revealed that the efficiency of pathogen clearance is mainly governed by the intrinsic flow velocity U_0 of MCCs. Significantly, the system exhibits remarkable functional robustness, as clearance efficiency remains high and largely insensitive to moderate variations in MCC density, spatial arrangement, and beat direction alignment. This finding suggests that the biological mechanism prioritises system resilience and reliable protection under non-ideal or compromised conditions over the optimisation of energy consump-

tion. The question of the role of the specific tissue patterning of MCCs remains open.

Materials and Methods

Xenopus laevis. All experiments were performed following the Directive 2010/63/EU of the European parliament and of the council of 22 September, 2010 on the protection of animals used for scientific purposes and approved by the "Direction départementale de la Protection des Populations, Pôle Alimentation, Santé Animale, Environnement, des Bouches du Rhône" (agreement number G 13055 21). Adult Xenopus laevis individuals were obtained from NASCO. Female frogs were injected with human chorionic gonadotropin a day before egg collection, and eggs were externally fertilized in a petri dish with male sperm. Embryos were then dejellied with a 2% cysteine hydrochloride solution (pH 8), rinsed, and kept in 0.1X MBS solution (Modified Barth's Saline) at 13 °C until they reached the desired developmental stage.

Foxj1 CRISPR/Cas9 injection. Following in vitro fertilization at 18 °C, eggs were dejellied with 2% cysteine hydrochloride solution (pH 8), rinsed a few times with tap water and once with 0.1X MBS. Eggs were then transferred to a 4% Ficoll solution at the 1-cell stage and injected with 5 nl of recombinant Cas9 (0.2 ng/nl, PNA Bio, #CP03) or Cas9 + foxJ1 single guide RNA (0.06 ng/nl; 5'-GCAGCTAATACGACTCACTATAGGGATACATACCTGCC AGGTGTTTTAGAGCTAGAAATAGCAAG-3') (29) near the centre of the animal pole. Injected embryos were kept at 23 °C for at least 3 hours before transferring them to 2% Ficoll at the desired temperature overnight. The next day, they were transferred back to 0.1X MBS for subsequent steps.

Flow measurement on embryos. Tadpoles were kept in anaesthetic (0.02% MS-222 in 0.1X MBS) lying on their sides. 10 μm fluorescent beads (Invitrogen) were then added near the top of the head using a 2 μl syringe (Hamilton) and videos of the moving beads were recorded using an epifluorescence microscope (Nikon SMZ18) at 3X magnification attached with a digital camera (Hamamatsu ORCA-fusion C14440) using 10 ms exposure. A small region of interest near the middle of the tadpoles was analysed using the TrackMate plugin (Kalman tracker) in ImageJ to quantify the mean speed of each bead.

Imaging multiciliated cells in embryos. To reveal MCCs, embryos were fixed at stage 30 in PFA 4% in PBS-Triton 0.1% during 30min at 23°C, rinsed 4 times in 1X PBS and blocked in BSA 3% in PBS for 1h at 23°C. Incubation with mouse anti-acetylated-Tubulin (Sigma T7451, 1/1000) was done overnight at 4°C. The next day, embryos were rinsed 4 times with 1X PBS and washed 3 times for 30 min in 1X PBS. Incubation with goat anti-mouse IgG2a-568 (Life Technologies, 1:800) was done 1h at 4°C. Embryos were rinsed 4 times in 1X PBS and washed 3 times in 1X PBS for 5min. Embryos were imaged using an epifluorescence microscope

(Nikon SMZ18) at 3X magnification attached with a digital camera (Hamamatsu ORCA-fusion C14440).

Quantification of infection rates by Aeromonas hydrophila. A. hydrophila (Chester Stanier, ATCC®7966; recombinant expressing GFP and kanamycin resistance) was kindly provided by Eamon Dubaissi (University of Manchester, UK). Stock colonies were maintained on LB agar plates supplemented with 50 µg/ml kanamycin at 4 °C for a month. A. hydrophila was freshly grown in Tryptic Soy Broth (TSB) liquid culture (50ml) supplemented with kanamycin (50 µg/ml) to OD600=0.8 to 1 (30°C, 232rpm). Bacteria were pelleted by centrifugation 10min at 3500xg, washed in MBS 0.1x to remove traces of TSB, pelleted again and resuspended in MBS 0.1x supplemented with Kanamycin 50 µg/ml to OD=0.8 (approximately 5.10^8 bacteria/ml). Stage 30 embryos were exposed to 3ml of bacteria suspension (5 embryos/well in 12-well dishes) for 9h at 23°C. At the end of the infection period, embryos were extensively washed to eliminate bacteria in suspension (5 times in 50ml tap water), and individually crushed in 500ml TBS in 1.5ml Eppendorf tubes with a tissue homogenizer. 100ml of embryo lysate was spread on a LB Agar plate supplemented with kanamycin 50mg/ml. Plates were incubated overnight at 37°C, after which colonies were manually counted to quantify colony-forming units per infected embryo. Individual embryos were arbitrarily considered infected at cfu>100 per embryo. Controls included plating of the initial bacterial suspension and of the culture medium at the end of the infection period to check bacterial health, and plating of the last embryo wash to verify washing efficiency.

Explant culture. The explantation protocol was adapted from established methods (30), to ensure that the tissue remained thin, flat, and transparent. Embryos kept at 13 °C were transferred to a Petri dish containing explant culture medium (Danilchik's for Amy (DFA): 53 mM NaCl2, 5 mM Na2CO3, 4.5 mM potassium gluconate, 32 mM sodium gluconate, 1mM CaCl2, 1mM MgSO4 buffered to pH 8.3 using 1M Bicine). At stage 16, vitelline membranes were carefully removed with tweezers, the neural cord was cut using an eyebrow knife, allowing to explant a piece of ectoderm, from which the adjoined mesodermal layer was peeled away. Explants were placed in imaging chambers made with collagen-coated coverslips and PDMS walls, bonded by plasma treatment. The chambers were pre-treated with fibronectin (20 mg/ml in 1X PBS) overnight at 4°C, blocked with 1X PBS + 1% BSA for one hour, and filled with explant culture medium containing antimycotic (100 µg/ml) to prevent contaminations. To ensure flattening of the tissue, sterilized 10×10 mm glass slides coated with silicone grease were gently placed on top of the explant for 12 h, then removed to avoid disrupting cilia growth. Sibling embryos were grown in parallel to monitor development and to stage explants. The culture medium was replaced daily, and all solutions and dissection tools were sterilized before use. Explants were cultured for 3–5 days before imaging, reaching stage 32.

Axitinib treatment.. We disrupted MCC distribution in Xenopus embryos and explants using Axitinib (Sigma), a Kit receptor chemical inhibitor known to block the cell migration responsible for their regular spacing (22). Embryos at NF stage 9-10 were incubated in 0.1x MBS containing 50 µM Axitinib diluted in DMSO, and allowed to develop until NF stage 29–30. For explants, embryos were treated with Axitinib at NF stage 9/10, then explanted at NF stage 16 and transferred to explant culture medium containing 50 µM Axitinib.

Imaging explants and Flow measurements. Observations were made using a Nikon Eclipse Ti inverted optical microscope equipped with an Andor Neo 5.5 sCMOS camera, offering a frame rate above 100 fps, sufficient to visualize ciliary beating. Images were captured using Micromanager 1.4 software with a 20X objective (CFI Plan Fluor LWD 20X; N.A. 0.70). To image flow fields, fluorescent beads (FluoSpheres Polystyrene Microspheres, 1.0 μm, red fluorescent (580/605), #F13083 thermofisher) were added to the medium at a 1:1000 dilution.

Image analysis. For every field of view, two videos were acquired: one in bright field to visualize the dynamics of MCC and the other in fluorescence, in presence of beads, for recording the flow fields associated with those MCCs. All the image processing was done using ImageJ, JPIV and custom-made Python scripts.

Quantification of MCC position and beat direction. MCC positions are identified from raw videos by computing the temporal standard deviation projection of the image stack, which highlights cilia activity as regions of maximum pixel intensity. These projections can be binarized in ImageJ, showing MCCs as white patches, with MCC position calculated as the centroid of these patches. Cilia-driven flow is confirmed to align with the underlying cilia's beating direction (31), allowing the average beating direction of each MCC to be obtained by correlating cilia activity videos with those of bead movement. The resulting beating directions are spatially averaged across all cilia in an MCC.

Quantification of MCC area and coverage fraction. The dynamic area of MCC is measured by segmenting regions of ciliary activity, identified as white patches in the standard deviation of video frames. The coverage fraction is calculated as the area with ciliary activity divided by the total tissue area analyzed. To compare coverage fraction of MCCs on explants (live microscopy) whith embryos (immunostaining), the dynamic area can be adjusted to reflect the apical membrane area of MCCs by applying a correction term based on cilia length, following the method outlined in (32).

Quantification of cilia beating frequency. We compute the CBF using an in-house routine developed in python. We perform image processing on movies acquired at 150 fps. For each pixel belonging to a ciliated cell, we extract the time evolution of the intensity of the pixel. Then, we compute

the power spectrum of this signal via the fast fourier transform. The CBF is the frequency with the highest energy in the power spectrum. The frequencies across pixels within a MCC are then averaged to obtain the mean ciliary beat frequency at the cell level.

Quantification of Geometrical order. To quantify the randomness in MCC distribution, a geometrical order parameter is defined based on the pairwise distances between MCCs, calculated using Delaunay triangulation (See Fig. 2 (B)). From the distribution of these distances, the mean and standard deviation are computed. The order parameter is then given by:

$$\text{GO} = 1 - \left(\frac{\text{Standard deviation of the lengths}}{\text{mean of the lengths}} \right)$$

A value near zero suggests a random, spread-out distribution, while a value near one indicates a more uniform, regular arrangement of MCCs.

Quantification of Orientational order. Orientational order quantifies the alignment of beat directions of MCCs. It is determined by calculating the length of the mean resultant vector of the unit orientation vectors p_i for each MCC. Using circular statistics, the average of these orientation vectors provides the orientational order, with higher values indicating a more aligned beating directions. The mean $\vec{P}$ of a set of N angles $(\gamma_1, \gamma_2, ... \gamma_N)$ is given by

$$\langle \vec{P} \rangle = \frac{1}{N} \sum_{i=1}^{N} \vec{P}_i = (\langle \cos \gamma \rangle, \langle \sin \gamma \rangle) \quad \textbf{(1)}$$

The order is then calculated as the length of the mean resultant vector

$$\mathbf{P} = || \langle \vec{P} \rangle || \quad \textbf{(2)}$$

(33)(34) The order parameter reaches one when all beating directions are perfectly parallel and approaches zero for randomly oriented vectors. To measure the correlation of polarity over distance, we calculate polarity at various length scales using circles of different radii. Starting with a specific radius, polarity is computed within each circle. The radius is then incremented in discrete steps, calculating polarity for each new radius. This approach provides an average polarity value at each distance, derived from multiple circles of the same radius.

Analysis of the flow. We quantified the flow fields via Particle Image Velocimetry (PIV), using the JPIV software. Briefly, each video frame is divided into interrogation windows, and displacement in each window is calculated as the peak of the autocorrelation function. For a 2048x2048 pixel field of view, we used 128-pixel interrogation windows with two iterations, providing a velocity vector for every 16 pixels. These fields were temporally averaged over 100 frames, equivalent to about 1 second of acquisition.

Numerical model. We developed an in-silico model of epithelial tissue was using a fluid solver, employing the three-dimensional Lattice Boltzmann method for the fluid domain and integrating ciliated cells through Lagrangian points with the Immersed Boundary Method (IBM). This combined approach has been validated for simulating rigid and deformable bodies (35)(36)(37). In this study, ciliated cells are represented as point forces at varying heights, comparable to cilia length. Input parameters include the positions (centroids of MCCs from experiments), directions (measured beating directions, and magnitudes of the point forces, derived from the characteristic velocities (Uo) measured above each MCC.

The numerical method to calculate this point force F is the immersed boundary method. The force is:

$$F(X,t) = \frac{2}{\Delta t}(I[\rho(x,t)]Uo(X,t) - I[\sum_i c_i f_i(x,t)]) \quad \textbf{(3)}$$

where X denotes the position of the MCC, ρ is the density of the fluid which is water here, I is an interpolating operator using numerical points surrounding the MCC to integrate values such as the density, Uo is the characteristic velocity measured above each MCC, and c_i and f_i are related to the LBM (Lattice Boltzmann Method), respectively the fluid particle velocity and the fluid distribution function.

The fluid domain features a solid wall at the bottom to simulate the epithelial surface, enforcing a no-slip condition, so the fluid flow is not interrupted by any barriers. Point forces are centered within the simulation domain to minimize boundary effects, with optimal distances between the open boundary and the epithelial domain determined through multiple trials with varying distances (Fig. S6). A grid size of dx=dy=dz=$20\mu m$ was chosen, reflecting the size of the MCCs and optimized using experimental data. The fluid used in the simulation, driven by ciliated cells in Xenopus embryos, is water. Therefore, the density of the medium is set at ρ = 1000 kg/m³ and the kinematic viscosity (μ) = 1×10^{-6} m²/s. The Reynolds number, calculated as $Re = \frac{V \cdot L}{\mu}$, is approximately 0.01, considering a velocity of 1 mm/s over a length scale of 10 μm. Such a low Re indicates that the flow is in the Stokes regime.

Bibliography

1. G A Deblandre, D A Wettstein, N Koyano-Nakagawa, and C Kintner. A two-step mechanism generates the spacing pattern of the ciliated cells in the skin of Xenopus embryos. *Development*, 126(21):4715–4728, 11 1999. ISSN 1477-9129. doi: 10.1242/dev.126.21.4715.
2. M Nokhbatolfoghahai, J R Downie, A K Clelland, and K Rennison. The surface ciliation of anuran amphibian embryos and early larvae: Patterns, timing differences and functions. *Journal of Natural History*, 39(12):887–929, 2005. ISSN 00222933. doi: 10.1080/00222930400002937.
3. Peter Walentek. Manipulating and analyzing cell type composition of the xenopus mucociliary epidermis. In *Xenopus: Methods and Protocols*, pages 251–263. Springer, 2018.
4. Gerd König and Peter Hausen. Planar Polarity in the Ciliated Epidermis of Xenopus Embryos. *Developmental Biology*, 160(2):355–368, 11 1993. ISSN 00121606. doi: 10.1006/dbio.1993.1312.
5. Brian Mitchell, Richard Jacobs, Julie Li, Shu Chien, and Chris Kintner. A positive feedback mechanism governs the polarity and motion of motile cilia. *Nature*, 447(7140):97–101, 11 2007. ISSN 0028-0836. doi: 10.1038/nature05771.
6. S Nishikawa, J Hirata, and F Sasaki. Fate of ciliated epidermal cells during early development of Xenopus laevis using whole-mount immunostaining with an antibody against chondroitin 6-sulfate proteoglycan and anti-tubulin: transdifferentiation or metaplasia of

amphibian epidermis. *Histochemistry*, 98(6):355–358, 1992. ISSN 0301-5564. doi: 10.1007/BF00271070.

7. Rosa Ventrella, Sun K Kim, Jennifer Sheridan, Aline Grata, Enzo Bresteau, Osama A Hassan, Eve E Suva, Peter Walentek, and Brian J Mitchell. Bidirectional multiciliated cell extrusion is controlled by Notch-driven basal extrusion and Piezo1-driven apical extrusion. *Development (Cambridge, England)*, 150(17), 9 2023. ISSN 1477-9129. doi: 10.1242/dev.201612.

8. Alexia Tasca, Martin Helmstädter, Magdalena Maria Brislinger, Maximilian Haas, Brian Mitchell, and Peter Walentek. Notch signaling induces either apoptosis or cell fate change in multiciliated cells during mucociliary tissue remodeling. *Developmental cell*, 56(4):525–539, 2021.

9. Richard R Strathmann and Chet Chaffee. Constraints on egg masses. II. Effect of spacing, size, and number of eggs on ventilation of masses of embryos in jelly, adherent groups, or thin-walled capsules. *Journal of Experimental Marine Biology and Ecology*, 84(1):85–93, 11 1984. ISSN 00220981. doi: 10.1016/0022-0981(84)90232-6.

10. R G Kessel, H W Beams, and C Y Shih. The origin, distribution and disappearance of surface cilia during embryonic development ofRana pipiens as revealed by scanning electron microscopy. *American Journal of Anatomy*, 141(3):341–359, 11 1974. ISSN 0002-9106. doi: 10.1002/aja.1001410306.

11. M Nokhbatolfoghahai, J R Downie, and V Ogilvy. Surface ciliation of anuran amphibian larvae: Persistence to late stages in some species but not others. *Journal of Morphology*, 267(10):1248–1256, 11 2006. ISSN 0362-2525. doi: 10.1002/jmor.10469.

12. Ingrid Reiten, Fazil Emre Uslu, Stephanie Fore, Robbrecht Pelgrims, Christa Ringers, Carmen Diaz Verdugo, Maximillian Hoffman, Pradeep Lal, Koichi Kawakami, Kerem Pekkan, Emre Yaksi, and Nathalie Jurisch-Yaksi. Motile-Cilia-Mediated Flow Improves Sensitivity and Temporal Resolution of Olfactory Computations. *Current Biology*, 27(2):166–174, 11 2017. ISSN 09609822. doi: 10.1016/j.cub.2016.11.036.

13. A Kemp. Role of epidermal cilia in development of the Australian lungfish,Neoceratodus forsteri (Osteichthyes: Dipnoi). *Journal of Morphology*, 228(2):203–221, 11 1996. ISSN 0362-2525. doi: 10.1002/(SICI)1097-4687(199605)228:2<203::AID-JMOR9>3.0.CO;2-5.

14. Aude Nommick, Camille Boutin, Olivier Rosnet, Claire Schirmer, Elsa Bazellières, Virginie Thomé, Etienne Loiseau, Annie Viallat, and Laurent Kodjabachian. Lrrcc1 and ccdc61 are conserved effectors of multiciliated cell function. *Journal of cell science*, 135(4):jcs258960, 2022.

15. M Nokhbatolfoghahai, J R Downie, and V Ogilvy. Surface ciliation of anuran amphibian larvae: Persistence to late stages in some species but not others. *Journal of Morphology*, 267(10):1248–1256, 11 2006. ISSN 0362-2525. doi: 10.1002/jmor.10469.

16. Jacques Robert and Yuko Ohta. Comparative and developmental study of the immune system in <i>Xenopus</i>. *Developmental Dynamics*, 238(6):1249–1270, 11 2009. ISSN 10588388. doi: 10.1002/dvdy.21891.

17. Eamon Dubaissi, Karine Rousseau, Gareth W Hughes, Caroline Ridley, Richard K Grencis, Ian S Roberts, and David J Thornton. Functional characterization of the mucus barrier on the xenopus tropicalis skin surface. *Proceedings of the National Academy of Sciences*, 115 (4):726–731, 2018.

18. Michael E Werner, Peter Hwang, Fawn Huisman, Peter Taborek, Clare C Yu, and Brian J Mitchell. Actin and microtubules drive differential aspects of planar cell polarity in multiciliated cells. *Journal of Cell Biology*, 195(1):19–26, 11 2011. ISSN 1540-8140. doi: 10.1083/jcb.201106110.

19. Brian Mitchell, Jennifer L Stubbs, Fawn Huisman, Peter Taborek, Clare Yu, and Chris Kintner. The PCP Pathway Instructs the Planar Orientation of Ciliated Cells in the Xenopus Larval Skin. *Current Biology*, 19(11):924–929, 11 2009. ISSN 09609822. doi: 10.1016/j.cub.2009.04.018.

20. Peter Walentek, Camille Boutin, Laurent Kodjabachian, and D Houston. Planar cell polarity in ciliated epithelia. In *Cell Polarity in Development and Disease*, pages 177–209. Elsevier, 2017.

21. Francesco Boselli, Jerome Jullien, Eric Lauga, and Raymond E Goldstein. Fluid Mechanics of Mosaic Ciliated Tissues. *Physical Review Letters*, 127(19):198102, 11 2021. ISSN 0031-9007. doi: 10.1103/PhysRevLett.127.198102.

22. Alexandre Chuyen, Charlotte Rulquin, Fabrice Daian, Virginie Thomé, Raphaël Clément, Laurent Kodjabachian, and Andrea Pasini. The Scf/Kit pathway implements self-organized epithelial patterning. *Developmental Cell*, 56(6):795–810, 11 2021. ISSN 15345807. doi: 10.1016/j.devcel.2021.02.026.

23. Xianwen Yu, Chee Peng Ng, Hermann Habacher, and Sudipto Roy. Foxj1 transcription factors are master regulators of the motile ciliogenic program. *Nature genetics*, 40(12): 1445–1453, 2008.

24. Jinyou Yang, Kenji Kikuchi, and Takuji Ishikawa. High shear flow prevents bundling of bacterial flagella and induces lateral migration away from a wall. *Communications Physics*, 6 (1):354, 2023.

25. Guanju Wei, Jessica-Jae S Palalay, Joseph E Sanfilippo, and Judy Q Yang. Flagellum-driven motility enhances pseudomonas aeruginosa biofilm formation by altering cell orientation. *Applied and Environmental Microbiology*, 91:e00821–25, 2025.

26. Jacinta C Conrad. Physics of bacterial near-surface motility using flagella and type iv pili: implications for biofilm formation. *Research in Microbiology*, 163(9-10):619–629, 2012.

27. Chieko Sakamoto, Nobuyasu Yamaguchi, and Masao Nasu. Rapid and simple quantification of bacterial cells by using a microfluidic device. *Applied and environmental microbiology*, 71 (2):1117–1121, 2005.

28. Romain Levayer. Staying away from the breaking point: Probing the limits of epithelial cell elimination. *Current Opinion in Cell Biology*, 86:102316, 2024.

29. EV Rachev, Karin Schuster-Gossler, Franziska Fuhl, Tim Ott, Lena Tveriakhina, Anja Beckers, Jan Hegermann, Karsten Boldt, Michaela Mai, Elisabeth Kremmer, et al. Cfap43 modulates ciliary beating in mouse and xenopus. *Developmental biology*, 459(2):109–125, 2020.

30. Michael E Werner and Brian J Mitchell. Using Xenopus Skin to Study Cilia Development and Function. *Methods in Enzymology*, 525:191–217, 11 2013. ISSN 0076-6879. doi: 10.1016/B978-0-12-397944-5.00010-9.

31. Regina Faubel, Christian Westendorf, Eberhard Bodenschatz, and Gregor Eichele. Cilia-based flow network in the brain ventricles. *Science*, 353(6295):176–178, 11 2016. ISSN 0036-8075. doi: 10.1126/science.aae0450.

32. Nicola Pellicciotta, Evelyn Hamilton, Jurij Kotar, Marion Faucourt, Nathalie Delgehyr, Nathalie Spassky, and Pietro Cicuta. Entrainment of mammalian motile cilia in the brain with hydrodynamic forces. *Proceedings of the National Academy of Sciences*, 117(15): 8315–8325, 11 2020. ISSN 0027-8424. doi: 10.1073/pnas.1910065117.

33. Philipp Berens. CircStat : A <i>MATLAB</i> Toolbox for Circular Statistics. *Journal of Statistical Software*, 31(10), 2009. ISSN 1548-7660. doi: 10.18637/jss.v031.i10.

34. Martin Schneiter, Sebastian Halm, Adolfo Odriozola, Helga Mogel, Jaroslav Rička, Michael H Stoffel, Benoît Zuber, Martin Frenz, and Stefan A Tschanz. Multi-scale alignment of respiratory cilia and its relation to mucociliary function. *Journal of Structural Biology*, 213 (1):107680, 11 2021. ISSN 10478477. doi: 10.1016/j.jsb.2020.107680.

35. Julien Favier, Alistair Revell, and Alfredo Pinelli. A Lattice Boltzmann–Immersed Boundary method to simulate the fluid interaction with moving and slender flexible objects. *Journal of Computational Physics*, 261:145–161, 11 2014. ISSN 00219991. doi: 10.1016/j.jcp.2013. 12.052.

36. Zhixiang Liu, Huichao Liu, Dongmei Huang, and Liping Zhou. The Immersed Boundary-Lattice Boltzmann Method Parallel Model for Fluid-Structure Interaction on Heterogeneous Platforms. *Mathematical Problems in Engineering*, 2020:1–13, 11 2020. ISSN 1024-123X. doi: 10.1155/2020/3913968.

37. Simon Gsell and Julien Favier. Direct-forcing immersed-boundary method: A simple correction preventing boundary slip error. *Journal of Computational Physics*, 435:110265, 11 2021. ISSN 00219991. doi: 10.1016/j.jcp.2021.110265.

Supplementary Figure S1

Images of Axitinib treated embryos with MCC denoted as white dots. Embryos 5-9 exhibit large regions devoided of MCC due to MCC disappearance

Supplementary Figure S2

Comparison of ciliated epithelium at day 5 and day 6 of explant maturation. Fig. (a) and (a') show ciliated cells from day 5 explants, while Fig. (b) and (b') show ciliated cells from day 6 explants. The ciliated cell area and beating frequency were analyzed across multiple tiles, revealing that ciliated cells on day 6 are larger and exhibit stronger beating compared to day 5. The corresponding velocity fields generated by day 5 and day 6 explants are shown in Fig. (e) and (e').

Supplementary Figure S3

Fig. 8. Images of untreated embryos with MCC denoted as white dots. These are confocal images where cilia are marked by staining Ac tub and binarised to see the MCC distribution.

Supplementary Figure S4

Images of explant tissues with MCC denoted as white dots

Supplementary Figure S5

(a) Uo distribution for explant 8 under untreated conditions. (b) Uo distribution for explant 9 under treated conditions. (c) Zoomed-in image showing the variation of Uo within a tile. (d) and (e) show Uo distributions across different explants. (d) Comparison of mean Uo between untreated and Axitinib-treated explants.

Supplementary Figure S6

(a) Periodic Boundaries (a') Open Boundaries (a)'' Open Boundaries with point forces at center

(c)

Tissue[a]	Height	Point force at 20µm (Error in Vmag)	Point force at 30µm (Error in Vmag)	Point force at 40µm (Error in Vmag)
Tissue 1	at height30	47.2µm/s	47.6µm/s	47.6µm/s
	at height60	47.8µm/s	47.6µm/s	53.7µm/s
	at height90	33.9µm/s	36.7µm/s	36.7µm/s
Tissue 2	at height30	60.6µm/s	61.4µm/s	65.1µm/s
	at height60	50.37µm/s	55.5µm/s	67.2µm/s
	at height90	40.3µm/s	40.7µm/s	50.4µm/s
Tissue 3	at height30	54.4µm/s	57.3µm/s	65.2µm/s
	at height60	58.1µm/s	65.7µm/s	64.8µm/s
	at height90	51.6µm/s	43.7µm/s	48µm/s

(a)Velocity profiles for different boundary conditions. The average velocity was compared across cases, with the configuration of centrally placed point forces and open boundaries showing the closest agreement with experimental results. The distance to the border was optimized through trial-and-error until boundary effects were eliminated.(b) Grid size in the Lattice Boltzmann method defined the effective size of a multiciliated cell (MCC); a grid size of 20 µm corresponded to an MCC area of 400 µm², comparable to biological values. Velocity profiles above each point force were examined for grid sizes of 20 µm and 10 µm. With a 20 µm grid and point forces placed at 20 µm, the maximum velocity occurred at 40 µm, whereas at a 10 µm grid with point forces at 20 µm, the maximum velocity shifted to 20 µm due to mesh refinement. Profiles obtained with a 20 µm grid resembled experimental data more closely, and this grid size was therefore adopted. (c) To mimic cilia height, point forces were positioned at different values of h. Three values of h were tested across three tissues, and while variation was minimal, h = 20 µm consistently yielded the lowest mean squared error (MSE) and was used in all subsequent analyses.

Supplementary Figure S7

Mean componant of velocity in x, y, z direction on a population of approx. 400 streamlines

Mean planar velocity, coefficient variation (CV=RSD/100), and linearity (R/L) for all U_0, density, geometrical order, and polarity conditions, plotted against the height z up to z=300 μm. Each parameter was varied startig from the baseline conditions ($U_0 = 350, \mu$m/s, a density of $3.2 \cdot 10^{-4}$ MCC/μm^2, an order parameter of 0.6 and a polarity order of 0.6), expect for polarity study where the baseline conditions where ($U_0 = 200, \mu$m/s, a density of $3.2 \cdot 10^{-4}$ MCC/μm^2, an order parameter of 0.6 and a polarity order of 0.6)

Supplementary Figure S9

RSD and Linearity with U_0, density, order and polarity variations, for two representative z-slices: near-field region ($z = 45$ μm) and far-field region ($z = 195$ μm). Error bars represent the standard deviation across the streamline population.

Supplementary Table S1

Characteristics of all embryos analyzed: Untreated. MCC area quantification for some embryos was not performed because ZO1 stained images were unavailable. Size is defined as overall area covered by the embryo.

Embryo ID	Size (mm^2)	Number of MCCs	Average MCC area (μm^2)	Coverage fraction of MCC	Number density (MCCs/mm
Embryo 1	5.01	1719	252	0.094	3.42
Embryo 2	5.10	1485	257	0.075	2.91
Embryo 3	4.67	1429	380	0.065	3.05
Embryo 4	4.16	1334	240	0.047	3.20
Embryo 5	4.31	1564	203	0.071	3.62
Embryo 6	4.80	1889	—	—	3.93
Embryo 7	4.11	1319	185	0.050	3.19
Embryo 8	4.36	840	240	0.044	1.92

Supplementary Table S2

Characteristics of all embryos analyzed: Axitinib-treated. MCC area quantification for some embryos was not performed because ZO1 stained images were unavailable. Size is defined as overall area covered by the embryo.

Embryo ID	Size(mm^2)	Number of MCCs	Average MCC area (μm^2)	Coverage fraction of MCC	Number density (MCCs/mm^2)
Embryo 1	4.85	1626	—	—	3.34
Embryo 2	3.91	1338	222	0.059	3.42
Embryo 3	3.97	1180	223	0.044	2.96
Embryo 4	4.21	1420	283	0.040	3.36
Embryo 5	5.03	796	188	0.035	1.57
Embryo 6	4.64	871	—	—	1.86
Embryo 7	4.35	854	170	0.028	1.96
Embryo 8	3.97	1635	160	0.071	4.10
Embryo 9	3.18	1359	228	0.040	4.30
Embryo 10	3.97	1413	304	0.046	3.54
Embryo 11	3.56	1157	—	—	3.25
Embryo 12	4.56	1437	—	—	3.15

Supplementary Table S3

Characteristics of all explant tissues: Untreated explants. For small explants, size is reported as "number of tiles x field of view"; for large explants, absolute area is shown.

Explant ID	Size	Number of MCCs	Average MCC area (μm^2)	Coverage fraction	Number density (MCCs/mm
Explant 1	$1 \times (680 \times 680)\,\mu m^2$	33	506	0.036	0.714
Explant 2	$1 \times (680 \times 680)\,\mu m^2$	21	697	0.031	0.454
Explant 3	$2 \times (680 \times 680)\,\mu m^2$	59	840	0.053	0.638
Explant 4	$2 \times (680 \times 680)\,\mu m^2$	81	710	0.062	0.876
Explant 5	$3 \times (680 \times 680)\,\mu m^2$	113	600	0.057	0.815
Explant 6	$2 \times (680 \times 680)\,\mu m^2$	277	497	0.041	0.833
Explant 7	$5.18\ mm^2$	262	675	0.046	0.506
Explant 8	$3.72\ mm^2$	300	580	0.033	0.827
Explant 9	$7.9\ mm^2$	540	735	0.043	0.683

Supplementary Table S4

Characteristics of all explant tissues: Treated explants. For small explants, size is reported as "number of tiles × field of view"; for large explants, absolute area is shown .

Explant ID	Size	Number of MCCs	Average MCC area (μm^2)	Coverage fraction	Number density (MCCs/mm^2)
Explant 1	$1 \times (680 \times 680)\,\mu$m^2	28	536	0.032	0.606
Explant 2	$1 \times (680 \times 680)\,\mu$m^2	16	581	0.020	0.346
Explant 3	$2 \times (680 \times 680)\,\mu$m^2	50	475	0.025	0.541
Explant 4	$1 \times (680 \times 680)\,\mu$m^2	25	540	0.029	0.540
Explant 5	$2 \times (680 \times 680)\,\mu$m^2	56	455	0.027	0.606
Explant 6	1.08 mm^2	88	400	0.037	0.816
Explant 7	1.97 mm^2	280	405	0.056	1.412
Explant 8	1.75 mm^2	194	407	0.044	1.103
Explant 9	2.05 mm^2	340	402	0.060	1.654

Supplementary Moovie S1

Beating cilia at the surface of an explant, 6 days after explantation. Immobile background has been substracted for a better visualisation of cilia.

Supplementary Moovie S2

Fluorescent beads transported at the surface of an explan at a distance from the epithelial surface z=20μm.