1 # Pooled CRISPRi screening reveals fungal-specific vulnerabilities across
2 # environments and genetic backgrounds
3
4

5 Lauren F Wensing*[1], Philippe C Després*[1,2,3], Desiree Francis[1], Meea Fogal[1], Anthony
6 Hendriks[1], Nicholas C Gervais[1], Clara Fikry[1], Abdul-Rahman Adamu Bukari[4], Aleeza C
7 Gerstein[4], Christina A Cuomo[2,3], Rebecca S Shapiro[1]
8
9 *These authors contributed equally to this work.
10
11   1. Department of Molecular and Cellular Biology, University of Guelph, Guelph, Canada
12   2. Infectious Disease and Microbiome Program, Broad Institute, Boston, United States
13   3. Department of Molecular Microbiology and Immunology, Brown University, Providence,
14   United States
15   4. Department of Microbiology, Faculty of Science, University of Manitoba, Winnipeg
16
17
18
19 *Correspondence: shapiror@uoguelph.ca
20

Abstract

The rising rate of drug-resistant fungal infections and the emergence of fungal pathogens with intrinsic resistance phenotypes are a growing concern. The close evolutionary distance between mammals and fungi complicates the design of new antifungals and increases the chances of toxic off-target effects. As such, antifungal drug development usually focuses on fungal-specific proteins when considering potential new targets. Ideal drug targets should mediate essential cell processes and be highly sensitive to inhibition. Targeted gene repression can serve as a model for drug-mediated inhibition and for determining the dosage-sensitivity profile of genes of interest. In the fungal pathogen *Candida albicans*, classical approaches for gene repression can be labour-intensive and limited to one genetic background due to low throughput. Here, we adapt pooled CRISPRi in *C. albicans* for the first time and exploit this technique for large-scale functional genomic analysis. Through pooled CRISPRi screening, we test the repression sensitivity of over a hundred essential genes conserved in fungi but absent in humans in a single flask, and successfully identify highly dosage-sensitive genes across multiple cell components and pathways. We further extended our analysis to ten diverse environmental conditions, allowing us to measure how the environment influences dosage-sensitivity profiles. Finally, we extend our experiments to two clinical drug-resistant *C. albicans* strain backgrounds and demonstrate that many of the fitness defects we observed are conserved in resistant clinical isolates. Together, our results highlight a set of genes that are highly dosage-sensitive across different genetic and environmental contexts, making them attractive targets for further investigation. By facilitating rapid, efficient large-scale functional genomics assays across diverse genetic backgrounds, CRISPRi pooled screening will open new frontiers in *C. albicans* biology.

Introduction

Over recent years, fungal infections have become more prevalent due to environmental changes and an increase in the immunocompromised population, a trend expected to continue in the coming decades (Fisher et al. 2022). One of the most medically relevant human fungal pathogens is *Candida albicans*, which is the fourth most common cause of hospital-acquired bloodstream infections and results in roughly a million deaths each year (Denning 2024). Current treatment options for invasive *Candida* infections are limited to three classes of antifungal drugs, with two of these, azoles and echinocandins, being dominant in therapeutic management. While these therapeutics have been broadly effective, the increased occurrence of antifungal resistance, as well as cases of cross-resistance (Bédard et al. 2025), has underscored the urgent need for new antifungal drugs. The close evolutionary relationship between fungi and humans has significantly complicated this process. For example, promising molecules with antifungal effects, such as rapamycin (a TOR complex inhibitor) and FK506 (a calcineurin inhibitor), were later found to have potent immunosuppressive effects in mammals due to the conservation of both complexes, precluding their use to treat fungal infections (Rivera and Heitman 2023). To avoid crosstalk with human cell components, fungal-specific cell components must therefore be prioritized as targets for future antifungal development.

In the genomes of microbial pathogens, essential genes have commonly been the focus of antimicrobial development efforts as they, by definition, directly affect the cell viability and reproduction (Fields et al. 2017). However, knowing that a given gene is essential provides no information about what fraction of its activity must be inhibited to significantly impede growth (Figure 1a). This relationship between gene dosage and fitness, often referred to as dosage-sensitivity in genetics, in turn mediates the dose-response profile to an inhibitor. Dosage sensitivity can vary immensely between genes (Keren et al. 2016) and has not been characterized for most of the *C. albicans* genome. The fraction of gene activity that must be reduced by mutations or inhibited by drugs to cause a fitness defect can be influenced by expression levels above the basic requirements for optimal fitness (Cisneros et al. 2023) and by downstream buffering by metabolic networks (Kacser and Burns 1981). Conversely, a gene for which only a slight reduction in activity translates to reduced fitness should be more sensitive to drug inhibition. For example, the dosage of haploinsufficient genes has low buffering capacity, so that reducing the effective concentration of the gene product by deleting one out of two alleles results in fitness defects or drug sensitization (Baetz et al. 2004). Importantly, dosage-sensitivity is not captured by *in vitro* enzymatic assays alone, which isolate the protein of interest outside of its cellular context. Finding and prioritizing highly dose-sensitive targets enables the use of lower drug doses, improving therapeutic safety and allowing for less stringent optimization of compounds during drug development. The current lack of information on dosage-sensitivity profiles for fungal pathogens' genes limits our ability to achieve this. Importantly, successful antifungal targets must remain effective across a range of environmental conditions and genetic backgrounds, including drug-resistant isolates, as these contexts can profoundly influence gene-

fitness relationships (Fu et al. 2025). However, for essential genes, filling this knowledge gap is
complicated by the inherent challenges of manipulating them genetically.
The advent of CRISPR technologies in *Candida* species presents new opportunities for a
deeper characterization of essential genes and their function beyond direct editing (Gervais et al.
2023; Maroc et al. 2024; Román et al. 2019). CRISPR interference (CRISPRi), in which a
transcriptional repressor is fused with a catalytically dead Cas9 (dCas9) enzyme, is targeted to a
promoter by a sgRNA and can be used to knock down gene expression of genes of interest in a
programmable manner (Wensing et al. 2019; Wensing and Shapiro 2022). This strategy provides
an alternative to locus-specific promoter replacement, enabling tunable gene repression while
simplifying strain construction and preserving the native genomic context (Roemer et al. 2003;
Noble et al. 2010; Homann et al. 2009). In addition, CRISPRi systems allow the design of unique
sgRNAs targeting the same promoter, each of which induces a different level of repression
depending on the guide and promoter sequence properties. Accordingly, genes that exhibit
fitness defects when repressed by multiple distinct sgRNAs can be interpreted as repression-
sensitive, based on their vulnerability to a broad range of decreases in expression, mimicking
drug inhibition and suggesting dosage-sensitivity (Figure 1b). Another advantage of CRISPR
approaches is their high scalability. High-throughput CRISPRi libraries have been used in a
diverse range of eukaryotic organisms to identify key cellular components through pooled
competition assays (Bock et al. 2022; Shapiro et al. 2018), but this approach has yet to be
extended to fungal pathogens. In these experiments, high-throughput sequencing of the sgRNA
locus can be used to infer the fitness effects of thousands of perturbations in a single test tube. In
C. albicans, such approaches could enable rapid characterization of genes of interest and are
well-suited to the study of essential cell components.
Here, we develop and validate a first-of-its-kind regulatable system for pooled CRISPRi
screening in *C. albicans*. Using this platform, we interrogate the repression sensitivity of 130
putative essential genes that are conserved in fungi but absent in humans, both in standard
laboratory conditions and across ten diverse environmental contexts. We further test the impact
of genetic background on repression-sensitivity profiles by repeating our screen with two clinical
antifungal-resistant isolates. We find that our pooled CRISPRi approach can robustly detect
genes that are sensitive to dosage perturbations. We identify a group of 16 genes that showed
significant repression sensitivity under all experimental and genetic contexts, nominating them as
high-priority candidates for future antifungal drug development. More broadly, our approach
provides a rapid, scalable, and portable method for creating large knockdown mutant libraries
that can be readily applied to elucidate gene function under different environmental conditions
and across different genetic backgrounds in *C. albicans*.

Results

Design and construction of the targeted essential gene library

To enable scalable, pooled repression of essential genes, we built on our previously established CRISPR interference (CRISPRi) framework in *C. albicans*, in which a catalytically dead Cas9 (dCas9) fused to the transcriptional repressor Mxi1 is directed to gene promoters by a single guide RNA (sgRNA). In this system, all CRISPRi components are encoded on a single integrating plasmid, simplifying strain construction and enabling pooled library generation (Wensing et al. 2019). In our previously published *C. albicans* CRISPRi system, dCas9–Mxi1 was expressed constitutively following integration at a safe-harbour locus. However, controlling the timing of gene repression is critical when interrogating essential genes, particularly to avoid premature depletion of strains before the start of competitive growth assays (Figure 1c). Thus, to create a regulatable gene repression system, we redesigned our vector to include a tetracycline (tet)-regulated transcription factor and an associated promoter (*tetO*) controlling dCas9 expression (Roemer et al. 2003). This *tetO* promoter is a *C. albicans*-optimized tet-off system in which the addition of tetracycline or an analog, such as anhydrotetracycline (ATc), represses dCas9 expression. In the absence of ATc, dCas9 is expressed, and the gene targeted by the sgRNA is repressed. We validated this new system using one non-essential gene, *KSP1* and one essential gene, *HSP90*. The regulatable expression system achieved comparable repression efficiency compared to constitutive expression (Figure 1d). In addition, the promoter exhibited low expression leakage, with minimal repression of *HSP90* when turned off and a strong effect when turned on (Figure 1e). Knockdown of *HSP90* only led to growth effects when ATc was absent over multiple generations, with minimal expression costs (Figure 1f). The regulatable system thus provides control over the timing of repression-induced effects, allowing for better decoupling of strain construction and fitness measurements. This modified CRISPRi system, which is well-suited to study essential genes, now enables us to scale up and develop an efficient strategy to generate a pooled *C. albicans* mutant library, supporting large-scale, targeted forward genetic screening of essential genes (Figure 1g).

145
146
Figure 1: Regulatable CRISPRi for pooled screening in *Candida albicans*. a) The dosage sensitivity
profile of a gene impacts druggability. For dosage-insensitive genes, even a large reduction in the
effective concentration of gene products can result in only minor fitness effects, while only a slight
reduction in the abundance of a dosage-sensitive product leads to large fitness effects. **b)** Gene
knockdown with CRISPRi reduces expression levels, with each different sgRNA potentially leading to
different levels of depletion and mimicking drug repression. As such, sgRNAs targeting the same gene at
different degrees of repression can identify repression-sensitive genes. **c)** The fitness of CRISPRi strains
with sgRNAs targeting essential genes of interest (GOI) is immediately impacted if dCas9-Mxi1
expression is constitutive, leading to difficulties and premature dropouts when constructing strains.
Placing *dCas9-Mxi1* under the control of a *tetO* promoter/repressor system prevents immediate fitness
effects, which occur only when switching cells to media without ATc, thereby lifting repression of the
CRISPRi system. **d)** CRISPRi repression of *KSP1* in strains transformed with either the constitutive
(cons.) or the regulatable system in its active state (ON) with the same sgRNA. Stars represent significant
differences between the *KSP1* targeting strains and the control strains (Cons.: p=0.0001, ON: p=0.0002,
Welch's t-test). **e)** CRISPRi repression of *HSP90* by the regulatable system when it is inactive (OFF) or

active (ON). Stars represent significant differences with the OFF control strain (*HSP90* OFF: p=0.03, *HSP90* ON: p=0.0002, Welch's t-test) **f)** Growth effect of *HSP90* repression using the inducible system after different ATc washout periods. Stars represent significant differences with the WT-12h control strain (CTRL OFF: p=0.019, *HSP90* ON-30h: p=0.00026, Welch's t-test). **g)** Workflow for pooled CRISPRi screening of fungal-specific essential genes in *C. albicans*. For each gene of interest, four sgRNAs were designed and cloned into the regulatable CRISPRi integration vector from an oligonucleotide pool. This plasmid library was then batch transformed into fungal cells under ATc repression. The resulting library can then be used for pooled screening, in which changes in relative sgRNA abundance measured by high-throughput sequencing are used to infer fitness effects and repression sensitivity.

We next sought to use our regulatable CRISPRi system to identify repression-sensitive essential genes in *C. albicans* at high-throughput. We focused on a list of 130 putatively essential genes identified in a transposon mutagenesis screen (TN-Seq) of a haploid *C. albicans* model (Table S1). In addition to being depleted for transposon insertions, these genes were conserved across three evolutionarily distant fungal pathogens (*Aspergillus fumigatus*, *Histoplasma gondii*, and *Cryptococcus neoformans*) and lacked close homologs in humans (Segal et al. 2018). We used the sgRNA design principles outlined in a previous CRISPRi screen in *Saccharomyces cerevisiae* (Momen-Roknabadi et al. 2020) to select sgRNA target sequences. Briefly, we selected four guides targeting the promoter of each gene, prioritizing sgRNAs with higher predicted efficiency scores (Peng and Tarleton 2015). We also included 30 randomized sgRNAs as negative controls (control sgRNAs). The library was synthesized as a pool of single-stranded oligonucleotides, which was then amplified and cloned *en masse* into the CRISPRi plasmid. We transformed the plasmid pool into *C. albicans* laboratory strain SC5314. For both bacterial and fungal library generation steps, we validated that library sequencing recovered the vast majority of designed sgRNAs, with ~85% of the designed sgRNAs being present in the fungal pool (Figure S1). Our optimized pipeline for pooled strain construction enables the generation of any screening-ready *C. albicans* CRISPRi library in as little as seven days (Supplementary Note 1). We have made the regulatable CRISPRi vector publicly available on Addgene (pRS479, #250493), thus providing a new, cost-effective and rapid option for high-throughput functional genomics in *C. albicans* research.

High-throughput screening of essential genes in standard laboratory conditions

Having built our CRISPRi strain library, we next established a pipeline to screen our mutant library for growth defects under standard laboratory conditions. We grew the library in rich media (YPD) with or without ATc over three 12-hour passages, spanning ~21 generations, and tracked changes in the relative abundance of mutant strains compared to the starting inoculum population using high-throughput sequencing of the sgRNA locus. Log$_2$ fold-change (Log$_2$FC) values were well correlated between replicates (ρ=0.82-0.84), showing that the pooled competition assay generated reproducible outputs (Figure S2). When CRISPRi repression was turned on (CRISPRi-ON), the relative abundance of some sgRNAs targeting essential gene

promoters decreased sharply compared to the randomized controls, while abundances remained
mostly unchanged when the system was kept inactive (CRISPRi-OFF) (Figure 2a, Figure S3).
Using the Log_2FC values of the random sgRNAs as a control distribution, we found 96 sgRNAs
were significantly depleted in abundance at the final time point, which we refer to as hits (Figure
2b, Figure S4). To validate that negative Log_2FCs were indeed associated with reduced fitness,
we constructed individual strains with sgRNAs targeting *MRLP33*, *MMM1,* or *ASK1*, as
candidate genes from this screen with three or more hit guides. Both *MRPL33* and *MMM1* are
linked with mitochondrial function, while *ASK1* is a subunit of the DASH complex involved in
chromosome segregation. In most cases (9/10), screen hit sgRNAs induced a measurable growth
defect in smaller-scale growth curve assays (Figure 2c), validating the phenotypes identified in
the large-scale assay. This demonstrates the ability of our novel, pooled CRISPRi screening
assay for rapid and scalable analysis of *C. albicans* strains libraries.
214

215
Figure 2: Pooled CRISPRi screening of essential genes in *C. albicans* identifies repression sensitivity
phenotypes. a) Changes in relative abundance of sgRNAs in the library throughout the timecourse assay,
with (ON) or without (OFF) CRISPRi repression. Randomized sgRNAs (n=30) are shown in green, and
essential gene targeting sgRNAs are shown in black. The red dotted line shows a hypothetical 1:1
relationship. **b)** Abundance change distribution at the final time point for the CRISPRi-ON condition. The
red line indicates the significance threshold corresponding to FDR=5%. **c)** Abundance changes of

sgRNAs targeting *MRPL33*, *MMM1* and *ASK1* in the timecourse assay, with the corresponding AUC
measurement obtained in growth curve assays. The grey bars show the mean of two biological replicates,
which are shown overlaid with error bars corresponding to the 95% confidence interval of 5 technical
replicates. Stars indicate strains with a significant change in growth ($p<0.05$, Welch's t-test). **d)** Number
of hit sgRNAs per gene in the timecourse assay, for genes with at least one hit. Empirical p-values were
obtained by sampling 96 hits over the 438 sgRNAs measured over 100 000 permutations to estimate the
expected frequency of genes with 3 or 4 hits. **e)** Frequency of mitochondrial localization annotation
(GO:0005739) across the *C. albicans* genome (n=6473) and within the fungal-specific gene set (n=131).
The difference in annotation frequency in both categories was tested using Fisher's exact test.

Amongst mutant strains with growth defects, we found additional instances in which
three or four of their four designed sgRNAs were significantly depleted, yielding a total of
twelve enriched genes. A higher fraction of sgRNAs that result in fitness defects is unlikely to
occur by chance, suggesting greater repression sensitivity for these targets (Figure 2d). These
twelve genes include five that are predicted by *S. cerevisiae* orthology to encode part of the
mitochondrial ribosome: *MRPL33*, *C1_12610W* (S.c.: *MRPL15*), *C2_07680W* (S.c.: *MRPL7*),
CR_01370C (S.c.: *MRPS28*), and *CR_04140W* (S.c.: *NAM9*), of which four remain
uncharacterized in *C. albicans*. These five proteins do not cluster spatially within the S.
cerevisiae mitoribosomal structure (Figure S5) but might represent different, potentially rate-
limiting components of complex assembly. In *S. cerevisiae*, *MRPL33*, *MRPL15*, and *MRPL7* are
essential components of the large subunit of the mitoribosome that join the complex in the early
to intermediate stages of assembly (Zeng et al. 2018). Two other highly repression-sensitive
genes, *MMM1* and *C1_00510W* (S.c.: *ECM31*), are also predicted to be localized to the
mitochondria (Perocchi et al. 2006). This reflects a high enrichment of the initial gene set for
mitochondrial genes and mitoribosomal components (Figure 2e). Collectively, these observations
suggest that mitochondrial pathways, particularly components of the mitoribosome, are among
the most repression-sensitive components in our screen and may therefore constitute especially
promising targets for antifungal inhibition.

Screening in multiple environments identifies a subset of core repression sensitive genes

Our initial pooled-library experiment enabled us to identify repression sensitive essential
genes under standard laboratory conditions. However, environmental factors can affect the
fitness contribution of genes, potentially leading to condition-dependent essentiality (Bosch-
Guiteras and van Leeuwen 2022; Fu et al. 2025). Ideally, candidate antifungal targets should
show strong repression sensitivity across all environments. To test this, we screened our
CRISPRi library across 11 alternative environments, spanning different nutrient sources and
stresses, and monitored changes in the relative abundance of mutant strains via sequencing
(Figure 3a). We found that changes in mutant strain abundance were highly correlated within
conditions (median ρ=0.96-0.80; Figure S6), supporting the high reproducibility of the approach.
Different signal-to-noise ratios across screening conditions resulted in variation in the number of

hits identified in each screen (Figure 3b), prompting us to exclude the tunicamycin condition from downstream analysis. Mapping CRISPRi-induced defects across conditions, we found that most phenotypes were broadly identical across conditions (e.g.: *MMM1*, *ASK1*, *C2_07680W*, *C2_03560C*: Figure S7), with a few cases of condition-dependent fitness defects (Figure 3c). Overall, these results demonstrate the applicability of our pooled CRISPRi approach for comparative environmental screening, which is crucial to resolving how essential genes contribute to fitness across diverse contexts.

Figure 3: Parallel screening identifies condition-independent repression-sensitive genes. a) The same CRISPRi library used in the timecourse assay was screened in 11 conditions in parallel. **b)** Number of significantly depleted sgRNAs in each screening condition. **c)** Number of depleted sgRNAs per gene in each condition, excluding tunicamycin. Only genes which had hits in at least two conditions are shown. **d)** Number of conditions with significant fitness decreases for hit genes in the experiment. **e)** Abundance

changes of the 4 sgRNAs targeting *FBA1* in media conditions with (YPD) or without (LAC: YP + lactate
2%, GLY: YP + glycerol 2%, ACE: YP + acetate 2%) a fermentable carbon source. The overall
distribution of Log$_2$FC for library sgRNAs is shown as a violin plot, and the randomized sgRNAs are
shown in green. The black lines show the significance threshold for each condition. **f)** Growth curve assay
of strains with *FBA1* targeting sgRNAs in YPD or GLY media. Each measurement was performed in
biological duplicates with three technical replicates. Error bars represent the 95% confidence interval
around each biological replicate.

From these screens, we identified 23 genes with repression-sensitivity phenotypes that
were universal across the 10 environments we analyzed (Figure 3d), representing the most
common scenario of effects across conditions. The consistent phenotypes observed for a subset
of genes across environments suggest they might be of particular interest for antifungal
development. Conversely, some hit genes from the first experiment were found to be repression
sensitive only in certain environments. One of the most striking cases was fructose-bisphosphate
aldolase FBA1, a key enzyme in both glycolysis and gluconeogenesis. In conditions where
glucose was the carbon source, *FBA1* showed strong repression sensitivity, which was abolished
when switching to non-fermentable nutrients (Figure 3e). We validated this phenotype in small-
scale assays, in which growth of FBA1 CRISPRi mutants was highly impaired relative to the
wild type in glucose but not in glycerol media (Figure 3f). Repression efficiency was comparable
in both conditions (Figure S8), Because the fungal fructose-bisphosphate aldolase is not
homologous to the human isozyme, there are currently efforts to develop specific inhibitors that
could be used as antifungals (Wen et al. 2022). The ability of *C. albicans* to efficiently perform
glycolysis is thought to be important for virulence traits like hypoxia adaptation and biofilm
formation (Bonhomme et al. 2011; Childers et al. 2016). On the other hand, the dispensability of
FBA1 on non-fermentable carbon sources suggests that the impact of reduced gluconogenesis
flux on fitness is much lower. Thus, the ability to screen CRISPRi strain libraries quickly and
efficiently in a large number of experimental conditions can facilitate the detection of
environmental effects such as these.

Many repression sensitivity phenotypes are conserved in antifungal-resistant isolates

In addition to environmental changes, genetic background differences can affect the
function and dosage sensitivity of genes (Bosch-Guiteras and van Leeuwen 2022; Larrimore and
Rancati 2019). This is especially important in the case of antifungal resistance mutations, which
can have both positive and negative genetic interactions with other cell components (Usher and
Haynes 2019; Halder et al. 2020; Gervais et al. 2025). One advantage of pooled library
construction and screening is that plasmid libraries can readily be transferred to different genetic
backgrounds, enabling parallelized screening and comparative functional genomic analysis
(Figure 4a). To validate this approach, we compared the effect of *HSP90* CRISPRi repression in
the laboratory strain SC5314 and two clinical isolates, resistant to either fluconazole (FLU[R])

(Perea et al. 2001) or caspofungin (CASP[R]) (Yadav et al. 2024). Both isolates have mutations
representative of their respective resistance class: an *ERG11* T299A F449S allele for FLU[R], and
a F641S *FKS1* allele for CASP[R] (Bédard et al. 2025). CRISPRi repression was efficient in both
strains (Figure 4b), although its efficiency varied despite no significant changes in baseline
HSP90 transcript levels compared to the wild-type (Figure S9), which might be explained by
differences in Cas9 expression (Chavez et al. 2016). Next, we used the same batch
transformation protocol to transform our CRISPRi plasmid library into both drug-resistant strain
backgrounds. After sequencing, we found that one transformation batch per background was
sufficient to obtain >90% representation of the designed sgRNA, which is higher than what we
obtained for the WT strain library (Figure 4c). Overall, this resulted in >80% of sgRNAs being
present in all three strain pools and thus directly comparable in downstream assays. Our pooled
CRISPRi approach is therefore readily portable, enabling rapid and efficient expansion of large-
scale strain libraries across multiple genetic backgrounds, a capability not previously possible in
C. albicans.
We next used these new libraries to compare the fitness effects of CRISPRi repression in
the three different strain backgrounds. We performed competition assays in rich media for the
three strains in parallel, and measured relative sgRNA abundance via sequencing. Across all
backgrounds, sgRNA abundance changes were highly correlated, indicating that our pooled
screening approach is highly reproducible in clinical isolates (FLU[R]: ρ=0.69-76, CASP[R]: ρ=85-
88, Figure S10 and S11). We used the distribution of random sgRNA abundance changes to set a
Log$_2$FC threshold for each strain (Figure 4d), and identified a similar number of mutant strains
with altered abundance in both the wild-type and CASP[R] strain backgrounds (WT: 107 vs.
CASP[R]: 119). We validated our screen's ability to detect fitness effects in the new strain
backgrounds by reconstructing several CRISPRi strains and measuring growth curves (Figure
S10). As we had observed for the wild-type, the growth curves generally recapitulated the pooled
screening results. Among unique genes with at least one CRISPRi fitness defect, we found that
44% (n=34) were represented in all genetic backgrounds (Figure 4e). The second-largest overlap
was between the wild-type and the CASP[R] strain, reflecting the greater sensitivity of the assay in
these two backgrounds. We further sequenced the genomes of all three strains to examine
whether polymorphisms in promoter regions could have altered sgRNA binding, which could
also explain strain-specific fitness effects via reduced or abolished CRISPRi activity. We found
that ~3% of sgRNAs bound promoter regions containing variants unique to clinical isolates
(Figure S12), a fraction much too low to explain the differences we observed between strains.
These results demonstrate that CRISPRi pooled screening functions reliably across multiple
distinct genetic backgrounds, thus extending the ability to perform these high-throughput
functional genomic screens to diverse isolates.

350
Figure 4: Portable CRISPRi screening of *C. albicans* **essential genes identifies a core set of**
repression-sensitive targets. a) The same CRISPRi plasmid library used to build the wild-type strain
pool was transformed into two new genetic backgrounds, resistant to either fluconazole (FLU^R) or
caspofungin (CASP^R). **b)** CRISPRi repression of *HSP90* using the same sgRNA in the three different
strains. **c)** Library coverage in each genetic background at the start of the screen. The sgRNAs present
across all three libraries are shown in black. **d)** Abundance change distribution for each genetic
background. The red line indicates the Log$_2$FC threshold corresponding to FDR=5%. **e)** Overlap between
repression-sensitive genes identified for each strain. **f)** Sensitive genes identified in all environmental
conditions and genetic backgrounds. **g)** Interaction between CRISPRi repression and Caspofungin
treatment. Only genes with at least one sgRNA showing antagonism or synergy of sufficient magnitude
and statistical significance (Benjamini-Hochberg corrected p-value <0.05) are shown as triangles and
colored. The direction of the triangle indicates the nature of the interaction, as indicated by the compass
(left: synergy in WT, right: antagonism in WT, down: synergy in CASP^R, up: antagonism in CASP^R). The
black lines show the thresholds for z-score change magnitude (see methods). Genes and sgRNAs not
showing synergy or antagonism are shown in grey.

Across these pooled CRISPRi screens, we investigated which genes from our set of
fungal specific essential genes have repression sensitivity phenotypes, and whether these
phenotypes are maintained in different environmental and genetic contexts. Out of the 23 genes

that were sensitive to growth in all environments in the wild-type, 16 were also hits in all three genetic backgrounds, allowing us to refine a shortlist of potential new targets to focus on for antifungal development (Figure 4f). As potential new therapeutics might have to be used in combination with existing compounds, understanding whether exposure to antifungals impact repression sensitivity profiles is important. In this scenario, increased CRISPRi fitness defects under treatment could indicate synergy with the drug, whereas reduced fitness defects could indicate antagonism. Ideally, the repression of potential therapeutic targets should be neutral, if not synergistic, to avoid reducing treatment efficiency. The portability and scalability of *C. albicans* CRISPRi pooled screening allow us to examine these relationships while also accounting for genetic background. Thus, we screened the wild-type and CASP[R] libraries with or without the antifungal caspofungin present at a concentration below the MIC, and compared the effects on sgRNA depletion in both cases. We found that the vast majority of mutant strains showed no changes, with only a few sgRNAs exhibiting large, significant changes in fitness in one but not both strain backgrounds (Figure 4g). For example, one sgRNA targeting *QCR8*, a structural component of respiratory complex III (shown in yellow, lower left quadrant), was more depleted in the wild-type, but did not change significantly in the CASP[R] background. Only the unsaturated fatty acid synthesis regulator *SPT23* showed a relatively consistent pattern of lower repression sensitivity under caspofungin treatment, with 1 sgRNA in the wild-type and 2 sgRNAs in the resistant strain having reduced effects. We observed a similar pattern with the FLU[R] strain, with no significant shifts compared to the wild-type (Figure S13). These results suggest that repressing the core repression-sensitive genes we identified should not interfere with at least one major antifungal, which could allow for combinatorial therapy options.

Discussion

Systematic genetic interrogation of essential genes has transformed our understanding of the core biological processes that regulate microbial life, yet such approaches have remained difficult to implement in fungal pathogens. Here, we establish a regulatable pooled CRISPRi platform for the fungal pathogen *C. albicans* that overcomes the major barriers to high-throughput perturbation of essential genes. By combining regulatable gene repression and rapid construction of high-coverage pooled mutant libraries, our system expands functional genomic capability in *C. albicans* and provides the first application of pooled CRISPRi screening in a fungal pathogen. This new toolkit is available to the community (Addgene #250493) and can be easily used to generate user-specific libraries for diverse functional genomic screening applications in *C. albicans*. Using this platform, we identified a substantial fraction of fungal-specific essential genes that exhibit strong repression sensitivity, assessed core vs. conditionally-essential genes, and demonstrated the portability of this system for comparative functional genomic analysis across diverse *C. albicans* strain backgrounds. Together, our findings underscore the importance of assessing gene essentiality in *C. albicans* as a complex trait that is influenced by gene repression sensitivity, the cellular environment, as well as the genetic context, all of which can impact the evaluation of potential antifungal targets.

Understanding the genetic architecture of essential cellular processes in fungal pathogens
remains a longstanding challenge. While foundational studies using deletion libraries and
repression libraries (Roemer et al. 2003; Fu et al. 2025, 2021), and transposon mutagenesis
(Segal et al. 2018) have provided critical early maps of gene essentiality, these approaches rarely
capture quantitative or qualitative relationships between gene dosage and fitness. Yet, an
accumulating body of work demonstrates that dosage sensitivity is a powerful predictor of
druggability (Baetz et al. 2004; Keren et al. 2016), and evolutionary constraint in the context of
resistance (Cisneros et al. 2023). CRISPRi approaches in the model yeast *S. cerevisiae* (Momen-
Roknabadi et al. 2020; Arita et al. 2021; Jaffe et al. 2019) has highlighted the potential of
expression-level perturbations to study critical phenotypes at a high-throughput scale, yet no
analogous high-throughput tool existed for major fungal pathogens. By developing a regulatable
CRISPRi system compatible with pooled screening, we extend these principles to *C. albicans*,
enabling quantitative interrogation of essential gene biology in a critical human fungal pathogen.
An increasing body of research has highlighted the importance of recognizing *C.*
albicans, and other fungal pathogen species more broadly, as genetically diverse species in
which diverse isolates differ markedly in genome structure, regulatory circuitry, stress tolerance,
and drug susceptibility (Hirakawa et al. 2015; Ropars et al. 2018; Gerstein and Berman 2020;
Iracane et al. 2024). These differences can fundamentally alter gene essentiality and genetic-
interaction networks, yet most functional genomics resources in *C. albicans* remain restricted to
a single laboratory background. Our findings demonstrate the utility of the CRISPRi system
developed here as a highly portable functional genomic tool, enabling scalable, and comparative
functional studies across the natural diversity of *C. albicans* strains. By introducing pooled
CRISPRi libraries into diverse *C. albicans* strains, including clinical isolates and drug-resistant
strains, we provide a route to systematically map background-specific repression sensitivity and
identify genetic interactions that are uniquely found in particular genomic contexts. In particular,
such analysis can allow us to identify cell components that can alter drug sensitivity uniquely in
drug-resistant genetic backgrounds. Previous studies have found that repressing factors such as
PKC1, *GIN4*, or *CCT8* sensitizes a drug-resistant *fks1** mutant to echinocandins (Caplan et al.
2018), illustrating the power of this type of approach at a small scale. Our system now extends
this capacity to a large scale, allowing scalable comparative analysis across drug-resistant fungal
strains. Integrating genetically diverse isolates into CRISPRi screens will therefore be essential
for uncovering strain-specific vulnerabilities, and ultimately designing antifungal strategies
robust across clinically relevant genetic backgrounds.
One advantage of our optimized *C. albicans* CRISPRi pooled screening system is its
ability to identify gene repression sensitivity at scale, thereby enabling the systematic
identification of essential genes for which even modest reductions in expression lead to
substantial growth defects. These highly repression-sensitive genes represent particularly
compelling antifungal targets, as therapeutic inhibition may require only partial suppression of
gene function, thus broadening the range of compounds that can be effective therapeutics at low
doses, and reducing the risk of off-target toxicity. By identifying fungal-specific, repression-

sensitive genes that are essential for fungal viability across diverse environmental and genetic
contexts, our study generates a prioritized list of targets for antifungal drug discovery. This
resource coincides with emerging AI-driven drug discovery pipelines, including protein
structure-prediction tools, generative molecular design, and machine-learning programs that
identify ligandable pockets and predict inhibitor binding and affinity (Ocana et al. 2025; Pun et
al. 2023; Liao et al. 2025). As protein structures for many of the identified targets can now be
rapidly modelled, AI-enhanced screening can accelerate the development of small molecules
tailored explicitly to these fungal vulnerabilities. Thus, our CRISPRi-based platform provides a
bridge between computational drug design efforts and the antifungal drug development pipeline.
Together, we present a fast, scalable, and portable CRISPRi system that enables high-
throughput functional genomics in *C. albicans*, including multi-environment screening,
comparative analysis across diverse genetic backgrounds, and interrogation of repression-
sensitive essential genes. This platform provides the first pooled CRISPRi screening framework
for a major fungal pathogen and reveals a core set of fungal-specific, repression-sensitive
essential genes with broad relevance for antifungal target discovery. Given its flexibility and
efficiency, this system may now be adapted to genome-scale libraries, additional fungal species,
and a wide range of biological questions, thus providing an important new strategy for dissecting
fungal biology.

Materials and methods

468 Strains, plasmids and oligonucleotides used in this study are shown in Table S2, S3 and S4, respectively.
469 Media recipes are available in Table S5.

Regulatable CRISPRi expression plasmid construction

To create the regulatable CRISPRi plasmid, the tetO promoter sequence and tetracycline
transactivator (TAR) were synthesized de novo by Synbio Technologies LLC and inserted into
the original CRISPRi plasmid in place of the original ACT1 promoter driving expression of
dCas9. Briefly, the backbone of the plasmid was taken from the original CRISPRi plasmid
(pRS159, Addgene # 122378) previously created by the Shapiro lab (Wensing *et al,* 2019). The
tetO promoter and TAR were synthesized using a *C. albicans* codon-optimized plasmid provided
by the Cowen lab (pRS250, Cowen lab reference pLC1031) as the sequence reference. The
ACT1 promoter that previously expressed dCas9 in pRS159 was removed and the TAR and
promoter amplicon was cloned in its place. The plasmid was transformed into *E. coli* DH5α cells
and grown at 30 °C in LB media containing both ampicillin (AMP) and nourseothricin (NAT).
This plasmid has been deposited on Addgene (on hold, #250493).

sgRNA library design.

The subset of essential genes we selected for this library was based on data from a recent
TN-Seq assay in a haploid *C. albicans* strain (Segal et al. 2018). Briefly, after filtering based on
putative essentiality (as measured by haploid TN-Seq data), absence of human homologs and
conservation in three other distant fungal pathogens (*Aspergillus fumigatus*, *Histoplasma gondii*,
and *Cryptococcus neoformans*), the authors identified a set of 130 fungal specific conserved
essential genes. SgRNA design was performed broadly as described, with some minor
modifications to adjust for the pooled library format (Wensing and Shapiro 2022). For each of
these genes, we searched for potential target sites in their promoter region (up to 300 bp
upstream from the start codon) with the EuPaGDT online tool and used the predicted score to
select the best four sgRNAs. Promoter sequences were retrieved from the *Candida* Genome
Database (CGD, (Lew-Smith et al. 2025)). Target sequences that contained Sap1 or Pac1
recognition sites were filtered out, as these enzymes are used in subsequent cloning steps. As a
negative control, we also generated four sgRNAs targeting the well-characterized molecular
chaperone, Hsp90, as a positive control and 30 randomly scrambled guide sequences with no
perfect matches in the *C. albicans* genome as a negative control. This resulted in a total of 554
guides in the library. The oligonucleotide sequences were designed as follows: 5'
GGTCTAATTT**GCTCTTCC**cga-N20-gtt**GGAAGAGC**TTTTAATTAAGCTT 3' (bold: SapI
recognition sequences, lowercase: ligation overhang for directional cloning, N20: 20 bp target
sequence). The resulting oligonucleotide library was then ordered from IDT as pooled single-
stranded DNA (ssDNA).

Primer extension

Pooled ssDNA sequences were converted to double-stranded DNA by PCR. Two PCR reactions were set up each with a 50 μL total volume. Each reaction consisted of 1 μL (1 μg/μL) of ssDNA (oligo pool), 1 μL of primer (pRS 596) 5' AAGCTTAATTAAAAGCTCTTC 3' (equimolar amount of primer, 318.18 ng/μL), 41 μL of Nuclease-free water and 5 μL of 10x NEBuffer 2.1 (NEB, cat. B7202S). The PCR conditions were set to: 94 $^{\circ}$C for 5 mins, slow ramp down (0.1 $^{\circ}$C/sec) to 46.4 $^{\circ}$C and hold for 5 mins, slow ramp down to 37 $^{\circ}$C and hold for 1 min. Then, 1 μL of 5,000 U/mL DNA polymerase I, Large (Klenow) fragment (NEB, cat. M0210S) and 1 μl of 10 mM dNTPs (NEB, cat. N0447L) was added to the tube. The thermocycler conditions were then set to 37 $^{\circ}$C for 1 hr, followed by a heat-inactivation step at 72 $^{\circ}$C for 20 mins, a slow ramp down to 42 $^{\circ}$C for 1 min, and then held at 4 $^{\circ}$C until ready to use. The 50 μL reaction was then processed using the DNA Clean and Concentrator-5 kit (Zymo research, cat. D4013) as per manufacturer's instructions. The cleaned product was quantified using a Tecan Infinite Pro with a NanoQuant plate.

Golden Gate cloning

The double-stranded oligo pool (dsN20) sequences were cloned into the inducible CRISPRi plasmid using a Golden Gate cloning strategy previously described (Wensing et al. 2019; Wensing and Shapiro 2022). Briefly, the following reaction was prepared in quadruplicate: 10 μl of miniprepped CRISPRi plasmid (180 ng/μl), 1 μl of dsN20 (diluted to a concentration that results in a 3:1 insert to vector ratio), 2 μl 10mM ATP (NEB, cat. P0756S), 2 μl 10x rCutsmart buffer (NEB, cat. B6004S), 1 μl 400,000 U/mL T4 DNA ligase (NEB, M0202S), 1 μl of 10,000 U/mL Sap1 (NEB, cat. R0569L) and 3 μl nuclease-free water. The thermocycler was then set for the following conditions: 37 $^{\circ}$C for 2 mins, 16 $^{\circ}$C for 5 mins, cycled 99x, 65 $^{\circ}$C for 15 mins, 80 $^{\circ}$C for 15 mins.

Bacterial transformation

Before transforming, the cloned plasmids were first treated with 1 μL of SapI and incubated for 1 hr at 37 $^{\circ}$C to linearize any remaining uncloned plasmids. Next, we transformed the cloned library into *E. coli* DH5α competent cells using a standard chemical transformation: 5 μl of cloned plasmid was added to 50 μL of commercial chemically competent cells (NEB, cat. C2987H) and incubated on ice for 30 mins, heat shocked at 42 $^{\circ}$C for 30 secs, and then incubated on ice for an additional 5 mins. After this, 950 μL of SOC media (NEB, cat. B9020S) was added. The cells were then incubated horizontally shaking for 1 hr and 30 mins at 30 $^{\circ}$C. From the tubes, 50 μl of culture and 150 μl of water were plated on LB AMP (100 μg/mL, BioShop, cat. AMP201.25) and NAT (50 μg/mL, Jena Bioscience, cat. AB-102L) plates to collect information on colony counts. To prepare the LB-semi-solid media, 0.35% seaprep agar (Lonza, cat. 50302) and LB media mix (BioShop, cat. LBL407.500) was autoclaved with the stir bar left in. The media was cooled to room temperature at which point AMP at a concentration of 100 μg/mL and

NAT at a concentration of 50 µg/mL was added to it. As per the methods from Momen-
Roknabadi et al 2020, 1 L of semi-solid agar can hold between ~1-5 million colonies and
therefore media should be prepared accordingly for the amount of transformation and the
efficiency of the transformations (Momen-Roknabadi et al. 2020). The remaining transformation
culture was added to the semi-solid agar and stirred for 10 mins. The flask was then put in an ice
water bath where the water level was above the media line in the flask for 1 hr to solidify. The
flask was then carefully moved to a 30 ^{o}C static incubator and left to grow for 24-36 hrs. From
the plates, 30 individual colonies were sequenced using Sanger sequencing as a quality control
check: all clones had a properly cloned sgRNA and did not contain any empty backbone
plasmids. Cells were then harvested from the agar by heating the media at 37 ^{o}C for 20 mins
while stirring with the magnetic stir bar to break apart the agar and then centrifuged to collect a
pellet which was then resuspended in 5-10 mL of LB-AMP-NAT. The pool was then frozen in 2
mL aliquots by combining 1 mL of culture (at an OD of ~30) with 1 mL of 40% glycerol.

Plasmid pool extraction

A maxi prep plasmid extraction (Nucleobond Xtra Maxi EF, Macherey-Nagel, cat.
740424.10) was performed to obtain a large quantity of plasmid DNA to be used in the fungal
transformation protocol. To prepare the culture for extraction, 600 mL of LB, AMP (100 µg/mL)
and NAT (50 µg/mL) was added to a 2 L flask. We then added 1 aliquot of the frozen bacteria
pool to the media, and cells were grown at 30 ^{o}C for 12-18 hours until an OD of > 2 was reached.
Extraction was then performed as per manufacturer's instructions. The extracted plasmids were
quantified using a Tecan Infinite Pro with a NanoQuant plate.

Candida albicans transformation

Plasmids were transformed into *C. albicans* using the LiAc-PEG chemical transformation
protocol, as previously described (Halder et al. 2019; Wensing and Shapiro 2022). Briefly, 30 µL
of plasmid DNA (~225 ng/µL) was combined with 1 µl of 10,000 U/mL PacI (NEB, cat.
R0547L), 5 µL of 10x rCutsmart and 14 µL of Nuclease-free water and was digested overnight at
37 ^{o}C static. A 50 mL overnight culture of the *C. albicans* strain of interest was grown at 30 ^{o}C
with shaking (250 rpm) for roughly 16-18 hours. From the overnight culture, calculate the
volume of culture needed to obtain roughly ~5x10^7 cells. Pellet cells using centrifugation and
remove the supernatant. Prepare a transformation master mix using the following recipe: 800 µl
of 50% polyethylene glycol (PEG), 100 µl of 1 M lithium acetate (pH 7.4) 100 µl of 10× Tris-
EDTA (TE) buffer, 40 µl of 10 mg/mL ssDNA (salmon sperm, ThermoFisher cat. 15632-011),
and 20 µL of 1 M dithiothreitol (DTT). The 1060 µl transformation mix and 45 µl (~6000 ng of
linearized plasmid) were then added to a cell pellet, and the cells were flicked to resuspend them.
The cells were incubated static at 30 °C for 1 hr and then heat-shocked at 42 °C for 45 mins,
before being washed twice with YPD and resuspended in 5 mL of YPD with ATc (250 ng/mL)
media for 4 hr at 30 °C with shaking (250 rpm). After outgrowth, these cells were then pelleted
and resuspended in 300 µL of YPD added to 1 L of YPD-semi-solid agar (0.35% seaprep agar)

containing 250 ng/mL of ATc (Alfa Aesar, cat. J66688) and 250□µg/ml NAT. To get a rough

estimation of the transformation efficiency 50 µL from the transformation culture was plated on

YPD-NAT-ATc plates and grown at 30 ^{o}C static for 48 hours. The 1 L of semi-solid agar was

divided between three fernbach flasks as we noticed that *C. albicans* cells grew poorly past a

~1.5 cm depth in the agar. Flasks were grown at 30 ^{o}C for 48-72 hours (to a point at which

colonies became visible). The cells were then recovered using the same method as described for

the bacterial transformation, pelleted, and frozen at -80^{o}C in 2 mL aliquots containing 1 mL of

pooled library and 1 mL of 40% glycerol.

Timecourse competitive growth assay

A timecourse assay was used to monitor the change in mutant abundance over time. For this test

an aliquot of the wildtype fungal pool was thawed on ice for 15 mins and then added to a 250 mL

flask containing 50 mL of YPD and ATc (final concentration 250 ng/mL). The flask was

incubated shaking (200 rpm) at 30 ^{o}C for 12 hrs. The flask was then removed from the incubator

and a subculture was prepared to an OD_{600} of 0.05 in 50 mL of YPD and incubated shaking (200

rpm) at 30 ^{o}C for 12 hrs. The culture was then removed from the incubator and a 1 mL sample of

the culture was labeled timepoint 0 (TP0) and frozen at -80 ^{o}C. To three 50 mL falcon tubes 15

mL of YPD was added and to another three tubes 15 mL YPD+ATc (250 ng/mL) was added.

The TP0 culture was subcultured into these tubes to an OD_{600} of 0.05 and incubated shaking (200

rpm) at 30 ^{o}C for 12 hrs. The six tubes were then subcultured again in their respective media

types and grown for 12 hrs under the same conditions. This was then repeated for a third and

final passage. At the end of each 12 hr out growth period (passages), 1 mL samples of each

culture were frozen at -80 ^{o}C.

Multi-environmental condition competitive growth assay

For this test an aliquot of the wildtype fungal pool was thawed on ice for 15 mins. Three 50 mL

falcon tubes were labeled TP0-YPD (time-Point 0) replicates (R)1-3 and were filled with 10 mL

of YPD. Another three falcon tubes were labeled TP0-ATC R1-3 and were filled with 10 mL of

YPD+ATc (250 ng/mL). To each tube, 200 µL of the thawed pool was added. The cultures were

incubated for 12 hrs at 37 ^{o}C shaking (250rpm). These cultures will be used as timepoint zero to

determine the initial frequency of each mutant in the pool. Media was prepared for each of the

conditions of interest (see Table S5 for media recipes). Media conditions were prepared with and

without ATc. To a 24 deep well plate (Fisher Scientific, cat. 14-222-349), 3 mL of each

condition (with and without ATc) were added to three wells. The TP0 cultures were subcultured

into each well at an OD_{600} of 0.1 with TP0-YPD R1 being subcultured into the well labeled R1

for each condition, following this for each of the other replicates and conditions. The subcultures

were grown at 37 ^{o}C shaking (550 rpm) for 12 hrs. After the first growth period the subculturing

method was repeated for another two rounds of passaging. After the third passage's outgrowth

was completed, samples were taken from each of the replicates. Sample volumes contained at

least ~ 5x10^7 cells and were frozen at -80 ^{o}C.

Competitive growth assay to evaluate changes in abundance in different genetic backgrounds

For this test an aliquot each of the three fungal pools was thawed on ice for 15 mins. For each fungal pool three 50 mL falcon tubes were labeled with the respective genetic background name and TP0-YPD (time-Point 0) replicates (R)1-3 and were filled with 10 mL of YPD. Another three falcon tubes were labeled TP0-ATC R1-3 and were filled with 10 mL of YPD+ATc (250 ng/mL). To each tube, 200 µL of the corresponding thawed pool was added. The cultures were incubated for 12 hours at 37 ^{o}C shaking (250rpm). These cultures will be used as timepoint zero to determine the initial frequency of each mutant in the pool. To a 24 deep well plate, 3 mL of YPD were added to three wells for each fungal pool and 3 mL of YPD+ATc were added to another three wells for each fungal pool. The TP0 cultures were subcultured into each well at an OD_{600} of 0.1 with TP0-YPD R1 being subcultured into the well labeled YPD R1, following this for each of the other replicates and pools. The subcultures were grown at 37 ^{o}C shaking (550 rpm) for 12 hrs. After the 12 hrs the subculturing method was repeated for another two rounds of passaging. After the third passage's growth period was completed, samples were taken from each of the replicates. Sample volumes contained at least ~5×10^{7} cells and were frozen at -80 ^{o}C.

Genomic DNA extraction

Genomic extractions were performed using the PureLink™ Genomic DNA Mini Kit (Invitrogen, cat. K182001) or with the phenol-chloroform method (Amberg 2005). With the kit, frozen samples were thawed, spun down, and the supernatant was removed. A master mix was prepared as follows: 4.55 g of D-Sorbitol (BioShop, cat. SOR508.1), 500 µL of 0.5 M EDTA (VWR, cat. E177-500ML), 0.5 M, 25 µL of β-mercaptoethanol (BioShop, cat. MER002.100), 0.0335 g Zymolyase (22,400 units/g, BioShop, cat. ZYM001.1), 24.48 mL nuclease-free water (final concentration of master mix 1 M sorbitol, 10 mM EDTA, 14 mM β-mercaptoethanol, 15 units/500 µL zymolyase). The cell pellet was resuspended in 500 µL of the master mix and incubated at 37 ^{o}C for 1-2 hrs (for cells from the clinical isolates backgrounds a longer incubation of 2 hrs was used). After incubation manufacturing instructions were followed to perform the kit genomic purification. After samples were extracted and purified, DNA concentrations were determined by Qubit analysis (analysed using a Qubit Flex Fluorometer with a high sensitivity DNA kit cat. Q33230) or by absorbance using a Tecan Infinite Pro with a NanoQuant plate, following the manufacturer's instructions.

For samples with low yields a phenol chloroform method was used in place of the kit. Briefly, frozen pellets were thawed and spun down into pellets and resuspended in 200 µL a homemade lysis buffer (2% Triton X-100, BioShop, cat. TRX777.500), 1% SDS, 100 mM NaCl, 10 mM Tris-Cl (pH 8.0), 1 mM Na2EDTA). To the resuspended sample 200 µL of phenol:chloroform:isoamyl alcohol (25:24:1) and a scoop (roughly the size of the original cell pellet) of acid washed glass beads (Sigma Aldrich, cat. G8772-100G) were added to each sample. Samples were beaten in a bullet blender for 4 mins. Samples were then pelleted at

13,000 xg for 5 mins. The top (aqueous) layer was transferred into a new "tough" micro-

centrifuge tube (Sigma, cat. EP022363344). To the separate aqueous layer, 200 µL of

phenol:chloroform:isoamyl alcohol (25:24:1) was added and the mixture was vortexed for 1 min.

Samples were then pelleted at 13,000 xg for 5 mins at room temperature. The top (aqueous) layer

was transferred to a standard microcentrifuge tube. A 1/10 volume of 3 M pH 5.2 sodium

acetate, and 2 volumes of chilled 95% ethanol was added to the tube and pipetted to mix.

Samples were then incubated at -20 ^{o}C for 15 mins. Samples were then centrifuged at 13,000 xg

for 15 mins at 4 ^{o}C. The supernatant was then discarded. The pellet was then re-suspended in 400

µL of pH 8.0 TE buffer with 3 µL of 10 mg/mL RNase and incubated for 5 mins at 37 ^{o}C. After

which 40 µL of 3 M pH 5.2 sodium acetate, and 800 µL of pre-chilled 95% ethanol and

incubated at -20 ^{o}C for 15 mins. Samples were then centrifuged at 13,000 xg for 15 mins and the

supernatant was discarded. The pellets were then dried in a 37 °C incubator for ~10 mins.

Pelleted DNA was then re-suspended in 50 µL of Tris-HCl (pH 8.0) or TE buffer (pH 8.0).

Nanodrop was then used to quantify the gDNA.

NGS library preparation and sequencing

NGS library preparation was performed in two successive PCR reactions. The first PCR involved

the use of amplicon specific primers attached to illumina sequencing adaptors. These primers

were designed to amplify the sgRNA region in the plasmid. Genomic DNA was diluted to 30

ng/µL. Briefly a master mix was prepared with 25µl of 2x NEBnext (NEB, cat. M0541L) and 5

µL of primers (see Table S4) diluted to 10 µM concentration per reaction. To a PCR tube 30 µL

of master mix and 20 µL diluted genomic DNA was added to PCR tubes for each reaction and

then run under the following thermocycler conditions 98^{o}C for 30 sec, cycle the following three

parameters 25 times: 98 ^{o}C for 10 sec, 61 ^{o}C for 30 sec, 72 ^{o}C for 30 sec, then a final step at 72^{o}C

for 2 mins. Following the PCR, each reaction was cleaned using the NucleoMag clean up kit

(Macherey-Nagel via D-MARK Biosciences, cat 744970.5) as per manufacturers instructions

using a 1.6:1 bead to DNA ratio. The clean-up resulted in 30 µL of cleaned amplicon product.

Qubit analysis was then used to quantify amplicon concentrations. For the second PCR, a set of

illumina index primers (see Table S4) were used. For this PCR a single reaction was set up for

each sample. A mastermix is prepared by adding 25 µL of 2x NEBnext and 15 µL of nuclease-

free water for each reaction. To each PCR tube, 40 µL of the master mix is added with 5 µL of 5

µM illumina index primer pairs and 5 µL of the cleaned amplicon (diluted to 3 ng/µL). The

following thermocycler conditions were then used: 98 ^{o}C for 30 sec, cycle the following three

parameters for 10 cycles: 98 ^{o}C for 10 sec, 55 ^{o}C for 30 sec, 72 ^{o}C for 30 sec, then a final step at

72^{o}C for 2 mins. From the reaction 45 µl of the PCR reactions was then cleaned using the

Nucleomag clean up kit with a bead to DNA ratio of 1:1. The final elution step resulted in 35 µL

of cleaned amplicon product. Qubit analysis was then performed on the samples to obtain the

DNA concentration of each sample. Samples were diluted to 5 ng/µL and were sent to the

University of Guelph Advanced Analysis center for quality control through RT-qPCR analysis to

identify if indices were correctly incorporated into the amplicon sequences. Libraries were

sequenced on either Illumina MiSeq or NovaSeq platforms. Samples were run on an Illumina
NovaSeq 6000 with a S4 flow cell for 150 cycles paired-end reads.

Competition assay data analysis

All data analysis and visualization was performed in Python 3 unless stated otherwise, using the
following packages: numpy v1.26.4 (Harris et al. 2020), matplotlib v3.9.2 (Hunter 2007), scipy
1.13.1 (Virtanen et al. 2020), pandas v2.2.2 (McKinney 2010), seaborn v0.13.2 (Waskom 2021).
Forward and reverse reads were first assembled using Pear v0.9.11 (Zhang et al. 2014), adjusting
overlap parameters depending on read length. The merged reads were then trimmed to keep only
a 80 bp region centered on the sgRNA, reverse complemented, and dereplicated using Fastx-
toolkit v0.0.14 (Gordon 2015), keeping the counts associated with each unique sequence. These
unique sequence clusters were then associated with library sgRNAs by perfect match of the
sgRNA sequence and an 8-nucleotide anchor on each side to obtain read counts. Counts were
normalized into frequencies using the following formula:

707
$$freq_{sgRNA} = \frac{counts_{sgRNA} + 1}{total\ counts}$$

709   From there, we measured changes in relative abundance using $\log_2$-fold change:

$$Log_2FC = Log_2(freq_{sgRNA}T_{end}) - Log_2(freq_{sgRNA}T_0)$$

We used a minimal read count threshold of 50 reads at T0 to filter out sgRNAs that were poorly
covered. We used the median of Log_2FC values across replicates for downstream analysis. To
determine what constituted a significant change in abundance, we fit a normal distribution to the
Log_2FC values of randomized sgRNAs. This allowed us to predict the expected false positive
rate as a function of Log_2FC values, and from there we set the significance threshold as the
Log2FC value at which the false discovery rate reached 5%. One of the random sgRNAs (#18,
AGCTCTACTGTTACAAGGGG) was found to induce CRISPRi specific fitness defects
reproducible across multiple screening conditions and genetic backgrounds, and as such was
excluded from the control distribution. GO annotations were retrieved from the *Candida*
Genome Database (Skrzypek et al. 2017). Thresholds were set separately by environmental
condition, genetic background, or background by environment combination. For Figure 4g and
Figure S12, we normalized the Log_2FC values of each condition using the parameters of the
randomized sgRNA distribution of each condition. The threshold for a significant difference in z-
score under caspofungin exposure was set to the mean of the normalized abundance change
thresholds at 5% FDRwith or without caspofungin 5% FDR.

Growth curve assays

Overnight cultures of *C. albicans* grown in YPD were diluted to an OD_{600} of 0.01 in YPD. From this dilution 100 μL was added to a 96 well plate containing 100 μL of YPD to obtain a final volume of 200 μL. Growth conditions were set to 37 ^oC with orbital shaking for 1000 sec at a 4 mm amplitude. Optical density measurements were taken at 20 min intervals over the course of 24 hours using a Tecan Infinite 200 PRO microplate reader.

RT-qPCR

Differential gene expression within mutants was performed through RNA-extractions and RT-qPCR. Overnight cultures of *C. albicans* were subcultured to an OD_{600} of 0.05 in 40 mL of YPD and grown at 37 ^oC shaking at 250 rpm to an OD_{600} between 0.2-0.5. Cultures were then centrifuged at 4000 rpm for 10 mins and the pellet was resuspended in ~ 200 μL of YPD and frozen at -80 ^oC. RNA extractions were performed using PuroSPIN™ Total RNA Purification Plus Kit (Luna Nanotech, cat. NK251-200). RT-qPCR was performed using a One-Step RT-qPCR kit (NEB, cat. E3005L). Real-time PCR was performed using a QuantStudio 3 Real-Time PCR system from Thermo Fisher Scientific Inc under the following cycling conditions 55 ^oC for 10 mins, 95 ^oC for 1 min, 40 cycles of a two-step qPCR (95^oC for 10 sec and 60 ^oC for 30 sec) and a Melt curve conditions were set for 95^oC for 15 s, 60^oC for 1 min and 95^oC for 1 sec. The C_T values for the experimental gene of interest were compared to the housekeeping gene to obtain a ΔC_T value for each strain. The ΔC_T for the mutants were compared to the ΔC_T of the control strain to obtain a $\Delta\Delta C_T$ value and used to calculate a fold difference in gene expression of the experimental gene of interest.

Whole Genome Sequencing (WGS)

Genomic extractions were performed using the Genomic extraction protocol described above, and samples were sent to Genscript for sequencing. WGS sequencing data was analyzed using a modified version of the Broad Institute Funpipe workflow with GATK v4.6.1.0 (van der Auwera and O'Connor 2020), Picard v3.1.0 (*Picard: A Set of Command Line Tools (in Java) for Manipulating High-Throughput Sequencing (HTS) Data and Formats such as SAM/BAM/CRAM and VCF*, n.d.), bwa199 v0.7.18 (Li and Durbin 2009), samtools v1.20 (Li and Durbin 2009), bcftools v1.19 (Danecek et al. 2021), bedtools v2.31.0 (Quinlan and Hall 2010). Briefly, adapter sequences were trimmed using GATK and Picard, and reads were aligned to the C. albicans SC5314 haploid reference genome (ASM18296v3203) using bwa-mem with default parameters. The alignments were then processed with GATK to prepare for variant calling using Haplotype caller with default settings except for -ploidy = 2. Variant calls were filtered using the following thresholds: for SNPs, QD > 12.5, QUAL > 30, SOR < 3, FS < 60, MQ > 40, and for indels QD > 20, QUAL > 30, FS < 200. After hard filtering, the variants were cross-referenced with a list of known heterozygous positions from a phased assembly (Skrzypek et al. 2017) and with sgRNA target sequence genome mapping generated using bowtie (Langmead et al. 2009). Potential

variant sites were examined manually using IGV (Robinson et al. 2011). We obtained high coverage for all three strains (WT: 83x coverage, ARB: 72x coverage, ERB: 71x coverage). No cases of aneuploidy, which are sometimes observed in antifungal-resistant strains, were detected. For the azole-resistant strain ARB, the presence of homozygous missense mutations T299A and F449S in *ERG11*, which are known to be sufficient to confer fluconazole resistance were validated. In ERB, the presence of a homozygous echinocandin resistance mutation (F641S) in *GSC1* (also known as *FKS1*) was validated .

Data availability

The sequencing data for pooled competition assays and whole-genome sequencing was deposited on the NCBI SRA (accession number:PRJNA1381690). Read counts and Log_2FC values of sgRNAs for each screen are provided as supplementary Tables S6, S7 and S8.

Code availability

Data analysis was performed using custom python 3.12.7 notebooks and scripts, available at https://github.com/TheShapiroLab/Calbicans_Pooled_CRISPRi_2025/.

Funding

P.C.D. was supported by a FRQS postdoctoral fellowship (https://doi.org/10.69777/376319). L.W., M.F., and N.C.G. were supported by the Natural Sciences and Engineering Research Council through PGS D awards. This work was supported by the Canadian Institutes of Health Research through grant PJT 162195 to R.S.S., the Canadian Institute for Advanced Research through a grant from the Fungal Kingdom: Threats & Opportunities program to both C.A.C. and R.S.S., and the National Institute of Allergy and Infectious Diseases through grant U19 AI110818. R.S.S. holds the Canada Research Chair in Microbial Functional Genomics and Synthetic Biology.

Acknowledgements

This research was enabled in part by support provided by SHARCNET (sharcnet.ca), Calcul Québec (calculquebec.ca) and the Digital Research Alliance of Canada (alliancecan.ca). The authors thank C. Landry for comments on the manuscript.

Competing interests

The authors declare no competing interests.

References

Amberg, David C. 2005. *Methods in Yeast Genetics : A Cold Spring Harbor Laboratory Course Manual*. Cold Spring Harbor Laboratory Press.

Arita, Yuko, Griffin Kim, Zhijian Li, et al. 2021. "A Genome-Scale Yeast Library with Inducible Expression of Individual Genes." *Molecular Systems Biology* 17 (6): e10207.

Auwera, Geraldine van der, and Brian D. O'Connor. 2020. *Genomics in the Cloud: Using Docker, GATK, and WDL in Terra*. O'Reilly Media.

Baetz, Kristin, Lianne McHardy, Ken Gable, et al. 2004. "Yeast Genome-Wide Drug-Induced Haploinsufficiency Screen to Determine Drug Mode of Action." *Proceedings of the National Academy of Sciences of the United States of America* 101 (13): 4525–4530.

Bédard, Camille, Alicia Pageau, Anna Fijarczyk, et al. 2025. "FungAMR: A Comprehensive Database for Investigating Fungal Mutations Associated with Antimicrobial Resistance." *Nature Microbiology* 10 (9): 2338–2352.

Bock, Christoph, Paul Datlinger, Florence Chardon, et al. 2022. "High-Content CRISPR Screening." *Nature Reviews. Methods Primers* 2 (1). https://doi.org/10.1038/s43586-022-00098-7.

Bonhomme, Julie, Murielle Chauvel, Sophie Goyard, Pascal Roux, Tristan Rossignol, and Christophe d'Enfert. 2011. "Contribution of the Glycolytic Flux and Hypoxia Adaptation to Efficient Biofilm Formation by Candida Albicans: Biofilm Formation and Adaptation to Hypoxia in C. Albicans." *Molecular Microbiology* 80 (4): 995–1013.

Bosch-Guiteras, Núria, and Jolanda van Leeuwen. 2022. "Exploring Conditional Gene Essentiality through Systems Genetics Approaches in Yeast." *Current Opinion in Genetics & Development* 76 (October): 101963.

Caplan, Tavia, Elizabeth J. Polvi, Jinglin L. Xie, et al. 2018. "Functional Genomic Screening Reveals Core Modulators of Echinocandin Stress Responses in Candida Albicans." *Cell Reports* 23 (8): 2292–2298.

Chavez, Alejandro, Marcelle Tuttle, Benjamin W. Pruitt, et al. 2016. "Comparison of Cas9 Activators in Multiple Species." *Nature Methods* 13 (7): 563–567.

Childers, Delma S., Ingrida Raziunaite, Gabriela Mol Avelar, et al. 2016. "The Rewiring of Ubiquitination Targets in a Pathogenic Yeast Promotes Metabolic Flexibility, Host Colonization and Virulence." *PLoS Pathogens* 12 (4): e1005566.

Cisneros, Angel F., Isabelle Gagnon-Arsenault, Alexandre K. Dubé, et al. 2023. "Epistasis between Promoter Activity and Coding Mutations Shapes Gene Evolvability." *Science Advances* 9 (5): eadd9109.

Danecek, Petr, James K. Bonfield, Jennifer Liddle, et al. 2021. "Twelve Years of SAMtools and BCFtools." *GigaScience* 10 (2): giab008.

Denning, David W. 2024. "Global Incidence and Mortality of Severe Fungal Disease." *The Lancet Infectious Diseases*, ahead of print, January 12. https://doi.org/10.1016/S1473-3099(23)00692-8.

Desai, Nirupa, Alan Brown, Alexey Amunts, and V. Ramakrishnan. 2017. "The Structure of the Yeast Mitochondrial Ribosome." *Science (New York, N.Y.)* 355 (6324): 528–531.

Fields, Francisco R., Shaun W. Lee, and Michael J. McConnell. 2017. "Using Bacterial Genomes and Essential Genes for the Development of New Antibiotics." *Biochemical Pharmacology* 134 (June): 74–86.

Fisher, Matthew C., Ana Alastruey-Izquierdo, Judith Berman, et al. 2022. "Tackling the Emerging Threat of Antifungal Resistance to Human Health." *Nature Reviews. Microbiology* 20 (9): 557–571.

Fu, Ci, Emily H. Xiong, Livia Kupczok, et al. 2025. "Expansion of the Functional Genomics GRACE Library Reveals Genes Relevant for Temperature-Dependent Fitness in Candida Albicans." *PLoS Biology* 23 (10): e3003409.

Fu, Ci, Xiang Zhang, Amanda O. Veri, et al. 2021. "Leveraging Machine Learning Essentiality Predictions and Chemogenomic Interactions to Identify Antifungal Targets." *Nature Communications* 12 (1): 6497.

Gerstein, Aleeza C., and Judith Berman. 2020. "Candida Albicans Genetic Background Influences Mean and Heterogeneity of Drug Responses and Genome Stability during Evolution in Fluconazole." *mSphere* 5 (3). https://doi.org/10.1128/mSphere.00480-20.

Gervais, Nicholas C., Alyssa A. La Bella, Lauren F. Wensing, et al. 2023. "Development and Applications of a CRISPR Activation System for Facile Genetic Overexpression in Candida Albicans." *G3 (Bethesda, Md.)* 13 (2). https://doi.org/10.1093/g3journal/jkac301.

Gervais, Nicholas C., Ruby K. J. Rogers, Madeleine R. Robin, and Rebecca S. Shapiro. 2025. "HyperdCas12a-Based Multiplexed Genetic Regulation in *Candida Albicans*." In *bioRxiv*. October 20. https://doi.org/10.1101/2025.10.20.683421.

Gordon, A. 2015. *Fastx_toolkit*. Github. https://github.com/agordon/fastx_toolkit/issues/7.

Halder, Viola, Brianna McDonnell, Deeva Uthayakumar, Jane Usher, and Rebecca S. Shapiro. 2020. "Genetic Interaction Analysis in Microbial Pathogens: Unravelling Networks of Pathogenesis, Antimicrobial Susceptibility and Host Interactions." In *FEMS Microbiology Reviews*. Preprint. https://doi.org/10.1093/femsre/fuaa055.

Halder, Viola, Caroline B. M. Porter, Alejandro Chavez, and Rebecca S. Shapiro. 2019. "Design, Execution, and Analysis of CRISPR-Cas9-Based Deletions and Genetic Interaction Networks in the Fungal Pathogen Candida Albicans." *Nature Protocols* 14 (3): 955–975.

Harris, Charles R., K. Jarrod Millman, Stéfan J. van der Walt, et al. 2020. "Array Programming with NumPy." *Nature* 585 (7825): 357–362.

Hirakawa, Matthew P., Diego A. Martinez, Sharadha Sakthikumar, et al. 2015. "Genetic and Phenotypic Intra-Species Variation in Candida Albicans." *Genome Research* 25 (3): 413–425.

Homann, Oliver R., Jeanselle Dea, Suzanne M. Noble, and Alexander D. Johnson. 2009. "A Phenotypic Profile of the Candida Albicans Regulatory Network." *PLoS Genetics* 5 (12): e1000783.

Hunter, John D. 2007. "Matplotlib: A 2D Graphics Environment." *Computing in Science & Engineering* 9 (3): 90–95.

Iracane, Elise, Cristina Arias-Sardá, Corinne Maufrais, Iuliana V. Ene, Christophe d'Enfert, and Alessia Buscaino. 2024. "Identification of an Active RNAi Pathway in Candida Albicans." *Proceedings of the National Academy of Sciences of the United States of America* 121 (17): e2315926121.

Jaffe, Mia, Adam Dziulko, Justin D. Smith, Robert P. St Onge, Sasha F. Levy, and Gavin Sherlock. 2019. "Improved Discovery of Genetic Interactions Using CRISPRiSeq across Multiple Environments." *Genome Research*, ahead of print, February 19. https://doi.org/10.1101/gr.246603.118.

Kacser, H., and J. A. Burns. 1981. "The Molecular Basis of Dominance." *Genetics* 97 (3-4): 639–666.

Keren, Leeat, Jean Hausser, Maya Lotan-Pompan, et al. 2016. "Massively Parallel Interrogation of the Effects of Gene Expression Levels on Fitness." *Cell* 166 (5): 1282–1294.e18.

Langmead, Ben, Cole Trapnell, Mihai Pop, and Steven L. Salzberg. 2009. "Ultrafast and Memory-Efficient Alignment of Short DNA Sequences to the Human Genome." *Genome Biology* 10 (3): R25.

Larrimore, Katherine E., and Giulia Rancati. 2019. "The Conditional Nature of Gene Essentiality." *Current Opinion in Genetics & Development* 58-59 (October): 55–61.

Lew-Smith, Jodi, Jonathan Binkley, and Gavin Sherlock. 2025. "The Candida Genome Database: Annotation and Visualization Updates." *Genetics* 229 (3). https://doi.org/10.1093/genetics/iyaf001.

Liao, Qian, Yu Zhang, Ying Chu, et al. 2025. "Application of Artificial Intelligence in Drug-Target Interactions Prediction: A Review." *NPJ Biomedical Innovations* 2 (1). https://doi.org/10.1038/s44385-024-00003-9.

Li, Heng, and Richard Durbin. 2009. "Fast and Accurate Short Read Alignment with Burrows-Wheeler Transform." *Bioinformatics (Oxford, England)* 25 (14): 1754–1760.

Maroc, Laetitia, Hajer Shaker, and Rebecca S. Shapiro. 2024. "Functional Genetic Characterization of Stress Tolerance and Biofilm Formation in Nakaseomyces (Candida) Glabrata via a Novel CRISPR Activation System." *mSphere*, January 24, e0076123.

McKinney, Wes. 2010. "Data Structures for Statistical Computing in Python." *Proceedings of the Python in Science Conference*, May 1, 56–61.

Momen-Roknabadi, Amir, Panos Oikonomou, Maxwell Zegans, and Saeed Tavazoie. 2020. "An Inducible CRISPR Interference Library for Genetic Interrogation of Saccharomyces Cerevisiae Biology." *Communications Biology* 3 (1): 723.

Noble, Suzanne M., Sarah French, Lisa A. Kohn, Victoria Chen, and Alexander D. Johnson. 2010. "Systematic Screens of a Candida Albicans Homozygous Deletion Library Decouple Morphogenetic Switching and Pathogenicity." *Nature Genetics* 42 (7): 590–598.

Ocana, Alberto, Atanasio Pandiella, Cristian Privat, et al. 2025. "Integrating Artificial Intelligence in Drug Discovery and Early Drug Development: A Transformative Approach." *Biomarker Research* 13 (1): 45.

Peng, Duo, and Rick Tarleton. 2015. "EuPaGDT: A Web Tool Tailored to Design CRISPR Guide RNAs for Eukaryotic Pathogens." *Microbial Genomics* 1 (4): e000033.

Perea, S., J. L. López-Ribot, W. R. Kirkpatrick, et al. 2001. "Prevalence of Molecular Mechanisms of Resistance to Azole Antifungal Agents in Candida Albicans Strains Displaying High-Level

Fluconazole Resistance Isolated from Human Immunodeficiency Virus-Infected Patients."
Antimicrobial Agents and Chemotherapy 45 (10): 2676–2684.

Perocchi, Fabiana, Lars J. Jensen, Julien Gagneur, et al. 2006. "Assessing Systems Properties of Yeast
Mitochondria through an Interaction Map of the Organelle." *PLoS Genetics* 2 (10): e170.

Picard: A Set of Command Line Tools (in Java) for Manipulating High-Throughput Sequencing (HTS)
Data and Formats such as SAM/BAM/CRAM and VCF. n.d. Github. Accessed December 2, 2025.
https://github.com/broadinstitute/picard.

Pun, Frank W., Ivan V. Ozerov, and Alex Zhavoronkov. 2023. "AI-Powered Therapeutic Target
Discovery." *Trends in Pharmacological Sciences* 44 (9): 561–572.

Quinlan, Aaron R., and Ira M. Hall. 2010. "BEDTools: A Flexible Suite of Utilities for Comparing
Genomic Features." *Bioinformatics (Oxford, England)* 26 (6): 841–842.

Rivera, Angela, and Joseph Heitman. 2023. "Natural Product Ligands of FKBP12: Immunosuppressive
Antifungal Agents FK506, Rapamycin, and beyond." *PLoS Pathogens* 19 (1): e1011056.

Robinson, James T., Helga Thorvaldsdóttir, Wendy Winckler, et al. 2011. "Integrative Genomics
Viewer." *Nature Biotechnology* 29 (1): 24–26.

Roemer, Terry, Bo Jiang, John Davison, et al. 2003. "Large-Scale Essential Gene Identification in
Candida Albicans and Applications to Antifungal Drug Discovery." *Molecular Microbiology* 50 (1):
167–181.

Román, Elvira, Ioana Coman, Daniel Prieto, Rebeca Alonso-Monge, and Jesús Pla. 2019.
"Implementation of a CRISPR-Based System for Gene Regulation in Candida Albicans." *mSphere* 4
(1). https://doi.org/10.1128/mSphere.00001-19.

Ropars, Jeanne, Corinne Maufrais, Dorothée Diogo, et al. 2018. "Gene Flow Contributes to
Diversification of the Major Fungal Pathogen Candida Albicans." *Nature Communications* 9 (1):
2253.

Segal, Ella Shtifman, Vladimir Gritsenko, Anton Levitan, et al. 2018. "Gene Essentiality Analyzed by In
Vivo Transposon Mutagenesis and Machine Learning in a Stable Haploid Isolate of Candida
Albicans." *mBio* 9 (5). https://doi.org/10.1128/mBio.02048-18.

Shapiro, Rebecca S., Alejandro Chavez, and James J. Collins. 2018. "CRISPR-Based Genomic Tools for
the Manipulation of Genetically Intractable Microorganisms." *Nature Reviews. Microbiology* 16 (6):
333–339.

Skrzypek, Marek S., Jonathan Binkley, Gail Binkley, Stuart R. Miyasato, Matt Simison, and Gavin
Sherlock. 2017. "The Candida Genome Database (CGD): Incorporation of Assembly 22, Systematic
Identifiers and Visualization of High Throughput Sequencing Data." *Nucleic Acids Research* 45
(D1): D592–D596.

Usher, Jane, and Ken Haynes. 2019. "Attenuating the Emergence of Anti-Fungal Drug Resistance by
Harnessing Synthetic Lethal Interactions in a Model Organism." *PLoS Genetics* 15 (8): e1008259.

Virtanen, Pauli, Ralf Gommers, Travis E. Oliphant, et al. 2020. "SciPy 1.0: Fundamental Algorithms for
Scientific Computing in Python." *Nature Methods* 17 (3): 261–272.

Waskom, Michael. 2021. "Seaborn: Statistical Data Visualization." *Journal of Open Source Software* 6
(60): 3021.

Wensing, Lauren, and Rebecca S. Shapiro. 2022. "Design and Generation of a CRISPR Interference
System for Genetic Repression and Essential Gene Analysis in the Fungal Pathogen Candida
Albicans." *Methods in Molecular Biology* 2377: 69–88.

Wensing, Lauren, Jehoshua Sharma, Deeva Uthayakumar, Yannic Proteau, Alejandro Chavez, and
Rebecca S. Shapiro. 2019. "A CRISPR Interference Platform for Efficient Genetic Repression in
Candida Albicans." *mSphere* 4 (1). https://doi.org/10.1128/mSphere.00002-19.

Wen, Wuqiang, Hongxuan Cao, Yunyuan Huang, et al. 2022. "Structure-Guided Discovery of the Novel
Covalent Allosteric Site and Covalent Inhibitors of Fructose-1,6-Bisphosphate Aldolase to
Overcome the Azole Resistance of Candidiasis." *Journal of Medicinal Chemistry* 65 (3): 2656–2674.

Yadav, Anshuman, Sudisht K. Sah, David S. Perlin, and Elena Rustchenko. 2024. "Candida Albicans
Genes Modulating Echinocandin Susceptibility of Caspofungin-Adapted Mutants Are Constitutively
Expressed in Clinical Isolates with Intermediate or Full Resistance to Echinocandins." *Journal of
Fungi (Basel, Switzerland)* 10 (3): 224.

Zeng, Rui, Erin Smith, and Antoni Barrientos. 2018. "Yeast Mitoribosome Large Subunit Assembly
Proceeds by Hierarchical Incorporation of Protein Clusters and Modules on the Inner Membrane."
Cell Metabolism 27 (3): 645–656.e7.

Zhang, Jiajie, Kassian Kobert, Tomáš Flouri, and Alexandros Stamatakis. 2014. "PEAR: A Fast and
Accurate Illumina Paired-End reAd mergeR." *Bioinformatics (Oxford, England)* 30 (5): 614–620.

Supplementary Figures 1-13

966
967 **Figure S1: Library coverage in the *C. albicans* wildtype strain pool. a)** Initial coverage
968 distribution in the library for the 551 sgRNAs. The red dotted line represents the minimal T0
969 abundance threshold for inclusion in Log$_2$FC calculations, 50 read counts. **b)** Proportion of genes
970 with partial or full sgRNA representation in the library.

Figure S2: Replicate correlation in the time-course competition assay.

a) Log$_2$FC in CRISPRi ON conditions after the third passage across replicates. Randomized sgRNAs (n=29) are shown in green, and those targeting essential gene promoters (n=438) are in black. The correlation between values was measured using Spearman's rank correlation. **b)** Spearman's rank correlation matrix across all time-course samples (n=469 sgRNAs per comparison).

978
979 **Figure S3: Significance threshold for sgRNA depletion. a)** False Discovery Rate (FDR) as a
980 function of Log_2FC threshold, based on the distribution of randomized sgRNAs abundance
981 change and the number of depleted sgRNAs at that threshold. **b)** FDR as a function of the
982 number of sgRNAs determined to be significantly depleted.
983
984
985

986
987 **Figure S4: Distribution of abundance changes across passages.** In all plots, the dotted red line
988 shows the significance threshold for depletion when comparing library and randomized sgRNAs.
989 **a)** Abundance changes for CRISPRi-ON (media without ATc) samples. **b)** Abundance change
990 for CRISPRi-OFF samples (media with ATc) **c)** Distributions of CRISPRi-ON/OFF Log$_2$FC
991 differences for the different passages.

Figure S5: Repression sensitive hit genes encoding mitochondrial ribosomal subunits do not cluster in space. The structure shown is the *Saccharomyces cerevisiae* mitochondrial subunit (pdb: 5MRC (Desai et al. 2017)**,** with orthologs corresponding to putative repression sensitive genes in *C. albicans* : *MRPL33* (*S.c. MRPL33*, red)*, C1_12610W* (*S.c.: MRPL15*, yellow)*, C2_07680W* (*S.c.: MRPL7*, purple)*, CR_01370C* (*S.c.: MRPS28*, blue)*, and CR_04140W* (*S.c.: NAM9*, green).

999
Figure S6: Inter-replicate and inter-condition correlation in the multi-condition screen.
Spearman's rank correlation matrix across all samples (n=469 sgRNAs per comparison) for
CRISPRi-OFF and CRISPRi-ON samples.

1003

1004 **Figure S7: Genes with constant repression sensitivity phenotypes.** For each gene, the overall

1005 distribution of Log₂FC for library sgRNAs for each condition is shown as a violin plot, with the

1006 randomized sgRNAs shown individually in green. The black lines indicate the significance

1007 threshold for each condition. Error bars show the 95% confidence interval around the mean

1008 Log₂FC. **a)** *MRPL33* **b)** *SPT23* **c)** *C2_07680W* **d)** *C2_03560C*.

1009
1010 **Figure S8: CRISPRi repression of FBA1 is comparable in glucose and glycerol conditions.**
1011 CRISPRi repression of *FBA1* in media with **a)** glucose or **b)** glycerol as the carbon source.

1012

1013 **Figure S9: Baseline *HSP90* expression levels in the three strains of interest.** Each dot

1014 represents a different biological replicate. Difference in expression compared with the wild-type

1015 was tested using a two-sided Welch's t-test: WT vs FLU[R]: p=0.22, WT vs CASP[R]: p=0.15.

1016

Figure S10: Pooled CRISPRi screening is reproducible in different genetic backgrounds. a)
Inter-replicate correlation for CRISPRi-ON samples of the wild-type strain library (R1 vs R2:
n=475, R1 vs R3: n=473, R2 vs R3: n=474). **b)** Inter-replicate correlation for CRISPRi-ON
samples of the FLU^R strain library (R1 vs R2: n=526, R1 vs R3: n=515, R2 vs R3: n=515). **c)**
Inter-replicate correlation for CRISPRi-ON samples of the CASP^R strain library (R1 vs R2:
n=521, R1 vs R3: n=519, R2 vs R3: n=519). d) Spearman's rank correlation matrix across all
backgrounds (n=436 sgRNAs per comparison) for CRISPRi-OFF and CRISPRi-ON samples.

1024
1025
Figure S11: Fitness effects of depleted sgRNAs can be validated across backgrounds Fitness
effects of sgRNAs in the pooled assay (top) and in small-scale growth curve assays (bottom).
The overall distribution of Log_2FC for library sgRNAs is shown as a violin plot, and the
randomized sgRNAs are shown individually in green. **a)** Effects for *MRPL33-2* (yellow) and
MRPL33-3 (purple) **b)** Effects for *ASK1-2* (yellow) and *ASK1-3* (purple).

1031
1032 **Figure S12: Few sgRNAs are affected by heterozygous or homozygous variants a)** Single
1033 nucleotide polymorphisms (SNPs) detected by whole genome sequencing for the three strains
1034 when compared to the SC5314 reference genome. **b)** Overlap between sgRNAs with target
1035 binding regions affected by SNPs. The lower left histogram shows the total number of sgRNAs
1036 affected for each strain.

1037

**Figure S13: CRISPRi antagonism or synergy with caspofungin in the FLU^R genetic
background.** Interaction between CRISPRi repression and Caspofungin treatment. Only genes
with at least one sgRNA showing antagonism or synergy of sufficient magnitude and statistical
significance (Benjamini-Hochberg corrected p-value <0.05) are shown as triangles and colored.
The direction of the triangle indicates the nature of the interaction, as indicated by the compass:
left: synergy in WT (*QCR8*); right: antagonism in WT (*MRPL33*, *SPT23*); down: synergy in
CASP^R (none); up: antagonism in CASP^R (*MMM1*, *MRPL33*, *QCR8*, *RSM22*, *SPT23*); upper
right: antagonism in both strains (*C2_035560C*). The black lines show the thresholds for the
magnitude of z-score change (see methods). Genes and sgRNAs not showing synergy or
antagonism are shown in grey.

1048 Supplementary Note 1: Outline of procedures to build *Candida albicans*
1049 CRISPRi pooled libraries

Pooled CRISPRi Library
Build Protocol

Day 0: Primer extension and Golden Gate
sgRNA target sequence insertion site
C. albicans CRISPRi plasmid
sgRNA target sequence
~2.5 hours
~13 hours

Day 1: Bacteria transformation
~4 hours
24-36 hours

Day 2: Harvest transformants and overnight cultures
~2.5 hours
12-18 hours

Day 3: Plasmid extraction and yeast transformation set up
~6.5 hours
12-18 hours

Optional Quality control check
NGS library preparation and sequencing

Day 4: Yeast transformation
~8 hours
48-72 hours

Day 6: Harvest yeast transformants
~2 hours

Day 7: gDNA extraction and NGS library preparation
~12 hours

Created in bioRender

1050
Before starting: Design a custom sgRNA library
1051
1. Design sgRNAs according to experimental goals (see methods: *sgRNA library design*).
1052
2. Order pooled oligo libraries from suppliers (In this work, we used IDT or Twist Bioscience oligonucleotide pools).
1053
1054
1055 ●**TIMING** 3-5 days to design; ~4-7 weeks for synthesis and delivery (dependent on pool size
1056 and supplier).
1057

Day 0: sgRNA cloning
1058
☐ Grow bacteria containing the plasmid backbone and extract the plasmid DNA.
1059
☐ Convert ssDNA oligo pool to dsDNA (see methods: *Primer extension*).
1060
☐ Clone pooled oligos into the CRISPRi plasmid backbone (see methods: *Golden Gate cloning*).
1061
1062
1063 ●**TIMING** *Recommended start time 2:00 pm.* Primer extension protocol ~1.5 hours; Golden
1064 Gate protocol ~1 hour; Thermocycler for ~13 hours.
1065

Day 1: Bacteria transformation
1066
☐ Transform the cloned plasmid pool into *E. coli* DH5α cells (see methods: *Bacteria Transformation*).
1067
1068

1069   ☐ During incubation steps in the bacteria transformation protocol, prepare the LB semi-
1070   solid agar for growing transformants.

1071 ●**TIMING** *Recommended start time 9:00 am.* Bacteria transformation protocol ~4 hours; Semi-
1072 solid agar preparation ~1.5 hours; Incubation period 24-36 hours or until colonies become
1073 visible.
1074

Day 2: Harvest Bacteria pooled library

1076   ☐ Harvest bacteria colonies (see methods: *Bacteria Transformation*).
1077   ☐ Prepare an overnight culture for plasmid extraction (see methods: *Plasmid pool*
1078   *extraction*).

1079 ●**TIMING** *Recommended start time 3:00 pm.* Harvesting bacteria cells ~2 hours; Starting
1080 overnight culture ~30 min; Incubation period ~12-18 hours.
1081

Day 3: Extract pooled plasmid Library and preparation for Fungal transformation

☐ Perform plasmid DNA extraction to obtain a large quantity of plasmid (see methods:
Plasmid pool extraction).
☐ Linearize pooled plasmids by restriction digest (see methods: *Candida albicans*
transformation).
☐ Prepare an overnight culture of the selected strain background (see methods: *Candida*
albicans transformation).

●**TIMING** *Recommended start time 9:00 am.* Plasmid extraction protocol ~5 hours; Overnight
culture set-up, and restriction enzyme digest ~1.5 hours; Incubation period ~12-18 hours.
1092

Optional (Highly recommended) Sequencing of the pooled plasmid library.

☐ After extracting the plasmid library, you can perform high-throughput sequencing to
determine the composition of your plasmid library and validate adequate representation
before moving on to fungal transformation steps (see methods: *NGS library preparation*
and sequencing).

●**TIMING** *Recommended start time 9:00 am.* Library preparation for sequencing ~8 hours;
Sequencing turnaround time dependent on the provider.
1100

Day 4: *Candida albicans* transformation

1102   ☐ Transform *C. albicans* with the pooled CRISPRi plasmid library (see methods: *Candida*
1103   *albicans transformation*).
1104   ☐ Prepare the YPD semi-solid agar during incubation steps.

1105 ●**TIMING** *Recommended start time 8:00 am.* Fungal transformation protocol ~8 hours; Semi-
1106 solid agar protocol ~1.5 hours; Incubation period 48-72 hours or until colonies become visible.
1107

Day 6: Harvest fungal pooled library

    ☐ Harvest fungal library (see methods: *Candida albicans transformation*).

    ☐ Prepare frozen aliquots of the fungal library for storage and downstream assays.

●**TIMING** *Recommended start time 4:00 pm.* Harvesting fungal cells ~2 hours.

Day 7-8: Sequencing of the pooled plasmid library.

    ☐ Extract Genomic DNA from a sample of your pooled fungal library (see methods: *Genomic DNA extraction*).

    ☐ Prepare samples for sequencing (see methods: *NGS library preparation and sequencing*).

●**TIMING** *Recommended start time 9:00 am.* Genomic extractions ~4 hours. Library preparation for sequencing ~8 hours.