B-vitamin requirements and phycosphere interactions for the HAB-forming

dinoflagellate, *Pyrodinium bahamense* var. *bahamense*

Lydia Ruggles Howe[1], Lauren D' Amore[1,2], Cary Lopez[2], Sugandha Shankar[2], Margaret Mars

Brisbin[1]*

[1] College of Marine Science, University of South Florida, St. Petersburg, FL, U.S.A.

[2] Florida Fish and Wildlife Conservation Commission, Fish and Wildlife Research Institute, St.

Petersburg, FL, U.S.A.

*Corresponding Author:

830 1st Street South, St. Petersburg, Florida, U.S.A.

mmarsbrisbin@usf.edu

Keywords: Cobalamin, dinoflagellate, *Pyrodinium*, Harmful Algal Bloom, HAB, Phycosphere,

phytoplankton-bacteria interactions, B-vitamins, B_{12}

Running title: *Pyrodinium* phycosphere and vitamin needs

Abstract:

Pyrodinium bahamense is a saxitoxin-producing dinoflagellate found in tropical and subtropical
estuarine waters globally, with the var. *bahamense* variety forming recurrent harmful algal
blooms (HABs) in Florida estuaries. The intensity and duration of bloom peaks in Florida
estuaries vary interannually, but life cycle transitions, estuarine water residence time, water
temperature, salinity, and nutrient concentrations appear to contribute to bloom formation.
However, the impact of B-vitamin availability on *P. bahamense* bloom formation has not been
investigated, despite the knowledge that many HAB dinoflagellates require B-vitamins for
growth and bloom development. In this study, we determined that of the three B-vitamins most
commonly required by marine phytoplankton—biotin (B_7), thiamine (B_1), and cobalamin (B_{12})—
P. bahamense strictly requires an exogenous B_{12} source to sustain growth. As bacterial
interactions can fulfill phytoplankton B_{12} requirements, we tested whether bacterial communities
recruited from two Florida estuaries could rescue *P. bahamense* from B_{12} limitation. We
demonstrate that bacteria from the Indian River Lagoon (IRL) estuary collected during a *P.*
bahamense bloom restored *P. bahamense* growth in media missing B_{12}, while bacteria present
in Old Tampa Bay (OTB) water collected when *P. bahamense* was not present did not restore
P. bahamense growth in media missing B_{12}. Putative bacterial B_{12} producers exclusively
recruited from IRL included Enterobacteraceae genera that likely have anthropogenic sources.
Our findings suggest B_{12} is essential for fueling *P. bahamense* blooms, and that specific
bacterial communities may contribute to bloom development through B_{12} production.

Introduction

Phytoplankton are critical to aquatic ecosystem functioning and global biogeochemical cycles [1–3], but phytoplankton can also negatively impact aquatic ecosystems through the formation of harmful algal blooms (HABs). During a HAB, phytoplankton species undergo mass proliferation and may also produce toxins, leading to water column hypoxia, benthic anoxia, wildlife mortalities, and human illnesses [4, 5]. Thus, it is imperative to fully understand the physiology of HAB-forming phytoplankton species to build predictive models for bloom initiation, duration, and intensity. This is increasingly important as changing climate and weather patterns are amplifying the effects of HABs on coastal ecosystems and communities [6].

B-vitamins—such as thiamine (B_1), biotin (B_7), and cobalamin (B_{12})—are integral to the growth of many phytoplankton species, as they are essential cofactors for key enzymes involved in cellular metabolism and reproduction [7]. Genomic surveys indicate that more than half of all phytoplankton and up to 90% of HAB dinoflagellates are only equipped with the B_{12}-dependent methionine synthase (MetH) [8], making them fully reliant on external B_{12} sources for essential amino acid synthesis. Through a comprehensive set of bioassays, Tang et al. (2012) demonstrated that 26 out of 27 HAB species (41 strains) required B_{12} and 20 required B_1 [9], further suggesting B-vitamin availability is especially critical to the growth of HAB-forming phytoplankton. However, the extent of the role B-vitamin availability plays in promoting or limiting HABs is not fully elucidated [10].

Despite reliance on B vitamins, many HAB species bloom in coastal waters where measurable dissolved B-vitamin concentrations are below phytoplankton growth requirements, suggesting that B vitamins may be limiting nutrients in coastal estuaries [10]. Moreover, studies in Long Island estuaries demonstrated a drawdown of vitamins B_1 and B_{12} during dinoflagellate blooms, and B_1 and B_{12} amendments in concurrent incubation experiments increased dinoflagellate biomass [11]. Bacterial production of B vitamins stands to be the major source of these critical resources to marine ecosystems. Roughly 25% of sequenced bacterial genomes contain complete B_{12}-synthesis pathways [12], but most B_{12} producers do not freely share the resource [13]. Thus, specific relationships between phytoplankton and bacteria inhabiting the phycosphere may be particularly important for fulfilling phytoplankton B_{12} requirements [14, 15]. Indeed, positive interactions between several HAB dinoflagellates and specific B-vitamin producing bacteria have been identified [16–18].

The saxitoxin-producing HAB dinoflagellate, *Pyrodinium bahamense* var. *bahamense,* forms dense assemblages ($> 100,000$ cells L^{-1}) in Florida estuaries, including in the northern

segment of Tampa Bay, Old Tampa Bay (OTB), on Florida's west coast and the Indian River

Lagoon (IRL) estuary on the east coast [19] (Fig. 1). B vitamins have long been hypothesized to

be contributing factors in supporting the high concentrations of *P. bahamense* observed in the

bioluminescent bays of Puerto Rico, where the dinoflagellate blooms carry ecological, cultural,

and economic significance [20]. However, the role of B-vitamin availability and bacterial

interactions in supporting *P. bahamense* blooms in Florida estuaries has yet to be directly

investigated, making it plausible that characterizing these interactions may improve our

understanding of bloom intensity and duration in the region. This study directly investigates the

B-vitamin requirements of a *P. bahamense* strain isolated from OTB and determines whether

bacterial communities recruited from OTB and IRL seawater can support *P. bahamens*e growth

in the absence of vitamin B_{12}.

Materials and Methods

P. bahamense *strains used in experiments*

P. bahamense strains FWC1006—originating from OTB in 2023—and FWC1003—

originating from the IRL in 2017—were acquired from the Florida Fish and Wildlife Conservation

Commission (FWC) Fish and Wildlife Research Institute (FWRI) for use in this study (Fig. 1).

Both strains were established by germinating resting cysts from sediment samples. Germinated

cells were then isolated and grown into unialgal cultures using aseptic techniques, but were not

deliberately rendered axenic. Upon transfer to our lab, both strains were maintained in L1/4

media prepared with sterile 25 PSU artificial seawater (ASW; Instant Ocean) [21] at 24ºC on a

12:12 h light: dark cycle with 50% illumination (~115 µM photons m^{-2} s^{-1}) in a Percival I-36 LLVL

incubator.

Fig. 1 Map depicting regions of Florida (U.S.A.) with recurrent *Pyrodinium bahamense* blooms and where samples were collected for this study. **A** Map of Florida with inset regions indicated (Tampa Bay and OTB; **B** and Indian River Lagoon; **C**). Red points in panels **B** and **C** indicate the locations where seawater was collected for bacterial community transfers to FWC1006.

Characterization of bacterial communities in P. bahamense *cultures*

Bacteria associated with phytoplankton can provide B vitamins to support phytoplankton [7, 16, 22], making it imperative to characterize bacteria co-cultured with *P. bahamense* before experimentation. The presence of bacteria in *P. bahamense* strains FWC1006 and FWC1003 was initially evaluated by transferring aliquots into two test broths for bacterial growth: Difco Marine Broth and an artificial seawater tryptone broth (ASWT) matching the salinity of algal culture media [23]. ASWT broth was prepared from 1 L of 25 PSU Instant Ocean ASW amended with 5 g tryptone, 3 g yeast extract, and 3 mL 50% glycerol (in water, RICCA Chemical Company), and sterilized by autoclaving and filtering with a 0.2-μm pore-size filter [23]. Aliquots of each isolate (100 μl) were transferred to triplicate wells containing 1 mL of test broth in a sterile 24-well plate. Triplicate negative controls for each test broth were included. Plates were incubated at room temperature, and bacterial growth was visually monitored after 72 h and checked under 400 x magnification with an inverted microscope.

Bacterial communities present in cultures of the two *P. bahamense* strains were further
characterized through metabarcode sequencing. Biomass was collected from each strain by
sequentially filtering cultures through 10.0-μm and 0.2-μm pore-size filters under gentle vacuum
pressure. *P. bahamense* cells and associated bacteria were retained on 10.0-μm pore-size
filters, and free-living bacteria were retained on 0.2-μm pore-size filters. RNA was extracted
from filters using the RNeasy PowerWater Kit (Qiagen) according to the manufacturer's protocol
with the suggested 10-minute 65°C incubation for improved cell lysis. RNA was reverse
transcribed following the manufacturer's protocols for the High-Capacity RNA-to-cDNA Kit
(Applied Biosystems). The V3–V4 hypervariable regions of the reverse-transcribed 16S rRNA
were amplified following the "16S Metagenomic Sequencing Library Preparation" Illumina
protocol, including the recommendations for primers [24] and PCR conditions. Metabarcode
libraries were sequenced on an Illumina NextSeq2000 2x300-bp P1 flow cell. Amplicons were
processed with the QIIME2 pipeline [25], utilizing the DADA2 denoising algorithm [26], and
representative sequences were taxonomically classified using a Naive Bayes classifier [27]
trained on the SILVA Ref 99 database v138.1 [28]. Amplicon Sequence Variants (ASVs) were
further processed using the phyloseq [29] package in R Studio [30]. Based on results from
bacterial growth tests and metabarcode analyses, the FWC1006 *P. bahamense* isolate from
OTB was selected for subsequent experiments.
135
Evaluating the B-vitamin requirements of P. bahamense
Given the difficulty of growing dinoflagellates axenically, dinoflagellate cultures with
reduced or knocked-down microbiomes have been used in experiments to minimize the impact
of bacterial processes on experimental results [17]*.* Here, we propose that the low-diversity
bacterial community associated with the FWC1006 *P. bahamense* strain from OTB represents a
reduced bacterial community and an opportunity for testing the B-vitamin requirements of *P.*
bahamense without subjecting cultures to extensive antibiotic treatment.
To determine the B-vitamin requirements for *P. bahamense*, the OTB strain was grown
in L1/4 media missing one of the three major B-vitamins—B_{12} (cobalamin), B_1 (thiamine), or B_7
(biotin)—and in replete L1/4 (i.e., containing all three B vitamins) as a positive control. All media
formulations were made using sterile 25 PSU ASW amended with $NaNO_3$, NaH_2PO_4, and trace
metal solution from the NCMA L1 Medium kit at 1/4 the concentration of L1 medium [21].
Individual B vitamins (Sigma-Aldrich) were added to media at 1/4 the concentration in L1
medium [21]. Experimental cultures were set up with 1 mL of the OTB isolate and 45 mL of each

media treatment in triplicate 50 mL cell culture flasks and incubated as described above [31].
Growth was monitored by measuring *in vivo* Chlorophyll-*a* fluorescence (RFU) with a Trilogy
fluorometer (Turner Designs) every 48 h. The experiment was maintained for four trials, with 1
mL serial transfers made in the late exponential growth phase or early stationary phase. The $-$
B_{12} treatment was not serially transferred because these cultures failed to grow. Instead, 1 mL
of the culture in replete media was also transferred to $-B_{12}$ media at the beginning of each new
trial. Growth rates were calculated from Chlorophyll-*a* fluorescence using the fit_easylinear
function from the growthrates R package (h = 5, quota = 1) [32].
To better characterize the minimum B_{12} requirement of *P. bahamense*, the OTB isolate
was grown in L1/4 media with varying final concentrations of vitamin B_{12} from 369 pM—the
concentration in L1 medium [21]—through 0.0369 pM, prepared through 10% serial dilutions.
Culture conditions matched those described above, and *in vivo* Chlorophyll-*a* fluorescence was
measured every 48 h for a single 40-day trial.
163
Evaluation of bacterial community potential to supplement B_{12} and support P. bahamense
blooms
Next, we tested the ability of bacterial communities from OTB and IRL to produce
vitamin B_{12} and promote *P. bahamense* growth. OTB surface seawater was collected on May 2,
2024 (27.9055ºN, −82.6046ºW) and IRL surface seawater was collected on July 10, 2024
(28.5136ºN, −80.6125ºW) by FWRI (Fig. 1). At the time of collection, no *P. bahamense* cells
were detected at the OTB site, whereas *P. bahamense* cells were detected at concentrations
exceeding the FWC-designated bloom level at the IRL site (113,333 cells L^{-1}, bloom-level:
>100,000 cells L^{-1}). Seawater from both sites and a *P. bahamense* culture (IRL FWC1003) were
sequentially filtered through 10.0-µm, 0.8-µm, and 0.45-µm pore-size filters to remove
eukaryotic phytoplankton, but particle-associated bacteria and larger or filamentous bacteria
were likely also removed or reduced during filtering. A 250 mL aliquot of the final filtrate for each
recruitment treatment was filtered through a 0.2-µm pore-size filter, and the remaining filtrate for
each recruitment treatment was diluted 50% with L1/4 media and used in recruitment trials. The
10.0-µm and 0.2-µm filters were flash frozen and stored at −80ºC for DNA extraction.
A 1-mL aliquot of OTB *P. bahamense* culture was transferred into 45 mL of each of the
three recruitment treatment filtrates (OTB seawater, IRL seawater, IRL culture) and incubated
for two weeks to allow equilibration of the introduced bacterial communities. Subsequently, 1 mL
aliquots of cultures with recruited bacterial communities were transferred in triplicate to 45 mL of

replete L1/4 and media without B_{12} (L1/4 $-B_{12}$), and grown and monitored as described above

for two trials. For the second trial, cultures were serially transferred if growth occurred, which

was the case for all cultures grown in replete media but only for the IRL seawater-treated culture

grown in L1/4 $-B_{12}$. The OTB seawater- and IRL culture-treated cultures in L1/4 $-B_{12}$ were

restarted from the matching treatment in replete media in trial 2, since they failed to grow in trial

1. All culture replicates from growth trial 2 were sequentially filtered through 10.0-μm and 0.2-

μm pore-size filters under gentle vacuum pressure when cultures began entering stationary

growth phase. Filters were flash frozen and stored at -80°C for DNA extraction.

Nanopore full-length 16S metabarcoding and bioinformatics

The bacterial communities introduced from seawater sources to OTB *P. bahamense* and

the established "recruited" communities from each source (IRL *P. bahamense* culture, IRL

seawater, and OTB seawater) were assessed through full-length 16S rRNA gene sequencing

on the Oxford Nanopore MinION platform. DNA was extracted from 10.0-μm and 0.2-μm pore-

sized filters with the Qiagen DNeasy PowerSoil Pro kit according to the manufacturer's protocol,

including the additional 65°C heating step. The full-length 16S rRNA gene was amplified using

forward primer S-D-008-c-S-20 (5′-AGRGTTYGATYMTGGCTCAG-3′) [24] and reverse primer

1492R (5′-RGYTACCTTGTTACGACTT-3′) [33] with KAPA2G Robust HotStart (Roche) in the

following PCR program: 95°C/ 3min, 35x (95°C/ 20sec, 57°C/ 20sec, 72°C/ 30sec), 72°C/ 1min,

and 4°C/ ∞ [33]. Sequencing libraries were prepared using the Oxford Nanopore Native

Barcoding Kit 24 v14 following the manufacturer's protocol and sequenced on two MinION v10.4

flow cells and one Flongle flow cell. Sequence data were processed (min. length = 1300 bp,

max. length = 2000 bp), quantified, and classified using Kraken v2.1.2 [34] within the wf-16s

nextflow pipeline in the EPI2ME platform [35] with the SILVA v138.1 taxonomic database [28].

Results were exported to R Studio for further processing.

Results

Assessment of bacterial communities co-cultured with P. bahamense *isolates.*

There was no visible bacterial growth (broth turbidity or observable cells by microscopy)

in Difco Marine Broth or ASWT broth 72 h after inoculation with FWC1006 *P. bahamense*

culture. Similarly, there was no visible bacterial growth in Difco Marine Broth or in salinity-

matched ASWT broth in the negative control wells after the 72-hour observation period. In
contrast, there was visible bacterial growth (high turbidity) in both Difco Marine Broth and
salinity-matched ASWT broth in wells inoculated with FWC1003 *P. bahamense* culture. Further
investigation into the bacterial community composition in cultures of both *P. bahamense* strains
through metabarcoding also revealed stark differences. The bacterial community in the OTB *P.*
bahamense culture had very low alpha diversity (Shannon Index) compared to the communities
in IRL *P. bahamense* cultures (Fig. 2A). The IRL *P. bahamense* culture hosted bacteria from 11
taxonomic orders, including typical phytoplankton-associated groups such as Alteromonadales,
Caulobacterales, and Rhodobacterales (Fig. 2B). Conversely, the OTB *P. bahamense* culture
hosted bacteria from just three orders: Microtrichales, Oceanospirillales, and Rhizobiales (Fig.
2B). At the amplicon sequence variant (ASV) level, only nine ASVs were detected among OTB
P. bahamense cultures and just six ASVs made up > 99% of sequencing reads (Table S1).
Further investigation of these ASVs through BLAST searches revealed that they were
previously found in soils, marine sediments, and aquatic plant rhizospheres (Table S1). Thus,
the OTB *P. bahamense* culture has a reduced bacterial community compared to the IRL *P.*
bahamense culture. Moreover, the bacteria associated with the OTB *P. bahamense* culture
likely do not represent water column bacteria that *P. bahamense* would interact with during
blooms of vegetative cells.
233

Fig. 2 Alpha diversity and composition of bacterial communities in *P. bahamense* cultures. **A** Shannon
diversity index for particle-associated (>10.0-µm size fraction; blue) and free-living (0.2–10.0-µm size

fraction; red) bacterial communities in cultures of FWC1003 and FWC1006 *P. bahamense* isolated from

Indian River Lagoon and Old Tampa Bay, respectively. **B** Relative abundance of bacterial orders in size-

fractioned communities in *P. bahamense* cultures.

B-vitamin requirements of OTB P. bahamense

Given the low-diversity bacterial community detected in the OTB *P. bahamense* culture,

this strain was used in experiments testing the B-vitamin requirements of *P. bahamense*. We

hypothesized that the low-diversity sediment-derived bacteria in the OTB culture would be less

likely to symbiotically provide B-vitamins than the higher-diversity community observed in the

IRL culture, which was more similar to communities observed in other phytoplankton cultures

(e.g., [36–38]).

Growth curves for OTB *P. bahamense* appeared similar across four serial transfers in

replete L1/4 media, L1/4 –thiamine, and L1/4 –biotin (Fig. 3A). In contrast, OTB *P. bahamense*

immediately ceased growing when transferred into L1/4 media missing vitamin B_{12} (L1/4 –B_{12},

Fig. 3A). As a result, serial transfers could not be made for this treatment, and culture from the

replete treatment was transferred to L1/4 –B_{12} in each transfer round. Results from a one-way

ANOVA indicated that the effect of media formulation on maximum chlorophyll-*a* fluorescence, a

proxy for biomass, was significant (p = .0000, F = 45.35), and a Tukey HSD multiple

comparisons of means test showed that the maximum fluorescence was significantly lower in

L1/4 –B_{12} media than in replete (*p-adj* = .0000), –thiamine (*p-adj* = .0000), and –biotin media (*p-*

adj = .0000). Of note, the maximum fluorescence reached in media missing thiamine was

significantly greater than in replete media (*p-adj* = .0000)or in –biotin media (*p-adj* = 0.0006).

Since *P. bahamense* ceased growing when transferred into media missing B_{12}, reliable growth

rates could not be determined for this treatment. Aside from the –B_{12} treatment, media

formulation did not have a large effect on growth rate (Fig. 3C, S1); a one–way ANOVA

excluding the –B_{12} treatment indicated that media formulation (i.e., –thiamine, –biotin, replete)

did not significantly affect growth rate (μ; p = 0.172, F = 1.861).

Fig. 3 Growth responses of *P. bahamense* when grown in replete L1/4 media and in media missing B_{12}, thiamine, or biotin. All data are plotted with color representing media treatment. **A** *In vivo* Chlorophyll *a* fluorescence (Chl-*a*) was used to track *P. bahamense* growth across four trials. Solid lines indicate the smoothed conditional mean, plotted with the geom_smooth function in the R package ggplot2 (method = 'loess'). **B** Maximum Chl-*a* values in each replicate by media formulation. Results from post-hoc comparisons to replete are shown with the following significance codes: *p-adj* > 0.05; 'n.s.' and *p-adj* < 0.001; '***'. **C** Growth rate of each replicate in –thiamine, –biotin, and replete media. Growth rates could not be calculated for replicates in –B_{12} media due to the lack of growth (Fig. S1). Means are plotted as black open circles, and error bars represent one standard deviation of the mean.

P. bahamense *growth response to varying B_{12} availability.*

Given that withholding B_{12} rapidly diminished *P. bahamense* growth (Fig. 3), we further evaluated the *P. bahamense* growth response to varying B_{12} concentrations using 10% serial dilutions of B_{12} in L1/4 media. *P. bahamense* growth showed a typical dose-response to increasing B_{12} availability (Fig. 4A). Maximum biomass (Chl-*a* fluorescence) increased with increasing B_{12} availability (Fig. 4B), with significant increases in maximum fluorescence between

0.0369 and 3.69 pM B_{12} and between 3.69 and 36.9 pM B_{12} (*p-adj* = 0.025 and 0.0000,

respectively; Tukey HSD). However, when enough B_{12} was available to support growth, the

growth rates were fairly consistent (Fig. 4C, S2). Growth rate was depressed in media with

0.0369 pM B_{12} (Fig 4C), but one-way ANOVA results did not indicate that B_{12} concentration

significantly affected growth rate (*p* = 0.196, *F* = 1.85, Fig 4C). Notably, Chl-*a* fluorescence in

0.0369 pM B_{12} media remained low and variable throughout the trial, leading to poor linear

model fits for estimating growth rate (*p* > 0.05, R^2 < 0.85, Fig. S2) that appear to overestimate

growth rates and skew the statistical test results.

Fig. 4 Growth responses of *P. bahamense* when grown in L1/4 media with a 10% dilution series of B_{12}

concentrations. All data are plotted with color representing the B_{12} concentration. *In vivo* Chlorophyll-*a*

fluorescence (filled circles) was used to track *P. bahamense* growth in L1/4 media with five

concentrations of B_{12}: 0.0369 pM, 0.369 pM, 36 pM, and 369 pM. Solid lines indicate the smoothed

conditional mean, plotted with the geom_smooth function in the R package ggplot2 (method = 'loess'). **B**

Maximum Chl-*a* fluorescence values and **C** growth rates (filled circles), shown for each replicate grouped

by media B_{12} concentration. Means are plotted as black open circles, and error bars represent one

standard deviation of the mean. Results from key post-hoc comparisons are shown with the following

significance codes: *p-adj* > 0.05; 'n.s.', *p-adj* < 0.05; '*', and *p-adj* < 0.001; '***'.

Effects of recruited bacterial communities on P. bahamense *growth*

Increased growth was detected in OTB *P. bahamense* (FWC1006) with bacteria

recruited from IRL seawater compared to *P. bahamense* with bacteria recruited from IRL culture

or OTB seawater when grown in L1/4 $-B_{12}$ media (Fig. 5A). One-way ANOVA indicated that the

recruited bacterial community significantly affected maximum Chl-*a* when B_{12} was withheld (Fig.

5B; $p = 0.0001$, $F = 50.27$). The maximum Chl-*a* in cultures with bacteria from IRL seawater was

significantly higher than in cultures with bacteria from OTB seawater, bacteria from the IRL

culture, or with the initial bacteria in the OTB culture (*p-adj* = 0.0000 for each comparison,

Tukey HSD). Growth rates for *P. bahamense* with bacteria from IRL seawater were not

significantly different when grown in replete media versus media missing B_{12} (two-sample t-test;

$p = 0.5678$, $t(10) = -0.5907$; Fig. 5C, S3). These results indicate that at least some of the

bacteria in the community recruited from IRL seawater are B_{12} producers and can support P.

bahamense growth.

Fig. 5 Growth responses of OTB *P. bahamense* (FWC1006) with bacterial communities recruited from
IRL culture (FWC1003), OTB seawater, and IRL seawater when grown in L1/4 media missing vitamin B_{12}
and replete L1/4 media. All data are plotted with color representing media treatment. **A** *In vivo*
Chlorophyll-*a* fluorescence (filled circles) was used to track the growth of OTB *P. bahamense* with
bacteria from three sources (IRL culture, OTB seawater, and IRL seawater) in replete L1/4 media and
L1/4 media missing B_{12} (L1/4 $-B_{12}$). Solid lines indicate the smoothed conditional mean, plotted with the
geom_smooth function in the R package ggplot2 (method = 'loess'). **B** Maximum Chl-*a* fluorescence
values (filled circles) in L1/4 $-B_{12}$ are shown for each replicate, grouped by bacterial community source.
Note, the maximum Chl-*a* reached in L1/4 $-B_{12}$ by OTB *P. bahamense* with its initial low-diversity
bacterial community is included for reference (from Fig. 3). Results from key post-hoc comparisons are
shown with the following significance code: *p-adj* < 0.001; '***'. **C** Growth rates (filled circles) for each
replicate with bacteria from IRL seawater, grouped by media treatment. Means are plotted as black open
circles, and error bars represent one standard deviation of the mean.

Sequencing results

Overall, this project produced 10,431,416 Nanopore full-length 16S reads that passed

quality filters. After applying a length filter (min = 1300 bp, max = 2000 bp), 8,526,020 reads

remained. Culture samples sequenced on MinION flow cells yielded 128,931–500,307 (median

= 322,769) reads per sample and retained 103,950–407,502 (median = 272,432) after applying

the length filter. Seawater samples (bacterial source cultures) sequenced on a Flongle flowcell

yielded 34,389–141,236 (median = 93,931) reads per sample and retained 12,221–53,990

(median = 37,987) reads per sample after length filtering. Alpha diversity (richness and

Shannon index, based on genus) was higher in the IRL seawater (richness = 1,695–1,805,

Shannon = 4.82–4.89) than the OTB seawater (richness = 1,056–1,208, Shannon = 3.24–4.69;

Fig. S5, S6). Alpha diversity was similar across recruited communities (Fig. S5, S6).

Comparison of bacterial communities recruited from different sources

Distinct bacterial communities were recruited to the FWC1006 *P. bahamense* cultures

from the three sources (FWC1003 *P. bahamense* cultures (IRL), IRL seawater, and OTB

seawater). Recruited bacterial communities formed three clusters in a principal coordinate

analysis based on Bray-Curtis distances (Fig. 6). PERMANOVA results further indicated that

bacterial source significantly affected recruited community composition (R^2 = 0.67542, F =

16.127, p = 0.001; 999 permutations), whereas size fraction and media formulation did not

significantly affect community composition (p > 0.05). Visualizing the relative abundance of

bacterial orders in each sample further demonstrates the distinct communities recruited from

each source (Fig. 7). Moreover, the FWC1006 *P. bahamense* culture clearly recruited bacteria

from the various sources; only three bacterial orders were detected in the initial FWC1006

culture microbiomes, despite deep Illumina metabarcode sequencing (394,399–621,150 reads

per sample after all filterings steps), whereas 10–12 bacterial orders were detected after

recruiting new bacterial communities.

Since only cultures with bacteria recruited from IRL seawater were able to grow in media

without vitamin B_{12}, we identified bacteria that were exclusively present in these cultures.

Alteromonadaceae, Saprospiraceae, and Enterobacteriaceae were consistently present in

communities recruited from IRL seawater but not from OTB seawater or IRL culture (Fig. 8).

Alteromonas spp., which typically require B_{12} but can not synthesize it [39], were abundant in

the smaller size fraction of communities recruited from IRL seawater when B_{12} was not included

in the media. *Klebsiella* and *Enterobacter*—both B_{12} producers [40, 41]—were present in all

communities recruited from the IRL seawater but not from OTB seawater or IRL culture.

Similarly, *Lewinella* was present in all communities recruited from IRL seawater but not from

OTB seawater or IRL culture. In contrast, *Ruegeria* and *Sulfitobacter* (Rhodobacteraceae),

genera often detected in phytoplankton cultures, were present in all cultures (Fig. 8).

Fig. 6 Principal coordinate analysis (PCoA) of Bray-Curtis distances between the bacterial communities

recruited by *P. bahamense* FWC1006 and the source surface seawater bacterial communities. FWC1006

cultures with each bacterial community in replete L1/4 media and L1/4–B_{12} media were sacrificed for 16S

rRNA gene sequencing at the end of round two in Fig. 5C. Three 95% confidence ellipses drawn with the

ggplot2 stat_ellipse function highlight the groupings within the data. The replicates with bacteria from OTB

seawater separate from all other samples on the primary axis, while the samples with bacteria from the

FWC1003 (IRL) culture and IRL seawater separate on the secondary axis. The bacterial communities in

the source seawater grouped most closely with the bacterial communities in culture that were recruited

from IRL seawater (bottom right quadrant), with the IRL seawater source communities within the 95%

confidence ellipse for the communities recruited from IRL seawater.

Fig. 7 Relative abundance of bacterial orders in source seawater samples and in cultures with bacterial communities recruited from IRL P. bahamense culture, OTB seawater, and IRL seawater. Orders that made up less than 1% in a given sample were combined to reduce the number of orders included in the plot.

Fig. 8 Relative abundance of genera in groups detected primarily in communities recruited from IRL seawater (Alteromonadaceae, Saprospiracea, and Enterobacteracea). Rhodobacteracea is included as a comparison. Rhodobacteracea are commonly found in association with phytoplankton and other algae. *Ruegeria* and *Sulfitobacter* were detected in all sample types.

Discussion

HAB-forming dinoflagellates generally require B_{12} for growth and often also require B_1 or

B_7 [9]. Despite this, HABs frequently form in regions with undetectable B_{12} concentrations,

suggesting that B_{12} may be being drawn down below detection limits or that cryptic B_{12} cycling

from bacteria is key to supporting HABs [10]. Here, we exclusively tested the cobalamin (B_{12}),

thiamine (B_1), and biotin (B_7) requirements of *P. bahamense* var. b*ahamense*—a dinoflagellate

that causes toxic HABs in Florida estuaries [19], but whose bioluminescence makes it culturally

and economically important in the Caribbean [42]—by utilizing a strain isolated from Florida

(OTB) with a reduced culture microbiome (Fig. 1). Our results clearly demonstrate that *P.*

bahamense has an absolute B_{12} requirement for growth and bloom formation. Only the bacterial

community recruited from IRL seawater collected during a *P. bahamense* bloom restored *P.*

bahamense growth in media without B_{12}. These results suggest that bacteria present during *P.*

bahamense blooms may contribute to fueling blooms through B_{12} production. While bacterial

communities that restored P. bahamense growth (from IRL seawater) when B_{12} was withheld

overlapped in composition with those that did not (IRL culture and OTB seawater), several

specific bacterial genera were found exclusively in the growth-restoring communities. Among

these, *Klebsiella* and *Enterobacter* are known B_{12} producers [41, 43] and may be introduced to

the environment through wastewater or runoff [44, 45]. Thus, anthropogenic inputs may be

contributing to *P. bahamense* blooms in Florida estuaries by introducing or otherwise supporting

B_{12}-synthesizing bacteria.

B-vitamin requirements of toxigenic dinoflagellates

Results from this study demonstrate that *P. bahamense* from OTB requires an external

source of vitamin B_{12}, but not necessarily B_1 or B_7, for growth. However, since the culture used

in the experiments had a reduced microbiome rather than being fully axenic, it remains possible

that the remaining bacteria supplied B_1 or B_7 and obscured *P. bahamense*'s requirement of

those vitamins. These results are in line with previous genomic surveys [8] and bioassays [9],

indicating that most HAB dinoflagellates are B_{12} auxotrophs. While *P. bahamense* was not

included in these studies, Santos and Azana [46] found that maximum biomass tripled in

antibiotic-treated cultures of *P. bahamense* var. *compressum* when the B-vitamin concentrations

in f/2 media [47] were also tripled. Together, these results strengthen the case that B_{12}

auxotrophy is a commonly shared trait among toxigenic dinoflagellates.

The *P. bahamense* strain used in this study required > 0.369 pM B_{12} to reach a

chlorophyll fluorescence commensurate with 100,000 cells L^{-1}, the minimum cell density

designated by Florida state regulatory agencies as a bloom (Fig. 4, S4). The *P. bahamense*

growth rate stabilized at 0.369 pM B_{12} and remained stable at higher B_{12} concentrations, even

though higher B_{12} concentrations further increased the maximum chlorophyll fluorescence

reached (Fig. 4). Similarly, the saxitoxin-producing dinoflagellates *Alexandrium pacificum* and

Gymnodinium catenatum required > 0.34 and $> 1. 52$ pM B_{12}, respectively, for measurable

growth [8]. The brevetoxin-producing *Karenia brevis* requires $> 0.01–1.0$ pM B_{12} for measurable

growth, depending on the strain tested [8, 9]. Thus, in addition to having an absolute B_{12}

requirement, toxigenic dinoflagellates may have especially high B_{12} needs due to their unique

physiologies.

Phytoplankton are generally considered auxotrophic for B_{12} if they lack the gene

encoding B_{12}-independent methionine synthase, but B_{12} may support dinoflagellate growth

through other mechanisms as well [48]. Vitamin B_{12} is a cofactor for methylmalonyl-CoA mutase

(MCM)—the enzyme that converts methylmalonyl-CoA to succinyl-CoA and is required for the

breakdown of odd-chain fatty acids and several amino acids. The metabolites from MCM-

catalyzed breakdown of amino acids are directed into the Krebs cycle, supporting energy

production in eukaryotic cells [48, 49]. Moreover, S-adenosylmethionine (SAM), the activated

form of methionine, acts as a universal methyl donor. SAM cycle methylation is key for

regulating gene expression through DNA and histone methylation, and is also central to

saxitoxin biosynthesis [8]. Given the large genomes characteristic of dinoflagellates [50],

extensive methylation may be required for gene regulation [51], which would increase usage of

the SAM cycle and elevate B_{12} demand.

Bacteria in seawater collected during a bloom supported P. bahamense *growth*

Many studies have demonstrated that bacteria can fulfill phytoplankton B_{12} needs (e.g.,

[23, 36, 52]), including those of HAB-forming dinoflagellates (e.g., [17, 22]). Therefore, we

tested whether bacteria recruited from several sources could rescue *P. bahamense* from B_{12}

limitation. Only the bacterial community recruited from IRL seawater collected during a *P.*

bahamense bloom was able to support *P. bahamense* growth in media without B_{12}, although

these cultures only reached about half the total biomass of cultures in replete media (Fig. 5).

The maximum Chl-*a* fluorescence in these cultures was 1,140 RFU, just shy of 1,457 RFU—the

value corresponding to the 100,000 cells L^{-1} bloom level (Fig. S4). Moreover, the growth rates in

cultures with bacteria from IRL seawater were similar in replete media and media missing B_{12}.

Together, these results suggest that bacteria present during the *P. bahamense* bloom in IRL

and transferred to our culture may supply B_{12} and support bloom development.

To evaluate which bacteria in the IRL seawater may provide B_{12}, we identified taxa that

were exclusively recruited from this source. Genera in the families Saprospiraceae (*Lewinella*)

and Enterobacteriaceae (*Klebsiella* and *Enterobacter*) were consistently present in cultures with

bacteria from IRL seawater and absent in cultures with bacteria from OTB seawater or IRL *P.*

bahamense culture. *Lewinella* is not a known B_{12}-producer, but is commonly associated with

phytoplankton [53] and macroalgae [54], and may be particularly suited to degrading algal

polysaccharides [53]. *Klebsiella* species are prolific B_{12} producers and are added to fermented

foods for B_{12} enrichment [41]. *Enterobacter* species are similarly B_{12} producers [43]. In addition

to being found in foods, *Klebsiella* and *Enterobacter* reside in mammalian small intestines [55]

and are found in wastewater [44, 45]. *Alteromonas spp.*, which are typically B_{12} auxotrophs [39],

were most abundant in the smaller size fraction of communities recruited from IRL seawater

when B_{12} was not included in the media, suggesting they may also benefit from B_{12} produced by

Enterobacteraceae or other bacteria recruited from IRL. In contrast, Rhodobacteraceae

genera—*Ruegeria* and *Sulfitobacter*—that are often found in phytoplankton cultures and

associated with phytoplankton blooms [56] were found in all of our cultures and were not

specific to those that grew without B_{12} added to media (Fig. 6, 7).

Surprisingly, bacteria recruited from the IRL culture did not support *P. bahamense*

growth in the absence of B_{12}, despite the IRL culture microbiome being fairly diverse and likely

representing the seawater community when the culture was initiated. The diversity of culture

microbiomes can diminish with culture age [56], and B_{12}-producing bacteria may be

outcompeted if the media is replete in B_{12}. The OTB seawater may have originally hosted B_{12}-

producing bacteria that were lost in the extensive filtering used to remove eukaryotic

phytoplankton from the inoculum, but it is also plausible that B_{12}-producing bacteria were absent

at the time of collection since *P. bahamense* was also absent. Notably, this study reveals that

distinct communities were recruited to *P. bahamense* from three sources, suggesting that *P.*

bahamense is opportunistic or a generalist in its associations with bacteria.

In some cases, bacteria associated with phytoplankton cultures or blooms compete for

essential nutrients [36, 57, 58]. Here, B_{12}-producing bacteria may have been present in the

communities recruited from OTB seawater, but other community members outcompeted *P.*

bahamense for the resource. For instance, *Caulobacter* can change morphology and motility to

survive oligotrophic conditions, making them especially good competitors [59]. Caulobacterales

were present in all recruited bacterial communities, but were most abundant in communities

recruited from OTB seawater. Thus, bacterial competition for essential resources may have

affected the growth of the *P. bahamense* grown with bacteria from OTB seawater.

Conclusions & future directions

The results from this study demonstrate that *P. bahamense* is auxotrophic for B_{12} and

that potentially anthropogenic bacteria present in the IRL may supply B_{12} to support bloom

formation. Bacteria from OTB seawater did not support *P. bahamense* growth when B_{12} was

withheld, but these bacteria were collected when *P. bahamense* was not detected in the

environment. Future work should investigate B_{12} sources that support *P. bahamense* bloom

formation by seeking to identify B_{12}-producing bacteria that co-bloom with *P. bahamense* or

determining if B_{12} is directly introduced to the ecosystem in wastewater. OTB and IRL are highly

urbanized and receive excess nutrients from wastewater treatment plants [60], and the

prolonged residence times [61] in these estuaries may also lead to an accumulation of B_{12}.

Given that increased B_{12} levels could exacerbate blooms, anthropogenic sources of B_{12} should

be investigated in tandem with B_{12}-synthesizing bacteria. Ultimately, measuring and tracing B_{12}

in OTB and the IRL could disentangle the sources of B_{12} to these ecosystems and the

relationship between anthropogenic B_{12} input and B_{12}-producing bacteria with HABs. Such an

effort may advance our ability to predict the severity of not just *P. bahamense* blooms, but other

HABs as well.

Acknowledgements

Research was funded by the University of South Florida College of Marine Science (USF CMS),

with additional support from the Phycological Society of America Grants-in-Aid of Research and

the Sanibel Captiva Shell Club grant. LRH was supported by the Abby Sallenger Endowed

Fellowship in Marine Science and the Anne and Werner Von Rosenstiel Fellowship in Marine

Science, awarded through USF CMS. We would like to thank the Florida Fish and Wildlife

Conservation Commission (FWC) Fish and Wildlife Research Institute (FWRI) for *P. bahamense*

culture loans and seawater sampling. Illumina metabarcode sequencing was performed by

Georgia Genomics and Bioinformatics Core at the University of Georgia (GGBC, UG Athens,

GA, RRID:SCR_010994).

Author Contributions

MMB planned the project. LRH, LD, and MMB performed research. LRH and MMB analyzed

data and wrote the manuscript. CL and SS reviewed and edited manuscript drafts.

Data Availability

All sequence data from this project is available on NCBI SRA under PRJNA1315153. Code,

data analysis pipelines, and raw data are available in a GitHub repository:

https://github.com/lruggles1/Pyrodinium_bahamense_growth.git. The reproducible R Markdown

analysis pipeline is further available as an interactive HTML file:

https://lruggles1.github.io/Pyrodinium_bahamense_growth/pyro_growth_meta.html.

Competing interests

The authors declare no competing interests.

References

1. Beardall J, Stojkovic S, Larsen S. Living in a high CO_2 world: impacts of global climate change on marine phytoplankton. *Plant Ecol Divers* 2009; **2**: 191–205.

2. Longhurst A, Sathyendranath S, Platt T, Caverhill C. An estimate of global primary production in the ocean from satellite radiometer data. *J Plankton Res* 1995; **17**: 1245–1271.

3. Sekerci Y, Ozarslan R. Oxygen-plankton model under the effect of global warming with nonsingular fractional order. *Chaos Solitons Fractals* 2020; **132**: 109532.

4. Glibert PM, Berdalet E, Burford MA, Pitcher GC, Zhou M. Global Ecology and Oceanography of Harmful Algal Blooms. 2018. Springer.

5. Anderson DM, Cembella AD, Hallegraeff GM. Progress in understanding harmful algal blooms: paradigm shifts and new technologies for research, monitoring, and management. *Ann Rev Mar Sci* 2012; **4**: 143–176.

6. Gobler CJ. Climate Change and Harmful Algal Blooms: Insights and perspective. *Harmful Algae* 2020; **91**: 101731.

7. Croft MT, Warren MJ, Smith AG. Algae need their vitamins. *Eukaryot Cell* 2006; **5**: 1175–1183.

8. Lin S, Hu Z, Song X, Gobler CJ, Tang YZ. Vitamin B_{12}-auxotrophy in dinoflagellates caused

by incomplete or absent cobalamin-independent methionine synthase genes (metE).

Fundamental Research 2022; **2**: 727–737.

9. Tang YZ, Koch F, Gobler CJ. Most harmful algal bloom species are vitamin B_1 and B_{12}

auxotrophs. *Proc Natl Acad Sci U S A* 2010; **107**: 20756–20761.

10. Sañudo-Wilhelmy SA, Cutter LS, Durazo R, Smail EA, Gómez-Consarnau L, Webb EA, et

al. Multiple B-vitamin depletion in large areas of the coastal ocean. *Proc Natl Acad Sci U S*

A 2012; **109**: 14041–14045.

11. Gobler CJ, Norman C, Panzeca C, Taylor GT, Sañudo-Wilhelmy SA. Effect of B-vitamins

(B_1, B_{12}) and inorganic nutrients on algal bloom dynamics in a coastal ecosystem. *Aquat*

Microb Ecol 2007; **49**: 181–194.

12. Fang H, Kang J, Zhang D. Microbial production of vitamin B_{12}: a review and future

perspectives. *Microb Cell Fact* 2017; **16**: 15.

13. Sultana S, Bruns S, Wilkes H, Simon M, Wienhausen G. Vitamin B_{12} is not shared by all

marine prototrophic bacteria with their environment. *ISME J* 2023; **17**: 836–845.

14. Wagner-Döbler I, Ballhausen B, Berger M, Brinkhoff T, Buchholz I, Bunk B, et al. The

complete genome sequence of the algal symbiont *Dinoroseobacter shibae*: a hitchhiker's

guide to life in the sea. *ISME J* 2010; **4**: 61–77.

15. Durham BP, Sharma S, Luo H, Smith CB, Amin SA, Bender SJ, et al. Cryptic carbon and

sulfur cycling between surface ocean plankton. *Proc Natl Acad Sci U S A* 2015; **112**: 453–

457.

16. Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. Algae acquire vitamin B_{12}

through a symbiotic relationship with bacteria. *Nature* 2005; **438**: 90–93.

17. Fei C, Booker A, Klass S, Vidyarathna NK, Ahn SH, Mohamed AR, et al. Friends and foes:

symbiotic and Algicidal bacterial influence on *Karenia brevis* blooms. *ISME Commun* 2024;

ycae164.

18. Ren C-Z, Gao H-M, Dai J, Zhu W-Z, Xu F-F, Ye Y, et al. Taxonomic and bioactivity

characterizations of *Mameliella alba* strain LZ-28 isolated from highly toxic marine

dinoflagellate *Alexandrium catenella* LZT09. *Mar Drugs* 2022; **20**: 321.

19. Phlips JE, Badylak S, Bledsoe E, Cichra M. Factors affecting the distribution of *Pyrodinium*

bahamense var. *bahamense* in coastal waters of Florida. *Mar Ecol Prog Ser* 2006; **322**: 99–

115.

20. Burkholder PR, Burkholder LM. Studies on B vitamins in relation to productivity of the Bahia

fosforescente, Puerto Rico. *Bull Mar Sci* 1958; **8**: 201–223.

21. Guillard RRL, Hargraves PE. *Stichochrysis immobilis* is a diatom, not a chrysophyte.
Phycologia . 1993. , **32**: 234–236

22. Cruz-López R, Maske H. The Vitamins B_1 and B_{12} required by the marine dinoflagellate
Lingulodinium polyedrum can be provided by its associated bacterial community in culture.
Front Microbiol 2016; **7**: 560.

23. Andrew S, Wilson T, Smith S, Marchetti A, Septer AN. A tripartite model system for
Southern Ocean diatom-bacterial interactions reveals the coexistence of competing
symbiotic strategies. *ISME Commun* 2022; **2**: 97.

24. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation of
general 16S ribosomal RNA gene PCR primers for classical and next-generation
sequencing-based diversity studies. *Nucleic Acids Res* 2013; **41**: e1.

25. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2.
Nat Biotechnol 2019; **37**: 852–857.

26. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. *Nat Methods* 2016; **13**: 581–583.

27. Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al. Optimizing
taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-
classifier plugin. *Microbiome* 2018; **6**: 90.

28. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal
RNA gene database project: improved data processing and web-based tools. *Nucleic Acids*
Res 2013; **41**: D590–6.

29. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. *PLoS One* 2013; **8**: e61217.

30. RStudio Team. RStudio: Integrated Development for R. RStudio, PBC.
http://www.rstudio.com/.

31. Guillard RRL, Morton SL. Culture methods. *Manual on harmful marine microalgae*
UNESCO, Paris 2003; 77–97.

32. Petzoldt T. *growthrates: Estimate Growth Rates from Experimental Data*. R package. 2025.

33. Matsuo Y, Komiya S, Yasumizu Y, Yasuoka Y, Mizushima K, Takagi T, et al. Full-length
16S rRNA gene amplicon analysis of human gut microbiota using MinION™ nanopore
sequencing confers species-level resolution. *BMC Microbiol* 2021; **21**: 35.

34. Lu J, Salzberg SL. Ultrafast and accurate 16S rRNA microbial community analysis using

Kraken 2. *Microbiome* 2020; **8**: 124.

35. Oxford Nanopore Technologies. Native Barcoding Kit 24 V14. *Oxford Nanopore*
Technologies.
https://community.nanoporetech.com/docs/prepare/library_prep_protocols/ligation-
sequencing-amplicons-native-barcoding-v14-sqk-nbd114-
24/v/nba_9168_v114_revm_15sep2022/overview-of-the-protocol?devices=minion.
Accessed 18 Aug 2024.

36. Mars Brisbin M, Mitarai S, Saito MA, Alexander H. Microbiomes of bloom-forming
Phaeocystis algae are stable and consistently recruited, with both symbiotic and
opportunistic modes. *ISME J* 2022; **16**: 2255–2264.

37. Câmara Dos Reis M, Romac S, Le Gall F, Marie D, Frada MJ, Koplovitz G, et al. Exploring
the phycosphere of *Emiliania huxleyi*: From bloom dynamics to microbiome assembly
experiments. *Mol Ecol* 2023; **32**: 6507–6522.

38. Isaac A, Francis B, Amann RI, Amin SA. Tight adherence (tad) pilus genes indicate putative
niche differentiation in phytoplankton bloom associated Rhodobacterales. *Front Microbiol*
2021; **12**: 718297.

39. Mars Brisbin M, Schofield A, McIlvin MR, Krinos AI, Alexander H, Saito MA. Vitamin B_{12}
conveys a protective advantage to phycosphere-associated bacteria at high temperatures.
ISME Commun 2023; **3**: 88.

40. Lawrence JG, Roth JR. Evolution of coenzyme B_{12} synthesis among enteric bacteria:
evidence for loss and reacquisition of a multigene complex. *Genetics* 1996; **142**: 11–24.

41. Keuth S, Bisping B. Vitamin B_{12} production by *Citrobacter freundii* or *Klebsiella pneumoniae*
during tempeh fermentation and proof of enterotoxin absence by PCR. *Appl Environ*
Microbiol 1994; **60**: 1495–1499.

42. Soler-Figueroa BM, Otero E. The influence of rain regimes and nutrient loading on the
abundance of two dinoflagellate species in a tropical bioluminescent bay, Bahía
fosforescente, La Parguera, Puerto Rico. *Estuaries Coast* 2015; **38**: 84–92.

43. Chung HJ, Fields ML. Production of Riboflavin and Vitamin B_{12} by *Bacillus megaterium*
ATCC 13639 and *Enterobacter aerogenes* in Corn Meal. *J Food Sci* 1986; **51**: 1514–1517.

44. Rajagopalan K. Studies on vitamin B_{12} production by strains of *Klebsiella pneumoniae* and
Bacillus pasteurii. *J Appl Bacteriol* 1976; **40**: 111–114.

45. Kvesić M, Šamanić I, Novak A, Fredotović Ž, Dželalija M, Kamenjarin J, et al. Submarine
outfalls of treated wastewater effluents are sources of extensively- and multidrug-resistant

KPC- and OXA-48-producing Enterobacteriaceae in coastal marine environment. *Front*
Microbiol 2022; **13**: 858821.

46. Santos MAG, Azanza RV. Responses of *Pyrodinium bahamense* var. *compressum* and
associated cultivable bacteria to antibiotic treatment. *J Appl Phycol* 2012; **24**: 825–835.

47. Guillard RR, Ryther JH. Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt,
and *Detonula confervacea* (cleve) Gran. *Can J Microbiol* 1962; **8**: 229–239.

48. Helliwell KE. The roles of B vitamins in phytoplankton nutrition: new perspectives and
prospects. *New Phytol* 2017; **216**: 62–68.

49. Forny P, Froese DS, Suormala T, Yue WW, Baumgartner MR. Functional characterization
and categorization of missense mutations that cause methylmalonyl-CoA mutase (MUT)
deficiency. *Hum Mutat* 2014; **35**: 1449–1458.

50. Wang H, Wu P, Xiong L, Kim H-S, Kim JH, Ki J-S. Nuclear genome of dinoflagellates: Size
variation and insights into evolutionary mechanisms. *Eur J Protistol* 2024; **93**: 126061.

51. Ho P, Kong KF, Chan YH, Tsang JSH, Wong JTY. An unusual S-adenosylmethionine
synthetase gene from dinoflagellate is methylated. *BMC Mol Biol* 2007; **8**: 87.

52. Costas-Selas C, Martínez-García S, Pinhassi J, Fernández E, Teira E. Unveiling
interactions mediated by B vitamins between diatoms and their associated bacteria from
cocultures. *J Phycol* 2024; **60**: 1456–1470.

53. Gómez-Letona M, Arístegui J, Riebesell U, Sebastián M. Consistent prokaryotic
successional dynamics across contrasting phytoplankton blooms. *Limnol Oceanogr* 2025;
70: 493–508.

54. Miranda LN, Hutchison K, Grossman AR, Brawley SH. Diversity and abundance of the
bacterial community of the red Macroalga *Porphyra umbilicalis*: did bacterial farmers
produce macroalgae? *PLoS One* 2013; **8**: e58269.

55. Albert MJ, Mathan VI, Baker SJ. Vitamin B_{12} synthesis by human small intestinal bacteria.
Nature 1980; **283**: 781–782.

56. Mars Brisbin M, Acord M, Davitt R, Bent S, Van Mooy BAS, Flaum E, et al. Exploring the
Phaeosphere: Characterizing the microbiomes of *Phaeocystis antarctica* colonies from the
coastal Southern Ocean and laboratory culture. *J Phycol* 2025; **61**: 1124–1139.

57. Cai G, Yu X, Wang H, Zheng T, Azam F. Nutrient-dependent interactions between a marine
copiotroph *Alteromonas* and a diatom *Thalassiosira pseudonana*. *MBio* 2023; **14**:
e0094023.

58. Calfee BC, Glasgo LD, Zinser ER. *Prochlorococcus* exudate stimulates heterotrophic

691 bacterial competition with rival phytoplankton for available nitrogen. *MBio* 2022; **13**:
692 e0257121.

693 59. Hallgren J, Koonce K, Felletti M, Mortier J, Turco E, Jonas K. Phosphate starvation
694 decouples cell differentiation from DNA replication control in the dimorphic bacterium
695 *Caulobacter crescentus*. *PLoS Genet* 2023; **19**: e1010882.

696 60. Greening H, Janicki A, Sherwood ET, Pribble R, Johansson JOR. Ecosystem responses to
697 long-term nutrient management in an urban estuary: Tampa Bay, Florida, USA. *Estuar*
698 *Coast Shelf Sci* 2014; **151**: A1–A16.

699 61. Meyers SD, Luther ME. A numerical simulation of residual circulation in Tampa bay. Part II:
700 Lagrangian residence time. *Estuaries Coast* 2008; **31**: 815–827.